Development and validation of Arc nanobodies:

new tools for probing Arc dynamics and function

Yuta Ishizuka[a,*], Tadiwos F. Mergiya[a,b], Rodolfo Baldinotti[a,b], Ju Xu[a], Erik I. Hallin[a], Sigurbjörn

Markússon[a], Petri Kursula[a,c,d], Clive R. Bramham[a,b**]

From:

[a]Department of Biomedicine, University of Bergen, Bergen, Norway.

[b]Mohn Center for Research on the Brain, University of Bergen, Norway

[c]Faculty of Biochemistry and Molecular Medicine, University of Oulu, Oulu, Finland.

[d]Biocenter Oulu, University of Oulu, Finland

*Corresponding author:

Yuta Ishizuka, PhD.

Department of Biomedicine, University of Bergen, Jonas Lies 91, 5009 Bergen, Norway

[Present Affiliation and Address]

Department of Pathophysiology and Metabolism, Kawasaki Medical School, 577 Matsushima,

Kurashiki, Okayama 701-0192, Japan

E-mail: yuta.ishizuka@med.kawasaki-m.ac.jp; yishizuka1110@gmail.com

**Corresponding author:

Clive R. Bramham, MD, PhD.

Department of Biomedicine, University of Bergen, Jonas Lies 91, 5009 Bergen, Norway

Phone: +47 55 58 60 32; E-mail: clive.bramham@uib.no

Abstract

Activity-regulated cytoskeleton-associated (Arc) protein plays key roles in long-term synaptic plasticity, memory, and cognitive flexibility. However, an integral understanding of Arc mechanisms is lacking. Arc is proposed to function as an interaction hub in neuronal dendrites and the nucleus, yet Arc can also form retrovirus-like capsids with proposed roles in intercellular communication. Here, we sought to develop anti-Arc nanobodies (ArcNbs) as new tools for probing Arc dynamics and function. Six ArcNbs representing different clonal lines were selected from immunized alpaca. Immunoblotting with recombinant ArcNbs fused to a small ALFA-epitope tag demonstrated binding to recombinant Arc as well as endogenous Arc from rat cortical tissue. ALFA-ArcNb also provided efficient immunoprecipitation of stimulus-induced Arc after carbachol-treatment of SH-SY5Y neuroblastoma cells and induction of long-term potentiation in the rat dentate gyrus *in vivo*. Epitope mapping showed that all Nbs recognize the Arc C-terminal region containing the retroviral Gag capsid homology domain, comprised of tandem N- and C-lobes. ArcNbs E5 and H11 selectively bound the N-lobe, which harbors a peptide ligand binding pocket specific to mammals. Four additional ArcNbs bound the region containing the C-lobe and terminal tail. For use as genetically encoded fluorescent intrabodies, we show that ArcNbs fused to mScarlet-I are uniformly expressed, without aggregation, in the cytoplasm and nucleus of HEK293FT cells. Finally, mScarlet-I-ArcNb H11 expressed as intrabody selectively bound the N-lobe and enabled co-immunoprecipitation of full-length intracellular Arc. ArcNbs are versatile tools for live-cell labeling and purification of Arc and analysis of capsid domain specific functions.

Keywords: Arc, N-lobe; C-lobe, nanobody, intrabody, chromobody, co-immunoprecipitation

1. Introduction

The immediate early gene encoding activity-regulated cytoskeleton-associated protein (Arc), also known as activity-regulated gene 3.1 (Arg3.1), is expressed mainly in excitatory neurons following neural activity to mediate functionally diverse actions [1]. In activity-dependent synaptic plasticity, such as long-term potentiation (LTP) and long-term depression, Arc protein is rapidly synthesized and degraded, indicating a transient and dynamic mode of action [2-5]. Stimulus-evoked expression of Arc is also required for memory, postnatal developmental plasticity of the visual cortex, and cognitive flexibility [3, 6-9]. These diverse functions are proposed to be mediated by distinct Arc-binding partner complexes in the postsynaptic compartment and the neuronal nucleus [10-14]. Arc protein is also capable for reversible self-association and formation of oligomers [15, 16]. Recently, Arc protein was shown to self-assemble into spheroid particles resembling retroviral Gag capsids [17-19]. Arc capsids are incorporated in extracellular vesicles and are capable of transferring RNA cargo from donor cells to recipient cells [17, 18]. However, an integral understanding that accounts for Arc hub signaling and capsid function is lacking.

Arc is a flexible protein comprised of two oppositely charged domains, N- and C-terminal domains (NTD and CTD), flanking a central disordered hinge region [15]. Structural studies and predictions indicate an antiparallel coiled-coil structure of the NTD [20]. The NTD mediates binding to phospholipid membrane [20, 21], and the NTD second coil harbors an oligomerization motif [113]MHVWREV[119] that is critical for Arc self-association including formation of higher-order oligomers and capsids, but is not required for dimer formation [19]. The CTD is a tandem domain comprised of two lobes, the N- and C-lobes (NL and CL), homologous to the capsid domain of retroviral Gag polyprotein [22, 23]. Arc CTD is important for formation of stable and double-shelled Arc capsid, which is dependent on NTD-mediated oligomerization [18, 19, 24]. In addition, the mammalian Arc NL has a unique hydrophobic peptide binding pocket that interacts with several postsynaptic proteins, including stargazin, guanylate kinase-associated protein (GKAP), Wiskott-Aldrich syndrome protein family member 1 (WAVE1), and GluN2A [14, 20, 22, 25]. Taken

together, this suggests that interactions between domains regulates oligomerization and protein-protein

interactions to determine Arc activity-state. However, tools that would allow labeling or manipulation

of endogenous Arc in situ are lacking.

Single-domain antibodies (sdAb), also known as variable domain of heavy chain of heavy chain

(VHH) antibody and Nanobody® (Nb), are derived from the heavy chain antibody of *Camelidae*

[26]. Nbs have unique properties relative to conventional antibodies that make them attractive for use

as research tools and development of therapeutics [27]. Nbs are small (15 kDa), have high chemical

stability, can be genetically-encoded for expression in mammalian cells as intrabodies, and they can

bind hidden epitopes in small cavities that are not recognized by antibodies [28-32].

Here, we sought to develop and validate experimental applications for six anti-Arc nanobodies

(ArcNbs: B5, B12, C11, D4, E5, and H11). ArcNbs fused to a small ALFA epitope were successfully

used for immunoblotting and immunoprecipitation of endogenous Arc. ArcNb E5 and H11

specifically recognized the capsid domain NL, while the other Nbs bound the segment containing the

CL and C-terminal tail. Genetically-encoded ArcNb fused to fluorescent mScarlet-I expressed

efficiently in mammalian cells as intrabodies and allowed immunoprecipitation of intracellular Arc.

The ArcNbs provide versatile tools for labeling, tracking, and purifying native intracellular Arc, with

further potential use as capsid domain-specific inhibitors.

93 2. Materials and Methods

94 2.1. Generation and modifications of ArcNbs

Procedures for generation and modification of ArcNbs were conducted by NanoTag

Biotechnologies GmbH (Göttingen, Germany) (**Fig. 1**). For detailed information on the purification,

structural characterization, and binding of the untagged ArcNbs, see Markússon et al. (2021). Briefly,

two alpacas were immunized six times starting with purified recombinant human wild-type (WT)

Arc protein (immunization 1 and 2), then a mixture of Arc WT and purified mutant Arc protein

(Arc$^{s113-119A}$: amino acid residues from 113 to 119 were substituted to alanine) with increasing

proportion of Arc$^{s113-119A}$ (immunization 3-5), and finally with injection of Arc$^{s113-119A}$ alone. A strong

general immune-response was confirmed using a serum-based ELISA in one of the animals. Total

RNA was extracted from peripheral blood mononuclear cell (PBMC) preparations obtained from

whole blood. cDNA encoding Nbs were reverse transcribed using a nested PCR that specifically

amplifies the coding regions for IgG2 and IgG3 VHH fragments. The complete PCR product was

cloned into a phagemid vector suitable for phage display followed by bio-panning with purified

mutant Arc$^{s113-119A}$ protein as antigen. Then, Nb sequences extracted from phages binding to antigen

were amplified by PCR and cloned in the screening vector. Ninety-six Nb clones were separately

expressed in *E. coli.* and validated by ELISA using Arc$^{s113-119A}$ as target protein. Six clones (B5, B12,

C11, D4, E5, and H11) were chosen based on their sequences and ELISA-based binding evaluations.

To generate recombinant Nbs suitable for biochemical analyses, such as immunoblotting and

immunoprecipitation, a small ALFA-epitope tag (SRLEEELRRRLTE) [33], which forms a stable

alpha-helix on target protein, was added to the Nb C-terminus. For expressing ArcNbs as

intrabody/chromobody, the Nb coding regions were subcloned from the bacterial expression vector

into a mammalian expression vector harboring mScarlet-I (mSc) driven by the CMV promoter. We

also subcloned the coding region of all ArcNbs into the AcGFP-N1 vector. AcGFP1-N1 was a gift

from Michael Davidson (Addgene plasmid # 54705; http://n2t.net/addgene:54705; RRID:Addgene

54705).

2.2. Purification of ALFA-tagged ArcNbs

ALFA-tagged ArcNbs (ALFA-ArcNbs) were expressed in BL21 (DE3) cells and purified

according to a general methods for protein purification [34] with minor modifications. Briefly, cells

were grown at 37ºC in LB medium overnight. The cultures were transferred to ZY5052 medium for

auto-induction of protein expression at 37ºC for 4 hours, then at 30ºC for 24 hours. The cells were

lysed in lysis/washing buffer containing 40 mM Hepes (pH 7.5), 100 mM NaCl, and 20 mM

Imidazole, by one freeze-thaw cycle followed by sonication. The lysates were centrifuged at 16,000

x g for 30 min at 4ºC and loaded onto a Ni-NTA resin. After washing resin with lysis/washing buffer,

ALFA-ArcNbs were eluted by elution buffer containing 40 mM Hepes (pH 7.5), 100 mM NaCl, and

300 mM Imidazole. His-tagged TEV protease was added to the elute, and the samples were dialyzed

against dialysis buffer containing 40 mM Hepes (pH 7.5), 100 mM NaCl and 1 mM dithiothreitol

(DTT) overnight at 4ºC. The dialyzed samples were passed through a Ni-NTA resin again to remove

the TEV protease and the cleaved His10-tag. The Ni-NTA flow-through was loaded on a

chromatography column (HiLoad® 16/60 Superdex® 75 PG, Cytiva, Marlborough, MA),

equilibrated with degassed buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl). Size exclusion

chromatography was run at 1 ml/min at 4ºC on a ÄKTA pure chromatography system (Cytiva). All

ALFA-ArcNbs gave one major peak in the chromatogram. The protein size and purity were analyzed

by SDS-PAGE, giving one strong Coomassie-stained band of the expected size (**Fig. S1**).

2.3. Animals, electrophysiology, and tissue collection

For naive cortical tissue collection, adult male Sprague-Dawley rats were anesthetized using

urethane (1.5 g/kg intraperitoneal) and decapitated before dissecting out cortical tissues. The tissues

were kept in −80ºC until experiments. LTP in the dentate gyrus (DG) was induced according to

previously described methods [35]. Briefly, a bipolar stimulating electrode (7.9 mm posterior to the

bregma, 4.2 mm lateral to the midline) and recording electrode (3.9 mm posterior to bregma, 2.3 mm

lateral to midline) were inserted to stimulate medial perforant path fibers and record of evoked field

potentials in the dentate hilus, respectively. Test pulses were given every 30 s throughout the

experiment except during the period of high-frequency stimulation (HFS). 20 min baseline

recordings were acquired before three sessions of HFS with five min interval were given. Each

session consisted of four, 400 Hz stimulus trains (8 pulses/train) and the interval between trains was

10 s. Changes in the field excitatory post-synaptic potential slope were expressed in percent of

baseline to quantify induction and maintenance of LTP. Two hours after HFS, rats were decapitated,

and the dentate gyri were dissected out and stored at -80^{o}C for further use.

2.4. Cell culture: transfection and pharmacological stimulation

Human embryonic kidney 293FT (HEK293FT) and SH-SY5Y neuroblastoma cells were

obtained from Thermo Fisher Scientific (Waltham, MA) and ATCC (Manassas, VA), respectively.

Cells were maintained in Dulbecco's modified Eagle medium (DMEM/high-glucose, Sigma-Aldrich,

St. Louis, MO) supplied with 10% FBS (Sigma-Aldrich) and 100 U/ml Penicilin-Streptomycin

(Thermo Fisher Scientific) at 37 ºC in a humidified incubator with 5% CO_2. HEK293FT cells were

transfected using Lipofectamine^{TM} 2000 Transfection Reagent (Thermo Fisher Scientific) according

to the manufacturer's instructions. For induction of endogenous Arc expression, SH-SY5Y cells

were stimulated with 100 μM carbachol (CCh) for 1 h at +37 ºC in a humidified incubator with 5%

CO_2 [36].

2.5. Antibodies

Primary antibodies are as follows: mouse anti-Arc (C-7) (Santa Cruz Biotechnology, Dallas, TX,

Cat# sc-17839, RRID:AB_626696), rabbit anti-Arc (Synaptic Systems, Göettingen, Germany, Cat#

156 003, RRID:AB_887694), rabbit anti-GFP (Santa Cruz Biotechnology Cat# sc-8334,

RRID:AB_641123), mouse anti-mCherry (Takara Bio, Siga, Japan, Cat# 632543,

RRID:AB_2307319), anti-ALFA HRP-coupled sdAb (NanoTag Biotechnologies, Cat#. N1501-HRP).

Secondary antibodies are as follows: Goat Anti-Mouse IgG, H & L Chain Antibody, Peroxidase

Conjugated (Merck, Darmstadt, Germany, Cat# 401253, RRID:AB_437779) and Goat Anti-Rabbit

IgG, H & L Chain Specific Peroxidase Conjugate antibody (Merck Cat# 401315,

RRID:AB_2617117). Secondary antibodies for detecting immunoprecipitated proteins are as

follows: Peroxidase AffiniPure Goat Anti-Mouse IgG, light chain specific (Jackson ImmunoResearch

Labs, West Grove, PA, Cat# 115-035-174, RRID:AB_2338512) and Peroxidase IgG Fraction

Monoclonal Mouse Anti-Rabbit IgG, light chain specific (Jackson ImmunoResearch Labs, Cat#

211-032-171, RRID:AB_2339149).

2.6. Electrophoresis and immunoblotting

Native-polyacrylamide gel electrophoresis (PAGE) was performed according to previously

described methods [37]. Briefly, different amounts of purified mutant $Arc^{s113-119A}$ protein were

dissolved in native sample buffer, then subjected to nativePAGE using 4−15% Tris-Glycine

acrylamide gel. SDS-PAGE was performed as described previously [38, 39]. Briefly, cultured cells

and brain tissues were lysed and sonicated in RIPA buffer containing cOmplete™, EDTA-free

Protease Inhibitor Cocktail Tablet (Sigma-Aldrich) and PhosSTOP™, Phosphatase Inhibitor Cocktail

Tablet (Sigma-Aldrich). After centrifugation, the supernatants were collected, and protein

concentration was determined using the Micro BCA Protein Assay Kit (Thermo Fisher Scientific).

Samples were mixed with 4X Laemmli sample buffer (Bio-Rad Laboratories, Inc., Hercules, CA)

containing 50 mM DTT at final concentration, then boiled and subjected to SDS-PAGE. Separated

proteins in the gel then were transferred to polyvinylidene difluoride membrane (Thermo Fisher

Scientific). The membranes were incubated with the appropriate primary and secondary antibodies

after incubation with 10% bovine serum albumin (BSA) in Tris-buffered saline with 0.05% Tween 20

(TBS-T) at room temperature for 1 h. For detecting Arc protein immunolabeled by ArcNb,

anti-ALFA HRP-coupled sdAb was used instead of conventional secondary antibody. Peroxidase

activity was detected using chemiluminescence regent (Clarity Western ECL substrate, Bio-Rad

Laboratories, Inc.) and visualized by an image analyzer using Image Lab™ Software (Gel Doc™

XR+, Bio-Rad Laboratories, Inc.). Figure presentations were performed using ImageJ/FIJI (RRID:

RRID:SCR_002285).

2.7. Immunoprecipitation of endogenous Arc protein with ALFA-ArcNbs

Endogenous Arc immunoprecipitation was performed using ALFA-ArcNbs and ALFA Selector[ST]

as carrier [33]. CCh-treated SH-SY5Y cells and DG tissue from HFS-treated rats were lysed in

modified RIPA (M-RIPA) buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% NP-40, 0.25% sodium

deoxycholate, 1 mM EDTA) supplemented with cOmplete™ (Sigma-Aldrich) and PhosSTOP™

(Sigma-Aldrich). First, 20 µl of ALFA Selector[ST] resin washed twice in 1 ml M-RIPA buffer and

centrifuged at 1000 g for 1 min at 4 ^{o}C. Then, the resin was incubated with 2 µg of ALFA-ArcNb

H11 for 1 hour at 4^{o}C with head-over-tail rotation. Following washing, the ALFA-ArcNb H11 and

selector resin complex was incubated with lysates for 30 min at room temperature. After washing

resin, the samples were mixed with 2X Laemmli sample buffer containing 100 mM DTT at final

concentration and denatured by boiling at 95 ^{o}C for 5 min. The supernatants were subjected to

SDS-PAGE followed by immunoblotting for analyzing precipitated complexes.

2.8. Fluorescence cell imaging

Preparation of coverslips for cell imaging were performed according to previously described

methods [39]. HEK293FT cells seeded on coverslips in 12-well plates were transfected with

mSc-ArcNb expression vectors according to the manufacturer's instructions. Transfected cells were

fixed with 4% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4) for 20 min at room temperature.

After washing coverslips with phosphate-buffered saline (PBS, pH 7.4), the coverslips were mounted

in Prolong™ Diamond Antifade Mountant with DAPI (Thermo Fisher Scientific). Fluorescence

imaging was performed on AxioImager Z1 microscopy (Carl Zeiss AG, Oberkochen, Germany).

Transfected cells were imaged using Plan-ApoChromat 20x/0.75 NA (Carl Zeiss AG), EC plan

Neofluar 40x/1.30 Oil M27 (Carl Zeiss AG) objectives, and Axiocam 503 mono digital camera

controlled by ZEN pro software (Carl Zeiss AG). The captured fluorescence images were analyzed

using MetaMorph® imaging software (Molecular Devices, San Jose, CA) and ImageJ/Fiji.

2.9. Co-immunoprecipitation

HEK293FT cells were co-transfected with plasmids expressing mSc-ArcNb H11 and an

mTurquoise2 (mTq2)-tagged Arc construct. mTq2 was N-terminally fused to Arc full-length (FL), 2)

N-terminal region (NTR 1-140), 3) linker (135-216), 4) N-lobe (NL 208-277), or 5) C-lobe +

C-terminal tail (CL+tail 278-396). Transfected cells were lysed in modified RIPA (M-RIPA) buffer.

First, 2 μg of normal rabbit IgG (Merck, Cat# 12-370, RRID:AB_145841), anti-mCherry (mCh)

antibody against mSc-ArcNb (H11), or anti-Arc rabbit polyclonal antibody were absorbed to Protein

G Sepharose 4 Fast Flow (Cytiva) for 1 h at room temperature. Then, antibody/bead complexes were

mixed with the cell lysates overnight at +4 ^{o}C for co-IP. The sepharose beads were washed four times

with M-RIPA buffer and resuspended in 2X Laemmli sample buffer containing 100 mM DTT before

denaturing at 95 ℃ for 5 min. The supernatants were subjected to SDS-PAGE followed by

immunoblot analysis.

3. Results

3.1. ALFA-ArcNbs recognize purified recombinant Arc and endogenous Arc protein

Two alpacas were immunized with a combination of purified recombinant human WT Arc and rat mutant $Arc^{s113-119A}$ [40]. WT Arc self-associates and oligomerizes *in vitro* [15, 16]. We have recently shown that amino acid residues 113-119 are critical for high-order oligomerization [19]. Interestingly, the substitution mutant protein $Arc^{s113-119A}$ failed to form high-order oligomers or capsids. A mixture of WT and $Arc^{s113-119A}$ mutant protein was used as an antigen cocktail because oligomerization of WT Arc protein might lead to epitope masking preventing proper immune response. Following bio-panning assay using $Arc^{s113-119A}$ as a target, 96 clones were expressed and evaluated by ELISA assay using mutant $Arc^{s113-119A}$ as an antigen. Six clones (B5, B12, C11, D4, E5, and H11) representing different clonal lines were chosen based on their sequences and ELISA-based binding to purified $Arc^{s113-119A}$.

Firstly, we examined whether ALFA-ArcNbs have ability to detect purified recombinant Arc and endogenous Arc from rat cortical lysates. In immunoblot assays, ALFA-ArcNb bound to Arc was detected by using anti-ALFA HRP-coupled sdAb. All six ArcNbs bound to purified $Arc^{s113-119A}$ both in native and denatured conditions (**Fig. S2**). On native-PAGE gels (**Fig. S2A**), several immunoreactive bands were detected as low-order oligomeric forms. In the denatured condition on SDS-PAGE (**Fig. S2B**), only one prominent immunoreactive band was detected around 50 kDa corresponding to Arc monomer. Also, immunoblotting performed in cortical lysates from naïve rats showed that all ArcNbs detect endogenous Arc as a single band (~50 kDa) consistent with conventional anti-Arc antibodies (**Fig. 2A**). We then tested whether ALFA-ArcNbs are suitable for immunoprecipitation of endogenous, stimulus-induced Arc. CCh treatment of SH-SY5Y cells and HFS-induced LTP in the rat dentate gyrus *in vivo* are both associated with enhanced Arc expression [3, 36, 41]. SH-SY5Y cells were stimulated with 100 μM CCh for 1 h and subjected to immunoprecipitation using ALFA-ArcNb H11 and ALFA SelectorST resin, followed by SDS-PAGE

and immunoblotting using anti-Arc rabbit polyclonal antibody. CCh treatment induced a prominent

increase in Arc expression in whole lysate, shown as input samples in figures, relative to untreated

cells (**Fig. 2B**, closed arrowhead), and immunoprecipitation using ALFA-ArcNb H11 massively

enriched for Arc (**Fig. 2B**, opened arrowhead). Similarly, HFS resulted in ipsilateral enhancement of

Arc expression in the DG (**Fig. 2C**, closed arrowhead), which was strongly enriched in the

immunoprecipitation (**Fig. 2C,** opened arrowhead). These results validate all ALFA-ArcNbs for

immunoblotting and ALFA-ArcNb H11 for immunoprecipitation of endogenous Arc.

3.2. ALFA-ArcNbs E5 and H11 specifically binds to Arc-NL

Next, we performed epitope mapping by expressing Arc-FL or truncation mutants in HEK293FT

cells, followed by SDS-PAGE and immunoblot analysis with the six ALFA-ArcNbs. Cells were first

transfected with mTq2-Arc-FL (residues 1-396, lane 1), NTR (1-140, lane 2), linker region (135-216,

lane 3), and CTR (208-396, lane 4). All Nbs detected Arc-FL and CTR, but not the NTR or linker

region (**Fig. 3A**). To further specific region of binding, cells were transfected with mTq2-Arc-NL

(208-277, lane 5) and CL+tail (278-396, lane 6), then subjected to immunoblotting. Intriguingly, two

clones E5 and H11 specifically bound to NL (**Fig. 3B**, opened arrowheads) while all other ArcNbs

bound to the CL+tail region (**Fig. 3B**).

3.3. Expression of mScarlet-I-tagged and AcGFP-tagged ArcNbs as intrabodies in HEK293FT

cells

We tested intracellular expression of ArcNbs in mammalian cells as genetically encoded

intrabodies. HEK293FT cells were transfected with mSc-ArcNb expression vectors, then subjected

to SDS-PAGE followed by immunoblotting using anti-mCh antibody to detect mSc. Expression of

mSc-ArcNbs was confirmed by immunoblotting (**Fig. 4A**) and fluorescence microscopy (**Fig. 4B**).

All six mSc-ArcNbs exhibited uniformly bright cytoplasmic and nucleoplasmic fluorescence without

selective accumulation in subcellular compartments, formation of aggregates, or signs of

morphological abnormalities. We also assessed the expression pattern of AcGFP-ArcNbs (**Fig. S3A**)

AcGFP fused to either E5, C11, or H11 showed uniform fluorescence similar to the corresponding

mSc-ArcNbs. However, in contrast to the mSc-ArcNbs, three of the AcGFP-ArcNbs (B5, B12, and

D4) showed strong accumulation and aggregation in the nucleus (**Fig. S3B**). These results indicate

that all mSc-ArcNbs and several AcGFP-ArcNbs can be expressed as intrabodies without

aggregation or deleterious effects.

3.4. mScarlet-I-tagged ArcNb H11 intrabody specifically binds the N-lobe and enables

immunoprecipitation of intracellular Arc

We examined whether ArcNb, when expressed as intrabody, is able to bind intracellular Arc. We

focused this analysis on ArcNb H11, which efficiently immunoprecipitated endogenous Arc (**Fig 2**)

and selectively bound the NL in the epitope mapping analysis (**Fig. 3**). HEK293FT cells were

co-expressed with mTq2-Arc (FL) together with mSc-ArcNb H11 (mSc-H11), then subjected to

co-IP assay using either anti-mCh or rabbit anti-Arc polyclonal antibodies to immunoprecipitate the

Arc/ArcNb complex (design illustrated in **Fig. 5A**). As expected, mSc-H11 was immunoprecipitated

with anti-mCh antibody (**Fig. 5B**, closed arrowhead). Importantly, mSc-H11 in complex with

mTq2-Arc (FL) was co-immunoprecipitated by anti-mCh antibody, as detected by both anti-Arc and

GFP antibodies (**Fig. 5B**, opened arrowheads). In cells co-transfected with mTq2-Arc NTR or

mTq2-Arc linker, no interaction of mSc-11 with these Arc regions was detected, although successful

immunoprecipitation of mSc-H11 with anti-mCh antibody was confirmed (**Fig. 5C and D**, closed

arrowheads). To further assess the intracellular specificity of binding, cells were co-transfected with

mTq2-Arc (NL) or mTq2-Arc (CL+tail) together with mSc-H11. We found that H11 forms an

intracellular complex with the Arc NL (**Fig. 5E**, opened arrowhead) but not with the CL+tail (**Fig.**

5F). We also did a reverse co-IP assay using rabbit anti-Arc polyclonal antibody (**Fig. S4**) and found

that mSc-H11 was co-immunoprecipitated with Arc FL and NL (**Figs. S4B and E**, opened

arrowheads), but not with Arc NTR, linker, or CL+tail (**Figs. S4C, D and F**). Taken together, we

conclude that ArcNb H11 expressed as intrabody specifically binds the Arc NL and can be used to

affinity-purify intracellular full-length Arc.

4. Discussion

We developed and validated the first ArcNbs as new tools for probing Arc expression and

function. For biochemical analyses, the small ALFA epitope tag was fused to the C-terminus of

ArcNbs [33]. All clones of ALFA-ArcNbs successfully detected purified recombinant Arc protein

and endogenous Arc protein from rat brain tissues (**Figs. S2 and 2A**). We show that ALFA-ArcNbs

combined with anti-ALFA HRP-coupled sdAbs can be used for immunoblotting of Arc protein in the

same manner as conventional anti-Arc antibodies. All six ArcNbs bound to the Arc CTR and not to

the NTR or linker. Nbs E5 and H11 selectively bound to the CTD/capsid domain NL, while the

remaining four Nbs bound to the segment containing the CL and C-terminal tail (**Fig. 3**).

Using a mammalian expression vector harboring mSc or AcGFP fluorescent proteins, we

showed that all ArcNbs are suitable for applications as genetically encoded chromobodies (**Figs. 4A**

and S3B). mSc is a bright monomeric red fluorescent protein [42, 43]. AcGFP1 is a monomeric

green fluorescent protein, obtained from *Aequorea coerulescens*, with similar spectral properties to

EGFP. Despite 94% amino acid sequence homology and equivalent brightness, AcGFP1 is superior

to EGFP for fusion applications because it is monomeric [44] unlike EGFP that forms dimers [45,

46]. Although all mSc-ArcNbs showed uniform cytosolic and nuclear expression, three of the

AcGFP-ArcNbs (B5, B12 and D4) aggregated in the nucleus (**Fig. S3B**). Further experiments are

needed to understand mechanism of nuclear aggregation of these AcGFP-ArcNbs.

In the study of Markússon *et al.* (2021) on the generation and structural characterization of

ArcNbs, purified untagged Nbs served as chaperones for crystallization of mammalian Arc. Atomic

structures were obtained of ArcNb H11 and C11 bound to NL and CL, respectively [40]. Moreover,

the complementarity determining region 3 (CDR3) of ArcNb H11 was shown to bind inside the

hydrophobic pocket of the NL [40]. The CDR3 in ArcH11 harbors amino acid sequences that

corresponds to the identified NL ligand binding motif [14, 20]. Isothermal titration calorimetry

assays showed that ArcNb H11 binds with high affinity (1 nM K_d) and competitively displaces

stargazin, an auxiliary subunit of AMPA-type glutamate receptor, which has the highest binding

affinity (34.9 μM) of the known Arc-NL ligands [40]. The present work shows that genetically

encoded ArcNb H11 expressed in mammalian cells binds specifically to the NL and enables

co-immunoprecipitation of intracellular Arc (**Figs. 5 and S3**). ArcNb H11 also enabled efficient

immunoprecipitation of endogenous Arc synthesized during the maintenance phase of LTP in the rat

dentate gyrus *in vivo* (**Figs. 2**). Taken together, these findings make ArcNb H11 attractive as a

high-affinity binder for labeling and tracking Arc inside cells, and a candidate competitive inhibitor

of Arc-NL function *in vivo*.

ArcNbs B5, B12, C11, and D4 were shown to bind to the Arc region encompassing the CL and

C-terminal tail (**Fig. 3B).** However, specific association with the CL is likely, as recombinant

purified ArcNbs bound to the isolated CTD (Markússon *et al.* (2021). Furthermore, crystal structure

analysis of untagged ArcNb C11 in complex with the CTD demonstrated binding to the CL in a

region that encompasses a conserved retroviral capsid dimerization motif [14, 24, 40]. Thus, ArcNb

C11 is of interest as a potential regulator of CL-mediated dimerization in the context of full-length

Arc oligomerization, and eventual capsid assembly. Although CL is structurally homologous to NL

[14], there is no evidence of similar peptide ligand binding [20, 25]. The Arc C-terminal tail

(370-396) has a PEST signal region (a sequence segment that is rich in P, E, S, T residues) [47]

implicated in Arc degradation [48].

In conclusion, we developed two lines of ArcNbs: one (clones E5 and H11) for binding to NL,

another (clones B5, B12, C11, and D4) for binding the CL and C-terminal region. These new tools

open a range of possibilities, including expression of intrabodies for labelling and tracking

endogenous Arc in live-cell imaging experiments and for biochemical isolation of native Arc

complexes. Given their stable expression and high-affinity binding, it will be important in future

work to evaluate ArcNbs as genetically-encoded inhibitors of Arc NL signaling and oligomerization

in neurons. ArcNbs can be used and further developed to elucidate Arc domain-specific mechanisms and functions in synaptic plasticity. The recombinant ALFA-tagged ArcNbs may also be useful in efforts to detect oligomers and isolate intact capsids from brain tissue and other sources.

CRediT author statement

Yuta Ishizuka: Conceptualization, Methodology, Validation, Investigation, Resources, Data Curation, Writing – Original Draft, Visualization, Supervision, Project administration. **Tadiwos F. Mergiya:** Validation, Investigation, Data Curation, Writing – Review & Editing. **Rodolfo Baldinotti:** Investigation, Validation. **Ju Xu:** Investigation, Resources. **Erik I. Hallin:** Resources. **Sigurbjörn Markússon;** Resources, **Petri Kursula:** Conceptualization, Resources, Project administration. Supervision, Funding acquisition. **Clive R. Bramham:** Conceptualization, Writing – Review & Editing, Supervision, Methodology, Resources, Data Curation, Funding acquisition, Project administration.

Declarations

Conflict of interest: All the authors declare that they have no conflict of interest related to this work.

Ethical Approval: All experimental procedures were approved by the Norwegian National Research Ethics Committee in compliance with EU Directive 2010/63/EU, ARRIVE guidelines. All methods were carried out in accordance with relevant guidelines and regulations. Persons involved in the animal experiments have approved Federation of Laboratory and Animal Science Associations (FELASA) C course certificates and training.

Consent for Publication: All authors reviewed and approved the manuscript.

Data availability

The datasets generated and analyzed during the current study are available from the corresponding

authors on reasonable request.

Acknowledgements

We would like to thank The Laboratory Animal Facility of University of Bergen for maintaining

experimental animals. This work was supported by a Research Council of Norway Toppforsk grant

(249951) to CRB.

References

1. Nikolaienko O, Patil S, Eriksen MS, Bramham CR (2018) Arc protein: a flexible hub for synaptic plasticity and cognition. Semin Cell Dev Biol 77:33-42. https://doi.org/10.1016/j.semcdb.2017.09.006

2. Bramham CR, Alme MN, Bittins M, Kuipers SD, Nair RR, Pai B, Panja D, Schubert M, Soule J, Tiron A, Wibrand K (2010) The Arc of synaptic memory. Exp Brain Res 200:125-140. https://doi.org/10.1007/s00221-009-1959-2

3. Messaoudi E, Kanhema T, Soule J, Tiron A, Dagyte G, da Silva B, Bramham CR (2007) Sustained Arc/Arg3.1 synthesis controls long-term potentiation consolidation through regulation of local actin polymerization in the dentate gyrus in vivo. J Neurosci 27:10445-10455. https://doi.org/10.1523/JNEUROSCI.2883-07.2007

4. Zhang H, Bramham CR (2021) Arc/Arg3.1 function in long-term synaptic plasticity: Emerging mechanisms and unresolved issues. Eur J Neurosci 54:6696-6712. https://doi.org/10.1111/ejn.14958

5. Waung MW, Pfeiffer BE, Nosyreva ED, Ronesi JA, Huber KM (2008) Rapid translation of Arc/Arg3.1 selectively mediates mGluR-dependent LTD through persistent increases in AMPAR endocytosis rate. Neuron 59:84-97. https://doi.org/10.1016/j.neuron.2008.05.014

6. Gao X, Castro-Gomez S, Grendel J, Graf S, Susens U, Binkle L, Mensching D, Isbrandt D, Kuhl D, Ohana O (2018) Arc/Arg3.1 mediates a critical period for spatial learning and hippocampal networks. Proc Natl Acad Sci U S A 115:12531-12536. https://doi.org/10.1073/pnas.1810125115

7. Guzowski JF, Lyford GL, Stevenson GD, Houston FP, McGaugh JL, Worley PF, Barnes CA (2000) Inhibition of activity-dependent arc protein expression in the rat hippocampus impairs the maintenance of long-term potentiation and the consolidation of long-term memory. J Neurosci 20:3993-4001.

8. Jenks KR, Kim T, Pastuzyn ED, Okuno H, Taibi AV, Bito H, Bear MF, Shepherd JD (2017) Arc restores juvenile plasticity in adult mouse visual cortex. Proc Natl Acad Sci U S A 114:9182-9187. https://doi.org/10.1073/pnas.1700866114

9. Plath N, Ohana O, Dammermann B, Errington ML, Schmitz D, Gross C, Mao X, Engelsberg A, Mahlke C, Welzl H, Kobalz U, Stawrakakis A, Fernandez E, Waltereit R, Bick-Sander A, Therstappen E, Cooke SF, Blanquet V, Wurst W, Salmen B, Bosl MR, Lipp HP, Grant SG, Bliss TV, Wolfer DP, Kuhl D (2006) Arc/Arg3.1 is essential for the consolidation of synaptic plasticity and memories. Neuron 52:437-444. https://doi.org/10.1016/j.neuron.2006.08.024

10. Zhang H, Bramham CR (2020) Bidirectional Dysregulation of AMPA Receptor-Mediated Synaptic Transmission and Plasticity in Brain Disorders. Front Synaptic Neurosci 12:26. https://doi.org/10.3389/fnsyn.2020.00026

11. Chowdhury S, Shepherd JD, Okuno H, Lyford G, Petralia RS, Plath N, Kuhl D, Huganir RL, Worley PF (2006) Arc/Arg3.1 interacts with the endocytic machinery to regulate AMPA receptor trafficking. Neuron 52:445-459. https://doi.org/10.1016/j.neuron.2006.08.033

12. Okuno H, Akashi K, Ishii Y, Yagishita-Kyo N, Suzuki K, Nonaka M, Kawashima T, Fujii H, Takemoto-Kimura S, Abe M, Natsume R, Chowdhury S, Sakimura K, Worley PF, Bito H (2012) Inverse synaptic tagging of inactive synapses via dynamic interaction of Arc/Arg3.1 with CaMKIIbeta. Cell 149:886-898. https://doi.org/10.1016/j.cell.2012.02.062

13. Nair RR, Patil S, Tiron A, Kanhema T, Panja D, Schiro L, Parobczak K, Wilczynski G, Bramham CR (2017) Dynamic Arc SUMOylation and Selective Interaction with F-Actin-Binding Protein Drebrin A in LTP Consolidation In Vivo. Front Synaptic Neurosci 9:8. https://doi.org/10.3389/fnsyn.2017.00008

14. Zhang W, Wu J, Ward MD, Yang S, Chuang YA, Xiao M, Li R, Leahy DJ, Worley PF (2015) Structural basis of arc binding to synaptic proteins: implications for cognitive disease. Neuron 86:490-500. https://doi.org/10.1016/j.neuron.2015.03.030

15. Myrum C, Baumann A, Bustad HJ, Flydal MI, Mariaule V, Alvira S, Cuellar J, Haavik J, Soule J, Valpuesta JM, Marquez JA, Martinez A, Bramham CR (2015) Arc is a flexible modular protein capable of reversible self-oligomerization. Biochem J 468:145-158. https://doi.org/10.1042/BJ20141446

16. Byers CE, Barylko B, Ross JA, Southworth DR, James NG, Taylor CAt, Wang L, Collins KA, Estrada A, Waung M, Tassin TC, Huber KM, Jameson DM, Albanesi JP (2015) Enhancement of dynamin polymerization and GTPase activity by Arc/Arg3.1. Biochim Biophys Acta 1850:1310-1318. https://doi.org/10.1016/j.bbagen.2015.03.002

460   17.   Ashley J, Cordy B, Lucia D, Fradkin LG, Budnik V, Thomson T (2018) Retrovirus-like Gag Protein
461   Arc1 Binds RNA and Traffics across Synaptic Boutons. Cell 172:262-274 e211.
462   https://doi.org/10.1016/j.cell.2017.12.022
463   18.   Pastuzyn ED, Day CE, Kearns RB, Kyrke-Smith M, Taibi AV, McCormick J, Yoder N, Belnap DM,
464   Erlendsson S, Morado DR, Briggs JAG, Feschotte C, Shepherd JD (2018) The Neuronal Gene Arc
465   Encodes a Repurposed Retrotransposon Gag Protein that Mediates Intercellular RNA Transfer. Cell
466   172:275-288 e218. https://doi.org/10.1016/j.cell.2017.12.024
467   19.   Eriksen MS, Nikolaienko O, Hallin EI, Grodem S, Bustad HJ, Flydal MI, Merski I, Hosokawa T,
468   Lascu D, Akerkar S, Cuellar J, Chambers JJ, O'Connell R, Muruganandam G, Loris R, Touma C, Kanhema
469   T, Hayashi Y, Stratton MM, Valpuesta JM, Kursula P, Martinez A, Bramham CR (2021) Arc
470   self-association and formation of virus-like capsids are mediated by an N-terminal helical coil
471   motif. FEBS J 288:2930-2955. https://doi.org/10.1111/febs.15618
472   20.   Hallin EI, Eriksen MS, Baryshnikov S, Nikolaienko O, Grodem S, Hosokawa T, Hayashi Y, Bramham
473   CR, Kursula P (2018) Structure of monomeric full-length ARC sheds light on molecular flexibility,
474   protein interactions, and functional modalities. J Neurochem 147:323-343.
475   https://doi.org/10.1111/jnc.14556
476   21.   Barylko B, Wilkerson JR, Cavalier SH, Binns DD, James NG, Jameson DM, Huber KM, Albanesi JP
477   (2018) Palmitoylation and Membrane Binding of Arc/Arg3.1: A Potential Role in Synaptic
478   Depression. Biochemistry 57:520-524. https://doi.org/10.1021/acs.biochem.7b00959
479   22.   Nielsen LD, Pedersen CP, Erlendsson S, Teilum K (2019) The Capsid Domain of Arc Changes Its
480   Oligomerization Propensity through Direct Interaction with the NMDA Receptor. Structure
481   27:1071-1081 e1075. https://doi.org/10.1016/j.str.2019.04.001
482   23.   Zhao WQ, Santini F, Breese R, Ross D, Zhang XD, Stone DJ, Ferrer M, Townsend M, Wolfe AL, Seager
483   MA, Kinney GG, Shughrue PJ, Ray WJ (2010) Inhibition of calcineurin-mediated endocytosis and
484   alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors prevents amyloid
485   beta oligomer-induced synaptic disruption. J Biol Chem 285:7619-7632.
486   https://doi.org/10.1074/jbc.M109.057182
487   24.   Zhang W, Chuang YA, Na Y, Ye Z, Yang L, Lin R, Zhou J, Wu J, Qiu J, Savonenko A, Leahy DJ, Huganir
488   R, Linden DJ, Worley PF (2019) Arc Oligomerization Is Regulated by CaMKII Phosphorylation of
489   the GAG Domain: An Essential Mechanism for Plasticity and Memory Formation. Mol Cell 75:13-25
490   e15. https://doi.org/10.1016/j.molcel.2019.05.004
491   25.   Hallin EI, Bramham CR, Kursula P (2021) Structural properties and peptide ligand binding of
492   the capsid homology domains of human Arc. Biochem Biophys Rep 26:100975.
493   https://doi.org/10.1016/j.bbrep.2021.100975
494   26.   Hamers-Casterman C, Atarhouch T, Muyldermans S, Robinson G, Hamers C, Songa EB, Bendahman N,
495   Hamers R (1993) Naturally occurring antibodies devoid of light chains. Nature 363:446-448.
496   https://doi.org/10.1038/363446a0
497   27.   Steeland S, Vandenbroucke RE, Libert C (2016) Nanobodies as therapeutics: big opportunities
498   for small antibodies. Drug Discov Today 21:1076-1113.
499   https://doi.org/10.1016/j.drudis.2016.04.003
500   28.   Li T, Vandesquille M, Koukouli F, Dudeffant C, Youssef I, Lenormand P, Ganneau C, Maskos U,
501   Czech C, Grueninger F, Duyckaerts C, Dhenain M, Bay S, Delatour B, Lafaye P (2016) Camelid
502   single-domain antibodies: A versatile tool for in vivo imaging of extracellular and
503   intracellular brain targets. J Control Release 243:1-10.
504   https://doi.org/10.1016/j.jconrel.2016.09.019
505   29.   Dong JX, Lee Y, Kirmiz M, Palacio S, Dumitras C, Moreno CM, Sando R, Santana LF, Sudhof TC,
506   Gong B, Murray KD, Trimmer JS (2019) A toolbox of nanobodies developed and validated for use
507   as intrabodies and nanoscale immunolabels in mammalian brain neurons. Elife
508   8https://doi.org/10.7554/eLife.48750
509   30.   Beghein E, Gettemans J (2017) Nanobody Technology: A Versatile Toolkit for Microscopic Imaging,
510   Protein-Protein Interaction Analysis, and Protein Function Exploration. Front Immunol 8:771.
511   https://doi.org/10.3389/fimmu.2017.00771
512   31.   Gerdes C, Waal N, Offner T, Fornasiero EF, Wender N, Verbarg H, Manzini I, Trenkwalder C,
513   Mollenhauer B, Strohaker T, Zweckstetter M, Becker S, Rizzoli SO, Basmanav FB, Opazo F (2020)
514   A nanobody-based fluorescent reporter reveals human alpha-synuclein in the cell cytosol. Nat
515   Commun 11:2729. https://doi.org/10.1038/s41467-020-16575-0
516   32.   Muyldermans S (2021) A guide to: generation and design of nanobodies. FEBS J 288:2084-2102.
517   https://doi.org/10.1111/febs.15515

33.   Gotzke H, Kilisch M, Martinez-Carranza M, Sograte-Idrissi S, Rajavel A, Schlichthaerle T, Engels
N, Jungmann R, Stenmark P, Opazo F, Frey S (2019) The ALFA-tag is a highly versatile tool for
nanobody-based bioscience applications. Nat Commun 10:4403.
https://doi.org/10.1038/s41467-019-12301-7

34.   Hallin EI, Markusson S, Bottger L, Torda AE, Bramham CR, Kursula P (2021) Crystal and solution
structures reveal oligomerization of individual capsid homology domains of Drosophila Arc. PLoS
One 16:e0251459. https://doi.org/10.1371/journal.pone.0251459

35.   Panja D, Bramham CR (2017) BDNF Function in Long-Term Synaptic Plasticity in the Dentate Gyrus
In Vivo: Methods for Local Drug Delivery and Biochemical Analysis of Translation. In: Duarte
C, Tongiorgi E (eds) Brain-Derived Neurotrophic Factor (BDNF). Humana, New York, NY, pp 241-256.
https://doi.org/10.1007/7657_2017_7

36.   Soule J, Alme M, Myrum C, Schubert M, Kanhema T, Bramham CR (2012) Balancing Arc synthesis,
mRNA decay, and proteasomal degradation: maximal protein expression triggered by rapid eye
movement sleep-like bursts of muscarinic cholinergic receptor stimulation. J Biol Chem
287:22354-22366. https://doi.org/10.1074/jbc.M112.376491

37.   Drab T, Kracmerova J, Ticha I, Hanzlikova E, Ticha M, Liberda J (2011) Native polyacrylamide
electrophoresis in the presence of Ponceau Red to study oligomeric states of protein complexes.
J Sep Sci 34:1692-1695. https://doi.org/10.1002/jssc.201000869

38.   Puspitasari A, Koganezawa N, Ishizuka Y, Kojima N, Tanaka N, Nakano T, Shirao T (2016) X
Irradiation Induces Acute Cognitive Decline via Transient Synaptic Dysfunction. Radiat Res
185:423-430. https://doi.org/10.1667/RR14236.1

39.   Ishizuka Y, Bramham CR (2020) A simple DMSO-based method for cryopreservation of primary
hippocampal and cortical neurons. J Neurosci Methods 333:108578.
https://doi.org/10.1016/j.jneumeth.2019.108578

40.   Markússon S, Hallin EI, Bustad HJ, Raasakka A, Xu J, Muruganandam G, Loris R, Martinez A, Bramham
CR, Kursula P (2021) High-affinity nanobodies as tools for structural and functional studies
on mammalian Arc. bioRxiv:2021.2012.2016.472929. https://doi.org/10.1101/2021.12.16.472929

41.   Waltereit R, Dammermann B, Wulff P, Scafidi J, Staubli U, Kauselmann G, Bundman M, Kuhl D (2001)
Arg3.1/Arc mRNA induction by Ca2+ and cAMP requires protein kinase A and mitogen-activated
protein kinase/extracellular regulated kinase activation. J Neurosci 21:5484-5493.

42.   Bindels DS, Haarbosch L, van Weeren L, Postma M, Wiese KE, Mastop M, Aumonier S, Gotthard G,
Royant A, Hink MA, Gadella TW, Jr. (2017) mScarlet: a bright monomeric red fluorescent protein
for cellular imaging. Nat Methods 14:53-56. https://doi.org/10.1038/nmeth.4074

43.   McCullock TW, MacLean DM, Kammermeier PJ (2020) Comparing the performance of mScarlet-I, mRuby3,
and mCherry as FRET acceptors for mNeonGreen. PLoS One 15:e0219886.
https://doi.org/10.1371/journal.pone.0219886

44.   Gurskaya NG, Fradkov AF, Pounkova NI, Staroverov DB, Bulina ME, Yanushevich YG, Labas YA,
Lukyanov S, Lukyanov KA (2003) A colourless green fluorescent protein homologue from the
non-fluorescent hydromedusa Aequorea coerulescens and its fluorescent mutants. Biochem J
373:403-408. https://doi.org/10.1042/BJ20021966

45.   Chen Y, Muller JD, Ruan Q, Gratton E (2002) Molecular brightness characterization of EGFP in
vivo by fluorescence fluctuation spectroscopy. Biophys J 82:133-144.
https://doi.org/10.1016/S0006-3495(02)75380-0

46.   Jain RK, Joyce PB, Molinete M, Halban PA, Gorr SU (2001) Oligomerization of green fluorescent
protein in the secretory pathway of endocrine cells. Biochem J 360:645-649.
https://doi.org/10.1042/0264-6021:3600645

47.   Boldridge M, Shimabukuro J, Nakamatsu K, Won C, Jansen C, Turner H, Wang L (2020)
Characterization of the C-terminal tail of the Arc protein. PLoS One 15:e0239870.
https://doi.org/10.1371/journal.pone.0239870

48.   Rao VR, Pintchovski SA, Chin J, Peebles CL, Mitra S, Finkbeiner S (2006) AMPA receptors regulate
transcription of the plasticity-related immediate-early gene Arc. Nat Neurosci 9:887-895.
https://doi.org/10.1038/nn1708

Figure legends

Fig. 1 Schematic of ArcNb generation and modification. *A,* Schematic of ArcNb generation pipeline. *B,* Modifications of ArcNb for biochemical and immunolabeling applications. Detailed information is described in **Materials and Methods** section.

Fig. 2 ALFA-ArcNbs for immunoblotting and immunoprecipitation analysis of endogenous Arc. *A,* Rat cortical tissues were lysed in RIPA buffer. Different amounts of lysates were subject to SDS-PAGE followed by immunoblotting using ALFA-ArcNbs (B5, B12, C11, D4, E5, and H11) and conventional anti-Arc antibodies as reference. Immunoreactive bands corresponding to Arc were detected around 50 kDa (lane 1, 30 µg; lane 2, 60 µg). *B,* CCh (100 µM, 1h)-treated SH-SY5Y cells were lysed and subjected to immunoprecipitation using ALFA-ArcNb H11 and ALFA Selector™ followed by SDS-PAGE and immunoblotting using anti-Arc antibody. NC, negative control (ALFA Selector[ST] resin without ALFA-ArcNb H11. *C,* HFS was applied unilaterally to the perforant path input to the dentate gyrus of anesthetized rats to induce LTP. At 2 hours post-HFS, DG tissue was collected from the ipsilateral (Ipsi) and contralateral (Contra) DG, and immunoprecipitation analysis was performed as described in panel B

Fig. 3 Epitope mapping of ArcNbs. *A,* HEK293FT cells were transfected with mTq2-Arc-FL (residues 1-396) and truncation mutants expressing the Arc NTR (1-140), linker (135-216), or CTR (208-396). *B,* Cells were transfected with mTq2-truncation mutants expressing the Arc CTR or the isolated NL and CL+tail. Cell lysates were subjected to SDS-PAGE followed by immunoblotting using ALFA-ArcNbs and anti-Arc antibody. 1, FL (residues 1-396); 2, NTR (residues 1-140); 3, linker region (residues 135-216); 4, CTR (residues 208-396); 5, NL (residues 208-277); 6, CL+tail (residues 278-370)

Fig. 4 mSc-ArcNbs expression in HEK293FT cells as intrabody. *A,* HEK293FT cells were

transfected with mSc control vector (Mock, closed arrowhead) and mSc-ArcNb expression vectors

(opened arrowhead). Cell lysates were subjected to SDS-PAGE followed by immunoblotting using

anti-mCh antibody against mSc-tag. 1, Mock; 2, B5; 3, B12; 4, C11; 5, D4; 6, E5; 7, H11. *B,*

HEK293FT cells transfected with mSc-ArcNbs were fixed and imaged by fluorescence microscopy.

Right images are high magnification images of the are in the squares in the left images. Two cells

were randomly selected

Fig. 5 Co-immunoprecipitation of NL and FL intracellular Arc using mSc-H11 intrabody. *A,* Schema

of co-IP assay. *B-F,* co-IP assay. HEK293FT cells were co-transfected with either mTq2-Arc-FL (*B*)

or NTR (*C*) or linker region (*D*) or NL (*E*) or CL (*F*) variants and mSc-H11 constructs. Cell lysates

were then subjected to co-IP assay using anti-mCh antibody. Following SDS-PAGE,

immunoprecipitants were probed using anti-mCh, anti-Arc, and anti-GFP antibodies. Closed and

opened arrowheads indicate immunoprecipitated mSc-H11 and co-immunoprecipitated Arc variant,

respectively

Supplemental Materials

Fig. S1. Quality analysis of purified ALFA-ArcNbs. All purified ALFA-ArcNbs were subjected to

SEC followed by SDS-PAGE (*A,* B5; *B,* B12; *C,* C11; *D,* D4; *E,* E5; *F,* H11). Upper images are acryl

amid gel stained by InstantBlue® Coomassie Protein Stain (Abcam plc, Cambridge, UK) after

SDS-PAGE. Opened arrowheads indicate purified ALFA-ArcNbs around 15 kDa. Lower images are

SEC chromatograms. Closed arrowheads indicate the fractions which were collected.

Fig. S2. All ArcNbs detect purified Arc recombinant protein both in native and denatured condition.

A, Purified rat Arc mutant protein (Arc$^{s113-119A}$) was subjected to native-PAGE followed by

immunoblotting using ALFA-ArcNbs and conventional anti-Arc antibodies. BSA was used as

reference for PAGE (~66 kDa in Ponceau S staining). Std., Molecular weight marker; 1, BSA (1 μg);

2, rArc mutant protein (0.1 μg); 3, rArc mutant protein (0.5 μg). *B,* Purified Arc$^{s113-119A}$ was

subjected to SDS-PAGE followed by immunoblotting. 4, rArc mutant protein (0.05 μg); 5, rArc

mutant protein (0.1 μg); 6, rArc mutant protein (0.5 μg).

Fig. S3. AcGFP-ArcNbs expression in HEK293FT cells as intrabody. *A,* HEK293FT cells were

transfected with AcGFP control vector (Mock, closed arrowhead) or AcGFP-ArcNb expression

vectors (opened arrowhead). Cell lysates were subjected to SDS-PAGE followed by immunoblotting

using anti-GFP antibody. 1, Mock; 2, B5; 3, B12; 4, C11; 5, D4; 6, E5; 7, H11. *B,* HEK293FT cells

transfected with AcGFP-ArcNbs were fixed and imaged by fluorescence microscopy. Right images

are high magnification images of the area in the squares in the left images. Two cells were randomly

selected.

Fig. S4. Reverse co-IP assay using anti-Arc antibody. *A,* Schema of Co-IP assay. ***B-F,*** Co-IP assay.

HEK293FT cells were co-transfected with either mTq2-Arc-FL (***B***) or NTD (***C***) or Linker (***D***) or NL

(***E***) or CL (***F***) variants and mSc- H11 constructs. Cell lysates were then subjected to co-IP assay

using anti-Arc antibody. Following SDS-PAGE, immunoprecipitants were probed using anti-mCh,

Arc, and GFP antibodies. Closed and opened arrowheads indicate immunoprecipitated mTq2-Arc

variants and co-immunoprecipitated mSc-H11, respectively.

Fig. 1

A

Immunization PBMC Phage display library Bio-panning ELISA 6 clones in bacterial expression vectors

B

Nb
bact. vector
Native ArcNb

Nb ALFA-tag
add ALFA-tag seq.

bacterial expression

purification

ALFA-tagged ArcNb

Validation for immunolabeling

mammalian expression vector

mSc AcGFP
add mScarlet-I/AcGFP seq.

transfection

intracellular expression

Validation as Intrabody/ Chromobody

174 mm (double column)

Fig. 2

A

B
SH-SY5Y

C
DG tissue

174 mm (double column)

Fig. 3

174 mm (double column)

Fig. 4

174 mm (double column)

Fig. 5

174 mm (double column)