Tropical Origin, Global Diversification and Dispersal in the Pond Damselflies (Coenagrionoidea) Revealed by a New Molecular Phylogeny

B. Willink[1,2*], J. Ware[3], E. I. Svensson[4]

[1] Department of Zoology, Stockholm University, Stockholm 106-91, Sweden

[2] School of Biology, University of Costa Rica, San José 11501-2060, Costa Rica

[*] Corresponding author email: bwillink@gmail.com

[3] Division of Invertebrate Zoology, American Museum of Natural History, NY, NY, 10024, USA

[4] Department of Biology, Evolutionary Ecology Unit, Ecology Building, Lund University, Lund 223-62, Sweden

Abstract

The processes responsible for the formation of Earth's most conspicuous diversity pattern, the

latitudinal diversity gradient (LDG), remain unexplored for many clades in the Tree of Life.

Here, we present a densely-sampled and dated molecular phylogeny for the most speciose clade

of damselflies worldwide (Odonata: Coenagrionoidea), and investigate the role of time,

macroevolutionary processes and biome-shift dynamics in shaping the LDG in this ancient insect

superfamily. We used process-based biogeographic models to jointly infer ancestral ranges and

speciation times, and to characterise within-biome dispersal and biome-shift dynamics across the

cosmopolitan distribution of Coenagrionoidea. We also investigated temporal and biome-

dependent variation in diversification rates. Our results uncover a tropical origin of pond

damselflies and featherlegs ~ 105 Ma, while highligthing uncertainty of ancestral ranges within

the tropics in deep time. Even though diversification rates have declined since the origin of this

clade, global climate change and biome-shifts have slowly increased diversity in warm- and

cold-temperate areas, where lineage turnover rates have been relatively higher. This study

underscores the importance of biogeographic origin and time to diversify as important drivers of

the LDG in pond damselflies and their relatives, while diversification dynamics have instead

resulted in the formation of ephemeral species in temperate regions. Biome-shifts, although

limited by tropical niche conservatism, have been the main factor reducing the steepness of the

LDG in the last 30 Myr. With ongoing climate change and increasing northward range

expansions of many damselfly taxa, the LDG may become less pronounced. Our results support

recent calls to unify biogeographic and macroevolutionary approaches to increase our

understanding of how latitudinal diversity gradients are formed and why they vary across time

and among taxa.

Keywords: biogeography, extinction, graphical models, latitudinal diversity gradient, niche

conservatism, paleogeography, speciation

One of the most striking features of species diversity is the non-uniformity of its global

distribution. Tropical regions of the world generally have higher species richness compared to

temperate regions, a pattern known as the Latitudinal Diversity Gradient (LDG) (Hillebrand

2004). The LDG has fascinated naturalists from centuries, including both Darwin (1859) and

Wallace (1876, 1878), who saw the climatic stability of the tropics as a factor hindering

extinction, and believed that stronger biotic interactions in the tropics could accelerate

speciation. Wallace also suggested that climatic fluctuations, such as Pleistocene glaciations,

were more pronounced in temperate regions, providing a limited opportunity for diversity to

accumulate in these areas compared to the tropics (Wallace 1878). These early biogeographic

hypotheses were further developed by ecologists and evolutionary biologists throughout the

twentieth century (Dobzhansky 1950; Fischer 1960; MacArthur 1969), and today, the

mechanisms responsible for the LDG remain a highly active area of research (e.g. Hanly et al.

2017; Rabosky et al. 2018; Gaboriau et al. 2019; Saupe et al. 2019; Meseguer and Condamine

2020).

Current hypotheses to explain the causes of the LDG can be broadly categorized into

those that emphasize ecological effects on diversification rates vs. those that emphasize time for

speciation in different regions and the contingencies of evolutionary history (Pianka 1966;

Mittelbach et al. 2007; Wiens 2011). At one end of this dichotomy, the tropics are portrayed as

ecology-fuelled species pumps, due to higher speciation rates (Wiens 2007), lower extinction

rates (Weir and Schluter 2007; Pyron 2014) or both (Jablonski et al. 2006; Pyron and Wiens

2013; Rolland et al. 2014). The tropics have also been suggested to be the sources of most

evolutionary novel lineages due to their greater geographical area (Rosenzweig and others 1995;

Belmaker and Jetz 2015), greater niche heterogeneity (Moritz et al. 2000; Salisbury et al. 2012),

stronger biotic interactions (Dobzhansky 1950; Schemske et al. 2009), or higher mean

temperature (Wright et al. 2006; Brown 2014). As Darwin and Wallace already anticipated,

tropical diversity might also be buffered from the extinction that occurs in temperate regions

because of greater climatic instability (Dynesius and Jansson 2000; Jablonski et al. 2006;

Hawkins et al. 2007). In contrast, studies at the other end of the dichotomy have proposed that

the amount of time available for diversification might be the predominant cause of variation in

regional species richness, whether aligned or contrary to the LDG (Stephens and Wiens 2003;

Stevens 2006; Economo et al. 2018; Miller and Román-Palacios 2021).

71   Understanding how ecological effects and historical contingencies shape regional

diversification dynamics is therefore fundamental for a full understanding of the LDG, and it is

crucial to critically evaluate the empirical evidence for these contrasting alternatives (Wiens

2011). However, regional biotas are not only shaped by varying speciation and extinction rates,

but also by dispersal (Salisbury et al. 2012; Smith et al. 2012; Antonelli et al. 2015; Jablonski et

al. 2017). A lineage may disperse to a distant region with a similar biome, maintaining the LDG,

or move to an ecologically novel area (i.e. a colder or warmer biome), thus altering the steepness

of the LDG. Moreover, diversification and dispersal can act interdependently (Goldberg et al.

2011). For example, ecological limits to diversity, if present, could potentially reduce both

speciation rates and incoming dispersal (Etienne et al. 2019). Similarly, ecological specialization

and niche partitioning may result in fast speciation locally, but trade-off with the evolution of

dispersal ability (Jocque et al. 2010; Salisbury et al. 2012; Polato et al. 2018). Despite a potential

key role of dispersal in molding the LDG, a prevailing focus on diversification dynamics may

have curbed our understanding of the relative roles of each of these processes shaping regional

biotas (Roy and Goldberg 2007).

To date, only a few studies have examined the combined effects of speciation, extinction

and dispersal in shaping diversity gradients (Antonelli et al. 2015; Rolland et al. 2015; Spano et

al. 2016; Jablonski et al. 2017; Owens et al. 2017). The majority of these studies target clades

with a relatively rich fossil record, which also enables process-based methods for inferring time-

calibrated phylogenies, such as the fossilized birth-death process and total evidence approaches

(Ronquist et al. 2012; Heath et al. 2014). However, most organisms, including insects, do not

fossilize well (Wills 2001; Kidwell and Holland 2002), and therefore other process-based

methods are necessary to calibrate molecular clocks and infer ancestral geographic ranges (e.g.

Tiley et al. 2020). Biogeographic dating approaches have been long used as an alternative to

fossil dating, but until recently, most of these methods relied on node calibrations

(Kodandaramaiah 2011; De Baets et al. 2016) and have consequently failed to consider the joint

probability of alternative diversification and dispersal scenarios (Ho et al. 2015).

Landis (2017) introduced a data-dependent generative model, in which speciation times

and ancestral geographic states are jointly estimated over discretely defined epochs. Crucially,

this approach used paleogeographical data to explicitly consider how the Earth's dynamic past,

through tectonic movements, has influenced the availability of dispersal routes for terrestrial

animals. More recently, Landis et al. (2021) implemented a similar time-stratified approach to

model within-biome dispersal and biome shifts, building on paleoecological literature and thus

examining the role of phylogenetic niche conservatism in shaping biogeographic histories. These

studies illustrate how empirical models of dispersal barriers can be used in lieu of fossils to

implement process-based calibrations. They also show how time-dependent process-based

models can be used to reconstruct ancestral diversity patterns, such as the LDG, while

accounting for paleoclimatic changes. Here, we capitalised on these developments to investigate

the phylogenetic and biogeographic history of a globally-distributed group of insects with a scant

insect fossil, the damselfly superfamily Coenagrionoidea (Carle et al. 2008; Bybee et al. 2021).

Our target clade includes two families: the pond damselflies (family Coenagrionidae) and

their close relatives the featherlegs (family Platycnemididae). Coenagrionidae is a large family,

distributed across all continents except Antarctica, and with species that are at extremes of trait

and niche diversity. For instance, the family includes the tropical-rainforest phytotelmata-

breeding helicopter damselflies, previously considered a separate family due to their

morphological distinctiveness (Dijkstra et al. 2014; Toussaint et al. 2019). At the other extreme,

Coenagrionidae also includes species are restricted to Subartic fens and marshes in the type

genus *Coenagrion* (Paulson 2009, 2011). In the Nearctic alone, where most systematic niche data

has been assembled, heterogeneity in habitat preference is striking. Species vary vastly in their

affinities to forested areas, range sizes and breeding habitats (Abbott et al. 2022).

Platycnemididae, the other family in Coenagrionoidea, is composed by primarily lotic

(i.e. inhabiting fast-moving water bodies) and paleotropical taxa, with some species reaching

Northern Europe and East Asia (Boudot and Kalkman 2015; Dijkstra 2017).

Combined, the two families in Coeangrionoidea account for over a fourth of all Odonata

species. Their ancient history (Kohli et al. 2021; Suvorov et al. 2021), rich ecological diversity

(Abbott et al. 2022) and high dispersal potential as relatively large volant insects (May 2019),

makes pond damselflies and featherlegs an excellent group to study the processes responsible for

global diversity patterns, such as the LDG. However, few molecular studies have examined the

phylogenetic relationships within Coenagrionoidea (Carle et al. 2008; Dijkstra et al. 2014;

Waller and Svensson 2017; Toussaint et al. 2019; Bybee et al. 2021; Kohli et al. 2021), and a

single dated and species-level phylogeny currently includes this clade (Waller and Svensson

2017). This species-level phylogeny nonetheless used phylogenetically-unsupported taxonomy

as backbone. There is clearly a need for an updated phylogeny of Coenagrionoidea with denser

taxon sampling, and for modern phylogenetic comparative analyses to uncover large-scale

diversity patterns and their drivers in this ancient group of insects.

Here, we present the most extensively sampled phylogeny of pond damselflies and their

relatives to date. We include over 35% of the approximately 1,805 species, and thereby

substantially increase the phylogenetic coverage of sampled taxa compared to previous studies

(approximately 7% of all species in Dijkstra et al. 2014; and 23% in Waller and Svensson 2017).

Recent phylogenomic studies have produced a robust and dated backbone for the order Odonata,

which we used to inform the origin time, over 100 Ma, of the most recent common ancestor of

pond damselflies and featherlegs (Kohli et al. 2021; Suvorov et al. 2021). We combine this root-

age prior with an empirical paleogeographic model (Landis 2017), genetic data, and distribution

range data of extant taxa, to simultaneously infer the divergence times and geographic

distributions of ancestral lineages. We also use this time-calibrated phylogeny to quantify

temporal and geographic patterns of diversification, and to jointly model how within-biome

dispersal and biome shifts have shaped regional diversity throughout the history of the clade. Our

results provide a comprehensive overview of the macroevolutionary history of this insect group

and shed new light on the question of how speciation, extinction and lineage movements

influence global diversity patterns.

MATERIALS AND METHODS

Here, we summarize the main features of the phylogenetic analyses performed in this study. For

a detailed description of data collection, sequence alignment and the statistical approach used for

biogeographic dating, diversification, and biome-shift analyses, see the Supplementary Material.

We mention here the most relevant prior distributions. For complete information on priors and

parameter proposals, see the Extended Methods and accompanying scripts. All novel sequence

data used in this study will be uploaded to NCBI Genbank. The sequence alignment, distribution

data and code necessary to reproduce our results will be uploaded on Dryad

(https://datadryad.org/stash) and are currently available on Github

(https://github.com/bwillink/Damsel_Phylo). Phylogenetic analyses were partly conducted on the

computer clusters at the LUNARC Center for Scientific and Technical Computing, Lund

University, Sweden, and the Uppsala Multidisciplinary Center for Advanced Computational

Science (UPPMAX).

Sequence Data and Tree Topology Inference

We sampled a total of 669 taxa in the Coenagrionoidea superfamily, 556 of which are currently

classified as Coenagrionidae and 113 as Platycnemididae (Table S1, S2). In a recent

phylogenomic study, the small family Isostictidae was recovered as sister to the clade including

both Coeangrionidae and Platycnemididae (Bybee et al. 2021). As we did not sample

Isostictidae, we hereafter use the superfamily name Coenagrionoidea to refer to Coenagrionidae

(pond damselflies) and Platycnemididae (featherlegs) only. Samples were collected in the field

(n = 142 species), or obtained from specimens in museum and private collections (n = 362

species), and publicly available data on NCBI (https://www.ncbi.nlm.nih.gov/genbank/) (n = 194

species). For each species, we obtained sequence data for up to five loci (cytochrome oxidase

subunit I, 16S ribosomal DNA, histone 3, phenyl-methyltransferase and 28S ribosomal DNA).

Our molecular data consists of 1,695 new sequences, 564 previously published sequences, and

105 unpublished sequences from the Naturalis Biodiversity Center, The Netherlands. We

conducted all phylogenetic analyses using probabilistic graphical models and Bayesian inference

for parameter estimation, using RevBayes v. 1.0.7, v. 1.0.12, and v. 1.1.1 (Höhna et al. 2014,

2016).

Our first analysis aimed to infer the topology of the Coenagrionoidea phylogeny. We

used a rooted uniform tree prior and a partitioned data scheme accounting for substitution rate

variation among loci and among codon positions of coding sequences (Table S3). We specified a

General Time-Reversible (GTR) model of molecular evolution for each partition, and accounted

for rate heterogeneity among sites by assuming gamma-distributed rates. Finally, we enforced

the monophyly of each of the two families within Coenagrionoidea (Coenagrionidae and

Platycnemididae), as supported by recent large-scale studies (Kohli et al. 2021; Bybee et al.

2021; Suvorov et al. 2021). To do this, we imposed a single topological constraint of the root

node of the tree, which was necessary as preliminary analyses did not adequately resolve this

well-established ancestral relationship. The maximum *a posteriori* probability (MAP) tree was

used to summarize phylogenetic inference.

Biogeographic Dating with Empirical Paleogeography

Next, we simultaneously inferred speciation times and ancestral distributions across the

Coenagrionoidea tree using the empirical paleogeographic model and statistical approach

developed by Landis (2017). Instead of relying on internal node calibrations, this approach

jointly models ancestral dispersal and speciation events as conditional on historical shifts in

oceanic barriers and the distribution ranges of extant taxa. To use biogeography in node-age

estimation, we recorded species distributions from the literature, museum collections, IUCN Red

List assessments and reputable web sources for all extant taxa included in this study (Table S4).

Due to the low spatial resolution and coarse-grained nature of available data, species

distributions were registered as presence or absence from administrative geographical regions

(countries and states/provinces/territories for countries spanning multiple biogeographic areas;

see below).

In the model developed by Landis (2017), dispersal can occur by three modes (short-

medium- and long-distance), which are differentially limited by oceanic barriers. The empirical

paleogeographic model captures historical changes in these water barriers due to continental drift

(Landis 2017). Thus, paleogeography determines the probability of transitions between

biogeographic states, for each dispersal mode and for each discrete time interval (epoch) since

the maximum plausible root age of the tree. This time-heterogeneous process describing the

probability of dispersal events in turn informs branch length estimation in absolute time (Landis

2017). For example, ancestral dispersal leading to a disjunct distribution in the present is inferred

to occur with higher probability during time intervals (epochs) in which the currently disjunct

land masses were adjacent, and hence connected by all three dispersal modes, than when they

were separated by small or large water barriers, which would have then required medium- and

long-distance dispersal respectively.

Damselflies are aquatic insects with a terrestrial adult phase. They require fresh water

habitats to breed and for larvae to develop (Corbet 1999). As damselflies cannot survive in or on

salt water, it is highly likely that their dispersal has been historically constrained by oceanic

barriers. Yet, Coenagrionoidea has a cosmopolitan distribution at present, ranging across all

continents except for Antarctica and Greenland (Table S4, Fig. 1). Unlike the larger dragonflies

(suborder Anisoptera), in which long-distance and sometimes trans-oceanic migration has been

described (Wikelski et al. 2006; Anderson 2009; Troast et al. 2016), the majority of damselflies

are small, weak fliers and tend to be more sedentary. For these biological reasons, we considered

that a global biogeographic model of changes in dispersal routes due to continental drift would

be a suitable approach to explore historical biogeography in these damselflies.

The empirical paleogeographic model in Landis (2017) consists of 25 biogeographic

areas (Fig. 1, Table S5) and 21 epochs between 240 Ma and the present (see Table S2 and Fig.

S8 in Landis 2017). We combined this model with a root-age prior based on recent large-scale

phylogenomic studies of major families in the whole order Odonata (Kohli et al. 2021; Suvorov

et al. 2021). Suvorov et al. (2021) inferred a backbone phylogeny for Odonata using a

supermatrix of 1,603 gene orthologs and fossil calibrations with 20 crown fossil constraints.

Kohli et al. (2021) used the sequences from nearly 3,000 protein-coding genes and 17 fossil

calibrations to reconstruct and date the phylogeny of Odonata. These studies date the origin of

Coenagrionoidea to ~107 and ~116 Ma respectively. We placed a normally distributed root-age

prior with a mean of 120 and standard deviation of 20 to reflect this prior knowledge on the

origin time of Coenagrionoidea. The normal distribution was truncated between 240 Ma, the

approximate age of the most recent common ancestor (MRCA) of all Odonata (Misof et al. 2014;

Suvorov et al. 2021) and 40 Ma, the minimum age according to the study with the youngest age

estimate for Coenagrionoidea (Waller and Svensson 2017).

To assess the impact of our strongly informed root-age prior and the empirical

paelogeographic model, we conducted two additional analyses to contrast our results. First, we

ran an identical model but with a weakly informed root-age prior, in which the MRCA of

Coenagrionoidea occurred with equal probability between 240 and 40 Ma. Second, we conducted

an analysis with a uniform prior on the root age, in this case bounded between 240 and 0 Ma, and

ignoring the empirical paleogeographic data. This latter analysis hence excluded all empirical

data (except the inferred origin time of Odonata) that could inform absolute divergence times and

time-heterogeneous dispersal rates between biogeographic areas, and was used to rule out the

possibility that divergence times were influenced by artefacts arising from model construction.

For a comparison of this paleogeographic dating approach to a dating analysis informed by fossil

constraints and a pure-birth model see the Supplementary Material.

FIGURE 1 HERE

Figure 1. Inferring the biogeographic history of pond damselflies and their relatives the

featherlegs (superfamily Coenagrionoidea). Speciation times and ancestral ranges were jointly

estimated using a time-heterogeneous continuous-time Markov model, based on empirical

paleogeography, an informed root-age prior and molecular and distribution data of 669 extant

taxa. The empirical paleogeographic model in Landis (2017) was used to determine dispersal

edges between 25 geographic areas across the globe (Greenland, Antarctica (E) and Antarctica

(W) not shown in the colour legend). Phylogenetic inference was summarized using a maximum

a posteriori (MAP) tree. The distribution areas occupied by each of the extant taxa are shown at

the tips of the tree. Pies show the proportion of each ancestral state in a random sample of 1000

posterior trees. The cladogram in the lower right shows the inferred ancestral ranges for the most

recent common ancestor of Coenagrionoidea, and for each of the three major clades in the

superfamily. The fraction of taxa sampled is shown in parenthesis. One representative species is

illustrated for each clade: *Acanthagrion adustum* ('core'), *Leptagrion elongatum* (ridge-face'),

Platycnemis pennipes (featherlegs). Photos: EIS.

Diversification through Time and Space

We quantified the temporal and geographic dynamics of diversification in Coenagrionoidea

using the MAP tree from the dating analyses above. We used an episodic birth-death (EBD)

model (Stadler 2011; Höhna 2015) to test for time-dependent diversification rate shifts since the

MRCA of Coeangrionoidea. EBD analyses assume that diversification rates change through time

in a piece-wise constant fashion, following a process of Brownian motion. Thus for each epoch,

rates are centred around the estimate in the previous epoch going backwards in time (Höhna

2015; Magee et al. 2020). We used an EBD model with 20 equal-length time intervals, and

accounted for missing taxa in each of the main clades of Coenagrionoidea (Fig. S4).

We used a Hidden State-Dependent Speciation and Extinction (HiSSE) model to estimate

the diversification rates in three latitude-delimited biomes (tropical, warm-temperate and cold-

temperate) while accommodating background diversification-rate heterogeneity across different

branches of the tree (Beaulieu and O'Meara 2016). Heterogeneity in background diversification

was taken into account by including a hidden trait with two states, which differed in their effects

on diversification. We first compared the relative fit of this biome-dependent HiSSE model

against a biome-independent model, which also accounted for diversification rate heterogeneity

using a hidden trait with two states. We then conducted three biome-dependent diversification

analyses under different data assumptions.

Of the 669 taxa in our phylogeny 31 had an unknown biome state, either because their

species identity was uncertain, or because their distribution range was not covered by the

paleobiome model used in the biome shift and within-biome dispersal analysis (see below). An

additional 99 species occupy ranges that extend across tropical and warm-temperate biomes (n =

46), across warm-temperate and cold-temperate biomes (n = 47) or across all three biome types

(n = 6). We treated these taxa in two ways in separated analyses. In the main text, we present the

results of our HiSSE model assigning taxa the coldest biome they inhabit. Because the common

ancestor of Coenagrionoidea was tropical (see Results), biome states here indicate the extent to

which a lineage has become established in a novel temperature niche. In the Supplementary

Material, we show these results are qualitatively unchanged if we treat the widely distributed

taxa as ambiguous between the biomes they currently occupy. Finally, we also conducted the

biome-dependent HiSSE analysis using the Coenagrionoidea tree from the weakly informed

dating analysis as input, and show these results are also qualitatively similar.

For all HiSSE analyses, biomes were assigned to each taxa based on present-day

geography, using distribution data at the level of administrative areas (Table S4), and biome

reconstructions for the last 5 Ma (Jolly et al. 1998; Salzmann et al. 2008; Otto-Bliesner et al.

2020). We note that the HiSSE model is time-constant and does not consider how paleoclimatic

change influence the geographic distribution of tropical and temperate biomes.

Biome shifts and within-biome dispersal

To investigate the role of lineage movements on the formation and dynamics of the LDG, we

jointly modelled dispersal events within biomes and biome shifts in Coenagrionoidea using the

approach developed in Landis et al. (2021). Similarly to our dating analysis, the likelihood of a

dispersal event or biome shift is informed by the availability of suitable routes, but here, routes

are potentially shaped by both land and biome continuity. For instance, if dispersal is limited by

biome affinities, a lineage adapted to tropical rainforests in South East Asia should have a better

chance of reaching Europe during periods of the Earth's climatic history in which tropical

rainforests were widespread in both regions, compared to more recent icehouse periods, when

tropical rainforests had withdrawn from Europe. In contrast, if dispersal is unconstrained by

ecological features, the same lineage movement would only depend on geographic continuity,

which has been largely maintained between Europe and South East Asia throughout the last 100

Myr. Crucially, because the extent to which these geographic and ecological structures influence

ancestral range shifts is unknown, Landis et al. (2021) treated them as free parameters that are

estimated from the data as part of the biome shift and dispersal process (see Extended Methods).

As in our dating analysis, the dynamics of biome availability and connectivity are

represented in this model by time-dependent graphs, informed by literature data. Here, we build

on the paleobiome graphs used in Landis et al. (2021), which include three broad biome

categories (tropical, warm-temperate, and cold-temperate) across six geographic regions

(Southeast Asia, East Asia, Europe, South America, Central America, and North America), and

eight time intervals over the last 106 Myr. We focus on the last 106 Myr, based on the inferred

origin time of Coenagrionoidea in our dating analysis (see Results). We nonetheless expand the

empirical model of Landis et al. (2021) to accommodate more of the global distribution of

Coenagrionoidea, by adding three geographic regions (Africa, Australia, and India) that harbour

a large fraction of pond damselfly and featherleg diversity (Fig. S2-S3). Species with

distributions outside these main geographic regions (e.g. Pacific islands and Central Asia

endemics, n = 32) were treated as missing data (i.e. included in the tree but lacking a biome-

region state). Similarly, extant species with ranges that span two or more geographic regions or

biomes (n = 151), were treated as ambiguous between their current states (e.g. a tropical

rainforest species ranging from Central America to South America was coded as ambiguous

between the Tropical+SAm and Tropica+CAm states, see also Table S7).

We used stochastic character mapping to sample lineage proportions through time and

evolutionary histories in the dispersal and biome shift model. Following Landis et al. (2021), we

also used these sampled histories to quantify the frequency of events (biome shifts and region

shifts) and event series (see below) throughout the evolution of Coenagrionoidea. *Event series*

represent different patterns of consecutive transitions between biomes and regions. First, we

quantified instances in which both biomes and regions shifted in a lineage and recorded whether

the biome shift preceded the dispersal event (*biome-first* series) or *vice versa* (*region-first* series).

Then, we examined consecutive changes in biomes and regions separately. A *reversal*

corresponds to a series of events with the same starting and ending state. For instance, a *biome*

reversal occurs when a tropical species shifts to a warm-temperate biome and then back to the

tropics. A *flight* is a series of two events across three different states. For example, a lineage that

originates in Africa and then disperses to India and South East Asia would represent a *region*

flight. We compute and report the proportion of each event and *event series* type across our

posterior sample of evolutionary histories (see Extended Methods).

In this and previous analyses, we report the posterior mean (PM) and 95% highest

posterior density (HPD) intervals for parameter estimates. For proportions of events and *event*

series we also report 80% HPD intervals. Phylogenetic inferences and parameter estimates were

summarized using R (R Core Team 2021). Plots we produced with the R packages *ggplot2*

(Wickham 2016) and *ggtree* (Yu et al. 2017).

RESULTS

Topology inference

The relationships reconstructed here are largely congruent with previous works (Dijkstra et al.

2014; Toussaint et al. 2019; Bybee et al. 2021). We fixed the relationship between the two

families of Coenagrionoidea, Platycnemididae (featherlegs) and Coenagrionidae (pond

damselflies) as resolved in previous genomic-scale backbone phylogenies (Suvorov et al. 2021;

Bybee et al. 2021; Kohli et al. 2021). Within Platycnemididae, our tree recovers the previously

well-supported relationships between the subfamilies Platycnemidinae, Disparoneurinae and

Calicnemiinae, while the phylogenetic placement of Allocnemidinae, Idiocnemidinae and

Onychargiinae (*sensu* Dijkstra et al. 2014) remains elusive across studies (Fig. S5a).

Nonetheless, the generic composition of each of these subfamilies (Fig. S6) is consistent with

Dijkstra et al. (2014), who sampled a relatively large fraction (~10%) of all fetherleg species.

Within Coenagrionidae, the relationships between four main clades, the 'core' pond

damselflies (*sensu* Dijkstra et al. 2014), the genus *Argia* (Rambur, 1842), the subfamily

Protoneurinae (*sensu* Dijkstra et al. 2014) and the strict-sense 'ridge-face' clade (Dijkstra et al.

2014), have remained contentious among recent studies (Fig S5b). Our results are congruent with

the most recent backbone phylogeny for Odonata, which used targeted genomics to resolve

ancestral relationships in the order (Bybee et al. 2021) (Fig S5b). In both phylogenies, *Argia* and

the strict-sense 'ridge-face' pond damselflies form a clade, in turn sister to Protoneurinae (Fig

S5b). We hereafter refer to this more inclusive clade as 'ridge-face' pond damselflies to

distinguish it from the the 'core' clade identified in Dijkstra et al. (2014).

Three large and well-sampled genera of 'core' pond damselflies (*Ischnura*, *Enallagma*

and *Acanthagrion*) form a clade that is sister to *Agriocnemis* (incl. *Mortonagrion* and

Argiocnemis, see Extended Results). We recovered these previously supported relationships in

our phylogenetic inference (Fig. S5c). We also recovered a close relationship between

Coenagrion and *Pseudagrion* (and allied genera) relatively to the clade described above and

consistently with previous studies (Fig. S5c). However, our results differ from previous

phylogenies in that it places other relatively small South Pacific genera (*Xanthocnemis*,

Austrocoenagrion, and *Stenagrion*) within the same clade as *Coenagrion* and *Pseudagrion*, albeit

with low support (Fig. S5c).

Finally, in the strict-sense 'ridge-face' pond damselflies (i.e. excluding *Argia* and

Protoneurinae), relationships between the most distinct genera are also largely congruent among

studies. The Neotropical *Telebasis* and the Paleotropical *Ceriagrion* form a clade separately from

the strict rainforest dwellers *Mecistogaster*, *Metaleptobasis* and *Teinobasis* (Fig. S5d). In both

our study and in Toussaint et al. (2019), who focused on ridge-face pond damselflies, the

Neotropical genus *Metaleptobasis* and the Neotropical helicopter damselflies (including

Mecistogaster) show closer affinity, while in Dijkstra et al. (2014), *Mecistogaster* and the

Paleotropical *Teinobasis* appear more closely related (Fig. S5d).

Out of 115 genera included in this study, 45 are represented by a single species, either

because the genus is monotypic (n = 22, i.e. having only one described species) or because

material was available for only one species (n = 23). Of the remaining 70 genera, 50 were

recovered as monophyletic, in most cases with high posterior probability (Table S6, median =

1.00, min = 0.55, max = 1.00). A total of 20 genera were thus recovered as paraphyletic. Of

these, five pose phylogenetic relationships that challenge current taxonomy supported by

morphology (*Acanthagrion*, *Cyanallagma*, and *Oxyagrion*) or earlier molecular studies

(*Erythromma*, *Platycnemis*), 10 are supported by recent studies that also inferred paraphyly

(*Agriocnemis*, *Coeliccia*, *Elattoneura*, *Indocnemis*, *Ischnura*, *Mecistogaster*, *Nesobasis*,

Pseudagrion, *Prodasineura*, and *Teinobasis*), and five lack previous studies with taxonomic

sampling that would enable detection of paraphyly (*Aciagrion*, *Enallagma*, *Mortonagrion*,

Psaironeura, *Telebasis*). We discuss these potentially paraphyletic genera in the Extended

Results.

Dated Phylogeny and Ancestral Biogeography

Our biogeographic dating analysis using a strongly informed root calibration returned a mean

age estimate for the MRCA of Coenagrionidae and Platycnemididae of 105 Ma (95% HPD

interval = 64 – 145 Ma; Fig. S26). In contrast, applying a broad uniform prior on the root age

resulted in a much younger estimate of 67 Ma (95% HPD interval = 40 – 118 Ma; Fig. S26). An

analysis excluding empirical paleogeographic data produced, as expected, a flat posterior root

age distribution between 240 and 0 Ma (Fig. S26). Finally, an analysis excluding

paleogeographic data and relying on fossil constraints for node-dating returned a similar root-age

estimate as the strongly informed biogeographic dating analysis (Table S10). Relatively shallow

nodes (i.e. at the level of genera) had older age estimates in the strongly informed biogeographic

dating analysis than in previous studies of comparable clades (Table S10). In contrast, our fossil-

based approach generally resulted in young age estimates for shallow nodes compared to other

large-scale studies, and in agreement with genus-level studies (Table S10). Despite large dating

differences between the two biogeographic analyses in this study, biogeographic trends were

qualitatively similar between strongly informed and weakly informed models (Fig. S27-S33).

Here, we focus on results under the strongly informed prior, and present the full results under the

weakly informed prior in the Supplementary Material.

Extant pond damselflies and featherlegs are globally distributed (Fig. 1). Our ancestral

biogeographic inference indicated that Coenagrionoidea most likely originated in either Northern

South America or Western Africa (Fig. 1, S27), as these two ancestral ranges combined

accounted for 88% of the posterior samples of the root state. The two extant lineages descending

from this common ancestor had contrasting biogeographic histories. The featherlegs (family

Platycnemididae) originated in Western Africa (PS = 0.959, Fig. 1, S27) and likely dispersed

throughout the Old World after multiple departures from Africa (Fig. 1, S28-S29). The pond

damselflies (family Coenagrionidae) are in turn composed of two clades with distinct dispersal

dynamics. The 'ridge-face' clade of pond damselflies originated with high probability in

Northern South America (PS = 0.999; Fig. 1, S27), and thereafter remained largely neotropical,

with a few successful dispersal events to the Nearctic (in *Argia* and *Nehalennia*), tropical Africa

and Asia (in *Ceriagrion*) and presumably also across the Pacific (*Melanesobasis*, *Teinobasis*,

Amphicnemis and allied genera) (Fig. 1, S30-S31). Finally, the 'core' pond damselflies have a

much more dynamic dispersal history, particularly in the genus *Ischnura* (Fig. 1, S32-S33), and

currently occupy all available biogeographic areas and many oceanic islands not considered in

this analysis (e.g. *Megalagrion* in Hawaii; Fig. S32-S33). While the origin of 'core' pond

damselflies is also tropical, the geographic range of their MRCA remained elusive (Fig. 1, S27).

Temporal Dynamics of Diversification

Our EBD analysis showed a decline in the rate of speciation since the origin of Coenagrionoidea,

particularly between 85 - 55 Ma (Fig. 2a). This analysis also detected a recent but modest

increase in extinction (Fig. 2b). The combined effects of speciation and extinction dynamics

resulted in a relatively constant net diversification rate in the early history of Coenagrionoidea,

followed by a period of declining diversification (~ 85 - 55 Ma) and another period of constant to

slowly declining diversification (~55 Ma to the present) (Fig. 2c). As a result of increasing

extinction over the last 5 Myr, the 95% HPD interval of diversification over this period includes

the possibility of net diversity loss near the present (Fig. 2c). EBD dynamics modelled on the

dated phylogeny under a weakly informed root-age prior are qualitatively similar to the patterns

described above (Fig. S34).

Willink et al.

FIGURE 2 HERE

Figure 2. Diversification rate changes since the MRCA of Coenagrionoidea: a) speciation, b)

extinction and c) net diversification. Diversification rates were estimated on the maximum *a*

posteriori (MAP) tree from the biogeographic phylogenetic dating analysis using a root age

calibration based of previous phylogenomic studies (Kohli et al. 2021; Suvorov et al. 2021).

Changes in diversification rates were modelled under an episodic birth-death (EBD) process,

over 20 equal-length time intervals and assuming autocorrelation among consecutive time

intervals (see Extended Methods).

Biome-dependent Diversification

The biome-dependent HiSSE model was very strongly supported over an alternative model with

only biome-independent rate heterogeneity (Table S11). As in the paleogeographic dating

analysis, the HiSSE model inferred a tropical distribution with high certainty for the MRCA of

pond damselflies and featherlegs (Fig. S35). Following dispersal from the tropics, both

speciation and extinction were accelerated in cold-temperate biomes, while warm-temperate

biomes prompted relatively low speciation and high extinction (Table S12; Fig. 3a,b). As a

result, net diversification rates were similar across biomes (Table S12; Fig. 3a,b). In contrast,

lineage turnover occurred at a faster pace in both temperate biomes compared to the tropics

(Table S12; Fig. 3d).

Background diversification rates distinguished the 'core' pond damselflies and the 'ridge-

face' genera *Argia* and *Ceriagrion*, all with relatively high diversification, from the remaining

'ridge-face' pond damselflies and the featherlegs, with with relatively low diversification (Fig.

S35). Results of the HiSSE analysis using the MAP tree under a weakly informed root-age prior

and using a character matrix that included ambiguous states for wide ranging taxa were

qualitatively similar (Table S13-S14; Fig. S36-S39). A model run without data recovered prior

distributions, as expected (Fig. S40).

FIGURE 3 HERE

Figure 3. Biome effects on diversification of pond damselflies and their relatives the featherlegs

(superfamily Coenagrionoidea). (a) Speciation and (b) extinction rates in tropical, warm-

temperate, and cold-temperate biomes were estimated using a Hidden-State Dependent

Speciation and Extinction (HiSSE) model that accounts for background heterogeneity in

diversification rates across the phylogeny, by including a hidden trait with two states. (c)

Diversification was calculated as the net difference between speciation and extinction and (d)

turnover was calculated as the ratio between extinction and speciation. The histograms show the

posterior distribution of parameter estimates.

Dispersal and biome shift dynamics

Estimated weighing parameters for geographical (w_G= 0.003) and biome features (w_B= 0.975),

indicated that lineage movements in Coenagrionoidea are strongly influenced by biome

affinities, while land connectivity independently of ecological features has a marginal effect on

shaping ancestral ranges. It is thus not surprising that ancestral state reconstructions differed

somewhat from those in our biogeographic dating model, which considered paleogeographic but

not paleoecological history. In both models, we obtained strong support for a tropical ancestor of

pond damselflies and featherlegs (Fig. 1, 4, S41). However, unlike the biogeographic dating

model, the ancestral region with highest posterior probability in the biome-shift model was South

East Asia (Fig. 4, S41). Featherlegs and 'core' pond damselflies were also inferred as most likely

originating in tropical South East Asia, in contrast to the biogeographic dating analysis, while

'ridge-face' pond damselflies originated with highest posterior probability in tropical South

America, consistent with the biogeographic dating analysis (Fig. S41).

Clade-wide biome-region dynamics over time point to three major themes in the ancestral

distribution of Coenagrionoidea. First, lineage diversity has been overwhelmingly tropical

throughout history, with a rapid increase of taxa in South America shortly after the origin of the

clade (Fig. 4). Second, a cooling trend during the Oligocene coincided with a slow and steady

increase in temperate diversity, which continues to the present and is most notable in warm-

temperate North America (Fig. 4). Finally, inference of relative biome-shift rates suggested a

dynamic diversity exchange between tropical and warm-temperate biomes, indicated by a

moderate rate of dispersal from tropical to warm-temperate biomes and fast dispersal back to the

tropics (Fig. S42). In contrast, cold-temperate biomes gain diversity and lose diversity at

comparable and slow rates (Fig. S42).

FIGURE 4 HERE

Figure 4. Proportion of Coenagrionoidea lineages in each biome-region state and through time.

Lineage proportions were computed from stochastic histories sampled in a time-heterogeneous

model of biome-region shifts (Landis et al. 2021). The dark-grey to light-grey bars on the right

show the proportions of the three biome states in extant taxa sampled in this study (669 out of

~1,800 species).

The dynamic diversity exchange between tropical and warm-temperate biomes also

emerged within regions, but the timing of these events seems to have varied across the planet. In

South America, increased shifts between tropical and warm-temperate biomes followed the

Oligocene cooling trend (Fig. 5a), as warm-temperate biomes became more available in the

region (Fig. S2-S3). In North America and Northern Eurasia (Fig. 5a,b), the exchange occurred

mainly before the end of the Eocene, after which tropical biomes began to decline in these

regions (Fig. S2-S3). Finally, in the rest of Afroeurasia (Africa, India and South East Asia) and

in Australia, shifts between tropical and warm-temperate biomes were comparatively stable over

time (Fig. 5c).

We note that stochastically mapped character histories were largely congruent with

regionally available biomes (Fig. S43, see also Extended Methods). The largest proportion of

lineages with mismatched states (i.e. lineages with biome states that are regionally unavailable)

occurred at the start of the Miocene, when nearly 6% of Coenagrionoidea lineages were inferred

to occupy tropical biomes in North America (Fig. 4), but tropical biomes in this region became

marginal during the transition between the Oligocene and Miocene, in the empirical paleobiome

model (Fig. S2-S3). As the proportion of tropical lineages in North America decreased during the

Miocene (Fig. 4), congruence between inferred and available states increased (Fig. S43).

Consistent with estimates of biome-shift rates (Fig. S42) and relatively frequent shifts

between tropical and warm-temperate biomes within regions (Fig. 5a-c), we found that *biome*

reversals (e.g. from tropical to warm-temperate and back to tropical) are by large the most

frequent *event series* across Coenagrionoidea (Fig. 5d). In contrast, *region reversals* and *biome*

flights are extremely rare, while *region flights* and series involving both biome shifts and

dispersal events occur at intermediate frequencies (Fig. 5d). Interestingly, the proportion of

region flights dropped to almost zero during the Oligocene (Fig. 5d), when our empirical

paleobiome model encodes a sharp decline in the continuity of tropical biomes across the Earth

(Fig S2-S3). Together, *event series* reconstructions further indicate that movements within

relatively warm areas of the globe (i.e. tropical and warm-temperate biomes) are the most

prominent features in the historical biogeography of Coenagrionoidea.

FIGURE 5 HERE

Figure 5. Proportion of biome-shift events (a-c) and event series (d) through time. Event

proportions were estimated from stochastic histories sampled in a time-heterogeneous model of

biome-region shifts. Circles represent posterior means, bars represent 80% HPD intervals, and

lines represent 95% HPD intervals. See Methods for descriptions of the event series types.

DISCUSSION

In this study, we used a combination of biogeographic and multi-locus sequence data from extant

taxa, empirical paleogeographic and paleobiome models (Landis 2017; Landis et al. 2021), and

dating priors from recent phylogenomic analyses (Kohli et al. 2021; Suvorov et al. 2021). Our

goal was to understand how biome-dependent diversification and dispersal shaped the current

latitudinal diversity gradient in the most speciose superfamily of damselflies. While inferences

about the age of Coenagrionoidea are notably sensitive to prior information (Fig. S26), our

ancestral biogeographic reconstructions based on empirical paleogeography are robust to node-

age uncertainty (Fig. S27-S33). Biogeographic inferences that additionally consider ecological

niche conservatism, by modelling the effects of biome continuity on lineage movements, were

partly in conflict with the paleogeographical-only model, especially in deeper nodes of the tree.

Nonetheless, all of our analyses concurred on a tropical origin of Coenagrionoidea, followed by

rapid dispersal throughout the world's tropics, and more recently into high latitudes.

Despite their early global presence, diversification rates in Coenagrionoidea have

declined since their evolutionary origin (Fig. 2). Diversification dynamics also vary between

biomes, with cold-temperate lineages undergoing the fastest speciation and extinction, but with

limited consequences for net diversification (Fig. 3). In contrast, biome-shifts, via lineage

movements and paleoclimatic change, have resulted in a growing representation of temperate

lineages, particularly as the Earth's climate became cooler during the Oligocene (Fig. 4). Taken

together, our results point to an early start of tropical diversification as the main driver of the

LDG in Coenagrionoidea. Diversity gains due to biome shifts and accelerated speciation in

temperate regions have counteracted the LDG, yet these effects may be at least partially

swamped by a tendency of damselfly lineages to stay in relatively warm areas of the planet and a

relatively fast pace of extinctions outside the tropics.

Our topology inference of Coenagrionoidea recovered phylogenetic relationships that are

largely congruent with other recent molecular phylogenetic studies (Dijkstra et al. 2014;

Toussaint et al. 2019; Bybee et al. 2021). However, our study also calls for the revision of

multiple generic classifications based exclusively on morphological evidence, and for further

phylogenetic studies of a few particularly understudied clades (see Extended Results). We note

that the root node of the tree, which marks the divergence between the pond damselfly and

featherleg families was not resolved in our preliminary analysis, despite being strongly supported

in multiple phylogenomic studies (Kohli et al. 2021; Suvorov et al. 2021; Bybee et al. 2021). In

our preliminary analysis, the featherlegs were recovered as a monophyletic group, but their

relationships to the main clades within the pond damselflies were uncertain. We therefore

imposed a topological constraint to enforce the monophyly of each family in our final

phylogenetic analysis. This resulted in strongly supported relationships between pond damselfly

clades, in agreement with the most comprehensive backbone phylogeny of Odonata to date (Fig.

S5c, Bybee et al. 2021). Nonetheless, enforcing the monophyly of pond damselflies may have

also contributed to underestimating branch lengths at the base of the family (see Table S10).

Ancestral introgression and incomplete lineage sorting in the early history of Coenagrionoidea

are likely causes for phylogenetic discordance, and their effects tend to be aggravated in analyses

based on concatenated loci (Kubatko and Degnan 2007). Moreover, both processes have been

pervasive in the deep history of Odonata, leaving a genetic footprint within and between

damselfly superfamilies (Suvorov et al. 2021).

Our biogeographic dating analysis, combining empirical paleogeography with prior

information on the root age of Coenagrionoidea, resulted in an estimated time for the MRCA of

105 Ma (Fig. S26). This estimate is only slightly younger, yet more uncertain, than the age of

Coenagrionoidea in the phylogenomic studies used for calibration (Kohli et al. 2021; Suvorov et

al. 2021). Our estimates of internal node ages under the strongly informed prior are also

consistent with most fossil records, except for specimens attributed to Platycnemididae, which

have been dated to ~ 99 Ma (Zheng et al. 2017), before the inferred origin of crown

Platycnemididae in this analysis (~ 80 Ma) and in previous studies (Table S10). Compared to

shallow (genus-level) nodes dated in previous studies, inferred ages in this analysis are relatively

older (Table S10). In contrast, the dating analysis with a weakly informed root-age prior,

returned a younger origin time for Coenagrionoidea (Fig. S26), which was also younger than

estimates in previous studies (Table S10). However, this latter analysis produced age estimates

for shallow nodes that were usually closer to estimates in previous studies (see for example

Ischnura, *Enallagma*, *Nesobasis*, *Nehalennia* and *Platycnemis* in Table S10).

These results underscore the challenges in dating phylogenies of groups of organisms

with a scant fossil record, and the limitations of simple models of dispersal and paleogeography.

The statistical approach developed by Landis (2017) is appealing for its ability to jointly sample

dispersal histories and diversification events. However, the computational challenges to the

likelihood calculation over a large space of states currently prohibits incorporating critical

complexity into the paleogeographic model (Ree and Sanmartín 2009). A terrestrial graph at a

global scale seems suitable for widely distributed damselflies, which require fresh water habitats

to breed (Corbet 1999). Nonetheless, many genera of pond damselflies are endemic to specific

archipelagos, while some taxa are widely distributed across continents (Fig. 1; Table S4). In the

present analysis, island endemics are subsumed into larger biogeographic areas or coded as

missing data (see Supplementary Material), and distribution ranges spanning two or more areas

are treated as character-state uncertainty in extant taxa. A model that incorporates branch

heterogeneity in dispersal rates and captures the more fine grained paleogeographic events that

shaped the distribution of island endemics would likely improve biogeographic dating in this and

other diverse groups of insects that vary in their ease of dispersal (e.g. Jones et al. 2016; Landis

et al. 2018) and tend to leave a relatively sparse fossil record (Wills 2001). Yet, such extensions,

as well as the use of large genomic data sets, under a fully Bayesian approach are currently

constrained by computational limitations.

Despite these challenges to node dating, both biogeographic dating analyses under

alternative root-age priors indicate that the MRCA of pond damselflies and featherlegs originated

in either Northern South America or Western Africa (Fig. 1; S27). These areas were separated by

narrower water barriers at the presumed origin of Coenagrionoidea (particularly under the

strongly informed root-age prior) and have preserved tropical rainforests throughout the

Cenozoic, notwithstanding climatic change (Morley 2007; Couvreur et al. 2011). In contrast, the

biome-shift and dispersal analysis inferred the ancestral range of Coenagrionoidea in tropical

South East Asia with relatively high (> 80%) posterior probability (Fig. S41). Two other deep

nodes in the Coenagrionoidea phylogeny, the MRCAs of the featherlegs and 'core' pond

damselflies, were also incongruent between models, while shallower nodes were generally

consistent. An important distinction between the analyses is that our dating model assumed that

all land habitats are equally suitable for a dispersing lineage, whereas the biome-shift model

allowed ecological affinities to influence lineage movements. Another feature that differentiated

the models is that the paleogeography of the Malaysian Archipelago was treated separately from

South East Asia in the dating analysis, while both areas were treated as a single region in the

biome-shift analysis. Our results therefore highlight the sensitivity of ancestral range

reconstructions to assumptions embedded in dispersal graphs. While more complex models

(e.g. with spatial and temporal resolution of dispersal graphs) could improve these inferences, we

think that improving our mechanistic understanding of range dynamics to inform these models

will be crucial to achieve robust inferences of ancestral ranges in ancient and widely distributed

clades such as Coenagrionoidea (Jønsson et al. 2016).

The early evolution of Coenagrionoidea was characterised by fast diversification,

followed by a decline in the rate of accumulation of new species (Fig. 2). This is a common

feature of dated phylogenies (Morlon et al. 2010), which has been subject to considerable

controversy and which can be attributed to several possible causes including statistical artefacts

(reviewed in Moen and Morlon 2014). One possibility is that regional diversification slows down

over time because niche space is reduced as species accumulate, eventually reaching some

ecological limit (Rabosky 2009; Etienne et al. 2011; Etienne and Haegeman 2012). While

plausible if contemporaneous taxa co-occur and compete for finite resources (Herrera-Alsina et

al. 2018), this explanation for a decline in diversification with time seems less likely for non-

adaptive radiations (Rundell and Price 2009; Czekanski-Moir and Rundell 2019). Some

relatively young radiations in pond damselflies and other odonates have resulted from non-

ecological speciation mechanisms, such as sexual selection or sexual conflict, and many closely

related taxa are therefore only weakly ecologically differentiated from each other (McPeek and

Brown 2000; Siepielski et al. 2010; Svensson 2012; Svensson et al. 2018). These features of

damselfly communities suggest that niche partitioning does not strongly constrain speciation,

although such weak niche differentiation might of course impact the longevity of newly formed

lineages (see below). Moreover, our finding of abundant range reversals from warm-temperate to

ancestral tropical biomes (Fig. 5) further argues against ecological diversity limits at a regional

scale.

An apparent slowdown in diversification may also arise from statistical artefacts or

sampling biases. For example, underparametrized models of molecular evolution can lead to

underestimation of branch lengths, particularly in older branches of the tree (Revell et al. 2005).

As diversification rate estimation ultimately relies on the distribution of branch lengths across

the tree, this type of model misspecification can result in an artificial slowdown of diversification

over time. An artificial slowdown can also be a product of incomplete lineage sampling,

especially when taxon sampling is non-random (Cusimano and Renner 2010; Brock et al. 2011).

However, in this study we have used methods that explicitly incorporate sampling fractions of

specific clades as a measure to counter this bias (Höhna 2015). Nonetheless, even when models

are correctly specified and sampling of extant taxa is accounted for, the "push of the past" can

produce an apparent slowdown of diversification over time (Budd and Mann 2018). This

survivorship bias occurs because we can only sample extant descendants from lineages which

had high enough speciation rates as to have survived to the present. Deep branches in the tree

that diversified more slowly, and would have contributed to a more constant estimate of

diversification rates over time, are more likely to have gone extinct over the 105 Myr of

Coenagrionoidea evolution, thereby leaving no descendants from which to infer their lower

diversification rates. The "push of the past" remains a challenge to estimating temporal

diversification dynamics when we lack information of whether and how many entire clades have

gone extinct (Budd and Mann 2018).

The present distribution of pond damselflies, as many other clades, is characterised by

higher species richness near the equator compared to temperate areas (Fig. 4) (Hillebrand 2004;

Kinlock et al. 2018). This LDG in extant Coenagrionoidea is striking, considering that tropical

lineages, particularly from India, South East Asia, and the Malaysian Archipelago, are probably

underrepresented in the largest collections that contributed to this study. With this potential

caveat in mind, our results indicated that none of the hypotheses based on accelerated speciation

or reduced extinction in the tropics (see Introduction) is a likely explanation for why most extant

damselfly species occur at low latitudes. While taking background diversification heterogeneity

into account (Beaulieu and O'Meara 2016), we found similar rates of net diversification between

the tropical and temperate regions of the globe (Fig. 3c). Nonetheless, speciation and extinction

dynamics do tend to differ across latitudes, with both rates being highest in cold-temperate

biomes, and thus cancelling each other out. As a consequence, both warm-temperate and cold-

temperate lineages tend to be relatively short lived and replace one another at a faster rate than

tropical lineages do, resulting in higher turnover rates outside of the tropics (Fig. 3d). Several

studies have uncovered a similar pattern of increased lineage turnover in the temperate region or

in harsher non-tropical environments (Weir and Schluter 2007; Botero et al. 2014; Pyron 2014;

Harvey et al. 2020).

Faster lineage turnover at higher latitudes may come about in at least two ways. First,

paleoclimatic fluctuations (e.g. DeConto et al. 2008) can cause the fragmentation of distribution

ranges, a phenomenon that is particularly pronounced at higher latitudes (Jansson and Dynesius

2002). By repeatedly creating isolated refugia, climate fluctuations may have episodically

fostered allopatric speciation (Weir and Schluter 2004; Milá et al. 2007; Sánchez-Ramírez et al.

2015; Morales-Barbero et al. 2018). However, the same recurrent climatic events can also

increase extinction rates, either because range fragmentation reduces population size, or because

hybridization and competitive exclusion drive extinction upon secondary contact (Dynesius and

Jansson 2000; Barnosky 2005; Botero et al. 2014). Furthermore, climate fluctuations at higher

latitudes might open up novel ecological opportunities that prompt speciation, by constantly

renewing the availability of unoccupied niches (Schluter 2016). High environmental harshness in

these novel environments at high latitudes might also result in the predominance of viability

selection driven by abiotic environmental conditions, potentially increasing extinction risk and

thereby accelerating lineage turnover (Haldane 1937; Gomulkiewicz and Holt 1995; Cutter and

Gray 2016).

Whether it is driven by ecological factors (Cutter and Gray 2016; Schluter 2016), or non-

ecological mechanisms (McPeek and Gavrilets 2006; Svensson et al. 2006; Siepielski et al.

2018), ephemeral speciation in temperate regions does not seem to have a net impact on

diversification. Our results therefore highlight the role of the historical origin of

Coenagrionoidea in the tropics as one of the main causes of the present-day LDG. This historical

early start of diversification in the tropics has maintained the LDG despite an overall decrease in

speciation rate (Fig. 2), and a net gain of lineages through biome shifts into temperate regions,

particularly over the last 30 Myr (Fig. 4). Our findings that biome affinities strongly influenced

ancestral distribution ranges, and that temperate lineages tend to return to the tropics (Fig. 5,

S42), suggest that dispersal out of the tropics has been limited by tropical niche conservatism,

even in periods when warm and cold-temperate biomes were widely available. These results

support previous studies finding stronger niche conservatism in tropical lineages (Smith et al.

2012; Pyron and Wiens 2013), and underscore how the potential for adaptive evolution into

colder biomes is likely a primary factor governing the steepness of the gradient (Rangel et al.

2018).

As a result of an early start of diversification in the tropics, followed by similar net

diversification rates across latitudinal regions, the Coenagrioniodea LDG is expected to persist,

unless an external input of diversity to the temperate areas reverses this general pattern.

Continuing the trend of the last 30 Myr, a trickle of tropical lineages reaching cold biomes could

slowly contribute to a shallower LDG (Fig. 4). However, in a much shorter term, the smoothing

of environmental gradients due of global warming (Loarie et al. 2009), might release adaptive

constraints for lineages dispersing into higher latitudes. Examples of such rapid northward

expansions have already been documented in Odonata (Paulson 2001; Grewe et al. 2013;

Lancaster et al. 2015). The role of dispersal shaping the future steepness of the LDG thus

warrants further investigation.

Conclusion

The damselfly superfamily Coenagrionoidea originated in tropical areas, where most of its

diversity is currently found (Fig. 1). An increasing number of studies have revealed the

importance of both ecology and historical biogeography in producing regional differences in

diversity across distinct clades of plants and animals (e.g. Rolland et al. 2014; Antonelli et al.

2015; Jablonski et al. 2017). Here, we have shown that in pond damselflies and their relatives,

early tropical diversification followed by relatively slow shifts into warm-temperate and later

cold-temperate biomes established the latitudinal diversity gradient observed today. Faster

ecological speciation in temperate regions is unlikely to compensate for this time lag in the

future, as speciation might be more ephemeral at higher latitudes (Fig. 3). However, climate

change and lineage movements have also contributed to an increase of Coenagrionoidea diversity

in cold biomes, currently at high latitudes (Fig. 4). Our results reveal a complex interplay

between history, macroevolutionary processes and dispersal shaping global diversity patterns.

ACKNOWLEDGEMENTS

We are grateful to Hanna Bensch, Robin Pranter, Tammy Ho, Chiara de Pasqual and Yassin

Tschinda for their ceaseless efforts to sample specimens in field and obtain sequence data in the

lab. This study would have been impossible without the generosity and time of all the avid

odonatologists and museum curators who made hundreds of specimens available to us and

helped us with taxonomic identifications. We are particularly grateful to Bill Mauffray and Paul

Skelly at the Florida State Collection of Arthropods, Oliver Flint and Floyd Shockley at the

Smithsonian Institute, Darren Ward at the New Zealand Arthropod Collection, Darío Lijtmaer at

the Argentinian Museum of Natural Sciences, and KD Dijkstra at the Naturalis Biodiversity

Center. Many others contributed with samples and/or taxonomic expertise (Table S2). Specimens

were collected with permits from the Guyana Environmental Protection Agency (Reference

No. 010715 BR001 and 012615 SP: 002), the Secretaría de Ambiente y Desarrollo Sustentable,

Argentina (Ref No. 11084/16) and from the Cameroon Ministry of Forestry and Wildlife (Ref.

No. 0000034). We acknowledge the logistic support from the Iwokrama International Centre for

Rain Forest Conservation and Development, Guyana, the Karanambu Trust, Guyana, the

National Park Administration, Argentina, and the Congo Basin Institute, Cameroon. Funding for

this study was provided by research grants from The Swedish Research Council (VR: grant no.

2016-03356), the Swedish Foundation for International Cooperation in Research and Higher

Education, and Erik Philip Sörensens Stiftelse to E.I.S and from The Royal Physiographic

Society of Lund, the Jörgen Lindströms Fund and "Lunds Djurskyddsfond" to B.W. We thank

Stephen De Lisle and Miguel Gómez-Llano for helpful comments on the manuscript and Brian

P. Hanotte for computational support.

LITERATURE CITED

Abbott J.C., Bota-Sierra C.A., Guralnick R., Kalkman V., González-Soriano E., Novelo-

Gutiérrez R., Bybee S., Ware J., Belitz M.W. 2022. Diversity of nearctic dragonflies and

damselflies (Odonata). Diversity. 14:575.

Anderson R.C. 2009. Do dragonflies migrate across the western Indian Ocean? J. Trop. Ecology.

25:347–358.

Antonelli A., Zizka A., Silvestro D., Scharn R., Cascales-Miñana B., Bacon C.D. 2015. An

engine for global plant diversity: Highest evolutionary turnover and emigration in the

American tropics. Front. Genet. 6:130.

Barnosky A.D. 2005. Effects of Quaternary climatic change on speciation in mammals. J.

Mamm. Evol. 12:247–264.

Beaulieu J.M., O'Meara B.C. 2016. Detecting hidden diversification shifts in models of trait-

dependent speciation and extinction. Syst. Biol. 65:583–601.

Belmaker J., Jetz W. 2015. Relative roles of ecological and energetic constraints, diversification

rates and region history on global species richness gradients. Ecol. Lett. 18:563–571.

Botero C.A., Dor R., McCain C.M., Safran R.J. 2014. Environmental harshness is positively

correlated with intraspecific divergence in mammals and birds. Mol. Ecol. 23:259–268.

Boudot J.-P., Kalkman V.J.(eds). 2015. Atlas of the European dragonflies and damselflies. The

Netherlands: KNNV Publishing.

Brock C.D., Harmon L.J., Alfaro M.E. 2011. Testing for temporal variation in diversification

rates when sampling is incomplete and nonrandom. Syst. Biol. 60:410–419.

Brown J.H. 2014. Why are there so many species in the tropics? J. Biogeogr. 41:8–22.

Budd G.E., Mann R.P. 2018. History is written by the victors: The effect of the push of the past

on the fossil record. Evolution. 72:2276–2291.

Bybee S.M., Kalkman V.J., Erickson R.J., Frandsen P.B., Breinholt J.W., Suvorov A., Dijkstra

K.-D.B., Cordero-Rivera A., Skevington J.H., Abbott J.C., Sánchez-Herrera M., Lemmon

A.L., Lemmon E.M., Ware J.L. 2021. Phylogeny and classification of Odonata using

targeted genomics. Mol. Phylogenet. Evol. 160:107115.

Carle F.L., Kjer K., May M. 2008. Evolution of Odonata, with special reference to

Coenagrionoidea (Zygoptera). Arthropod Syst. 66:37–44.

Corbet P.S. 1999. Dragonflies: Behavior and ecology of Odonata. Colchester, UK: Harley.

Couvreur T.L.P., Forest F., Baker W.J. 2011. Origin and global diversification patterns of

tropical rain forests: Inferences from a complete genus-level phylogeny of palms. BMC

Biol. 9:44.

Cusimano N., Renner S.S. 2010. Slowdowns in diversification rates from real phylogenies may

not be real. Syst. Biol. 59:458–464.

Cutter A.D., Gray J.C. 2016. Ephemeral ecological speciation and the latitudinal biodiversity

gradient. Evolution. 70:2171–2185.

Czekanski-Moir J.E., Rundell R.J. 2019. The ecology of nonecological speciation and

nonadaptive radiations. Trends Ecol. Evol. 34:400–415.

Darwin C.R. 1859. The origin of species by means of natural selection, or the preservation of

favoured races in the struggle for life. London, UK: John Murray.

De Baets K., Antonelli A., Donoghue P.C. 2016. Tectonic blocks and molecular clocks. Philos.

Trans. R. Soc. Lond., B. 371:20160098.

DeConto R.M., Pollard D., Wilson P.A., Pälike H., Lear C.H., Pagani M. 2008. Thresholds for

816   Cenozoic bipolar glaciation. Nature. 455:652–657.

817   Dijkstra K.-D.B. 2017. African Dragonflies and Damselflies. Online: Http://addo.adu.org.za/

818   Dijkstra K.D.B., Kalkman V.J., Dow R.A., Stokvis F.R., Van Tol J.A.N. 2014. Redefining the

819   damselfly families: A comprehensive molecular phylogeny of Zygoptera (Odonata). Syst.

820   Entomol. 39:68–96.

821   Dobzhansky T. 1950. Evolution in the tropics. Amer. Sci. 38:209–221.

822   Dynesius M., Jansson R. 2000. Evolutionary consequences of changes in species' geographical

823   distributions driven by Milankovitch climate oscillations. Proc. Natl. Acad. Sci. U.S.A.

824   97:9115–9120.

825   Economo E.P., Narula N., Friedman N.R., Weiser M.D., Guénard B. 2018. Macroecology and

826   macroevolution of the latitudinal diversity gradient in ants. Nat. Commun. 9:1778.

827   Etienne R.S., Cabral J.S., Hagen O., Hartig F., Hurlbert A.H., Pellissier L., Pontarp M., Storch

828   D. 2019. A minimal model for the latitudinal diversity gradient suggests a dominant role

829   for ecological limits. Am. Nat. 194:E122–E133.

830   Etienne R.S., Haegeman B. 2012. A conceptual and statistical framework for adaptive radiations

831   with a key role for diversity dependence. Am. Nat. 180:E75–E89.

832   Etienne R.S., Haegeman B., Stadler T., Aze T., Pearson P.N., Purvis A., Phillimore A.B. 2011.

833   Diversity-dependence brings molecular phylogenies closer to agreement with the fossil

834   record. Proc. R. Soc. B. 279:1300–1309.

Fischer A.G. 1960. Latitudinal variations in organic diversity. Evolution. 14:64–81.

Gaboriau T., Albouy C., Descombes P., Mouillot D., Pellissier L., Leprieur F. 2019. Ecological

constraints coupled with deep-time habitat dynamics predict the latitudinal diversity

gradient in reef fishes. Proc. R. Soc. B. 286:20191506.

Goldberg E.E., Lancaster L.T., Ree R.H. 2011. Phylogenetic inference of reciprocal effects

between geographic range evolution and diversification. Syst. Biol. 60:451–465.

Gomulkiewicz R., Holt R.D. 1995. When does evolution by natural selection prevent extinction?

Evolution. 49:201–207.

Grewe Y., Hof C., Dehling D.M., Brandl R., Brändle M. 2013. Recent range shifts of European

dragonflies provide support for an inverse relationship between habitat predictability and

dispersal. Global Ecol. Biogeogr. 22:403–409.

Haldane J.B.S. 1937. The effect of variation of fitness. Am. Nat. 71:337–349.

Hanly P.J., Mittelbach G.G., Schemske D.W. 2017. Speciation and the latitudinal diversity

gradient: Insights from the global distribution of endemic fish. Am. Nat. 189:604–615.

Harvey M.G., Bravo G.A., Claramunt S., Cuervo A.M., Derryberry G.E., Battilana J., Seeholzer

G.F., McKay J.S., O'Meara B.C., Faircloth B.C., Edwards S.V., Pérez-Emán J., Moyle

R.G., Sheldon F.H., Aleixo A., Smith B.T., Chesser R.T., Silveira L.F., Cracraft J.,

Brumfield R.T., Derryberry E.P. 2020. The evolution of a tropical biodiversity hotspot.

Science. 370:1343–1348.

Hawkins B.A., Diniz-Filho J.A.F., Jaramillo C.A., Soeller S.A. 2007. Climate, niche

conservatism, and the global bird diversity gradient. Am. Nat. 170:S16–S27.

Heath T.A., Huelsenbeck J.P., Stadler T. 2014. The fossilized birth–death process for coherent

calibration of divergence-time estimates. Proc. Natl. Acad. Sci. U.S.A. 111:E2957–

E2966.

Herrera-Alsina L., Pigot A.L., Hildenbrandt H., Etienne R.S. 2018. The influence of ecological

and geographic limits on the evolution of species distributions and diversity. Evolution.

72:1978–1991.

Hillebrand H. 2004. On the generality of the latitudinal diversity gradient. Am. Nat. 163:192–

211.

Ho S.Y.W., Tong K.J., Foster C.S.P., Ritchie A.M., Lo N., Crisp M.D. 2015. Biogeographic

calibrations for the molecular clock. Biol. Lett. 11:20150194.

Höhna S. 2015. The time-dependent reconstructed evolutionary process with a key-role for mass-

extinction events. J. Theor. Biol. 380:321–331.

Höhna S., Heath T.A., Boussau B., Landis M.J., Ronquist F., Huelsenbeck J.P. 2014.

Probabilistic graphical model representation in phylogenetics. Syst. Biol. 63:753–771.

Höhna S., Landis M.J., Heath T.A., Boussau B., Lartillot N., Moore B.R., Huelsenbeck J.P.,

Ronquist F. 2016. RevBayes: Bayesian phylogenetic inference using graphical models

and an interactive model-specification language. Syst. Biol. 65:726–736.

Jablonski D., Huang S., Roy K., Valentine J.W. 2017. Shaping the latitudinal diversity gradient:

New perspectives from a synthesis of paleobiology and biogeography. Am. Nat. 189:1–

12.

Jablonski D., Roy K., Valentine J.W. 2006. Out of the tropics: Evolutionary dynamics of the

latitudinal diversity gradient. Science. 314:102–106.

Jansson R., Dynesius M. 2002. The fate of clades in a world of recurrent climatic change:

Milankovitch oscillations and evolution. Annu. Rev. Ecol. Evol. Syst. 33:741–777.

Jocque M., Field R., Brendonck L., De Meester L. 2010. Climatic control of dispersal–ecological

specialization trade-offs: A metacommunity process at the heart of the latitudinal

diversity gradient? Global Ecol. Biogeogr. 19:244–252.

Jolly D., Harrison S.P., Damnati B., Bonnefille R. 1998. Simulated climate and biomes of Africa

during the Late Quaternary: Comparison with pollen and lake status data. Quaternary

Science Reviews. 17:629–657.

Jones H.B.C., Lim K.S., Bell J.R., Hill J.K., Chapman J.W. 2016. Quantifying interspecific

variation in dispersal ability of noctuid moths using an advanced tethered flight

technique. Ecol. Evol. 6:181–190.

Jønsson K.A., Tøttrup A.P., Borregaard M.K., Keith S.A., Rahbek C., Thorup K. 2016. Tracking

animal dispersal: From individual movement to community assembly and global range

dynamics. Trends in Ecology & Evolution. 31:204–214.

Kidwell S.M., Holland S.M. 2002. The quality of the fossil record: Implications for evolutionary

analyses. Annu. Rev. Ecol. Evol. Syst. 33:561–588.

Kinlock N.L., Prowant L., Herstoff E.M., Foley C.M., Akin-Fajiye M., Bender N., Umarani M.,

Ryu H.Y., Şen B., Gurevitch J. 2018. Explaining global variation in the latitudinal

diversity gradient: Meta-analysis confirms known patterns and uncovers new ones.

Global Ecol. Biogeogr. 27:125–141.

Kodandaramaiah U. 2011. Tectonic calibrations in molecular dating. Curr. Zool. 57:116–124.

Kohli M., Letsch H., Greve C., Béthoux O., Deregnaucourt I., Liu S., Zhou X., Donath A.,

Mayer C., Podsiadlowski L., Gunkel S., Machida R., Niehuis O., Rust, J., Wappler T., Yu

X., Misof B., Ware J.L. 2021. Evolutionary history and divergence times of Odonata

(dragonflies and damselflies) revealed through transcriptomics. Iscience. 24:103324.

Kubatko L.S., Degnan J.H. 2007. Inconsistency of phylogenetic estimates from concatenated

data under coalescence. Syst. Biol. 56:17–24.

Lancaster L.T., Dudaniec R.Y., Hansson B., Svensson E.I. 2015. Latitudinal shift in thermal

niche breadth results from thermal release during a climate-mediated range expansion. J.

Biogeogr. 42:1953–1963.

Landis M., Edwards E.J., Donoghue M.J. 2021. Modeling phylogenetic biome shifts on a planet

with a past. Syst. Biol. 70:86–107.

Landis M.J. 2017. Biogeographic dating of speciation times using paleogeographically informed

processes. Syst. Biol. 66:128–144.

Landis M.J., Freyman W.A., Baldwin B.G. 2018. Retracing the Hawaiian silversword radiation

despite phylogenetic, biogeographic, and paleogeographic uncertainty. Evolution.

72:2343–2359.

Loarie S.R., Duffy P.B., Hamilton H., Asner G.P., Field C.B., Ackerly D.D. 2009. The velocity

of climate change. Nature. 462:1052–1055.

MacArthur R.H. 1969. Patterns of communities in the tropics. Biol. J. Linn. Soc. 1:19–30.

Magee A.F., Höhna S., Vasylyeva T.I., Leaché A.D., Minin V.N. 2020. Locally adaptive

Bayesian birth-death model successfully detects slow and rapid rate shifts. PLoS Comput.

Biol. 16:e1007999.

May M.L. 2019. Aquatic insects: Behavior and ecology. Aquatic Insects: Behavior and Ecology.

Springer. p. 35–73.

McPeek M.A., Brown J.M. 2000. Building a regional species pool: Diversification of the

Enallagma damselflies in eastern North America. Ecology. 81:904–920.

McPeek M.A., Gavrilets S. 2006. The evolution of female mating preferences: Differentiation

from species with promiscuous males can promote speciation. Evolution. 60:1967–1980.

Meseguer A.L., Condamine F.L. 2020. Ancient tropical extinctions at high latitudes contributed

to the latitudinal diversity gradient. Evolution. 74:1966–1987.

Milá B., McCormack J.E., Castañeda G., Wayne R.K., Smith T.B. 2007. Recent postglacial

range expansion drives the rapid diversification of a songbird lineage in the genus *Junco*.

Proc. R. Soc. B. 274:2653–2660.

Miller E.C., Román-Palacios C. 2021. Evolutionary time best explains the latitudinal diversity

gradient of living freshwater fish diversity. Global Ecol. Biogeogr. 30:749–763.

Misof B., Liu S., Meusemann K., Peters R.S., Donath A., Mayer C., Frandsen P.B., Ware J.,

Flouri T., Beutel R.G., Niehuis O., Petersen M., Izquierdo-Carrasco F., Wappler T., Rust

J., Aberer A.J., Aspöck U., Aspöck H., Bartel D., Blanke A., Berger S., Böhm A.,

Buckley T.R., Calcott B., Chen J., Friedrich F., Fukui M., Fujita M., Greve C., Grobe P.,

Gu S., Huang Y., Jermiin L.S., Kawahara A.Y., Krogmann L., Kubiak M., Lanfear R.,

Letsch H., Li Y., Li Z., Li J., Lu H., Machida R., Mashimo Y., Kapli P., McKenna D.D.,

Meng G., Nakagaki Y., Navarrete-Heredia J.L., Ott M., Ou Y., Pass G., Podsiadlowski

L., Pohl H., Reumont von B.M., Schütte K., Sekiya K., Shimizu S., Slipinski A.,

Stamatakis A., Song W., Su X., Szucsich N.U., Tan M., Tan X., Tang M., Tang J.,

Timelthaler G., Tomizuka S., Trautwein M., Tong X., Uchifune T., Walzl M.G.,

Wiegmann B.M., Wilbrandt J., Wipfler B., Wong T.K.F., Wu Q., Wu G., Xie Y., Yang

S., Yang Q., Yeates D.K., Yoshizawa K., Zhang Q., Zhang R., Zhang W., Zhang Y.,

Zhao J., Zhou C., Zhou L., Ziesmann T., Zou S., Li Y., Xu X., Zhang Y., Yang H., Wang

947   J., Wang J., Kjer K.M., Zhou X. 2014. Phylogenomics resolves the timing and pattern of

948   insect evolution. Science. 346:763–767.

949   Mittelbach G.G., Schemske D.W., Cornell H.V., Allen A.P., Brown J.M., Bush M.B., Harrison

950   S.P., Hurlbert A.H., Knowlton N., Lessios H.A. McCain C., McCune A.R., McDade

951   L.A., McPeek M.A., Near T.J., Price T.D., Ricklefs R.E., Roy K. Sax D.F., Schluter D.,

952   Sobel J.M., Turelli M. 2007. Evolution and the latitudinal diversity gradient: Speciation,

953   extinction and biogeography. Ecol. Lett. 10:315–331.

954   Moen D., Morlon H. 2014. Why does diversification slow down? Trends Ecol. Evol. 29:190–

955   197.

956   Morales-Barbero J., Martinez P.A., Ferrer-Castán D., Olalla-Tárraga M.Á. 2018. Quaternary

957   refugia are associated with higher speciation rates in mammalian faunas of the Western

958   Palaearctic. Ecography. 41:607–621.

959   Moritz C., Patton J.L., Schneider C.J., Smith T.B. 2000. Diversification of rainforest faunas: An

960   integrated molecular approach. Annu. Rev. Ecol. Evol. Syst. 31:533–563.

961   Morley R.J. 2007. Tropical rainforest responses to climatic change. In: Bush M., Flenley J.,

962   Gosling W., editors. Tropical rainforest responses to climatic change. Berlin, Germany:

963   Springer. p. 1–31.

964   Morlon H., Potts M.D., Plotkin J.B. 2010. Inferring the dynamics of diversification: A coalescent

965   approach. PLoS Biol. 8:e1000493.

966   Otto-Bliesner B.L., Brady E.C., Tomas R.A., Albani S., Bartlein P.J., Mahowald N.M., Shafer

967   S.L., Kluzek E., Lawrence P.J., Leguy G., others. 2020. A comparison of the CMIP6

968   midHolocene and lig127k simulations in CESM2. Paleoceanogr. Paleoclimatol.

969   35:e2020PA003957.

Owens H.L., Lewis D.S., Dupuis J.R., Clamens A.L., Sperling F.A., Kawahara A.Y., Guralnick

R.P., Condamine F.L. 2017. Latitudinal diversity gradient in New World swallowtail

butterflies is caused by contrasting patterns of out-of- and into-the-tropics dispersal.

Global Ecol. Biogeogr.

Paulson D. 2009. Dragonflies and Damselflies of the West. Princeton University Press.

Paulson D. 2011. Dragonflies and Damselflies of the East. Princeton University Press.

Paulson D.R. 2001. Recent Odonata records from southern Florida-effects of global warming?

Int. J. Odonatol. 4:57–69.

Pianka E.R. 1966. Latitudinal gradients in species diversity: A review of concepts. Am. Nat.

100:33–46.

Polato N.R., Gill B.A., Shah A.A., Gray M.M., Casner K.L., Barthelet A., Messer P.W.,

Simmons M.P., Guayasamin J.M., Encalada A.C., Kondratieff B.C., Flecker A.S.,

Thomas S.A., Ghalambor C.K., Poff L., Funk W.C., Zamudio K. 2018. Narrow thermal

tolerance and low dispersal drive higher speciation in tropical mountains. Proc. Natl.

Acad. Sci. U.S.A.. 115:12471–12476.

Pyron R.A. 2014. Temperate extinction in squamate reptiles and the roots of latitudinal diversity

gradients. Global Ecol. Biogeogr. 23:1126–1134.

Pyron R.A., Wiens J.J. 2013. Large-scale phylogenetic analyses reveal the causes of high

tropical amphibian diversity. Proc. R. Soc. B. 280:20131622.

R Core Team. 2021. R: A language and environment for statistical computing. Vienna, Austria:

R Foundation for Statistical Computing.

Rabosky D.L. 2009. Ecological limits and diversification rate: Alternative paradigms to explain

the variation in species richness among clades and regions. Ecol. Lett. 12:735–743.

Rabosky D.L., Chang J., Title P.O., Cowman P.F., Sallan L., Friedman M., Kaschner K., Garilao

C., Near T.J., Coll M., Alfaro M.E. 2018. An inverse latitudinal gradient in speciation

rate for marine fishes. Nature. 559:392–395.

Rangel T.F., Edwards N.R., Holden P.B., Diniz-Filho J.A.F., Gosling W.D., Coelho M.T.P.,

Cassemiro F.A.S., Rahbek C., Colwell R.K. 2018. Modeling the ecology and evolution of

biodiversity: Biogeographical cradles, museums, and graves. Science. 361:eaar5452.

Ree R.H., Sanmartín I. 2009. Prospects and challenges for parametric models in historical

biogeographical inference. J. Biogeogr. 36:1211–1220.

Revell L.J., Harmon L.J., Glor R.E. 2005. Under-parameterized model of sequence evolution

leads to bias in the estimation of diversification rates from molecular phylogenies. Syst.

Biol. 54:973–983.

Rolland J., Condamine F.L., Beeravolu C.R., Jiguet F., Morlon H. 2015. Dispersal is a major

driver of the latitudinal diversity gradient of Carnivora. Global Ecol. Biogeogr. 24:1059–

1071.

Rolland J., Condamine F.L., Jiguet F., Morlon H. 2014. Faster speciation and reduced extinction

in the tropics contribute to the mammalian latitudinal diversity gradient. PLoS Biol.

12:e1001775.

Ronquist F., Klopfstein S., Vilhelmsen L., Schulmeister S., Murray D.L., Rasnitsyn A.P. 2012. A

total-evidence approach to dating with fossils, applied to the early radiation of the

Hymenoptera. Syst. Biol. 61:973–999.

Rosenzweig M.L., others. 1995. Species diversity in space and time. Cambridge University

Press.

Roy K., Goldberg E.E. 2007. Origination, extinction, and dispersal: Integrative models for

understanding present-day diversity gradients. Am. Nat. 170:S71–S85.

Rundell R.J., Price T.D. 2009. Adaptive radiation, nonadaptive radiation, ecological speciation

and nonecological speciation. Trends Ecol. Evol.. 24:394–399.

Salisbury C.L., Seddon N., Cooney C.R., Tobias J.A. 2012. The latitudinal gradient in dispersal

constraints: Ecological specialisation drives diversification in tropical birds. Annu. Rev.

Ecol. Evol. Syst. 15:847–855.

Salzmann U., Haywood A.M., Lunt D.J., Valdes P.J., Hill D.J. 2008. A new global biome

reconstruction and data-model comparison for the Middle Pliocene. Global Ecol.

Biogeogr. 17:432–447.

Saupe E.E., Myers C.E., Peterson A.T., Soberón J., Singarayer J., Valdes P., Qiao H. 2019.

Spatio-temporal climate change contributes to latitudinal diversity gradients. Nat. Ecol.

Evol. 3:1419–1429.

Sánchez-Ramírez S., Tulloss R.E., Guzmán-Dávalos L., Cifuentes-Blanco J., Valenzuela R.,

Estrada-Torres A., Ruán-Soto F., Díaz-Moreno R., Hernández-Rico N., Torres-Gómez

M., León H., Moncalvo J.-M. 2015. In and out of refugia: Historical patterns of diversity

and demography in the North American Caesar's mushroom species complex. Mol. Ecol.

24:5938–5956.

Schemske D.W., Mittelbach G.G., Cornell H.V., Sobel J.M., Roy K. 2009. Is there a latitudinal

gradient in the importance of biotic interactions? Annu. Rev. Ecol. Evol. Syst. 40:245–

269.

Schluter D. 2016. Speciation, ecological opportunity, and latitude: (American Society of

Naturalists Address). Am. Nat. 187:1–18.

Siepielski A.M., Hasik A.Z., Ousterhout B.H. 2018. An ecological and evolutionary perspective

on species coexistence under global change. Curr. Opin. Insect Sci. 29:71–77.

Siepielski A.M., Hung K.-L., Bein E.E.B., McPeek M.A. 2010. Experimental evidence for

neutral community dynamics governing an insect assemblage. Ecology. 91:847–857.

Smith B.T., Bryson Jr R.W., Houston D.D., Klicka J. 2012. An asymmetry in niche conservatism

contributes to the latitudinal species diversity gradient in New World vertebrates. Ecol.

Lett. 15:1318–1325.

Spano C.A., Hernández C.E., Rivadeneira M.M. 2016. Evolutionary dispersal drives the

latitudinal diversity gradient of stony corals. Ecography. 39:836–843.

Stadler T. 2011. Mammalian phylogeny reveals recent diversification rate shifts. Proc. Natl.

Acad. Sci. U.S.A. 108:6187–6192.

Stephens P.R., Wiens J.J. 2003. Explaining species richness from continents to communities:

The time-for-speciation effect in emydid turtles. Am. Nat. 161:112–128.

Stevens R.D. 2006. Historical processes enhance patterns of diversity along latitudinal gradients.

Proc. R. Soc. B. 273:2283–2289.

Suvorov A., Scornavacca C., Fujimoto M.S., Bodily P., Clement M., Crandall K.A., Whiting

M.F., Schrider D.R., Bybee S.M. 2021. Deep ancestral introgression shapes evolutionary

history of dragonflies and damselflies. Syst. Biol.:syab063.

Svensson E.I. 2012. Non-ecological speciation, niche conservatism and thermal adaptation: How

are they connected? Org. Divers. Evol. 12:229–240.

Svensson E.I., Eroukhmanoff F., Friberg M. 2006. Effects of natural and sexual selection on

adaptive population divergence and premating isolation in a damselfly. Evolution.

60:1242–1253.

Svensson E.I., Gómez-Llano M.A., Torres A.R., Bensch H.M. 2018. Frequency dependence and

ecological drift shape coexistence of species with similar niches. Am. Nat. 191:691–703.

Tiley G.P., Poelstra J.W., Dos Reis M., Yang Z., Yoder A.D. 2020. Molecular clocks without

rocks: New solutions for old problems. Trends Genet.

Toussaint E.F.A., Bybee S.M., Erickson R.J., Condamine F.L. 2019. Forest giants on different

evolutionary branches: Ecomorphological convergence in helicopter damselflies.

Evolution. 73:1045–1054.

Troast D., Suhling F., Jinguji H., Sahlén G., Ware J. 2016. A global population genetic study of

Pantala flavescens. PLoS One. 11:e0148949.

Wallace A.R. 1876. The geographical distribution of animals. Macmillan; Company.

Wallace A.R. 1878. Tropical nature, and other essays. Macmillan; Company.

Waller J.T., Svensson E.I. 2017. Body size evolution in an old insect order: No evidence for

Cope's Rule in spite of fitness benefits of large size. Evolution. 71:2178–2193.

Weir J.T., Schluter D. 2004. Ice sheets promote speciation in boreal birds. Proc. R. Soc. B

271:1881–1887.

Weir J.T., Schluter D. 2007. The latitudinal gradient in recent speciation and extinction rates of

birds and mammals. Science. 315:1574–1576.

Wickham H. 2016. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.

Wiens J.J. 2007. Global patterns of diversification and species richness in amphibians. Am. Nat.

170:S86–S106.

Wiens J.J. 2011. The causes of species richness patterns across space, time, and clades and the

role of "ecological limits." Q. Rev. Biol. 86:75–96.

Wikelski M., Moskowitz D., Adelman J.S., Cochran J., Wilcove D.S., May M.L. 2006. Simple

rules guide dragonfly migration. Biol. Lett.. 2:325–329.

Wills M.A. 2001. How good is the fossil record of arthropods? An assessment using the

stratigraphic congruence of cladograms. Geol. J. 36:187–210.

Wright S., Keeling J., Gillman L. 2006. The road from Santa Rosalia: A faster tempo of

evolution in tropical climates. Proc. Natl. Acad. Sci. U.S.A. 103:7718–7722.

Yu G., Smith D., Zhu H., Guan Y., Lam T.T.-Y. 2017. ggtree: An R package for visualization

and annotation of phylogenetic trees with their covariates and other associated data.

Methods Ecol. Evol. 1:28–36.

Zheng D., Wang B., Chang S.-C. 2017. *Palaeodisparoneura cretacica* sp. Nov., A new

damselfly (Odonata: Zygoptera: Platycnemididae) from mid-Cretaceous Burmese amber.

Comptes Rendus Palevol. 16:235–240.

Figure Legends

Figure 1. Inferring the biogeographic history of pond damselflies and their relatives the

featherlegs (superfamily Coenagrionoidea). Speciation times and ancestral ranges were jointly

estimated using a time-heterogeneous continuous-time Markov model, based on empirical

paleogeography, an informed root-age prior and molecular and distribution data of 669 extant

taxa. The empirical paleogeographic model in Landis (2017) was used to determine dispersal

edges between 25 geographic areas across the globe (Greenland, Antarctica (E) and Antarctica

(W) not shown in the colour legend). Phylogenetic inference was summarized using a maximum

a posteriori (MAP) tree. The distribution areas occupied by each of the extant taxa are shown at

the tips of the tree. Pies show the proportion of each ancestral state in a random sample of 1000

posterior trees. The cladogram in the lower right shows the inferred ancestral ranges for the most

recent common ancestor of Coenagrionoidea, and for each of the three major clades in the

superfamily. The fraction of taxa sampled is shown in parenthesis. One representative species is

illustrated for each clade: *Acanthagrion adustum* ('core'), *Leptagrion elongatum* (ridge-face'),

Platycnemis pennipes (featherlegs). Photos: EIS.

Figure 2. Diversification rate changes since the MRCA of Coenagrionoidea: a) speciation, b)

extinction and c) net diversification. Diversification rates were estimated on the maximum *a*

posteriori (MAP) tree from the biogeographic phylogenetic dating analysis using a root age

calibration based of previous phylogenomic studies (Kohli et al. 2021; Suvorov et al. 2021).

Changes in diversification rates were modelled under an episodic birth-death (EBD) process,

over 20 equal-length time intervals and assuming autocorrelation among consecutive time

intervals (see Extended Methods).

Figure 3. Latitudinal effects on diversification of pond damselflies and their relatives the

featherlegs (superfamily Coenagrionoidea). (a) Speciation and (b) extinction rates at tropical and

temperate latitudes were estimated using a Hidden-State Dependent Speciation and Extinction

(HiSSE) model that accounts for background heterogeneity in diversification rates across the

phylogeny, by including a hidden trait with two states. (c) Diversification was calculated as the

net difference between speciation and extiction and (d) turnover was calculated as the ratio

between extinction and speciation. The histograms show the posterior distribution of parameter

estimates.

Figure 4. Proportion of Coenagrionoidea lineages in each biome-region state and through time.

Lineage proportions were computed from stochastic histories sampled in a time-heterogeneous

model of biome-region shifts (Landis et al. 2021). The dark-grey to light-grey bars on the right

show the proportions of the three biome states in extant taxa sampled in this study (669 out of

~1,800 species).

Figure 5. Proportion of biome-shift events (a-c) and event series (d) through time. Event

proportions were estimated from stochastic histories sampled in a time-heterogeneous model of

biome-region shifts. Circles represent posterior means, bars represent 80% HPD intervals, and

lines represent 95% HPD intervals. See Methods for descriptions of the event series types.