Delayed Oligodendrocyte Progenitor Cell Recruitment and Maturation in Large Volume

Demyelination

James J. M. Cooper[1], Jessie J. Polanco[2], Darpan Saraswat[1], Jennifer Peirick[3], Fraser J. Sim[1,2]

Department of Pharmacology and Toxicology[1], Neuroscience Program[2], Lab Animal Facilities[3],

Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY.

Address correspondence to:

Fraser Sim, Ph.D.

Department of Pharmacology and Toxicology

Jacobs School of Medicine and Biomedical Sciences

University at Buffalo

955 Main Street

Buffalo, NY 14203

716-829-2151

fjsim@buffalo.edu

Keywords: demyelination, remyelination, oligodendrocyte progenitor.

Running title: Oligodendrocyte progenitors and large volume demyelination

Abstract, 278; Introduction, 742; Discussion, 1491; Tables: 0, Figures: 8

Acknowledgements: This work was supported by NINDS grant (R01NS104021), National

Multiple Sclerosis Society (RG-1701-26750), the Kalec Multiple Sclerosis Foundation, the

Change MS Foundation. JJP received additional support from NIGMS (R25GM09545902),

NCATS (UL1TR001412-S1), and New York State Department of Health (Empire State Stem Cell

Fund) through the Stem Cells in Regenerative Medicine Fellowship (NYSTEM C30290GG). We

acknowledge support provided by the Center for Computational Research at the University at

Buffalo for use of the academic compute cluster for training the Keras-based convolution network.

Conflict of Interest: The authors declare no competing financial interests.

Abstract

Remyelination-inducing therapies have the potential to restore neurologic function and prevent

progression in multiple sclerosis (MS). However, preclinical therapies capable of driving

oligodendrocyte differentiation and enhancing remyelination in murine models have not yet

yielded meaningful clinical improvement. These setbacks suggest that small animal models may

be inadequate for the development and identification of promising regenerative therapies. Unlike

MS, rodent models of demyelination undergo highly efficient remyelination, have typically very

small volumes and exhibit abundant oligodendrocyte recruitment rapidly following demyelination.

We hypothesized that oligodendrocyte progenitor cell (OPC) recruitment becomes rate-limiting as

lesion volume increases leading to a failure of oligodendrocyte differentiation and endogenous

remyelination. In this study, we utilized lysolecithin injection into the large white matter tracts of

the rabbit to create demyelinated lesions that have cross-sectional areas 20-fold larger than those

previously reported in mice, and volumes 8-fold larger. Unlike rodent lesions which show robust

supernumerary OPC recruitment (representing a 3-10 fold increase in Olig2$^+$ cell density),

recruitment in the rabbit never exceeds normal levels. Importantly, the density of mature CC1$^+$

oligodendrocytes failed to return to NAWM levels by 56 dpl and the percentage of differentiated

oligodendrocytes drastically lagged behind that of rodent lesions. These results suggest that as

lesion volume increases the requirement for increased OPC proliferation and migration becomes

limiting. As oligodendrocyte differentiation from human OPCs is dependent on the local density

of progenitors, the reduced density of progenitors in large lesions may contribute to failed

differentiation and remyelination. Together, we show that large volume demyelination in the white

matter of the rabbit recapitulates several key features of human MS lesions, and represents an

important preclinical model for the assessment of remyelination therapies in human demyelinating

disease.

INTRODUCTION

Primary demyelination is most commonly associated with multiple sclerosis (MS), where half of

patients will require walking assistance after a decade of disease duration [1]. Immune-mediated

demyelination and loss of oligodendrocytes (OL) results in loss of neurological function. In

experimental models, remyelination relies on restoration of the OL population through the

recruitment (both migration and proliferation) and subsequent differentiation and maturation of

oligodendrocyte precursor cells (OPCs) [2]. OPCs are a multipotent stem cell population that are

distributed throughout the adult CNS. While the role of adult OPCs in normal brain remains

elusive, OPCs are responsible for the replacement of lost OLs in the context of demyelination [3].

Following a demyelinating event, OPCs migrate into the damaged region from adjacent healthy

tissue where they become activated [4] and begin to proliferate. The expression of activation

markers, such as Sox2, are downregulated as OPCs begin to differentiate and mature [4]. Newly

generated OLs engage and ensheath denuded axons and upregulate myelin proteins resulting in

the production of a thin myelin sheath with shorter internodes that is capable of restoring saltatory

conduction and neurological function [5].

While remyelination is a common feature of early relapsing-remitting MS, remyelination

failure becomes more common with increasing disease duration and age. Remyelination failure

is thought to be due to the malfunction of one or more stages of oligodendrocyte regeneration:

namely OPC recruitment, differentiation, and myelin formation [2]. While quiescent OPCs are

associated with chronic demyelination in MS [6], the prevailing theory is that remyelination fails in

MS due to a failure of differentiation. In response, a number of drug candidates aimed at boosting

endogenous remyelination through the promotion of OPC differentiation have been developed

and advanced to clinical trials after successful preclinical results, notably opicinumab (anti-Lingo1)

and clemastine [7-9]. Unfortunately, while both were able to significantly improve visual evoked

potential latency indicative of remyelination in the optic nerve, neither was able to significantly

impact clinical outcome or imaging endpoints in initial trials [10-12]. The paradoxical success of

differentiation-promoting therapies in rodent models, but little success in clinical trials suggests

fundamental differences in the rate-limiting steps between remyelination in animal models and

human MS.

In rodent models of demyelination, there is little evidence that the rate of OPC recruitment

is rate-limiting. For example, when the number of OPCs within the lesion is experimentally

increased via overexpression of PDGF-A there is no change in the time course or extent of

remyelination [13]. Additionally, relatively few rounds of division are required to restore OPC density

in rodent lesions (1-3 cycles) [14], and when nearly all OPCs are pharmacogenetically depleted in

the rodent brain OPC densities are restored in less than 10 days [15]. Indeed, it is only after long-
term administration of cuprizone to chronically deplete of OPCs over the period of several months
that recruitment fails and remyelination is inefficient [16]. As OPCs are recruited from the adjacent
NAWM, the burden of both migration and proliferation disproportionately increases with increased
lesion volume. In MS, individual demyelinating lesions have volumes of several thousand cubic
millimeters [17]. In contrast, the volume of focal demyelinated lesions modelled in rodents is very
small, typically less than 1 mm^3 [18, 19]. Thus, we hypothesized that OPC recruitment becomes rate-
limiting as lesion volume increases leading to a failure of OL differentiation and endogenous
remyelination.

In this study we sought to develop a model that more accurately recapitulates the burden
of recruitment seen in MS lesions. To do this we utilized the large white matter tracks of the rabbit
to create demyelinating lesions with cross-sectional areas 20 times that produced in mice. Along
with increased lesion size, OPC recruitment was substantially inferior to corresponding rodent
models and closely resembled the pattern observed in MS. Large volume rabbit lesions exhibited
distinct regional differences in the pattern of gliotic response, axonal injury, OPC recruitment,
activation and proliferation that evolved with time following induction of demyelination. Together,
we describe lysolecithin-induced lesions in the rabbit that recapitulate several key features of
demyelination in MS which may more accurately predict the outcome of novel remyelination-
inducing therapies in clinical trials.

MATERIALS AND METHODS

Animals and surgery

All experiments were performed according to protocols approved by the University at

Buffalo's Institutional Animal Care and Use Committee. New Zealand White Rabbits were

purchased as needed (Female, average weight 2.96 ± 0.13 kg and average age 15.79 ± 0.53

weeks) (Charles Rivers Laboratories, Wilmington, MA). Induction anesthesia was accomplished

via intramuscular injection of ketamine (35 mg/kg) and xylazine (5 mg/kg) into epaxial muscles.

After loss of consciousness, supplemental heating began and corneas were lubricated. The

proximal 3 cm of the tail was shaved to provide an optimal location for the pulse oximetry probe

(far preferable to use of ear or tongue). ECG leads and rectal temperature probe were placed.

After pre-oxygenation via face mask, rabbits were intubated in ventral recumbency with heads

and necks held in extension. A size 3.0 endotracheal tube was threaded over an 8-French urinary

catheter. After placement of the catheter through the glottis, the ET tube was advanced over the

catheter and into the trachea. Successful intubation was verified via immediate capnography.

Upon success, isoflurane was delivered via Bain Circuit (1-3%). An IV catheter was placed

(marginal ear vein), and the surgical site was shaved and prepped prior to moving the rabbit to

the operating room (OR). Pre-emptive, initial analgesics were administered (0.05 mg/kg

buprenorphine, subcutaneously (SC) and 1.5 mg/kg carprofen, SC).

Once moved onto the OR table, rabbits were immediately reconnected to isoflurane to ensure a

suitable anesthetic plane. Then, they were transiently disconnected from the Bain Circuit to allow

loose movement of the head into the stereotaxic frame (Kopf Model 902 Small Animal Stereotaxic

Instrument + Kopf Model 1240 Rabbit Adaptor + Rabbit Risers, Kopf Instruments, Tujunga, CA).

The endotracheal (ET) tube, was carefully moved to rest between the arms of the Kopf Rabbit

Adaptor. It was important that the ET tube was not advanced beyond 12 cm to avoid bronchial

obstruction. Once positioned, the capnograph and Bain Circuit were quickly reconnected. At this

time further careful positioning of the rabbit's head was possible via tooth bar, zygoma clamps

and nose clamp. General anesthesia could be maintained for the remainder of the stereotaxic

procedure with isoflurane. Spontaneous respirations were augmented by manual intermittent

positive pressure ventilations (IPPV). 0.1 mL bupivacaine (0.25%) or lidocaine (2%, diluted 1:1

with 0.9% NaCl) was injected SC. A 7-8 cm incision was made alone the midline of the scalp to

expose both bregma (intersection of the sagittal and coronal sutures) and lambda (intersection of

the sagittal and lambdoid sutures). Both left and right arm Hamilton needles were filled with

distilled H_2O to prevent occlusion and ensure flow prior to lysolecithin loading. Placing the needles

point on either bregma or lambda, the dorsal-ventral (DV) relationship was adjusted using the

Kopf Rabbit Adaptor until bregma was 1.5 mm above lambda. The arms of the Kopf frame were

adjusted to a 20° angle, and the needle points were placed on bregma. The anterior-posterior

(AP), lateral-medial (LM), and DV of each arm was recorded. The following adjustments were

made to these values: AP= +0.5 mm, LM= +7.7 mm, DV= -5.7 mm. Injection site was confirmed

via Evans Blue injection (**Supplemental Figure 3A**).Drill bits (Burrs for Micro Drill 19007-29, 2.9

mm, Foster City, CA) were sterilized via dry micro bead bench side sterilizer for 1 minute and

cooled with room temp saline. In advance of drilling, topical articaine (4%) was dripped onto the

skull and allowed to absorb and dry over drilling skull sites. Drill holes were made through the

skull to the level of the meninges with constant saline stream to prevent parenchymal heat

damage. A 30G needle was then used to puncture the meninges to allow for free passage of the

Hamilton needle. 1% lysolecithin was then injected at a rate 0.1 µl/min, with 5 minute wait times

between microliters to allow for diffusion. Once injection was completed, the needles were left in

place for 20 additional minutes to allow for diffusion.

Post-operatively, buprenorphine was administered (0.05 mg/kg, SC) 4 hours after the initial dose,

and carprofen (1.5 mg/kg, SC) was continued twice daily for a total of 4 consecutive days. Rabbits

were monitored daily after surgery and assessed for: mentation, weight loss from baseline,

appetite, water consumption, amount and character of feces, amount of urination, and presence

of neurologic signs. Post-surgical weight loss peaked at 28 dpl, followed by a rapid recovery

(**Supplemental Figure 3B-C**).

Tissue processing

Animals were sacrificed at 7, 14, 21, and 56 days post-lesion (dpl) by transcardial

perfusion of saline followed by 4% paraformaldehyde under deep anesthesia. Following

decapitation, whole brain tissue was extracted and post-fixed for 30 minutes in 4%

paraformaldehyde. For cryopreservation, the tissue was first left in 1x PBS overnight, then

transferred into a 7.5% sucrose solution overnight, and finally 15% sucrose overnight.

Cryoprotected tissue was then frozen in optimal cutting temperature medium (Tissue-Tek). Serial,

16 µm-thick coronal sections were cut using a Leica cryostat, and stored in -80 °C freezer until

processing.

To identify lesion location and volume, every 10[th] section was washed with 1x PBST (x3

by 5 min), stained with FloroMyelin for 1 hour at room temperature (1:300, Thermo Fisher

Scientific) and 4′,6-Diamidino-2-phenylindole dihydrochloride (DAPI) for 3 minutes (1:5000,

Sigma Aldrich), and mounted in prolong gold (Thermo Fisher Scientific). Lesions were identified

as regions with lesser FloroMyelin signal and corresponding hypercellularity by DAPI, compared

to surrounding white matter. Cross-sectional lesion areas were calculated for both left and right

hemisphere lesions using the NIH ImageJ software, and the largest cross-sectional area for each

lesion was identified as the lesion center.

Estimation of lesion volume

Lesion volume was calculated from the measured cross-sectional areas determined by

FloroMyelin staining using Cavalieri's estimator of morphometric volume [56]:

$$V_c = d \left(\sum_{i=0}^{n} (y_i) \right) - (t) y_{MAX}$$

Where V_c is the Cavalieri's estimated volume, d is the distance between sections, y_i is the cross-

sectional area of section i, t is the thickness of the sections, and y_{MAX} is the largest cross-

sectional area of the lesion. Since every 10th section was stained, $d = 160$ μm, and $t = 16$ μm.

Immunohistochemistry (IHC)

Slides immediately adjacent to the lesion centers, identified by FloroMyelin staining, were

used for all immunohistochemical procedures. In general, sections were permeabilized for 1 hour

with 1% Triton X-100 (Alfa Aesar) and 0.25% Tween 20 (Calbiochem). Sections were blocked for

1 hour with a solution containing 0.5% Triton X-100, 5% bovine serum albumin (BSA), and 5%

normal donkey serum. Primary antibodies utilized were goat anti-Olig2 (1:100, R&D Systems),

mouse anti-CC1 (1:50, Millipore), mouse IgG2a anti-SOX2 (1:100, R&D Systems), mouse IgG1

anti-Ki67 (1:25, Fisher Scientific), rat IgG2a anti-MBP (1:300, Abcam Inc.), goat IgG Smi311 &

Smi312 (1:1000, BioLegend), rabbit IgG anti-App (1:500, Thermo Fisher), rabbit IgG anti-Iba1

(1:300, Wako Chemicals), and mouse IgG1 anti-Gfap (1:400, Sigma-Aldrich). Alexa 488, 594,

and 647-conjugated secondary antibodies (Invitrogen) were used at 1:500. Lesion areas were

captured at 10X and 20X magnification using an Olympus IX83 with IX3-SSU motorized stage

and image tiles stitched via cellSens software. Images of immunostained sections were aligned

with their corresponding FloroMyelin stained section to accurately mark lesion boundaries, with

exclusion of any portion of the lesion which extends into the gray matter. Cells counts were

determined via a deep learning (machine learning) augmented process (code available upon

request), validated with multi-counter manual counts. The Keras based convolution network was

trained on a set of ~3250 CC1 positive and ~3250 CC1 negative pre-classified images

(supervised learning).

Statistics

All statistical testing was completed using GraphPad Prism Version 8.4.0 (455). The

control group for all analyses of cell number and phenotype was NAWM. Groups comprised either

time-points following injection of lysolecithin and/or regional analysis within the lesion. For all

reporting, an experimental unit comprised a single lesion. Given the novelty of the method, sample

size was based on previous experience with gliotoxin lesions established in mouse and rat. A

minimum of 4 experimental units (lesions) per group was set *a priori* and along with a requirement

for at least three animals per group. Criteria for lesion inclusion were established *a priori*. Only

lesions located within internal capsule white matter were included in the analysis. No white matter

lesions regardless of size were excluded from the analysis. Rabbits were allocated to each group

in a randomized fashion based on availability from the vendor and surgical schedule. Manual cell

counting for all quantitative measures was performed in a blinded fashion. Normality of data was

tested in GraphPad using the D'Agostino-Pearson omnibus test. For analysis of time-series data,

one-way ANOVA with Tukey's post-hoc testing was performed. For analysis of time-series and

region delimited data, two-way ANOVA with Sidak's post-hoc testing was performed.

Data availability

The data that support the findings of this study are available upon reasonable request.

RESULTS

Abundant OPC recruitment is universal in rodent models but is relatively uncommon in
MS lesions.

To compare OPC recruitment between experimental models and human disease, we performed
a meta-analysis of absolute and relative OPC densities following demyelination. While the
recruitment of OPCs occurs following demyelination in MS, the density of OPCs found within
lesions does not exceed that of normal appearing white matter (NAWM) (**Supplemental Figure**
1A). In contrast, rodent models of toxin-induced demyelination are associated with
supernumerary OPC recruitment and highly efficient remyelination [20, 21](**Supplemental Figure**
1B). Likewise, following inflammatory demyelination or viral-mediated encephalomyelitis with
attendant axonal injury and immune cell responses, OPC recruitment is similarly robust
(**Supplemental Figure 1C**). As such, we hypothesized that supernumerary densities of OPCs
are necessary for efficient remyelination, and that MS lesion tissue displays relatively weak OPC
recruitment because OPC recruitment becomes rate-limiting as lesion volume increases.

Lysolecithin-induced lesions in the rabbit were larger and more persistent than those
created in smaller rodent models.

In order to test the hypothesis that lesion volume influences OPC recruitment we sought an
experimental species which had white matter structures that would allow for the production of
lesions significantly larger than those induced in mice and rats. We selected the rabbit brain as
the accessible white matter volume was 6 to 14 times larger [22] then current rodent models, and
we predicted that lesions could be generated with cross-sectional areas between 10 and 40 times
greater than rat and mouse brain (**Fig. 1A**, ovals). To induce demyelination, we stereotactically
injected lysolecithin into the internal capsules of adult NZW rabbits (bilateral injections of 1%
lysolecithin, 5 µl). We first examined lesion formation at 14 days post lesion (dpl). Similar to murine
lesions, lysolecithin induced profound focal demyelination at the site of injection and a clearly
demarcated region of hypercellularity that precisely matched the region of myelin loss (**Fig. 1B**,
white dashed line). Volumetric analysis by serial section reconstruction of the lesion indicated a
comet-like appearance with a relatively spherical region of demyelination corresponding to the
site of injection and a tapered region extending in the posterior-ventral direction, likely due to
diffusion within the fascicles of the internal capsule. The mean cross-sectional area and volume
peaked at 14 dpl (area: 2.21 ± 0.3 mm^2, volume: 3.29 ± 0.3 mm^3, $n = 8$) (**Fig. 1C-D**). Compared
to mouse, rabbit lesions were over 8-fold larger in volume and a 20-fold increase in maximal
cross-sectional area (**Fig. 1B-C**). Intriguingly, the 8-fold increased volume in rabbit exceeded the

5-fold increase in volume of lysolecithin injected. Following the peak in lesion area at 14 dpl, lesion

area gradually declined with time (21 vs 14 days, Tukey's p = 0.0280). In stark contrast to both

mouse and rat models, large regions of demyelination persisted in the rabbit even at 56 dpl by

which time rodent models have typically completed remyelination (**Fig. 1E**).

Rabbit lesions represent areas of demyelination with preserved axons and prolonged

gliotic response.

Next, we sought to characterize the extent of the gliotic response in these lesions, as well as

determine the extent of axonal injury and loss in this new model. MBP staining showed abundant

myelin debris at 7 dpl, with substantial clearance by 14 dpl (**Fig. 2A**). At 56 dpl, there was a partial

recovery of MBP immunoreactivity, but lacked the structure observed in NAWM (**Fig. 2A**).

Consistent with primary demyelination, a large number of neurofilament (NF)-labelled axons

persisted following demyelination (7-56 dpl) (**Fig. 2B**). We observed a gradient of axonal density

with greater numbers of axons at the lesion edge (**Fig. 2B**) (**Fig. 3**). Interestingly, low axonal

densities in the lesion core is similarly observed in MS lesions [23]. Additionally, axonal density did

not decrease further beyond 7 dpl, based on a qualitative analysis (**Fig. 3**). Consistent with an

initial injury, we observed evidence of axonal dysfunction by accumulation of App

immunoreactivity at 7 dpl and a decline in the overall proportion of App$^+$NF$^+$ axons thereafter (**Fig.**

2B). Retraction bulbs, a common feature of MS lesion tissue [17], were noted at all timepoints in

and around the lesions (**Fig. 3**).

All lesions exhibited increased numbers of astrocytes (Gfap) and microglia/macrophages

(Iba1), with the greatest number of microglia localized within the lesion core (**Fig. 2C-D**). Within

the core, ameboid microglia commonly formed vesicular structures that were surrounded by Gfap$^+$

astrocytic processes (**Supplemental Figure 2**). In addition to astrocytes and microglia, fibroblasts

are recruited into mouse spinal cord lysolecithin and EAE lesions [24]. However, we observed few

Col1$^+$ (fibroblast) cells within the lesion at 7 dpl and no Col1$^+$ cells at 56 dpl (data not shown).

Next, we quantified the overall gliotic (microglial and astrocytic) response by determining the

density of nonoligodendrocyte lineage cell population (nonOligo, Olig2$^-$DAPI$^+$) within the lesion at

each time point. The density of nonOligos increased 10-fold following injury, with densities

peaking at 14 dpl (NAWM (n = 20) vs 14dpl (n = 7), Tukey's p <0.0001) (**Fig. 2F**). While there

was a statistically significant decrease in NonOligo cell density between 14 and 56 dpl (14dpl (n

= 7) vs 56dpl (n = 5), Tukey's p < 0.0001), the percentage of nonOligos remained constant (14dpl

(n = 7) vs 56dpl (n = 5), Tukey's p = 0.64) (**Fig 2F-G**). However, there was a statistically significant

negative trend from 7dpl to 56dpl in the percentage of nonOligos (Test for linear trend, 7dpl to

56dpl, slope = -2.7, F (1, 17) = 8.70, p = 0.009). Together, this suggests that the gliotic response

is persistent and remains active for at least the first two months following demyelination.

Demyelinated areas show insufficient OPC recruitment and failed to restore normal

oligodendrocyte cell density.

As recruitment varies greatly between experimental models and MS lesions, we next sought to

characterize OPC recruitment and differentiation following demyelination in this new model. We

found that total oligodendrocyte lineage (Olig2$^+$) cell density was decreased relative to NAWM at

all timepoints and remained so even at 56 dpl (NAWM (n = 22) vs 56dpl (n = 5), Tukey's p =

0.003) (**Fig 4A,4E**). There were no statistically significant differences between timepoints in Olig2$^+$

cell density (7-56dpl) (One-way ANOVA, F (3, 18) = 0.98, p = 0.42). Importantly, the density of

OPCs and immature oligodendrocytes defined as CC1$^-$Olig2$^+$ (**Fig 4D**) was maintained, showing

no statistically significant differences between groups (One-way ANOVA, F (4, 28) = 0.88, p =

0.49) (**Fig 4F**), a similar observation to that of MS lesion tissue (**Fig 1A**). However, the proportion

of OPCs increased significantly at 7dpl (7 dpl (n = 4) vs NAWM (n = 16), Tukey's p <0.0001), but

returned to that of NAWM by 56 dpl (n = 5; Tukey's p =0.79) (**Fig 4G**). Sox2 expression is

upregulated by Olig2-defined OPCs following demyelination and corresponds with OPC

activation, acting to maintain these cells in a proliferative state and priming them for differentiation

[4]. In rodents, OPC activation peaked at 5 dpl [4]. Comparatively, Sox2$^+$Olig2$^+$ activated OPCs in

the rabbit peaked at 14 dpl (Tukey's multiple comparisons test, NAWM (n = 19) vs 14dpl (n = 4),

p = <0.0001), but this peak did not correspond with an increase in total OPC density at 14, 21 or

56 dpl (**Fig 4B-4C**). Likewise, the density of Ki67$^+$ proliferating Olig2$^+$ OPCs peaked at 14 dpl and

declined thereafter, returning to NAWM levels by 56 dpl (**Fig 4J-4K**).

In rodent models, the loss of Sox2 in OPCs and exit from cell cycle correlates with a

recovery of mature OL cell density [4]. We analyzed oligodendrogenesis using CC1$^+$Olig2$^+$ at each

time point. Unlike the rodent, the density of mature OLs did not return to NAWM levels. Such that

at 56 dpl, the density of post-mitotic OLs remained less than half that of oligodendrocytes in

NAWM (Tukey's p = 0.0013). However, we noted a significant increase in OL density between 7

and 56 dpl (test for linear trend, 7dpl to 56 dpl, F (1, 13) = 4.8, p = 0.047) (**Fig 4D, 4L**). Intriguingly,

while the overall density of mature OLs was dramatically reduced in the rabbit, the percentage of

mature OLs returned to that of NAWM by 56 dpl (NAWM (n = 16) vs 56dpl (n = 5), Tukey's p =

0.79) (**Fig 4M**). These results indicate that while rabbit OPCs remain capable of

activation/proliferation and subsequent oligodendrocyte differentiation, the number of OPC

recruited following demyelination is insufficient to restore the density of OLs to NAWM levels.

Compared to rodent models, rabbit lesions exhibited inferior OPC recruitment and delayed OL generation.

We directly compared the dynamics of OPC recruitment and differentiation in the rabbit to analogous rodent focal models of gliotoxin-induced demyelination. The rabbit internal capsule (IC) contained higher density of Olig2$^+$ cells compared to mouse spinal cord (SC) (**Fig 5A-B**). Following demyelination, recruitment of Olig2$^+$ cells was far more efficient in mouse SC than that observed in rabbit IC (**Fig 5C-D**). To quantitatively compare OPC recruitment and differentiation across models, we compared our results to those of previous lysolecithin models of demyelination in the mouse [4, 25, 26] and ethidium bromide in the rat caudal cerebral peduncle (CCP)[4, 9, 27, 28]. Following induction of demyelination, both mouse and rat exhibit increased density of Olig2$^+$ following recruitment of OPCs at 5 dpl and thereafter (black and brown lines, **Fig 5E**). However, the relative density of Olig2$^+$ oligodendroglia in rabbit does not increase following demyelination and is much reduced compared to mouse and rat (gray line, **Fig 5E-5F**). Similarly, while the density of OPCs in NAWM is greater in the rabbit compared to rat and mouse (**Fig 5G**), the spike in relative OPC density that corresponds with successful recruitment after injury is absent in the rabbit (**Fig 5G-5H**). Both mouse and rat display between two and ten-fold increase overall Olig2 and OPC density relative to NAWM. In contrast, the rabbit OPC response is muted and delayed relative to the other species. Intriguingly, rabbit lesions also display slower OPC differentiation (both absolute and relative) compared to mouse and rat (**Fig 5A-5D, 5I-5J**). Thus, abundant OPC recruitment seems to be a characteristic of the remyelination-efficient mouse and rat models, while less-abundant OPC recruitment is a shared characteristic between the rabbit model and observations of MS lesions (**Supplemental Fig 1A vs Fig 5H**).

Regional differences in axonal injury correspond to chronic gliosis and prolonged OPC activation.

Unlike rodent models with relatively small cross-sectional area, we observed clearly distinct sub-regions within the rabbit lesion. A common feature of rabbit IC lesions was a central core region that displayed a reduced density of axons relative to the rest of the lesion (**Fig 3**). This appears to replicate a common feature of MS lesions which are often found to possess lower axonal density in a central core region [23]. We defined the remainder of the lesion as lesion periphery (**Fig 6A-C**). The area of the core region remained stable even at 56 dpl while the area of the lesion periphery significantly reduced with time (**Fig 6G-H**), which could be due to resolution of inflammation, tissue edema, and remyelination.

At 14 dpl, the lesion core possessed high numbers of both Iba1$^+$ microglia and Gfap$^+$
astrocytes (**Fig 6D-F**). At this point, the astrocytic and microglial responses were qualitatively
similar in the lesion core and periphery. In contrast, at 56 dpl, hypertrophic and reactive astrocytes
were highly enriched around the lesion core, often forming a distinct border indicative of glial scar
formation (**Fig 2C**). The density of microglia/macrophages (Iba1$^+$) was greatest at earlier
timepoints and was somewhat reduced in the periphery at later timepoints (**Fig 6D-F, 2D**).
Quantification of the density of NonOligo cells reflected these observations indicating that more
cell nuclei were present in the core at each stage following demyelination and that cell density
gradually decreased following 14 dpl (Two-way ANOVA, location factor, $F_{(1, 38)} = 27.47$, p
<0.0001) (**Fig 6I**).
We analyzed the relative recruitment and differentiation of OPCs in both lesion core and periphery
(**Fig 7**). The overall Olig2$^+$ cell density was significantly greater in the lesion periphery then the
core regardless of time (Two-way ANOVA, region main effect: $F_{(1, 41)} = 9.30$, $p = 0.004$) (**Fig
7E**). This difference was seemingly not due to absolute OPC numbers as there was no statistically
significant difference between CC1$^-$ OPC and immature OL density of the core and periphery
across time ($F_{(1, 32)} = 3.01$, $p = 0.85$) (**Fig 7F**). Furthermore, the density of Sox2$^+$ activated
OPCs (**Fig 7H**) or Ki67$^+$ proliferating OPCs (**Fig 7J**) did not differ significantly between regions.
However, the proportion of OPCs among the Olig2 fraction consistently exceeded that of NAWM
and was greatest in the lesion core ($F_{(1, 25)} = 8.484$, $p = 0.008$)(**Fig 7G, I, K**). At 56 dpl, Sox2$^+$
OPCs remained highly abundant representing over half of all OL lineage cells in the core, while
the proportion of activated OPCs in periphery had reduced to NAWM levels ($F_{(1, 32)} = 33.24$, p
<0.0001). Likewise, while OPC proliferation peaked at 14 dpl in the lesion periphery and decline
thereafter, in the lesion core the rate of OPC proliferation continued to rise until 21 dpl and
remained substantially higher than NAWM even at 56 dpl ($F_{(1, 31)} = 9.754$, $p = 0.004$). In
contrast, the density of CC1$^+$Olig2$^+$ OL was significantly higher in the lesion periphery across time
points ($F_{(3, 28)} = 3.379$, $p = 0.032$) (**Fig 7L**). The proportion of OLs dramatically increased from
7 to 21 dpl in both lesion core and periphery but also trended toward an overall increase in the
lesion periphery ($p = 0.058$) (**Fig 7M**). Together, these results suggest that the core environment
could be driving OPCs to a state of chronic activation and proliferation following demyelination
while the lesion periphery supported OL differentiation after an initial period of activation and
proliferation.

Decreased OL and preserved OPC density in the NAWM directly adjacent to the lesion.

As the larger rabbit lesion afforded a closer examination of regional differences in OPC response

to demyelination, we examined the density and proportion of OL lineage cells in and around the

border of demyelination. The lesion border was defined by reduced solochrome staining

compared to NAWM (**Fig 8A-B**). A region extending 100 µm outside the lesion border was defined

as perilesion, while the analogous region within the lesion was classified as the lesion edge (**Fig**

8C). While the perilesion region was not hypercellular, with DAPI density not exceeding NAWM

at 7 dpl, the lesion edge exhibited clear hypercellularity that was reduced compared to lesion

periphery and core (**Fig 8D, 8I**). Furthermore, the perilesion did not exhibit detectable

demyelination as we confirmed that both MBP and FluoroMyelin were not reduced in this region

(data not shown). The remaining lesion was divided into inner periphery and lesion core as

described above. Intriguingly, the different regions exhibited significantly distinct temporal

increases in cellular density (p = <0.0001), with the greatest density of cells found in the lesion

core at between 14 and 21 dpl (**Fig 8D**). Also corresponding with these demarcations, we

observed a distinct pattern of OPC and OL lineage fate between each region (**Fig 8E, 8H**). The

density of CC1⁻Olig2⁺ OPC and immature OL cells returned to normal density by 7 dpl within the

lesion but was substantially lower in the perilesion region than the NAWM (**Fig 8J**). This suggests

that a depletion of OPCs occurs around the site of the lesion as cells are recruited. The density

of activated Sox2⁺ OPCs peaked at 14 dpl across regions and was greatest in the central lesion

regions (core and inner periphery) (**Fig 8F, 8K**). While the lesion core possessed the greatest

proportion of Sox2⁺ OPCs exceeding 50% of Olig2⁺ cells, up to 25% of OL lineage cells expressed

Sox2 in the perilesion region (p = 0.005) (**Fig 8F, Fig 8L**). This suggests that OPCs outside the

region of demyelination upregulate Sox2 in response to nearby demyelination. Intriguingly, the

perilesion exhibited a substantial loss of Olig2⁺CC1⁺ OL. In contrast, the perilesion region in

mouse spinal cord lesions, which was defined in a similar manner based on the lack of

hypercellularity, did not demonstrate OL loss [29]. As expected, OL loss was apparent within the

bulk of the rabbit lesion and the relatively small increase in OL density did not differ significantly

between regions (p = 0.012) (**Fig 8M-8N**).

DISCUSSION

The inadequacies of current rodent models of demyelination used to study and develop
remyelination-inducing therapies could contribute to their lack of successful translation into the
clinic. Here, we show that we are able to produce lesions which can more closely approximate
the size of lesions and several key pathological features found in MS patients. Among these,
rabbit lesions display insufficient oligodendrocyte progenitor cell (OPC) recruitment where OPC
density never exceeds that of NAWM. In stark contrast, regardless of the method of demyelination
or species used, small rodent models of demyelination universally display OPC densities between
two and ten times above that of NAWM shortly following demyelination. Large volume rabbit
lesions were more persistent than those produced in small rodents, with areas of demyelination
visible for at least two months. This is consistent with previous observations in the rabbit which
noted scant remyelination at two weeks post injection of lysolecithin and chronic demyelination
persisting for at least six months [30-32]. Additionally, we found region-specific differences in the
response of OPCs to demyelination, notably chronic activation of OPCs in the core and relatively
higher OL generation in the periphery.

In small rodent models, OPC recruitment is not considered rate-limiting for remyelination
following lysolecithin-induced demyelination [13, 33]. OL differentiation is dependent on the local
density of OPCs both *in vitro* [34] and following transplantation into hypomyelinated *shiverer* mice
[35]. This suggests that below a certain threshold local density of OPCs, the process of OL
differentiation will be limited. Consist with this hypothesis, remyelinated (RM) lesions and active
lesions in MS, which have a higher propensity for remyelination, tend to have higher densities of
OPCs then chronic demyelinated lesions [36]. Unlike rodent models, the large volume rabbit lesion
is associated with inefficient and slow OPC recruitment. Together with the huge difference in
volume between rabbit and mouse, this suggests that recruitment may become rate-limiting with
increasing lesion volume and that improving recruitment could be a vital and previously
underappreciated step in remyelination therapeutics.

While we have observed insufficient OPC recruitment in the setting of larger lesion
production, we cannot entirely rule out whether failed OPC recruitment is due to volume or
species-dependent differences. There are several well described differences in the behavior of
human and mouse OPCs both *in vitro* and following transplantation. For example, adult mouse
OPCs are more than twice as migratory compared to adult human OPCs [37]. Developmentally,
there are vast differences in the volume and proportion of myelination required in the human
compared to rodents, with developmental myelination completing over years in the human but
weeks in rodents [35]. These differences are also reflected at a cell autonomous level, as, following

transplantation human OPCs take 8-12 weeks to generate myelin-forming OLs [38], while equivalent

rodent cells typically complete the process of differentiation in 3-4 weeks [39]. In contrast, the

majority of signaling cascades and transcriptional networks appear to be largely conserved

between primary human and mouse OPCs [40-43]. Regardless of the contribution of OPC species-

differences, the failure of OPC recruitment in the rabbit recapitulates that observed in the majority

of MS lesions and thereby provides a suitable test bed for therapeutic approaches to improve the

migration and proliferation of OPCs.

Several other causes of remyelination failure have been proposed. Among these, age is

a critical factor determining the success for a number of regenerative processes [44]. While young

adult animals efficiently remyelinate following focal demyelination and demonstrate widespread

OPC recruitment, aged adult animals display substantially impaired rate of remyelination [27]. In

aged adult rodent, OPC recruitment and differentiation are delayed and intriguingly OPC

recruitment exhibits a gradient with greatest density in the lesion periphery vs central core [27].

Future studies will need to investigate the role of aging in the rabbit model to address whether

aging similarly influences both recruitment and differentiation in large volume lesions. In addition

to age, the innate immune response and extent of axonal injury are critical components that

determining the success or failure or remyelination.

Axonal degeneration and transection in MS occurs in the context of inflammatory

processes associated with (1) primary demyelination, (2) following the loss of trophic support by

myelinating OL and, in some cases, (3) independently of overt demyelination [45]. While loss of

trophic support by myelinating OLs results in axonal injury [46], it is likely that axonal damage in

active lesions is due to the intense proinflammatory and toxic microenvironment [17, 47]. Interestingly

in experimental models, axonal damage peaks at 6 days following induction of demyelination and

decreases substantially thereafter, corresponding to the differentiation of OPCs into myelinating

OLs [48]. Further supporting the association of axonal injury and myelin regeneration, induction of

OL differentiation and remyelination following ablation of the muscarinic receptor in OPCs

prevents axonal loss in EAE [49]. In contrast, an increase in pro-inflammatory signaling such as

IFN-γ administered at the time of demyelination prevents efficient OPC recruitment and

exacerbates axonal injury [50]. Likewise, in the rabbit model we observed axonal loss and injury

marked by App1 following demyelination at 7 dpl. Axonal loss was greatest in the lesion core and

the distribution of axonal loss in rabbit lesions was similar to that observed in MS [23]. Axonal loss

in the rabbit was largely associated with the initial gliotoxin insult as the size of core region

remained stable out to 56 dpl and we noted a substantial reduction in App1 staining after 7 dpl.

Interestingly, the microenvironment of the core region likely influences OPC state. Within the core

we observed many more OPCs that remained activated and proliferative for at least two months

following demyelination. Prolonged activation of OPCs within the core correlated with increased

microglial density suggesting that regional differences in microglial signaling alter OPC state.

Recent studies have increasingly highlighted the role that the innate immune response

plays in influencing remyelination [51]. OPC recruitment and differentiation is dependent on the

polarization of microglia/macrophages, whereby a proinflammatory so-called M1 state promotes

OPC recruitment while an immunoregulatory M2 state promotes OPC differentiation and

remyelination [52]. The transition of microglia/macrophages to M2 polarization can be blocked by

preventing necroptosis of M1 microglia and blocking this transition impairs remyelination of

demyelinated cerebellar explants [53]. While type 1 interferon may recruit microglia into regions of

demyelination, type 2 interferon (IFN-γ) produced by Th1 cells is associated with and found at

high levels during active lesion formation in MS [54]. Interestingly, IFN-γ directly drives OPC

quiescence via upregulation of PRRX1 in human OPCs [55]. This can be reversed by modulation

of pharmacological modulation of the local heparanome within regions of demyelination rescuing

both remyelination and axonal injury [50]. It is interesting to speculate that in large volume lesions,

the transition of the microglia/macrophage response from proinflammatory to immunomodulatory

phenotype occurs before the lesion achieves a sufficient density of OPC and thereby instead of

promoting remyelination acts instead to limit OPC recruitment resulting in an overall failure of

myelin regeneration. Whether or not this occurs in the complex inflammatory milieu of MS lesions

will need to be determined.

Remyelination in gliotoxin models provides a simplified model to study the cellular

mechanisms of myelin regeneration. While several molecular mechanisms have been described

that regulate and are rate-limiting for the repair of mouse and rat lesions, similar studies have not

yet been performed in large volume lesions. In the present study, we have shown that aberrations

in OPC recruitment and differentiation, akin to those observed in MS, occur following the induction

of white matter lesions in the rabbit. These lesions were over 20-fold larger than those produced

in mice and exhibit axon injury/loss as well as prolong gliosis that correspond with similar active

lesions in MS lesions. OPC recruitment is profoundly different in the rabbit model compared to

small rodents as the density of OPCs never exceeds that of NAWM. This corresponds well with

observations of OPC density in human MS lesions. Unlike other large species (cat, dog, sheep or

pig) with larger volumes of available white matter, rabbits are cost-effective and stereotaxic

techniques are routinely accessible in most neuroscience laboratories. As such, we propose that

the simplicity and accessibly of the rabbit model along with the clear differences in lesion

environment and patterns of OPC recruitment and differentiation will provide a vital

complementary approach for preclinical testing of remyelination therapeutics and aid in the

successful development of clinical interventions aimed at promoting myelin regeneration.

REFERENCES

1.   Confavreux C, Vukusic S, Moreau T, Adeleine P. Relapses and progression of disability in multiple sclerosis. *The New England journal of medicine*. Nov 16 2000;343(20):1430-8. doi:10.1056/NEJM200011163432001

2.   Franklin RJM, Ffrench-Constant C. Regenerating CNS myelin - from mechanisms to experimental medicines. *Nat Rev Neurosci*. Nov 16 2017;18(12):753-769. doi:10.1038/nrn.2017.136

3.   Hughes EG, Kang SH, Fukaya M, Bergles DE. Oligodendrocyte progenitors balance growth with self-repulsion to achieve homeostasis in the adult brain. *Nature neuroscience*. Jun 2013;16(6):668-76. doi:10.1038/nn.3390

4.   Zhao C, Ma D, Zawadzka M, et al. Sox2 Sustains Recruitment of Oligodendrocyte Progenitor Cells following CNS Demyelination and Primes Them for Differentiation during Remyelination. *J Neurosci*. Aug 19 2015;35(33):11482-99. doi:10.1523/JNEUROSCI.3655-14.2015

5.   Lubetzki C, Sol-Foulon N, Desmazieres A. Nodes of Ranvier during development and repair in the CNS. *Nat Rev Neurol*. Aug 2020;16(8):426-439. doi:10.1038/s41582-020-0375-x

6.   Wolswijk G. Chronic stage multiple sclerosis lesions contain a relatively quiescent population of oligodendrocyte precursor cells. *J Neurosci*. Jan 15 1998;18(2):601-9.

7.   Mi S, Miller RH, Lee X, et al. LINGO-1 negatively regulates myelination by oligodendrocytes. *Nature neuroscience*. Jun 2005;8(6):745-51. doi:10.1038/nn1460

8.   Zhang M, Zhan XL, Ma ZY, Chen XS, Cai QY, Yao ZX. Thyroid hormone alleviates demyelination induced by cuprizone through its role in remyelination during the remission period. *Exp Biol Med (Maywood)*. Sep 2015;240(9):1183-96. doi:10.1177/1535370214565975

9.   Neumann B, Baror R, Zhao C, et al. Metformin Restores CNS Remyelination Capacity by Rejuvenating Aged Stem Cells. *Cell Stem Cell*. Oct 3 2019;25(4):473-485 e8. doi:10.1016/j.stem.2019.08.015

10.   Green AJ, Gelfand JM, Cree BA, et al. Clemastine fumarate as a remyelinating therapy for multiple sclerosis (ReBUILD): a randomised, controlled, double-blind, crossover trial. *Lancet*. Dec 2 2017;390(10111):2481-2489. doi:10.1016/S0140-6736(17)32346-2

11.   Mellion M, Edwards KR, Hupperts R, et al. Efficacy Results from the Phase 2b SYNERGY Study: Treatment of Disabling Multiple Sclerosis with the Anti-LINGO-1 Monoclonal Antibody Opicinumab. *Neurology*. Apr 18 2017;88

12.   Klistorner A, Chai Y, Leocani L, et al. Assessment of Opicinumab in Acute Optic Neuritis

Using Multifocal Visual Evoked Potential. *CNS Drugs*. Dec 2018;32(12):1159-1171.

doi:10.1007/s40263-018-0575-8

13.   Woodruff RH, Fruttiger M, Richardson WD, Franklin RJ. Platelet-derived growth factor

regulates oligodendrocyte progenitor numbers in adult CNS and their response following CNS

demyelination. *Mol Cell Neurosci*. Feb 2004;25(2):252-62. doi:10.1016/j.mcn.2003.10.014

14.   Foerster S, Neumann B, McClain C, et al. 2020;doi:10.1101/2020.05.21.108373

15.   Xing YL, Chuang BHA, Poh J, et al. 2021;doi:10.1101/2021.05.13.443012

16.   Mason JL, Toews A, Hostettler JD, et al. Oligodendrocytes and progenitors become

progressively depleted within chronically demyelinated lesions. *The American journal of*

pathology. May 2004;164(5):1673-82. doi:10.1016/S0002-9440(10)63726-1

17.   Trapp BD, Peterson J, Ransohoff RM, Rudick R, Mork S, Bo L. Axonal transection in the

lesions of multiple sclerosis. *The New England journal of medicine*. Jan 29 1998;338(5):278-85.

doi:10.1056/NEJM199801293380502

18.   Jean I, Lavialle C, Barthelaix-Pouplard A, Fressinaud C. Neurotrophin-3 specifically

increases mature oligodendrocyte population and enhances remyelination after chemical

demyelination of adult rat CNS. *Brain Res*. May 16 2003;972(1-2):110-8.

19.   Rusielewicz T, Nam J, Damanakis E, John GR, Raine CS, Melendez-Vasquez CV.

Accelerated repair of demyelinated CNS lesions in the absence of non-muscle myosin IIB. *Glia*.

Apr 2014;62(4):580-91. doi:10.1002/glia.22627

20.   Shields SA, Gilson JM, Blakemore WF, Franklin RJ. Remyelination occurs as extensively

but more slowly in old rats compared to young rats following gliotoxin-induced CNS demyelination.

Glia. Oct 1999;28(1):77-83.

21.   Arnett HA, Fancy SP, Alberta JA, et al. bHLH transcription factor Olig1 is required to repair

demyelinated lesions in the CNS. *Science*. Dec 17 2004;306(5704):2111-5.

doi:10.1126/science.1103709

22.   Zhang K, Sejnowski TJ. A universal scaling law between gray matter and white matter of

cerebral cortex. *Proc Natl Acad Sci U S A*. May 9 2000;97(10):5621-6.

doi:10.1073/pnas.090504197

23.   Ferguson B, Matyszak MK, Esiri MM, Perry VH. Axonal damage in acute multiple sclerosis

lesions. *Brain*. Mar 1997;120 (Pt 3):393-9. doi:10.1093/brain/120.3.393

24.   Dorrier CE, Aran D, Haenelt EA, et al. CNS fibroblasts form a fibrotic scar in response to

immune cell infiltration. *Nature neuroscience*. Feb 2021;24(2):234-244. doi:10.1038/s41593-020-

00770-9

25. Kucharova K, Stallcup WB. NG2-proteoglycan-dependent contributions of
oligodendrocyte progenitors and myeloid cells to myelin damage and repair. *J Neuroinflammation*.
Sep 4 2015;12:161. doi:10.1186/s12974-015-0385-6

26. Fancy SP, Baranzini SE, Zhao C, et al. Dysregulation of the Wnt pathway inhibits timely
myelination and remyelination in the mammalian CNS. *Genes Dev*. Jul 1 2009;23(13):1571-85.
doi:10.1101/gad.1806309

27. Sim FJ, Zhao C, Penderis J, Franklin RJ. The age-related decrease in CNS remyelination
efficiency is attributable to an impairment of both oligodendrocyte progenitor recruitment and
differentiation. *J Neurosci*. Apr 1 2002;22(7):2451-9. doi:20026217

28. Syed YA, Hand E, Mobius W, et al. Inhibition of CNS remyelination by the presence of
semaphorin 3A. *J Neurosci*. Mar 9 2011;31(10):3719-28. doi:10.1523/JNEUROSCI.4930-10.2011

29. Crawford AH, Tripathi RB, Foerster S, et al. Pre-Existing Mature Oligodendrocytes Do Not
Contribute to Remyelination following Toxin-Induced Spinal Cord Demyelination. *Am J Pathol*.
Mar 2016;186(3):511-6. doi:10.1016/j.ajpath.2015.11.005

30. Blakemore WF. Observations on remyelination in the rabbit spinal cord following
demyelination induced by lysolecithin. *Neuropathology and applied neurobiology*. Jan-Feb
1978;4(1):47-59. doi:10.1111/j.1365-2990.1978.tb00528.x

31. Waxman SG, Kocsis JD, Nitta KC. Lysophosphatidyl choline-induced focal demyelination
in the rabbit corpus callosum. Light-microscopic observations. *J Neurol Sci*. Dec 1979;44(1)(45-
53):45-53. doi:10.1016/0022-510x(79)90221-1

32. Foster RE, Kocsis JD, Malenka RC, Waxman SG. Lysophosphatidyl choline-induced focal
demyelination in the rabbit corpus callosum. Electron-microscopic observations. *J Neurol Sci*. Nov
1980;48(2):221-31. doi:10.1016/0022-510x(80)90202-6

33. Woodruff RH, Franklin RJ. Demyelination and remyelination of the caudal cerebellar
peduncle of adult rats following stereotaxic injections of lysolecithin, ethidium bromide, and
complement/anti-galactocerebroside: a comparative study. *Glia*. Feb 1 1999;25(3):216-28.
doi:10.1002/(SICI)1098-1136(19990201)25:3<216::AID-GLIA2>3.0.CO;2-L [pii]

34. Rosenberg SS, Kelland EE, Tokar E, De la Torre AR, Chan JR. The geometric and spatial
constraints of the microenvironment induce oligodendrocyte differentiation. *Proc Natl Acad Sci U
S A*. Sep 23 2008;105(38):14662-7. doi:10.1073/pnas.0805640105

35. Dietz KC, Polanco JJ, Pol SU, Sim FJ. Targeting human oligodendrocyte progenitors for
myelin repair. *Exp Neurol*. Sep 2016;283(Pt B):489-500. doi:10.1016/j.expneurol.2016.03.017

36. Boyd A, Zhang H, Williams A. Insufficient OPC migration into demyelinated lesions is a
cause of poor remyelination in MS and mouse models. *Acta Neuropathol*. Jun 2013;125(6):841-
59. doi:10.1007/s00401-013-1112-y
37. Bribian A, Medina-Rodriguez EM, Josa-Prado F, et al. Functional Heterogeneity of Mouse
and Human Brain OPCs: Relevance for Preclinical Studies in Multiple Sclerosis. *J Clin Med*. Jun
2 2020;9(6)doi:10.3390/jcm9061681
38. Sim FJ, McClain CR, Schanz SJ, Protack TL, Windrem MS, Goldman SA. CD140a
identifies a population of highly myelinogenic, migration-competent and efficiently engrafting
human oligodendrocyte progenitor cells. *Nat Biotechnol*. Sep 25 2011;29(10):934-41.
doi:10.1038/nbt.1972
39. Baron-Van Evercooren A, Blakemore WF. Chapter 6 - Remyelination through
Engraftment. In: Lazzarini RA, Griffin JW, Lassman H, Nave K-A, Miller R, Trapp BD, eds. *Myelin*
Biology and Disorders. Academic Press; 2004:143-172.
40. Conway GD, O'Bara MA, Vedia BH, Pol SU, Sim FJ. Histone deacetylase activity is
required for human oligodendrocyte progenitor differentiation. *Glia*. Dec 2012;60(12):1944-53.
doi:10.1002/glia.22410
41. Abiraman K, Pol SU, O'Bara MA, et al. Anti-muscarinic adjunct therapy accelerates
functional human oligodendrocyte repair. *J Neurosci*. Feb 25 2015;35(8):3676-88.
doi:10.1523/JNEUROSCI.3510-14.2015
42. Pol SU, Polanco JJ, Seidman RA, et al. Network-Based Genomic Analysis of Human
Oligodendrocyte Progenitor Differentiation. *Stem Cell Reports*. Aug 8 2017;9(2):710-723.
doi:10.1016/j.stemcr.2017.07.007
43. Saraswat D, Shayya HJ, Polanco JJ, et al. Overcoming the inhibitory microenvironment
surrounding oligodendrocyte progenitor cells following experimental demyelination. *Nat Commun*.
Mar 26 2021;12(1):1923. doi:10.1038/s41467-021-22263-4
44. Ruckh JM, Zhao JW, Shadrach JL, et al. Rejuvenation of regeneration in the aging central
nervous system. *Cell Stem Cell*. Jan 6 2012;10(1):96-103. doi:10.1016/j.stem.2011.11.019
45. Haines JD, Inglese M, Casaccia P. Axonal damage in multiple sclerosis. *Mt Sinai J Med*.
Mar-Apr 2011;78(2):231-43. doi:10.1002/msj.20246
46. Nave KA. Myelination and the trophic support of long axons. *Nature reviews*
Neuroscience. Apr 2010;11(4):275-83. doi:10.1038/nrn2797
47. Kornek B, Storch MK, Weissert R, et al. Multiple sclerosis and chronic autoimmune
encephalomyelitis: a comparative quantitative study of axonal injury in active, inactive, and

remyelinated lesions. *The American journal of pathology*. Jul 2000;157(1):267-76.
doi:10.1016/S0002-9440(10)64537-3

48. Schultz V, van der Meer F, Wrzos C, et al. Acutely damaged axons are remyelinated in
multiple sclerosis and experimental models of demyelination. *Glia*. Aug 2017;65(8):1350-1360.
doi:10.1002/glia.23167

49. Mei F, Lehmann-Horn K, Shen YA, et al. Accelerated remyelination during inflammatory
demyelination prevents axonal loss and improves functional recovery. *Elife*. Sep 27
2016;5:e18246. doi:10.7554/eLife.18246

50. Saraswat D, Welliver RR, Ravichandar R, et al. Heparanome-Mediated Rescue of
Oligodendrocyte Progenitor Quiescence following Inflammatory Demyelination. *J Neurosci*. Mar
10 2021;41(10):2245-2263. doi:10.1523/JNEUROSCI.0580-20.2021

51. Lloyd AF, Miron VE. The pro-remyelination properties of microglia in the central nervous
system. Review. *Nat Rev Neurol*. Aug 2019;15(8):447-458. doi:10.1038/s41582-019-0184-2

52. Miron VE, Boyd A, Zhao JW, et al. M2 microglia and macrophages drive oligodendrocyte
differentiation during CNS remyelination. *Nat Neurosci*. Sep 2013;16(9):1211-8.
doi:10.1038/nn.3469

53. Lloyd AF, Davies CL, Holloway RK, et al. Central nervous system regeneration is driven
by microglia necroptosis and repopulation. *Nat Neurosci*. Jul 2019;22(7):1046-1052.
doi:10.1038/s41593-019-0418-z

54. Panitch HS. Interferons in Multiple Sclerosis. *Drugs*. 1992/12/01 1992;44(6):946-962.
doi:10.2165/00003495-199244060-00004

55. Wang J, Saraswat D, Sinha AK, et al. Paired Related Homeobox Protein 1 Regulates
Quiescence in Human Oligodendrocyte Progenitors. *Cell reports*. Dec 18 2018;25(12):3435-3450
e6. doi:10.1016/j.celrep.2018.11.068

56. Rosen GD, Harry JD. Brain volume estimation from serial section measurements: a
comparison of methodologies. *J Neurosci Methods*. Nov 1990;35(2):115-24. doi:10.1016/0165-
0270(90)90101-k

57. Swanson LW. Brain maps 4.0-Structure of the rat brain: An open access atlas with global
nervous system nomenclature ontology and flatmaps. *J Comp Neurol*. Apr 15 2018;526(6):935-
943. doi:10.1002/cne.24381

58. Kuhlmann T, Miron V, Cui Q, Wegner C, Antel J, Bruck W. Differentiation block of
oligodendroglial progenitor cells as a cause for remyelination failure in chronic multiple sclerosis.
Brain. Jul 2008;131(Pt 7):1749-58. doi:10.1093/brain/awn096

59.   Chang A, Nishiyama A, Peterson J, Prineas J, Trapp BD. NG2-positive oligodendrocyte
progenitor cells in adult human brain and multiple sclerosis lesions. *J Neurosci*. Sep 1
2000;20(17):6404-12. doi:10.1523/jneurosci.20-17-06404.2000

60.   Yu Q, Hui R, Park J, et al. Strain differences in cuprizone induced demyelination. *Cell*
Biosci. 2017;7:59. doi:10.1186/s13578-017-0181-3

61.   Lin W, Kemper A, Dupree JL, Harding HP, Ron D, Popko B. Interferon-gamma inhibits
central nervous system remyelination through a process modulated by endoplasmic reticulum
stress. *Brain*. May 2006;129(Pt 5):1306-18. doi:10.1093/brain/awl044

62.   Zhang J, Zhang ZG, Li Y, et al. Fingolimod treatment promotes proliferation and
differentiation of oligodendrocyte progenitor cells in mice with experimental autoimmune
encephalomyelitis. *Neurobiol Dis*. Apr 2015;76:57-66. doi:10.1016/j.nbd.2015.01.006

63.   Garay L, Tungler V, Deniselle MC, Lima A, Roig P, De Nicola AF. Progesterone attenuates
demyelination and microglial reaction in the lysolecithin-injured spinal cord. *Neuroscience*. Sep
29 2011;192:588-97. doi:10.1016/j.neuroscience.2011.06.065

64.   Ulrich R, Seeliger F, Kreutzer M, Germann PG, Baumgartner W. Limited remyelination in
Theiler's murine encephalomyelitis due to insufficient oligodendroglial differentiation of nerve/glial
antigen 2 (NG2)-positive putative oligodendroglial progenitor cells. *Neuropathol Appl Neurobiol*.
Dec 2008;34(6):603-20. doi:10.1111/j.1365-2990.2008.00956.x

FIGURE LEGENDS

Figure 1. Large regions of demyelination following injection of lysolecithin into rabbit

subcortical white matter. (**A**) Diagram representing injection sites, and white matter tract size

for comparison with rat and mouse, drawn to scale (5 mm) (mouse, overlay of lab section; rat

overlay of rat brain atlas [57]). (**B**) Lysolecithin-induced lesion in the rabbit stained for FloroMyelin

(FM, red) and 4',6'-diamidino-2-phenylindole (DAPI, blue). White dotted line indicates lesion

border. (**C-D**) Maximal cross-sectional area (**C**) and lesion volume (**D**) of rabbit lesions at 14 dpl

(peak lesion volume, $n = 8$), compared to that seen in lysolecithin mouse models [18, 19]. (**E**) Maximal

cross-sectional area of lesions following lysolecithin injection in rabbit ($n = 6$-8) and rat [33] (dpl,

days post-lesion). Mean ± SEM shown. Scale: 500 μm.

Figure 2. Evolution of rabbit demyelinating lesions with preserved axon integrity and

prolonged gliotic response. (**A-D**) Representative confocal images of normal appearing white

matter (NAWM) stained for myelin, axonal, astrocytic and microglial/macrophage markers.

Following demyelination at 7 to 56 days post lesion (dpl), lesions were imaged by widefield

epifluorescence (outlined with dotted white line) shown in the left panel. Higher power confocal

images of the area outlined with the white box are shown in the right panel. (**A**) At 7 dpl, myelin

basic protein (Mbp, gray) staining demonstrated myelin loss and degeneration within the lesion,

and Mbp levels remain depressed over time. (**B**) Axons, stained with neurofilament (NF, green),

persist within the lesion bounds even at 56 dpl. Axonal injury (APP, red) is evident at 7dpl but is

reduced thereafter following demyelination. (**C-D**) Robust neuroinflammatory response,

astrocytes were stained with glial fibrillary acidic protein (Gfap, green) and amyloid microglial cells

were stained with ionized calcium binding adaptor molecule 1 (IBA1, red). (**E**) Total DAPI cell

density within the lesion was quantified ($n = 4$-19 per group, mean ± SEM). (**F-G**) The non-

oligodendrocyte lineage cells (nonOligos) was defined as DAPI$^+$Olig2$^-$ cells. Density (**F**) and

proportion of total cells (**G**) was quantified ($n = 4$-19). The density of cells increases dramatically

following lysolecithin injection and peaks at 14 days. This is largely due to an increase in the

density of non-oligodendrocyte lineage cells which likely represents microglia and at later time-

points astrocytic influx and proliferation (**G**). Lesion image scale: 500 μm. Confocal scale: 25 μm.

Figure 3 (related to Figure 3). Lesions were composed of an axon deficient core, with

higher axon preservation in the periphery. Representative 60x oil immersion objective +

constrained iterative deconvolution stitched images (0.8 μm steps over 4μm depth) of NAWM (**A**)

or following lysolecithin injection (**B-E**) stained for neurofilament (NF, green) and/or DAPI (blue).

White dotted lines indicate lesion borders and green dotted lines represents lesion cores.

Retraction bulbs were observed in and around the lesions at all timepoints (small frame images

found in **B-E**). Scale bar: 100um.

Figure 4. Insufficient OPC recruitment and oligodendrocyte generation following

demyelination in rabbit white matter. (**A**) Olig2 (red) was used to identify total oligodendrocyte

lineage cell density (NAWM, 14dpl, and 56 dpl shown). Insert shows high magnification of labelled

cells. Activated Sox2$^+$ OPCs (**B**, cyan), proliferative Ki67$^+$ OPCs (**C**, gray) and mature CC1$^+$

oligodendrocyte (**D**, green) were identified by colocalization with Olig2. Additionally, Sox2 has

been noted to be expressed by a subpopulation of astrocytes [4], which are identified as Sox2$^+$Olig2$^-$

cells. (**E**) Oligodendrocyte lineage cell density was significantly depressed relative to NAWM

following demyelination (*n* = 4-22, one-way ANOVA). The density and proportion of OPCs defined

as either Olig2$^+$CC1$^-$ (**F-G**), Sox2$^+$Olig2$^+$ (**H-I**) or Ki67$^+$Olig2$^+$ (**J-K**) increased following

demyelination and peaked between 7-14 dpl. While the proportion of mature oligodendrocytes

(Olig2$^+$CC1$^+$) (**M**) increased following demyelination, the overall density remained significantly

depressed relative to NAWM (**L**). Mean ± SEM. Tukey's multiple comparisons post-test vs.

NAWM. **, ***, and **** indicate p < 0.01, 0.001, and 0.0001, respectively. Large frame scale: 50

μm. Small frame scale; 20 μm.

Figure 5. Rabbit lesions show delayed recruitment of OPCs, abundant Sox2 activation, and

slower maturation of oligodendrocytes compared to rodent models. Representative images

of normal appearing white matter (NAWM) of mouse spinal cord (**A**) and rabbit internal capsule

(**B**). Mature oligodendrocytes and OPCs identified as Olig2$^+$CC1$^+$ and Olig2$^+$CC1$^-$, respectively.

Comparative images of lysolecithin induced lesions in mouse (**C**) and rabbit (**D**) at 14 dpl. White

dashed line indicates lesion border. *L* indicates lesion area when border is visible. (**E-J**)

Comparison of oligodendrocyte lineage cell response following demyelinating lesions between

rabbit (gray), mouse (black), and rat (brown). Density and relative change in density compared to

NAWM of that species. (**E-F**) While Olig2$^+$ cell density increases relative to NAWM following lesion

production in mouse [4] and rats [4], it remains depressed in rabbits. (**G-H**) Relative to NAWM, OPC

density in rat [27] (Pdgfra mRNA$^+$) and mouse [25] (NG2$^+$) models increases greatly following lesion

production. Conversely, OPC density in the rabbit (CC1$^-$Olig2$^+$) does not significantly change from

NAWM. (**I-J**) Mature OLs determined by Olig2$^+$PLP$^+$ (mouse [26], black), or Olig2$^+$CC1$^+$ (rat [9, 28],

rabbit). Rat mature OL density curve was determined from two separate studies, with NAWM

(original data: 326.4 ± 27.1 cells/mm^2) and 10dpl (original data: 102.2 ± 14.7 cells/mm^2) data from

Syed et al. 2011 normalized to the results of Neumann et al. 2019 (7, 21, 49 dpl). Mean ± SEM

shown. Scale bar: 100 µm.

Figure 6. Axonal deficient lesion core does not expand with time following demyelination.

Representative images following demyelination at 14 days post lesion (dpl), stained for axons

with neurofilament (NF, green), axonal injury with beta amyloid precursor protein (App, Red), and

nuclei with DAPI (blue)(**A**) or astrocytes (Gfap, green), and Iba1$^+$ microglia/macrophages (red)(**D**).

Lesion border was defined by hypercellularity and FluoroMyelin staining (blue dashed line) and

core was defined as a region containing sparse NF$^+$ intact axons (green dashed line). (**B-C, E-F**)

High power confocal images of the area outlined with the white boxes in **A** and **D**. (**G-H**) Core

lesion area did not significantly change through 56 dpl, while peripheral lesion area decreased

with time (n = 5-8 per group, two-way ANOVA). (**I**) Quantification of non-oligodendrocyte lineage

cells (DAPI$^+$Olig2$^-$ cells). Two-way ANOVA across time points comparing core and periphery

indicated a significant difference in the NonOligo cells between core and periphery (p < 0.0001),

and over time (p < 0.0001). The gliotic response was significantly greater in the core compared

to the periphery. Sidak's multiple comparisons comparing means of periphery and core at each

timepoint. *, **, ***, and **** indicate p < 0.05, 0.01, 0.001, and 0.0001, respectively. Mean ± SEM

shown. Lesion scale bar: 250 µm. Confocal scale bar: 25 µm.

Figure 7. OPCs in the axon-sparse core remained activated and proliferative. (**A-D**)

Representative images following demyelination at 14 days post lesion (dpl), stained for axons

with neurofilament (NF, green), activated Olig2$^+$Sox2$^+$ OPCs (**B**, cyan), proliferative Olig2$^+$Ki67$^+$

OPCs (**C**, gray) and mature Olig2$^+$CC1$^+$ oligodendrocytes (OL) (**D**, green). (**E**) Quantification of

Olig2$^+$ cell density in the core and periphery. Quantification of cell density and proportion of total

oligodendrocyte linage cells for OPCs (**F-G**), activated OPCs (**H-I**), proliferative OPCs (**J-K**), and

mature OLs (**L-M**). Dotted lines represent cellular density of NAWM. While few intact axons

remained, this did not significantly affect the recruitment of OPCs to the lesion core. However, a

significantly greater proportion of OPCs were activated and proliferating in the lesion core

compared to the periphery. Two-way ANOVA was performed across all time points (TME: time

main effect) and comparing core and periphery regions (RME: regional main effect). Sidak's post-

test comparing regions at each time point. *, **, ***, and **** indicate p < 0.05, 0.01, 0.001, and

0.0001, respectively (NAWM: n = 19-22, lesion: n = 4-7). Mean ± SEM reported. Scale bar: 200

µm.

Figure 8. Decreased density of mature OLs and no change in OPC density in NAWM directly
adjacent to demyelinating lesion. Representative images of NAWM (**A**) and lysolecithin
induced lesion at 56 dpl (**B-C**) in the rabbit stained with solochrome cyanine (myelin, black). Blue
dashed line indicates lesion border. PeriLesion region (purple) was defined as an area extending
~100 μm past the lesion border. Lesion edge (blue) was defined as an area ~100 μm smaller than
the lesion border. Inner periphery (red) was defined as the area between the outer periphery and
the core. Core defined as a central area of the lesion where few intact axons were observed.
Scale bar: 200 μm. Lesion cross-sectional columns covering the various regions stained for (**D**)
DAPI, (**E**) Olig2, (**F**) activated OPCs (Olig2$^+$Sox2$^+$), (**G**) proliferative OPCs (Olig2$^+$Ki67$^+$) (**H**) and
mature OLs (Olig2$^+$CC1$^+$). Scale bar, 500 μm. (**I**) Total DAPI$^+$ cell density was quantified in all
regions. Quantification of cell density and/or proportion of total oligodendrocyte linage cells
between regions for OPCs (Olig2$^+$CC1$^-$, **J**), Activated OPCs (Olig2$^+$Sox2$^+$, **K-L**), and mature OLs
(Olig2$^+$CC1$^+$, **M-N**). Dotted lines represent cellular densities of NAWM. Two-way ANOVA was
performed across all time points (TME: time main effect) and comparing regions (RME: regional
main effect). Mean ± SEM reported (NAWM: n = 19-22, lesion: n = 4-7).

Supplemental Figure 1. OPC density in MS lesions rarely exceeds that of NAWM, but is a
universal feature of rodent models. (**A**) In human MS lesions, OPCs were identified as either
Olig2$^+$Nogo$^-$ (green)[36], Olig2^{strong} (blue)[58] or NG2$^+$ cells (orange)[59]. Gray bar indicates the range of
observed OPC density from control human white matter (mean ± 1 SD, n = 5 patients)[36]. OPC
density rarely surpasses that of NAWM. RM = remyelinated lesions. In experimental
demyelination, relative OPC density in toxin (**B**) and immune-mediated models (**C**) was plotted
against time following the induction of demyelination (lysolecithin and ethidium bromide), or
appearance of demyelinated lesions (Cuprizone, EAE, Theiler's virus). OPCs were identified as
NG2$^+$ (Cuprizone [60], red; EAE [61, 62], purple; lysolecithin [25, 63], blue; Theiler's virus [64], orange), or
Pdgfra mRNA$^+$ (Rat EB [27], green), n = 3-5. Horizontal dotted line indicates the normalized OPC
density of the mouse. Robust OPC recruitment evidenced by increased OPC density relative to
normal white matter was a common feature of all rodent models, regardless of method of
demyelination. Rat EB = ethidium bromide induced lesions in the rat. EAE = experimental
autoimmune encephalomyelitis. Mean ± SEM shown.

Supplemental Figure 2 (related to Figure 3). Amoeboid microglia in lesion core form large
vesicles surrounded by astrocytic processes. (**A-B**) Confocal images of NAWM stained for
Iba1 (red), Gfap (green), and DAPI (blue). Cells at the center of the frame represent an astrocyte

(**A,** green arrowhead) or microglial cell (**B**, red arrowhead), respectively. (**C**) Orthogonal views of

an amoeboid microglial cell within the core of a lesion at 56 dpl. Scale bar: 25 µm.

Supplemental Figure 3 (related to Methods). Significant post-operative weight loss in the

rabbit, and depiction of injection site. (**A**) Dye injection (Evans Blue in PBS) showing injection

site (left hemisphere). Ruler, 1 mm between dashes. (**F**) Averaged weight of nonlesioned (blue)

and lesioned (gray) animals.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supplemental 1

Supplemental 3

Supplemental 2