Time-resolved parameterization of aperiodic and periodic brain activity

3 Luc E. Wilson[1], Jason da Silva Castanheira[1], Sylvain Baillet[1]†

4 [1] McConnell Brain Imaging Centre, Montreal Neurological Institute, McGill University,

5 Montreal, QC H3A 2B4, Canada

6 † Corresponding authors: sylvain.baillet@mcgill.ca.

Abstract

Macroscopic neural dynamics comprise both aperiodic and periodic signal components. Recent

advances in parameterizing neural power spectra offer practical tools for evaluating these features

separately. Although neural signals vary dynamically and express non-stationarity in relation to

ongoing behaviour and perception, current methods yield static spectral decompositions. Here, we

introduce Spectral Parameterization Resolved in Time (SPRiNT) as a novel method for

decomposing complex neural dynamics into periodic and aperiodic spectral elements in a time-

resolved manner. First, we demonstrate with naturalistic synthetic data SPRiNT's capacity to

reliably recover time-varying spectral features. We emphasize SPRiNT's specific strengths

compared to other time-frequency parameterization approaches based on wavelets. Second, we

use SPRiNT to illustrate how aperiodic spectral features fluctuate across time in empirical resting-

state electroencephalography data ($n = 178$), and relate the observed changes in aperiodic

parameters over time to participants' demographics and behaviour. Lastly, we use SPRiNT to

demonstrate how aperiodic dynamics relate to movement behaviour in intracranial recordings in

rodents. We foresee SPRiNT responding to growing neuroscientific interests in the

parameterization of time-varying neural power spectra and advancing the quantitation of complex

neural dynamics at the natural time scales of behaviour.

Keywords: aperiodic, periodic, time-resolved, parameterization, power spectra, neural

oscillations, electrophysiology.

Significance Statement

The new method and reported findings address a growing interest in neuroscience for

research tools that can reliably decompose brain activity at the mesoscopic scale into interpretable

components. We show that the new approach proposed is capable of tracking transient, dynamic

spectral (aperiodic and periodic) components across time, both in synthetic data and in *in vivo*

experimental data. We anticipate that this novel technique, SPRiNT, will enable new neuroscience

inquiries that reconcile multifaceted neural dynamics with complex behaviour.

Introduction

The brain constantly expresses a repertoire of complex dynamics related to behaviour in health

and disease. Neural oscillations, for instance, are rhythmic (periodic) components of brain activity

that emerge from a background of arrhythmic (aperiodic) fluctuations recorded with a range of

electrophysiological techniques at the mesoscopic scale (Buzsáki, 2006). Brain oscillations and

their rhythmic dynamics have been causally linked to individual behaviour and cognition (Albouy

et al., 2017), and shape brain responses to sensory stimulation (Samaha et al., 2020).

Current methods for measuring the time-varying properties of neural fluctuations include

several time-frequency decomposition techniques such as Hilbert, wavelet, and short-time Fourier

signal transforms (Bruns, 2004; Cohen, 2014), and more recently, empirical mode decompositions

(EMD; Huang et al., 1998) and time-delay embedded hidden Markov models (TDE-HMM; Quinn

et al., 2018). Following spectral decomposition, rhythmic activity within the empirical bands of

electrophysiology manifests as peaks of signal power (Buzsáki & Watson, 2012; Cohen, 2014).

However, time-resolved signal power decompositions (spectrograms) do not explicitly account for

the presence of aperiodic signal components, which challenge both the detection and the

interpretability of spectral peaks as genuine periodic signal elements (Donoghue et al., 2020). This

is critical as aperiodic and periodic components of neural activity represent distinct, although

possibly interdependent physiological mechanisms (Gao et al., 2017).

Aperiodic neural activity is characterized by a reciprocal distribution of power with

frequency (1/f), which can be parameterized with two scalars: *exponent* and *offset*. These

parameters are physiologically meaningful: current constructs consider the offset as reflecting

neuronal population spiking, and the exponent as related to the integration of synaptic currents

(Voytek & Knight, 2015) and reflecting the balance between excitatory (E) and inhibitory (I)

currents (i.e., the larger the exponent, the stronger the inhibition; Chini et al., 2021; Gao et al.,

2017; Waschke et al., 2021). Aperiodic neural activity is ubiquitous throughout the brain (He,

2014); it differentiates healthy aging (Cellier et al., 2021; Donoghue et al., 2020; Hill et al., 2022;

Ostlund et al., 2022; Schaworonkow & Voytek, 2021; Voytek et al., 2015), and is investigated as

a potential marker of neuropsychiatric conditions (Molina et al., 2020) and epilepsy (van Heumen

et al., 2021). Though the study of aperiodic neural activity has recently advanced, unanswered

questions remain about its functional relevance, which requires an expanded toolkit to track their

evolution across time and the broadest possible expressions of behaviour.

One little studied aspect of aperiodic activity is its fluctuations, both spontaneously over

time, and in association with task and mental states. Baseline normalization is a common approach

to compensate for aperiodic contributions to spectrograms (Cohen, 2014), with the underlying

assumption, however, that characteristics of aperiodic activity (exponent and offset) remain

unchanged throughout the data length—an assumption that is challenged by recent empirical data

that demonstrated their meaningful temporal fluctuations (Van Heumen et al., 2021; Waschke et

al., 2021). Akin to the motivations behind aperiodic/periodic spectral parameterization and signal

decomposition techniques (Donoghue et al., 2020, Wen & Liu, 2016), undetected temporal

variations within the neural spectrogram may conflate fluctuations in aperiodic activity with

modulations of periodic signals, hence distorting data interpretation (Donoghue et al., 2020).

Recent methodological advances have contributed practical tools to decompose and parameterize

the neural power spectrum (periodogram) into aperiodic and periodic components (Donoghue et

al., 2020; Wen & Liu, 2016; He, 2014). One such practical tool (*specparam*) sequentially fits

aperiodic and parametric components to the empirical neural power spectrum (Donoghue et al.,

2020). The resulting model for the aperiodic component is represented with exponent and offset

scalar parameters; periodic elements are modelled with a series of Gaussian-shape functions

characterized with three scalar parameters (centre frequency, amplitude, and standard deviation).

specparam accounts for static spectral representations and as such, does not account for the non-

stationary contents of neural time series.

We introduce SPRiNT (Spectral Parameterization Resolved in Time) as a novel approach

to identify and model dynamic shifts in aperiodic and periodic brain activity, yielding a time-

resolved parameterization of neurophysiological spectrograms. We validate the method with an

extensive set of naturalistic simulations of neural data, with benchmark comparisons to

parameterized wavelet signal decompositions. Using SPRiNT, we also show that aperiodic

fluctuations of the spectrogram can be related to meaningful behavioural and demographic

variables from human EEG resting-state data and electrophysiological recordings from free-

moving rodents.

Results

SPRiNT consists of the following methodological steps. First, short-time Fourier transforms

(STFT) are derived from overlapping time windows that slide over data time series. Second, the

resulting STFT coefficients are averaged over consecutive time windows to produce a smooth

estimate of the power spectral density of the recorded data. Third, the resulting periodogram is

parameterized into aperiodic and periodic components with *specparam* (see Methods). As the

procedure is repeated over the entire length of the data, SPRiNT produces a time-resolved

parameterization of the data's spectrogram (Figure 1A). The resulting parameters are then

compiled into fully parameterized time-frequency representations for visualization and further

derivations. A fourth step consists of an optional post-processing procedure meant to prune outlier

transient manifestations of periodic signal components (Figure 1 – figure supplement 1).

Figure 1: Methods Synopsis.

(**A**) Overview of the SPRiNT approach: At each time bin along a neurophysiological time series

(black trace) *n* overlapping time windows are Fourier-transformed to yield an estimate of

spectral contents, which is subsequently parameterized using *specparam* (Donoghue et al.,

2020). The procedure is replicated across time over sliding, overlapping windows to generate a

parameterized spectrogram of neural activity. (**B**) Simulation challenge I: We simulated 10,000

time series composed of the same time-varying spectral (aperiodic and periodic) features, with

different realizations of additive noise. (**C**) Simulation challenge II: We simulated another

10,000 time series, each composed of different time-varying spectral (aperiodic and periodic)

ground-truth features with additive noise. All simulated time series were used to evaluate the

respective performances of SPRiNT and the wavelet-*specparam* alternative.

Figure supplement 1. Overview of the outlier peak removal procedure.

We generated a total of 21,000 naturalistic synthetic neural time series comprising non-

stationary aperiodic and periodic signal components, using scripts adapted from the NeuroDSP

toolbox (Cole et al., 2019) with MATLAB (R2020a; Natick, MA). We first tested SPRiNT's

ability to detect and track transient and chirping periodic elements (with time-changing aperiodic

components) and benchmarked its performance against parameterized wavelet signal

decompositions and parameterized periodograms (Figure 1B). A second validation challenge

focused on simulations derived from randomly generated sets of realistic aperiodic and periodic

parameters; this challenge served to assess SPRiNT's performance across naturalistic

heterogenous signals (Figure 1C; see Methods). Further below, we describe the application of

SPRiNT to a variety of empirical data from human and rodent electrophysiology.

Methods benchmarking (synthetic data).

We first simulated 10,000 time series (60 s duration each) with aperiodic components transitioning

linearly between t=24 s and t=36 s, from an initial exponent of 1.5 Hz^{-1} and offset of -2.56

[arbitrary units, a.u.] towards a final exponent of 2.0 Hz^{-1} and offset of -1.41 a.u. The periodic

components of the simulated signals included transient activity in the alpha band (centre

frequency: 8 Hz; amplitude: 1.2 a.u.; standard deviation: 1.2 Hz) occurring between 8-40 s, 41-46

s, and 47-52 s, and a down-chirping oscillation in the beta band (centre frequency decreasing from

18 to 15 Hz; amplitude: 0.9 a.u.; standard deviation: 1.4 Hz, between 15-25 s (Figure 1B). We

applied SPRiNT on each simulated time series, post-processed the resulting parameter estimates

to remove outlier transient periodic components, and derived goodness-of-fit statistics of the

SPRiNT parameter estimates with respect to their ground-truth values. We compared SPRiNT's

performances to parameterized periodograms (*specparam)*, as well as the parameterization of

temporally smoothed spectrograms obtained from Morlet wavelets time-frequency

decompositions of the simulated time series (smoothed using a 4-s Gaussian kernel, standard

deviation = 1 s). We refer to the latter approach as wavelet-*specparam* (see Methods). We assessed

the respective performances of SPRiNT and wavelet-*specparam* with measures of Mean Absolute

Error (MAE) on their respective estimates of a/periodic spectrogram profiles and of the parameters

of their a/periodic components across time.

Figure 2: **SPRiNT *vs.* wavelet-*specparam* performances (Simulation Challenge I).**

(A) Ground truth spectrogram (left) and averaged modelled spectrograms produced by the

wavelet-*specparam* approach (middle) and SPRiNT (right; *n* = 10,000). (**B**) Aperiodic parameter

estimates (lines: median; shaded regions: first and third quartiles, *n* = 10,000) across time from

wavelet-*specparam* (left) and SPRiNT (right; black: ground truth; blue: exponent; yellow:

offset). (**C**) Absolute error (and detection performance) of alpha and beta-band rhythmic

components for wavelet-*specparam* (left) and SPRiNT (right). Violin plots represent the sample

distributions (*n* =10,000; blue: alpha peak; yellow: beta peak; white circle: median, grey box:

first and third quartiles; whiskers: range).

Figure supplement 1. Periodic parameter estimates across time (*n* = 10,000 simulations).

Figure supplement 2. Wavelet-*specparam* performance at varying spectral/temporal resolutions.

Figure supplement 3. SPRiNT performances at varying spectral/temporal resolutions.

Figure supplement 4. SPRiNT and wavelet-*specparam* without temporal smoothing and outlier

peak removal.

Overall, we found that SPRiNT parameterized spectrograms were better fits to ground

truth (MAE = 0.04, SEM [standard error of the mean] = 2.9×10^{-5}) than those from wavelet-

specparam (MAE = 0.58, SEM = 5.1×10^{-6}; Figure 2a). The data showed marked differences in

performance between SPRiNT and wavelet-*specparam* in the parameterization of aperiodic

components (error on aperiodic spectrogram: wavelet-*specparam* MAE = 0.60, SEM = 6.7×10^{-6};

SPRiNT MAE = 0.06, SEM = 4.0×10^{-5}). The performances of the two methods in parameterizing

periodic components were strong and similar (wavelet-*specparam* MAE = 0.05, SEM = 4.0×10^{-6};

SPRiNT MAE = 0.03, SEM = 2.7×10^{-5}).

SPRiNT errors on exponents (MAE = 0.11, SEM = 7.8×10^{-5}) and offsets (MAE = 0.14,

SEM = 1.1×10^{-4}) were substantially less than those from wavelet-*specparam* (exponent MAE =

0.19, SEM = 1.5×10^{-5}; offset MAE = 0.78, SEM = 2.6×10^{-5}; Figure 2A). SPRiNT detected periodic

alpha activity with higher sensitivity (99% at time bins with maximum peak amplitude) and

specificity (96%) than wavelet-*specparam* (95% sensitivity, 47% specificity). SPRiNT estimates

of alpha peak parameters were also closer to ground truth (centre frequency, amplitude, bandwidth

MAE [SEM] = 0.33 [3.6×10^{-4}] Hz, 0.20 [1.7×10^{-4}] a.u., 0.42 [4.8×10^{-4}] Hz, respectively) than

wavelet-*specparam*'s (MAE [SEM] = 0.41 Hz [4.8×10^{-5}], 0.24 [2.6×10^{-5}] a.u., 0.64 [6.4×10^{-5}] Hz,

respectively; Figure 2C). SPRiNT detected and tracked down-chirping beta periodic activity with

higher sensitivity (95% at time bins with maximum peak amplitude) and specificity (98%) than

wavelet-*specparam* (62% sensitivity, 90% specificity). SPRiNT's estimates of beta peak

parameters were also closer to ground truth (centre frequency, amplitude, bandwidth MAE = 0.43

[9.4×10^{-4}] Hz, 0.17 [3.6×10^{-4}] a.u., 0.48 [1.1×10^{-3}] Hz, respectively) than with wavelet-*specparam*

(centre frequency, amplitude, bandwidth MAE = 0.58 [1.4×10^{-4}] Hz, 0.16 [4.2×10^{-5}] a.u., 1.05

[1.2×10^{-4}] Hz, respectively; Figure 2C). We noted that both methods tended to overestimate peak

bandwidths (Figure 2 – figure supplement 1), and the effect was more pronounced with wavelet-

specparam (Figure 2C). While SPRiNT and wavelet-*specparam* performances varied with the

chosen parameters (i.e., spectral/temporal resolutions; Figure 2 – figure supplement 2 and Figure

2 – figure supplement 3; see Supplemental Materials), the optimal settings for SPRiNT

outperformed the optimal settings for wavelet-*specparam*. We report SPRiNT performances prior

to the removal of outlier peaks, as well as wavelet-*specparam* performances without temporal

smoothing in Supplemental Materials (Figure 2 – figure supplement 4).

We also parameterized the periodogram of each time series of the first simulation challenge

with *specparam* to assess the outcome of a biased assumption of stationary spectral contents across

time. The PSDs were computed using the Welch approach over 1-s time windows with 50%

overlap. The average recovered aperiodic exponent was 1.94 Hz^{-1} (actual = 1.5 to 2 Hz^{-1}) and offset

was -1.64 a.u. (actual = -2.56 to -1.41 a.u.). The only peak detected by *specparam* (99% sensitivity)

was the alpha peak, with an average center frequency of 8.09 Hz (actual = 8 Hz), amplitude of

0.79 a.u. (actual max = 1.2 a.u., and peak frequency standard deviation of 1.21 Hz (actual = 1.2

Hz). No beta peaks were detected across all spectra processed with *specparam*.

Generalization of SPRiNT across generic aperiodic and periodic fluctuations (synthetic

data).

We simulated 10,000 additional time series consisting of aperiodic and periodic components,

whose parameters were sampled continuously from realistic ranges (Figure 1C). The generators of

each trial time series comprised: i) one aperiodic component whose exponent and offset parameters

were shifted linearly over time, and ii) 0 to 4 periodic components (see Methods for details).

SPRiNT, followed by outlier peak post-processing, recovered 69% of the simulated periodic

components, with 89% specificity (70% sensitivity and 73% specificity prior to outlier removal as

shown in Figure 3 – figure supplement 3). Aperiodic exponent and offset parameters were

recovered with MAEs of 0.12 and 0.15, respectively. The centre frequency, amplitude, and

frequency width of periodic components were recovered with MAEs of 0.45, 0.23, and 0.49,

respectively (Figure 3B). We evaluated whether the detection and accuracy of parameter estimates

of periodic components depended on their frequency and amplitude (Figure 3C). The synthetized

data showed that overall, SPRiNT accurately detects up to two simultaneous periodic components

(Figure 3D). We also found that periodic components of lower frequencies were more challenging

to detect (Figure 3C & 3D; Figure 3 – figure supplement 1B) because their peak spectral

component, when present, tended to be masked by the aperiodic component of the power spectrum.

We also observed that lower amplitude peaks were more challenging to detect (Figure 3C).

However, the detection rate did not depend on peak bandwidth (Figure 3 – figure supplement 1A).

We found that when two or more peaks were present simultaneously, the detection of either or

both peaks depended on their spectral proximity (Figure 3 – figure supplement 1C). Model fit

errors (MAE = 0.032) varied significantly with the number of simultaneous periodic components,

but this effect was small (β = -0.0001, SE = 6.7×10^{-6}, 95% CI [-0.0001 -0.0001], p = 8.6 $\times 10^{-85}$; R^2

= 0.0003; Figure 3E).

Finally, we simulated 1000 additional time series comprising two periodic components

(within the 3-30 Hz and 30-80 Hz ranges, respectively) and a static knee frequency. We used

SPRiNT to parameterize the spectrograms of these times series over the 1-100 Hz frequency range

(Figure 3 – figure supplement 2). SPRiNT did not converge to fit aperiodic exponents in the range

[-5, 5] Hz^{-1} only on rare occasions (<2% of all time points). We removed these data points from

further analysis. The simulated aperiodic exponents and offsets were recovered with MAEs of 0.22

and 0.42, respectively; static knee frequencies were recovered with a MAE of 3.55×10^4 (inflated

by large outliers in absolute error; median absolute error = 11.72). Overall, SPRiNT detected the

peaks of the simulated periodic components with 56% sensitivity and 99% specificity. The spectral

parameters of periodic components were recovered with equivalent performances in the lower (3-

30 Hz) and respectively, higher (30-80 Hz) frequency ranges: MAEs for centre frequency (0.32,

resp. 0.32), amplitude (0,27, resp. 0.22), and standard deviation (0,35, resp. 0.29).

Figure 3: **SPRiNT** *performances (Simulation Challenge II).*

(A) SPRiNT parameterized spectrogram for a representative simulated time series with time-varying aperiodic (offset, exponent) and transient periodic (centre frequency, amplitude, standard

deviation) components. The red arrow indicates a cross-sectional view of the spectrogram at 14

s. (**B**) Absolute error in SPRiNT parameter estimates across all simulations (*n* = 10,000). (**C**)

Detection probability of spectral peaks (i.e., rhythmic components) depending on simulated

centre frequency and amplitude (light blue: 3-8 Hz theta; yellow: 8-13 Hz alpha; orange: 13-18

Hz beta; brown:18-35 Hz). (**D**) Number of fitted vs. simulated periodic components (spectral

peaks), across all simulations and time points. The underestimation of the number of estimated

spectral peaks is related to centre frequency: 3-8 Hz simulated peaks (light blue) account for

proportionally fewer of recovered peaks between 3-18 Hz (light blue, yellow, orange) than from

the other two frequency ranges. Samples sizes by number of simulated peaks: 0 peaks = 798,753,

1 peak = 256,599, 2 peaks = 78,698, 3 peaks =14,790, 4 peaks = 1,160. (**E**) Model fit error is not

affected by number of simulated peaks. Violin plots represent the full sample distributions (white

circle: median, grey box: first and third quartiles; whiskers: range).

Figure supplement 1. Performances of SPRiNT across a range of peak standard deviations,

frequency bands, and spectral separations between peaks.

Figure supplement 2. Performances of SPRiNT on broad-range spectrograms comprising

spectral knees.

Figure supplement 3. Performances of SPRiNT (without outlier peak removal).

Aperiodic and periodic fluctuations in resting-state EEG dynamics with eyes-closed, eyes-

open behaviour (empirical data).

We applied SPRiNT and *specparam* to resting-state EEG data from the openly available LEMON

dataset (Babayan et al., 2019). Participants (*n* = 178) were instructed to open and close their eyes

(alternating every 60 s). We used *Brainstorm* (Tadel et al., 2011) to preprocess EEG time series

from electrode Oz and obtained parameterized spectra with *specparam* and parameterized

spectrograms with SPRiNT in both behavioural conditions (eyes open or closed). We also

generated time-frequency decompositions of the same preprocessed EEG time series using Morlet

wavelets (with default parameters; see Methods and Supplemental Materials).

As expected, the group averaged periodograms showed increased Oz signal power in the

alpha range (6-14 Hz) in the eyes-closed behavioural condition with respect to eyes-open (Figure

4A). A logistic regression of *specparam* outputs (aperiodic exponent, alpha peak centre frequency,

and alpha peak amplitude entered as fixed effects) identified alpha peak power (β = -2.73, SE =

0.33, 95% CI [-3.42, -2.11]; BF = 3.21×10^{-21}) and aperiodic exponent (β = 1.14, SE = 0.42, 95%

CI [0.33, -1.99]; BF = 0.20) as predictors of eyes-open or eyes-closed behaviour (Table 1). The

resulting model suggests that both lower alpha power and larger aperiodic exponents characterize

the eyes-open condition.

Using SPRiNT, we found time-varying fluctuations of both aperiodic and alpha-band

periodic components as participants opened or closed their eyes (Figure 4 – figure supplement 1).

We observed sharp changes in aperiodic exponent and offset at the transitions between eyes-open

and eyes-closed (Figure 4 – figure supplement 1), which are likely to be artifactual residuals of

eye movements. We discarded these segments from further analysis. We ran a logistic regression

model with SPRiNT parameter estimates as fixed effects (mean and standard deviation of alpha

centre frequency, alpha power, and the aperiodic exponent across time) and found a significant

effect of mean alpha power (β = -6.31, SE = 0.92, 95% CI [-8.23, -4.61]), standard deviation of

alpha power (β = 4.64, SE = 2.03, 95% CI [0.76, 8.73]), and mean aperiodic exponent (β = 2.55,

SE = 0.53, 95% CI [1.55, 3.63]) as predictors of the behavioural condition (Table 2). According

to this model, lower alpha power, larger aperiodic exponents, and stronger fluctuations of alpha-

band activity over time are signatures of the eyes-open resting condition. A Bayes factor analysis

confirmed the evidence of effects of mean alpha power (BF = 4.39x10^{-13}) and mean aperiodic

exponent (BF = 1.62x10^{-4}), and indicated mild evidence against the temporal variability of alpha

power (BF = 3.81; Table 2). Although model fit error was slightly higher in the eyes-closed

condition, it did not affect condition relationships when included in a logistic regression (see

Supplemental Materials; Table 3). In summary, both *specparam* and SPRiNT analyses confirmed

alpha power and aperiodic exponent as neurophysiological markers of eyes-closed vs. eyes-open

behaviour. Wavelet analyses confirmed that mean alpha-band activity predicted behavioural

condition (β = -2.05, SE = 0.31, 95% CI [-2.67, -1.47]; BF = 1.08x10^{-11}; Table 4). We emphasize

that SPRiNT's spectrogram parameterization was uniquely able to reveal time-varying changes in

alpha power related to eyes-closed vs. eyes-open behaviour, although the Bayes factor for this

effect suggests it to be marginal.

Figure 4: SPRiNT *parameterization of resting-state electroencephalography.*

(A) Mean periodogram and *specparam* models for eyes-closed (blue) and eyes-open (yellow)

resting-state EEG activity (from electrode Oz; *n* = 178). (B) *specparam*-derived eyes-closed

alpha peak amplitude was predictive of age group, but mean eyes-closed alpha peak amplitude

derived from SPRiNT was not. (C) Example of intrinsic dynamics in alpha activity during the

eyes-closed period leading to divergent SPRiNT and *specparam* models (participant sub-016). In

a subset of participants (<10%), we observed strong intermittence of the presence of an alpha

peak. Since an alpha peak was not consistently present in the eyes-closed condition, *specparam*-

derived alpha peak amplitude (0.77 a.u.; light blue) is lower than SPRiNT-derived mean alpha

peak amplitude (1.06 a.u.; dark blue), as the latter includes only time samples featuring a

detected alpha peak. (D) Temporal variability in eyes-open alpha centre frequency predicts age

group. Left: mean SPRiNT spectrogram (n = 178) and sample distribution of eyes-open alpha

centre frequency (participant sub-067). Right: variability (standard deviation) in eyes-open alpha

centre frequency separated by age group. Note: no alpha peaks were detected in the eyes-open

period for one participant (boxplot line: median; boxplot limits: first and third quartiles;

whiskers: range). Sample sizes: younger adults (age: 20-40 years): 121; older adults (age: 55-80

years): 56.

Figure supplement 1. SPRiNT model parameters in resting-state EEG.

Prediction of biological age group from aperiodic and periodic components of the resting-

state EEG spectrogram (empirical data).

Using the same dataset, we tested the hypothesis that SPRiNT parameter estimates are

associated with participants' age group (i.e., younger [n = 121] vs older [n = 57] adults). Extant

literature reports slower alpha rhythms and smaller aperiodic exponents in healthy aging

(Donoghue et al., 2020). We performed a logistic regression based on SPRiNT parameter estimates

of the mean and standard deviation of alpha centre frequency, alpha power, and aperiodic exponent

as fixed effects in the eyes-open condition. We found significant effects of mean aperiodic

exponent (β = -3.31, SE = 0.75, 95% CI [-4.88, -1.91]) and standard deviation of alpha centre

frequency (β = 1.30, SE = 0.53, 95% CI [0.28, 2.39]; Table 5). We therefore found using SPRiNT

that the EEG spectrogram of older participants decreased less rapidly with frequency

(characterized by a smaller exponent) and revealed stronger time-varying fluctuations of alpha-

peak centre frequency. A Bayes factor analysis showed strong evidence for the effect of the

aperiodic exponent (BF = 5.14×10^{-5}) and for the variability of the alpha peak centre frequency (BF

= 0.20; Table 5).

We replicated the same SPRiNT parameter analysis with the data in the eyes-closed

condition. We found that mean aperiodic exponent (β = -4.34, SE = 0.84, 95% CI [-6.10, -2.79])

and mean alpha centre frequency (β = -0.74, SE = 0.27, 95% CI [-1.28, -0.24]) were predictors of

participants' age group, with older participants again showing a flatter spectrum and a slower alpha

peak (lower centre frequency; Table 6). A Bayes factor analysis provided strong evidence for the

effect of mean aperiodic exponent (BF = 1.10×10^{-7}) and for the effect of mean alpha centre

frequency (BF = 0.07; Table 6).

We performed an additional logistic regression to predict age group using the mean and

variability (standard deviation) of individual alpha peak frequency (between 6-14 Hz) from the

STFT as fixed effects. We found that only variability in eyes-open individual alpha peak frequency

predicted age group (β = 0.63, SE = 0.30, 95% CI [0.04, 1.24]), though a Bayes factor analysis

showed anecdotal evidence for this effect (BF = 0.59; Table 7). Measures of individual alpha peak

frequency can be distorted by aperiodic activity (Donoghue et al., 2020) and by the absence of a

clear peak in the spectrum. In that regard, SPRiNT can help clarify the underlying dynamical

structure of the observed effects by systematically decomposing spectrograms into explicitly

detected time-varying aperiodic and periodic components.

We also tested whether the observed differences in mean spectral parameters could be

replicated from the parameterization of the periodograms using *specparam*. We performed a

logistic regression based on *specparam* parameter estimates of alpha centre frequency, alpha

power, and aperiodic exponent as fixed effects from the average periodogram, in both behavioural

conditions. We confirmed significant effects in all the same predictors as detected by SPRiNT:

eyes-open aperiodic exponent (β = -3.30, SE = 0.85, 95% CI [-5.08, -1.74]; Table 9), eyes-closed

aperiodic exponent (β = -2.67, SE = 0.61, 95% CI [-3.94, -1.54]), and eyes-closed alpha centre

frequency (β = -0.85, SE = 0.25, 95% CI [-1.38, -0.39]; Table 10). However, we found significant

effects for eyes-open alpha centre frequency (β = -0.35, SE = 0.16, 95% CI [-0.68, -0.05]; Table

9) and eyes-closed alpha power (β = -0.96, SE = 0.37, 95% CI [-1.72, -0.24]; Table 10), which

were not observed using SPRiNT (Figure 4B). We also observed intrinsic dynamics in the alpha

band of a subset of participants (<10%) contributing to diverging measures of alpha peak

amplitude between *specparam* and SPRiNT (Figure 4C).

Finally, we performed a logistic regression using mean alpha power from the wavelet

spectrogram as a fixed effect and found that mean alpha power discriminated between age groups

only in the eyes-closed condition (β = -1.13, SE = 0.38, 95% CI [-1.90 -0.41]; Table 12). Because

wavelet spectrograms are not readily decomposed into aperiodic and periodic components, these

findings may be biased by age-related effects on aperiodic exponent, alpha-peak centre frequency

(Scally et al., 2018), and the absence of an actual periodic component in the alpha range.

Transient changes in aperiodic brain activity are associated with locomotor behaviour

(empirical data).

We used intracranial data from two Long-Evans rats recorded in layer 3 of entorhinal cortex while

they moved freely along a linear track (Mizuseki et al., 2009; https://crcns.org). Rats travelled

alternatively to either end of the track to receive a water reward, resulting in behaviours of

recurring bouts of running and resting. Power spectral density estimates revealed substantial

broadband power increases below 20 Hz during rest relative to movement (except for spectral

peaks around 8 Hz and harmonics; Figure 5; Samiee & Baillet, 2017). We therefore tested for the

possible expression of two alternating modes of aperiodic neural activity associated with each

behaviour. SPRiNT parameterization found in the two subjects that resting bouts were associated

with larger aperiodic exponents and more positive offsets than during movement bouts (Figure 5

– figure supplement 2). We ran SPRiNT parameterizations over eight-second epochs proximal to

transitions between movement and rest; we observed dynamic shifts between aperiodic modes

associated with behavioural changes (Figure 5). We tested whether changes in aperiodic exponent

proximal to transitions of movement and rest were related to movement speed and found a negative

linear association in both subjects for both transition types (EC012 transitions to rest: $\beta = -9.6 \times 10^{-3}$,

SE $= 4.7 \times 10^{-4}$, 95% CI [-1.1×10^{-2} -8.6×10^{-3}], $p < 0.001$, $R^2 = 0.29$; EC012 transitions to

movement: $\beta = -7.3 \times 10^{-3}$, SE $= 4.3 \times 10^{-4}$, 95% CI [-8.1×10^{-3} -6.4×10^{-3}], $p < 0.001$, $R^2 = 0.18$; EC013

transitions to rest: $\beta = -1.1 \times 10^{-2}$, SE $= 2.3 \times 10^{-4}$, 95% CI [-1.2×10^{-2} -1.1×10^{-2}], $p < 0.001$, $R^2 = 0.32$;

EC013 transitions to movement: $\beta = -1.2 \times 10^{-2}$, SE $= 3.2 \times 10^{-4}$, 95% CI [-1.3×10^{-2} -1.2×10^{-2}], $p <$

0.001, $R^2 = 0.26$; Figure 5 – figure supplement 3). We emphasize that the periodic features of the

recordings were non-sinusoidal and therefore were not explored further with the methods

discussed herein (Donoghue et al., 2021; Figure 5 – figure supplement 1).

Figure 5: **SPRiNT** *captures aperiodic dynamics related to locomotion.*

(A) We derived the data periodograms collapsed across rest (green) and movement (purple)

periods for subject EC012, and observed broad increases in signal power during rest compared to

movement, below 20 Hz. A representative SPRiNT spectrogram is shown. The time series of the

subject's position is shown in the top plot (green: rest; purple: movement). We observed gradual

shifts of aperiodic exponent around the occurrence of locomotor transitions (right plot), with

increasing exponents at the onset of rest (green curve) and decreasing exponents at movement

onset (purple curve). Solid lines indicated trial mean, with shaded area showing the 95% CI. **(B)**

Same data as (A), but for subject EC013. The data samples consisted of, for EC012, 62 epochs of

rest onset, 81 epochs of movement onset; for EC013, 303 epochs of rest onset and 254 epochs of

movement onset.

Figure supplement 1. Examples of sawtooth rhythms from two representative electrodes in

entorhinal cortex layer 3 from both subjects.

Figure supplement 2. Empirical distributions of SPRiNT aperiodic exponent and offset

parameters.

Figure supplement 3. Temporal variability of aperiodic exponent during transitions between

movement and rest is partially explained by movement speed.

Discussion

We introduce SPRiNT as a new method to parameterize dynamic fluctuations in the spectral

contents of neurophysiological time series. SPRiNT extends recent practical tools that determine

aperiodic and periodic parameters from static power spectra of neural signals to their spectrograms.

Aperiodic spectral components may confound the detection and interpretation of narrow-band

power changes as periodic, oscillatory signal elements. Given the scientific prominence of

measures of neural oscillations in (causal) relation to behaviour (e.g., Albouy et al., 2017) and

clinical syndromes (e.g., Ostlund et al., 2021), it is essential that their characterization in time and

frequency be contrasted with that of the underlying aperiodic background activity at the natural

time scale of behaviour and perception.

SPRiNT expands the neural spectrogram toolkit.

Recent empirical studies show that the spectral distribution of neural signal power with frequency

can be decomposed into low-dimensional aperiodic and periodic components (Donoghue et al.,

2020), and that these latter are physiologically (Gao et al., 2019), clinically (Molina et al., 2020;

Van Heumen et al., 2021), and behaviourally (Ouyang et al., 2020; Waschke et al., 2021)

meaningful.

SPRiNT extends the approach to the low-dimensional time-resolved parameterization of

neurophysiological spectrograms. The method combines the simplicity of the *specparam* spectral

decomposition approach with the computational efficiency of short-time Fourier transforms across

sliding windows. The present results demonstrate its technical concept and indicate that SPRiNT

unveils meaningful additional information from the data, beyond established tools such as wavelet

time-frequency decompositions.

Using realistic simulations of neural time series, we demonstrate the strengths and current

limitations of SPRiNT. We show that SPRiNT decompositions provide a comprehensive account

of the neural spectrogram (Figure 2A), tracking the dynamics of periodic and aperiodic signal

components across time (Figure 2B and Figure 2 – figure supplement 1). We note that the

algorithm performs optimally when the data features narrow-band oscillatory components that can

be characterized as spectral peaks (Figure 3C). The algorithm performs best when the data contains

two or fewer salient periodic components concurrently (Figure 3D). We found that these current

limitations are inherent to *specparam*, which is challenged by the dissociation of spectral peaks

from background aperiodic activity at the lower edge of the power spectrum (Donoghue et al.,

2020).

Our synthetic data also identified certain limitations of the SPRiNT approach. The

algorithm tends to overestimate the bandwidth of spectral peaks, which we discuss as related to

the frequency resolution of the spectrogram (mostly 1 Hz in the present study). The frequency

resolution of the spectrogram at 1Hz, for example, may be too low to quantify narrower band-

limited components. The intrinsic noise level present in short-time Fourier transforms (i.e., spectral

power not explained by periodic or aperiodic components) may also challenge bandwidth

estimation. Increasing STFT window length augments spectral resolution and reduces intrinsic

noise, although to the detriment of temporal specificity. We also found that SPRiNT may

underestimate the number of periodic components, though this can be interpreted as the joint

probability of SPRiNT detecting multiple independent oscillatory peaks (where the probability of

detecting a given peak is between 65-75%; approximating a binomial distribution). We found that

a peak was more likely to be detected if its amplitude is stronger and centre frequency is above 8

Hz (Figure 3C and Figure 3 – figure supplement 1B), and if separated from other peaks by at least

8 Hz (Figure 3 – figure supplement 1D). Finally, SPRiNT's performances were slightly degraded

when spectrograms comprised an aperiodic knee (Figure 3 – figure supplement 2). This is due to

the specific challenge of estimating knee parameters. Nevertheless, the spectral knee frequency is

related to intrinsic neuronal timescales and cortical microarchitecture (Gao et al., 2021), which are

expected to be stable properties within each individual and across a given recording. Thus, we

recommend estimating (and reporting) aperiodic knee frequencies from the power spectrum of the

data with *specparam,* and specifying the estimated value as a SPRiNT parameter.

SPRiNT's optional outlier peak removal procedure increases the specificity of detected

spectral peaks by emphasizing the detection of periodic components that develop over time. This

feature is controlled by threshold parameters that can be adjusted along the time and frequency

dimensions. So far, we found that applying a semi-conservative threshold for outlier removal (i.e.,

if less than 3 more peaks are detected within 2.5 Hz and 3 s around a given peak of the spectrogram)

reduced the false detection rate by 50%, without affecting the true detection rate substantially (a

<5% reduction; Figure 3 and Figure 3 – figure supplement 3). Setting these threshold parameters

too conservatively would reduce the sensitivity of peak detection.

Practical mitigation techniques have been proposed to account for the presence of

background aperiodic activity when estimating narrow-band signal power changes. For instance,

baseline normalization is a common approach used to isolate event-related signals and prepare

spectrograms for comparisons across individuals (Cohen, 2014). However, the resulting relative

measures of event-related power increases or decreases do not explicitly account for the fact that

behaviour or stimulus presentations may also induce rapid changes in aperiodic activity. Therefore,

baseline normalization followed by narrow-band analysis of power changes is not immune to

interpretation ambiguities when aperiodic background activity also changes dynamically. Further,

the definition of a reference baseline can be inadequate for some study designs, as exemplified

herein with the LEMON dataset.

SPRiNT decomposition of EEG data tracks and predicts behaviour & demographics.

We found in the LEMON dataset that measuring narrow band power changes without accounting

for concurrent variations of the aperiodic signal background challenges the interpretation of effects

manifested in the spectrogram (Scally et al., 2018). Spectral parameterization with SPRiNT or

specparam enables this distinction, showing that both periodic and aperiodic changes in neural

activity are associated with age and behaviour. We found strong evidence for decreases in alpha-

peak power and increases in aperiodic exponent during eyes-open resting-state behaviour

(compared to eyes-closed; Figure 4A). However, it remains unclear whether these effects are

independent or related. A recent analysis of the same dataset showed that the amplitude of alpha

oscillations around a non-zero mean voltage influence baseline cortical excitability (Studenova et

al., 2021)—an effect observable in part through variations of the aperiodic exponent (Gao et al.,

2017). Using both SPRiNT and *specparam*, we also observed both slower alpha-peak centre

frequencies and smaller aperiodic exponents in the older age group, in agreement with previous

literature on healthy aging (Cellier et al., 2021; Donoghue et al., 2020; Hill et al., 2022; Ostlund

et al., 2022; Schaworonkow & Voytek, 2021).

Using *specparam*, we found lower alpha-band peak amplitudes in older individuals in the

eyes-closed condition. We could not replicate this effect from spectrograms parameterized with

SPRiNT (Figure 4B). This apparent divergence may be due to the challenge of detecting low-

amplitude peaks in the spectrogram of older individuals. Periodograms are derived from averaging

across time windows, which augments signal-to-noise ratios, and therefore the sensitivity of

specparam to periodic components of lower amplitude. In a subset of participants (<10%), we also

observed considerable differences between the alpha peak amplitudes extracted from *specparam*

and SPRiNT, which we explained by unstable expressions of alpha activity over time in these

participants (Figure 4C). The average alpha peak amplitude estimated with SPRiNT is based only

on time segments when an alpha-band periodic component is detected. With *specparam*, this

estimate is derived across all time windows, regardless of the presence/absence of a bona fide

alpha component at certain time instances. The consequence is that the estimate of the average

alpha peak amplitude is larger with SPRiNT than with *specparam* in these participants. Therefore,

differences in alpha power between SPRiNT and *specparam* may be explained, at least in some

participants, by differential temporal fluctuations of alpha band activity (Donoghue et al., 2021).

This effect is reminiscent of recent observations that beta-band power suppression during motor

execution is due to sparser bursting activity, not to a sustained decrease of beta-band activity

(Sherman et al., 2016).

We also emphasize how the variability of spectral parameters may relate to demographic

features, as shown with SPRiNT's prediction of participants' age from the temporal variability of

eyes-open alpha-peak centre frequency (Figure 4D). This could account for the interpretation

derived from the periodogram, where eyes-open alpha-peak centre frequency is predictive of age

instead. Previous studies explored similar effects of within-subject variability of alpha peak centre

frequency (Haegens et al., 2014) and their clinical relevance (Larsson & Kostov 2005). These

findings augment the recent evidence that neural spectral features are robust signatures proper to

an individual (da Silva Castanheira et al., 2021) and open the possibility that their temporal

variability is neurophysiologically significant.

We also report time-resolved fluctuations in aperiodic activity related to behaviour in

freely moving rats (Figure 5). SPRiNT aperiodic parameters highlight larger spectral exponents

in rats during rest than during movement. Time-resolved aperiodic parameters can also be

tracked with SPRiNT as subjects transition from periods of movement to rest and vice-versa. The

smaller aperiodic exponents observed during movement may be indicative of periods of general

cortical disinhibition (Gao et al., 2017). Previous work on the same data has shown how

locomotor behaviour is associated with changes in amplitude and centre frequency of entorhinal

theta rhythms (Mizuseki et al., 2009; Samiee & Baillet, 2017). We also note that strong theta

activity may challenge the estimation of aperiodic parameters (Gao et al., 2017). Changes in

aperiodic exponent were partially explained by movement speed (Figure 5 – figure supplement

3), which could reflect increased processing demands from additional spatial information

entering entorhinal cortex (Keene et al., 2017) or increased activity in cells encoding speed

directly (Iwase et al., 2020). Combined, the reported findings support the notion that aperiodic

background neural activity changes in relation to a variety of contexts and subject types

(Donoghue et al., 2020; Gao et al., 2017; Molina et al., 2020; Ostlund et al., 2021; Pathania et al.,

2021; Waschke et al., 2021; van Heumen et al., 2021). Gao et al. (2017) established a link

between aperiodic exponent and the local balance of neural excitation vs. inhibition. How this

balance adjusts dynamically, potentially over a multiplicity of time scales, and relates directly or

indirectly to individual behaviour, demographics, and neurophysiological factors remains to be

studied.

Practical recommendations for using SPRiNT.

SPRiNT returns goodness-of-fit metrics for all spectrogram parameters. However, these metrics

cannot account entirely for possible misrepresentations or omissions of certain components of

the spectrogram. Visual inspections of original spectrograms and SPRiNT parameterizations are

recommended e.g., to avoid fitting a 'fixed' aperiodic model to data with a clear spectral knee, or

to ensure that the minimum peak height parameter is adjusted to the peak of lowest amplitude in

the data. Most of the results presented here were obtained with similar SPRiNT parameter

settings. Below are practical recommendations for SPRiNT parameter settings, in mirror and

complement of those provided by Ostlund et al. (2022) and Gerster et al. (2022) for *specparam*:

-    *Window length* determines the frequency and temporal resolution of the spectrogram.

This parameter needs to be adjusted to the expected timescale of the effects under study

so that multiple overlapping SPRiNT time windows cover the expected duration of the

effect of interest; see for instance, the 2-s time windows with 75% overlap designed to

detect the effect at the timescale characterized in Figure 5.

-    *Window overlap ratio* is a companion parameter of *window length* that also determines

the temporal resolution of the spectrogram. While a greater overlap ratio increases the

rate of temporal sampling, it also increases the redundancy of the data information

collected within each time window and therefore smooths the spectrogram estimates over

the time dimension. A general recommendation is that longer time windows (>2 s) enable

larger overlap ratios (>75%). We recommend a default setting of 50% as a baseline for

data exploration.

-   *Number of windows averaged in each time bin* enables to control the signal-to-noise ratio

(SNR) of the spectrogram estimates (higher SNR with more windows averaged), with the

companion effect of increasing the temporal smoothing (i.e., decreased temporal

resolution) of the spectrogram. We recommend a baseline setting of 5 windows.

Learning from the *specparam* experience, we expect that more practical (and critical)

recommendations will emerge and be shared by more users adopting SPRiNT, with the pivotal

expectation, as with all analytical methods in neuroscience (Salmelin and Baillet, 2009), that

users carefully and critically review the sensibility of the outcome of SPRiNT parameterization

applied to their own data and to their own neuroscience questions (Ostlund et al., 2022).

Future directions.

We used the short-time Fourier transform as the underlying time-frequency decomposition

technique for SPRiNT. A major asset of STFT is computational efficiency, but with sliding time

windows of fixed duration, the method is less sophisticated that wavelet alternatives in terms of

trading-off between temporal specificity and frequency resolution (Cohen, 2014). Combining

specparam with STFT yields rapid extraction of spectral parameters from time-frequency data. In

principle, spectral parameterization should be capable of supplementing any time-frequency

decomposition technique, such as wavelet transforms (Pietrelli et al., 2021), though at the expense

of significantly greater computational cost. However, we have shown that the wavelet-*specparam*

alternative to SPRiNT underperformed to recover aperiodic signal components. Further, the

temporal smoothing necessary to reduce wavelet-*specparam* parameter estimation errors to levels

similar to SPRiNT's (4-s Gaussian kernel; Figure 2), yields substantial redundancy of the spectral

parameterization following wavelet decompositions. Another alternative to using STFT would be

the recent *superlet* approach (Moca et al., 2021), which was designed to preserve a fixed resolution

across time and frequency. Combining *superlets* with *specparam* is to be explored, although

reduced computational cost remains a very practical benefit of STFT.

Scientific interest towards aperiodic neurophysiological activity has recently intensified,

especially in the context of methodological developments for the detection of transient oscillatory

activity in electrophysiology (Brady and Bardouille, 2022; Seymour et al., 2022). These methods

first remove the aperiodic component from power spectra using *specparam,* before detecting

oscillatory bursts from wavelet spectrograms. SPRiNT's outlier peak removal procedure also

detects burst-like spectrographic components, although for a different purpose. SPRiNT is one

methodological response for measuring and correcting for aperiodic spectral components and as

such, could contribute to improve tools for detecting oscillatory bursts, as suggested by Seymour

et al. (2022).

Future ameliorations for SPRiNT to determine the parameters of periodic components

(number of peaks, peak amplitude) may be driven by a model selection approach based, e.g., on

the Bayesian Information Criterion (Schwarz, 1978), which would advantage models with the most

parsimonious number of periodic components in the data.

In   conclusion,   the   SPRiNT   algorithm   enables   the   parameterization   of   the

neurophysiological spectrogram. We validated the time tracking of periodic and aperiodic spectral

features with a large sample of ground-truth synthetic time series, and empirical data including

human resting-state and rodent intracranial electrophysiological recordings. We showed that

SPRiNT provides estimates of dynamic fluctuations of aperiodic and periodic neural activity that

are related to meaningful demographic or behavioural outcomes. We anticipate that SPRiNT and

future related developments will augment the neuroscience toolkit and enable new advances in the

characterization of complex neural dynamics.

Methods

SPRiNT runs on individual time series and returns a parameterized representation of the

spectrogram. The algorithm first derives short-time fast Fourier transforms (STFTs) over time

windows that slide on the time series. Second, the modulus of STFT coefficients is averaged over

n consecutive time windows to produce smoothed power spectral density (PSD) estimates at each

time bin. Third, each of the resulting PSDs are parameterized into periodic and aperiodic

components, using the *specparam* algorithm. A fourth optional step consists of the removal of

outlier periodic components from the raw SPRiNT spectrograms. We developed SPRiNT as a

plug-in library that interoperates with *Brainstorm* (Tadel et al., 2011), and therefore is open-source

and accessible to everyone.

Parameterization of short-time periodograms.

Short-time Fourier transforms are computed iteratively on sliding time windows (default window

length = 1 s; tapered by a Hann window) using MATLAB's Fast Fourier Transform (FFT; R2020a;

Natick, MA). Each window overlaps with its nearest neighbours (default overlap = 50%). The

modulus of Fourier coefficients of the running time window is then averaged locally with those

from preceding and following time windows, with the number of time windows included in the

average, *n*, determined by the user (default is *n* = 5; Figure 1A). The resulting periodogram is then

parameterized with *specparam*. The resulting spectrogram is time-binned based on time points

located at the centre of each sliding time window.

Tracking periodic and aperiodic components across time.

We used the MATLAB implementation of *specparam* in *Brainstorm* (Tadel et al., 2011), adapted

from the original Python code (Version 1.0.0) by Donoghue et al. (2020). The aperiodic component

of the power spectrum is typically represented using two parameters (exponent and offset); an

additional knee parameter is added when a bend is present in the aperiodic component (Donoghue

et al., 2020; Gao et al., 2020). Periodic components are parameterized as peaks emerging from the

aperiodic component using Gaussian functions controlled with three parameters (mean [centre

frequency], amplitude, and standard deviation).

For algorithmic speed optimization purposes, in each iteration of *specparam* across time*,* the

optimization of the aperiodic exponent is initialized from its *specparam* estimate from the

preceding time bin. All other parameter estimates are initialized using the same data-driven

approaches as *specparam* (Donoghue et al., 2020).

Pruning of periodic component outliers.

We derived a procedure to remove occasional peaks of periodic activity from parameterized

spectrograms and emphasize expressions of biologically plausible oscillatory components across

successive time bins. This procedure removes peaks with fewer than a user-defined number of

similar peaks (by centre frequency; default = 3 peaks within 2.5 Hz) within nearby time bins

(default = 6 bins). This draws from observations in synthetic data that non-simulated peaks are

parameterized in isolation (few similar peaks in neighbouring time bins; Figure 1 – figure

supplement 1). Aperiodic parameters are refit at time bins where peaks have been removed, and

models are subsequently updated to reflect changes in parameters. This post-processing procedure

is applied on all SPRiNT outputs shown but remains optional (albeit recommended).

Study 1: Time Series Simulations.

We simulated neural time series using in-house code based on the NeuroDSP toolbox (Cole et al.,

2019) with MATLAB (R2020a; Natick, MA). The time series combined aperiodic with periodic

components (Donoghue et al., 2020). Each simulated 60-s time segment consisted of white noise

sampled at 200 Hz generated with MATLAB's coloured noise generator (R2020a; Natick, MA).

The time series was then Fourier-transformed (frequency resolution = 0.017 Hz) and convolved

with a composite spectrogram of simulated aperiodic and periodic dynamics (temporal resolution

= 0.005 s). The final simulated time series was generated as the linear combination of cosines of

each sampled frequency (with random initial phases), with amplitudes across time corresponding

to the expected power from the spectrogram.

Simulations of transient and chirping periodic components.

The aperiodic exponent was initialized to 1.5 Hz^{-1} and increased to 2.0 Hz^{-1}, offset was initialized

to -2.56 a.u. and increased to -1.41 a.u.; both linearly increasing between the 24 s and 36 s time

stamps of the time series. Periodic activity in the alpha band (centre frequency = 8 Hz, amplitude

= 1.2 a.u., standard deviation = 1.2 Hz) was generated between time stamps 8s and 40 s, as well as

between 41-46 s and 47-52 s. Periodic activity in the beta band (centre frequency = 18 Hz,

amplitude = 0.9 a.u., standard deviation = 1.4 Hz) was generated between 15-25 s and down-

chirped linearly from 18-15Hz between 18-22 s. Peak amplitude was calculated as the relative

height above the aperiodic component at every sampled frequency and time point. The signal-to-

noise ratio for peaks is reflected in their respective amplitudes, with peaks of lower amplitude

exhibiting lower signal-to-noise ratios. All amplitudes of periodic activity were tapered by a Tukey

kernel (cosine fraction = 0.4). Aperiodic and periodic parameters (and their dynamics) were

combined to form a spectrogram of simulated activity.

All simulations ($n = 10,000$) were unique as each was generated from a unique white-noise

time series seed, and the cosine waves to simulate periodic components were each assigned a

random initial phase.

Each simulated time series was analyzed with SPRiNT using 5x1s sliding time windows

with 50% overlap (frequency range: 1-40 Hz). Settings for *specparam* were: peak width limits:

[0.5 6]; maximum number of peaks: 3; minimum peak amplitude: 0.6 a.u.; peak threshold

(minimum peak signal-to-noise ratio): 2.0 standard deviations; proximity threshold: 2 standard

deviations; aperiodic mode: fixed. Settings for peak post-processing were: number of neighbouring

peaks: 3; centre frequency: 2.5 Hz; time bin: 6 bins (= 3 s). Periodic alpha activity was identified

using the highest amplitude peak parameterized in each time bin between 5.5-10.5 Hz, while

periodic beta activity was identified using the highest amplitude peak in each time bin between

13.5-20.5 Hz.

We also parameterized Morlet wavelet spectrograms of each simulated time series using

specparam (Donoghue et al., 2020; MATLAB version). Wavelet transforms were computed with

Brainstorm (Tadel et al., 2011; 1-40 Hz, in 1 Hz steps) using default settings (central frequency =

3 Hz, FWHM = 1 s). Before parameterizing wavelet transforms, we applied a 4-s temporal

smoothing filter (Gaussian kernel, standard deviation = 1 s; time range: 3.5-56.5 s, in 0.005 s steps)

to increase signal-to-noise ratio (results prior to this step are shown for the first 1,000 simulations

in Supplemental Materials). Settings for *specparam* were: peak width limits: [0.5 6]; maximum

number of peaks: 3; minimum peak amplitude: 0.6 a.u.; peak threshold: 2.0 standard deviations;

proximity threshold: 2 standard deviations; aperiodic mode: fixed. Periodic alpha activity was

identified using the highest amplitude peak parameterized in each time bin between 5.5-10.5 Hz.

Periodic beta activity was identified using the highest amplitude peak in each time bin between

13.5-20.5 Hz.

Model fit error was calculated as the mean absolute error (MAE) between expected and

modelled spectral power by each component across simulations and times. Algorithmic

performances were assessed by calculating MAE in parameter estimates across simulations and

time points relative to expected parameters. Peak-fitting probability in the alpha (5.5-10.5 Hz) and

beta (13.5-20.5 Hz) bands were calculated for each time bin as the fraction of simulations with at

least one oscillatory peak recovered in the frequency band of interest.

Generic time series simulations.

For each time series generation, we sampled the parameter values of their a/rhythmic components

uniformly from realistic ranges. Aperiodic exponents were initialized between 0.8-2.2 Hz^{-1}.

Aperiodic offsets were initialized between -8.1 and -1.5 a.u. Within the 12-36 s time segment into

the simulation (onset randomized), the aperiodic exponent and offset underwent a linear shift of

magnitude in the ranges -0.5 to 0.5 Hz^{-1} and -1 to 1 a.u. (sampled continuously, chosen randomly),

respectively. The duration of the linear shift was randomly selected for each simulated time series

between 6 and 24 s. Between 0-4 oscillatory (rhythmic) components were added to each trial with

parameters randomly sampled within the following ranges: centre frequency: 3-35 Hz; amplitude:

0.6-1.6 a.u.; standard deviation: 1-2 Hz. The onset (5-40 s) and duration (3-20 s) of each of the

rhythmic components were also randomized across components and across trials, with the

constraint that they would not overlap both in time and frequency; they were allowed to overlap

in either dimension. If a rhythmic component overlapped temporally with another one, its centre

frequency was set at least 2.5 peak standard deviations from the other temporally overlapping

rhythmic component(s). The magnitude of each periodic component was tapered by a Tukey kernel

(cosine fraction = 0.4).

Each simulation was analyzed with SPRiNT using 5x1s STFT windows with 50% overlap

(frequency range: 1-40 Hz). Settings for *specparam* were: peak width limits: [0.5 6]; maximum

number of peaks: 6; minimum peak amplitude: 0.6 a.u.; peak threshold: 2.0 standard deviations;

proximity threshold: 2 standard deviations; aperiodic mode: fixed. Settings for peak post-

processing were: number of neighbouring peaks: 3; centre frequency: 2.5 Hz; time bin: 6 bins (=

3 s). The spectrogram outcome of SPRiNT was analyzed to identify rhythmic components as

correct (i.e., present in ground truth signal) or incorrect components. Rhythmic SPRiNT

components were labelled as correct if their centre frequency was within 2.5 peak standard

deviations from any of the ground truth rhythmic components. In the event of multiple SPRiNT

rhythmic components meeting these conditions, we selected the one with the largest amplitude

peak (marking the other as incorrect).

Errors on parameter estimates were assessed via MAE measures with respect to their

ground truth values. The peak-fitting probability for each simulated rhythmic component was

derived as the fraction of correct peaks recovered when one was expected. Model fit error was

calculated for each time bin as the MAE between empirical and SPRiNT spectral power. We

used a linear regression model (*fitlm* MATLAB function; 2020a; Natick, MA) to predict model

fit errors at each time bin, using number of simulated peaks as a predictor.

$$MAE = intercept + B * number\ of\ simulated\ peaks$$

We also simulated 1000 time series with aperiodic activity featuring a static knee (Figure

3 – figure supplement 2). Aperiodic exponents were initialized between 0.8-2.2 Hz^{-1}. Aperiodic

offsets were initialized between -8.1 and -1.5 a.u., and knee frequencies were set between 0 and

30 Hz. Within the 12-36 s time segment into the simulated time series (onset randomized), the

aperiodic exponent and offset underwent a linear shift and a random magnitude in the range of -

0.5 to 0.5 Hz^{-1} and -1 to 1 a.u., respectively. The duration of the linear shift was randomly

selected for each simulated time series between 1 and 20 s; the knee frequency was constant for

each simulated time series. We added two oscillatory (rhythmic) components (amplitude: 0.6-1.6

a.u.; standard deviation: 1-2 Hz) of respective peak centre frequencies between 3-30 Hz and

between 30-80 Hz, with the constrain of minimum peak separation of at least 2.5 peak standard

deviations. The onset of each periodic component was randomly assigned between 5-25 s, with

an offset between 35-55 s.

We analyzed each simulated time series with SPRiNT using 5x1s STFT windows with

50% overlap over the 1-100 Hz frequency range. Parameter settings for *specparam* were: peak

width limits: [0.5 6]; maximum number of peaks: 3; minimum peak amplitude: 0.6 a.u.; peak

threshold: 2.0 standard deviations; proximity threshold: 2.0 standard deviations; aperiodic mode:

knee. Settings for peak post-processing were: number of neighbouring peaks: 3; centre

frequency: 2.5 Hz; time bin: 6 bins (= 3 s). The identification of periodic components was

registered as correct or incorrect using the methods described above. We discarded the time bins

(<2%) where aperiodic exponent estimations did not converge within the expected range.

Study 2: Resting-state electrophysiology data.

We used open-access resting-state electroencephalography (EEG) and demographics data

collected for 212 participants from the Leipzig Study on Mind-Body-Emotion Interactions

(LEMON; Babayan et al., 2019). Data from the original study was collected in accordance with

the Declaration of Helsinki and the study protocol was approved by the ethics committee at the

medical faculty of the University of Leipzig. Participants were asked to alternate every 60 seconds

between eyes-open and eyes-closed resting-state for 16 minutes. Continuous EEG activity (2500-

Hz sampling rate) was recorded from 61 Ag/AgCl active electrodes placed in accordance with the

10-10 system. An electrode below the right eye recorded eye-blinks (ActiCap System, Brain

Products). Impedance of all electrodes was maintained below 5 kΩ. EEG recordings were

referenced to FCz during data collection (Babayan et al., 2019), and re-referenced to an average

reference during preprocessing.

Preprocessing was performed using *Brainstorm* (Tadel et al., 2011). Recordings were

resampled to 250 Hz before being high-pass filtered above 0.1 Hz using a Kaiser window. Eyeblink

EEG artifacts were detected and attenuatedusing signal-space projection (SSP). Data was visually

inspected for bad channels and artifacts exceeding 200 µV. 20 participants were excluded for not

following task instructions, 2 for failed recordings, 1 for data missing event markers, and 11 were

excluded for EEG data of poor quality (> 8 bad sensors). The results herein are from the remaining

178 participants (average number of bad sensors = 3). We extracted the first 5 minutes of

consecutive quality data, beginning with the eyes-closed condition, from electrode Oz for each

participant. We removed 2.5 seconds of data centred at transitions between eyes-open and eyes-

closed from further analyses due to sharp changes observed in aperiodic parameters when

participants transitioned between eyes-open and eyes-closed (Figure 4 – figure supplement 1),

likely to be artifactual residuals of eye movements.

Spectrogram analysis.

Each recording block was analyzed with SPRiNT using 5x1s sliding time windows with 50%

overlap (frequency range: 1-40 Hz). We ran SPRiNT using *Brainstorm* with the following settings:

peak width limits: [1.5 6]; maximum number of peaks: 3; minimum peak amplitude: 0.5 a.u.; peak

threshold: 2.0 standard deviations; proximity threshold: 2.5 standard deviations; aperiodic mode:

fixed. Peak post-processing was run on SPRiNT outputs (number of neighbouring peaks 3; centre

frequency: 2.5 Hz; time bin: 6 bins (= 3 s). Alpha peaks were defined as all periodic components

detected between 6-14 Hz. To capture variability in alpha peak centre frequency across time, mean

and standard deviations of alpha peak centre frequency distributions were computed across both

the eyes-open and eyes-closed conditions and by age group (defined below).

We computed spectrograms from Morlet wavelet time-frequency decompositions (1-40

Hz, in 1 Hz steps) using *Brainstorm* (with default parameters; central frequency = 1 Hz, FWHM

= 3 s; Tadel et al., 2011). We also parameterized periodograms across eyes-open and eyes-closed

time segments using *specparam* with *Brainstorm*, with the following settings: frequency range: 1-

40 Hz; peak width limits: [0.5 6]; maximum number of peaks: 3; minimum peak amplitude: 0.2

a.u.; peak threshold: 2.0 standard deviations; proximity threshold: 1.5 standard deviations;

aperiodic mode: fixed.

Contrast between eyes-open and eyes-closed conditions.

All regression analyses were performed in R (V 3.6.3; R Core Team, 2020). We ran a logistic

regression model whereby we predicted the condition (i.e., eyes-open vs eyes-closed) from the

mean and standard deviation of the following SPRiNT parameters: alpha centre frequency, alpha

power, and aperiodic exponent. All model predictors were entered as fixed effects. Significance of

each beta coefficient was tested against zero (i.e., $B_n = 0$). We quantified the evidence for each

predictor in our models with a Bayes factor analysis where we systematically removed one of the

predictors and computed the Bayes factor using the *BayesFactor* library (Morey & Rouder, 2018).

We compared the most complex model (i.e., the full model) against all models formulated by

removing a single predictor. Evidence in favor of the full model (i.e., BF < 1) indicated that a given

predictor improved model fit, whereas evidence for the model without the predictor (i.e., BF > 1)

showed limited improvement in terms of model fit.

We also fitted a logistic regression model to predict experimental condition (i.e., eyes -

open & -closed; dummy coded) from mean alpha-band power (6-14 Hz) entered as a fixed effect.

Alpha-band power was computed as the mean log-power between 6-14 Hz for each condition

extracted from the Morlet wavelets spectrograms. Significance of each beta coefficient was tested

against zero (i.e., $B_n = 0$). Finally, we adjusted a logistic regression model to predict behavioural

condition (eyes -open vs eyes-closed) from *specparam* parameters (aperiodic exponent, alpha peak

centre frequency, alpha peak power) as fixed effects, where significance of each beta coefficient

was tested against zero (i.e., $B_n = 0$).

Predicting age from resting-state activity.

Participants were assigned to two groups based on their biological age: younger adults (age: 20-

40 years, *n* = 121) and older adults (age: 55-80 years, *n* = 57). The SPRiNT-modelled alpha peaks

and aperiodic parameters were collapsed across time to generate condition-specific distributions

of model parameters per participant. We used these distributions to examine the mean and standard

deviation of alpha centre frequency, alpha power, and aperiodic exponent. We fitted two logistic

regression models using the *glm* function in R (R Core Team, 2020) for the eyes-open and eyes-

closed conditions:

861
$\text{age} = \text{intercept} + B_1 * \text{mean alpha center frequency} + B_2$
$* \text{standard deviation alpha center frequency} + B_3 * \text{mean alpha power} + B_4$
$* \text{standard deviation alpha power} + B_5 * \text{mean aperiodic slope} + B_6$
$* \text{standard deviation aperiodic slope}$
$\text{age} = \text{intercept} + B_1 * \text{mean alpha center frequency} + B_2$
$* \text{standard deviation alpha center frequency} + B_3 * \text{mean alpha power} + B_4$
$* \text{standard deviation alpha power} + B_5 * \text{mean aperiodic slope} + B_6$
$* \text{standard deviation aperiodic slope}$
870
All predictors were entered as fixed effects. Significance of each beta coefficient was tested

against zero (i.e., $B_n = 0$). We also quantified the evidence for each predictor in our models with

a Bayes factor analysis. We performed similar logistic regressions using data from Morlet wavelets

spectrograms and *specparam*-modelled power spectra (using the same parameters as those used

for predicting behavioural condition). Finally, we performed logistic regressions using the mean

and temporal variability (standard deviation) of individual alpha peak frequency (the frequency

corresponding to the maximum power value between 6-14 Hz; Klimesch, 1999) derived from the

short-time Fourier transform in both conditions to predict age group.

Study 3: Intracranial rodent data.

Local field potential (LFP) recordings and animal behaviour, originally published by Mizuseki et

al. (2009), were collected from 2 Long-Evans rats (data retrieved from https://crcns.org). Animals

were implanted with eight-shank multi-site silicon probes (200 μm inter-shank distance) spanning

multiple layers of dorsocaudal medial entorhinal cortex (entorhinal cortex, dentate gyrus,

hippocampus). Neurophysiological signals were recorded while animals traversed to alternating

ends of an elevated linear track (250 x 7 cm) for 30 µL water reward (animals were water deprived

for 24 hrs prior to task). All surgical and behavioural procedures in the original study were

approved by the Institutional Animal Care and Use Committee of Rutgers University. Recordings

were acquired continuously at 20 kHz (RC Electronics) and bandpass-filtered (1 Hz-5 kHz) before

being down-sampled to 1250 Hz. In two rats (EC012, EC013), nine recording blocks of activity in

entorhinal cortex layer 3 (EC3) were selected for further analysis (16 electrodes in EC012, 8

electrodes in EC013). Electrodes in EC012 with consistent isolated signal artifacts were removed

(average number of bad electrodes = 2; none in EC013). Movement-related artifacts (large

transient changes in local field potential across all electrodes, either positive or negative) were

identified by visual inspection and data coinciding with these artifacts were later discarded from

further analysis. Animal head position was extracted from video recordings (39.06 Hz) of two

head-mounted LEDs and temporally interpolated to align with SPRiNT parameters across time

(piecewise cubic Hermite interpolative polynomial; *pchip* in MATLAB; 2020a; Natick, MA).

Spectrogram analysis.

Each recording block was analyzed with SPRiNT using 5x2 s sliding time windows with 75%

overlap (frequency range: 2-40 Hz). The 1 Hz frequency bin was omitted from spectral analyses

due to its partial attenuation by the bandpass filter applied to the data. Two-second time windows

were used to increase frequency resolution, with an overlap ratio of 75% to preserve the temporal

resolution of 0.5 s and to increase the temporal specificity of the spectrogram windows. Settings

for *specparam* were set: peak width limits: [1.5 5]; maximum number of peaks: 3; minimum peak

amplitude: 0.5 a.u.; peak threshold: 2.0 standard deviations; proximity threshold: 2.0 standard

deviations; aperiodic mode: fixed. Settings for peak post-processing were set as: number of

neighbouring peaks: 3; centre frequency bounds: 2.5 Hz; time bin bounds: 6 bins (= 3 s). Aperiodic

parameters were averaged across electrodes and aligned with behavioural data.

Tracking aperiodic dynamics during movement transitions.

Time bins were categorized based on whether animals were resting at either end of the track or

moving toward opposite ends of the track ('rest' or 'movement', respectively) using animal

position (and speed). Rest-to-movement and movement-to-rest transitions were defined as at least

four consecutive seconds of rest followed by four consecutive seconds of run ($t = 0$ s representing

the onset of movement), or vice-versa ($t = 0$ s representing the onset of rest), respectively. In both

subjects, we also fit separate linear regression models (Matlab's *fitlm*) of the relation between

aperiodic exponents and movement speed at the transitions between movement and rest.

Software and code availability

The SPRiNT algorithm and all code needed to produce the figures shown are available from

GitHub (https://github.com/lucwilson/SPRiNT). The SPRiNT algorithm is also available from the

Brainstorm distribution (Tadel et al., 2011).

Data availability

The simulated data are openly available on OSF (https://osf.io/c3gn4/). Resting-state EEG data

was obtained from the open repository LEMON (Babayan et al., 2019;

https://openneuro.org/datasets/ds000221/versions/00002). Intracranial rodent data (study HC3) is

openly available from Mizuseki et al. (2009; data retrieved from https://crcns.org). SPRiNT-

processed data is available upon request from the corresponding author.

Author contributions

All authors conceptualized the study, L.E.W. and J.D.S.C. preformed the analyses, S.B. provided

guidance with data interpretation, all authors contributed to the writing and editing of the

manuscript.

Competing interests

The authors declare no competing financial interest.

Acknowledgements

L.E.W. acknowledges the support of an NSERC Undergraduate Student Research Award.

J.D.S.C. acknowledges the support of the Alexander Graham-Bell Doctoral NSERC fellowship.

S.B. is grateful for the support received from the NIH (R01 EB026299), a Discovery Grant from

the Natural Science and Engineering Research Council of Canada (436355-13), the CIHR Canada

Research Chair in Neural Dynamics of Brain Systems, the Brain Canada Foundation with support

from Health Canada, and the Innovative Ideas program from the Canada First Research Excellence

Fund, awarded to McGill University for the Healthy Brains for Healthy Lives initiative. This

research was undertaken thanks in part to funding from the Canada First Research Excellence

Fund, awarded to McGill University for the Healthy Brains for Healthy Lives initiative.

References

Albouy, P., Weiss, A., Baillet, S., & Zatorre, R. J. (2017). Selective Entrainment of Theta

Oscillations in the Dorsal Stream Causally Enhances Auditory Working Memory

Performance. *Neuron*, *94*(1), 193-206.e195. doi:10.1016/j.neuron.2017.03.015

Alhourani, A., Wozny, T. A., Krishnaswamy, D., Pathak, S., Walls, S. A., Ghuman, A. S.,

. . . Niranjan, A. (2016). Magnetoencephalography-based identification of functional

connectivity network disruption following mild traumatic brain injury. *J Neurophysiol,*

116(4), 1840-1847. doi:10.1152/jn.00513.2016

Babayan, A., Erbey, M., Kumral, D., Reinelt, J. D., Reiter, A. M. F., Röbbig, J., . . . Villringer,

A. (2019). A mind-brain-body dataset of MRI, EEG, cognition, emotion, and peripheral

physiology in young and old adults. *Scientific Data*, *6*(1), 180308.

doi:10.1038/sdata.2018.308

Brady, B., & Bardouille, T. (2022). Periodic/Aperiodic parameterization of transient oscillations

(PAPTO)-Implications for healthy ageing. *NeuroImage*, *251*, 118974.

doi:10.1016/j.neuroimage.2022.118974

Bruns, A. (2004). Fourier-, Hilbert- and wavelet-based signal analysis: are they really

different approaches? *Journal of Neuroscience Methods, 137*(2), 321-332.

doi:10.1016/j.jneumeth.2004.03.002

Buzsáki, G. (2006). *Rhythms of the brain*. Oxford: Oxford University Press.

Buzsáki, G., & Watson, B. O. (2012). Brain rhythms and neural syntax: implications for efficient

coding of cognitive content and neuropsychiatric disease. *Dialogues in clinical*

neuroscience, *14*(4), 345-367. doi: 10.31887/DCNS.2012.14.4/gbuzsaki

Cellier, D., Riddle, J., Petersen, I., & Hwang, K. (2021). The development of theta and alpha neural

oscillations from ages 3 to 24 years. *Developmental Cognitive Neuroscience*, *50*, 100969.

doi:10.1016/j.dcn.2021.100969

Chini, M., Pfeffer, T., & Hanganu-Opatz, I. L. (2021). Developmental increase of inhibition drives

decorrelation of neural activity. *bioRxiv*, 2021.2007.2006.451299.

doi:10.1101/2021.07.06.451299

Cohen, M. X. (2014). *Analyzing neural time series data: Theory and practice*. Boston:

MIT Press.

Cole, S., Donoghue, T., Gao, R., & Voytek, B. (2019). NeuroDSP: A package for neural

digital signal processing. *Journal of Open Source Software, 4*(36), 1272.

doi:10.21105/joss.01272

da Silva Castanheira, J., Orozco Perez, H. D., Misic, B., & Baillet, S. (2021). Brief segments of

neurophysiological activity enable individual differentiation. *Nature Communications*,

12(1), 5713. doi:10.1038/s41467-021-25895-8

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., . . . Knight, R. T. (2020).

Parameterizing neural power spectra into periodic and aperiodic components. *Nature

Neuroscience, 23*(12), 1655-1665. doi:10.1038/s41593-020-00744-x

Donoghue, T., Schaworonkow, N., & Voytek, B. (2021). Methodological considerations for studying

neural oscillations. *Eur J Neurosci*. doi:10.1111/ejn.15361

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance

from field potentials. *NeuroImage, 158*, 70-78. doi:10.1016/j.neuroimage.2017.06.078

Gerster, M., Waterstraat, G., Litvak, V., Lehnertz, K., Schnitzler, A., Florin, E., . . . Nikulin, V.

(2022). Separating Neural Oscillations from Aperiodic 1/f Activity: Challenges and

Recommendations. *Neuroinformatics*. doi:10.1007/s12021-022-09581-8

Haegens, S., Cousijn, H., Wallis, G., Harrison, P. J., & Nobre, A. C. (2014). Inter- and intra-

individual variability in alpha peak frequency. *NeuroImage, 92*, 46-55.

doi:10.1016/j.neuroimage.2014.01.049

He, B. J. (2014). Scale-free brain activity: Past, present, and future. *Trends in Cognitive Sciences, 18*(9),

480-487. doi:10.1016/j.tics.2014.04.003

Hill, A. T., Clark, G. M., Bigelow, F. J., Lum, J. A. G., & Enticott, P. G. (2022). Periodic and aperiodic

neural activity displays age-dependent changes across early-to-middle childhood. *Developmental*

Cognitive Neuroscience, 54, 101076. doi:10.1016/j.dcn.2022.101076

Huang, N. E., Shen, Z., Long, S. R., Wu, M. C., Shih, H. H., Zheng, Q., . . . Liu, H. H. (1998). The

empirical mode decomposition and the Hilbert spectrum for nonlinear and non-stationary time

series analysis. *Proceedings of the Royal Society of London. Series A: Mathematical, Physical*

and Engineering Sciences, 454(1971), 903-995. doi:10.1098/rspa.1998.0193

Iwase, M., Kitanishi, T., & Mizuseki, K. (2020). Cell type, sub-region, and layer-specific speed

representation in the hippocampal–entorhinal circuit. *Scientific Reports, 10*(1), 1407.

doi:10.1038/s41598-020-58194-1

Keene, C. S., Bladon, J., McKenzie, S., Liu, C. D., O'Keefe, J., & Eichenbaum, H. (2016).

Complementary Functional Organization of Neuronal Activity Patterns in the Perirhinal, Lateral

Entorhinal, and Medial Entorhinal Cortices. *The Journal of neuroscience: the official journal of*

the Society for Neuroscience, 36(13), 3660-3675. doi:10.1523/jneurosci.4368-15.2016

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory

performance: a review and analysis. *Brain Research Reviews, 29*(2), 169-195.

doi:10.1016/S0165-0173(98)00056-3

Kucewicz, M. T., Cimbalnik, J., Matsumoto, J. Y., Brinkmann, B. H., Bower, M. R., Vasoli, V., . . .

Worrell, G. A. (2014). High frequency oscillations are associated with cognitive processing in

human recognition memory. *Brain, 137*(8), 2231-2244. doi:10.1093/brain/awu149

Larsson, P. G., & Kostov, H. (2005). Lower frequency variability in the alpha activity in EEG

among patients with epilepsy. *Clinical Neurophysiology, 116*(11), 2701-2706.

doi:10.1016/j.clinph.2005.07.019

Mizuseki, K., Sirota, A., Pastalkova, E., & Buzsáki, G. (2009). Theta Oscillations Provide Temporal

Windows for Local Circuit Computation in the Entorhinal-Hippocampal Loop. *Neuron*, *64*(2),

267-280. doi:10.1016/j.neuron.2009.08.037

Moca, V. V., Bârzan, H., Nagy-Dăbâcan, A., & Mureşan, R. C. (2021). Time-frequency super-

resolution with superlets. *Nature Communications, 12*(1), 337. doi:10.1038/s41467-020-

20539-9

Molina, J. L., Voytek, B., Thomas, M. L., Joshi, Y. B., Bhakta, S. G., Talledo, J. A., . . . Light, G. A.

(2020). Memantine Effects on Electroencephalographic Measures of Putative

Excitatory/Inhibitory Balance in Schizophrenia. *Biol Psychiatry Cogn Neurosci Neuroimaging,*

5(6), 562-568. doi:10.1016/j.bpsc.2020.02.004

Morey, R. D., & Rouder, J. N. (2018). BayesFactor: Computation of Bayes factors for common designs.

R package version 0.9.12-4.2. Retrieved from https://CRAN.R-project.org/package=BayesFactor

Ostlund, B. D., Alperin, B. R., Drew, T., & Karalunas, S. L. (2021). Behavioral and cognitive correlates

of the aperiodic (1/f-like) exponent of the EEG power spectrum in adolescents with and without

ADHD. *Developmental Cognitive Neuroscience, 48*, 100931. doi:10.1016/j.dcn.2021.100931

Ostlund, B., Donoghue, T., Anaya, B., Gunther, K. E., Karalunas, S. L., Voytek, B., & Pérez-Edgar, K.

E. (2022). Spectral parameterization for studying neurodevelopment: How and why.

Developmental Cognitive Neuroscience, 54, 101073. doi:10.1016/j.dcn.2022.101073

Ouyang, G., Hildebrandt, A., Schmitz, F., & Herrmann, C. S. (2020). Decomposing alpha and 1/f

brain activities reveals their differential associations with cognitive processing

speed. *NeuroImage*, *205*, 116304. doi:10.1016/j.neuroimage.2019.116304

Pathania, A., Schreiber, M., Miller, M. W., Euler, M. J., & Lohse, K. R. (2021). Exploring

the reliability and sensitivity of the EEG power spectrum as a biomarker. *International*

Journal of Psychophysiology, 160, 18-27. doi:10.1016/j.ijpsycho.2020.12.002

Pietrelli, M., Samaha, J., & Postle, B. R. (2021). Spectral distribution dynamics across different

attentional priority states. *bioRxiv*, 2021.2012.2002.470964.

doi:10.1101/2021.12.02.470964

Quinn, A. J., Vidaurre, D., Abeysuriya, R., Becker, R., Nobre, A. C., & Woolrich, M. W. (2018).

Task-Evoked Dynamic Network Analysis Through Hidden Markov Modeling. *Frontiers*

in neuroscience, 12. doi:10.3389/fnins.2018.00603

R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation

for Statistical Computing, Vienna, Austria. Retrieved from https://www.R-project.org/

Salmelin, R., & Baillet, S. (2009). Electromagnetic brain imaging. *Human Brain Mapping,*

30(6), 1753-1757. doi:10.1002/hbm.20795

Samaha, J., Iemi, L., Haegens, S., & Busch, N. A. (2020). Spontaneous Brain Oscillations and

1063   Perceptual Decision-Making. *Trends in Cognitive Sciences*, *24*(8), 639-653.

1064   doi:10.1016/j.tics.2020.05.004

1065   Samiee, S., & Baillet, S. (2017). Time-resolved phase-amplitude coupling in neural oscillations.

1066   *NeuroImage*, *159*, 270-279. doi:10.1016/j.neuroimage.2017.07.051

1067   Scally, B., Burke, M. R., Bunce, D., & Delvenne, J.-F. (2018). Resting-state EEG power and

1068   connectivity are associated with alpha peak frequency slowing in healthy aging.

1069   *Neurobiology of Aging*, *71*, 149-155. doi:10.1016/j.neurobiolaging.2018.07.004

1070   Schaworonkow, N., & Voytek, B. (2021). Longitudinal changes in aperiodic and periodic

1071   activity in electrophysiological recordings in the first seven months of life.

1072   *Developmental Cognitive Neuroscience, 47*, 100895. doi:10.1016/j.dcn.2020.100895

1073   Schwarz, G. (1978). Estimating the Dimension of a Model. *The Annals of Statistics*, *6*(2), 461-

1074   464. doi:10.1214/aos/1176344136

1075   Seymour, R. A., Alexander, N., & Maguire, E. A. (2022). Robust Estimation of 1/f Activity

1076   Improves Oscillatory Burst Detection. *bioRxiv*, 2022.2003.2024.485674.

1077   doi:10.1101/2022.03.24.485674

1078   Sherman Maxwell, A., Lee, S., Law, R., Haegens, S., Thorn Catherine, A., Hämäläinen Matti, S.,

1079   . . . Jones Stephanie, R. (2016). Neural mechanisms of transient neocortical beta rhythms:

1080   Converging evidence from humans, computational modeling, monkeys, and mice.

1081   *Proceedings of the National Academy of Sciences, 113*(33), E4885-E4894.

1082   doi:10.1073/pnas.1604135113

Studenova, A. A., Villringer, A., & Nikulin, V. V. (2021). Baseline shift in neuronal oscillations

and its implications for the interpretation of evoked activity obtained with EEG/MEG.

bioRxiv, 2021.2012.2001.470793. doi:10.1101/2021.12.01.470793

Tadel, F., Baillet, S., Mosher, J. C., Pantazis, D., & Leahy, R. M. (2011). Brainstorm: A User-

Friendly Application for MEG/EEG Analysis. *Computational Intelligence and*

Neuroscience, 2011, 879716. doi:10.1155/2011/879716

van Heumen, S., Moreau, J. T., Simard-Tremblay, E., Albrecht, S., Dudley, R. W., & Baillet, S. (2021).

Case Report: Aperiodic Fluctuations of Neural Activity in the Ictal MEG of a Child With Drug-

Resistant Fronto-Temporal Epilepsy. *Frontiers in Human Neuroscience, 15*(101).

doi:10.3389/fnhum.2021.646426

Voytek, B., & Knight, R. T. (2015). Dynamic Network Communication as a Unifying Neural Basis for

Cognition, Development, Aging, and Disease. *Biological Psychiatry*, *77*(12), 1089-1097.

doi:10.1016/j.biopsych.2015.04.016

Voytek, B., Kramer, M. A., Case, J., Lepage, K. Q., Tempesta, Z. R., Knight, R. T., & Gazzaley, A.

(2015). Age-Related Changes in 1/f Neural Electrophysiological Noise. *The Journal of*

Neuroscience, 35(38), 13257. doi:10.1523/JNEUROSCI.2332-14.2015

Waschke, L., Donoghue, T., Fiedler, L., Smith, S., Garrett, D. D., Voytek, B., & Obleser, J. (2021).

Modality-specific tracking of attention and sensory statistics in the human electrophysiological

spectral exponent. *eLife, 10*, e70068. doi:10.7554/eLife.70068

Wen, H., & Liu, Z. (2016). Separating fractal and oscillatory components in the power spectrum of

neurophysiological signal. *Brain Topography, 29*(1), 13-26. doi:10.1007/s10548-015-0448-0

1104

1105

Supplemental Materials

Detection and removal of spectrogram outlier components.

A known issue of *specparam* is the fitting of spurious, outlier spectral peaks (Donoghue et al.,

2020). With SPRiNT, these peaks often appear as transient episodes of periodic activity in the

spectrogram. We propose a post-processing option in SPRiNT to detect and remove fast, transient

periodic activity (Figure 1 – figure supplement 1). In short, the procedure searches for clusters of

spectral peaks over a user-defined maximum time period (see Methods). Once an outlier peak is

detected at a given time bin and removed from the model, aperiodic parameters are refit with

specparam to account for the variance previously attributed to the spurious peak and models are

subsequently updated to reflect changes in parameters.

1116

1117

Figure 1 – figure supplement 1. Overview of the outlier peak removal process.

(A) Original SPRiNT spectrogram (top) and time-frequency distribution of peak centre frequencies

(bottom). In this example, the detected peaks (red and blue dots) are sequentially removed if they

are not part of a cluster of contiguous peaks within a time-frequency region of 3 s by 2.5 Hz (red

dots; size of minimum cluster is adjustable by user). **(B)** Resulting SPRiNT spectrogram (top) and

peak centre frequencies (bottom) after outlier removal and the update of aperiodic parameters.

Dashed rectangular areas show time-frequency regions where periodic activity was simulated in

the synthesized signal.

1126

Wavelet-*specparam* with alternative wavelet parameters (synthetic data challenge I).

In addition to the wavelet settings used in the main text, we parameterized Morlet wavelet

spectrograms of the first 1,000 simulated time series from Challenge I using alternative full width

at half-maximum (FWHM) settings for the wavelet transforms. This resulted in lower FWHM

yielding wavelet spectrograms of higher temporal and lower spectral resolution (Cohen, 2014). As

with other analyses of this dataset, settings for *specparam* were: peak width limits: [0.5 6];

maximum number of peaks: 3; minimum peak amplitude: 0.6 a.u.; peak threshold: 2.0 standard

deviations; proximity threshold: 2.0 standard deviations; aperiodic mode: fixed.

1135    Wavelet settings of finer resolution in time and coarser in frequency (time range: 3-57 s,

in 0.005 s steps; central frequency = 1 Hz, FWHM = 2 s; frequency range: 1-40 Hz, in 1 Hz steps)

yielded lower estimation errors of exponent (MAE = 0.12) and offset (MAE = 0.35) compared to

original settings (exponent, offset MAE = 0.19, 0.78). Alpha peaks were recovered with higher

sensitivity (97% at time bins with maximum peak amplitude, original 95%) and specificity (32%

spurious detections, original 47%), although with greater errors in centre frequency (MAE = 0.61),

amplitude (MAE = 0.25), and bandwidth (MAE = 0.94) compared to original settings (centre

frequency, amplitude, bandwidth MAE = 0.41, 0.24, 0.64, respectively). Down-chirping beta

oscillations were detected with lower sensitivity (29% sensitivity at time bins with maximum peak

amplitude, original 62%) but higher specificity (97%, original 90%), and with greater errors in

centre frequency (MAE = 0.63), amplitude (MAE = 0.17), and bandwidth (MAE = 1.59) relative

to original settings (centre frequency, amplitude, bandwidth MAE = 0.58, 0.16, 1.05, respectively).

When wavelet settings prioritized resolution in frequency over time (time range: 4-56 s, in

0.005 s steps; central frequency = 1 Hz, FWHM = 4 s; frequency range: 1-40 Hz, in 1 Hz steps)

relative to original settings, the errors in estimates of exponent (MAE = 0.16) and offset (MAE =

0.47) parameters were reduced (original exponent, offset MAE = 0.19, 0.78, respectively). Alpha

peaks were recovered with higher sensitivity (99% at time bins with maximum peak amplitude,

original 95%) and similar specificity (46% spurious detections, original 47%), although with larger

errors in centre frequency (MAE = 0.33), amplitude (MAE = 0.20), and bandwidth (MAE = 0.43)

compared to original settings (centre frequency, amplitude, bandwidth MAE = 0.41, 0.24, 0.64,

respectively). In contrast, down-chirping beta oscillations were detected with slightly higher

sensitivity (79% at time bins with maximum peak amplitude, original 62%) and specificity (91%,

original 90%), and with lower errors on centre frequency (MAE = 0.37), amplitude (MAE = 0.14),

and bandwidth (MAE = 0.71) compared to original settings (centre frequency, amplitude,

bandwidth MAE = 0.58, 0.16, 1.05, respectively).

1160

SPRiNT with alternative STFT parameters (synthetic data challenge I).

In addition to the primary SPRiNT settings used in the main text (i.e., 5x1s windows with 50%

overlap), we parameterized STFT spectrograms of the first 1,000 simulated time series from

Challenge I using alternative settings for the short-time Fourier transforms (Figure 2 – figure

supplement 3). One setting enabled higher temporal resolution (5x1s with 75% overlap), while the

other enabled higher frequency resolution (5x2s with 75% overlap). As with other analyses of this

dataset, settings for *specparam* were: peak width limits: [0.5 6]; maximum number of peaks: 3;

minimum peak amplitude: 0.6 a.u.; peak threshold: 2.0 standard deviations; proximity threshold:

2.0 standard deviations; aperiodic mode: fixed.

SPRiNT settings for higher temporal resolution (time range: 1-59 s, in 0.25 s steps;

frequency range: 1-40 Hz, in 1 Hz steps) provided slightly larger estimation errors of exponent

(MAE = 0.15) and offset (MAE = 0.20) relative to original settings (exponent, offset MAE = 0.11,

0.14, respectively). Alpha peaks were recovered with slightly lower sensitivity (98% at time bins

with maximum peak amplitude; original 99%) and specificity (9% spurious detections; original

4%), and with greater errors in centre frequency (MAE = 0.43), amplitude (MAE = 0.24), and

bandwidth (MAE = 0.53) compared to original settings (centre frequency, amplitude, bandwidth

MAE = 0.33, 0.20, 0.42, respectively). Down-chirping beta oscillations were detected with lower

sensitivity (93% sensitivity at time bins with maximum peak amplitude, original 98%; 86%

specificity, original 98%), and with greater errors in centre frequency (MAE = 0.57), amplitude

(MAE = 0.22), and bandwidth (MAE = 0.57) compared to original settings (centre frequency,

amplitude, bandwidth MAE = 0.43, 0.17, 0.48, respectively).

SPRiNT settings for higher frequency resolution (time range: 2-58 s, in 0.5 s steps;

frequency range: 1-40 Hz, in 0.5 Hz steps) provided comparable estimation errors of exponent

(MAE = 0.13) and offset (MAE = 0.16) relative to original settings (exponent, offset MAE =

0.11, 0.20, respectively). Alpha peaks were recovered with similar sensitivity (99% at time bins

with maximum peak amplitude; original 99%) but lower specificity (21% spurious detections;

original 4%), and with comparable errors in centre frequency (MAE = 0.35), amplitude (MAE =

0.23), and bandwidth (MAE = 0.41) to original settings (centre frequency, amplitude, bandwidth

MAE = 0.33, 0.20, 0.42, respectively). Down-chirping beta oscillations were detected with

comparable sensitivity (99% sensitivity at time bins with maximum peak amplitude, original

98%) but lower specificity (78%, original 98%), and with greater errors in centre frequency

(MAE = 0.50), amplitude (MAE = 0.21), and bandwidth (MAE = 0.59) relative to original

settings (centre frequency, amplitude, bandwidth MAE = 0.43, 0.17, 0.48, respectively).

1194

Wavelet-*specparam* without temporal smoothing (synthetic data Challenge I).

We parameterized Morlet wavelet spectrograms (central frequency = 1 Hz, FWHM = 3 s; 1-40

Hz, in 1 Hz steps) of the first 1,000 simulated time series consisting of transient alpha and down-

chirping beta periodic activity (time range: 1.5-58.5 s, in 0.005 s steps). In the main text, we discuss

results from temporally smoothed wavelet spectrograms (see Methods). Here, we show results

without temporal smoothing (Figure 2 – figure supplement 4).

Error in estimates from unsmoothed parameterized wavelet spectrograms of exponent

(MAE = 0.41) and offset (MAE = 0.83) parameters were worse than those obtained from smoothed

wavelet decompositions (exponent MAE = 0.19; offset MAE = 0.78). Alpha peaks were recovered

with lower sensitivity (84% at time bins with maximum peak amplitude) and specificity (41%

spurious detections), and with greater errors on centre frequency (MAE = 0.82), amplitude (MAE

= 0.53), and bandwidth (MAE = 0.91). The down-chirping beta oscillation was also detected with

lower sensitivity (53% at time bins with maximum peak amplitude) and specificity (74%), and

with greater errors on centre frequency (MAE = 1.23), amplitude (MAE = 0.60), and bandwidth

(MAE = 1.10).

1210

SPRiNT without outlier peak removal (synthetic data Challenge 1).

Here, we present the results of SPRiNT from synthetic data Challenge I without outlier peak

removal (Figure 2 – figure supplement 4). For the aperiodic component, SPRiNT accurately

recovered both ground truth exponent (MAE = 0.11) and offset (MAE = 0.15). It also detected the

occurrences of alpha peaks with high sensitivity (99% at time bins with maximum peak amplitude)

and specificity (6% spurious detections), and with low errors on their centre frequency (MAE =

0.33) and amplitude (MAE = 0.20) parameters, but overestimated the width of the periodic peak

components (MAE = 0.42). SPRiNT detected and tracked the down-chirping beta periodic

components with high sensitivity (95% at time bins with maximum peak amplitude) but lower

specificity (95%) than with outlier peak removal (98%). Time-collapsed errors on centre frequency

(MAE = 0.44) and amplitude (MAE = 0.17) parameters were low, with a tendency to overestimate

the width in frequency of the periodic component (MAE = 0.48). Results following outlier peak

removal are presented and discussed in the main text.

1224

Figure 2 – figure supplement 1. Periodic parameter estimates across time.

(A) Results from the temporally smoothed wavelet-*specparam* approach for the alpha (top) and

beta (bottom) rhythmic components for each estimated parameter (from left to right: centre

frequency, spectral peak amplitude, and standard deviation). Grey dashed line: ground truth;

coloured line: median; shaded region: first and third quartiles. Bar plots in left panels: probability

1231   of detecting an oscillatory peak within respective frequency ranges at each time bin. **(B)** Same

1232   display for the results obtained with SPRiNT. All with *n* = 10,000 simulations.

1233

1234

Figure 2 – figure supplement 2. Wavelet-*specparam* performances at varying

spectral/temporal resolutions.

(A) Aperiodic parameter estimates (lines: median; shaded regions: first and third quartiles, *n* =

1,000) across time from wavelet-*specparam* with FWHM = 2 s (left) and wavelet-*specparam* with

FWHM = 4 s (right; black dash: ground truth; blue: exponent; yellow: offset). **(B)** Absolute error

(and detection performance) of alpha and beta-band rhythmic components for wavelet-*specparam*

with FWHM = 2 s (left) and wavelet-*specparam* with FWHM = 4 s (right). Violin plots represent

the sample distributions (n =1,000; blue: alpha peak; yellow: beta peak; white circle: median, grey

box: first and third quartiles; whiskers: range).

1244

1245

Figure 2 – figure supplement 3. SPRiNT performances at varying spectral/temporal

resolutions.

(A) Aperiodic parameter estimates (lines: median; shaded regions: first and third quartiles, n =

1,000) across time from SPRiNT spectrograms obtained from 5 1-s time windows with 75%

overlap (left) vs. 5 2-s windows with 75% overlap (right; black dash: ground truth; blue: exponent;

yellow: offset). **(B)** Absolute error (and detection performance) of alpha and beta-band periodic

components for SPRiNT using 5 1-s time windows with 75% overlap (left) and 5 2-stime windows

with 75% overlap (right). Violin plots represent the sample distributions ($n =$1,000; blue: alpha

peak; yellow: beta peak; white circle: median, grey box: first and third quartiles; whiskers: range).

Figure 2 - figure supplement 4. Raw performances of SPRiNT and wavelet-*specparam*

(without temporal smoothing and outlier peak removal).

(A) Aperiodic parameter estimates (lines: median; shaded regions: first and third quartiles, n =

1,000) across time from unsmoothed wavelet-*specparam* (left) and SPRiNT without outlier peak

removal (right; black dash: ground truth; blue: exponent; yellow: offset). **(B)** Absolute error (and

detection performance) of alpha and beta-band periodic components for unsmoothed wavelet-

specparam (left) and SPRiNT without outlier peak removal (right). Violin plots represent the

sample distributions (n =1,000; blue: alpha peak; yellow: beta peak; white circle: median, grey

box: first and third quartiles; whiskers: range).

1266

Generalization of SPRiNT across generic aperiodic and periodic fluctuations without

outlier removal (synthetic data Challenge II).

We present the results of the second synthetic data challenge without outlier peak removal

(Figure 3 – figure supplement 3). SPRiNT recovered 70% of the simulated periodic components,

with 73% specificity. Dynamic aperiodic exponents were recovered with a MAE of 0.13, while

dynamic offsets were recovered with a MAE of 0.16. Centre frequency, amplitude, and standard

deviation parameters were recovered with MAEs of 0.46, 0.23, and 0.49, respectively.

1274

1275

Figure 3 – figure supplement 1. Performances of SPRiNT across a range of peak standard

deviation, frequency band, and spectral separation between peaks.

(A) Detection probability of spectral peaks (i.e., rhythmic components) did not depend on

simulated standard deviation (bandwidth). **(B)** Confusion matrix of estimated centre frequency

range against simulated centre frequency range. 3-8 Hz peaks were more challenging to detect than

other frequency ranges. **(C)** Sample estimated shape (yellow: median; shaded areas: first and third

quartiles) of two simulated spectral peaks (blue; peak 1 [centre frequency, amplitude, standard

deviation]: 8.8 Hz, 1.1 a.u, 1.8 Hz; peak 2: 13.8 Hz, 1.2 a.u., 1.7 Hz) overlapping in frequency

space. **(D)** Individual (any peak; yellow) and joint (both peaks; blue) peak detection rate as a

function of the proximity of two peaks simultaneously present. Individual and joint peak detection

probability was lower when two peaks were within 8 Hz of one another.

1287

1288

Figure 3 – figure supplement 2. Performances of SPRiNT on broad-range spectrograms

comprising spectral knees.

(A) SPRiNT parameterized spectrogram for a representative simulated time series with time-

varying aperiodic (offset, exponent) and two periodic (centre frequency, amplitude, standard

deviation) components, with a static knee component. One peak is in a lower frequency range (3-

30 Hz), while the other is in a higher frequency range (30-80 Hz). The red arrow indicates a cross-

sectional view of the spectrogram at 20 s (in log-frequency space). **(B)** Absolute error in SPRiNT

aperiodic parameter estimates across all simulations (n = 1,000) **(C)** Absolute error in SPRiNT

periodic parameter estimates across all simulations (blue: 3-30 Hz; yellow: 30-80 Hz; n = 1,000).

Violin plots represent the full-sample distributions (white circle: median, grey box: first and third

quartiles; whiskers: range).

1300

1301

Figure 3 – figure supplement 3. Performances of SPRiNT (without outlier peak removal).

(A) SPRiNT parameterized spectrogram for a representative simulated time series with time-

varying aperiodic (offset, exponent) and transient periodic (centre frequency, amplitude, standard

deviation) components. The red arrow indicates a cross-sectional view of the spectrogram at 14 s.

(B) Absolute error in SPRiNT parameter estimates across all simulations (n = 10,000). **(C)**

Detection probability of spectral peaks (i.e., rhythmic components) depending on simulated centre

frequency and amplitude (light blue: 3-8 Hz theta; yellow: 8-13 Hz alpha; orange: 13-18 Hz beta).

(D) Number of fitted vs. simulated periodic components, across all simulations and time points.

The underestimation of the number of estimated spectral peaks is related to centre frequency: 3-8

Hz simulated peaks (blue) account for proportionally fewer of recovered peaks between 3-18 Hz

(blue, yellow, orange) than from the other two frequency ranges. (Samples sizes by number of

simulated peaks: 0 peaks = 798,753, 1 peak = 256,599, 2 peaks = 78,698, 3 peaks =14,790, 4 peaks

= 1,160) **(E)** Model fit error is not affected by number of simulated peaks. Violin plots show the

full sample distributions (white circle: median, blue box: first and third quartiles; whiskers: range).

1316

SPRiNT model fit error does not affect condition associations.

We performed t-tests of model fit errors (MAE) between conditions and age groups. While there

were no age-related effects on model fit error (eyes-open: p = 0.09; eyes closed: p = 0.69), we

observed slightly lower model fit errors in the eyes-open condition (mean = 0.032) compared to

the eyes-closed condition (mean = 0.033; t(354) = -3.17, p = 0.002, 95% CI [$3.0x10^{-4}$ $1.3x10^{-3}$]).

The size of this effect was small-to-medium (Cohen's d = 0.34).

To determine whether model fit error would affect our SPRiNT logistic regression model

for condition, we included it as a fixed effect in a new logistic regression model (Table 3). Here,

we observed the same effects for predicting condition as the original model (mean aperiodic

exponent, mean alpha power, variability of alpha power; see Table 2), with no significant effect

of model fit error (p = 0.45).

1328

1329

1330 **Figure 4 – figure supplement 1. SPRiNT model parameters in resting-state EEG.**

1331 SPRiNT aperiodic parameters (top panel; line: group mean ($n = 178$); shaded region: 95% CI) and

1332 SPRiNT periodic activity averaged across participants (bottom panel).

1333

	Condition			
Predictors	*Log-Odds*	*CI*	*p*	*BF*
(Intercept)	0.86	-1.85 – 3.64	0.537	
Alpha centre frequency (*specparam*)	0.00	-0.23 – 0.23	0.990	7.97
Alpha amplitude (*specparam*)	-2.73	-3.42 – -2.11	**<0.001**	**3.21 e-21**
Aperiodic exponent (*specparam*)	1.14	0.33 – 1.99	**0.007**	**0.20**

Observations	323		
R^2 Tjur	0.284		

Table 1. Logistic regression model of *specparam* parameters for predicting condition (eyes-closed vs eyes-open).

	Condition			
Predictors	*Log-Odds*	*CI*	*p*	*BF*
(Intercept)	0.10	-3.75 – 4.02	0.959	
Mean alpha center frequency	0.24	-0.04 – 0.52	0.101	1.58
Std alpha center frequency	-0.06	-0.97 – 0.86	0.898	4.39
Mean alpha power	-6.31	-8.23 – -4.61	**<0.001**	**4.51e-13**
Std alpha power	4.64	0.76 – 8.73	**0.022**	**3.81**
Mean aperiodic slope	2.55	1.55 – 3.63	**<0.001**	**1.62e-4**
Std aperiodic slope	-2.74	-8.54 – 3.38	0.362	4.32
Observations	355			
R^2 Tjur	0.432			

Table 2. Logistic regression model of SPRiNT parameters for predicting condition (eyes-closed vs eyes-open).

	Condition		
Predictors	*Log-Odds*	*CI*	*p*
(Intercept)	-1.37	-8.83 – 4.07	0.620

Mean alpha center frequency	0.23	-0.05 – 0.51	0.115
Std alpha center frequency	-0.15	-1.08 – 0.79	0.751
Mean alpha power	-6.62	-8.73 – -4.73	**<0.001**
Std alpha power	5.15	1.05 – 9.46	**0.016**
Mean aperiodic slope	2.63	1.60 – 3.73	**<0.001**
Std aperiodic slope	-3.79	-10.21 – 2.89	0.253
Model fit MAE	59.96	-95.00 – 215.29	0.447
Observations	355		
R^2 Tjur	0.433		

Table 3. Logistic regression model of SPRiNT parameters for predicting condition (eyes-closed vs eyes-open), with model fit error (MAE) as a predictor.

	Condition			
Predictors	*Log-Odds*	*CI*	*p*	*BF*
(Intercept)	-25.98	-33.87 – -18.65	**<0.001**	
Alpha power (Morlet wavelets)	-2.05	-2.67 – -1.47	**<0.001**	**1.08e-11**
Observations	356			
R^2 Tjur	0.148			

Table 4. Logistic regression model parameters for predicting condition (eyes-closed vs eyes-open) from Morlet wavelet spectrograms.

Age

Predictors	Log-Odds	CI	p	BF
(Intercept)	1.92	-2.82 – 6.80	0.428	
Eyes-open mean alpha center frequency	-0.05	-0.39 – 0.29	0.789	3.43
Eyes-open std alpha center frequency	1.30	0.28 – 2.39	**0.015**	**0.20**
Eyes-open mean alpha power	0.41	-2.69 – 3.27	0.784	2.97
Eyes-open std alpha power	-3.81	-9.47 – 1.54	0.172	1.14
Eyes-open mean aperiodic slope	-3.31	-4.88 – -1.91	**<0.001**	**5.14e-05**
Eyes-open std aperiodic slope	3.44	-4.83 – 11.06	0.388	2.66
Observations	177			
R^2 Tjur	0.216			

Table 5. Eyes-open logistic regression model parameters for predicting age group, SPRiNT.

		Age		
Predictors	Log-Odds	CI	p	BF
(Intercept)	11.23	4.63 – 18.50	**0.001**	
Eyes-closed mean center frequency	-0.74	-1.28 – -0.24	**0.006**	**0.07**
Eyes-closed std center frequency	1.01	-0.48 – 2.56	0.188	1.65
Eyes-closed mean alpha power	-0.15	-1.76 – 1.43	0.852	3.90
Eyes-closed std alpha power	-0.51	-5.32 – 4.22	0.831	3.61
Eyes-closed mean aperiodic slope	-4.34	-6.10 – -2.79	**<0.001**	**1.10e-07**
Eyes-closed std aperiodic slope	0.54	-9.66 – 9.45	0.910	3.93
Observations	178			
R^2 Tjur	0. 272			

Table 6. Eyes-closed logistic regression model parameters for predicting age group, SPRiNT.

Predictors	Age			
	Log-Odds	CI	p	BF
(Intercept)	-0.44	-3.04 – 2.11	0.734	
Eyes-open mean individual alpha peak frequency (STFT)	-0.17	-0.45 – 0.11	0.233	2.33
Eyes-open std individual alpha peak frequency (STFT)	0.63	0.04 – 1.24	**0.040**	**0.59**
Observations	178			
R^2 Tjur	0.026			

1352   **Table 7.** Eyes-open logistic regression model parameters for predicting age group, STFT.

1353

Predictors	Age			
	Log-Odds	CI	p	BF
(Intercept)	1.83	-1.98 – 5.75	0.350	
Eyes-closed mean individual alpha peak frequency (STFT)	-0.31	-0.70 – 0.07	0.113	1.28
Eyes-closed std individual alpha peak frequency (STFT)	0.30	-0.22 – 0.81	0.256	2.32
Observations	178			
R^2 Tjur	0.024			

1354   **Table 8.** Eyes-closed logistic regression model parameters for predicting age group, STFT.

1355

Predictors	Age			
	Log-Odds	CI	p	BF
(Intercept)	7.61	3.63 – 12.09	**<0.001**	
Eyes-open aperiodic exponent (*specparam)*	-3.30	-5.08 – -1.74	**<0.001**	**4.61 e-4**

Eyes-open alpha center frequency (*specparam*)	-0.35	-0.68 – -0.05	**0.028**	**0.26**
Eyes-open alpha amplitude (*specparam*)	-1.34	-2.86 – 0.02	0.066	0.72
Observations	147		
R² Tjur	0.207		

Table 9. Eyes-open logistic regression model parameters for predicting age group, *specparam*.

	Age			
Predictors	*Log-Odds*	*CI*	*p*	*BF*
(Intercept)	12.40	7.11 – 18.50	**<0.001**	
Eyes-closed aperiodic exponent (*specparam*)	-2.67	-3.94 – -1.54	**<0.001**	**3.22e-5**
Eyes-closed alpha center frequency (*specparam*)	-0.85	-1.38 – -0.39	**0.001**	**3.61e-3**
Eyes-closed alpha amplitude (*specparam*)	-0.96	-1.72 – -0.24	**0.010**	**0.11**
Observations	176			
R² Tjur	0.246			

Table 10. Eyes-closed logistic regression model parameters for predicting age group,

specparam.

	Age			
Predictors	*Log-Odds*	*CI*	*p*	*BF*
(Intercept)	-9.04	-21.73 – 3.13	0.152	
Eyes-open alpha power (Morlet wavelets)	-0.64	-1.63 – 0.30	0.189	2.74
Observations	178			
R² Tjur	0.010			

1361 **Table 11.** Eyes-open logistic regression model parameters for predicting age group, Morlet

1362 wavelets.

1363

Predictors	Age			
	Log-Odds	CI	p	BF
(Intercept)	-14.93	-24.68 – -5.89	**0.002**	
Eyes-closed alpha power (Morlet wavelets)	-1.13	-1.90 – -0.41	**0.003**	**0.07**
Observations	178			
R^2 Tjur	0.053			

1364 **Table 12.** Eyes-closed logistic regression model parameters for predicting age group, Morlet

1365 wavelets**.**

1366

1368   **Figure 5 – figure supplement 1. Examples of sawtooth rhythms from two representative**

1369   **electrodes in entorhinal cortex layer 3 from both subjects.**

1370   **(A)** Example time-series of prominent sawtooth rhythms from two representative electrodes during

1371   a movement bout for subject EC012 (top), producing harmonics of activity in the average modelled

1372   spectrogram (bottom). **(B)** Same as (A), for subject EC013.

1373

1374

Figure 5 – figure supplement 2. Empirical distributions of SPRiNT aperiodic exponent and

offset parameters.

(A) Empirical distributions of aperiodic exponent and offset estimates for subject EC012 at time

bins associated with rest (green) and movement (purple) bouts (boxplot line: median; boxplot

limits: first and third quartiles; whiskers: range). **(B)** Same as (A), for subject EC013. Sample

sizes: EC012 rest (movement) = 3,584 (4,325) bins; EC013 rest (movement) = 9,238 (6,672) bins.

1375

1376

1377

1378

1379

1380

1381

Figure 5 – figure supplement 3. Temporal variability of aperiodic exponent during

transitions between movement and rest is partially explained by movement speed.

(A) Empirical distributions of aperiodic exponent plotted across movement speeds for subject

EC012 during transitions to rest. **(B)** Same as (A) for transitions to movement. **(C)** Same as (A),

for subject EC013. **(D)** Same as (B), for subject EC013. Line: linear model fit; shaded area: model

95% CI). Sample sizes: EC012 transition to rest (movement) = 1,054 (1,377); EC013 rest

(movement) = 5,151 (4,318).

1393 **References**

1394 Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., . . . Knight, R. T. (2020).

1395   Parameterizing neural power spectra into periodic and aperiodic components. *Nature*

1396   *Neuroscience, 23*(12), 1655-1665. doi:10.1038/s41593-020-00744-x