Within and among population differences in cuticular hydrocarbons in the

seabird tick *Ixodes uriae*.

Marlène Dupraz[a]*, Chloe Leroy[b], Thorkell Lindberg Thórarinsson[c], Patrizia d'Ettorre[b], Karen D.

McCoy[a].

[a] MIVEGEC, IRD, CNRS, Univ. Montpellier, Domaine La Valette - 900, rue Jean François BRETON

34090 Montpellier, France

[b] Laboratoire d'Ethologie Expérimentale et Comparée UR 4443 (LEEC), Université Sorbonne Paris

Nord, F-93430 Villetaneuse, France

[c] Northeast Iceland Nature Research Centre, 640 Húsavík, Iceland

*Corresponding author: marlenedupraz@gmail.com; Tel. +33 4 67 41 61 78;

Abstract

The hydrophobic layer of the arthropod cuticle acts to maintain water balance, but can also serve to transmit chemical signals via cuticular hydrocarbons (CHC), essential mediators of arthropod behavior. CHC signatures typically vary qualitatively among species, but also quantitatively among populations within a species, and have been used as taxonomic tools to differentiate species or populations in a variety of taxa. Most work in this area to date has focused on insects, with little known for other arthropod classes such as ticks. The worldwide distribution and extensive host-range of the seabird tick *Ixodes uriae* make it a good model to study the factors influencing CHC composition. Genetically differentiated host-races of *I. uriae* have evolved across the distribution of this species but the factors promoting sympatric population divergence are still unknown. To test for a potential role of host-associated CHC in population isolation, we collected *I. uriae* specimens from two of its seabird hosts, the Atlantic puffin (*Fratercula arctica)* and the common guillemot (*Uria aalge*) in different colonies in Iceland. Using gas-chromatography and mass-spectrometry, we detected a complex cuticular mixture of 22 hydrocarbons, including *n*-alkanes, methyl-alkanes and alkenes ranging from 17 to 33 carbons in length. We found that each population had a distinct CHC profile. The host group explained the greatest amount of population divergence, with long-chain hydrocarbons being more abundant in puffin tick populations compared to guillemot tick populations. Future work will now be required to test whether the different CHC signals reinforce assortative mating, and thus play a role in generating *I. uriae* population divergence patterns, and to evaluate diverse hypotheses on the origin of distinct population signatures.

Keywords: Host race formation; GC-MS; colonial seabirds; Ixodidae; environmental variation

Introduction

The arthropod cuticle acts as both an exoskeleton and a barrier from the external environment (Andersen, 1979). The outermost layer, the epicuticle, is covered by a lipid layer (Lockey, 1988) made-up of esters, carboxylic acids, alcohols, carbonyls and long-chain hydrocarbons (Andersen, 1979) that protects the organism from desiccation (Filshie, 1982). However, cuticular hydrocarbons (CHC) are also involved in chemical communication serving as sex pheromones, kairomones and/or signature mixtures allowing recognition of social identity (van Zweden and d'Ettorre, 2010; Wyatt, 2010). Using analytic techniques such as gas chromatography-mass spectrometry (GC-MS) and MALDI-TOF mass spectrometry, researchers have described hydrocarbons of up 70 carbons in chain length in insects, principally *n*-alkanes, methyl-branched alkanes and alkenes (Blomquist and Bagnères, 2010). The array of CHC on the cuticle constitute a species-specific chemical signature, varying qualitatively between species and quantitatively within species (Lockey, 1988). CHC patterns are genetically controlled, but the relative abundance of particular components can be linked to environmental conditions (Estrada-Peña et al., 1993; Gibbs et al., 1991). CHC patterns have been used as taxonomic tools to characterize hundreds of arthropod species (Howard and Blomquist, 2005), and to discriminate closely-related populations (Bagnères et al., 1991; Bartelt et al., 1986; Jallon and David, 1987; Kruger et al., 1991; Simmons et al., 2014).

In ticks, hematophagous arthropods widely distributed across the globe and parasitizing a diverse array of vertebrate species (McCoy and Boulanger, 2015), little work has been performed to describe CHC profiles and their variation among species and populations. The only studies to date have used CHC profiles in an attempt to differentiate closely-related taxa (Estrada-Peña et al., 1992, 1994, 1996; Hunt, 1986; Estrada-Peña and Dusbabek, 1993). However, tick CHCs may play essential roles in several aspects of tick life histories. First, the tick epicuticle is perforated with numerous channels providing

a large surface for exchange with the external environment. As most tick species spend the major

part of their life cycle in the off-host environment, maintaining water balance across this surface

under different environmental conditions will directly dictate survival (Randolph and Storey, 1999);

the presence of CHCs likely plays an important role in this. Second, as obligate parasites, access to

the vertebrate host for the bloodmeal is a key aspect of the tick life cycle and, as such, ticks have

adapted key traits to locate and successfully exploit their host (McCoy and Boulanger, 2015), as for

example Haller's organ, a unique chemosensory structure located on the 1st pairs of legs (Leonovich,

2004). Shimshoni et al. (2013) found *Rhipicephalus* tick species had different cuticular fatty acid

compositions in relation to host use. However, whether these differences are linked to adaptive

survival or by-product variation due to the host resource is unknown as yet. Finally, ticks aggregate

both on hosts and in the off-host environment (Randolph, 1998). This behavior is thought to facilitate

blood feeding on the host and increase survival in the off-host environment. It may also enable ticks

to find appropriate mates for reproduction. Some ticks, such as the tropical bont tick *Amblyomma*

variegatum, produce a multicomponent pheromone to provoke aggregation (Schöni et al., 1984) but

the potential role of these pheromones in assortative mating has never been examined.

Seabird ticks are generally nidicolous, exploiting different local host species that use diverse micro-

habitats within the colony (Dietrich et al., 2011). Due to this diversity, these ticks may experience

diverse selective pressures coming from both the hosts and from the temperature and humidity

conditions of the nest micro-habitat. In particular, *Ixodes uriae*, a tick associated with seabird colonies

in the polar regions of both hemispheres, is known to form host-specific races that show genetic

(McCoy et al., 2005, 2003, 2001) and morphological (Dietrich et al., 2013) differences in relation to

host use. Differential performance on alternative hosts has also been experimentally demonstrated

(Dietrich et al., 2014). Nevertheless, the factors driving divergence within this species have yet to be

specifically identified. A potential role for isolating mechanisms that lead to assortative mating based

on host use have been suggested (McCoy et al., 2013).

Here, we examine the degree of cuticular hydrocarbon diversity in *I. uriae* and the factors influencing

these chemical signatures. Based on current knowledge, we predicted that cuticular hydrocarbon

patterns could vary in ticks 1) exploiting different host species and 2) sampled in different geographic

locations. If host exploitation modifies CHC diversity and abundance, we expected that signatures in

ticks from the same seabird host in different locations should be more similar than signatures in ticks

from different host species in the same geographic location. To test these predictions, we collected *I.*

uriae specimens in the nesting area of two seabird host species, the common guillemot (CG) *Uria*

aalge, and the Atlantic puffin (PF) *Fratercula arctica,* at three locations in Iceland. We then extracted

cuticular hydrocarbons and analyzed them using gas chromatography-mass spectrometry (GC-MS).

Material and methods

Population samples

Flat adult female ticks were collected off-host in three sites in Iceland in June 2016: common

guillemot ticks (CG) were collected under rocks in the middle of the colony in Langanes (L)

(66°22'07.2"N 14°38'33.0"W) and on Grimsey Island (GRI) (66°32'57.3"N 17°59'31.1"W); puffin ticks

(PF) were collected in burrows on Lundey Island (LI) (66°06'53.2"N 17°22'13.1"W) and on Grimsey

Island (GRI) (66°32'39.1"N 18°01'12.9"W) (Fig.1). Ticks were kept alive in plastic tubes before

hydrocarbon extraction. Four replicates of 10 flat ticks were extracted for each host species at each

site.

I. uriae ♀   PF   CG

Fig.1: Map showing the sampling locations for *Ixodes uriae* ticks and hosts in Iceland in June 2016. PF refers to the Atlantic puffin (*Fratercula arctica*) and CG

to the common guillemot (*Uria aalge*). The acronyms of the sampling locations are as follows: Grimsey Island - G; Lundey Island – Li; Langanes - L.

Hydrocarbon extraction

Several extraction attempts were made with ticks kept alive or stored in alcohol and with only one individual or in batches of five, but the quantities of extracted cuticular compounds were too low to use in analyses. Cuticular compounds were therefore extracted by immersing 10 living flat female ticks in 200 µl of pentane (HPLC grade, Sigma-Aldrich) in glass vials. The vials were agitated for 1 minute, set down to rest for 5 minutes and then re-agitated for 1 minute. Ticks were then removed from the vials and preserved in 70% ethanol. The pentane was then evaporated off over 30 minutes to obtain dried eluent and the vials were closed and kept at 4°C until analyses. A negative control containing only 200 µl of pentane was added to each extraction session to control for potential contamination. We did not use an internal standard because the aim of our analysis was not to quantify the absolute amount of CHCs, but to compare the relative proportions of CHCs among samples.

Chemical analyses

Cuticular hydrocarbons are not very volatile, but to reduce variation during the evaporation process, all samples were treated in the same way, placed under the same fume hood in a temperature controlled room. Samples were re-diluted in 40µl of pentane and 3µl were injected into an Agilent Technologies 7890A gas chromatograph (capillary column: Agilent HP-5MS, 30 m × 0.25 mm × 0.25 µm; split–splitless injector; carrying helium gas at 1 mL/min) coupled with an Agilent 5975C mass spectrometer with 70 eV electron impact ionization. The oven temperature was programmed at 70°C for 1min, and was increased at 30°C/min to 200°C, then to 320°C at 5°C/min and held for 5 min. Compounds were then identified (raw data available on Zenodo at https://doi.org/10.5281/zenodo.6497483) and sorted on the basis of their mass spectra and retention time by comparison with standards (alkane standard solutions, Sigma-Aldrich, Saint Louis,

MO, USA) and published spectra (NIST Library). The abundance of each detected cuticular

hydrocarbon was used to build a box plot using the ggplot 2 package with R software (version 4.1.1).

The areas under the peaks were extracted using the Agilent MSD ChemStation software

(E.02.01.1177) and were used to calculate the relative amount of each hydrocarbon to normalize data

coming from different origins and of different qualities. The mean peak area of the four population

replicates was calculated for each hydrocarbon to obtain a colony average.

Multivariate analyses

For analytical purposes, if the relative abundance of a CHC was 0, the value was replaced by 0.00001

which is several times lower than the lowest quantity found for a CHC in our dataset. The data were

then transformed using a centered log ratio procedure (data available in supplementary materials,

Table S1 and on Zenodo at https://doi.org/10.5281/zenodo.5889077) and analyzed by Partial Least

Squares coupled with a Discriminant analysis (PLS-DA) using the R software (v 3.4.3) and the

"RVAidememoire"   package   (Hervé,   2014)   (https://cran.r-

project.org/web/packages/RVAideMemoire/index.html). PLS-DA is an appropriate analysis when the

dataset contains fewer groups than explanatory variables, as in the case of the present quantitative

dataset. PLS-DA is a supervised technique, meaning that class memberships of the CHCs need to be

predefined. Here, we only used the first eight axes produced by the PLS to perform two PLS-DA tests:

the first analysis considered the four population samples as four different classes (LCG, GCG, LiPF,

GPF, outlined on Fig. 1). The second analysis was based on two classes only, representing the two

host types (PF, CG). The number of significant PLS components was determined by cross model

validation. We also calculated a numerical value representing the importance of each CHC variable in

the projection (VIP), i.e. VIP values larger than 1 are most influential (Hervé, 2014).

Results

Hydrocarbon profiles

We detected a complex pattern of 22 hydrocarbons and two non-identified components (I1 and I2)

in the cuticle of *I. uriae* ticks. The hydrocarbon mixture was composed of: 14 *n*-alkanes ranging from

17 to 33 carbon atoms in length, 4 monomethyl-alkanes with 17 to 24 carbon atoms and 4 alkenes

with 21 to 27 carbon atoms (Table 1).

Table 1: List of detected compounds in the cuticle of *I. uriae* adult female ticks, including name and
type.

Cuticular component	Name	Type
C_{17}	*n*-heptadecane	alkane
meC_{17}	methylheptadecane	methyl-branched alkane
C_{18}	*n*-octadecane	alkane
C_{20}	*n*-eicosane	alkane
$C_{21:1}$	heneicosene	alkene
C_{21}	*n*-heneicosane	alkane
$C_{22:1}$	docosene	alkene
C_{22}	*n*-docosane	alkane
$2meC_{22}$	2methyldocosane	methyl-branched alkane
C_{23}	*n*-tricosane	alkane
C_{24}	*n*-tetracosane	alkane
$9meC_{24}$	9methyltetracosane	methyl-branched alkane
$2meC_{24}$	2methyltetracosane	methyl-branched alkane
$C_{25:1}$	pentacosene	alkene
C_{25}	*n*-pentacosane	alkane
C_{26}	*n*-hexacosane	alkane
$C_{27:1}$	heptacosene	alkene
C_{27}	*n*-heptacosane	alkane
C_{28}	*n*-octacosane	alkane
C_{29}	*n*-nonacosane	alkane
C_{31}	*n*-hentriacontane	alkane
C_{33}	*n*-tritriacontane	alkane
I1	Non-identified	-
I2	Non-identified	-

All hydrocarbons were shared by the four tick populations, except $2meC_{24}$ and C_{33} which were only

found in puffin ticks (Fig.2).

Fig.2: Boxplot representing the mean abundance of each detected cuticular hydrocarbon from *I. uriae* ticks.
Compounds ranged from 17 to 33 carbon atoms in length. Mean abundance was calculated based on values
from four replicate tick pools. First and third quartiles, median and data points are shown. See Table 1 for CHC

abbreviations. PF refers to the Atlantic puffin (*Fratercula arctica*) and CG to the Common guillemot (*Uria*
aalge).

A high degree of variation among samples in mean abundance of cuticular components was obvious

(Fig.2). Hydrocarbons $C_{22:1}$, n-C_{29} and n-C_{31} were the most predominant, whereas many hydrocarbons

were detected in low quantities, as for example n-C_{20}, $2meC_{22}$ and n-C_{26}.

The abundance pattern of cuticular hydrocarbons varied between ticks of the two host species (Fig.3),

but also among the same host in different colony sites. Nevertheless, results of all pairwise profile

comparisons were non-significant. No CHC was specific to CG samples, although n-C_{17} and $C_{22:1}$ were

highly abundant in the GRI colony (Fig.2 and 3). In contrast, long chain cuticular hydrocarbons tended

to be present in both PF samples: n-C_{27}, n-C_{29} and n-C_{31}. CHC components tended to be in lower

overall abundance in samples from Langanes (Langanes CG), whereas samples from Lundey Island

(Lundey PF) had the highest overall abundance (Fig. 2).

Fig.3: Gas-chromatograms showing the comparison of cuticular hydrocarbon profiles for pools of 10 female ticks from Atlantic puffins (PF: black line) and
Common guillemots (CG: orange line) on Grimsey Island, Iceland.

194
195

PLS-DA analyses

Population samples

Cross model validation for population samples showed that 50.6% of samples were assigned to the

population of origin. PLS-DA analysis showed a discriminating population effect (p=0.008) separating

the two host species on first axis (Fig.4). The three first axes explained respectively 11.94, 7.99 and

3.99 % of the total variance among samples.

Fig.4: Graphical representation of the four population samples on the two first axes of the PLS-DA analysis. The
first and second axes explained respectively 11.94 and 7.99% of the total variance among samples (p=0.008).
The acronyms of the sampling locations are as follows: Grimsey - G; Langanes - L; Lundey Island - Li. PF refers
to Atlantic puffin (*Fratercula arctica*) and CG to Common guillemot (*Uria aalge*).

Hydrocarbons n-C_{33}, 2meC$_{24}$ and n-C_{29} appeared as the most influential components separating the

four population samples (Fig.5).

Fig.5: VIP classification showing the influence of CHC in discriminating the four population samples. VIP values
larger than 1 (dashed red line) are most influential. See Table 1 for CHC abbreviations.

Host types

The cross-validation test for host type revealed that 75% of the tick pools were assigned to the host

of origin (PF, CG). However, the PLS-DA analysis revealed no significant difference in CHC profiles

between the host types, although a tendency was evident (PLS-DA: p=0.064).

Discussion

Cuticle composition and function

Cuticular hydrocarbons, including linear and methyl-branched alkanes and alkenes, are

biosynthesized from an acetyl-CoA molecule in specialized secretory cells, the oenocytes which are

mainly located in the epidermis of insects (Howard and Blomquist, 2005). They are shuttled through

the hemolymph to the epicuticular surface via specialized pore canals penetrating the cuticular layers

(Holze et al, 2021). Once fixed on the arthropod cuticle within a complex mixture of alcohols, esters,

aldehydes, fatty acids, etc, hydrocarbons help prevent desiccation and serve in chemical

communication, constituting essential mediators of insect behavior (Blomquist and Bagnères, 2010).

Using GC-MS techniques to analyze extracts from female *I. uriae* ticks, we detected a complex mixture

of cuticular hydrocarbons containing linear and monomethyl-alkanes and alkenes ranging from 17 to

33 carbon atoms. The qualitative composition is similar to that reported for many other arthropods,

comprising predominantly linear alkanes (n-C_{23}, n-C_{25}, n-C_{27}, n-C_{29} and n-C_{31}) (Howard and Blomquist,

2005; Lockey, 1988). A majority of the detected cuticular hydrocarbons were already reported in

other hard ticks: *I. persulcatus* (Tkachev et al., 2000), *Amblyomma variegatum* (Estrada-Peña et al.,

1994) and *Rhipicephalus* spp. (Estrada-Peña et al., 1992). Nevertheless, alkenes were only detected

once in low quantities in *Rhipicephalus* spp. (Estrada-Peña et al., 1992). Here, we report the presence

of alkenes as heneicosene, pentacosene, heptacosene and a high quantity of docosene, particularly

in one of the CG samples. Alkenes were demonstrated to act as sex pheromones in the Alfalfa leaf-

cutter bee *Megachile rotundata* and the rove beetle *Aleochara curtula* (Paulmier et al., 1999; Peschke

and Metzler, 1987). Heneicosene is also described as an aggregation pheromone in *Drosophila* species

(Bartelt et al., 1988; Bartelt and Jackson, 1984). Pentacosene is implicated in the mating process of

different fly species as a stimulant pheromone (Uebel et al., 1978). The presence of these alkenes in

I. uriae could corresponds to the biological state of the female ticks when they were collected.

However, as mating in this tick species frequently takes place prior to feeding on the host, and in

particular when nymphal ticks emerge as flat females (McCoy and Tirard, 2002), we cannot conclude

on the mating status of collected females. Indeed, some of the profile variation in our data may be

due to the inclusion of females in different reproductive states. Experimental studies will now be

necessary to determine the potential role of the observed CHC compounds in mating behavior.

We found also large amount of long-chain hydrocarbons in PF samples (n-C_{29}, n-C_{31} and n-C_{33}). Mixed

with other compounds, n-C_{31} was demonstrated to induce copulation in males of the stable fly

Stomoxys calcitrans (Uebel et al., 1975). Other long chain hydrocarbons, including n-C_{23} to n-C_{31},

produced in large quantities and acting in combination, were found to serve in colony recognition by

bumblebees *Bombus terrestris* (Rottler et al., 2013). Studies have also highlighted the importance of

the environment in the acquisition of new hydrocarbons (d'Ettorre et al., 2006; d'Ettorre et al., 2002),

acting for example as a recognition signal of their own nest for the social wasp *Polistes metricus* Say

(1981) (Singer and Espelie, 1996; Espelie et al., 1990). Moreover, 2meC_{24}, that we only detected in PF

CHC profiles, has been shown to serve as a contact pheromone in peach twig borers *Anarsia lineatella*

(Schlamp, 2005). As puffin burrows are often deep, and densely distributed (Harris and Wanless,

2011), the production of these cuticular hydrocarbons in *I. uriae* may enable ticks to find conspecifics

in the host nesting environment. The use of this particular blend of CHCs may also help ticks to find

individuals that smell similar, favoring assortative mating (van Zweden and d'Ettorre, 2010). The

production of a specific CHC blend may not be necessary for CG ticks because guillemots breed in

extremely dense numbers on cliff ledges with no constructed nest; ticks tend to aggregate under and

around rocks on the cliff ledge such that finding a mate from the same host type may be easier

compared to ticks exploiting puffin hosts. More generally, the presence of complex compounds in the

CHC pattern of *I. uriae* suggest that chemical communication may be important in this tick species,

potentially enabling ticks to find a suitable mating partner that would enable them to maintain host-

adapted gene complexes (Sonenshine and Roe, 2014).

Alternative hypotheses to explain the origin of CHC patterns

As expected, chemical analyses revealed that each tick population had a distinct CHC profile, but that

specific CHCs were also associated with different host types. In particular, n-C$_{33}$, 2meC$_{24}$ and n-C$_{29}$

were most frequently or exclusively detected in PF populations. The quantitative variability among

detected hydrocarbons could be related to different factors.

First, aging and development have been demonstrated to impact cuticular hydrocarbon patterns in

different taxa (Desena et al., 1999; Ichinose and Lenoir, 2009). For example, aging favors the

production of longer hydrocarbon chains and decreases sexual attractiveness in *Drosophila*

melanogaster (Kuo et al., 2012). The high quantities of long-chain hydrocarbons (n-C$_{27}$, n-C$_{29}$, n-C$_{31}$)

observed in the puffin tick samples could be a consequence of age variability between host-associated

tick samples. Samples used in the present study were collected in the field and the relative age of the

specimens could not be determined. However, as tick activity is relatively synchronous with seabird

breeding, we did not expect the overall timing in adult female activity to differ in a systematic way

between ticks exploiting the different host species, nor among distinct colony locations within

Iceland.

Second, climatic conditions can also shape CHC composition. Differences in cuticular hydrocarbon composition were observed in geographically distant tick populations of *Amblyomma variegatum* (Estrada-Peña et al., 1994) and some extreme climatic parameters were shown to be correlated with methyled-alkanes in *Rhipicephalus sanguineus* (Estrada-Peña, 1993). These results suggest a potential link between variation in CHC compounds and adaptation to environmental temperatures. The different types of off-host environments used by *I. uriae* (rock or burrow) may display significant variation in terms of temperature and relative humidity due to differential exposure to climatic factors such as sun, rain and/or snow (Buckley and Buckley, 1980). These selective factors can impact tick survival and could lead to the quantitative variation in cuticular components observed in this study. Howard et al (1978) argued that the effectiveness of CHCs in preventing desiccation depends on the quantity of CHCs and that saturated CHCs are more effective components in protecting against water loss. By positioning temperature and humidity captors during one year in the off-host environment in a heterospecific seabird breeding colony in northern Norway (70°22' N, 31°10' E), we observed that the average temperature and relative humidity (HR) ranged respectively from -7.5 to 18°C and from 0 to 110% HR in breeding sites of Common guillemots, and from -7.5 to 10°C and from 80 to 105% HR in those occupied by puffins (data not shown). The micro-habitat used by CG ticks may therefore be more exposed to temperature and humidity variation than the more stable deep burrows used by PF ticks. However, the presence and abundance of saturated CHCs, such as $9meC_{24}$ and $2meC_{24}$, were not more frequent in CG samples compared to PF samples. More detailed environmental data from each location and more sampled locations are therefore necessary to more fully evaluate this hypothesis.

Third, diet also appears to be an important factor shaping cuticular hydrocarbon profiles. In the Argentine ant *Linepithema humile*, colonies eating different prey items present particular hydrocarbon profiles that include components coming directly from the prey (Liang and Silverman,

2000). In the same way, Geiselhardt et al., (2012) showed that males of the phytophagous mustard

leaf beetle *Phaedon cochleariae* preferred to mate with females reared on the same host plant

compared to females from a different host plant, even though they originated from the same

laboratory stock population. Studies have also shown that the ectoparasite *Varroa destructor* can

mimic its host's cuticular hydrocarbons, enabling the parasite to escape the hygienic behavior of the

host honeybee (Le Conte et al., 2015; Nazzi and Le Conte, 2016). In ticks, it was demonstrated that

the cuticular fatty acid profiles of *Rhipicephalus* spp. presented significant differences in abundance

according to host use (Shimshoni et al., 2013). In our system, avian erythrocyte membrane lipids

might provide a direct pathway for the tick to synthesize cuticular hydrocarbons from pre-existing

shorter-chain fatty acids and could result in the acquisition of specific hydrocarbon mixtures. This

type of acquisition could explain both the variation among ticks from different host species, and

variation among colony locations, if seabird diets shift among locations. Futures studies could

therefore be expanded to look for correlations between avian erythrocyte membrane lipid

composition and tick cuticular signatures by comparing ticks and host blood samples. In addition,

looking at patterns in tick cuticular lipids, such as fatty acids and steroids, which have been reported

as pheromones in metastriate ticks (Sonenshine, 2004; Sonenshine et al, 1985), would also be of

particular interest.

Fourth, CHCs can be altered by infection with different micro-organisms. For example, both the corn

borer (*Ostrinia nubilalis*) and the common cockchafer (Melolontha melolontha) show altered CHC

profiles when infected by fungi of the genus Beauveria (Lecuona et al, 1991). Moreover, Yoder and

Domingus (2003) demonstrated that production of the CHCs n-C_{20} and n-C_{24} by *Dermacentor variabilis*

ticks could protect against predatory behavior by ants. The abundances of n-C_{20}, n-C_{21}, n-C_{23} and n-

C_{24} in the puffin tick pools of the present study may represent different infection statuses and/or that

puffin burrows provide a less protective environment against predators, such as ants or spiders, than

stones in common guillemot colonies.

In general, making inferences on environmental versus host effects will require the examination of

ticks from additional sites and host types. However, regardless of the origin of host-associated

differences in CHC profiles of *I. uriae*, what is important to now determine is whether these

differences reinforce assortative mating patterns, favoring the divergence of sympatric populations

and the rapid formation of local host races. Future analyses should focus on the characterization and

isolation of the main components of the cuticular mixture from ticks of each seabird host type to test

for their biological activity and potential role in tick mating behavior.

Acknowledgements

We would like to thank Yann Kolbeinsson from Northeast Iceland Nature Research Centre, Iceland for sampling help. We thank "Tiques et Maladies à Tiques" Working Group of the Societé Française d'Ecologie et d'Evolution (SFE2) for stimulating discussions and support.

Funding

Funding for this study was provided by the ANR grant ESPEVEC (ANR-13-BSV7-0018-01) to KDM. MD was supported by a fellowship from the French Ministry for National Education and Research at University of Montpellier.

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article.

Supplementary data

Table S1: Relative abundance of the CHC extracts of each population replicate after log centered ratio transformation (also available on Zenodo at https://doi.org/10.5281/zenodo.5889077).

	C17	meC17	C18	C20	C21.1	C21
CHC_GPF1	1,258262	-1,675478	-1,225611	-1,358771	-0,456587	-1,093444
CHC_GPF2	-0,983004	-0,983004	-0,783199	-1,164937	-1,056892	-0,184697
CHC_GPF3	4,151361	3,580785	4,724183	4,617548	3,807260	3,540713
CHC_GPF4	3,597384	3,540178	2,810370	-5,405227	2,969631	3,222111
CHC_LIPF1	1,518413	-6,017122	1,043470	-1,007387	0,183513	-1,120153
CHC_LIPF2	0,687396	-0,588309	-0,053014	-0,504584	0,166784	-1,747215
CHC_LIPF3	2,953938	1,854215	2,570707	-4,252033	-4,252033	2,319610
CHC_LIPF4	3,334867	2,899295	1,532183	2,736876	-4,685930	2,599998
CHC_GCG1	4,385533	1,306893	1,157818	0,389237	1,944246	0,212800
CHC_GCG2	1,730616	1,703967	1,991403	1,042497	2,791342	0,217889
CHC_GCG3	5,149559	5,171603	4,695299	-4,141914	-4,141914	-4,141914
CHC_GCG4	6,607656	6,507890	-2,764376	-2,764376	-2,764376	6,667177
CHC_LCG1	4,213065	2,588977	1,956074	2,221551	-5,462891	1,390069
CHC_LCG2	1,933896	1,561455	1,352674	1,168282	2,599910	1,493046
CHC_LCG3	5,173815	-3,295400	-3,295400	-3,295400	-3,295400	5,004318
CHC_LCG4	-2,395630	-2,395630	-2,395630	-2,395630	-2,395630	6,592884

	C22.1	C22	meC22	C23	C24	9meC24
CHC_GPF1	1,50623611	-0,50555381	-1,70555949	0,12096462	-0,42768644	-1,2206834
CHC_GPF2	1,18425441	-0,49071836	-0,98066082	0,94030298	-0,28189781	-1,02914483
CHC_GPF3	4,32918786	4,60171239	4,02667532	-4,77602802	-4,77602802	-4,77602802
CHC_GPF4	2,85991037	2,24131129	3,95590755	3,57761114	-5,4052267	2,55126443
CHC_LIPF1	2,6847376	0,88062159	-6,01712163	0,22189045	0,56501804	-0,13693536
CHC_LIPF2	2,96343585	0,37650416	-5,92683159	0,07138213	0,62140857	-0,18069582
CHC_LIPF3	-4,25203345	-4,25203345	-4,25203345	2,48114139	2,91217821	-4,25203345
CHC_LIPF4	-4,68592953	2,30293014	-4,68592953	2,17284599	2,95775455	-4,68592953
CHC_GCG1	4,72183226	2,16796305	-5,97456355	0,50141786	1,61476737	1,69619514
CHC_GCG2	4,63765818	1,96470376	-6,27678757	0,9386865	1,89458577	1,80575829
CHC_GCG3	-4,14191373	-4,14191373	4,50899999	4,55481651	5,23921344	4,3019409
CHC_GCG4	-2,76437623	-2,76437623	-2,76437623	6,56677188	-2,76437623	-2,76437623
CHC_LCG1	5,14840786	2,88187094	-5,46289066	2,15974193	2,64600173	1,30457536
CHC_LCG2	5,19919581	2,44288367	-5,83634374	0,96429749	1,55031557	1,00827119
CHC_LCG3	-3,2954001	-3,2954001	-3,2954001	4,10398495	5,06658076	-3,2954001
CHC_LCG4	-2,39563008	-2,39563008	-2,39563008	6,38876341	-2,39563008	-2,39563008

	2meC24	C25.1	C25	I1	C26	I2
CHC_GPF1	-1,49255919	0,710483	1,09996476	0,56276083	-0,17627985	0,07387322
CHC_GPF2	-0,85129625	0,41550683	1,72274199	0,15326967	-0,25663413	-0,00255114
CHC_GPF3	3,04100103	-4,77602802	-4,77602802	3,46602618	3,89103385	-4,77602802
CHC_GPF4	2,74237448	-5,4052267	4,27702093	3,00740146	-5,4052267	3,29174679
CHC_LIPF1	-6,01712163	-0,1488945	1,29293382	1,12637653	0,69030705	-6,01712163
CHC_LIPF2	-5,92683159	-0,64688596	1,31627079	0,90987066	0,41085836	-5,92683159
CHC_LIPF3	-4,25203345	-4,25203345	4,71776728	3,3029661	2,56104558	-4,25203345
CHC_LIPF4	-4,68592953	-4,68592953	4,6771672	-4,68592953	2,68595365	-4,68592953
CHC_GCG1	-5,97456355	-0,42396916	1,22349549	1,72540467	1,38243536	-1,39612756
CHC_GCG2	-6,27678757	-0,2065063	1,72213495	1,06728457	1,33585569	-0,21747187
CHC_GCG3	-4,14191373	-4,14191373	5,71433159	-4,14191373	4,75085336	-4,14191373
CHC_GCG4	-2,76437623	-2,76437623	7,51121475	7,01077959	-2,76437623	-2,76437623
CHC_LCG1	-5,46289066	-5,46289066	3,01341895	2,82230157	2,69452068	-5,46289066
CHC_LCG2	-5,83634374	2,0586822	1,72477962	2,25712355	1,39064679	-5,83634374
CHC_LCG3	-3,2954001	-3,2954001	7,60591058	5,62063417	5,2526945	-3,2954001
CHC_LCG4	-2,39563008	-2,39563008	7,96176069	-2,39563008	7,19765085	-2,39563008

	C27.1	C27	C28	C29	C31	C33
CHC_GPF1	-0,00016215	1,30711449	0,33827026	2,04171998	1,59307887	0,72564782
CHC_GPF2	-0,19069859	1,69885837	-0,1473274	1,99452145	1,07445349	0,20275287
CHC_GPF3	-4,77602802	-4,77602802	4,75882151	-4,77602802	-4,77602802	-4,77602802
CHC_GPF4	4,00281861	-5,4052267	-5,4052267	-5,4052267	-5,4052267	-5,4052267
CHC_LIPF1	1,02468304	2,57346606	1,2522136	4,63157554	4,23145935	2,56117937
CHC_LIPF2	-1,29244538	2,91653471	0,74631061	4,80493688	4,29222175	2,5097294
CHC_LIPF3	4,42572948	-4,25203345	-4,25203345	6,29373406	6,22187233	4,15746354
CHC_LIPF4	3,97386234	-4,68592953	-4,68592953	5,79395404	5,57504998	3,61655868
CHC_GCG1	-2,31093524	1,17815423	0,96870906	1,45238521	-5,97456355	-5,97456355
CHC_GCG2	-2,63802842	1,37672655	1,18012447	0,76792459	-6,27678757	-6,27678757
CHC_GCG3	4,78876251	-4,14191373	-4,14191373	-4,14191373	4,9694991	-4,14191373
CHC_GCG4	6,12290594	-2,76437623	-2,76437623	-2,76437623	-2,76437623	-2,76437623
CHC_LCG1	-5,46289066	3,03426816	2,96262862	2,66565265	-5,46289066	-5,46289066
CHC_LCG2	-5,83634374	1,89920237	2,48119674	1,9322039	-5,83634374	-5,83634374
CHC_LCG3	6,05677419	-3,2954001	-3,2954001	-3,2954001	5,54628953	-3,2954001
CHC_LCG4	7,47336812	-2,39563008	-2,39563008	-2,39563008	7,50691468	-2,39563008

References

Andersen, S.O., 1979. Biochemistry of Insect Cuticle. Annu. Rev. Entomol. 24, 29–59. https://doi.org/10.1146/annurev.en.24.010179.000333

Bagnères, A.-G., Killian, A., Clement, J.-L., Lange, C., 1991. Interspecific recognition among termites of the genusReticulitermes: Evidence for a role for the cuticular hydrocarbons. J. Chem. Ecol. 17, 2397–2420. https://doi.org/10.1007/BF00994590

Bartelt, R.J., Armold, M.T., Schaner, A.M., Jackson, L.L., 1986. Comparative analysis of cuticular hydrocarbons in the Drosophila virilis species group. Comp. Biochem. Physiol. Part B Comp. Biochem. 83, 731–742. https://doi.org/10.1016/0305-0491(86)90138-0

Bartelt, R.J., Jackson, L.L., 1984. Hydrocarbon Component of the Drosophila virilis (Diptera: Drosophilidae) Aggregation Pheromone: (Z)-10-Heneicosene. Ann. Entomol. Soc. Am. 77, 364–371. https://doi.org/10.1093/aesa/77.4.364

Bartelt, R.J., Schaner, A.M., Jackson, L.L., 1988. Aggregation pheromones inDrosophila borealis andDrosophila littoralis. J. Chem. Ecol. 14, 1319–1327.

Blomquist, G.J., Bagnères, A.-G., 2010. Insect Hydrocarbons: Biology, Biochemistry, and Chemical Ecology. Cambridge University Press.

Buckley, F.G., Buckley, P.A., 1980. Habitat Selection and Marine Birds, in: Behavior of Marine Animals. Springer, Boston, MA, pp. 69–112. https://doi.org/10.1007/978-1-4684-2988-6_3

Desena, M.L., Edman, J.D., Clark, J.M., Symington, S.B., Scott, T.W., 1999. Aedes aegypti (Diptera: Culicidae) Age Determination by Cuticular Hydrocarbon Analysis of Female Legs. J. Med. Entomol. 36, 824–830. https://doi.org/10.1093/jmedent/36.6.824

D'Ettorre, P., Mondy, N., Lenoir, A., Errard, C., 2002. Blending in with the crowd: social parasites integrate into their host colonies using a flexible chemical signature. Proc. R. Soc. Lond. B Biol. Sci. 269, 1911–1918. https://doi.org/10.1098/rspb.2002.2110

D'Ettorre, P., Wenseleers, T., Dawson, J., Hutchinson, S., Boswell, T., Ratnieks, F.L.W., 2006. Wax combs mediate nestmate recognition by guard honeybees. Anim. Behav. 71, 773–779. https://doi.org/10.1016/j.anbehav.2005.05.014

Dietrich, M., Beati, L., Elguero, E., Boulinier, T., McCoy, K.D., 2013. Body size and shape evolution in host races of the tick Ixodes uriae. Biol. J. Linn. Soc. 108, 323–334. https://doi.org/10.1111/j.1095-8312.2012.02021.x

Dietrich, M., Gómez-Díaz, E., McCoy, K.D., 2011. Worldwide Distribution and Diversity of Seabird Ticks: Implications for the Ecology and Epidemiology of Tick-Borne Pathogens. Vector-Borne Zoonotic Dis. 11, 453–470. https://doi.org/10.1089/vbz.2010.0009

Dietrich, M., Lobato, E., Boulinier, T., McCoy, K.D., 2014. An experimental test of host specialization in a ubiquitous polar ectoparasite: a role for adaptation? J. Anim. Ecol. 83, 576–587. https://doi.org/10.1111/1365-2656.12170

Espelie, K.E., Wenzel, J.W., Chang, G., 1990. Surface lipids of social waspPolistes melricus say and its nest and nest pedicel and their relation to nestmate recognition. J. Chem. Ecol. 16, 2229–2241. https://doi.org/10.1007/BF01026933

Estrada-Peña, A., 1993. Climate and cuticular hydrocarbon variation inRhipicephalus sanguineus ticks (Acari: Ixodidae). Parasitol. Res. 79, 512–516. https://doi.org/10.1007/BF00931594

Estrada-Peña, A., CASTELLÁ, J., MOREL, P.C., 1994. Cuticular Hydrocarbon Composition, Phenotypic Variability, and Geographic Relationships in Allopatric Populations of Amblyomma variegatum (Acari: Ixodidae) from Africa and the Caribbean. J. Med. Entomol. 31, 534–544.

Estrada-Peña, A., Estrada-Peña, R., Peiró, J.M., 1992. Differentiation of Rhipicephalus Ticks (Acari: Ixodidae) by Gas Chromatography of Cuticular Hydrocarbons. J. Parasitol. 78, 982. https://doi.org/10.2307/3283217

Estrada-Peña, A., Gray, J.S., Kahl, O., 1996. Variability in cuticular hydrocarbons and phenotypic
discrimination of Ixodes ricinus populations (Acarina: Ixodidae) from Europe. Exp. Appl. Acarol. 20,
457–466. https://doi.org/10.1007/BF00053309
Estrada-Peña, A., Guglielmone, A.A., Mangold, A.J., Castellá, J., 1993. Patterns of cuticular hydrocarbon
variation and genetic similarity between natural populations of Amblyomma cajennense (Acari:
Ixodidae). Acta Trop. 55, 61–78. https://doi.org/10.1016/0001-706X(93)90049-H
Filshie, B.K., 1982. Fine Structure of the Cuticle of Insects and Other Arthropods, in: Insect Ultrastructure.
Springer, Boston, MA, pp. 281–312. https://doi.org/10.1007/978-1-4615-7266-4_10
Geiselhardt, S., Otte, T., Hilker, M., 2012. Looking for a similar partner: host plants shape mating preferences
of herbivorous insects by altering their contact pheromones. Ecol. Lett. 15, 971–977.
https://doi.org/10.1111/j.1461-0248.2012.01816.x
Gibbs, A., Mousseau, T.A., Crowe, J.H., 1991. Genetic and acclimatory variation in biophysical properties of
insect cuticle lipids. Proc. Natl. Acad. Sci. 88, 7257–7260. https://doi.org/10.1073/pnas.88.16.7257
Harris, M.P., Wanless, S., 2011. The Puffin. Bloomsbury Publishing.
Hervé, M., 2014. Aide-mémoire de statistique appliquée à la biologie. Constr Son Étude Anal Résultats À Aide
Logiciel R Version 5.
Holze, H., Schrader, L., Buellesbach, J., 2021. Advances in deciphering the genetic basis of insect cuticular
hydrocarbon biosynthesis and variation. Heredity 126, 219–234. https://doi.org/10.1038/s41437-
020-00380-y
Howard, R.W., Blomquist, G.J., 2005. Ecological, Behavioral, and Biochemical Aspects of Insect Hydrocarbons.
Annu. Rev. Entomol. 50, 371–393. https://doi.org/10.1146/annurev.ento.50.071803.130359
Howard, R.W., McDaniel, C.A., Blomquist, G.J., 1978. Cuticular hydrocarbons of the eastern subterranean
termite,Reticulitermes flavipes (Kollar) (Isoptera: Rhinotermitidae). J. Chem. Ecol. 4, 233–245.
https://doi.org/10.1007/BF00988058
Hunt, L.M., 1986. Differentiation between three species of Amblyomma ticks (Acari: Ixodidae) by analysis of
cuticular hydrocarbons. Ann. Trop. Med. Parasitol. 80, 245–249.
Ichinose, K., Lenoir, A., 2009. Ontogeny of hydrocarbon profiles in the ant Aphaenogaster senilis and effects
of social isolation. C. R. Biol. 332, 697–703. https://doi.org/10.1016/j.crvi.2009.04.002
Jallon, J.-M., David, J.R., 1987. Variation in Cuticular Hydrocarbons Among the Eight Species of the Drosophila
melanogaster Subgroup. Evolution 41, 294–302. https://doi.org/10.2307/2409139
KRUGER, E.L., PAPPAS, C.D., HOWARD, R.W., 1991. Cuticular Hydrocarbon Geographic Variation Among
Seven North American Populations of Aedes albopictus (Diptera: Culicidae). J. Med. Entomol. 28,
859–864.
Kuo, T.-H., Yew, J.Y., Fedina, T.Y., Dreisewerd, K., Dierick, H.A., Pletcher, S.D., 2012. Aging modulates cuticular
hydrocarbons and sexual attractiveness in Drosophila melanogaster. J. Exp. Biol. 215, 814–821.
https://doi.org/10.1242/jeb.064980
Le Conte, Y, Huang, Y. Z., Roux, M., Zeng, Z.-J., Christidès, J.-P., Bagnères, A.-G., 2015. Varroa destructor
changes its cuticular hydrocarbons to mimic new hosts. Biol. Lett. 11, 20150233.
https://doi.org/10.1098/rsbl.2015.0233
Lecuona, R., Riba, G., Cassier, P., Clement, J.L., 1991. Alterations of insect epicuticular hydrocarbons during
infection with Beauveria bassiana or B. brongniartii. Journal of Invertebrate Pathology 58, 10–18.
https://doi.org/10.1016/0022-2011(91)90156-K
Léger, E., Vourc'h, G., Vial, L., Chevillon, C., McCoy, K.D., 2013. Changing distributions of ticks: causes and
consequences. Exp. Appl. Acarol. 59, 219–244. https://doi.org/10.1007/s10493-012-9615-0
Leonovich, S.A., 2004. Phenol and lactone receptors in the distal sensilla of the Haller's organ in Ixodes
ricinus ticks and their possible role in host perception. Exp. Appl. Acarol. 32, 89–102.
https://doi.org/10.1023/B:APPA.0000018200.24760.78
Liang, D., Silverman, J., 2000. "You are what you eat": Diet modifies cuticular hydrocarbons and nestmate
recognition in the Argentine ant, Linepithema humile. Naturwissenschaften 87, 412–416.
https://doi.org/10.1007/s001140050752

Lockey, K.H., 1988. Lipids of the insect cuticle: origin, composition and function. Comp. Biochem. Physiol.
Part B Comp. Biochem. 89, 595–645.
McCoy, K.D., Boulanger, N. (Eds.), 2015. Tiques et maladies à tiques : biologie, écologie évolutive,
épidémiologie, Didactiques. IRD, Marseille.
McCoy, K.D., Boulinier, T., Tirard, C., Michalakis, Y., 2001. Host specificity of a generalist parasite: genetic
evidence of sympatric host races in the seabird tick Ixodes uriae. J. Evol. Biol. 14, 395–405.
https://doi.org/10.1046/j.1420-9101.2001.00290.x
McCoy, K.D., Chapuis, E., Tirard, C., Boulinier, T., Michalakis, Y., Bohec, C.L., Maho, Y.L., Gauthier-Clerc, M.,
2005. Recurrent evolution of host-specialized races in a globally distributed parasite. Proc. R. Soc. B
Biol. Sci. 272, 2389–2395. https://doi.org/10.1098/rspb.2005.3230
McCoy, K.D., Léger, E., Dietrich, M., 2013. Host specialization in ticks and transmission of tick-borne diseases:
a review. Front. Cell. Infect. Microbiol. 3. https://doi.org/10.3389/fcimb.2013.00057
McCoy, K.D., Tirard, C., 2002. Reproductive Strategies of the Seabird Tick Ixodes uriae (Acari: Ixodidae). J.
Parasitol. 88, 813–816. https://doi.org/10.1645/0022-3395(2002)088[0813:RSOTST]2.0.CO;2
McCoy, K.D., Tirard, C., Michalakis, Y., 2003. Spatial genetic structure of the ectoparasite Ixodes uriae within
breeding cliffs of its colonial seabird host. Heredity 91, 422–429.
https://doi.org/10.1038/sj.hdy.6800339
Nazzi, F., Le Conte, Y., 2016. Ecology of *Varroa destructor*, the Major Ectoparasite of the Western Honey Bee,
Apis mellifera. Annu. Rev. Entomol. 61, 417–432.
https://doi.org/10.1146/annurev-ento-010715-023731
Paulmier, I., Bagnères, A.-G., Afonso, C.M., Dusticier, G., Rivière, G., Clément, J.-L., 1999. Alkenes as a sexual
pheromone in the alfalfa leaf-cutter bee Megachile rotundata. J. Chem. Ecol. 25, 471–490.
Peschke, K., Metzler, M., 1987. Cuticular hydrocarbons and female sex pheromones of the rove beetle,
Aleochara curtula (Goeze) (Coleoptera:Staphylinidae). Insect Biochem. 17, 167–178.
https://doi.org/10.1016/0020-1790(87)90157-0
Randolph, S.E., 1998. Ticks are not Insects: Consequences of Contrasting Vector Biology for Transmission
Potential. Parasitol. Today 14, 186–192. https://doi.org/10.1016/S0169-4758(98)01224-1
Randolph, S.E., Storey, K., 1999. Impact of Microclimate on Immature Tick-Rodent Host Interactions (Acari:
Ixodidae): Implications for Parasite Transmission. J Med Entomol 36, 741–748.
https://doi.org/10.1093/jmedent/36.6.741
Rottler, A.-M., Schulz, S., Ayasse, M., 2013. Wax Lipids Signal Nest Identity in Bumblebee Colonies. J. Chem.
Ecol. 39, 67–75. https://doi.org/10.1007/s10886-012-0229-0
Schlamp, K.K., 2005. Contact pheromone components and diel periodicity of sexual communication in peach
twig borers, Anarsia lineatella (Lepidoptera: Gelechiidae) (PhD Thesis). Biological Sciences
Department-Simon Fraser University.
Schöni, R., Hess, E., Blum, W., Ramstein, K., 1984. The aggregation-attachment pheromone of the tropical
bont tick Amblyomma variegatum Fabricius (Acari, Ixodidae): Isolation, identification and action of its
components. J. Insect Physiol. 30, 613–618. https://doi.org/10.1016/0022-1910(84)90045-3
Shimshoni, J.A., Erster, O., Rot, A., Cuneah, O., Soback, S., Shkap, V., 2013. Cuticular fatty acid profile analysis
of three Rhipicephalus tick species (Acari: Ixodidae). Exp. Appl. Acarol. 61, 481–489.
https://doi.org/10.1007/s10493-013-9713-7
Simmons, L.W., Thomas, M.L., Gray, B., Zuk, M., 2014. Replicated evolutionary divergence in the cuticular
hydrocarbon profile of male crickets associated with the loss of song in the Hawaiian archipelago. J.
Evol. Biol. n/a-n/a. https://doi.org/10.1111/jeb.12478
Singer, T.L., Espelie, K.E., 1996. Nest surface hydrocarbons facilitate nestmate recognition for the social
wasp,Polistes metricus Say (Hymenoptera: Vespidae). J. Insect Behav. 9, 857–870.
https://doi.org/10.1007/BF02208974
Sonenshine, D.E., 2004. Pheromones and other semiochemicals of ticks and their use in tick control.
Parasitology 129, S405–S425. https://doi.org/10.1017/S003118200400486X
Sonenshine, D.E., Roe, R.M. (Eds.), 2014. Biology of ticks, 2nd ed. ed. Oxford University Press, New York.

Sonenshine, D.E., Silverstein, R.M., Brossut, R., Davis, E.E., Taylor, D., Carson, K.A., Homsher, P.J., Wang, V.B.,
1985. Genital sex pheromones of ixodid ticks: 1. Evidence of occurrence in anterior reproductive tract
of American dog tick,Dermacentor variabilis (Say) (Acari: Ixodidae). J Chem Ecol 11, 1669–1694.
https://doi.org/10.1007/BF01012120
Tkachev, A.V., Dobrotvorsky, A.K., Vjalkov, A.I., Morozov, S.V., 2000. Chemical composition of lipophylic
compounds from the body surface of unfed adult Ixodes persulcatus ticks (Acari: Ixodidae). Exp. Appl.
Acarol. 24, 145–158. https://doi.org/10.1023/A:1006430323587
Uebel, E.C., Schwarz, M., Sonnet, P.E., Miller, R.W., Menzer, R.E., 1978. Evaluation of the mating stimulant
pheromones of Fannia canicularis, F. pusio, and F. femoralis as attractants. Fla. Entomol. 139–143.
Uebel, E.C., Sonnet, P.E., Bierl, B.A., Miller, R.W., 1975. Sex pheromone of the stable flyl: Isolation and
preliminary identification of compounds that induce mating strike behavior. J. Chem. Ecol. 1, 377–
385.
van Zweden, J.S., d'Ettorre, P., 2010. Nestmate recognition in social insects and the role of hydrocarbons.
Insect Hydrocarb. Biol. Biochem. Chem. Ecol. 11, 222–243.
Wyatt, T.D., 2010. Pheromones and signature mixtures: defining species-wide signals and variable cues for
identity in both invertebrates and vertebrates. J. Comp. Physiol. A 196, 685–700.
https://doi.org/10.1007/s00359-010-0564-y
Yoder, J.A., Domingus, J.L., 2003. Identification of hydrocarbons that protect ticks (Acari: Ixodidae) against
fire ants (Hymenoptera: Formicidae), but not lizards (Squamata: Polychrotidae), in an allomonal
defense secretion. International Journal of Acarology 29, 87–91.
https://doi.org/10.1080/01647950308684326