2   Are Pacific Chorus Frogs (*Pseudacris regilla*) Resistant to Tetrodotoxin (TTX)?

3   Characterizing Potential TTX Exposure and Resistance in an Ecological Associate of

4   Pacific Newts (*Taricha*)

Katherine O. Montana[1,2,3], Valeria Ramírez Castañeda[1], Rebecca D. Tarvin[1,3]

[1]*Museum of Vertebrate Zoology and Department of Integrative Biology, University of*

California, Berkeley, Berkeley, California, USA 94720

[2]*Present address: Department of Biology, San Francisco State University, San*

Francisco, California, USA 94132 and *Institute for Biodiversity Science and*

Sustainability, California Academy of Sciences, San Francisco, California, USA 94118

[3]*Corresponding authors. kmontana@berkeley.edu, rdtarvin@berkeley.edu*

Abstract.—Animals that frequently encounter toxins often select for mechanisms

of toxin resistance. Both predators that consume toxic prey and organisms in physical

contact with a toxin or pollutant in their environment may experience natural selection

for resistance. Based on field observations that Pacific Chorus Frogs (*Pseudacris*

regilla) sometimes eat and mistakenly amplect tetrodotoxin (TTX)-defended *Taricha*

newts, we predicted that *P. regilla* may possess resistance to TTX. We tested this

prediction by comparing the amino acid sequences of the molecular target of TTX, the

muscle voltage-gated sodium channel gene SCN4A (*Na$_V$1.4*), in populations of *P. regilla*

that are sympatric and allopatric with *Taricha*. We identified a single substitution in

Na$_V$1.4 of *P. regilla* in a conserved site near the pore loop where TTX binds. Although

the role of this site in TTX resistance has not been functionally assessed, both allopatric

and sympatric *P. regilla* had this substitution, suggesting that it may be unrelated to TTX

exposure from *Taricha*. Thus, there is no conclusive evidence that *P. regilla* has

selected for TTX resistance encoded by amino acid substitutions in this domain. In

addition, California occurrence data from the last 50 years indicate that *Taricha* activity

peaks in January while the activity of *P. regilla* peaks in April. These relatively distinct

activity patterns suggest that *P. regilla* may not be exposed to levels of TTX from

Taricha that are high enough to select for mutations in the sodium channel.

Nevertheless, other unidentified mechanisms of TTX resistance could be present in *P.*

regilla and other species that are sympatric with toxic newts.

Key words: Toxin resistance; California; Environmental toxins; Target-site

insensitivity; Salamanders; Chemical ecology

Resumen.— Los animales que tienen contacto frecuente con toxinas suelen

desarrollar mecanismos de resistencia a las mismas. Tanto los depredadores que

consumen presas tóxicas como los organismos en contacto cercano con una toxina o

contaminante en su entorno pueden experimentar una presión de selección que los

lleva a evolucionar resistencia a toxinas. Basándose en las observaciones de que las

ranas coro del Pacífico (*Pseudacris regilla*) a veces comen por error y/o amplexan

salamandras del género *Taricha* que poseen tetrodotoxina (TTX), se planteó la

hipótesis de que *P. regilla* podría poseer resistencia a la TTX. Esta predicción fue

probada comparando las secuencias de aminoácidos en el loop del poro del dominio IV

en el gen del canal de sodio voltaje dependiente muscular SCN4A (proteína *Nav1.4*) en

poblaciones de *P. regilla* que son simpátricas y alopátricas con *Taricha*. Se identificó

una única sustitución en el *Nav1.4* de P. regilla en un sitio conservado cerca del loop

del poro donde se une la TTX. Aunque el papel de este sitio en la resistencia a la TTX

no ha sido evaluado funcionalmente, tanto el *P. regilla* alopátrico como el simpátrico

tienen esta sustitución, lo que sugiere que no está relacionado con la exposición a la

TTX secretada por *Taricha*. Por lo tanto, no hay evidencias concluyentes de que *P.*

regilla haya evolucionado resistencia a la TTX por medio de sustituciones de

aminoácidos en este dominio. Por otro lado, los datos de ocurrencia en California de la

actividad de *Taricha* en los últimos 50 años indican alcanza su máximo en enero,

mientras que la de *P. regilla* lo hace en abril. Estos patrones de actividad relativamente

distintos sugieren que *P. regilla* puede no estar expuesta a niveles de TTX provenientes

de *Taricha* que sean lo suficientemente altos como para inducir la evolución de

mutaciones en el canal de sodio. Sin embargo, otros mecanismos no identificados de

resistencia a la TTX podrían estar presentes en *P. regilla* y en otras especies

simpáticas a los salamandras tóxicas.

Palabras clave: Resistencia a las toxinas; California; Toxinas ambientales;

Insensibilidad en el sitio de union; Salamandras; Ecología química

Newts of the genus *Taricha* (*Taricha torosa*, *T. granulosa*, *T. sierrae,* and *T.*

rivularis, herein collectively referred to as *Taricha*) wield tetrodotoxin (TTX) on their skin

(Brodie et al., 1974; Mailho-Fontana et al., 2019; Reimche et al., 2020) that likely

functions as an anti-predator defense (Brodie et al., 2002; Williams et al., 2010;

Bucciarelli et al., 2017). TTX is a neurotoxin that blocks voltage-gated sodium channels

by binding to highly conserved regions of the channel via hydrophobic forces and

hydrogen bonds, fitting in the opening of the channel and blocking proper transport of

Na$^+$ ions (Kao and Levinson, 1986). TTX binding prevents action potentials from

propagating, resulting in cessation of nerve signals (Kao and Levinson, 1986). In

vertebrate animals lacking TTX resistance, ingested TTX can temporarily paralyze or, in

some cases, kill the organism. The poison usually becomes fatal when respiratory

muscles fail, resulting in suffocation (Brodie, 1968). The San Francisco Bay Area in

California is one of two known hotspots of the arms race between newts and garter

snake predators (Brodie et al., 2002; but see Bucciarelli et al. 2022). Here we explore

the idea that high levels of TTX defenses in newts may select for resistance in other

sympatric organisms outside the predator-prey dynamic, namely the Pacific Chorus

Frog, *Pseudacris regilla.*

Taricha and their relatives possess amino acid changes in the muscle voltage-

gated sodium channel $Na_V1.4$ that help them resist their own TTX defenses (Hanifin and

Gilly, 2015; Gendreau et al., 2021). Amino acid substitutions in voltage-gated sodium

channels (VGSCs) alter the shape of the pore loop (p-loop) and the surrounding area

which prevents TTX from binding, especially in domain IV (DIV) of $Na_V1.4$ (Geffeney et

al., 2005; Tikhonov and Zhorov, 2005; Jost et. al, 2008; Hanifin and Gilly, 2015). Amino

acid substitutions in the p-loop can affect the function of voltage-gated sodium channels

by altering sodium ion selectivity or gating properties; hence, trade-offs may exist

between TTX resistance and organismal function (Chiamvimonvat et al., 1996; Lee et

al., 2011; Hague et al., 2018; but see Moniz et al., 2021). Thus, amino acid changes are

often confined to specific conserved regions that can provide resistance while allowing

the organism to maintain sodium channel functionality (Feldman et al., 2012). The

presence of amino acid substitutions that alter toxin binding is a mechanism known as

target-site insensitivity and is the focus of this study.

TTX from *Taricha* skin has been shown to enter waterways and affect the

behavior of organisms present in the same aquatic systems. Larval *T. torosa* use TTX

as a cue to detect and avoid adult *T. torosa* that sometimes cannibalize larvae (Zimmer

et al., 2006)*.* TTX in water can also slow dragonfly nymph consumption of *T. torosa*

larvae (Bucciarelli and Kats, 2015). The New Zealand Mud Snail (*Potamopyrgus*

antipodarum) moved out of its stream environment when encountering biologically

relevant levels of TTX or chemical cues from *T. torosa* containing TTX (Ota et al.,

2018). Water containing TTX also increases newt trematode parasite mortality (Calhoun

et al., 2017). Thus, naturally occurring levels of TTX produced by *T. torosa* can affect

other organisms in the same aquatic environment.

109   While *P. regilla* do not regularly prey upon *Taricha* (or vice versa), they likely

come into contact with TTX in pond water shared with *Taricha* or via direct contact with

Taricha, sometimes in mistaken amplexus behavior or even attempted predation (Fig.

1). Given that *P. regilla* have permeable skin, we expected that they would be exposed

to TTX through physical contact with newts or by occupying pond water that contains

newts. We hypothesized that TTX exposure has selected for TTX resistance in *P. regilla*

populations that are microsympatric with *Taricha*. We aimed to test whether the

sympatry of *Taricha* newts with *Pseudacris regilla* has selected for target-site

insensitivity in the domain IV of the skeletal muscle voltage-gated sodium channel

($Na_V1.4$) of *P. regilla*.

119   Our study centered on California populations, examining *Taricha* and *P. regilla*

that do and do not overlap geographically in the present. We conducted demographic

surveys of *Taricha granulosa, T. torosa,* and *P. regilla* at three San Francisco Bay Area

ponds where the species co-occur to estimate the extent to which *P. regilla* may

encounter TTX. We supplemented our surveys with occurrence data available through

the Global Biodiversity Information Facility (GBIF) to better estimate overlap of annual

activity patterns and to determine the degree of sympatry between *Taricha* spp. and *P.*

regilla. To determine whether *P. regilla* has structural modifications in the muscle

voltage-gated sodium channel *$Na_V1.4$* that may provide TTX resistance, we sequenced

and reviewed amino acid residues encoded by the SCN4A gene. Some amino acid

substitutions in domain IV of SCN4A are known to confer resistance to TTX and have

been found in many organisms that either have TTX defense or consume TTX-

defended prey (Geffeney et al., 2005; Venkatesh et al., 2005; Jost et al., 2008; Feldman

et al., 2012; Hanifin and Gilly, 2015; Gendreau et al., 2021). We hypothesized that if the

presence of *Taricha* selects for TTX resistance in *P. regilla,* populations sympatric with

Taricha would have mutations in the domain IV p-loop of *Na$_V$1.4* while populations that

are allopatric would not. Our work provides insight into the potential cascading

ecological consequences and unintended effects of chemical defenses.

MATERIALS AND METHODS

Field Work.—We studied *Taricha* and *P. regilla* at four San Francisco Bay Area

ponds. We conducted surveys at three of these sites where *Taricha granulosa, T.*

torosa, and *P. regilla* co-occur: two sites in Briones Regional Park in Martinez, Contra

Costa County, CA (Old Briones Road Trail 1 [OBRT 1] at 37.94571, -122.13395, and

OBRT 2 at 37.94352, -122.14102, named after the trail closest to the sites) and one site

at the Japanese Pool (37.87440, -122.23760) at the University of California Botanical

Garden (UCBG) in Berkeley, Alameda County, CA (Fig. S1). OBRT 1 and OBRT 2 are

in an unshaded area of grazed open grassland that lies within an oak woodland habitat,

and they connect to each other when water levels are high. The UCBG Japanese Pool

is a human-constructed pool with a concrete floor and sides, located under tree cover

within the visitor area of the UCBG (Fig. S1). We selected these three ponds because

we knew them to contain populations of *P. regilla* as well as *Taricha torosa* and *Taricha*

granulosa. Close to our field sites, *P. regilla* and *T. torosa* were known to be prey of

Thamnophis (Preston and Johnson, 2012), supporting cohabitation in the San Francisco

Bay Area. We also collected specimens from a fourth pond (Mountain Lake) located in

the Presidio of San Francisco (37.788496, -122.468605) that has historically contained

P. regilla but no *Taricha* (records show absence of *Taricha* since at least 1966; pers.

comm. J. Young, Presidio Trust). As of 2021 (shortly after our surveys), efforts were

being made to reintroduce *Taricha* to the Presidio waterways. Mountain Lake is a

natural lake surrounded by grass and receives run-off from the nearby Presidio Golf

Course (Fig. S1). From October 2019 to February 2020, we conducted six surveys at

the two Briones ponds and five surveys at UCBG, with surveys at each site occurring

three to four weeks apart from one another. We were unable to gather abundance data

for more than this time period because the parks were closed from February until

October 2020 during the early months of the COVID-19 pandemic.

Two researchers (usually K. Montana and V. Ramírez-Castañeda) conducted

surveys in the morning or early afternoon (between 8:00 AM and 1:30 PM). During each

survey, surveyors split the pond perimeter into 7.62-m (25-feet) sections, and circled the

pond on foot, counting how many individuals of each species were visible in the water

from the shore within each transect. We note that surveyors were only able to count the

number of visible newts. OBRT 1 and OBRT 2 were murky with plants and algae in

which newts tended to hide. Although the surface of the Japanese Pool at UCBG was

glassy, the pool was filled with plants that likely obscured observations. Surveyors also

counted the numbers of visible, individual adult *Pseudacris regilla*; *P. regilla* that were

calling but not visible were not counted because surveyors could not ascertain if

multiple calls were coming from a single or a few individuals. For the purposes of this

study, we did not differentiate between *T. torosa* and *T. granulosa* when we recorded

their occurrences because they both contribute to the presence of TTX in pond water,

and we could not easily identify individuals to species from the pond shores. We

counted large (~5 cm or longer) *Taricha* larvae, adult *Taricha,* and *Taricha torosa* egg

masses, which are conspicuous groups of eggs known to possess TTX via vertical

transfer from the parent (Gall et al., 2012). The much smaller *T. granulosa* eggs are

typically hidden in the substrate and are more difficult to see; hence we did not include

them in our survey. To simplify our survey protocol, we counted larval *Taricha*, which

also have some amount of TTX (Gall et al., 2011), together with adults. We plotted

survey results using R version 4.1.1 and the packages ggplot2 v.3.3.5 and ggpubr

v.0.4.0 (Wickham, 2016; Kassambara, 2020; R Core Team, 2021) (Fig. S2).

After COVID-19 conditions permitted field work to continue in October 2020, we

collected one *P. regilla* from the dry bed of OBRT 2 and eight *P. regilla* from the east

arm of Mountain Lake in the Presidio of San Francisco. We surveyed OBRT 1 and

UCBG at the same time but did not find any *P. regilla* at either location; OBRT 1 was

dry.

GBIF Occurrences and Sympatry.—We estimated seasonal patterns of *P. regilla*

and *Taricha* activity across the state of California and within the San Francisco Bay

Area (N 38.66226, S 37.09298, E -121.12701, W -123.23579) by reviewing annual

patterns of GBIF occurrence data of *P. regilla* and *Taricha* from 1 January 1971 to 23

November 2021 (GBIF) using the following R packages: RColorBrewer v.1.1.2, ggplot2

v.3.3.5, chron v.2.3.56, cowplot v.1.1.1, and dplyr v.1.0.7 (Neuwirth, 2014; Wickham,

2016; James and Hornik, 2020; Wilke, 2020; Wickham et al., 2021). In our search terms

for GBIF, we took into account that the taxonomy of *Pseudacris regilla* has changed

several times over the last few decades (Recuero et al., 2006; Fouquette and Dubois,

2014; Duellman et al., 2016). iNaturalist categorizes some *Pseudacris regilla* (e.g., in

the San Francisco Bay Area) as *Pseudacris sierra*. However, these distinctions are not

universally accepted by the herpetology community (Barrow et al., 2014; Crother et al.,

2017). Here we consider *P. sierra*, *P. hypochondriaca*, and *Hyla regilla* synonyms of

Pseudacris regilla. We downloaded occurrence data from GBIF for the *P. regilla*

synonyms listed here and for all currently recognized species of *Taricha* (*T. granulosa*,

T. rivularis, *T. sierrae*, *T. torosa*), as well as those no longer recognized (*T. lindoei*, *T.*

miocenica, and *Palaeotaricha oligocenica*) and individuals identified only to the genus

Taricha. GBIF data include Research Grade iNaturalist observations in addition to

museum and specimen records.

We aimed to sequence *Na$_V$1.4* from *P. regilla* that were sympatric and allopatric

with *Taricha* (see below). We originally classified *P. regilla* tissue samples as sympatric

if they were collected within 35 km or 500 m of elevation from the nearest *Taricha*

occurrence documented on GBIF as of 23 November 2021; this included 11 individuals

and the single *P. regilla* collected at OBRT 2. We classified allopatric *P. regilla* samples

as those individuals collected at least 35 km or 500 m in elevation away from the

nearest *Taricha* observation; this included 18 individuals and the eight *P. regilla* from

the Mountain Lake field site in the Presidio (Table S1).

To review general patterns in geographic range overlap between species, we

plotted *P. regilla* and *Taricha* GBIF occurrence data from the western USA, as well as

localities of museum samples we sequenced using ggplot2 v.3.3.5 and tidyverse v.1.3.1

(Wickham, 2016; Wickham et al., 2019; GBIF). We plotted the occurrence data of these

species to confirm that the museum tissues were from locations where *P. regilla* and

Taricha are allopatric or sympatric. Specimens that were collected from localities within

overlapping hexagons were considered to be sympatric; none of our categorizations

changed. The binwidth of each hexagon was 0.05 in the vertical and horizontal

directions. We note that this visualization, though qualitatively useful, is limited in

several ways. For example, our designations of sympatry versus allopatry are not

indicative of species distributions prior to 1971. Additionally, iNaturalist data included in

the GBIF data can be biased; for instance, individuals are unlikely to post multiple

images of a single species at a site and observations can be more highly concentrated

in regions where the iNaturalist is well-known (Hochmair et al., 2020). Future studies

may consider estimating more accurate species distribution models.

DNA Sequencing and Analysis.—We selected 29 liver tissues of *Pseudacris*

regilla from the Museum of Vertebrate Zoology at the University of California, Berkeley

to broaden our geographic sampling beyond the San Francisco Bay Area (Table S1). In

total, we sequenced $Na_V1.4$ from 11 *P. regilla* that were sympatric with *Taricha* and 18

that were allopatric. We predicted that $Na_V1.4$ sequences from sympatric, but not

allopatric, *P. regilla* would show amino acid substitutions potentially conferring TTX

resistance. We also sequenced $Na_V1.4$ from two close relatives of *P. regilla, Pseudacris*

cadaverina and *Acris crepitans* (Table S1). The individuals of *P. cadaverina* and *A.*

crepitans are from locations far from *Taricha* and, thus, were likely not exposed to TTX

from *Taricha*. However, the range of *A. crepitans* overlaps with another TTX-bearing

newt, *Notophthalmus viridescens* (Baird, 1854; Mecham, 1967), so it may be exposed

to TTX as well. Thus, if TTX from newts selects for mutations in $Na_V1.4$ in sympatric

frog species, we would expect to find amino acid substitutions in *Nav1.4* of *A. crepitans*

as well as in *Nav1.4* of *P. regilla* sympatric with *Taricha*. In contrast, we expected to find

no resistance-conferring amino acid substitutions in allopatric *P. regilla* or in *P.*

cadaverina.

In the Evolutionary Genetics Lab (UC Berkeley), we extracted DNA from tissues

according to the Miller et al. (1988) salt-extraction protocols, with modifications (see

Supporting Methods for details). The primers used for *P. regilla* samples targeted exon

24 (DIV segment 6 and p-loop) of the gene SCN4A: 2268F_SCN4A 5'-

TCTCCCGGCCCTCTTCAATA-3' and 2681R_SCN4A 5'-

TCGTCCTCGCATAAAGGCTC-3' (Tarvin et al., 2016). We added 0.5 µL of $MgCl_2$ to

each PCR sample. PCR protocols were: an initial denaturation for 2 min at 94 ºC; 35

cycles of: 94 ºC at 30 s for denaturation, an annealing temperature of 54 ºC for 30 s,

and an elongation temperature of 72 ºC for 45 s; and a final extension time of 7 min at

72 ºC. PCR products were run on an agarose gel, purified with ExoSAP-IT PCR Product

Cleanup Reagent (ThermoFisher Scientific, 78201.1.ML) and sequenced using the

reverse primer 2681R_SCN4A at the UC Berkeley DNA Sequencing Facility. For PCR

and sequencing samples of *Pseudacris cadaverina* and *Acris crepitans*, we used the

same protocol and forward primer, but the reverse primer was redesigned based on the

DIV SCN4A *P. regilla* sequences we produced: P.regilla_SCN4AD4_R 5'-

ACTGCTTTCTTCTGTGGCCAC-3'.

Using BLAST (Altschul et al., 1990) with the *P. regilla Nav1.4* sequence as a

query, we obtained 180 previously published representative sequences of SCN4A from

other herpetofauna including salamandrids, dendrobatids, snakes, and lizards with

available data on GenBank and DataDryad (Table S2; Gendreau et al., 2021; Gendreau

et al., 2021). We compare these SCN4A sequences, as well as those of *P. cadaverina*

and *A. crepitans*, to SCN4A of our *P. regilla* samples. Of the additional 180 sequences

added, *Erythrolamprus epinephelus, Rhabdophis tigrinus*, *Thamnophis elegans*,

Thamnophis sirtalis, *Thamnophis atratus*, *Taricha torosa*, *Taricha granulosa, S.*

salamandra, Tylototriton shanjing, and other salamandrids are either confirmed or

hypothesized to possess target-site insensitivity to TTX via mutations in *Na$_V$1.4*

(Feldman et al., 2009; Feldman et al., 2012; Hanifin and Gilly, 2015; Ramírez-

Castañeda, 2017; Gendreau et al., 2021; Vaelli et al. 2021). We use *Rattus norvegicus*

as a reference (Uniprot #P15390) of a species that has not been documented to be

resistant to TTX.

We identified the location of the DIV p-loop in *Na$_V$1.4* based on protein domains

inferred by Tikhonov and Zhorov (2005), and we numbered the residues according to

SCN4A of Rattus norvegicus (UniProt accession #P15390). We aligned sequences

using Clustal Omega in Geneious (Sievers et al., 2011, Geneious Prime 2020.1.1). We

reviewed the alignment by eye for amino acid differences between sequences, focusing

on residues that have been shown to provide TTX resistance. We estimated a

phylogeny of the *Na$_V$1.4* DIV sequences using maximum likelihood on the IQTree web

server, with two partitions (one for codon positions 1 and 2, another for codon position

3), the GTR + G substitution model, and support assessed using 1000 ultrafast

bootstraps and the SH-aLRT test (Nguyen et al., 2015; Chernomor et al., 2015;

Trifinopoulos et al., 2016; Hoang et al., 2018; IQ-Tree web server

http://iqtree.cibiv.univie.ac.at/). We edited and visualized the phylogeny using the

Interactive Tree of Life (Letunic and Bork, 2021).

292   RESULTS

Field Surveys.—While we did not observe *P. regilla* at OBRT 2 or UCBG, their

abundance at OBRT 1 peaked early in the observation period at 32 individuals on the

first field trip date (20 October 2019). For OBRT field sites, *Taricha* abundance peaked

at the beginning of February; for the UCBG, abundance was low, and we may not have

observed the seasonal peak (surveys were suspended at the end of February 2020

because of the COVID-19 pandemic). *Taricha torosa* egg masses were most abundant

at the beginning of February 2020 at OBRT 1 and OBRT 2, while they were most

abundant at UCBG at the end of February (though we cannot be certain these are true

peaks because our surveys ended before March) (Fig. S2).

GBIF.—California-wide GBIF observations for *P. regilla* and *Taricha* from 1

January 1971 to 23 November 2021 indicate two seasonal peaks of *P. regilla*

abundance in April and August, while *Taricha* peaks in January (Fig. S3). Though the

abundance peaks do not occur at precisely the same time, *P. regilla* and *Taricha* do

overlap with one another and, thus, the species have opportunities to interact such that

Taricha could expose *P. regilla* to TTX. These data are summarized across California

and thus represent average activity across a range of habitat types. Nevertheless, when

the data are constrained to a square roughly representing the San Francisco Bay Area

(N 38.66226, S 37.09298, E -121.12701, W -123.23579), the results are similar,

indicating a peak for *Taricha* in January and a peak for *P. regilla* in March (Fig. S4).

Along the west coast of the USA, patterns of geographic sympatry estimated from GBIF

occurrence data vary substantially (Fig. 2). Hot spots of sympatry were found in

Humboldt County, the San Francisco Bay Area, Los Angeles area, and northern

Oregon, although much of each taxon's range does not overlap with that of the other.

Amino Acid Changes.— We found one amino acid change (I1519M, numbered

according to the *Rattus norvegicus* UniProt accession #P15390) within the DIV p-loop of

the sodium channel protein $Na_V1.4$ in *Pseudacris regilla* compared to a sequence of

Rattus norvegicus (a common species used in comparative TTX studies [e.g., Tikhonov

and Zhorov, 2005; Tarvin et al. 2016] as it represents the mammalian TTX-sensitive

$Na_V1.4$ [Chahine et al., 1994]) (Fig. 3). Importantly, there were no observed differences

in amino acid sequences between *P. regilla* specimens that were sympatric and

allopatric with *Taricha*. Close relatives of *P. regilla* (*P. cadaverina* or *A. crepitans*) did

not possess the I1519M substitution, suggesting that the mutation arose within *P. regilla*

or its ancestor.

326   DISCUSSION

The *Taricha-Thamnophis* system is a well-studied example of predator prey

interactions involving a toxin. Here we extend the system to include an ecological

associate, *P. regilla*. We note that *Taricha* are well known to vary the level of their toxic

defenses over time, complicating the arms race story (Bucciarelli et al., 2021; Bucciarelli

et al., 2022). Furthermore, while TTX is known to be produced by bacterial symbionts in

several marine organisms and in *Taricha torosa* (Magarlamov et al., 2017; Vaelli et al.,

2020; Bucciarelli et al., 2021), the biosynthesis pathways of TTX remain unresolved,

and endogenous production of TTX in newts is also plausible (Kudo et al., 2020; Kudo

et al., 2021; but see Bucciarelli et al. 2022). Here we document ecologically relevant

interactions between *Taricha* and *P. regilla* (Fig. 1) with new field observations and an

assessment of GBIF data (Fig. 2; Figs. S2–S4). We show the absence of any obvious

TTX-resistance-conferring amino acid mutations in the $Na_V1.4$ DIV p-loop of *P. regilla*

(Fig. 3). It remains unknown how *P. regilla* survives living in such close proximity with

toxic *Taricha* newts and whether it encounters high enough levels of TTX to evolve

resistance. Our data have added to the complex story involving newt toxicity and its

effects on members of the surrounding community.

Abundance data indicate that *P. regilla* and *Taricha* annual activity patterns

overlap but that peak breeding times are not synchronized (Figs. S2–S4). When we

surveyed the field sites in Briones Regional Park and the UC Botanical Garden from

October 2019 to March 2020, California was experiencing drought. The area of the

ponds fluctuated due to changes in precipitation and, in the case of the Japanese Pool,

some drainage for maintenance (UCBG personnel, pers. com.). Thus, weather and

pond conditions may have influenced our survey observations.

Observations of mistaken amplexus and predation of newts by *P. regilla* indicate

that the species do come into contact (Fig. 1). Still, TTX exposure may not be frequent

or potent enough for *P. regilla* to undergo strong selection for resistance to TTX. In

cases where environmental selective regimes are not predictable, more plastic

responses like upregulation of detoxifying enzymes may be the primary way animals

deal with toxin exposure (Zanger and Schwab, 2013). Nevertheless, even occasional

exposure to TTX can result in maintenance of target-site insensitivity (Durso et al.,

2021).

We hypothesized that we would find TTX-resistance-conferring amino acid

substitutions in *Na$_V$1.4* in *P. regilla* that are sympatric with *Taricha* and no substitutions

in *P. regilla* that are allopatric with *Taricha*, with a greater number of substitutions

indicating a higher level of resistance as has been shown various Na$_V$ channels in

salamanders and snakes (Geffeney et al., 2005; Feldman et al., 2010; Hanifin and Gilly,

2015; McGlothlin et al., 2016). Compared to the TTX-sensitive *Rattus norvegicus*

Na$_V$1.4 DIV, we found one amino acid change, I1519M, at a conserved hydrophobic

residue located in proximity to TTX-sensing residues in the p-loop (Tikhonov and

Zhorov, 2005; Geffeney et al., 2005; Tikhonov and Zhorov, 2011; Feldman et al. 2012;

Gendreau et al., 2021). This substitution was present in *P. regilla* that were sympatric

and allopatric with *Taricha.* Amino acid changes in important regions of the *Na$_V$1.4*

protein do not always translate to a resistant phenotype (e.g., see Abderemane-Ali et

al., 2021). Furthermore, the presence of this substitution in all *P. regilla* samples does

not support our hypothesis that substitutions would only be found where *P. regilla* is

exposed to TTX from sympatric *Taricha* (at least based on occurrence data from the last

50 years). We recognize that our hypothesis made a simplifying assumption that the

designation of *P. regilla* populations as sympatric or allopatric with *Taricha* is relevant

and valid during the time in which selection may have occurred for TTX resistance, and

that gene flow does not occur between sympatric and allopatric *P. regilla* populations.

Nonetheless, our study suggests that the I1519M substitution evolved in the ancestor of

P. regilla, which likely rules out contemporary selection from *Taricha.* Furthermore, our

survey of SCN4A sequences in reptiles and amphibians (Fig. 3) suggests that this site

is relatively less conserved than other adjacent sites.

In addition to *P. regilla*, the I1519M substitution is present in *Nanorana parkeri*,

Bufo bufo, Bufo gargarizans, Pogona vitticeps, Charina bottae, and *Rhabdophis tigrinus*

(Fig. 3)*.* None of these species are known for having a strong association with TTX;

though, *R. tigrinus* has been observed consuming TTX-defended treefrogs (Feldman et

al., 2012). Furthermore, *Nanorana parkeri*, which also possesses the I1519M transition,

occurs within the range of the toxic and relatively TTX-resistant newt *Tylototriton*

verrucosus, though its toxin is yet to be identified (Brodie et al., 1984; Hanifin and Gilly,

2015, which studies a relative *T. shanjing*). Similarly, *A. crepitans* co-occurs with the

TTX-defended newt *Notophthalmus viridescens* (Mebs et al., 2010)*,* but this species

does not possess the I1519M substitution.

The close proximity of the I1519M substitution to other changes implicated in

TTX resistance may support its importance in TTX resistance (Feldman et al. 2012).

However, the hypothesis that a I1519M substitution contributes to TTX resistance has

been suggested to be incorrect because the same substitution is found in other TTX-

sensitive voltage-gated sodium channel paralogs (McGlothlin et al., 2016; noted as

I1555M in their study). Mutagenesis studies suggest that substituting methionine for

isoleucine has limited effect on a protein (Ohmura et al., 2001). Another hydrophobic

change in the same position (I1519L) is found in several genera of TTX-resistant newts

(Fig. 3: *Triturus dobrogicus, Cynops pyrrhogaster, Pachytriton labiatus, Notophthalmus*

viridescens); however, it is hypothesized that this change is not directly related to TTX

resistance either (Hanifin and Gilly, 2015). Because there is disagreement whether the

I1519M mutation (or the site itself) provides resistance, we cannot say with certainty

whether it provides TTX resistance in *P. regilla* without further experimental evidence.

While there is not strong evidence that *P. regilla* possesses target-site

insensitivity in the p-loop of DIV in *Na$_V$1.4*, it is plausible that *P. regilla* may have other

ways of avoiding the effects of TTX. Other regions of *Na$_V$1.4*, such as domain III, and

other voltage-gated sodium channels, such as *Na$_V$1.7*, also harbor amino acid

substitutions implicated in TTX resistance (Feldman et al., 2012; McGlothlin et al.,

2016). *Pseudacris regilla* could possess the ability to smell TTX, as larval *Taricha* can,

in order to avoid it (Zimmer et al., 2006). Alternatively, they could possess a diffusion

barrier in the skin or the gut that would prevent high levels of TTX exposure, as occurs

in mantids (Mebs et al., 2016). It is also possible that *P. regilla* possesses a binding

protein that can scavenge TTX. Other frogs have saxiphillin, a protein that binds

saxitoxin, which is structurally similar to TTX. However, saxiphilin is incapable of binding

TTX, at least in the animals that have been tested (Mahar et al., 1991; Llewellyn et al.,

1997; Abderemane-Ali et al., 2021). Similarly, little is known about the potential

interactions between *Acris crepitans* and *Notophthalmus viridescens*, or between

Nanorana parkeri and *Tylototriton verrucosus,* which could also select for toxin

resistance in the frogs. To date, none of these alternative resistance mechanisms have

been studied in *Pseudacris, Acris, Nanorana,* or salamandrids.

A number of reasons could explain why *P. regilla* would lack target-site

insensitivity in *Na$_V$1.4*. Evolving TTX resistance could be costly to *P. regilla* due to

possible alterations in the selectivity filter of sodium channels that frequently occurs

when amino acid substitutions are present (Chiamvimonvat et al., 1996; Lee et al.,

2011; Feldman et al., 2012; Hague et al., 2018; but see Moniz et al., 2021 for evidence

against potential impact of mutations on metabolism of the organism). Evolving costly

mutations may require a stronger selective pressure than is currently occurring. Given

the relative quantities of TTX on newt skin (up to 5.48 mg/newt [Reimche et al., 2020] or

even one recording of 8.551 mg/newt [Stokes, 2015]) compared to TTX levels in newt-

containing water (~1×10^{-7} mol/L [Zimmer et al., 2006]), the environmental toxicity of TTX

is likely to cause a weaker selection for TTX resistance than exposure via chemical

defenses from prey. The cost of exhibiting TTX resistance combined with the low

potential for *P. regilla* to interact with high concentrations of TTX in its environment may

explain why we did not unequivocally identify target-site insensitivity in *P. regilla* DIV

Na$_V$1.4. Future work exploring the effects of TTX as an environmental toxin could

include measuring TTX concentrations in newt-containing ponds, performing whole

organism or *in vitro* assays of *P. regilla* to determine TTX sensitivity, and testing for

other modes of TTX resistance.

Conclusions.—We do not find strong evidence for target-site insensitivity in DIV

of the skeletal muscle voltage-gated sodium channel *Na$_V$1.4* of *Pseudacris regilla*.

Although it is possible that the I1519M substitution in the p-loop of *Na$_V$1.4* may be

related to TTX resistance, *P. regilla* may not encounter TTX from *Taricha* with high

enough frequency or magnitude to select for TTX resistance. Future seasonal

measurements of pond-water concentrations will help evaluate this hypothesis. If *P.*

regilla are indeed exposed to significant levels of TTX, they may have behavioral or

physiological mechanisms to survive TTX exposure other than target-site insensitivity in

Na$_V$1.4 DIV. Investigating the environmental toxicity of TTX in the relationship between

P. regilla and *Taricha* is an opportunity to explore the nexus of ecological interactions,

neurotoxins, and resistance mechanisms. The likely absence of target-site insensitivity

in *P. regilla* indicates that exposure to tetrodotoxin outside of trophic interactions may

not strongly select for fixed mechanisms of toxin resistance.

Acknowledgments.—We affirm the sovereignty of the Karkin, Me-Wuk (Bay

Miwok), Muwekma, Chochenyo, Ramaytush, and Ohlone people and the Confederated

Villages of Lisjan; we thank them for stewarding the land where we worked and, thus,

serving an integral role in the production of this study. We thank L. Smith in the

Evolutionary Genetics Lab, as well as C. Spencer and the support of the Museum of

Vertebrate Zoology collections staff. We also thank those who supported our work at

field sites: UC Botanical Garden staff, including H. Forbes, J. Fong, E. Hupperts, and C.

Loughran; East Bay Regional Parks District staff, including T. Lin and J. Miller; and J.

Young with the Presidio Trust. We thank the other members of the Tarvin Lab for their

editing prowess and general support. We thank A. Shabel for his advice in the early

stages of this project, and K. Sanko, J. Toman, and L. Galindo who helped as field

assistants. We appreciate the newts and frogs who were the focus of this project and

who were sacrificed to offer the knowledge they hold. K. Montana thanks her family and

ancestors for their guidance. R. Tarvin was supported by UC Berkeley start-up funds.

The East Bay Regional Parks District authorized the surveys at Briones under

permit #19-1044, and staff at the UCBG gave written consent for surveys conducted on

the premises. All animals were collected under approved collection permits and

protocols: for the National Parks Service Scientific Research and Collecting Permit 2

number PRSF-2020-SCI-0002 within study PRSF-00035, with a written extension after

expiration of the original permit, University of California IACUC AUP-2019-08-12457,

and California Department of Fish and Wildlife Scientific Collection Permit S-

190980001-19111-001.

The Spanish translation was done using www.DeepL.com/Translator (free

version), which was then edited by Valeria Ramírez-Castañeda.

478   LITERATURE CITED

ABDEREMANE-ALI, F., N.D. ROSSEN, M.E. KOBIELA, R.A. CRAIG II, C.E. GARRISON, Z. CHEN,

C.M. COLLERAN, L.A. O'CONNELL, J. DU BOIS, J.P. DUMBACHER, AND D.L. MINOR, JR.

2021. Evidence that toxin resistance in poison birds and frogs is not rooted in

sodium channel mutations and may rely on 'toxin sponge' proteins. Journal of

General Physiology 153:e202112872. https://doi.org/10.1085/jgp.202112872.

ALTSCHUL, S. F., W. GISH, W. MILLER, E. W. MYERS AND D.J. LIPMAN. 1990. Basic local

alignment search tool. Journal of Molecular Biology 215:403–410.

BAIRD, S.F. 1854. Descriptions of new genera and species of North American frogs.

Proceedings of the Academy of Natural Sciences of Philadelphia 7:59–62.

BARROW, L.N., H.F. RALICKI, S.A. EMME, AND E.M. LEMMON. 2014. Species tree estimation

of North American chorus frogs (Hylidae: *Pseudacris*) with parallel tagged

amplicon sequencing. Molecular Phylogenetics and Evolution 75:78–90.

BRODIE, E.D., JR. 1968. Investigations on the skin toxin of the adult rough-skinned

newt, *Taricha granulosa*. Copeia 1968:307–313.

BRODIE, E.D., JR., J.L. HENSEL, AND J.A. JOHNSON. 1974. Toxicity of the urodele

amphibians *Taricha*, *Notophthalmus*, *Cynops* and *Paramesotriton*

(Salamandridae). Copeia 1974:506–511.

BRODIE, E.D., JR. B.J. RIDENHOUR, AND E. D. BRODIE III. 2002. The evolutionary

response of predators to dangerous prey: hotspots and coldspots in the

geographic mosaic of coevolution between garter snakes and newts.

Evolution 56:2067–2082.

BRODIE, E.D., R.A. NUSSBAUM, AND M. DIGIOVANNI. 1984. Antipredator

Adaptations of Asian Salamanders (Salamandridae). Herpetologica 40:56–68.

BUCCIARELLI, G.M., AND L.B. KATS. 2015. Effects of newt chemical cues on the

distribution and foraging behavior of stream macroinvertebrates. Hydrobiologia

749:69–81.

BUCCIARELLI, G.M., H.B. SHAFFER, D.B. GREEN, AND L.B. KATS. 2017. An amphibian

chemical defense phenotype is inducible across life history stages. Scientific

Reports 7:8185. https://doi.org/10.1038/s41598-017-08154-z.

BUCCIARELLI, G.M., M. LECHNER, A. FONTES, L.B. KATS, H.L. EISTHEN, AND H.B. SHAFFER.

2021. From poison to promise: the evolution of tetrodotoxin and its potential as a

therapeutic. Toxins 13:517.

BUCCIARELLI, G.M., F. ALSALEK, L.B. KATS, D.B. GREEN, AND H.B. SHAFFER. 2022. Toxic

relationships and arms-race coevolution revisited. Annual Review of Animal

Biosciences 10:63–80.

CALHOUN, D.M., G.M. BUCCIARELLI, L.B. KATS, R.K. ZIMMER, AND P.T.J. JOHNSON. 2017.

Noxious newts and their natural enemies: experimental effects of tetrodotoxin

exposure on trematode parasites and aquatic macroinvertebrates. Toxicon

137:120–127.

CHAHINE, M., P.B. BENNETT, A.L. GEORGE, AND R. HORN. 1994. Functional expression and

properties of the human skeletal muscle sodium channel. European Journal of

Physiology 427:136–142.

CHERNOMOR, O., A. VON HAESELER, AND B.Q. MINH. 2016. Terrace aware data structure

for phylogenomic inference from supermatrices. Systematic Biology 65:997–

1008.

CHIAMVIMONVAT, N., M.T. PÉREZ-GARCÍA, G.F. TOMASELLI, AND E. MARBAN. 1996.

Control of ion flux and selectivity by negatively charged residues in the outer

mouth of rat sodium channels. The Journal of Physiology 491:51–59.

CROTHER, B.I., SOCIETY FOR THE STUDY OF AMPHIBIANS AND REPTILES, COMMITTEE ON

STANDARD ENGLISH AND SCIENTIFIC NAMES, AMERICAN SOCIETY OF ICHTHYOLOGISTS

AND HERPETOLOGISTS, CANADIAN ASSOCIATION OF HERPETOLOGISTS, PARTNERS IN

AMPHIBIAN AND REPTILE CONSERVATION, SOCIETY FOR THE STUDY OF AMPHIBIANS

AND REPTILES, AND HERPETOLOGISTS' LEAGUE. 2017. Scientific and Standard

English Names of Amphibians and Reptiles of North America North of Mexico,

with Comments Regarding Confidence in Our Understanding, Eighth Edition.

DUELLMAN, W.E., A.B. MARION, AND S. B. HEDGES. 2016. Phylogenetics, classification,

and biogeography of the treefrogs (Amphibia: Anura: Arboranae). Zootaxa

4104:1–109.

DURSO, A.M., L.A. NEUMAN-LEE, G.R. HOPKINS, AND E.D. BRODIE, JR. 2021. Stable

isotope analysis suggests that tetrodotoxin-resistant Common Gartersnakes

(*Thamnophis sirtalis*) rarely feed on newts in the wild. Canadian Journal of

Zoology 99:331–338.

FELDMAN, C.R., E.D. BRODIE, JR., E.D. BRODIE III, AND M.E. PFRENDER. 2009.

The evolutionary origins of beneficial alleles during the repeated adaptation of

garter snakes to deadly prey. Proceedings of the National Academy of

Sciences 106:13415–3420.

FELDMAN, C.R., E.D. BRODIE, JR., E.D. BRODIE III, AND M.E. PFRENDER. 2010. Genetic

architecture of a feeding adaptation: garter snake (*Thamnophis*) resistance to

tetrodotoxin bearing prey. Proceedings of the Royal Society B 277.

https://doi.org/10.1098/rspb.2010.0748.

FELDMAN, C.R., E.D. BRODIE, JR., E.D. BRODIE III, AND PFRENDER, M.E. 2012.

Constraint shapes convergence in tetrodotoxin-resistant sodium channels of

snakes. Proceedings of the National Academy of Sciences 109:4556–4561.

FOUQUETTE JR., M. J., AND A. DUBOIS. 2014. A checklist of North American amphibians

and reptiles: the United States and Canada. Xlibris Corporation.

GALL, B.G., A.N. STOKES, S.S. FRENCH, E.A. SCHLEPPHORST, E.D. BRODIE III, AND E.D.

BRODIE, JR. 2011. Tetrodotoxin levels in larval and metamorphosed newts

(*Taricha granulosa*) and palatability to predatory dragonflies. Toxicon 57:978–

983.

GALL, B.G., A.N. STOKES, S.S. FRENCH, E.D. BRODIE III, AND E.D. BRODIE, JR. 2012.

Predatory caddisfly larvae sequester tetrodotoxin from their prey, eggs of the

rough-skinned newt (*Taricha granulosa*). Journal of Chemical Ecology 38:1351–

1357.

GBIF.org (23 November 2021) GBIF Occurrence Download

https://doi.org/10.15468/dl.qzrq4k.

GBIF.org (23 November 2021) GBIF Occurrence Download

https://doi.org/10.15468/dl.z9tw6r.

GBIF.org (30 November 2021) GBIF Occurrence Download

https://doi.org/10.15468/dl.dwz9yn.

GBIF.org (30 November 2021) GBIF Occurrence Download

https://doi.org/10.15468/dl.ed2uqn.

GBIF.org (23 May 2022) GBIF Occurrence Download

https://doi.org/10.15468/dl.wzbmv6

GBIF.org (06 June 2022) GBIF Occurrence Download

https://doi.org/10.15468/dl.fy2m4g

GEFFENEY, S.L., E. FUJIMOTO, E.D. BRODIE III, E.D. BRODIE, JR., AND P.C. RUBEN. 2005.

Evolutionary diversification of TTX-resistant sodium channels in a predator–prey

interaction. Nature 434:759–763.

GENDREAU, K.L., A.D. HORNSBY, M.T.J. HAGUE, AND J.W. MCGLOTHLIN. 2021. Gene

conversion facilitates the adaptive evolution of self-resistance in highly toxic

newts. Molecular Biology and Evolution 38:4077–4094.

GENDREAU, K.L., J.W. MCGLOTHLIN, A.D. HORNSBY, AND M.T.J. HAGUE. 2021. Gene conversion

facilitates the adaptive evolution of self-resistance in highly toxic newts." Dryad.

https://doi.org/10.5061/DRYAD.KPRR4XH4V.

Geneious Prime 2020.1.1. https://www.geneious.com.

HAGUE, M.T.J., G. TOLEDO, SHANA L. GEFFENEY, C.T. HANIFIN, E.D. BRODIE, JR., AND E.D.

BRODIE, III. 2018. Large-effect mutations generate trade-off between predatory

and locomotor ability during arms race coevolution with deadly prey. Evolution

Letters 2:406–416.

HANIFIN, C.T., AND W.F. GILLY. 2015. Evolutionary history of a complex adaptation:

tetrodotoxin resistance in salamanders. Evolution; International Journal of

Organic Evolution 69:232–244.

HOANG, D.T., O. CHERNOMOR, A. VON HAESELER, B.Q. MINH, AND L.S. VINH. 2018.

UFBoot2: improving the ultrafast bootstrap approximation. Molecular Biology and

Evolution 35:518–522.

HOCHMAIR, H.H., R.H. SCHEFFRAHN, M. BASILLE, AND M. BOONE. 2020.

Evaluating the data quality of iNaturalist termite records. PLOS ONE

15:e0226534. https://doi.org/10.1371/journal.pone.0226534.

JAMES, D., AND K. HORNIK. 2020. chron: chronological objects which can handle dates

and times. R package version 2.3-56.

JOST, M.C., D. M. HILLIS, Y. LU, J.W. KYLE, H.A. FOZZARD, AND H.H. ZAKON. 2008.

Toxin-resistant sodium channels: parallel adaptive evolution across a complete

gene family. Molecular Biology and Evolution, 25:1016–1024.

KAO, C.Y., AND S.R. LEVINSON. 1986. Preface—tetrodotoxin, saxitoxin, and the

molecular biology of the sodium channel. Annals of the New York Academy of

Sciences 479:ix-x.

KASSAMBARA, A. 2020. ggpubr: 'ggplot2' Based Publication Ready Plots. R package

version 0.4.0. https://CRAN.R-project.org/package=ggpubr.

KUDO, Y., C.T. HANIFIN, AND M. YOTSU-YAMASHITA. 2021. Identification of tricyclic

guanidino compounds from the tetrodotoxin-bearing newt *Taricha granulosa*.

Organic Letters 23:3513–3517.

KUDO, Y., C.T. HANIFIN, Y. KOTAKI, AND M. YOTSU-YAMASHITA. 2020. Structures of

n-hydroxy-type tetrodotoxin analogues and bicyclic guanidinium compounds

found in toxic newts. Journal of Natural Products 83:2706–2717.

LEE, C.H., D.K. JONES, C. AHERN, M.F. SARHAN, AND P.C. RUBEN. 2011. Biophysical costs

associated with tetrodotoxin resistance in the sodium channel pore of the garter

snake, *Thamnophis sirtalis*. Journal of Comparative Physiology A 197:33–43.

LETUNIC, I., AND P. BORK. 2021. Interactive tree of life (ITOL) v5: an online tool for

phylogenetic tree display and annotation. Nucleic Acids Research 49:W293–

W296.

LLEWELLYN, L.E., P.M. BELL, AND E.G. MOCZYDLOWSKI. 1997. Phylogenetic survey of

soluble saxitoxin-binding activity in pursuit of the function and molecular evolution

of saxiphilin, a relative of transferrin. Proceedings of the Royal Society B:

Biological Sciences 264:891–902.

MAGARLAMOV, T.Y., D.I. MELNIKOVA, AND A.V. CHERNYSHEV. 2017. Tetrodotoxin-

producing bacteria: detection, distribution and migration of the toxin in aquatic

systems. Toxins:166.

MAHAR, J., G.L. LUKÁCS, L. LI, S. HALL, AND E. MOCZYDLOWSKI. 1991. Pharmacological

and biochemical properties of saxiphilin, a soluble saxitoxin-binding protein from

the bullfrog (*Rana catesbeiana*). Toxicon 29:53–71.

MAILHO-FONTANA, P.L., C. JARED, M.M. ANTONIAZZI, J. MOZER SCIANI, D. CARVALHO

PIMENTA, A.N. STOKES, T. GRANT, E.D. BRODIE III, AND E.D. BRODIE, JR. 2019.

Variations in tetrodotoxin levels in populations of *Taricha granulosa* are

expressed in the morphology of their cutaneous glands. Scientific Reports

9:18490. https://doi.org/10.1038/s41598-019-54765-z.

McGLOTHLIN, J.W., M.E. KOBIELA, C.R. FELDMAN, T.A. CASTOE, S.L. GEFFENEY, C.T.

HANIFIN, G. TOLEDO, F.J. VONK, M.K. RICHARDSON, E.D. BRODIE, JR., M.E.

PFRENDER, E.D. BRODIE, III. 2016. Historical contingency in a multigene family

facilitates adaptive evolution of toxin resistance. Current Biology 26:1616–1621.

MEBS, D., M. YOTSU-YAMASHITA, AND O. ARAKAWA. 2016. The praying mantis (*Mantodea*)

as predator of the poisonous red-spotted newt *Notophthalmus viridescens*

(Amphibia: Urodela: Salamandridae). Chemoecology 26:121–126.

MEBS, D., O. ARAKAWA, AND M. YOTSU-YAMASHITA. 2010. Tissue distribution of

tetrodotoxin in the red-spotted newt *Notophthalmus viridescens*. Toxicon 55:

1353–1357.

MECHAM, J.S. 1967. *Notophthalmus viridescens.* Catalogue of American Amphibians

and Reptiles:53.1–53.4.

MILLER, S.A., D.D. DYKES, AND H.F. POLESKY. 1988. A simple salting out procedure for

extracting DNA from human nucleated cells. Nucleic Acids Research 16:1215.

MONIZ, H.A., M.A. RICHARD, C.M. GIENGER, AND C.R. FELDMAN. 2021. Every breath you

take: assessing metabolic costs of toxin resistance in garter snakes

(*Thamnophis*). Integrative Zoology. https://doi.org/10.1111/1749-4877.12574.

NEUWIRTH, E. 2014. RColorBrewer: ColorBrewer palettes. R package version1.1-2.

https://CRAN.R-project.org/package=RColorBrewer.

NGUYEN, L., H.A. SCHMIDT, A. VON HAESELER, AND B.Q. MINH. 2015. IQ-TREE: A fast and

654   effective stochastic algorithm for estimating maximum likelihood phylogenies.

655   Molecular Biology and Evolution 32:268–274.

656 OHMURA, T., T. UEDA, Y. HASHIMOTO, AND T. IMOTO. 2001. Tolerance of

657   point substitution of methionine for isoleucine in hen egg white lysozyme.

658   Protein Engineering, Design and Selection 14:421–425.

659 OTA, W.M., B. OLSEN, G.M. BUCCIARELLI, AND L.B. KATS. 2018. The effect of newt toxin

660   on an invasive snail. Hydrobiologia 817:341–348.

661 PRESTON, DANIEL L., AND PIETER T. J. JOHNSON. 2012. Importance of native amphibians in

662   the diet and distribution of the aquatic gartersnake (*Thamnophis atratus*) in the

663   San Francisco Bay Area of California. Journal of Herpetology 46:221–227.

664 RAMÍREZ-CASTAÑEDA, V. 2017. Origin of mutations in the NaV gene family associated

665   with resistance to neurotoxins in snake (*Erythrolamprus sp.*) predators of dart

666   poison frogs (Dendrobatidae). Master's thesis. University of the Andes, Bogota,

667   Colombia.

668 R CORE TEAM. 2021. R: A language and environment for statistical computing. R

669   Foundation for Statistical Computing, Vienna, Austria.

670   https://www.R-project.org.

671 RECUERO, E., Í. MARTÍNEZ-SOLANO, G. PARRA-OLEA, AND M. GARCÍA-PARÍS. 2006.

672   Phylogeography of *Pseudacris regilla* (Anura: Hylidae) in western North America,

673   with a proposal for a new taxonomic rearrangement. Molecular Phylogenetics

674   and Evolution 39:293–304.

675 RECUERO, E., Í. MARTÍNEZ-SOLANO, G. PARRA-OLEA, AND M. GARCÍA-PARÍS. 2006.

676   Corrigendum to 'Phylogeography of *Pseudacris regilla* (Anura: Hylidae) in

western North America, with a proposal for a new taxonomic rearrangement'

[Mol. Phylogenet. Evol. 39 (2006) 293–304]. Molecular Phylogenetics and

Evolution 41:511.

REIMCHE, J.S., E.D. BRODIE JR., A.N. STOKES, E.J. ELY, H.A. MONIZ, V.L. THILL, J.M.

HALLAS, M.E. PFRENDER, E.D. BRODIE III, AND C.R. FELDMAN. 2020. The

geographic mosaic in parallel: matching patterns of newt tetrodotoxin levels and

snake resistance in multiple predator–prey pairs. Journal of Animal Ecology

89:1645–1657.

SIEVERS F., A. WILM, D.G. DINEEN, T.J. GIBSON, K. KARPLUS, W. LI, R. LOPEZ, H.

MCWILLIAM, M. REMMERT, J. SÖDING, J.D. THOMPSON, D.G. HIGGINS. 2011. Fast,

scalable generation of high-quality protein multiple sequence alignments using

Clustal Omega. Molecular Systems Biology 7:539.

STOKES, A.N. 2015. Otter predation on Taricha granulosa and variation in tetrodotoxin

levels with elevation. Northwestern Naturalist 96:13–21.

TARVIN, R.D., J.C. SANTOS, L.A. O'CONNELL, H.H. ZAKON, AND D.C. CANNATELLA. 2016.

Convergent substitutions in a sodium channel suggest multiple origins of toxin

resistance in poison frogs. Molecular Biology and Evolution 37:607.

TIKHONOV, D.B., AND B.S. ZHOROV. 2005. Modeling p-loops domain of sodium channel:

homology with potassium channels and interaction with ligands. Biophysical

Journal 88:184–197.

TIKHONOV, D.B., AND B.S. ZHOROV. 2011. Possible roles of exceptionally conserved

residues around the selectivity filters of sodium and calcium channels. Journal of

Biological Chemistry 286:2998–3006.

TRIFINOPOULOS, J., L.T. NGUYEN, A. VON HAESELER, AND B.Q. MINH. 2016. W-IQ-TREE: a

fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids

Research 44:W232-W235.

VAELLI, P.M., K.R. THEIS, J.E. WILLIAMS, L.A. O'CONNELL, J.A. FOSTER, AND H.L. EISTHEN.

2020. The skin microbiome facilitates adaptive tetrodotoxin production in

poisonous newts. ELife 9:e53898. https://doi.org/10.7554/eLife.53898.

VENKATESH, B., S.Q. LU, N. DANDONA, S.L. SEE, S. BRENNER, AND T.W. SOONG. 2005.

Genetic basis of tetrodotoxin resistance in pufferfishes. Current Biology 15:2069–

2072.

WICKHAM, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

York. United States of America.

WICKHAM, H., M. AVERICK, J. BRYAN, W. CHANG, L.D. MCGOWAN, R. FRANÇOIS, G.

GROLEMUND, A. HAYES, L. HENRY, J. HESTER, M. KUHN, T.L. PEDERSEN, E. MILLER,

S.M. BACHE, K.MÜLLER, J. OOMS, D. ROBINSON, D.P. SEIDEL, V. SPINU, K.

TAKAHASHI, D. VAUGHAN, C. WILKE, K. WOO, AND H. YUTANI. 2019. Welcome to the

tidyverse. Journal of Open Source Software 4:1686.

https://doi.org/10.21105/joss.01686.

WICKHAM, H., R. FRANÇOIS, L. HENRY, AND K. MÜLLER. 2021. dplyr: a grammar of data

manipulation. R package version 1.0.7.

https://CRAN.R-project.org/package=dplyr.

WILKE, C.O. 2020. cowplot: streamlined plot theme and plot annotations for 'ggplot2'.

R package version 1.1.1. https://CRAN.R-project.org/package=cowplot.

WILLIAMS, B.L., C.T. HANIFIN, E.D. BRODIE, AND E.D. BRODIE III. 2010. Tetrodotoxin affects

survival probability of rough-skinned newts (*Taricha granulosa*) faced with TTX-

resistant garter snake predators (*Thamnophis sirtalis*). Chemoecology 20:285–

290.

ZANGER, U.M., AND M. SCHWAB. 2013. Cytochrome P450 enzymes in drug metabolism:

regulation of gene expression, enzyme activities, and impact of genetic variation.

Pharmacology & Therapeutics 138:103–41.

ZIMMER, R.K., D.W. SCHAR, R.P. FERRER, P.J. KRUG, L.B. KATS, AND W.C. MICHEL. 2006.

The scent of danger: tetrodotoxin (TTX) as an olfactory cue of predation risk.

Ecological Monographs 76:585–600.

733   SUPPLEMENTARY ONLINE MATERIAL

Supporting Methods.—DNA Extraction Procedures, Adapted from Miller et al.,

1988.

Table S1.—Metadata for tissues obtained from the Museum of Vertebrate

Zoology (UC Berkeley) and from newly collected specimens (indicated by asterisks) that

were used to sequence SCN4A exon 24 in this study.

Table S2.—Sequences of *Nav1.4* (SCN4A exon 24) obtained from GenBank

used in Fig. 3.

Figure S1.—Field Sites: A. Old Briones Road Trail 1 (OBRT 1) pond, 37.94441, -

122.13377; B. Old Briones Road Trail 2 (OBRT 2) pond, 37.94352, -122.14102; C.

Japanese Pool at the UC Botanical Garden (UCBG), 37.87440, -122.23760; D. Presidio

site, 37.788496, -122.468605; all photos taken by K. Montana.

Figure S2.—Numbers of A. *P. regilla* adults, B. *Taricha torosa* and *T. granulosa*

adults and larvae, and C. *T. torosa* egg masses observed over field trips to the Old

Briones Road Trail 1 (OBRT 1) (blue circle line), OBRT 2 (red triangle line), and U.C.

Botanical Garden (yellow square line) ponds; 20 October 2019 to late February 2020.

Figure S3.—Percentage of total abundance of GBIF-recorded individuals of

Pseudacris regilla (green) and *Taricha* (blue) in California by month, over the course of

the last 50 years. Peak abundances for *P. regilla* occur around April and August each

year, and peak abundances for *Taricha* occur in January each year. Samples included

were from 1 January 1971 to 23 November 2021.

Figure S4.—Percentage of total abundance of GBIF-recorded individuals of

Pseudacris regilla (green) and *Taricha* (blue) in the San Francisco Bay Area (N

38.66226, S 37.09298, E -121.12701, W -123.23579) by month, over the course of the

last 50 years. Peak abundance for *P. regilla* occurs around March each year, and peak

abundance for *Taricha* occurs in January each year. Samples included were from 1

January 1971 to 23 November 2021

Fig. 1. Examples of *Pseudacris regilla* and *Taricha torosa* interactions: A. Amplexus in

Briones Regional Park, Martinez, CA, 1 March 2019, photo credit K. Montana, user

kmontana on iNaturalist (CC BY-NC 4.0; observation 100725148); B. Another instance

of amplexus in Briones Regional Park, Martinez, CA, 21 February 2009, photo credit

user ap2il on iNaturalist (CC BY-NC 4.0; observation 550227); C. Attempted predation

on *Taricha torosa* by *Pseudacris regilla* in Mark West Springs, CA, 1 March 2014, photo

credit user tiwane on iNaturalist (CC BY-NC 4.0; observation 2312). The frogs did not

appear to be harmed in any of these scenarios, though the tongue posture in C

suggests some numbing effects from newt exposure.

Fig. 2. Map showing overlap of *P. regilla* and *Taricha* occurrences on the west coast of

the USA. Data are from GBIF, collected through 23 May 2022 for *Taricha* and through 6

June 2022 for *P. regilla*. *P. regilla* museum specimens that are sympatric with *Taricha*

are indicated by pink triangles; *P. regilla* museum specimens allopatric with *Taricha* are

indicated by blue triangles; field sites are indicated by black squares. Geographic

regions with only *Taricha* occurrences are indicated by blue hexagons; those with only

P. regilla are indicated by orange hexagons; regions where *Taricha* and *P. regilla*

overlap are indicated by brown hexagons.

783 Fig. 3. Alignment of the p-loop of domain IV of the voltage-gated sodium channel

784 *Na$_V$1.4* in several reptile and amphibian species including new sequences from

785 *Pseudacris regilla,* mapped onto a phylogeny inferred using 381 bp of the domain IV in

786 *Na$_V$1.4* and IQTREE software (support values are SH-aLRT/ultrafast bootstrap values

787 from 1000 iterations each). Amino acid positions are numbered according to *Rattus*

788 *norvegicus,* UniProt accession #P15390. The amino acid substitution I1519M is found in

789 *P. regilla* as well as *N. parkeri*, *Bufo gargarizans*, *Bufo bufo*, *Pogona vitticeps*, *Charina*

790 *bottae*, and *R. tigrinus*.

A.

B.

C.