JOURNAL OF HERPETOLOGY

Characterizing Potential Tetrodotoxin Resistance in Domain IV of the Voltage-Gated

Sodium Channel Na$_v$1.4 of Pacific Chorus Frogs, *Pseudacris regilla*

Katherine O. Montana[1,2,3], Valeria Ramírez Castañeda[1], Rebecca D. Tarvin[1,3]

[1]*Museum of Vertebrate Zoology and Department of Integrative Biology, University of*

California, Berkeley, Berkeley, California, USA 94720

[2]*Present address: Department of Biology, San Francisco State University, San*

Francisco, California, USA 94132 and *Institute for Biodiversity Science and*

Sustainability, California Academy of Sciences, San Francisco, California, USA 94118

[3]*Corresponding authors. kmontana@berkeley.edu, rdtarvin@berkeley.edu*
11

LRH: K.O. Montana et al.

RRH: Tetrodotoxin resistance in *Pseudacris regilla*

Abstract.—Animals that frequently encounter toxins often evolve mechanisms of

toxin resistance. Both predators that consume toxic prey and organisms in physical

contact with a toxin or pollutant in their environment may experience natural selection

for resistance. Based on observations that Pacific Chorus Frogs (*Pseudacris regilla*)

sometimes eat and mistakenly amplect tetrodotoxin (TTX)-defended *Taricha* newts, we

hypothesized that *P. regilla* may possess resistance to TTX. We tested this prediction

by comparing the amino acid sequences of the domain IV pore loop of the muscle

voltage-gated sodium channel gene SCN4A (Na$_v$1.4 protein) in populations of *P. regilla*

that are sympatric and allopatric with *Taricha*. We identified a single substitution in the

domain IV pore loop of Na$_v$1.4 in *P. regilla* that has been hypothesized to provide TTX

resistance based on computational modeling, though its role in resistance has been

challenged. Nonetheless, both allopatric and sympatric *P. regilla* had this substitution,

suggesting that it may be unrelated to TTX exposure from *Taricha*. Thus, there is no

conclusive evidence that *P. regilla* has evolved TTX resistance encoded by amino acid

substitutions in this domain. In addition, California occurrence data from the last 50

years indicate that *Taricha* activity peaks in January while the activity of *P. regilla* peaks

in April. These relatively distinct activity patterns suggest that *P. regilla* may not be

exposed to levels of TTX from *Taricha* that are high enough to require mutations in

SCN4A. Nevertheless, other unidentified mechanisms of TTX resistance could be

present in *P. regilla* and other species that are sympatric with toxic newts.

Key words: Toxin resistance; California; Environmental toxins; Target-site

insensitivity; Salamanders; Taricha

Resumen.— Los animales que tienen contacto frecuente con toxinas suelen

desarrollar mecanismos de resistencia a las mismas. Tanto los depredadores que

consumen presas tóxicas como los organismos en contacto cercano con una toxina o

contaminante en su entorno pueden experimentar una presión de selección que los

lleva a evolucionar resistencia a toxinas. Basándonos en las observaciones de que las

ranas coro del Pacífico (*Pseudacris regilla*) a veces comen por error y/o amplexan las

salamandras del género *Taricha* que poseen tetrodotoxina (TTX), planteamos la

hipótesis de que *P. regilla* podría poseer resistencia a la TTX. Probamos esta

predicción comparando las secuencias de aminoácidos en el loop del poro del dominio

IV en el gen del canal de sodio voltaje dependiente muscular SCN4A (proteína $Na_v1.4$)

en poblaciones de *P. regilla* que son simpátricas y alopátricas con *Taricha*.

Identificamos una única sustitución en el loop del poro del dominio IV de $Na_v1.4$ en *P.

regilla* que se ha hipotetizado utilizando un modelo computacional que está relacionado

con la resistencia a TTX. Sin embargo, tanto el *P. regilla* alopátrico como el simpátrico

tienen esta sustitución, lo que sugiere que esta puede no estar relacionada con la

resistencia a la TTX proveniente de *Taricha*. Por lo tanto, no hay pruebas concluyentes

de que *P. regilla* haya evolucionado resistencia a la TTX causada sustituciones de

aminoácidos en este dominio de la proteina. Adicionalmente, los datos de ocurrencia en

California en los últimos 50 años indican una alternancia en la actividad de ambos

grupos, *Taricha* alcanza su punto máximo en enero, mientras que la de *P. regilla* lo

hace en abril. Estos patrones de actividad relativamente distintos sugieren que *P.

regilla no está expuesta constantemente a niveles de TTX lo suficientemente altos

como para requerir mutaciones en SCN4A. Sin embargo, otros mecanismos de

resistencia a la TTX no identificados hasta ahora, podrían estar presentes en *P. regilla*

y otras especies simpáticas a las salamandras tetrodotóxicas.

Palabras clave: Resistencia a las toxinas; California; Toxinas ambientales;

Insensibilidad en el sitio de union; salamandras; Taricha

Many animals can synthesize or bioaccumulate toxic chemicals as an anti-predator

defense. One common toxin is tetrodotoxin (TTX), which is employed as a defense by a

variety of organisms including pufferfish and amphibians (e.g., dendrobatid frogs*,*

Atelopus, Polypedates, Brachycephalus, salamandrid newts) (Daly et al., 1994; Tanu et

al., 2001; Pires et al., 2002; Hanifin, 2010), as well as an offensive prey-capture

technique by blue-ringed octopuses and flatworms (Ritson-Williams et al., 2006; Hwang

et al., 2007; Williams, 2010). Bacterial symbionts of toxic newts and marine organisms

produce TTX for their hosts, suggesting a bacterial origin in most if not all of these

organisms (Magarlamov et al., 2017; Vaelli et al., 2020). TTX is a neurotoxin that blocks

voltage-gated sodium channels by binding to highly conserved regions of the channel

via hydrophobic forces and hydrogen bonds, fitting in the opening of the channel and

blocking proper transport of Na$^+$ ions (Kao and Levinson, 1986; Tikhonov and Zhorov,

2011). TTX binding prevents action potentials from propagating, resulting in cessation of

nerve signals (Kao and Levinson, 1986; Tikhonov and Zhorov, 2011). In vertebrate

animals lacking TTX resistance, ingested TTX can temporarily paralyze or, in some

cases, kill the organism. The poison usually becomes fatal when respiratory muscles

fail, resulting in suffocation (Brodie, 1968).

While TTX can incapacitate or kill an animal that attempts to consume it, several

species are able to come into contact with TTX and survive. Resistance to TTX

intoxication usually evolves via one of three common mechanisms. In the first, amino

acid substitutions in voltage-gated sodium channels (VGSCs), especially in domain IV

(DIV) of $Na_v1.4$, alter the shape of the pore loop (p-loop) and the surrounding area,

which prevents TTX from binding (Geffeney et al., 2005; Tikhonov and Zhorov, 2005).

Other regions of $Na_v1.4$, such as domain III, and other voltage-gated sodium channels,

such as $Na_v1.7$, also harbor amino acid substitutions implicated in TTX resistance

(Feldman et al., 2012; McGlothlin et al., 2016). Changes in the p-loops are particularly

significant because they not only alter the sensitivity of the sodium channel to TTX but

also to sodium ions; hence, there is often a tradeoff between TTX resistance and

protein function (Chiamvimonvat et al., 1996; Geffeney et al., 2005; Lee et al., 2011;

Tikhonov and Zhorov, 2011; Hague et al., 2018; but see Moniz et al., 2021). Thus,

amino acid changes are usually confined to specific conserved regions that can provide

resistance while allowing the organism to maintain sodium channel functionality

(Feldman et al., 2012). The presence of amino acid substitutions that alter toxin binding

is a mechanism known as target-site insensitivity and is the focus of this study.

Other mechanisms of TTX resistance include toxin-binding proteins, which have

evolved in gastropods, pufferfish, horseshoe crabs, and shore crabs (Ho et al., 1994;

Nagashima et al., 2002; Hwang et al., 2007; Hashiguchi et al. 2015), and diffusion

barriers, which are present in some invertebrates (Mebs et al., 2016). Some amphibians

possess the toxin-binding, blood-plasma protein saxiphilin, but saxiphilin is incapable of

binding TTX, at least in the animals that have been tested (Mahar et al., 1991; Llewellyn

et al., 1997; Abderemane-Ali et al., 2021). Some species of mantids (*Tenodera*

sinensis, *Sphodromantis viridis*, *Hierodula membranacea*, and *Miomantis caffra*) can

survive tetrodotoxin ingestion but not toxin injection, suggesting the presence of a gut

diffusion barrier (Mebs et al., 2016). Crayfish appear to have a similar mechanism

(Wilson et al., 2014). Because we do not directly test for toxin-binding proteins or

diffusion barriers, we cannot say whether *P. regilla* possesses these mechanisms of

resistance.

Taricha newts (*Taricha torosa*, *T. granulosa*, *T. sierrae,* and *T. rivularis*, herein

collectively referred to as *Taricha*) have TTX present on their skin (Brodie et al., 1974;

Reimche et al., 2020) that likely functions as an anti-predator defense (Brodie et al.

2002). *Taricha* (and other salamandrids) possess amino acid changes in $Na_v1.4$ that

help them resist their own TTX defenses (Hanifin and Gilly, 2015; Gendreau et al.,

2021). The San Francisco Bay Area in California is one of two known hotspots of the

arms race between newts and garter snake predators (Brodie et al., 2002). Here we

explore the idea that high levels of TTX defenses in newts may select for resistance in

other sympatric organisms outside the predator-prey dynamic, namely the Pacific

Chorus Frog, *Pseudacris regilla*.

TTX from *Taricha* skin has been shown to enter waterways and affect the

behavior of organisms present in the same aquatic systems. Larval *T. torosa* use TTX

as a cue to detect and avoid adult *T. torosa* that sometimes cannibalize larvae (Zimmer

et al., 2006). TTX in water can also slow predatory patterns of dragonfly nymphs, which

are macroinvertebrate prey of *T. torosa* (Bucciarelli and Kats, 2015). The New Zealand

Mud Snail (*Potamopyrgus antipodarum*) moved out of its natural stream environment

when encountering biologically relevant levels of TTX or chemical cues from *T. torosa*

containing TTX (Ota et al., 2018). Water containing TTX also negatively increases newt

trematode parasite mortality (Calhoun et al., 2017). Thus, naturally occurring levels of

TTX produced by *T. torosa* can affect other organisms in the same aquatic environment.

Given that *P. regilla* have permeable skin, we hypothesized that they would be

exposed to TTX through physical contact with newts or by occupying pond water that

contains newts. We predicted that TTX exposure has selected for TTX resistance in *P.*

regilla populations that are microsympatric with *Taricha*. We recognize that our

hypothesis makes a simplifying assumption that the designation of *P. regilla* populations

as sympatric or allopatric with *Taricha* is relevant and valid during the time in which

selection may have occurred for TTX resistance, and that gene flow does not occur

between sympatric and allopatric *P. regilla* populations. Although both of these

assumptions may be violated by changes in climate and species distributions over time,

evidence from the evolutionary arms race between garter snakes and TTX-bearing

newts suggest that TTX potency and the degree of TTX resistance can change over a

short period of time (Brodie et al., 2002), which aligns with our study that examines

approximately the past 50 years. Occasional exposure to TTX may also be sufficient to

select for target-site insensitivity (Durso et al., 2021).

We aimed to test whether the sympatry of *Taricha* newts with *Pseudacris regilla*

has selected for target-site insensitivity in the skeletal muscle voltage-gated sodium

channel of *P. regilla*. While *P. regilla* do not regularly prey upon *Taricha* (or vice versa),

they likely come into contact with TTX in pond water shared with *Taricha* or via direct

contact with *Taricha*, sometimes in mistaken amplexus behavior or even attempted

predation (Fig. 1).

Our study centered on California populations, examining *Taricha* and *P. regilla*

that do and do not overlap geographically in the present. We conducted demographic

surveys of *Taricha granulosa, T. torosa,* and *P. regilla* at three San Francisco Bay Area

ponds where the species co-occur to estimate the extent to which *P. regilla* may

encounter TTX. Because these surveys were suspended after only a few months due to

the COVID-19 pandemic, we then reviewed occurrence data available through the

Global Biodiversity Information Facility (GBIF) to better estimate overlap of annual

activity patterns. To determine whether *P. regilla* has structural modifications in the

protein muscle voltage-gated sodium channel $Na_v1.4$ that may provide TTX resistance,

we sequenced and reviewed amino acid residues encoded by the SCN4A gene. Amino

acid substitutions in this domain are known to confer resistance to TTX and have been

found in several organisms that either have TTX defense or consume TTX-defended

prey (Tikhonov and Zhorov, 2005; Venkatesh et al., 2005; Feldman et al., 2012; Hanifin

and Gilly, 2015). We hypothesized that if the presence of *Taricha* selects for TTX

resistance in *P. regilla,* populations sympatric with *Taricha* would have mutations in the

p-loop of $Na_v1.4$ while populations that are allopatric would not. Our work provides

insight into the potential cascading ecological consequences of chemical defenses, as

we evaluate the unintended effects of animals wielding chemical defenses.

172   MATERIALS AND METHODS

Field Work.—We studied *Taricha* and *P. regilla* at four San Francisco Bay Area

ponds. We conducted surveys at three of these sites where *Taricha granulosa, T.*

torosa, and *P. regilla* co-occur: two in Briones Regional Park in Martinez, Contra Costa

County, CA (Old Briones Road Trail 1 [OBRT 1] at 37.94571, -122.13395, and OBRT 2

at 37.94352, -122.14102, named after the trail closest to the sites) and the Japanese

Pool (37.87440, -122.23760) at the University of California Botanical Garden (UCBG) in

Berkeley, Alameda County, CA (Fig. 2). OBRT 1 and OBRT 2 are in an unshaded area

of grazed open grassland that lies within an oak woodland habitat, and they connect to

each other when water levels are high. The UCBG Japanese Pool is a man-made pool

with a concrete floor and sides, located under tree cover within the visitor area of the

UCBG (Fig. 2). We selected these three ponds because they were known to contain

populations of *P. regilla* as well as *Taricha torosa* and *Taricha granulosa* (Katherine

Montana, Rebecca Tarvin, pers. obs.). We also collected specimens from a fourth pond

located in the Presidio of San Francisco (37.788496, -122.468605) that has historically

contained *P. regilla* but no *Taricha* (records show absence of *Taricha* since at least

1966; pers. comm. J. Young, Presidio Trust). Mountain Lake is a natural lake

surrounded by grass and receives run-off from the nearby Presidio Golf Course (Fig. 2).

From October 2019 to February 2020, we conducted six surveys at the two Briones

ponds and five surveys at UCBG, with surveys at each site occurring three to four

weeks apart from one another. We were unable to gather abundance data for more than

this time period because the parks were closed from February until October 2020 during

the early months of the COVID-19 pandemic.

195   Two researchers (usually K. Montana and V. Ramírez Castañeda) conducted

surveys in the morning or early afternoon (between 8:00 AM and 1:30 PM). During each

survey, surveyors split the pond perimeter into 7.62-m (25-feet) sections, and circled the

pond on foot, counting how many individuals of each species were visible in the water

from the shore within each transect. We note that surveyors were only able to count the

number of visible newts; OBRT 1 and OBRT 2 were murky with plants and algae in

which newts tended to hide and although the surface of the Japanese Pool at UCBG

was glassy, the pool was filled with plants that could have obscured observations.

Surveyors also counted the numbers of visible, individual adult *Pseudacris regilla*; *P.*

regilla that were calling but not visible were not counted because surveyors could not

ascertain if multiple calls were coming from a single or a few individuals. Though the

TTX defenses of *T. granulosa* are thought to be more potent than that of *T. torosa*

(Brodie et al., 1974), for the purposes of this study, we did not differentiate between the

two species when we recorded their occurrences because they both contribute to the

presence of TTX in pond water, and we could not easily identify individuals to species

from the pond shores. We counted large (~5 cm or longer) *Taricha* larvae, adult *Taricha,*

and *Taricha torosa* egg masses, which are conspicuous groups of eggs known to

possess TTX via vertical transfer from the parent (Gall et al., 2012). The much smaller

T. granulosa eggs are typically hidden in the substrate and are more difficult to see;

hence we did not include them in our survey. To simplify our survey protocol, we

counted larval *Taricha*, which also have some amount of TTX (Gall et al., 2011),

together with adults. We plotted survey results using R version 4.1.1 and the packages

ggplot2 v.3.3.5 and ggpubr v.0.4.0 (Wickham, 2016; Kassambara, 2020; R Core Team,

2021) (Fig. 3). We compared survey results to GBIF occurrence data of *P. regilla* and

Taricha from 1 January 1971 to 23 November 2021 by plotting the GBIF data using the

following R packages: RColorBrewer v.1.1.2, ggplot2 v.3.3.5, chron v.2.3.56, cowplot

v.1.1.1, and dplyr v.1.0.7 (Neuwirth, 2014; Wickham, 2016; James and Hornik, 2020;

Wilke, 2020; Wickham et al., 2021).

After COVID-19 conditions permitted field work to continue in October 2020, we

collected one *Pseudacris regilla* from the dry bed of OBRT 2 and eight *P. regilla* from

the east arm of Mountain Lake in the Presidio of San Francisco. We surveyed OBRT 1

and UCBG at the same time but did not find any *P. regilla* at either location; OBRT 1

was dry.

Precipitation Data.—To better interpret amphibian abundance data from field

surveys, we searched for precipitation data from NOAA for the weather stations closest

to our field sites in Berkeley (UCBG) and Martinez (OBRT 1 and OBRT 2): Berkeley, CA

USC00040693 and Martinez 2.2 SW, CA US1CACC0011 (NOAA,

https://www.ncdc.noaa.gov/cdo-web/search). We measured pond size during each

survey event by adding the number of 7.62-m transects we walked during each survey

to get the circumference and calculate the area, assuming the ponds were circular.

Notes on Taxonomy.—The taxonomy of *Pseudacris regilla* has changed several

times over the last decades (Recuero et al., 2006; Fouquette and Dubois, 2014;

Duellman et al., 2016). iNaturalist categorizes some *Pseudacris regilla* (e.g., in the San

Francisco Bay Area) as *Pseudacris sierra*. However, these distinctions are not

universally accepted by the herpetology community (Barrow et al., 2014; Crother et al.,

2017). Here we consider *P. sierra*, *P. hypochondriaca*, and *Hyla regilla* synonyms of

Pseudacris regilla. We downloaded occurrence data from GBIF for the *P. regilla*

synonyms listed here and for all currently recognized species of *Taricha* (*T. granulosa*,

T. rivularis, *T. sierrae*, *T. torosa*), as well as those no longer recognized (*T. lindoei*, *T.*

miocenica, and *Palaeotaricha oligocenica*) and individuals identified only to the genus

Taricha.

DNA Sequencing.—We selected 29 liver tissues of *Pseudacris regilla* from the

Museum of Vertebrate Zoology at the University of California, Berkeley to broaden our

geographic sampling (Table S1). We classified *P. regilla* samples as sympatric with

Taricha if they were collected within 35 km or 500 m of elevation from the nearest

Taricha occurrence documented on GBIF as of 23 November 2021; this included 11

individuals and the single *P. regilla* collected at OBRT 2. We classified allopatric *P.*

regilla samples as those individuals collected at least 35 km or 500 m in elevation away

from the nearest *Taricha* observation; this included 18 individuals and the eight *P.*

regilla from the Mountain Lake field site in the Presidio (Table S1). We predicted that

Na$_v$1.4 sequences from sympatric, but not allopatric, *P. regilla* would show amino acid

substitutions potentially conferring TTX resistance. We utilized the R packages ggplot2

v.3.3.5 and tidyverse v.1.3.1 to map GBIF occurrences of *Taricha* throughout California,

P. regilla museum samples that were sympatric with *Taricha*, *P. regilla* museum

samples that were allopatric with *Taricha*, and our field sites (Wickham, 2016; Wickham

et al., 2019; GBIF).

Our preliminary results suggested that a single amino acid substitution is present

in *P. regilla.* Thus, to estimate when this substitution evolved, we also sequenced

Na$_v$1.4 from two close relatives of *P. regilla, Pseudacris cadaverina* and *Acris crepitans*

(Table S1). The individuals of *P. cadaverina* and *A. crepitans* are from locations far from

Taricha and, thus, were likely not exposed to TTX from *Taricha*. However, the range of

A. crepitans overlaps with another TTX-bearing newt, *Notophthalmus viridescens*

(Baird, 1854; Mecham, 1967), so it may be exposed to TTX as well. Thus, if TTX from

newts selects for mutations in Na$_v$1.4 in sympatric frog species, we would expect to find

amino acid substitutions in Na$_v$1.4 of *A. crepitans* as well as in Na$_v$1.4 of *P. regilla*

sympatric with *Taricha*. In contrast, we expected to find no resistance-conferring amino

acid substitutions in allopatric *P. regilla* or in *P. cadaverina*.

In the Evolutionary Genetics Lab (UC Berkeley), we extracted DNA from tissues

according to the Miller et al. (1988) salt-extraction protocols, with modifications (see

Supplementary Online Material for details). The primers used for *P. regilla* samples

targeted exon 24 (DIV segment 6 and p-loop) of the gene SCN4A: 2268F_SCN4A 5'-

TCTCCCGGCCCTCTTCAATA-3' and 2681R_SCN4A 5'-

TCGTCCTCGCATAAAGGCTC-3' (Tarvin et al., 2016). We added 0.5 µL of $MgCl_2$ to

each PCR sample. PCR protocols were: an initial denaturation for 2 min at 94 ºC; 35

cycles of: 94 ºC at 30 s for denaturation, an annealing temperature of 54 ºC for 30 s,

and an elongation temperature of 72 ºC for 45 s; and a final extension time of 7 min at

72 ºC. PCR products were run on an agarose gel, purified with ExoSAP-IT PCR Product

Cleanup Reagent (ThermoFisher Scientific, 78201.1.ML) and sequenced using the

reverse primer 2681R_SCN4A at the UC Berkeley DNA Sequencing Facility. For PCR

and sequencing samples of *Pseudacris cadaverina* and *Acris crepitans*, we used the

same protocol and forward primer, but the reverse primer was redesigned based on the

DIV SCN4A *P. regilla* sequences we produced: P.regilla_SCN4AD4_R 5'-

ACTGCTTTCTTCTGTGGCCAC-3'.

We also obtained previously published SCN4A sequences from species that are

resistant to TTX, including *Erythrolamprus epinephelus*, which is potentially resistant to

TTX from *Atelopus*, as well as *Rhabdophis tigrinus*, *Thamnophis elegans*, *Thamnophis*

sirtalis, *Taricha torosa*, and *Taricha granulosa*, which are either confirmed or

hypothesized to possess target-site insensitivity to TTX via mutations in $Na_V1.4$

(Feldman et al., 2009; Feldman et al., 2012; Hanifin and Gilly, 2015). We also include

sequences from *Nanorana parkeri* and *Rattus norvegicus,* which have not been

documented as being resistant to TTX (Table S2). We identified the location of the DIV

p-loop in $Na_V1.4$ based on protein domains inferred by Tikhonov and Zhorov (2005). We

aligned sequences using Clustal Omega in Geneious (Sievers et al., 2011, Geneious

Prime 2020.1.1). We reviewed the alignment by eye for amino acid differences between

sequences, focusing on residues that have been shown to provide TTX resistance. We

estimated a phylogeny of the species that we compared to *P. regilla* using 11 COI

sequences (Table S2) and maximum likelihood on the IQTree web server, with two

partitions (one for codon positions 1 and 2, another for codon position 3), the GTR + G

substitution model, and support assessed using 1000 ultrafast bootstraps and the SH-

aLRT test (Nguyen et al., 2015; Chernomor et al., 2015; Trifinopoulos et al., 2016;

Hoang et al., 2018; IQ-Tree web server http://iqtree.cibiv.univie.ac.at/). We transformed

the resulting phylogeny into an ultrametric chronogram using the function *chronos* in

phytools v.0.7.90 and ape v.5.5 (Revell, 2012; Paradis and Schliep, 2019) (Fig. 6). We

made all figures in R and Adobe Illustrator (Adobe, Inc., 2019; R Core Team, 2021).

313   RESULTS

Demographic surveys indicated that *P. regilla* may encounter TTX from *Taricha* in their

environment. DNA sequencing identified one amino acid substitution in domain IV of

$Na_v1.4$ in *P. regilla* that could potentially be connected to TTX resistance, but the

degree to which the mutation affects resistance is inconclusive.

Field Surveys.—While we never observed *P. regilla* at OBRT 2 or UCBG, their

abundance at OBRT 1 peaked early in the observation period at 32 individuals on the

first field trip date (20 October 2019). For OBRT field sites, *Taricha* abundance peaked

at the beginning of February; for the UCBG, abundance was low, and we may not have

observed the seasonal peak (surveys were suspended at the end of February 2020

because of the COVID-19 pandemic). *Taricha torosa* egg masses were most abundant

at the beginning of February 2020 at OBRT 1 and OBRT 2, while they were most

abundant at UCBG at the end of February (though we cannot be certain these are true

peaks because our surveys ended before March) (Fig. 3). California-wide GBIF

observations for *P. regilla* and *Taricha* from 1 January 1971 to 23 November 2021

indicate two peaks of *P. regilla* abundance in April and August, while *Taricha* peaks in

January (Fig. 4). Though the abundance peaks do not occur at precisely the same time,

P. regilla and *Taricha* do overlap with one another and, thus, the species have

opportunities to interact such that *Taricha* could expose *P. regilla* to TTX. These data

are summarized across California and thus represent average activity across a range of

habitat types. Nevertheless, when the data are constrained to a square roughly

representing the San Francisco Bay Area (N 38.66226, S 37.09298, E -121.12701, W -

123.23579), the results are similar, indicating a peak for *Taricha* in January and a peak

for *P. regilla* in March (Supplementary Fig. 1).

When we surveyed the field sites in Briones Regional Park and the UC Botanical

Garden from October 2019 to March 2020, California was experiencing drought. During

this period, rain began in mid-November 2019 and stopped at the end of February 2020.

The most rain recorded in one day during the six-month period in Berkeley, where

UCBG is located, was 30.23 mm, the average recorded rainfall during that period was

1.76 mm/day, and the total number of recorded days with rain was 40 days (NOAA field

station Berkeley, CA USC00040693, elevation 94.49 m, 37.8744° N, 122.2605° W)

(NOAA, 2022). The most rain recorded in one day during the six-month period in

Martinez, where Briones is located, was 89.15 mm, the average recorded rainfall during

that period was 1.61 mm/day, and the total number of recorded days with rain was 33

days (NOAA field station Martinez 2.2 SW, CA US1CACC0011, elevation 210.92 m,

37.9744° N, 122.1384° W) (NOAA, 2022). The area of the ponds fluctuated due to

changes in precipitation and, in the case of the Japanese Pool, some drainage for

maintenance (UCBG personnel, pers. com.). The area of OBRT 1 ranged between 319

and 912 m^2 with a survey perimeter of about 84-122 m. The area of OBRT 2 ranged

between 1550 and 1733 m^2 with a perimeter of about 130-145 m. During our study

period, OBRT 1 and OBRT 2 did not merge. The area of the Japanese Pool ranged

between 114 and 137 m^2 with a perimeter of about 46-53 m.

Amino Acid Changes.— We found one amino acid change (I1519M) within the

DIV p-loop of the sodium channel protein Na$_v$1.4 in *Pseudacris regilla* compared to a

sequence of *Rattus norvegicus* (a common species used in comparative TTX studies

[e.g., Tikhonov and Zhorov, 2005; Tarvin et al. 2016] as it represents the mammalian

TTX-sensitive Na$_v$1.4 [Chahine et al., 1994]) (Fig. 6) (Tables S1 and S2). Importantly,

there were no observed differences in amino acid sequences between *P. regilla*

specimens that were sympatric and allopatric with *Taricha*. The I1519M substitution has

been previously predicted by computational modeling to provide some amount of TTX

resistance (Tikhonov and Zhorov, 2005; Tikhonov and Zhorov, 2011). The I1519M

substitution is also present in the snake *Rhabdophis tigrinus,* which consumes TTX-

defended treefrogs, and is inferred to be associated with TTX resistance (Feldman et

al., 2012). However, it has been suggested that the characterization of the I1519M

substitution as contributing to TTX resistance is erroneous because the same

substitution is found in TTX-sensitive voltage-gated sodium channels of other snakes

(McGlothlin et al., 2016). Additionally, we found that the amino acid change I1519M

occurs in the frog *Nanorana parkeri,* which is not known for being resistant to or in

contact with tetrodotoxin; though, it does occur within the range of TTX-defended

salamandrids such as *Tylotritron verrucosus* (Che et al., 2010; Vitt and Caldwell, 2013;

Wanninger et al., 2018). Close relatives of *P. regilla* (*P. cadaverina* or *A. crepitans*) did

not possess the I1519M substitution, suggesting that the mutation arose within *P. regilla*

or its ancestor.

379   DISCUSSION

Our results provide conflicting evidence regarding whether $Na_v1.4$-based TTX

resistance is present in *Pseudacris regilla*. In support of resistance is the presence of a

single amino acid substitution in *P. regilla* $Na_v1.4$ (Fig. 6) hypothesized to provide TTX

resistance based on computational models (Tikhonov and Zhorov, 2005; Tikhonov and

Zhorov, 2011), and evidence that *P. regilla* and *Taricha* overlap somewhat in annual

activity patterns (Fig. 4). However, amino acid changes in important regions of the

$Na_v1.4$ protein do not always translate to a resistant phenotype (e.g., see Abderemane-

Ali et al., 2021). Furthermore, both allopatric and sympatric *P. regilla* possessed the

substitution, but *A. crepitans* did not, although we expected that it might possess some

resistance due to its overlap in range with *N. viridescens*. The presence of an identical

substitution in *Nanorana parkeri* was unexpected and could either present evidence in

support of resistance (the species does live near the TTX-defended salamandrid

Tylototriton verrucosus), or against it, potentially indicative of functionally permissive

amino acid diversity at that site in anurans. Below we review in more detail these lines

of evidence.

395   Abundance data indicate that *P. regilla* and *Taricha* annual activity patterns

overlap but that peak breeding times are not synchronized (Figs. 3, 4; Supplementary

Fig. 1). Observations of mistaken amplexus and predation of newts by *P. regilln* indicate

that the species do come into contact (Fig. 1). Still, TTX exposure may not be frequent

or potent enough for *P. regilla* to undergo strong selection for resistance to TTX. In

cases where environmental selective regimes are not predictable, more plastic

responses like upregulation of detoxifying enzymes may be the primary way animals

deal with toxin exposure (Zanger and Schwab, 2013). Nevertheless, even occasional

exposure to TTX can result in maintenance of target-site insensitivity (Durso et al.,

2021).

405   We hypothesized that we would find TTX-conferring amino acid substitutions in

Na$_v$1.4 in *P. regilla* that are microsympatric with *Taricha* and no substitutions in *P. regilla*

that are allopatric with *Taricha*, with a greater number of substitutions indicating a higher

level of resistance as has been shown in salamanders and snakes (Geffeney et al.,

2005; Lee et al., 2011; Hanifin and Gilly, 2015). Compared to *Rattus norvegicus* DIV,

we found one amino acid change at a site hypothesized to provide TTX resistance,

I1519M, which is a TTX-sensing residue located in the p-loop (Tikhonov and Zhorov,

2005; Tikhonov and Zhorov, 2011; Feldman et al. 2012). This substitution was present

in *P. regilla* both sympatric and allopatric with *Taricha* and in *Nanorana parkeri*. It is also

found in *Rhabdophis tigrinus*, a species of snake that has been hypothesized to be

resistant to TTX (Feldman et al., 2012), although no empirical tests have been

undertaken and other authors have disputed the role of I1519M in TTX resistance

(McGlothlin et al., 2016; noted as I1555M in these studies). The presence of this

substitution in all *P. regilla* samples does not support our hypothesis that substitutions

would only be found where *P. regilla* is exposed to TTX from sympatric *Taricha* (at least

based on occurrence data from the last 50 years). Ultimately, our study suggests that

the I1519M substitution evolved in the ancestor of *P. regilla*, which likely rules out

contemporary selection from *Taricha.* The presence of the substitution in *N. parkeri*

could also support that the mutation is unrelated to newt exposure. However, *N. parkeri*

overlaps with other toxin-defended salamandrids like *T. verrucosus,* making it difficult to

rule out resistance in *N. parkeri* or, consequently, in *P. regilla*. Similarly, *A. crepitans* co-

occurs with the toxic newt *Notophthalmus viridescens,* but this species does not

possess the I1519M substitution. Because there is disagreement whether the I1519M

mutation provides resistance, we cannot say with certainty whether it provides TTX

resistance in *P. regilla* without further experimental evidence.

While there is not strong evidence that *P. regilla* possesses target-site

insensitivity in the p-loop of DIV in $Na_v1.4$, it is plausible that *P. regilla* may have other

ways of avoiding the effects of TTX, including amino acid mutations in other domains of

Na$_v$1.4 or in other voltage-gated sodium channels (Feldman et al., 2012; McGlothlin et

al., 2016). *Pseudacris regilla* could also possess the ability to smell TTX, as larval

Taricha can, in order to avoid it (Zimmer et al., 2006). Alternatively, they could possess

a diffusion barrier in the skin or the gut that would prevent high levels of TTX exposure,

as occurs in mantids (Mebs et al., 2016). It is also possible that *P. regilla* possesses a

binding protein that can scavenge TTX; other frogs have saxiphillin, a protein that binds

saxitoxin, which is structurally similar to TTX (Mahar et al., 1991; Llewellyn et al., 1997).

Similarly, little is known about the potential interactions between *Acris crepitans* and

Notophthalmus viridescens, which could also select for TTX resistance in the frogs. To

date, none of these alternative resistance mechanisms have been studied in

Pseudacris, Acris, or *Notophthalmus.*

A number of reasons could explain why *P. regilla* would lack target-site

insensitivity in Na$_v$1.4. Evolving TTX resistance could be costly to *P. regilla* due to

possible alterations in the selectivity filter of sodium channels that frequently occurs

when amino acid substitutions are present (Chiamvimonvat et al., 1996; Geffeney et al.,

2005; Lee et al. 2011; Tikhonov and Zhorov, 2011; Moniz et al., 2021). Evolving costly

mutations may require a stronger selective pressure than is currently occurring. Given

the relative quantities of TTX on newt skin [up to 4.69 mg/newt (Hanifin et al., 2008)]

compared to TTX levels in newt-containing water [~1×10^{-7} mol/L (Zimmer et al.,2006)],

the environmental toxicity of TTX is likely to cause a weaker selection for TTX

resistance than exposure via chemical defenses from prey. The cost of exhibiting TTX

resistance combined with the low potential for *P. regilla* to interact with high

concentrations of TTX in its environment may explain why we did not unequivocally

identify target-site insensitivity in *P. regilla* DIV Na$_v$1.4. Future work exploring the effects

of TTX as an environmental toxin could include measuring TTX concentrations in newt-

containing ponds, performing whole organism or *in vitro* assays of *P. regilla* to

determine TTX sensitivity, and testing for other modes of TTX resistance in animals that

live in sympatry with salamandrid newts.

461
Conclusions.—We do not find strong evidence for target-site insensitivity in DIV

of the skeletal muscle voltage-gated sodium channel Na$_v$1.4 of *Pseudacris regilla*.

Although it is possible that the I1519M substitution in the p-loop of Na$_v$1.4 may be

related to TTX resistance, *P. regilla* may not encounter TTX from *Taricha* with high

enough frequency or magnitude to select for TTX resistance. Thus, these animals may

have behavioral or alternative physiological mechanisms to survive TTX exposure, or

TTX exposure may not be severe enough to require adaptation by *P. regilla*.

Investigating the environmental toxicity of TTX in the relationship between P. *regilla* and

Taricha is an opportunity to explore the nexus of ecological interactions, neurotoxins,

and resistance mechanisms. The likely absence of target-site insensitivity in *P. regilla*

indicates that exposure to tetrodotoxin outside of trophic interactions may not strongly

select for fixed mechanisms of toxin resistance.

Acknowledgments.—We affirm the sovereignty of the Karkin, Me-Wuk (Bay

Miwok), Muwekma, Chochenyo, Ohlone, and Ramaytush people and the Confederated

Villages of Lisjan; we thank them for stewarding the land where we worked and, thus,

serving an integral role in the production of this study. We thank L. Smith in the

Evolutionary Genetics Lab, as well as C. Spencer and the support of the Museum of

Vertebrate Zoology collections staff. We also thank those who supported our work at

field sites: UC Botanical Garden staff, including H. Forbes, J. Fong, E. Hupperts, and C.

Loughran; East Bay Regional Parks District staff, including T. Lin and J. Miller; and J.

Young with the Presidio Trust. We thank the other members of the Tarvin Lab for their

editing prowess and general support. We thank A. Shabel for his advice in the early

stages of this project, and K. Sanko, J. Toman, and L. Galindo who helped as field

assistants. We appreciate the newts and frogs who were the focus of this project and

who were sacrificed to offer the knowledge they hold. K. Montana thanks her family and

ancestors for their guidance. R. Tarvin was supported by UC Berkeley start-up funds.

The East Bay Regional Parks District authorized the surveys at Briones under

permit #19-1044, and staff at the UCBG gave written consent for surveys conducted on

the premises. All animals were collected under approved collection permits and

protocols: for the National Parks Service Scientific Research and Collecting Permit 2

number PRSF-2020-SCI-0002 within study PRSF-00035, with a written extension after

expiration of the original permit, University of California IACUC AUP-2019-08-12457,

and California Department of Fish and Wildlife Scientific Collection Permit S-

190980001-19111-001.

Spanish translation was done using www.DeepL.com/Translator (free version),

which was then edited by Valeria Ramírez Castañeda.

500   LITERATURE CITED

501   (see attached document)

503   SUPPLEMENTARY ONLINE MATERIAL

Supporting Methods.—Specimens sequenced in this study, SCN4A exon 24

Table S1.—DNA Extraction Procedures, Adapted from Miller et al., 1988.

Table S2.— Na$_v$1.4 (SCN4A exon 24) and COI sequences from GenBank

Supplementary Figure 1.—San Francisco Bay Area GBIF Observations of

Pseudacris regilla and *Taricha*, 1971-2021 (N 38.66226, S 37.09298, E -121.12701, W -

123.23579)

F IGURES

Fig. 1. Examples of *Pseudacris regilla* and *Taricha torosa* interactions: A. Amplexus in

Briones Regional Park, Martinez, CA, 1 March 2019, photo credit K. Montana, user

kmontana on iNaturalist (CC BY-NC 4.0; observation 100725148); B. Another instance

of amplexus in Briones Regional Park, Martinez, CA, 21 February 2009, photo credit

user ap2il on iNaturalist (CC BY-NC 4.0; observation 550227); C. Attempted predation

on *Taricha torosa* by *Pseudacris regilla* in Mark West Springs, CA, 1 March 2014, photo

credit user tiwane on iNaturalist (CC BY-NC 4.0; observation 2312). The frogs did not

appear to be harmed in any of these scenarios, though the tongue posture in C

suggests some numbing effects from newt exposure.

Fig. 2. Field sites: A. Old Briones Road Trail 1 (OBRT 1) pond, 37.94441, -122.13377;

B. Old Briones Road Trail 2 (OBRT 2) pond, 37.94352, -122.14102; C. Japanese Pool

at the UC Botanical Garden (UCBG), 37.87440, -122.23760; D. Presidio site,

37.788496, -122.468605; all photos taken by K. Montana.

Fig. 3. Numbers of A. *P. regilla* adults, B. *Taricha torosa* and *T. granulosa* adults and

larvae, and C. *T. torosa* egg masses observed over field trips to the Old Briones Road

Trail 1 (OBRT 1) (blue circle line), OBRT 2 (red triangle line), and U.C. Botanical

Garden (yellow square line) ponds; 20 October 2019 to late February 2020.

Fig. 4. Percentage of total abundance of GBIF-recorded individuals of *Pseudacris regilla*

(green) and *Taricha* (blue) in California by month, over the course of the last 50 years.

Taricha includes *T. granulosa*, *T. rivularis*, *T. sierrae*, *T. torosa*, as well as those *Taricha*

species no longer recognized (*T. lindoei*, *T. miocenica*, and *Palaeotaricha oligocenica*)

plus individuals just identified to the genus *Taricha* (GBIF). Peak abundances for *P.*

regilla occur around April and August each year, and peak abundances for *Taricha*

occur in January each year. Samples included were from 1 January 1971 to 23

November 2021.

Fig. 5. Map of the state of California, USA showing location of *Pseudacris regilla*

museum specimens sympatric with *Taricha* (green squares), *P. regilla* museum

specimens allopatric with *Taricha* (orange triangles), and field sites (black squares) in

relation to occurrences of *Taricha* (blue circles).

Fig. 6. Alignment of the p-loop of domain IV of the voltage-gated sodium channel Na$_v$1.4

in several species including *Pseudacris regilla,* mapped onto a COI phylogeny inferred

using IQTREE (support values are SH-aLRT/ultrafast bootstrap values from 1000

iterations each). Amino acid positions are numbered according to *Rattus norvegicus,*

UniProt accession #P15390. Species marked with an asterisk* are known to be

resistant to TTX, and the AA substitutions known or hypothesized to confer TTX

resistance are shown in bold (Feldman et al., 2012; Hanifin and Gilly, 2015). The amino

acid substitution I1519M is found in *P. regilla* as well as *N. parkeri* and *R. tigrinus. R.*

tigrinus consumes *Polypedates leucomystax*, which wields tetrodotoxin and

bufadienolides; the I1519M substitution is thought to be linked to TTX resistance

(Hutchinson et al., 2007; Tikhonov and Zhorov, 2005; Tikhonov and Zhorov, 2011). *N.*

parkeri has the I1519M substitution but is not known to consume or be resistant to any

toxins. Close relatives of *P. regilla*, *P. cadaverina* and *A. crepitans,* do not have the

I1519M substitution or any others. *E. epinephelus* has AA substitutions that are thought

to be connected to resistance to other toxins in their diets; they eat *Atelopus* toads that

have guanidine alkaloids including tetrodotoxin as well as dendrobatid poison dart frogs

that wield batrachotoxin and pumiliotoxin (Myers et al., 1978; Feldman et al., 2012;

Ramírez Castañeda et al., 2017).

ABDEREMANE-ALI, F., N.D. ROSSEN, M.E. KOBIELA, R.A. CRAIG II, C.E. GARRISON, Z. CHEN,

C.M. COLLERAN, L.A. O'CONNELL, J. DU BOIS, J.P. DUMBACHER, AND D.L. MINOR, JR.

2021. Evidence that toxin resistance in poison birds and frogs is not rooted in

sodium channel mutations and may rely on 'toxin sponge' proteins." Journal of

General Physiology 153:e202112872. https://doi.org/10.1085/jgp.202112872.

Adobe Inc. 2019. Adobe Illustrator. Retrieved from

https://adobe.com/products/illustrator.

BAIRD, S.F. 1854. Descriptions of new genera and species of North American frogs.

Proceedings of the Academy of Natural Sciences of Philadelphia 7:59–62.

BARROW, L.N., H.F. RALICKI, S.A. EMME, AND E.M LEMMON. 2014. Species tree estimation

of North American chorus frogs (Hylidae: *Pseudacris*) with parallel tagged

amplicon sequencing. Molecular Phylogenetics and Evolution 75:78–90.

BRODIE, E.D., JR. 1968. Investigations on the skin toxin of the adult rough-skinned

newt, *Taricha granulosa*. Copeia 1968:307–313.

BRODIE, E.D., JR., J.L. HENSEL, AND J.A. JOHNSON. 1974. Toxicity of the urodele

amphibians *Taricha*, *Notophthalmus*, *Cynops* and *Paramesotriton*

(Salamandridae). *Copeia* 1974:506-511.

BRODIE, E.D., JR. B.J. RIDENHOUR, AND E. D. BRODIE III. 2002. The evolutionary

response of predators to dangerous prey: hotspots and coldspots in the

geographic mosaic of coevolution between garter snakes and newts.

Evolution 56:2067–2082.

BUCCIARELLI, G.M., AND L.B. KATS. 2015. Effects of newt chemical cues on the

distribution and foraging behavior of stream macroinvertebrates. Hydrobiologia

749:69–81.

CALHOUN, D.M., G.M. BUCCIARELLI, L.B. KATS, R.K. ZIMMER, AND P.T.J. JOHNSON. 2017.

Noxious newts and their natural enemies: experimental effects of tetrodotoxin

exposure on trematode parasites and aquatic macroinvertebrates. Toxicon:

Official Journal of the International Society on Toxinology 137:120–127.

CHAHINE, M., P.B. BENNETT, A.L. GEORGE, AND R. HORN. 1994. Functional expression and

properties of the human skeletal muscle sodium channel. Pflugers Archive:

European Journal of Physiology 427:136–142.

CHE, J., W.W. ZHOU, J.S. HU, F. YAN, T.J. PAPENFUSS, D.B. WAKE, AND Y.P. ZHANG.

2010. Spiny frogs (*Paini*) illuminate the history of the Himalayan region and

southeast Asia. Proceedings of the National Academy of Sciences 107:13765–

13770.

CHERNOMOR, O., A. VON HAESELER, AND B.Q. MINH. 2016. Terrace aware data structure

for phylogenomic inference from supermatrices. Systematic Biology 65:997-

1008.

CHIAMVIMONVAT, N., M.T. PÉREZ-GARCÍA, G.F. TOMASELLI, AND E. MARBAN. 1996.

Control of ion flux and selectivity by negatively charged residues in the outer

mouth of rat sodium channels. The Journal of Physiology 491:51–59.

CROTHER, B.I., SOCIETY FOR THE STUDY OF AMPHIBIANS AND REPTILES, COMMITTEE ON

STANDARD ENGLISH AND SCIENTIFIC NAMES, AMERICAN SOCIETY OF ICHTHYOLOGISTS

AND HERPETOLOGISTS, CANADIAN ASSOCIATION OF HERPETOLOGISTS, PARTNERS IN

AMPHIBIAN AND REPTILE CONSERVATION, SOCIETY FOR THE STUDY OF AMPHIBIANS

AND REPTILES, AND HERPETOLOGISTS' LEAGUE. 2017. Scientific and Standard

English Names of Amphibians and Reptiles of North America North of Mexico,

with Comments Regarding Confidence in Our Understanding, Eighth Edition.

DALY, J.W., F. GUSOVSKY, C.W. MYERS, M. YOTSU-YAMASHITA, AND T. YASUMOTO. 1994.

First occurrence of tetrodotoxin in a dendrobatid frog (*Colostethus inguinalis*),

with further reports for the bufonid genus *Atelopus*. Toxicon 32:279–285.

DUELLMAN, W.E., A.B. MARION, AND S. B. HEDGES. 2016. Phylogenetics, classification,

and biogeography of the treefrogs (Amphibia: Anura: Arboranae). Zootaxa

4104:1–109.

DURSO, A.M., L.A. NEUMAN-LEE, G.R. HOPKINS, AND E.D. BRODIE, JR. 2021. Stable

isotope analysis suggests that tetrodotoxin-resistant Common Gartersnakes

(*Thamnophis sirtalis*) rarely feed on newts in the wild. Canadian Journal of

Zoology 99:331–338.

FELDMAN, C.R., E.D. BRODIE, JR., E.D. BRODIE III, AND M.E. PFRENDER. 2009.

The evolutionary origins of beneficial alleles during the repeated adaptation of

garter snakes to deadly prey. Proceedings of the National Academy of

Sciences 106: 13415-13420.

FELDMAN, C.R., E.D. BRODIE, JR., E.D. BRODIE III, AND PFRENDER, M.E. 2012.

Constraint shapes convergence in tetrodotoxin-resistant sodium channels of

snakes. Proceedings of the National Academy of Sciences 109: 4556-4561.

FOUQUETTE JR., M. J., AND A. DUBOIS. 2014. A checklist of North American amphibians

and reptiles: the United States and Canada. Xlibris Corporation.

GALL, B.G., A.N. STOKES, S.S. FRENCH, E.A. SCHLEPPHORST, E.D. BRODIE III, AND E.D.

BRODIE, JR. 2011. Tetrodotoxin levels in larval and metamorphosed newts

(*Taricha granulosa*) and palatability to predatory dragonflies. Toxicon 57:978–

983.

GALL, B.G., A.N. STOKES, S.S. FRENCH, E.D. BRODIE III, AND E.D. BRODIE, JR. 2012.

Predatory caddisfly larvae sequester tetrodotoxin from their prey, eggs of the

rough-skinned newt (*Taricha granulosa*). Journal of Chemical Ecology 38: 1351–

1357.

GBIF.org (23 November 2021) GBIF Occurrence Download

https://doi.org/10.15468/dl.qzrq4k.

GBIF.org (23 November 2021) GBIF Occurrence Download

https://doi.org/10.15468/dl.z9tw6r.

GBIF.org (30 November 2021) GBIF Occurrence Download

https://doi.org/10.15468/dl.dwz9yn.

GBIF.org (30 November 2021) GBIF Occurrence Download

https://doi.org/10.15468/dl.ed2uqn.

GEFFENEY, S.L., E. FUJIMOTO, E.D. BRODIE III, E.D. BRODIE, JR., AND P.C. RUBEN. 2005.

Evolutionary diversification of TTX-resistant sodium channels in a predator–prey

interaction. Nature 434: 759–763.

GENDREAU, K.L., A.D. HORNSBY, M.T.J. HAGUE, AND J.W. MCGLOTHLIN. 2021. Gene

conversion facilitates the adaptive evolution of self-resistance in highly toxic

newts. Molecular Biology and Evolution:4077–4094.

Geneious Prime 2020.1.1. https://www.geneious.com.

HAGUE, M.T.J., G. TOLEDO, SHANA L. GEFFENEY, C.T. HANIFIN, E.D. BRODIE, JR., AND E.D.

BRODIE, III. 2018. Large-effect mutations generate trade-off between predatory

and locomotor ability during arms race coevolution with deadly prey. Evolution

Letters 2:406–416.

HANIFIN, C.T. 2010. The chemical and evolutionary ecology of tetrodotoxin (TTX) toxicity

in terrestrial vertebrates. Marine Drugs 8:577–593.

HANIFIN, C.T., E.D. BRODIE JR, AND E.D. BRODIE III. 2008. Phenotypic mismatches reveal

escape from arms-race coevolution. PLOS Biology 6:e60.

https://doi.org/10.1371/journal.pbio.0060060.

HANIFIN, C.T., AND W.F. GILLY. 2015. Evolutionary history of a complex adaptation:

tetrodotoxin resistance in salamanders. Evolution; International Journal of

Organic Evolution 69: 232–244.

HASHIGUCHI, Y., J. M. LEE, M. SHIRAISHI, S. KOMATSU, S. MIKI, Y. SHIMASAKI, N. MOCHIOKA,

T. KUSAKABE, AND Y. OSHIMA. 2015. Characterization and evolutionary analysis of

tributyltin-binding protein and pufferfish saxitoxin and tetrodotoxin-binding protein

genes in toxic and nontoxic pufferfishes. Journal of Evolutionary Biology

28:1103–1118.

HO, B., D.S. YEO, AND J.L. DING. 1994. A tetrodotoxin neutralizing system in the

haemolymph of the horseshoe crab, *Carcinoscorpius rotundicauda*. Toxicon:

Official Journal of the International Society on Toxinology 32:755–762.

HOANG, D.T., O. CHERNOMOR, A. VON HAESELER, B.Q. MINH, AND L.S. VINH. 2018.

UFBoot2: improving the ultrafast bootstrap approximation. Molecular Biology and

Evolution 35:518–522.

HUTCHINSON, D.A., A. MORI, A.H. SAVITZKY, G.M. BURGHARDT, X. WU, J. MEINWALD, AND

F.C. SCHROEDER. 2007. Dietary sequestration of defensive steroids in nuchal

glands of the Asian snake *Rhabdophis tigrinus*. Proceedings of the National

Academy of Sciences 104:2265–2270.

HWANG, P.A., Y.H. TSAI, H.P. LIN, AND D.F. HWANG. 2007. Tetrodotoxin-binding proteins

isolated from five species of toxic gastropods. Food Chemistry 103:1153–1158.

JAMES, D., AND K. HORNIK. 2020. chron: chronological objects which can handle dates

and times. R package version 2.3-56.

KAO, C.Y., AND S.R. LEVINSON. 1986. Preface—tetrodotoxin, saxitoxin, and the

molecular biology of the sodium channel. Annals of the New York Academy of

Sciences 479:ix-x.

KASSAMBARA, A. 2020. ggpubr: 'ggplot2' Based Publication Ready Plots. R package

version 0.4.0. https://CRAN.R-project.org/package=ggpubr.

LEE, C.H., D.K. JONES, C. AHERN, M.F. SARHAN, AND P.C. RUBEN. 2011. Biophysical costs

associated with tetrodotoxin resistance in the sodium channel pore of the garter

snake, *Thamnophis sirtalis*. Journal of Comparative Physiology A 197:33–43.

LLEWELLYN, L.E., P.M. BELL, AND E.G. MOCZYDLOWSKI. 1997. Phylogenetic survey of

soluble saxitoxin-binding activity in pursuit of the function and molecular evolution

of saxiphilin, a relative of transferrin. Proceedings of the Royal Society B:

Biological Sciences 264:891–902.

MAGARLAMOV, T.Y., D.I. MELNIKOVA, AND A.V. CHERNYSHEV. 2017. Tetrodotoxin-

producing bacteria: detection, distribution and migration of the toxin in aquatic

systems. Toxins:166.

MAHAR, J., G.L. LUKÁCS, L. LI, S. HALL, AND E. MOCZYDLOWSKI. 1991. Pharmacological

and biochemical properties of saxiphilin, a soluble saxitoxin-binding protein from

the bullfrog (*Rana catesbeiana*). Toxicon 29:53–71.

MCGLOTHLIN, J.W., M.E. KOBIELA, C.R. FELDMAN, T.A. CASTOE, S.L. GEFFENEY, C.T.

HANIFIN, G. TOLEDO, F.J. VONK, M.K. RICHARDSON, E.D. BRODIE, JR., M.E.

PFRENDER, E.D. BRODIE, III. 2016. Historical contingency in a multigene family

facilitates adaptive evolution of toxin resistance. Current Biology 26:1616–1621.

MEBS, D., M. YOTSU-YAMASHITA, AND O. ARAKAWA. 2016. The praying mantis (*Mantodea*)

as predator of the poisonous red-spotted newt *Notophthalmus viridescens*

(Amphibia: Urodela: Salamandridae). Chemoecology 26:121–126.

MECHAM, J.S. 1967. *Notophthalmus viridescens.* Catalogue of American Amphibians

and Reptiles: 53.1-53.4.

MILLER, S.A., D.D. DYKES, AND H.F. POLESKY. 1988. A simple salting out procedure for

extracting DNA from human nucleated cells. Nucleic Acids Research 16:1215.

MONIZ, H.A., M.A. RICHARD, C.M. GIENGER, AND C.R. FELDMAN. 2021. Every breath you

take: assessing metabolic costs of toxin resistance in garter snakes

(*Thamnophis*). Integrative Zoology. https://doi.org/10.1111/1749-4877.12574.

MYERS, C.W., J.W. DALY, AND B. MALKIN. 1978. A dangerously toxic new frog

(*Phyllobates*) used by Emberá Indians of western Colombia, with discussion of

blowgun fabrication and dart poisoning. Bulletin of the AMNH 161:article 2.

NAGASHIMA, Y., K. YAMAMOTO, K. SHIMAKURA, AND K. SHIOMI. 2002. A tetrodotoxin-binding

protein in the hemolymph of shore crab *Hemigrapsus sanguineus*: purification

and properties. Toxicon: Official Journal of the International Society on

Toxinology 40:753–760.

NEUWIRTH, E. 2014. RColorBrewer: ColorBrewer palettes. R package version1.1-2.

https://CRAN.R-project.org/package=RColorBrewer.

N GUYEN, L., H.A. S CHMIDT, A. VON H AESELER, AND B.Q. M INH. 2015. IQ-TREE: A fast and

effective stochastic algorithm for estimating maximum likelihood phylogenies.

Molecular Biology and Evolution 32:268-274.

NOAA (National Oceanic and Atmospheric Association). 2022. Climate data online

search—data tables for Berkeley, CA and Martinez CA.

https://www.ncdc.noaa.gov/cdo-web/search.

O TA, W.M., B. O LSEN, G.M. B UCCIARELLI, AND L.B. K ATS. 2018. The effect of newt toxin

on an invasive snail. Hydrobiologia 817:341–348.

P ARADIS E., AND K. S CHLIEP. 2019. ape 5.0: an environment for modern phylogenetics

and evolutionary analyses in R. Bioinformatics 35:526-528.

P IRES, O.R., A. S EBBEN, E.F. S CHWARTZ, S.W.R. L ARGURA, C. B LOCH, R.A.V. M ORALES,

AND C.A. S CHWARTZ. 2002. Occurrence of tetrodotoxin and its analogues in the

Brazilian frog *Brachycephalus ephippium* (Anura: Brachycephalidae). Toxicon:

Official Journal of the International Society on Toxinology 40:761–766.

R AMÍREZ-C ASTAÑEDA, V. 2017. Origin of mutations in the NaV gene family associated

with resistance to neurotoxins in snake (*Erythrolamprus sp.*) predators of dart

poison frogs (Dendrobatidae). Master's thesis. University of the Andes, Bogota,

Colombia.

R Core Team. 2021. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

https://www.R-project.org.

R ECUERO, E., Í. M ARTÍNEZ-S OLANO, G. P ARRA-O LEA, AND M. G ARCÍA-P ARÍS. 2006.

Phylogeography of *Pseudacris regilla* (Anura: Hylidae) in western North America,

with a proposal for a new taxonomic rearrangement. Molecular Phylogenetics

and Evolution 39:293–304.

RECUERO, E., Í. MARTÍNEZ-SOLANO, G. PARRA-OLEA, AND M. GARCÍA-PARÍS. 2006.

Corrigendum to 'Phylogeography of *Pseudacris regilla* (Anura: Hylidae) in

western North America, with a proposal for a new taxonomic rearrangement'

[Mol. Phylogenet. Evol. 39 (2006) 293–304]. Molecular Phylogenetics and

Evolution 41:511.

REIMCHE, J.S., E.D. BRODIE JR., A.N. STOKES, E.J. ELY, H.A. MONIZ, V.L. THILL, J.M.

HALLAS, M.E. PFRENDER, E.D. BRODIE III, AND C.R. FELDMAN. 2020. The

geographic mosaic in parallel: matching patterns of newt tetrodotoxin levels and

snake resistance in multiple predator–prey pairs. Journal of Animal Ecology

89:1645–1657.

REVELL, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and

other things). Methods in Ecology and Evolution 3:217-223.

RITSON-WILLIAMS, R., M. YOTSU-YAMASHITA, AND V.J. PAUL. 2006. Ecological functions of

tetrodotoxin in a deadly polyclad flatworm. Proceedings of the National Academy

of Sciences 103:3176–3179.

SIEVERS F., A. WILM, D.G. DINEEN, T.J. GIBSON, K. KARPLUS, W. LI, R. LOPEZ, H.

MCWILLIAM, M. REMMERT, J. SÖDING, J.D. THOMPSON, D.G. HIGGINS. 2011. Fast,

scalable generation of high-quality protein multiple sequence alignments using

Clustal Omega. Molecular Systems Biology 7:539.

TANU, M.B., Y. MAHMUD, K. TSURUDA, O. ARAKAWA, AND T. NOGUCHI. 2001. Occurrence of

tetrodotoxin in the skin of a rhacophoridid frog *Polypedates* sp. from Bangladesh.

Toxicon: Official Journal of the International Society on Toxinology 39:937–941.

TARVIN, R.D., J.C. SANTOS, L.A. O'CONNELL, H.H. ZAKON, AND D.C. CANNATELLA. 2016.

Convergent substitutions in a sodium channel suggest multiple origins of toxin

resistance in poison frogs. Molecular Biology and Evolution 37:607.

TIKHONOV, D.B., AND B.S. ZHOROV. 2005. Modeling p-loops domain of sodium channel:

homology with potassium channels and interaction with ligands. Biophysical

Journal 88:184–197.

TIKHONOV, D.B., AND B.S. ZHOROV. 2011. Possible roles of exceptionally conserved

residues around the selectivity filters of sodium and calcium channels. Journal of

Biological Chemistry 286:2998–3006.

TRIFINOPOULOS, J., L.T. NGUYEN, A. VON HAESELER, AND B.Q. MINH. 2016. W-IQ-TREE: a

fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids

Research 44:W232-W235.

VAELLI, P.M., K.R. THEIS, J.E. WILLIAMS, L.A. O'CONNELL, J.A. FOSTER, AND H.L. EISTHEN.

2020. The skin microbiome facilitates adaptive tetrodotoxin production in

poisonous newts. ELife 9:e53898. https://doi.org/10.7554/eLife.53898.

VENKATESH, B., S.Q. LU, N. DANDONA, S.L. SEE, S. BRENNER, AND T.W. SOONG. 2005.

Genetic basis of tetrodotoxin resistance in pufferfishes. Current Biology 15:2069–

2072.

VITT, L. J., AND J. P. CALDWELL. 2013. Herpetology: An Introductory Biology of

Amphibians and Reptiles. United Kingdom and United States of America.

WANNINGER, M., T. SCHWAHA, AND E. HEISS. 2018. Form and function of the skin glands

in the Himalayan newt *Tylototriton verrucosus*. Zoological Letters 4:15.

WICKHAM, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

York. United States of America.

WICKHAM, H., M. AVERICK, J. BRYAN, W. CHANG, L.D. MCGOWAN, R. FRANÇOIS, G.

GROLEMUND, A. HAYES, L. HENRY, J. HESTER, M. KUHN, T.L. PEDERSEN, E. MILLER,

S.M. BACHE, K.MÜLLER, J. OOMS, D. ROBINSON, D.P. SEIDEL, V. SPINU, K.

TAKAHASHI, D. VAUGHAN, C. WILKE, K. WOO, AND H. YUTANI. 2019. Welcome to the

tidyverse. Journal of Open Source Software 4:1686.

https://doi.org/10.21105/joss.01686.

WICKHAM, H., R. FRANÇOIS, L. HENRY, AND K. MÜLLER. 2021. dplyr: a grammar of data

manipulation. R package version 1.0.7.

https://CRAN.R-project.org/package=dplyr.

WILKE, C.O. 2020. cowplot: streamlined plot theme and plot annotations for 'ggplot2'.

R package version 1.1.1. https://CRAN.R-project.org/package=cowplot.

WILLIAMS, B.L. 2010. Behavioral and chemical ecology of marine organisms with respect

to tetrodotoxin. Marine Drugs 8:381–398.

WILSON, N.J., A.N. STOKES, G.R. HOPKINS, E.D. BRODIE, JR., AND C.R. WILLIAMS. 2014.

Functional and physiological resistance of crayfish to amphibian toxins:

tetrodotoxin resistance in the white river crayfish (*Procambarus acutus*).

Canadian Journal of Zoology 92:939–945.

ZANGER, U.M., AND M. SCHWAB. 2013. Cytochrome P450 enzymes in drug metabolism:

regulation of gene expression, enzyme activities, and impact of genetic variation.

Pharmacology & Therapeutics 138:103-41.

https://doi.org/10.1016/j.pharmthera.2012.12.007

ZIMMER, R.K., D.W. SCHAR, R.P. FERRER, P.J. KRUG, L.B. KATS, AND W.C. MICHEL. 2006.

The scent of danger: tetrodotoxin (TTX) as an olfactory cue of predation risk.

Ecological Monographs 76:585–600.

A *Pseudacris regilla* adults

B *Taricha* spp., including larvae

C *Taricha torosa* egg masses

Pond — OBRT1 — OBRT2 — UCBG

DIV S5-S6 p-loop

1519

FGNSIICLFEITTSAGWDGLLNP

Rattus norvegicus
Erythrolamprus epinephelus •••••••••••••••••••SD••••
Rhabdophis tigrinus ••••M•••••••••••••••••••
*Thamnophis elegans** ••••••••••••••••••••••••
*Thamnophis sirtalis** •••••L••••V••••NV•••••
*Taricha torosa** ••S••••••QS••••••SD••I•
*Taricha granulosa** ••S•I••••QS••••••SD••I•
Nanorana parkeri ••••M••••Q••••••••••
Acris crepitans ••••••••••••••••••••••••
Pseudacris cadaverina ••••••••••••••••••••••••
Pseudacris regilla ••••M••••••••••••••••••

0.2 substitutions/site