Early transition in foraging mode promoted greater reproductive effort in

widely foraging lizards

Dylan J. Padilla Perez[1,*]

dpadil10@asu.edu

Dale F. DeNardo[1]

denardo@asu.edu

Michael J. Angilletta Jr.[1]

ma@asu.edu

[1]School of Life Sciences, Arizona State University, Tempe, AZ 85287, USA.

*Corresponding author: dpadil10@asu.edu, Phone: +1 (480) 646-7769

Statement of authorship: DJPP, DD and MJA conceived the study. DJPP performed data
analyses. All authors drafted, revised, and approved the final version of the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Data accessibility statement: Supporting data, including a rmarkdown file with code and good
quality figures from the statistical analyses, will be archived in an online repository.
21
22 **Article type:** Letter
23
Keywords: fecundity, clutch size, maternal size, phenotypic plasticity, predation, offspring size,
reproduction.

This manuscript includes:

Abstract: 196 words
Main Text: 3650 words
Figures: 4
Tables: 1
References: 60

32 Abstract

Life-history theory predicts that the optimal reproductive effort of an organism is affected by

factors such as energy acquisition and predation risk. Although this is a major focus of study in

evolutionary ecology, the empirical evidence consists of conflicting data on a few organisms. For

instance, theoretical models within the foraging mode paradigm suggest that widely foraging

females have evolved low reproductive effort, because a heavy reproductive load decreases their

ability to escape from predators. By contrast, a long-standing prediction of evolutionary theory

indicates that organisms subject to high mortality due to predation, as suggested for widely

foraging species, should increase their reproductive investment. Here, we revise the available

literature on the relationship between foraging mode and reproductive effort of lizards. In doing

so, we present evidence that widely foraging species have evolved greater reproductive effort

than sit-and wait species. This is the largest comparative analysis of foraging mode and

reproductive effort to date: 485 species grouped in 32 families. Based on our findings, we

propose a theoretical model derived from the optimal foraging theory that potentially explains the

observed patterns in lizards, paving the way for ecologist to test mechanistic hypotheses at the

intraspecific level.

Main Text

Introduction

The foraging behaviors of vertebrates lie along a continuum, ranging from the energetically

demanding strategy of foraging actively to the energetically conservative strategy of sit-and-wait

foraging (Pianka 1966; Perry 1999). Over the past 50 years, ecologists have developed a set of

hypotheses about how an organism's foraging mode relates to its life history (Vitt & Congdont

1978; Vitt & Price 1982; Dunham *et al.* 1988; Webb *et al.* 2003). These relationships stem from

two major assumptions (Fig. 1). First, widely foraging animals spend more energy to forage than

a sit-and-wait foragers, but might also consume enough food to have more surplus energy

(Anderson & Karasov 1981, 1988; Huey & Pianka 1981; Nagy *et al.* 1984; Stuginski *et al.* 2018;

Bury 2021). However, sit-and-wait foragers might exhibit wider diet breadths, because they

encounter prey less frequently compared to active foragers (Glaudas *et al.* 2019). Second, a

widely foraging animal could suffer a greater risk of mortality because its movements are

conspicuous to predators (Cooper & Perez-Mellado 2004). Both a greater energy supply and a

greater mortality risk would select for genotypes that allocate more energy to reproduction,

manifested as more or larger offspring (Bonnet *et al.* 2001; Silva *et al.* 2020). The evolution of

greater reproductive effort could feed back on the energy gain and mortality risk if a female

carrying a greater mass of offspring tends to chase prey (Shine 1980). Similarly, the load

associated with food consumption may compromise the locomotion of individuals, increasing the

vulnerability to predators (Werner & Anholt 1993; Cooper 2000). In this model, a suite of traits

associated with foraging mode would coadapt to the spatiotemporal distributions of prey and

predators.

Despite the wealth of conjecture, these hypothetical relationships among foraging mode

and other traits have been assessed in only a handful of cases, as described above. Ideally, one

would isolate each relationship and conduct experiments to quantify the evolution of traits in

controlled environments (Huey & Bennet 1986). However, such data are difficult to gather for

many species and will probably remain rare. For instance, limited circumstantial evidence exists

for the putative relationship between foraging mode and predation risk. Data on stomach contents

of field-collected vipers suggested that widely foraging lacertids are more vulnerable to predators

than sedentary species (Huey & Pianka 1981), but a number of confounding factors can explain

this observation as well as foraging mode can. Alternatively, researchers have evaluated the

relationship between mortality and life-history traits with no emphasis on foraging behavior. For

example, experimental evolution with guppies and fruit flies revealed that genotypes that evolved

in risky environments developed more rapidly, matured at a smaller size, and reached their peak

of fecundity faster than did genotypes that evolved in safe environments (Stearns 2000; Reznick

et al. 2001). From these results, we expect that a high rate of predation selects for greater

reproductive effort, but given the scarce evidence, it is still uncertain whether foraging mode

generally affects predation risk or net energy gain.

Although experimental data are lacking, comparative methods have been used to explore

how foraging mode and life-history traits have evolved. These interspecific analyses have

focused mostly on the reproductive effort of lizards. The earliest analysis of 22 species revealed

an unexpected pattern: sit-and-wait species had a greater reproductive effort than widely foraging

species (Vitt & Congdont 1978). The authors suggested that widely foraging species might be

forced to carry fewer or smaller offspring, because moving long distances with a voluminous

clutch decreases the chance of escaping from a predator. A subsequent analysis of data for 50

species of lizards supports this result by testing a model in which predation risk increased with

increasing reproductive effort (Vitt & Price 1982). Consecutively, Roff (2002) extended support for

this model in a comparative analysis of 130 lizard species. However, none of these early

analyses controlled for potential phylogenetic correlations that might generate spurious

relationships between foraging mode and reproductive effort (Felsenstein 1985), especially

because foraging could vary greatly among but little within families. A more recent analysis, using

phylogenetic comparative methods, failed to detect a significant relationship between foraging

mode and reproductive effort (Mesquita *et al.* 2016). Thus, despite several attempts to establish a

relationship between foraging mode and the life history, we still lack support for a long-standing

prediction of the foraging-mode paradigm.

Here, we present the first evidence that widely foraging species have evolved greater

reproductive effort than sit-and wait species. This evidence comes from the largest comparative

analysis of foraging mode and reproductive effort to date: 485 species of lizards representing 32

families. In this analysis, we inferred the evolutionary history of foraging modes, complementing

on past reconstructions of ancestral states (Miles *et al.* 2007). In contrast to previous analyses,

our study supports the prediction of theoretical models of the optimal reproductive effort, paving

the way for ecologists to test mechanistic hypotheses at the intraspecific level.

Materials and Methods

Data source and description of variables

We used published estimates of life history and foraging behavior for 485 species of lizards

grouped in 32 families, excluding amphisbaenians and snakes. These data represent a subset of

a data assembled from primary and secondary literature on lizards' traits published during the last

12 years (Meiri 2018). The reproductive effort of the species was defined as the product of the

mean snout-vent length of hatchlings or neonates (mm) and the mean clutch or litter size. Given

the limited data available for clutch frequency in widely foraging and sit-and-wait species, we

could not examine the lifetime reproductive effort of lizards. Instead, our measure represents the

investment of females in a single reproductive event. Our measure of maternal size consisted of

the mean snout-vent length of adult females. We used the length of hatchlings rather than the

mass of eggs, because the latter might reflect high water content rather than energy content in

lizards that lay poorly calcified eggs (Deeming 2004; Meiri *et al.* 2015). Moreover, hatchling size

can be measured for viviparous species as well as oviparous species.

Although relative clutch mass is commonly reported as an index of reproductive effort, we

used a different index to avoid statistical issues associated with analyzing a ratio. Ratios of

random numbers regressed against their denominator will automatically yield negative

correlations (Atchley *et al.* 1876; Packard & Boardman 1988). For example, a comparative study

of 551 species of lizards revealed a negative effect of maternal size on relative clutch mass (Meiri

et al. 2012). This result is difficult to interpret as it contrasts with most theoretical models, which

predict that relative clutch mass should increase with maternal size (Roff 2002)—a pattern

observed in many organisms (e.g., Primack 1979; Barneche et al. 2018; Marshall et al. 2021).

We defined the foraging mode of a species based on whether it has been reported as an

ambush predator ("sit-and-wait"), an active forager ("widely foraging"), or uses a mixed strategy

("mixed"). Although this categorization seems somewhat artificial and simplistic, numerous

species of lizards clearly belong to one of these categories (Huey & Pianka 1981; Perry 1999).

We focused our analyses only on carnivores, because herbivores do not fit into the classical

paradigm of foraging modes (Pianka & Vitt 2003).

Ancestral character and phylogenetic signal estimates

To determine the appropriate model of evolution, we used a set of continuous-time discrete-state

Markov chain models to sample the character histories from their posterior probability distribution

(Huelsenbeck *et al.* 2003), and a time-calibrated phylogeny of squamate reptiles (Zheng & Wiens

2016). We rooted the tree with the Tuatara (*Sphenodon punctatus*) as the outgroup for our study

taxa. Based on previous analyses, we coded this outgroup as a sit-and-wait forager (Pianka & Vitt

2003; Vitt *et al.* 2003). We fitted three different models to our data, using the function

make.simmap from the R package "phytools" version 0.7.80 (Revell 2012; R Core Team 2021).

These models were as follows: 1) an equal rates model (ER), in which the rate of change

between all three states of the character are assumed to be equivalent; 2) an all-rates-different

model (ARD), which allows the transitions among states to have different rates; 3) and a

symmetrical model (SYM), which allows different rates of change between pairs of states but

changes between all states are theoretically possible. For each model, we estimated the

stationary distribution by numerical solving for *pi* (*pi* = 'estimated'), and this was used as prior

distribution on the root node of the tree. These models sample the character histories conditioned

on the transition matrix (Q matrix) and use the phylogenetic tree with annotated tips to create

stochastic simulation maps for the potential evolutionary transitions between foraging modes. To

describe the variation in reproductive effort across the study taxa, we plotted bars adjacent to the

tips of the phylogeny representing values of species reproductive effort. We selected the most

likely model of evolution based on the Akaike Information Criterion (AIC). Lastly, we generated

10000 trees of the most-likely model and used the *summary* function to count the total number of

transitions that occurred between character states (see supporting material for detailed

information).

We computed our measure of phylogenetic signal (or character dispersion on a

phylogeny) by using the Fritz and Purvis' $D\ test$, available through the function *phylo.d* in the R

package "caper" version 1.0.1 (Fritz & Purvis 2010; Orme *et al.* 2018; R Core Team 2021). This

parameter is calculated as follows:

$$D = \frac{[\sum d_{obs} - mean(\sum d_b)]}{[mean(\sum d_r) - mean(\sum d_b)]};$$

where d_{obs} equals the number of character-state changes required to get the observed

distribution of character states at the tips of the phylogeny. The d_{obs} is then scaled using two null

distributions. The first distribution, d_r, comprises d values obtained from permutations where the

number of species with each character state is kept constant, but the values are shuffled on the

tips of the phylogeny. Thus, d_r is the expected distribution of d values if character states are

randomly distributed among species without respect to phylogeny. The second distribution, d_b,

comprises the d values expected if character states are distributed among species under the

expectations of the Brownian motion model of evolution. We generated d_b by simulating a

continuous trait along the phylogeny, then defining the character state at each tip according to

some threshold value of the continuous trait (Fritz & Purvis 2010). The value of D equals 1 if the

distribution of the binary trait is random with respect to phylogeny, and exceeds 1 if the

distribution of the trait is more overdispersed than the random expectation. The value of D equals

0 if the binary trait is distributed as expected under the Brownian motion model of evolution, and

is less than 0 if the binary trait is more phylogenetically conserved than the Brownian expectation.

The distributions d_r and d_b were used to assign $p-values$ to d_{obs}. Accordingly, if d_{obs} is larger

than 95% of d_r values, the distribution of the trait would be significantly more overdispersed than

the random expectation, if d_{obs} is less than 95% of d_b values, the character would be significantly

more clumped than the Brownian expectation.

Effects of body length and foraging mode on reproductive effort

We fitted Phylogenetic Generalized Least Squares models (PGLS) to test our prediction on the

relationship among body length, foraging mode, and the reproductive effort of lizards. In

comparative biology, normal regression models cannot be used because the assumption of

independence of residuals is violated (Felsenstein 1985; Harvey & Pagel 1991). This problem can

be corrected using phylogenetic comparative methods. We fitted the models using the function

gls from the R package "nlme" version 3.1-152 (Pinheiro *et al.* 2021; R Core Team 2021). We

adopted the reproductive effort of lizards as our response variable. Similarly, maternal length and

foraging mode corresponded to our predictor variables. We fitted the models assuming two ways

in which the tree structure was expected to affect the covariance in trait values across taxa

(corBrownian and corPagel —value = 0— error structures, using the R package "ape"). We

selected the best-fit model based on the AIC values (see supporting material for detailed

information). To avoid confounding results due to a potential large variation in maternal length

among species, we not only tested our prediction for all lizards in our data set (Fig. 3A), but also

examined the data at lower taxonomic scales (e.g., within the superfamily Scincoidea; Fig. 3B).

Analyzing the data at different taxonomic scales enabled us to better investigate the level of

support for the prediction tested.

Results

The evolutionary transitions in foraging modes among lizards were best described by a model in

which rates of evolution differ among modes, referred to as the all-rates-different model. Despite

a slightly difference with the widely foraging state, our analysis revealed that sit-and-wait foraging

is the most likely ancestral state of all lacertilians (Fig. 2). Sit-and-wait foraging has evolved in two

major clades of lizards; near the root of the tree, we found strong evidence suggesting that the

ancestor of Gekkota was a sit-and-wait predator. Likewise, the ancestor of iguanians was a sit-

and-wait predator. Iguanians included in our analysis are part of both the Acrodonta and

Pleurodonta clades. During the early Jurassic (≈ 200 mya), a major transition from sit-and-wait to

widely foraging occurred in the ancestor of Anguimorpha, Lacertoidea, and Scincoidea. Since

then, a reverse transition from widely foraging to sit-and-wait foraging occurred within Scincoidea,

in the ancestor of spiny-tailed lizards and night lizards (Cordylidae and Xantusiidae). The

estimate of the $D\ test$ for phylogenetic signal indicated that foraging mode is phylogenetically

conserved as expected under a Brownian motion model of evolution ($D = -0.14,\ p[D < 1] = 0,$

$p[D > 0] = 0.80$).

Our analyses show that widely foraging species have evolved greater reproductive effort

than sit-and-wait species. However, the mechanism underlying this pattern is mediated by a

complex interaction between foraging mode and maternal length (Table 1). In Lacertilia, widely

foraging species have greater reproductive effort for two reasons: 1) they are larger in size, and

2) on average, they give birth to a higher number of offspring for a given size (Fig. 3A). In the

superfamily Scincoidea, widely foraging species evolved greater reproductive effort despite

having a smaller mean body length (Fig. 3B). Interestingly, skinks that adopt a mixed foraging

behavior have also evolved greater reproductive effort than sit-and-wait skinks.

Discussion

Our results show that widely foraging species have evolved a greater reproductive effort than sit-

and-wait species. Presumably, the ability of widely foraging lizards to harvest and assimilate more

resources might explain this pattern. Studies on field metabolic rate of free-ranging lizards

suggest that an average widely foraging species spends 32% more daily energy than a sit-and-

wait lizard, but this extra energy expenditure is probably paid off with greater daily food

consumption (Brown & Nagy 2007). Food consumption may play an important role in determining

the reproductive effort of lizards in three ways: 1) it promotes follicular growth during the

reproductive season; 2) in the long term, it increases energy stores to initiate reproduction (e.g.,

vitellogenesis). and 3) it may also reduce age at first reproduction. Consistent with these ideas,

Bonnet and colleagues found that female vipers that had good body condition early in

vitellogenesis produced large litters (Bonnet *et al.* 2001). Similarly, vipers that gained more mass

during follicular growth produced larger offspring. Early reproduction gives offspring sufficient time

to mature in the same year that they hatched, which enables them to participate as adults in the

subsequent breeding season (Hahn & Tinkle 1965). Therefore, the foraging behavior of a widely

foraging female can increase the amount of gametic resources gathered before and during each

reproductive bout, potentially resulting in more energy to invest in reproduction.

Additionally, maternal size also plays an important role in determining the reproductive

effort of lizards. Among many ectotherms and plants, a common pattern is that reproductive effort

increases with size (Tinkle 1967; Primack 1979; Bownds *et al.* 2010; Barneche *et al.* 2018;

Marshall *et al.* 2018, 2021). This observation indicates that increased size might evolve by natural

selection, because it increases the fitness of individuals. Interestingly, our results paint a more

complex picture of the effect of body size on reproductive effort. We found that the relationship

between maternal length and reproductive effort depends on foraging mode. The most complex

form of relationship occurs when small widely-foraging females have a greater reproductive effort

than large sit-and-wait females (Fig. 3B). A model of the evolution of optimal reproductive effort

predicts this pattern (Parker & Begon 1986). This model suggests that the total energy

accumulated for reproduction (m) depends on the time spent foraging (t) and maternal size.

Assuming that females have the same foraging efficiency, smaller females would reach their

maximum capacity to accumulate resources at lower values of m, producing fewer or smaller

offspring than larger females (Fig. 4A). However, if a widely foraging female has a smaller body

size but a higher foraging efficiency, she may have greater reproductive investment than a large

sit-and-wait female (Fig. 4B). The same outcome should be observed if a widely foraging female

is both a larger and more efficient forager than a sit-and-wait female (Fig. 4C). Furthermore, if

widely foraging females are more efficient at foraging, they may have both a shorter foraging time

and a reduced interclutch interval. Therefore, widely foraging females might not only produce

larger or more offspring in a single clutch, but they might also increase the number of clutches

throughout their lifetime.

Because the risk of predation presumably depends on the time spent foraging, individuals

that require less time to accumulate the optimal amount of resources should incur a lower

probability of death. Based on the theory of foraging by Schoener, the energy accumulated from

foraging increases monotonically toward an asymptote (Schoener 1971). In such cases, the law

of diminishing returns implies that foraging for twice as long would not result in twice the energetic

return (Fig. 4B). Therefore, greater foraging efficiency (more energy gain per foraging time)

should enable an organism to survive better in environments with high predation risk. Consistent

with this idea, a study on foraging efficiency (defined as ratio of metabolizable energy gained

while foraging to the energy spent while foraging) partially indicates that widely foraging lizards

spent less time foraging but grew larger than did sit-and-wait lizards (Nagy *et al.* 1984). Indeed,

our results show that widely foraging lizards are, on average, larger than sit-and-wait lizards (Fig.

3A). The ability of widely foraging species to rapidly outgrow a sit-and-wait species may also

enable them to outgrow the gape limitations of predators (Lynch 1980; Reznick & Endler 1982).

Furthermore, when competition for mates is crucial, the winner of a mating-contest between two

competitors is generally the larger individual (Roff 2002). Therefore, if body size is critical to

outperform predators and outcompete conspecifics, the optimal pattern may be to grow to the

maximal size that leads to the greatest reproductive effort.

The foraging mode paradigm is mostly focused on dichotomous variation, yet plastic

variation in foraging mode may drive the evolution of the reproductive effort in some species.

Recent studies have revealed that the foraging mode of an organism depends on the ecological

context, such as presence of predators, abundance of prey, or habitat degradation (Greef &

Whiting 2000; Hawlena & Perez-Mellado 2009; Wasiolka *et al.* 2009; Donihue 2016). Such

plasticity could precede rapid evolutionary change and local adaptation of the life history

(Richardson *et al.* 2014). Our results show a strong presence of species that adopt a mixed

foraging strategy in the superfamilies Anguimorpha, Gekkota, Iguania, and Scincoidea (Fig. 2).

Interestingly, these lizards have greater reproductive effort than do sit-and-wait lizards,

specifically for species of relative small size (Fig. 3B). Because mixed foraging species are often

exposed to a wide range of environments with different selective pressures, these species might

actively select habitats that maximize their reproductive effort, indirectly resulting in local

adaptation (Richardson *et al.* 2014). Our analysis, which provides the first evidence that a mixed

foraging strategy can lead to greater reproductive effort, should encourage others to address the

questions raised by this discovery.

Our study presents the first evidence that the early shift in foraging mode—from sit-and-

wait to widely foraging—in the evolutionary history of lacertilians was accompanied by the

evolution of a greater reproductive effort in most widely foraging species (Fig. 3). Based on the

theory of optimal foraging, we propose a theoretical model that potentially helps to better

understand the evolution of the optimal life-history strategy. This model is emphasized on a

complex covariation between energy gain, maternal size, and foraging mode. Investigating how

much energy an individual obtains from the environment is crucial in life-history analysis because

the flux of energy within an organism not only determines its survival, but also whether it can

engage in reproduction, a very evolutionarily important activity. Currently, analyses of the

energetics of foraging modes stem from measures of a few species in short periods of a season.

Such data overlook the physiological and ecological constraints of seasonality on some

organisms. For example, widely foraging lizards in the Kalahari Desert consume more food during

the summer, but stop eating during winter as they hibernate. By contrast, sit-and-wait lizards

probably forage during both seasons as they remain active during winter (Huey *et al.* 2021).

Evidence of this nature is still rare in the current literature, revealing the need for long-term

studies on the energetics of foraging modes. Similarly, direct comparisons of mortality rates in

widely foraging versus sit-and-wait species are required, as the existing data prevent us from

making convincing conclusions on the relationship between foraging mode and vulnerability to

predation. Finally, our study captured the effects of foraging plasticity on the evolution of the

reproductive effort of lizards. Overall, lizards that adopt a mixed foraging strategy also have

greater reproductive effort than do sit-and-wait species. This result suggests that foraging

plasticity can result in differential reproductive effort and opens the question for future studies of

whether the foraging efficiency and predation risk of each mode influence the evolution of

fecundity in species that forage plastically.

Acknowledgments

We thank Mark O'Shea for providing us with lizard photos for data visualization. We also thank

Gustavo Burin for his valuable comments on an early draft of our manuscript.

References

Anderson, R. & Karasov, W. (1981). Contrasts in Energy Intake and Expenditure in Sit-and-Wait
and Widely Foraging Lizards. *Oecologia*, 49, 67–72.

Anderson, R. & Karasov, W. (1988). Energetics of the Lizard *Cnemidophorus Tigris* and Life
History Consequences of Food- Acquisition Mode. *Ecological Monographs*, 58, 79–110.

Atchley, W.R., Gaskins, C.T. & Anderson, D. (1876). Statistical Properties of Ratios . I . Empirical
Results. *Systematic zoology*, 25, 137–148.

Barneche, D.R., Robertson, † D Ross, White, C.R. & Marshall, D.J. (2018). Fish reproductive-
energy output increases disproportionately with body size. *Science*, 360, 642–645.

Bonnet, X., Naulleau, G., Shine, R. & Lourdais, O. (2001). Short-term versus long-term effects of
food intake on reproductive output in a viviparous snake, *Vipera aspis*. *Oikos*, 92, 297–308.

Bownds, C., Wilson, R. & Marshall, D.J. (2010). Why do colder mothers produce larger eggs? An
optimality approach. *Journal of Experimental Biology*, 213, 3796–3801.

Brown, T.K. & Nagy, K.A. (2007). Lizard Ecology. In: *Lizard Ecology* (eds. Reilly, S., McBrayer, L.
& Miles, D.B.). Cambridge University Press, New York, pp. 120–140.

Bury, S. (2021). Energy expenses on prey processing are comparable, but paid at a higher
metabolic scope and for a longer time in ambush vs active predators: a multispecies study
on snakes. *Oecologia*, 197, 61–70.

Cooper, W.E.Jr. (2000). Tradeoffs between Predation Risk and Feeding in a Lizard, the Broad-
Headed Skink (*Eumeces laticeps*). *Behaviour*, 137, 1175–1189.

Cooper, W.E.Jr. & Perez-Mellado, V. (2004). Tradeoffs between Escape Behavior and Foraging
Opportunity by the Balearic Lizard (*Podarcis lilfordi*), 60, 321–324.

Deeming, D. (2004). *Reptilian Incubation: Environment, Evolution and Behaviour*. Nottingham
University Press, Nottingham.

Donihue, C.M. (2016). Aegean wall lizards switch foraging modes, diet, and morphology in a
human-built environment. *Ecology and Evolution*, 6, 7433–7442.

Dunham, A., Miles, D. & Reznick, D. (1988). Life history patterns in squamate reptiles. In: *Biology
of the Reptilia* (eds. Gans, C. & Huey, R.B.). A.R. Liss., New York, pp. 441–511.

Felsenstein, J. (1985). Phylogenies and the Comparative Method. *The American naturalist*, 125,
1–15.

Fritz, S.A. & Purvis, A. (2010). Selectivity in mammalian extinction risk and threat types: A new
measure of phylogenetic signal strength in binary traits. *Conservation Biology*, 24, 1042–
1051.

Glaudas, X., Glennon, K.L., Martins, M., Luiselli, L., Fearn, S., Trembath, D.F., *et al.* (2019).
Foraging mode, relative prey size and diet breadth: A phylogenetically explicit analysis of
snake feeding ecology. *Journal of Animal Ecology*, 88, 757–767.

Greef, J.M. & Whiting, M.J. (2000). Foraging-Mode Plasticity in the Lizard *Platysaurus broadleyi*.
Herpetologica, 56, 402–407.

Hahn, W.E. & Tinkle, D.W. (1965). Fat body cycling and experimental evidence for its adaptive
significance to ovarian follicle development in the lizard *Uta stansburiana*. *Journal of
Experimental Zoology*, 158, 79–85.

Harvey, P. & Pagel, M. (1991). *The Comparative Method in Evolutionary Biology*. Oxford
University Press, United States.

Hawlena, D. & Perez-Mellado, V. (2009). Change Your Diet or Die: Predator-Induced Shifts in
Insectivorous Lizard Feeding Ecology. *Oecologia*, 161, 411–419.

Huelsenbeck, J.P., Nielsen, R. & Bollback, J.P. (2003). Stochastic mapping of morphological
characters. *Systematic Biology*, 52, 131–158.

Huey, R.B. & Pianka, E.R. (1981). Ecological Consequences of Foraging Mode. *Ecology*, 62,
991–999.

Huey, R.B. & Bennet, A.F. (1986). A comparative approach to field and laboratory studies in
evolutionary ecology. In: *Predator-prey relationships in lower vertebrates* (eds. Feder, M.E.
& Lauder, G. v.). Uiversity of Chicago, Chicago, IL, pp. 82–89.

Huey, R.B., Miles, D.B. & Pianka, E.R. (2021). Seasonality in kgalagadi lizards: Inferences from
legacy data. *American Naturalist*, 198, 759–771.

Lynch, M. (1980). The Evolution of Cladoceran Life Histories. *The Quarterly Review of Biology*,
55, 23–42.

Marshall, D.J., Barneche, D.R. & White, C.R. (2021). How does spawning frequency scale with
body size in marine fishes? *Fish and Fisheries*, 1–8.

Marshall, D.J., Pettersen, A.K. & Cameron, H. (2018). A global synthesis of offspring size
variation, its eco-evolutionary causes and consequences. *Functional Ecology*, 32, 1436–
1446.

Meiri, S. (2018). Traits of lizards of the world: Variation around a successful evolutionary design.
Global Ecology and Biogeography, 27, 1168–1172.

Meiri, S., Brown, J.H. & Sibly, R.M. (2012). The ecology of lizard reproductive output. *Global
Ecology and Biogeography*, 21, 592–602.

Meiri, S., Feldman, A. & Kratochvíl, L. (2015). Squamate hatchling size and the evolutionary
causes of negative offspring size allometry. *Journal of Evolutionary Biology*, 28, 438–446.

Mesquita, D.O., Costa, G.C., Colli, G.R., Costa, T.B., Shepard, D.B., Vitt, L.J., *et al.* (2016). Life-
history patterns of lizards of the world. *The American Naturalist*, 187, 689–705.

Miles, D.B., Losos, J.B. & Irschick, D. (2007). Morphology, performance, and foraging mode. In:
Lizard Ecology (eds. Reilly, S., McBrayer, L. & Miles, D.B.). Cambridge University Press,
New York, pp. 49–93.

Nagy, K.A., Huey, R.B. & Bennett, A.F. (1984). Field Energetics and Foraging Mode of Kalahari
Lacertid Lizards. *Ecology*, 65, 588–596.

Orme, D., Freckleton, R., Thomas, G., Petzoldt, T., Fritz, S., Isaac, N., *et al.* (2018). caper:
Comparative Analyses of Phylogenetics and Evolution in R.

Packard, G.C. & Boardman, T.J. (1988). The Misuse of Ratios, Indices, and Percentages in
Ecophysiological Research. *Source: Physiological Zoology*, 61, 1–9.

Parker, G.A. & Begon, M. (1986). Optimal Egg Size and Clutch Size : Effects of Environment and
Maternal Phenotype. *The American naturalist*, 128, 573–592.

Perry, G. (1999). The evolution of search modes: Ecological versus phylogenetic perspectives.
American Naturalist, 153, 98–109.

Pianka, E.R. (1966). Convexity , Desert Lizards , and Spatial Heterogeneity. *Ecology*, 47, 1055–
1059.

Pianka, E.R. & Vitt, L.J. (2003). *Lizards: windows to the evolution of diversity*. University of
California Press; First edition (May 15, 2006).

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R Core Team. (2021). nlme: Linear and
Nonlinear Mixed Effects Models.

Primack, R.B. (1979). Reproductive Effort in Annual and Perennial Species of Plantago
(Plantaginaceae). *The American Naturalist*, 114, 51–62.

R Core Team. (2021). R: A Language and Environment for Statistical Computing.

Revell, L.J. (2012). phytools: An R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution*, 3, 217–223.

Reznick, D.N., Buckwalter, G., Groff, J. & Elder, D. (2001). The evolution of senescence in
natural populations of guppies (*Poecilia reticulata*): a comparative approach. *Experimental
Gerontology*, 36, 791–812.

Reznick, D.N. & Endler, J.A. (1982). The impact of predation on life history evolution in
Trinidadian guppies: genetic basis of observed life history patterns. *Evolution*, 36, 1236–
1250.

Richardson, J.L., Urban, M.C., Bolnick, D.I. & Skelly, D.K. (2014). Microgeographic adaptation
and the spatial scale of evolution. *Trends in Ecology and Evolution*, 29, 165–176.

Roff, D. (2002). *Life History Evolution*. Sinauer Associates is an imprint of Oxford University
Press.

Schoener, T.W. (1971). Theory of Feeding Strategies. *Annual Review of Ecology and
Systematics*, 2, 369–404.

Shine, R. (1980). "Costs" of Reproduction in Reptiles. *Oecologia*, 92–100.

Silva, K.M.P., Braz, H.B., Kasperoviczus, K.N., Marques, O.A. v & Almeida-Santos, S.M. (2020).
Reproduction in the pitviper *Bothrops jararacussu*: large females increase their reproductive
output while small males increase their potential to mate. *Zoology*, 142, 125816.

Stearns, S.C. (2000). Life history evolution: Successes, limitations, and prospects.
Naturwissenschaften, 87, 476–486.

Stuginski, D.R., Navas, C.A., de Barros, F.C., Camacho, A., Bicudo, J.E.P.W., Grego, K.F., *et al.*
(2018). Phylogenetic analysis of standard metabolic rate of snakes: a new proposal for the
understanding of interspecific variation in feeding behavior. *Journal of Comparative
Physiology B: Biochemical, Systemic, and Environmental Physiology*, 188, 315–323.

Tinkle, D.W. (1967). The Life and Demography of the Side-Blotched Lizard. *Miscellaneous
Publications, Museum of Zoology, University of Michigan*, 182.

Vitt, L.J. & Congdont, J. (1978). Body shape, reproductive effort, and relative clutch mass in
lizards: resolution of a paradox. *The American Naturalist*, 112, 595–608.

Vitt, L.J. & Price, H. (1982). Ecological and Evolutionary Determinants of Relative Clutch Mass in
Lizards. *Herpetologica*, 38, 237–255.

Vitt, L.J., Pianka, E.R., Cooper, W.E.Jr. & Schwenk, K. (2003). History and the global ecology of
squamate reptiles. *American Naturalist*, 162, 44–60.

Wasiolka, B., Blaum, N., Jeltsch, F. & Henschel, J. (2009). Behavioural responses of the lizard
Pedioplanis l. lineoocellata to overgrazing. *Acta Oecologica*, 35, 157–162.

Webb, J.K., Brook, B.W. & Shine, R. (2003). Does foraging mode influence life history traits? A
comparative study of growth, maturation and survival of two species of sympatric snakes
from south-eastern Australia. *Austral Ecology*, 28, 601–610.

Werner, E.E. & Anholt, B.R. (1993). Ecological consequences of the trade-off between growth
and mortality rates mediated by foraging activity. *Am. Nat*, 142, 242–272.

Zheng, Y. & Wiens, J.J. (2016). Combining phylogenomic and supermatrix approaches, and a
time-calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes
and 4162 species. *Molecular Phylogenetics and Evolution*, 94, 537–547.

Figures and Tables

Fig. 1. Conceptual model depicting potential relationships between foraging behavior, energetics,

predation risk, and reproductive effort. The predicted relationships derived from long-standing

models of life-history theory (see text).

Fig. 2. Random sample of stochastic character maps depicting the evolution of foraging mode in

485 species of lizards (see supporting material for estimates of ancestral states in each internal

node). Bars at the tips of the phylogeny represent log-transformed values of reproductive effort

for all lizards, but not the outgroup *Sphenodon punctatus*. Major clades are enumerated as

follows: 1) Gekkota, 2) Scincoidea, 3) Lacertoidea, 4) Anguimorpha, 5) Toxicofera, and 6)

Iguania. Lizard photos by Mark O'Shea.

Fig. 3. Effects of maternal length and foraging mode on the evolution of the reproductive effort of

lizards, as determined by phylogenetic generalized least squares analysis. A) In Lacertilians,

widely foraging species have evolved the greatest reproductive effort. However, our analysis

revealed a right-skewed distribution of body length in this clade. B) The same pattern holds in

lizards with small and normal-distributed body length, such as skinks. According to the estimated

parameters of the model, sit-and-wait species have the lowest reproductive effort

$(\beta = -1.026, Std. Error = 0.507, t - value = -2.021, p = 0.046)$.

Fig. 4. Theoretical model relating the amount of energy gain for reproduction, *m*, as a function of

time spent foraging, *t*, and maternal size. A) Larger females reach their maximum capacity at a

higher value of *m* than small females. B) Widely foraging females that are smaller but more

efficient foragers may produce a greater reproductive effort than larger sit-and-wait females. C)

Widely foraging females may also produce a greater reproductive effort than sit-and-wait females

if they are both larger and more efficient foragers. $t_{wf} \; and \; t_{sw}$ in (B) represent the optimal

foraging time of widely foraging females and sit-and-wait females, respectively.

517   **Table 1.** Summary statistics for the most likely model of evolution of the reproductive effort,

518   based on the ranking of AICc for potential candidate models included in our analyses.

Terms	numDF	F-value	p-value
(Intercept)	1	27.560	<.0001
female.SVL	1	533.959	<.0001
foraging.mode	2	0.533	0.587
female.SVL:foraging.mode	2	19.910	<.0001