A brain atlas of the camouflaging dwarf cuttlefish, *Sepia bandensis*

Tessa G. Montague[1,2,*], Isabelle J. Rieth[1], Sabrina Gjerswold-Selleck[1], Daniella Garcia-Rosales[1], Sukanya Aneja[3], Dana Elkis[3], Nanyan Zhu[1], Sabrina Kentis[1], Frederick A. Rubino[4], Adriana Nemes[1], Katherine Wang[1], Luke A. Hammond[1], Roselis Emiliano[1], Rebecca A. Ober[1], Jia Guo[1], Richard Axel[1,2,*].

[1]The Mortimer B. Zuckerman Mind Brain Behavior Institute, Department of Neuroscience, Columbia University, New York, NY 10027, USA
[2]Howard Hughes Medical Institute, Columbia University, New York, NY 10027, USA
[3]Interactive Telecommunications Program, New York University, New York, NY 10003, USA
[4]Department of Cell Biology, NYU Grossman School of Medicine, New York, NY 10016, USA

*Correspondence: tessa.montague@columbia.edu and ra27@columbia.edu

Summary

The coleoid cephalopods (cuttlefish, octopus, and squid) are a group of soft-bodied marine mollusks that exhibit an array of interesting biological phenomena, including dynamic camouflage, complex social behaviors, prehensile regenerating arms, and large brains capable of learning, memory, and problem-solving [1–10]. The dwarf cuttlefish, *Sepia bandensis*, is a promising model cephalopod species due to its small size, substantial egg production, short generation time, and dynamic social and camouflage behaviors [11]. Cuttlefish dynamically camouflage to their surroundings by changing the color, pattern and texture of their skin. Camouflage is optically-driven, and is achieved by expanding and contracting hundreds of thousands of pigment-filled saccules (chromatophores) in the skin, which are controlled by motor neurons emanating from the brain. We generated a dwarf cuttlefish brain atlas using magnetic resonance imaging (MRI), deep learning, and histology, and we built an interactive web tool (cuttlebase.org) to host the data. Guided by observations in other cephalopods [12–20], we identified 32 brain lobes, including two large optic lobes (75% the total volume of the brain), chromatophore lobes whose motor neurons directly innervate the chromatophores of the color-changing skin, and a vertical lobe that has been implicated in learning and memory. This brain atlas provides a valuable tool for exploring the neural basis of cuttlefish behavior.

Results

An MRI-based 3D brain atlas of the dwarf cuttlefish

The dwarf cuttlefish, *Sepia bandensis,* is a small, tropical species from the Indo-Pacific (Figure 1A) that exhibits a rich repertoire of behaviors. The skin of the dwarf cuttlefish is covered in hundreds of thousands of chromatophores (pigment-filled saccules surrounded by radial muscles) whose expansion states are controlled by motor neurons projecting from the brain [21]. During camouflage (Figure 1B), conspecific communication (Figures 1C,D), and deimatic behavior, dwarf cuttlefish dynamically alter the color, pattern and texture of their skin using chromatophores and subcutaneous papillae. This process is driven by visual circuits in the brain, and is therefore extremely rapid, occurring in milliseconds [22, 23]. Thus, the skin patterning of cuttlefish reflects both the organism's perception of the external world and its internal state.

The dwarf cuttlefish brain is located posterior and medial to the eyes and is encased on the posterior side by cranial cartilage (Figure 1E). We performed *ex vivo* magnetic resonance imaging (MRI) of the brains of 8 adult dwarf cuttlefish (4 males, 4 females) at 50 μm isotropic resolution and then combined manual segmentation with deep learning techniques to extract each brain from its surrounding tissue (see Methods, Figure S1). Using 6 independent annotators, we segmented the brain lobes using prior neuroanatomical descriptions as a guide [12, 18, 19]. Finally, we co-registered the 8 brains to create a merged, annotated template brain (Figure 1F).

The *ex vivo* dwarf cuttlefish brain is 94% the volume of the *ex vivo* mouse brain (Table S1, Methods), which is consistent with the cuttlefish possessing a deep behavioral repertoire, akin to that of a small mammal. The cuttlefish brain surrounds the esophagus, and can been coarsely divided into a supraesophageal mass and a subesophageal mass [24, 25] that we further divided into 32 discrete lobes (Table 1). This value is in accord with previous anatomical studies in other species, but it remains possible that the annotated lobes can be subdivided further. The largest brain lobes, the optic lobes (O), receive direct projections from the retina and are engaged in visual processing [13] (Figures 1Fi,iii). They comprise 75% of the total volume of the brain (Table S1), consistent with the dominant role of vision in

Figure 1. 3D brain atlas of the dwarf cuttlefish, *Sepia bandensis*. A) Adult dwarf cuttlefish (~8 cm in length). B) Cuttlefish camouflage to their surroundings by changing the color and texture of their skin. C) The innate skin pattern and posture adopted by male dwarf cuttlefish during aggressive interactions. D) An innate skin pattern frequently adopted by dwarf cuttlefish in the presence of conspecifics. E) Anatomy of a cuttlefish. The brain is encased on its posterior side by rigid cartilage. F) 3D template brain, based on MR imaging of 8 brains (4 males, 4 females) and manual annotations of 32 brain lobes and 12 nerve tracts. For abbreviations, see Table 1. (i) Anterior view, (ii) Right view (shown without the optic lobes) and (iii) Posterior view.

guiding many cephalopod behaviors [26]. One relevant projection of the optic lobe is the lateral basal lobe (LB), an intermediate station in the camouflage pathway [12, 13, 18, 20]. The lateral basal lobe may use processed visual information from the optic lobes to compute the most appropriate skin pattern components needed for camouflage. The vertical lobe complex lies on the dorsal side of the supraesophageal mass (Figure 1F). This complex is considered to be the learning and memory center of the brain [27] and may facilitate short-term [28], spatial [29], and visual [30] learning in cephalopods, including working [31] and episodic-like memory [32]. The vertical lobe (V) is directly underneath the dorsal cranial cartilage – an ideal structure for anchoring an electrode or GRIN lens for neural recordings (Figure 3B).

The subesophageal mass comprises multiple structures that elicit simple motor actions. There is a rough mapping of lobe position in the brain to motor target in the body. For instance, the lobes in the anterior portion of the subesophageal mass control movements of structures in the anterior portion of the cuttlefish: the brachial lobe (B) controls the arms, the lateral pedal lobes (LP) control movements of the eyes (Figure 1Fii), and the anterior chromatophore lobes control the chromatophores of the head and arms [18]. Connections between the anterior and posterior chromatophore lobes are thought to coordinate skin patterns between the head, arms and body.

The posterior region of the subesophageal mass (the posterior subesophageal mass, PSM), projects through an opening in the cranial cartilage (Figure 3B). Structures in this region control motor actions in the more posterior aspect of the cuttlefish. For instance, the posterior chromatophore lobes (PCL) control the chromatophores of the mantle, the fin lobes (F) control movement of the fins, and the pallioviseral lobe (PV) controls escape movements and inking, which are mediated by the funnel [18] (Figure 1Fiii). Some cephalopod brain lobes have been assigned function through lesion [29, 33–37] and electrophysiological studies [18, 38–43]. However, the functions of most brain regions are unknown.

A histological brain atlas for the dwarf cuttlefish

We complemented the anatomical description of the cuttlefish brain with histological examination of the

entire brain to obtain cellular resolution. We sectioned the brain in the transverse, horizontal and sagittal

planes (Figure 2A) and stained the sections with Phalloidin, an F-actin peptide that labels axons, and

NeuroTrace, a Nissl stain that labels neuronal cell bodies and glia. Our annotated 3D MRI datasets and

prior neuroanatomical descriptions [12, 18, 19] were used to describe the histological organization of 32

brain lobes and 12 nerve tracts (Figures 2B-D).

Phalloidin and NeuroTrace staining confirmed previously described organizational features of the

cephalopod brain. The majority of lobes are discrete and bounded, a feature not uniformly observed in

vertebrate brains: each lobe contains an outer perikaryal layer of cell bodies that surrounds a dense,

inner neuropil (Figures 2B-D). Dwarf cuttlefish neuronal cell bodies range in diameter from ~5 um (e.g.

the vertical and optic lobes) to <80 um in some motor areas (e.g. the fin and pedal lobes) (Figures 2E-G).

Similar to other coleoid cephalopods, the *Sepia bandensis* optic lobe consists of two ordered outer layers,

the inner and outer granular layers, apposing a zone of fibers (Figure 2E) [13, 20, 44]. Because the

cephalopod retina contains only photoreceptors, the outer layers of the optic lobe may function like the

vertebrate retina and perform the initial stages of visual processing [44, 45]. Beneath the outer layers, the

cephalopod optic lobe features a novel neural organization: a central medulla that features cell 'islands'

(Figure 2F) connected in an elaborate tree-like structure [13, 20, 46]. Cell bodies near the cortex (the

'branches') appear to form columns (Figures 2B,C), which may reflect the presumed retinotopic order of

the outer layers [13, 20, 47].

Five pairs of nerves project from the cuttlefish brain that are easily distinguishable in histological slices

and contain many of the brain's major efferents. The largest nerves, the pallial nerves, exit the posterior

side of the brain and contain motor fibers that innervate the chromatophores of the mantle skin and the

muscles of the mantle and fin (Figure 2B) [12]. The motor neurons that originate in the posterior

chromatophore lobe exit via the pallial nerve, pass through the stellate ganglion, and directly innervate

the radial muscles of the chromatophores [48]. Severing the pallial nerve results in blanched skin, loss of

Figure 2. Histological brain atlas of the dwarf cuttlefish, *Sepia bandensis.* A) The dwarf cuttlefish brain was sectioned in the transverse, horizontal and sagittal planes. B-D) Representative histological slices of the cuttlefish brain stained with NeuroTrace, a Nissl stain that labels cell bodies, and annotated with 32 brain lobes and 12 nerve tracts. Scale bar, 1 mm. For abbreviations see Table 1. E-G) Confocal images of cuttlefish tissue stained with NeuroTrace (purple) and Phalloidin (an F-actin peptide that labels axons, green). Boxed areas in C show approximate locations of confocal images within the brain. E) Optic lobe layered cortex, F) Optic lobe medulla, G) Anterior pedal lobe. Confocal scale bars, 50 μm.

the skin's textural control, and a limp fin on the ipsilateral side of the body [49]. Closely apposed to the

pallial nerves are the collar nerves, which innervate the collar muscles and valves of the mantle (Figure

2B) [12]. The visceral nerves are positioned near the midline on the posterior side of the brain and

innervate the heart, ink sac muscles, digestive, reproductive and excretory systems [12]. Finally, the head

retractor nerves, positioned ventral to the lateral basal lobes, innervate the muscles that connect the head

to the mantle (Figure 2B) [12]. When dwarf cuttlefish sense danger, they can retract their head inside their

body using these muscles.

Cuttlebase - a web tool for visualizing the cuttlefish brain

We built an interactive web tool, Cuttlebase, to maximize the utility of our brain atlas (Figure 3A).

Cuttlebase features multiple tools for the dwarf cuttlefish, including the annotated histological sections of

the brain in 3 planes (Figure 3A), the 3D brain model described above (Figure 3B), and a 3D model of an

entire adult cuttlefish labeled with 26 organs, including the animal's three hearts, gills, ink sac, beak,

cuttlebone and digestive system (Figure 3C). Furthermore, the 3D body model features the eight brachial

nerves that innervate the cuttlefish's arms, and the two pallial nerves, which exit the posterior aspect of

the brain, circumscribe the digestive gland, and then pass through a hole in the mantle musculature to

reach the stellate ganglion [50]. Cuttlebase is easy-to-use, with an array of features including responsive

color-coded labels for each brain region, a dynamic scale bar, the ability to zoom, rotate and screenshot

the data, and synchronized graphics that denote the brain's orientation.

Cuttlebase was designed for both educational and research settings. For its use in education, we added

features that increase accessibility, including: (i) A landing page that introduces cuttlefish behavior, brain

anatomy, evolution and animal care; (ii) An info page that shows the cuttlefish's gross anatomy and the

sectioning planes used in the histological atlas; (iii) An option to show the 3D brain model in the context of

the cuttlefish's body; and (iv) Mouseover descriptions of the function of each brain lobe. For expert users,

we included an array of more advanced features. First, we included a dynamic scale bar in the histology

mode to help users measure distances within the brain and devise brain coordinates for stereotactic

Figure 3. Cuttlebase is a scientific toolkit for the dwarf cuttlefish. A) Cuttlebase hosts multiple dwarf cuttlefish tools including an interactive histological brain atlas, which features color-coded and responsive labels, a dynamic scale bar, and the ability to screenshot, zoom and rotate the data. B) The 3D brain atlas can be visualized with the cranial cartilage and features a synchronized graphic that denotes the brain's orientation. C) Cuttlebase also features a body atlas, labeled with 26 organs and nerves.

surgeries. Second, we imaged the cranial cartilage — the only rigid tissue around the brain — and

provided its interactive graphic (Figure 3B) to facilitate the development of methods to head-fix cuttlefish,

which is an important step for stabilizing the brain for two-photon imaging or electrophysiology. Finally, we

made the MRI data, brain model and full-resolution histology data available to download, so users can

carry out their own MRI analyses, annotate the histological images with the results of tracing experiments,

and import the to-scale 3D cuttlefish into modeling and 3D-printing software.

Discussion

The 3D and histological brain atlases presented here serve multiple functions. First, they provide a

comprehensive resource for comparative neuroanatomical analyses. Little is currently known about the

biology of the dwarf cuttlefish (*Sepia bandensis),* whereas the common cuttlefish (*Sepia officinalis)* has

been studied extensively to understand behavior [8], learning and memory [1] and the neural control of

skin patterning [4, 23]. Comparison of our dataset with the neuroanatomy of the common cuttlefish [18,

51] and mourning cuttlefish, *Sepia plangon* [52], reveals substantial neuroanatomic resemblance despite

the species' different sizes, behaviors and habitats [11, 53]. Every cuttlefish species has evolved a

different ensemble of skin pattern components for camouflage and social communication. *Sepia*

bandensis, which live in Indo-Pacific coral reefs, assemble skin patterns with predominantly high

frequency patterns, likely reflecting the visual statistics of their tropical habitat. By contrast, *Sepia*

officinalis, which live in the rocky waters of Europe, include in their repertoire a set of large, disruptive

patterns that resemble rocks [53]. During male social encounters, although *Sepia officinalis* and *Sepia*

bandensis both exhibit a high-contrast, stereotyped aggression pattern, their precise manifestations differ:

Sepia officinalis create a zebra pattern [53], whereas the *Sepia bandensis* aggression pattern is stippled

(Figure 1B) [11]. Additionally, although both species are capable of creating dynamic waves across their

skin, they do so in opposite directions (Video 1) [10]. Such fine-scale differences as these are unlikely to

be reflected at the coarse anatomical level. Spatial transcriptomics, fine anatomical tracing and

developmental tracking may reveal how the neural circuits diverged to facilitate different skin patterns.

Furthermore, the investigation of the neural circuits underlying camouflage may reveal how the optic lobe-

lateral basal lobe-chromatophore lobe pathway creates an internal representation of a visual scene, and

then recreates an approximation of the scene on the skin. Recording neural activity in the camouflage

circuit, as well as precise anterograde and retrograde tracing studies, may answer questions such as: *Is*

the cuttlefish capable of only a limited number of patterns that when combined recapitulate the natural

scene? Is the neural network underlying camouflage capable of exhibiting experience-dependent

plasticity? Are the activity patterns in the chromatophore lobe topographic with patterns on the skin, or are

they an abstraction? How is an internal, abstract representation of the visual world transformed to a more

figurative representation on the skin?

Comparison of the dwarf cuttlefish brain with other decapodiformes (10-limbed cuttlefish and squid) and

octopodiformes (8-limbed octopus and vampire squid) reveals substantial organizational similarities and a

few key differences. First, unlike the octopodiformes, the decapodiformes possess a giant-fiber system,

which is used for funnel-based jet propulsion during escape behaviors. The giant cells are located in the

ventral magnocellular lobe, and receive input from the optic lobes and statocysts (the cephalopod

vestibular system) [54, 55]. Second, octopodiformes employ sophisticated use of their arms and

demonstrate tactile learning; their brachial, pedal and inferior frontal lobes are larger than the

decapodiformes [56]. Third, the octopus vertical lobe is folded into gyri, creating a larger surface area

[12], whereas the cuttlefish vertical lobe is dome-shaped and lacks gyri. Finally, the decapodiformes, but

not octopodiformes, possess a pair of fin lobes used for fin locomotion [18]. Most decapodiformes use two

forms of locomotion: jet propulsion using the funnel, and swimming using the funnel and fins [57].

Interestingly, the dwarf cuttlefish uses an additional mode: it can walk bipedally using its ventral arms

when combined with water ejection through the funnel (Video 2). The gross morphology of the dwarf

cuttlefish brachial and pedal lobes are not readily distinguishable from other cuttlefish species, but deeper

analysis of the circuitry might uncover the evolution of a control system for coordinated locomotion.

This atlas also provides a valuable resource for comparisons across phyla. For instance, despite ~600

million years of separation between cuttlefish and fruit flies [58], the brains of these species share coarse

organizational features. The esophagus runs through the center of the brain, two large optic lobes flank

the central brain mass, and the learning and memory centers - the mushroom body in *Drosophila* and the

vertical lobe in cuttlefish - lie on the dorsal side of the brain. Furthermore, the two outer layers of the

cuttlefish optic lobe may share properties with the lamina and medulla of the dipteran visual system [59].

Interestingly, recent single-cell gene expression profiling of a developing octopus brain has revealed

molecular resemblances between the Kenyon cells of the *Drosophila* mushroom body and the vertical

lobe cells of the octopus brain [60].

The dwarf cuttlefish brain atlas presented here serves as both an educational resource and a tool for the

investigation of the neural basis of cephalopod behavior. Users can employ Cuttlebase to plan the target

angle and depth of electrode insertion or to design head plates that anchor GRIN lenses for calcium

imaging. Following the completion of *in vivo* experiments, the histological atlas affords users the ability to

identify the experimentally targeted brain regions by directly comparing post-operative tissue slices with

the atlas. Beyond the interrogation of neural circuits by imaging, recording or tracing, the histological atlas

can be used to assign function to brain lobes by mapping immediate early gene expression following a

behavioral paradigm, and to map spatially-restricted gene expression to neuroanatomy. The recent

generation of squid and octopus cell atlases by single-cell RNA-seq complement the generation of

neuroanatomical atlases, providing a more comprehensive view of brain organization [60–63].

Finally, it is important to note that this atlas represents a first attempt to demarcate the brain lobes of the

dwarf cuttlefish, but these demarcations will likely need to be adjusted as we learn more about the

functional organization of the cephalopod brain. At present, the vast majority of functions assigned to the

cuttlefish brain lobes originated from a single study, in which brain lobes were systematically stimulated

by hand [18]. The application of new technologies, such as transgenic tools, will likely lead to a more

nuanced view of lobe function, and the development and application of retrograde and anterograde

tracing methods will improve our understanding of connectivity within the brain.

In summary, Cuttlebase was designed to aid the investigation of cuttlefish biology, as well as foster the

curiosity of scientists and non-scientists alike. The generation of the neuroanatomical and histological

atlases for the dwarf cuttlefish may inform studies in other cephalopods, for which atlases do not exist,

and now renders *Sepia bandensis* a more facile system for the study of the neural control of camouflage.

Acknowledgements

We wish to thank Connor Gibbons, Sonia Thomas, Telicia Lewis, Sarah Wilson and Josh Barber for

cuttlefish care, the Zuckerman Institute Cellular Imaging platform for instrument use and technical

support, Lokke Highstein for IT support, and Gwenyth Card and Carrie Albertin for valuable discussions.

This project was funded by a Zuckerman Institute MRI Seed Grant (T.G.M. and R.A.), the Howard

Hughes Medical Institute Hanna H. Gray Fellowship (T.G.M.), the Zuckerman Institute BRAINYAC

Program (R.E.) and the Howard Hughes Medical Institute (R.A.)

Author contributions

T.G.M., R.A.O. and J.G. conceived the project. T.G.M., I.J.R. and J.G. performed MRI experiments.

T.G.M. performed histology experiments. S.G-R. conducted MRI data processing steps, with input from

N.Z. and J.G. The MRI data was annotated by I.G.R., T.G.M., D.G-R., S.K., F.A.R., A.N., K.W., and R.E.

The histology data was annotated by I.J.R. and T.G.M. L.A.H. developed the BrainJ histology data

processing software. D.E. designed and S.A. developed the Cuttlebase website, respectively. J.G. and

R.A. provided supervision. T.G.M. wrote the original draft of the manuscript. T.G.M. and R.A. reviewed

and edited the manuscript. T.G.M. and R.A. acquired funding for the project.

Declaration of interests

257 The authors have no competing interests.

Methods

Ethics statement

263 The use of cephalopods in laboratory research is currently not regulated in the USA. However, Columbia

264 University has established strict policies for the ethical use of cephalopods, including operational

265 oversight from the Institutional Animal Care and Use Committee (IACUC). All of the cuttlefish used in this

266 study were handled according to approved IACUC protocols (AC-AABE3564), including the use of deep

267 anesthesia and the minimization and prevention of suffering.

3D brain atlas

MRI acquisition

8 adult cuttlefish (4 male, 4 female) were euthanized in 10% ethanol, and the brain and connected eyes

were surgically removed and fixed overnight in a solution of 4% paraformaldehyde (PFA) in filtered

artificial seawater (FASW) at 4°C. The fixed brains were washed in PBS, incubated in 0.2% OMNISCAN

(gadodiamide) for 2 days at 4°C to increase contrast, and then suspended in fomblin in a 15 mL conical

tube. Imaging was performed on a Bruker BioSpec 94/30 horizontal small animal MRI scanner (field

strength, 9.4 T; bore size, 30 cm) equipped with a CryoProbe and ParaVision 6.0.1 software (Bruker). A

23 mm 1H circularly polarized transmit/receive-capable mouse head volume coil was used for imaging.

For each cuttlefish, one scan was acquired of the brain and eyes (to obtain a scan of the entire cranial

cartilage), and a higher resolution scan was acquired of the brain only. T1-weighted images were

acquired with a Fast Low Angle Shot (FLASH) sequence (brain/eye scan: TR = 50 ms, TE = 10 ms, FOV

= 30 x 24 x 15 mm^3, voxel size = 100 x 100 x 100 μm^3, scan time = 33 m 12 s; brain-only scan: TR = 50

ms, TE = 8.5 ms, FOV = 14 x 12 x 11 mm^3, voxel size = 60 x 60 x 60 μm^3, scan time = 4 hr 28 m).

MRI processing & brain extraction

All scans underwent N4 bias field correction [64]. Whole brain scans were isotropically upsampled to 50

μm isotropic resolution with cubic B-spline interpolation. To computationally extract the brains from their

surrounding tissue, brain masks were generated by an in-house deep learning model [65], which was pre-

trained with brain masks manually annotated in 3D Slicer [66]. The deep learning brain masks were

manually polished using the 3D Slicer Segment Editor and then used to extract the brain from each brain-

only MRI scan ("brain-extracted images") (Figure S1). For the brain/eye scans, the cranial cartilage was

manually segmented in 3D Slicer using the Segment Editor. The cartilage masks were used to extract the

cartilage from each brain/eye MRI scan ("cartilage-extracted images").

Segmentation

Two of the whole brain scans (1 male, 1 female) were manually segmented in 3D Slicer by 6 independent

annotators guided by prior neuroanatomical descriptions [18, 19]. The brain label maps for each subject

were merged using pixel-level majority voting, transformed to the remaining 6 subjects, and then

manually corrected, resulting in 8 brain label maps corresponding to the 8 subjects. Lobe locations were

determined by prior annotations [18, 19] and precise lobe boundaries were estimated using the location of

an outer cell body layer, which is a feature of most cephalopod brain lobes. The combined use of high-

resolution histological data and three-dimensional MRI data aided the discrimination of lobes in 3D space.

Generation of the template brain

The final MRI atlas was built by merging the brain template, built from the high-resolution brain-only scans

and their mirror images, with the cranial cartilage template, built from the brain/eye scans. First, to

generate each template, a population average of brain-extracted or cartilage-extracted images was

constructed through an iterative process by averaging the co-registered images over multiple cycles using

a symmetric diffeomorphic registration algorithm [67]. The brain label map (annotations) for each subject

was diffeomorphically transformed to the whole brain template space and combined through pixel-level

majority voting. The brain/eye template was isotropically upsampled to match the 50 μm resolution of the

whole brain scans, and the whole brain template underwent rigid registration to align it with the brain/eye

template. Finally, the brain regions of the whole brain template were combined with the cartilage regions

of the whole head template to build a single atlas. In the regions where the two templates overlapped,

pixel values of the high-resolution whole brain template were selected. The cuttlefish brain label map was

smoothed by taking a majority vote in a local neighborhood with a 3 x 3 x 3 kernel size.

Brain volume calculation

A mouse MRI brain model (15 µm resolution, average of 18 *ex vivo* subjects) was downloaded from the

Australian Mouse Brain Mapping Consortium (nonsymmetric version, NiFTI format,

https://imaging.org.au/AMBMC/Model), and then downsampled to 50 µm resolution and imported into 3D

Slicer. Using the Editor tool at a threshold of zero, a mask was generated of the entire mouse brain, and

the spinal cord was manually removed using the eraser tool. The volume of the mouse brain mask was

calculated using the Label Statistics tool (total volume: 332.7 mm^3). To calculate the volume of the

cuttlefish brain, the volume of each brain lobe in the merged, template brain was calculated using the

Label Statistics tool and summed (total volume: 312.1 mm^3), see Table S1.

<u>**Histological atlas**</u>

Adult cuttlefish were euthanized in 10% ethanol, and the brain and connected eyes were surgically

removed and fixed overnight in a solution of 4% paraformaldehyde (PFA) in filtered artificial seawater

(FASW) at 4°C. Note that fixing brains in a solution of PBS (instead of FASW) created abnormal cell

morphologies. The fixed brains were washed in PBS, the eyes were removed with a scalpel, and the

brains were incubated in 10% sucrose overnight followed by 30% sucrose overnight at 4°C. The brains

were embedded in OCT on dry ice and stored at -80°C. Each brain was sliced in 100 µm sections on a

cryostat (Leica CM3050 S), and the sections were dried overnight at room temperature. The sections

were stained with Phalloidin (Life Technologies #A12379, 1/40 dilution) and NeuroTrace (Life

Technologies #N21482, 1/20 dilution) and then imaged on a custom-built Nikon AZ100 Multizoom Slide

Scanner. Images were registered using BrainJ (http://github.com/lahammond/BrainJ). Brightness and

contrast were adjusted uniformly across each image, and surrounding tissue was removed manually from

the image in FIJI and Photoshop. The data was manually segmented in 3D Slicer by two annotators using

a neuroanatomical study [18] and our 3D brain data for reference.

Whole body atlas

An adult male cuttlefish was anesthetized in $MgCl_2$ (17.5g/L) and then euthanized in 10% ethanol, fixed

for 2 days in 4% PFA/FASW at 4°C, and then transferred to 0.2% OMNISCAN (gadodiamide) for 2 days

at 4°C to increase contrast. The fixed specimen was suspended in fomblin in a custom-made vessel and

imaged on a Bruker BioSpec 94/30 horizontal small animal MRI scanner (field strength, 9.4 T; bore size,

30 cm) equipped with a CryoProbe and ParaVision 6.0.1 software (Bruker). A 112/86-mm 1H circularly

polarized transmit/receive-capable volume coil was used for imaging. T1-weighted images were acquired

with a FLASH sequence (TR = 55 ms, TE = 17 ms, FOV = 90 x 48 x 38 mm^3, voxel size = 100 x 100 x

100 μm^3, scan time = 2 hr 47 m). The scan underwent N4 bias field correction [64] and was manually

segmented in 3D Slicer using anatomical descriptions [68].

Cuttlebase

The Cuttlebase web content is delivered using React, a Javascript front-end framework for dynamic

websites. React-three-fiber (a Three.js wrapper for React) is used to assist with interactions in the 3D

view.

3D brain

In 3D Slicer, each segment (brain lobe or tract) of the final, template brain was exported as an STL file

and then labelled with a unique identifier in Blender, a 3D authoring software. The 3D model was

exported as a GLB, and then imported into a webpage using Three.js - a Javascript library for handling

3D content on the web. A custom web-interface was created to assign colors to each of the region

meshes, and this data was exported as a JSON file.

Histology

To convert the histological annotations to high-resolution images that could be toggled on Cuttlebase, each segment (brain lobe or tract) for each brain (horizontal, sagittal and transverse) was converted to a binary label map in 3D Slicer and saved as a TIFF stack. Each TIFF stack was resized to the canvas size of the original image in Fiji [69], and saved as a JPEG Stack in monochrome. The images were then inverted and processed with Potrace (through a custom Node.js script), to create SVGs for each region of each layer. The SVGs for all regions in a single layer were combined, and each region was assigned the color corresponding to the 3D atlas. These images, along with the originals, were resized and cropped for more efficient web delivery. Additionally, the cartilage in each Phalloidin section was isolated (by outlining) using Adobe Illustrator, and exported as a PNG. These images were used as masks on the NeuroTrace layers with the command-line tool ImageMagick, to automate this process for the remaining images. Custom Bash scripts were used to manage and organize the large amounts of data and the processing steps required.

Data availability

All data generated in this study is available to download from cuttlebase.org/downloads.

Video legends

Video 1. Dwarf cuttlefish produce waves (of unknown function) on their skin.

Video 2. Dwarf cuttlefish can walk bipedally using their ventral arms when coupled with water ejection through the funnel.

Figure legends

Figure 1. 3D brain atlas of the dwarf cuttlefish, Sepia bandensis. A) Adult dwarf cuttlefish (~8 cm in length). B) Cuttlefish camouflage to their surroundings by changing the color and texture of their skin. C)

The innate skin pattern and posture adopted by male dwarf cuttlefish during aggressive interactions. D)
An innate skin pattern frequently adopted by dwarf cuttlefish in the presence of conspecifics. E) Anatomy
of a cuttlefish. The brain is encased on its posterior side by rigid cartilage. F) 3D template brain, based on
MR imaging of 8 brains (4 males, 4 females) and manual annotations of 32 brain lobes and 12 nerve
tracts. For abbreviations, see Table 1. (i) Anterior view, (ii) Right view (shown without the optic lobes) and
(iii) Posterior view.

Figure 2. Histological brain atlas of the dwarf cuttlefish, Sepia bandensis. A) The dwarf cuttlefish
brain was sectioned in the transverse, horizontal and sagittal planes. B-D) Representative histological
slices of the cuttlefish brain stained with NeuroTrace (a Nissl stain that labels cell bodies) and annotated
with 32 brain lobes and 12 nerve tracts. Scale bar, 1 mm. For abbreviations, see Table 1. E-G) Confocal
images of cuttlefish tissue stained with NeuroTrace (purple) and Phalloidin (an F-actin peptide that labels
axons, green). Boxed areas in C show approximate locations of confocal images within the brain. E) Optic
lobe layered cortex, F) Optic lobe medulla, G) Anterior pedal lobe. Confocal scale bars, 50 µm.

Figure 3. Cuttlebase is a scientific toolkit for the dwarf cuttlefish. A) Cuttlebase hosts multiple dwarf
cuttlefish tools including an interactive histological brain atlas, which features color-coded and responsive
labels, a dynamic scale bar, and the ability to screenshot, zoom and rotate the data. B) The 3D brain atlas
can be visualized with the cranial cartilage and features a synchronized graphic that denotes the brain's
orientation. C) Cuttlebase also features a body atlas, labeled with 21 organs and tissues.

Figure S1. Deep learning pipeline for improving manually-generated brain masks.

Table S1. Cuttlefish brain lobe volumes.

References

1.   Turchetti-Maia A., Shomrat T., and Hochner B. (2019). The Vertical Lobe of Cephalopods. 558-
574.
2.   Fiorito G., von Planta C., and Scotto P. (1990). Problem solving ability of Octopus vulgaris
Lamarck (Mollusca, Cephalopoda). Behav Neural Biol. 53, 217-230.
3.   Richter J.N., Hochner B., and Kuba M.J. (2016). Pull or Push? Octopuses Solve a Puzzle
Problem. PLoS One. 11, e0152048.
4.   Reiter S., Hülsdunk P., Woo T., Lauterbach M.A., Eberle J.S., Akay L.A., Longo A., Meier-Credo
J., Kretschmer F., Langer J.D. et al. (2018). Elucidating the control and development of skin patterning in
cuttlefish. Nature. 562, 361-366.
5.   Schnell A.K., and Clayton N.S. (2019). Cephalopod cognition. Curr Biol. 29, R726-R732.
6.   Imperadore P., and Fiorito G. (2018). Cephalopod tissue regeneration: consolidating over a
century of knowledge. Frontiers in physiology. 9, 593.
7.   Amodio P., Boeckle M., Schnell A.K., Ostojíc L., Fiorito G., and Clayton N.S. (2019). Grow Smart
and Die Young: Why Did Cephalopods Evolve Intelligence. Trends Ecol Evol. 34, 45-56.
8.   Hanlon R.T., and Messenger J.B. (2018). Cephalopod Behaviour, Cambridge University Press).
9.   Osorio D., Ménager F., Tyler C.W., and Darmaillacq A.S. (2022). Multi-level control of adaptive
camouflage by European cuttlefish. Curr Biol. S0960-9822(22)00599.
10.   How M.J., Norman M.D., Finn J., Chung W.S., and Marshall N.J. (2017). Dynamic Skin Patterns
in Cephalopods. Front Physiol. 8, 393.
11.   Montague T.G., Rieth I.J., and Axel R. (2021). Embryonic development of the camouflaging dwarf
cuttlefish, Sepia bandensis. Dev Dyn. 250, 1688-1703.
12.   Young J.Z. (1971). The Anatomy of the Nervous System of Octopus Vulgaris, Oxford University
Press, USA).
13.   Young J.Z. (1974). The central nervous system of Loligo. I. The optic lobe. Philos Trans R Soc
Lond B Biol Sci. 267, 263-302.
14.   Young J.Z. (1976). The nervous system of Loligo. II. Suboesophageal centres. Philos Trans R
Soc Lond B Biol Sci. 274, 101-167.
15.   Young J.Z. (1977). The nervous system of Loligo, III. Higher motor centres: the basal
supraoesophageal lobes. Philosophical Transactions of the Royal Society of London. B, Biological
Sciences. 276, 351-398.
16.   Young J.Z. (1979). The nervous system of Loligo. V. The vertical lobe complex. Philosophical
Transactions of the Royal Society of London. B, Biological Sciences. 285, 311-354.
17.   Messenger J.B. (1979). The nervous system of Loligo IV. The peduncle and olfactory lobes.
Philosophical Transactions of the Royal Society of London. B, Biological Sciences. 285, 275-309.
18.   Boycott B.B. (1961). The functional organization of the brain of the cuttlefish Sepia officinalis.
Proceedings of the Royal Society of London. Series B, Biological Sciences. 153, 503-534.
19.   Chung W.S., Kurniawan N.D., and Marshall N.J. (2020). Toward an MRI-Based Mesoscale
Connectome of the Squid Brain. iScience. 23, 100816.
20.   Young J.Z. (1962). The optic lobes of Octopus vulgaris. Philosophical Transactions of the Royal
Society of London. Series B, Biological Sciences. 245, 19-58.
21.   Messenger J.B. (2001). Cephalopod chromatophores: neurobiology and natural history. Biol Rev
Camb Philos Soc. 76, 473-528.
22.   Hill A.V., and Solandt D.Y. (1935). Myograms from the chromatophores of Sepia. J. Physiol.
Lond. 83, 13P-14P.
23.   Hanlon R. (2007). Cephalopod dynamic camouflage. Curr Biol. 17, R400-4.
24.   Dietl M.J. (1878). Untersuchungen über die Organisation des Gehirns wirbelloser Thiere
(Cephalopoden, Tethys). Sitzungsberichte der Akademie der Wissenschaften in Wien. 77, 481-533.
25.   Hillig R. (1912). Das Nervensystem von Sepia officinalis L. Z. wiss. Zoo. 101, 736-800.

26.   Darmaillacq A.S., Mezrai N., O'Brien C.E., and Dickel L. (2017). Visual Ecology and the
Development of Visually Guided Behavior in the Cuttlefish. Front Physiol. 8, 402.
27.   Boycott B.B., and Young J.Z. (1955). A memory system in Octopus vulgaris Lamarck. Proc R Soc
Lond B Biol Sci. 143, 449-480.
28.   Shomrat T., Zarrella I., Fiorito G., and Hochner B. (2008). The octopus vertical lobe modulates
short-term learning rate and uses LTP to acquire long-term memory. Curr Biol. 18, 337-342.
29.   Graindorge N., Alves C., Darmaillacq A.S., Chichery R., Dickel L., and Bellanger C. (2006).
Effects of dorsal and ventral vertical lobe electrolytic lesions on spatial learning and locomotor activity in
Sepia officinalis. Behav Neurosci. 120, 1151-1158.
30.   Young J.Z. (1991). Computation in the Learning System of Cephalopods. Biol Bull. 180, 200-208.
31.   Sanders F.K., and Young J.Z. (1940). Learning and other functions of the higher nervous centres
of Sepia. Journal of Neurophysiology. 3, 501-526.
32.   Jozet-Alves C., Bertin M., and Clayton N.S. (2013). Evidence of episodic-like memory in
cuttlefish. Curr Biol. 23, R1033-5.
33.   Wells M.J., and Young J.Z. (1975). The subfrontal lobe and touch learning in the octopus. Brain
Res. 92, 103-121.
34.   Boycott B.B., and Young J.Z. (1957). Effects of interference with the vertical lobe on visual
discriminations in Octopus vulgaris Lamarck. Proc R Soc Lond B Biol Sci. 146, 439-459.
35.   Chichery M.P., and Chichery R. (1987). The anterior basal lobe and control of prey-capture in the
cuttlefish (Sepia officinalis). Physiol Behav. 40, 329-336.
36.   Fiorito G., and Chichery R. (1995). Lesions of the vertical lobe impair visual discrimination
learning by observation in Octopus vulgaris. Neurosci Lett. 192, 117-120.
37.   Messenger J.B. (1967). The effects on locomotion of lesions to the visuo-motor system in
octopus. Proceedings of the Royal Society of London. Series B. Biological Sciences. 167, 252-281.
38.   Chichery R., and Chanelet J. (1976). Motor and behavioral responses obtained by stimulation
with chronic electrodes of the optic lobe of Sepia officinalis. Brain Res. 105, 525-532.
39.   Dubas F., Hanlon R.T., Ferguson G.P., and Pinsker H.M. (1986). Localization and stimulation of
chromatophore motoneurones in the brain of the squid, Lolliguncula brevis. J Exp Biol. 121, 1-25.
40.   Bullock T.H., and Budelmann B.U. (1991). Sensory evoked potentials in unanesthetized
unrestrained cuttlefish: a new preparation for brain physiology in cephalopods. J Comp Physiol A. 168,
141-150.
41.   Zullo L., Sumbre G., Agnisola C., Flash T., and Hochner B. (2009). Nonsomatotopic organization
of the higher motor centers in octopus. Curr Biol. 19, 1632-1636.
42.   Hochner B., Brown E.R., Langella M., Shomrat T., and Fiorito G. (2003). A learning and memory
area in the octopus brain manifests a vertebrate-like long-term potentiation. J Neurophysiol. 90, 3547-
3554.
43.   Shomrat T., Graindorge N., Bellanger C., Fiorito G., Loewenstein Y., and Hochner B. (2011).
Alternative sites of synaptic plasticity in two homologous "fan-out fan-in" learning and memory networks.
Curr Biol. 21, 1773-1782.
44.   y Cajal S.R. (1930). Contribución Al Conocimiento De La Retina Y Centros Ópticos De Los
Cefalópodos, Unión Internacional de Ciencias Biológicas, Comité Español).
45.   Young J.Z. (1960). The Visual System of Octopus: Regularities in the Retina and Optic Lobes of
Octopus in Relation to Form Discrimination. Nature. 186, 836-839.
46.   Liu Y.C., Liu T.H., Su C.H., and Chiao C.C. (2017). Neural Organization of the Optic Lobe
Changes Steadily from Late Embryonic Stage to Adulthood in Cuttlefish Sepia pharaonis. Front Physiol.
8, 538.
47.   Pungor J.R., Allen V.A., Songco-Casey J.O., and Niell C.M. (2023). Functional organization of
visual responses in the octopus optic lobe. bioRxiv. 2023.02.16.528734.

48. Sereni E., and Young J.Z. (1932). Nervous degeneration and regeneration in cephalopods.
Pubbl. Staz. Zool. Napoli. 12, 173-208.
49. Gonzalez-Bellido P.T., Scaros A.T., Hanlon R.T., and Wardill T.J. (2018). Neural control of
dynamic 3-dimensional skin papillae for cuttlefish camouflage. iScience. 1, 24-34.
50. Bühler A., Froesch D., Mangold K., and Marthy H.J. (1975). On the motor projection of the stellate
ganglion in Octopus vulgaris. Brain Res. 88, 69-72.
51. Wild E., Wollesen T., Haszprunar G., and Heß M. (2015). Comparative 3D microanatomy and
histology of the eyes and central nervous systems in coleoid cephalopod hatchlings. Organisms Diversity
& Evolution. 15, 37-64.
52. Chung W.-S., López-Galán A., Kurniawan N.D., and Marshall N.J. (2022). The brain structure and
the neural network features of the diurnal cuttlefish Sepia plangon. iScience. 105846.
53. Hanlon R.T., and Messenger J.B. (1988). Adaptive coloration in young cuttlefish (Sepia officinalis
L.): the morphology and development of body patterns and their relation to behaviour. Philosophical
Transactions of the Royal Society of London. B, Biological Sciences. 320, 437-487.
54. Nixon M., and Young J.Z. (2003). The Brains and Lives of Cephalopods, Oxford University
Press).
55. Abbott N.J., Williamson R., and Maddock L. (1995). Cephalopod Neurobiology, Oxford University
Press, USA).
56. Ponte G., Taite M., Borrelli L., Tarallo A., Allcock A.L., and Fiorito G. (2020). Cerebrotypes in
Cephalopods: Brain Diversity and Its Correlation With Species Habits, Life History, and Physiological
Adaptations. Front Neuroanat. 14, 565109.
57. Russell F.S., and Steven G.A. (1930). The swimming of cuttlefish. Nature. 125, 893-893.
58. dos Reis M., Thawornwattana Y., Angelis K., Telford M.J., Donoghue P.C., and Yang Z. (2015).
Uncertainty in the Timing of Origin of Animals and the Limits of Precision in Molecular Timescales. Curr
Biol. 25, 2939-2950.
59. Shinomiya K., Horne J.A., McLin S., Wiederman M., Nern A., Plaza S.M., and Meinertzhagen I.A.
(2019). The Organization of the Second Optic Chiasm of the Drosophila Optic Lobe. Front Neural
Circuits. 13, 65.
60. Styfhals R., Zolotarov G., Hulselmans G., Spanier K.I., Poovathingal S., Elagoz A.M., De Winter
S., Deryckere A., Rajewsky N., Ponte G. et al. (2022). Cell type diversity in a developing octopus brain.
Nat Commun. 13, 7392.
61. Duruz J., Sprecher M., Kaldun J.C., Al-Soudy A.S., Lischer H.E.L., van Geest G., Nicholson P.,
Bruggmann R., and Sprecher S.G. (2023). Molecular characterization of cell types in the squid Loligo
vulgaris. Elife. 12, e80670.
62. Songco-Casey J.O., Coffing G.C., Piscopo D.M., Pungor J.R., Kern A.D., Miller A.C., and Niell
C.M. (2022). Cell types and molecular architecture of the Octopus bimaculoides visual system. Curr Biol.
S0960-9822(22)01623.
63. Gavriouchkina D., Tan Y., Künzli-Ziadi F., Hasegawa Y., Piovani L., Zhang L., Sugimoto C.,
Luscombe N., Marlétaz F., and Rokhsar D.S. (2022). A single-cell atlas of bobtail squid visual and
nervous system highlights molecular principles of convergent evolution. bioRxiv. 2022.05. 26.490366.
64. Tustison N.J., Avants B.B., Cook P.A., Zheng Y., Egan A., Yushkevich P.A., and Gee J.C. (2010).
N4ITK: improved N3 bias correction. IEEE Trans Med Imaging. 29, 1310-1320.
65. Gjerswold-Selleck S., Zhu N., Sun H., Sikka D., Shi J., Liu C., Nuriel T., Small S.A., and Guo J.
(2021). DL-BET-A deep learning based tool for automatic brain extraction from structural magnetic
resonance images in mice. International Society for Magnetic Resonance in Medicine.
66. Fedorov A., Beichel R., Kalpathy-Cramer J., Finet J., Fillion-Robin J.C., Pujol S., Bauer C.,
Jennings D., Fennessy F., Sonka M. et al. (2012). 3D Slicer as an image computing platform for the
Quantitative Imaging Network. Magn Reson Imaging. 30, 1323-1341.

67.   Avants B.B., Tustison N.J., Song G., Cook P.A., Klein A., and Gee J.C. (2011). A reproducible
evaluation of ANTs similarity metric performance in brain image registration. Neuroimage. 54, 2033-2044.
68.   Gestal C., Pascual S., Guerra Á., Fiorito G., and Vieites J.M. (2019). Handbook of Pathogens and
Diseases in Cephalopods, Springer).
69.   Schindelin J., Arganda-Carreras I., Frise E., Kaynig V., Longair M., Pietzsch T., Preibisch S.,
Rueden C., Saalfeld S., Schmid B. et al. (2012). Fiji: an open-source platform for biological-image
analysis. Nat Methods. 9, 676-682.

Brain region	Abbreviation	Function
Supraesophageal mass		
Vertical lobe complex		**Learning and memory (Young, 1991)**
Vertical lobe	V	
Subvertical lobe	SV	
Superior frontal lobe	SF	
Inferior frontal lobe	IF	
Posterior frontal lobe	PF	
Basal lobe complex		**Higher motor control (Boycott, 1961)**
Anterior anterior basal lobe	AAB	Movement of head, arms and eyes
Anterior posterior basal lobe	APB	
Precommissural lobe	PC	
Dorsal basal lobe	DB	
Interbasal lobes	IB	Movement of feeding tentacles
Median basal lobe	MB	Movement of mantle and funnel during swimming and breathing; protraction and retraction of head; movement of fins, movement of buccal mass, and expansion and contraction of chromatophores
Lateral basal lobes	LB	Control of chromatophores and papillae
Subesophageal mass		
Pedal lobe complex		**Intermediate and lower motor control of movement (Boycott, 1961)**
Anterior pedal lobe	AP	Movement of arms and tentacles
Posterior pedal lobe	PP	Movement of funnel, fins and tentacles; head retraction
Lateral pedal lobes	LP	Movement of eyes
Anterior dorsal chromatophore lobes	ADC	Control of chromatophores and papillae on the head and arms
Anterior ventral chromatophore lobes	AVC	Control of chromatophores and papillae on the head and arms
Magnocellular lobe complex		**Giant fiber response (breathing) (Boycott, 1961)**
Dorsal magnocellular lobes	DM	
Ventral magnocellular lobes	VM	
Posterior magnocellular lobes	PM	
Palliovisceral lobe complex		**Lower motor control of locomotion (Boycott, 1961)**
Palliovisceral lobe	PV	Control of escape movements and ink ejection
Lateral ventral palliovisceral lobes	LPV	
Fin lobes	F	Movement of fins
Posterior chromatophore lobes	PCL	Control of chromatophores and papillae on mantle, fin and visceral mass
Dorsal vasomotor lobe	DV	
Ventral vasomotor lobe	VV	
Brachial lobe complex		**Motor control of arms and feeding**
Brachial lobe	B	Intermediate motor control of arms
Superior buccal lobe	SB	Biting movements of the buccal mass
Inferior buccal lobe	IB	Biting movements of the buccal mass
Periesophageal mass		
Optic-tract complex		
Optic lobes	O	Visual processing (Boycott, 1961)
Peduncle lobes	P	Visuo-motor control (Messenger, 1967)
Dorsolateral lobes	DL	
Optic glands	OG	Neurosecretory center (Messenger, 1967)
Olfactory lobes	OL	Unclear (Messenger, 1979)
Nerve fibers		
Anterior magnocellular commissure	AMC	
Brachio-palliovisceral connectives	BPC	
Collar nerves	C	Innervate the collar muscles and valves of the mantle (Young, 1971)
Head retractor nerves	HR	Innervate the muscles that connect the head to the mantle (Young, 1971)

Lateral basal to posterior chromatophore lobe tracts	LBPC	
Optic to anterior basal lobe tracts	OAB	
Optic to dorsal magnocellular lobe tracts	ODM	
Optic to vertical lobe tracts	OV	
Pallial nerves	PN	Innervate the muscles and skin of the mantle and visceral mass (Young, 1971)
Subvertical to optic tracts	SOT	
Ventral optic commissure	VOC	
Visceral nerves	VS	Innervate retractor muscles of the head and funnel, ink sac, heart, main blood vessels and digestive, excretory and reproductive systems (Young 1971)

Table 1. The lobes of the cuttlefish brain.