Observing non-covalent interactions in experimental electron

density for macromolecular systems: A novel perspective for

protein–ligand interaction research

Kang Ding[†], Shiqiu Yin[†], Zhongwei Li[†], Shiju Jiang, Yang Yang, Wenbiao Zhou[*], Yingsheng

Zhang, and Bo Huang[*]

Beijing StoneWise Technology Co Ltd., Haidian street #15, Haidian District, Beijing 100080, China

[†]Equal contributors

[*]Corresponding authors:

huangbo@stonewise.cn, zhouwenbiao@stonewise.cn

ABSTRACT

We report for the first time the use of experimental electron density (ED) in the Protein Data Bank

for modeling non-covalent interactions (NCIs) for protein–ligand complexes. Our methodology is

based on the reduced electron density gradient (RDG) theory describing intermolecular NCI by

ED and its first derivative. We established a database called the Experimental NCI Database

(ExptNCI; http://ncidatabase.stonewise.cn/#/nci) containing ED saddle points, indicating

~200,000 NCIs from over 12,000 protein–ligand complexes. We also demonstrated the use of the

database for depicting amide–π interactions in a protein–ligand binding system. In summary, the

database provides details on experimentally observed NCIs for protein–ligand complexes and can

support future studies, including studies on rarely documented NCIs and the development of

artificial intelligent models for protein–ligand binding prediction.

INTRODUCTION

Non-covalent interactions (NCIs) govern protein–ligand interactions and are critical for

understanding the determinants affecting ligand-binding affinity. To achieve a deep understanding

of NCIs, many protein–ligand interaction databases have been established in the last decade.[1-7]

Two types of technologies are primarily applied to build such databases: (i) structure-based data

mining and (ii) quantum mechanical (QM) methods-powered computation. For the first type,

protein–ligand complex structures in the Protein Data Bank (PDB) are used as the main source,

and different indices, such as distance, angle, exposed surface, and line-of-sight statistics, are used

to depict the possibility of NCIs between a pair or two groups of atoms.[8-10] For the second type,

different levels of QM methods, ranging from semiempirical to coupled-cluster singles-doubles-

and-triples wave function (CCSD(T)), are used to quantify the interaction energy of small model

complexes.[5, 11] The two technologies together have contributed greatly to the development of rules

for the recognition of classical NCIs, such as hydrogen bonds, halogen bonds, salt bridges, and π-

π stacking. To further expand the ability to recognize and quantify the entire spectrum of NCIs in

highly complicated polarization environments such as protein–ligand binding systems and

protein–protein interaction systems, we need to address the gap in the direct evidence of NCI

between two proximal atoms in macromolecule systems. The gap is caused by the limitation of

applying quantum mechanics for large systems and by the uncertainty of atom positions in the

structures in PDB: e.g., the absence of hydrogen atoms and errors induced during structure building.

A potential solution for this gap can be found in the field of materials research[12] in studies

applying the reduced electron density gradient (RDG) theory[13] in analyzing experimental electron

density (ED) derived from the X-ray diffraction of small molecular crystals.[14, 15] Stating the RDG

theory in simple terms, NCI can be observed by pinpointing the ED saddle point, i.e. (3,-1) critical

points and further quantified by measuring ED deviation from a homogeneous electron distribution

using density and its first derivative ($s = [1/(2(3\pi^2)^{1/3})]|\nabla\rho|/\rho^{4/3}$). Some researchers have even

proved that experimental ED can contribute to optimizing functions for density functional theory

(DFT), given the fact that experimental ED is inherently time-averaged, whereas DFT ED

represents pure ground-state.[12]

Inspired by research on small molecule crystals,[12, 14, 15] we have developed a potentially path-

breaking procedure to extract critical points from experimental ED for protein–ligand complexes

deposited in the PDB. We processed >12,000 protein–ligand complexes and extracted ~200,000

saddle points. These data were subjected to noise reduction by varying the ED resolution and then

consolidated into a database called the ExptNCI (Experimental NCI Database), available at

http://ncidatabase.stonewise.cn/#/nci. In addition to database construction, we also present a case

of using such data for empirical NCI mining. ED saddle points indicating amide–π interactions are

extracted and used to support the QM interaction energy landscape scan. The QM result is well

aligned with the observed points: 85% of the observed points are covered by the region with energy

lower than -1.44 kcal/mol (semiempirical level). In addition to the attractive interaction of NH/π,

which is consistent with previous research,[16] we also found a -2.65 kcal/mol interaction (DFT level)

between the edge of the aromatic ring and the amide plane when they interact in a perpendicular

"edge-on" geometry.

RESULTS

NCI observed in experimental ED of the protein–ligand complex

X-ray diffraction (XRD) detects the electron distribution of the target molecule and generates an

ED map. By searching for the ED saddle points, we can not only recognize classical NCIs, such

as hydrogen bonds, π stacking, and halogen bonds, but also find relatively rare NCIs such as

fluorine interacting with sulfur and methyl interacting with pyridine, as shown in Figure 1a–e.

Additionally, because experimental ED represents a time-averaged density, some dynamics of the

NCIs can also be observed (Figure 1f and Figure 2).

Figure 1. Observing NCI in X-ray diffraction-derived electron density maps (2Fo–Fc). Blue mesh

indicates 2.5-Å resolution. All the maps are sigma-scaled and presented at specified counter levels.

Saddle points are indicated by red arrows. a) Hydrogen bond interactions (PDB: 1S38, map counter

level 0.2 sigma); b) Interaction between methyl and aromatic ring (PDB: 1Q8T, map counter level

0.2 sigma); c) interaction between F and methylthio (PDB: 2P4Y, map counter level 0 sigma); d)

Weak π stacking revealed in low-resolution electron density map (PDB: 3LDQ; blue mesh

indicates a 2.5-Å resolution map countered at 1.0 sigma; sand yellow mesh indicates a 3.5-Å

resolution map countered at 1.0 sigma); e) Sulfur-involved NCI (PDB: 4I1R, map counter level

0.3 sigma); f) Observing NCIs under dynamic context caused by the rotation of threonine side

chain (PDB:1XKK, map counter level 0 sigma).

Another benefit of using XRD ED for NCI detection is that we can clearly observe the signals

of weak NCI; this can be done by checking them in low-resolution ED maps generated by only

including XRDs at low-resolution. The intensity of XRD decreases as the ED map resolution

increases, resulting in a relatively high signal-to-noise ratio for low-resolution ED maps, which

enables us to confirm weak NCIs by checking them in ED maps at different resolutions (Figure 2).

Doing so enables the identification of weak NCIs and helps distinguish them from false-positive

signals. Our research conducted such a check in a low-resolution (3.5 Å) map for every NCI

associated with saddle points having sigma-scaled intensity at 2.5 Å less than 0, i.e., less than the

average.

Figure 2. Emphasizing NCI signal in low-resolution ED maps. All the maps are sigma-scaled and presented at counter level 0 sigma. Red dashed circles indicate the relatively weak NCIs, which are clearly observed in low-resolution ED maps. Hydrogen bonds in a dynamic environment are shown in panels a, b, and c, with 2Fo-Fc maps for PDB 1A28 at 2.5-Å, 3.0-Å, and 3.5-Å resolution, respectively. Red arrows indicate the rotation of the groups causing the dynamics. π stacking contacts are shown in panels d, e, and f, with 2Fo-Fc maps for PDB 1Z6E at 1.8-Å, 2.5-Å, and 3.0-Å resolution, respectively.

In addition to using saddle points as a general indicator for recognizing NCI, we also used RDG in experimental ED as a more comprehensive NCI descriptor. Both repulsive and attractive

interactions can be identified and visualized, as shown in Figure 3. Specifically, a spike in the

RDG vs. sign(λ_2)ρ plot indicates the presence of NCIs (Figure 3b), with the location of the spike

on the negative side of the horizontal axis indicating an attractive interaction and that on the

positive side of the axis indicating a repulsive interaction.[13]

Figure 3. Depicting NCI with RDG in experimental ED for protein–ligand complex (PDB:

2WNC). a) Saddle points detected in 2fo–fc map (counter level 1.0 sigma). Three saddle points

indicating three hydrogen bonds are respectively indicated by yellow, purple, and cyan arrows; b)

plots of RDG versus electron density multiplied by the sign of the second Hessian eigenvalue for

NCIs indicated in the panel (a) because the ρ here is sigma-scaled, to avoid negative value, all the

ρ values used for calculating RDG and sign(λ_2)ρ have their value added by 3. Spikes indicating

three hydrogen bonds are indicated by arrows. All the dots on the scatter plot are colored according

to their positions in real space. In detail, the dots within 1 Å of the saddle points 1, 2, and 3 are

colored in yellow, purple, and cyan, respectively; c) RDG-based NCI isosurface showing the

ligand–pocket interaction. Regions inside the RDG isosurface at a value of 0.2 (arbitrary unit) are

indicated with dots, and the dots are colored based on sign(λ_2)ρ using the rainbow scheme, in

which blue depicts large negative values indicating strong, attractive interactions, and red depicts

large positive values indicating repulsive interactions.

However, one limitation of using experimental ED for RDG analysis needs to be mentioned.

Because of the lack of experimental measures on the forward-scattered reflection swamped by the

transmitted beam, which is known as F_{000}, the absolute value of ED is not available for

macromolecule crystals. Therefore, the ED maps are contoured on a relative scale, and we had to

use a sigma-scaled ρ for calculating RDG and $\text{sign}(\lambda_2)\rho$. Consequently, the plot in Figure 3b has

scales on the horizontal and vertical axes in arbitrary units. However, the spikes appearing in low-

density regions still can indicate the occurrence of NCIs.

ExptNCI database content

The current version of ExptNCI contains a total of 215,397 saddle points extracted from the

experimental ED of 12,589 ligand–pocket complex structures in the PDB. The ED maps used for

saddle points extraction have resolutions ranging from 2.5 to 4.5 Å, and 83% of them have a

resolution greater than 2.5 Å. The ED topology information of the saddle points (such as sigma-

scaled ρ, RDG, λ_1, λ_2, λ_3, and Laplacian) as well as the structural information of atoms at both

ends of the saddle points (such as residual name, element, and its hybridization in the Mol2/Sybyl

atom format)[17] are included in the database (Table 1). We also included ρ at a low-resolution (3.5

Å) at the position of saddle points in a 2.5-Å ED map to use it to distinguish noise from signals of

weak NCIs. As discussed in the first part of the results section, blurring the map by only including

low-resolution data with a relatively high signal-to-noise ratio can present weak NCIs more clearly.

Here, we filtered out false-positive saddle points in a 2.5-Å resolution ED map to check if such

points have negative sigma ρ in a 3.5-Å resolution ED map.

Table 1. List of fields in ExptNCI database

Field	Description
NCI ID	NCI ID in ExptNCI Database
Complex ID	Complex ID composed by PDBID and ligand name
PDB Code	PDB Code
L_type	Ligand atom type in Mol2/Sybyl format
R_type	Receptor atom type in Mol2/Sybyl format
NCI_atom_pair	NCI classified based on Mol2/Sybyl formatted type of atoms at both ends of the saddle point
NCI_intuitive	NCI classified based on the properties of the atoms at both ends of the saddle points
NCI_ODDT	ODDT based NCI type (hydrogen bond, salt bridge, halogen bond, pi stacking, pi cation)
Resolution	The high-resolution limit of electron density map available in PDB
CP_type	Type of critical point: (3,-1) indicates saddle points while (3,+1) indicates ring CPs
RDG	$[1/(2(3\pi^2)^{1/3})]|\nabla\rho|/\rho^{4/3}$, where ρ indicates the modified intensity of electron density. RDG is used as an indicator for NCI.
slr	$sign(\lambda_2)\rho$, where ρ indicates the modified intensity of electron density at the critical point; λ_2 indicates the second larger value of the three eigenvalues of the electron density Hessian (second derivative) matrix. The value of slr is ρ when λ_2 is positive, indicating repulsive interaction; it is $-\rho$ when λ_2 is negative, indicating attractive interaction.
λ_1	Three eigenvalues of the electron density Hessian (second derivative) matrix, such that ($\lambda_1 <= \lambda_2 <= \lambda_3$)
λ_2	Three eigenvalues of the electron density Hessian (second derivative) matrix, such that ($\lambda_1 <= \lambda_2 <= \lambda_3$), a positive λ_2 indicates repulsive interaction, while a negative λ_2 indicates attractive interaction
λ_3	Three eigenvalues of the electron density Hessian (second derivative) matrix, such that ($\lambda_1 <= \lambda_2 <= \lambda_3$)
Laplacian	$\nabla^2\rho = \lambda_1 + \lambda_2 + \lambda_3$
ED_2.5A_modified	$\rho+3$, where r is sigma-scaled ED intensity of the saddle point in 2Fo-Fc map at 2.5 Å. Such modified ρ is used to calculate $sign(\lambda_2)$ ρ and RDG.
ED_2.5A	Sigma-scaled ED intensity of the saddle point in 2Fo-Fc map at 2.5 Å

ED_3.5A	Sigma-scaled ED intensity of the saddle point in 2Fo-Fc map at 3.5 Å
Distance	Distance between ligand atom and receptor atom
is_backbone	Boolean: 1 if NCI occurs on protein backbone
is_ED_based	Boolean: 1 if NCI is defined based on an ED saddle point
is_ODDT	Boolean: 1 if NCI is recognized by ODDT
ResName	Name of the receptor residue involved in the NCI
ResAtomName	Name of the receptor atom involved in the NCI
ResID	Residue ID of the receptor residue involved in the NCI
ChainID	Chain ID of the receptor residue involved in the NCI
LigName	Name of the ligand involved in the NCI
LigAtomName	Name of the ligand atom involved in the NCI
Lig_ResID	Residue ID of the ligand residue involved in the NCI
Lig_ChainID	Chain ID of the ligand residue involved in the NCI
Group	Type of atom in terms of protein, water, or heteroatoms

By including strong saddle points with sigma-scaled intensity at 2.5-Å resolution above 0 and

weak saddle points which can pass the false-positive test by using the low-resolution map of 3.5

Å (the method described above), we selected 95,532 saddle points, accounting for 51% of the

originally labeled points (Figure 4a). Among them, 32% were also recognized as NCIs by rules

embedded in the widely used software ODDT[18], with hydrogen bonds accounting for the majority

(Figure 4b). For the 68% that were not recognized by ODDT, we made a rough classification based

on the properties of the atoms at both ends of the saddle points, as shown in Figure 1c, in which

polar interactions (hydrophilic–hydrophilic), aliphatic C…hydrophilic (N/O) interactions, and

aromatic …hydrophilic (N/O) interactions accounted for the majority.

Figure 4. Database construction and dataset profile. a) Database construction workflow. *AND* and

OR in the select box are logical operators; b) distribution of interaction type for NCIs recognized

by both ODDT and ED saddle points; c) distribution of interaction type for NCIs recognized by

ED saddle points but not ODDT.

Usage Case: Depicting amide–π interactions in the ligand–protein binding system

Amide–π interactions[16] have been increasingly studied for their involvement in the binding of

drug molecules to target proteins.[19-21] Most previous studies focused on how the plane of the arene

ring interacts with the amide,[16, 20-22] which can be classified as focusing on face-on geometry, a

configuration with γ approximately 0° in a coordinate system, as shown in Figure 5a. To check

whether such face-on geometry represents the majority of amide–π interactions in the protein–

ligand binding system, we extracted 3,162 amide-π pairs from the ExptNCI database (details of

the list provided in supplementary information). The amide–π pairs were extracted based on the

fulfillment of the following requirements: (i) it must have ED saddle points between the aromatic

carbon and any atom of the amide group and (ii) the ED map must have a resolution better than

2.5 Å (examples shown in Figure 5b). Those with saddle points between C=O and hetero atoms in

the aromatic ring were excluded so that classical hydrogen bonds are not included in the analysis.

The spatial distribution of the aromatic ring center relative to the carbon atom of the amide plane

was plotted and colored with a γ-related color scheme. Interestingly, most of the interactions

displayed γ values of approximately 90°, indicating an edge-on geometry (Figure 5c). Notably,

face-on and edge-on interactions occur on two ellipsoids with different radii (Figure 5d and 5e).

Figure 5. Using experimental ED data to support the profiling of amide–π interaction. a) Examples

of amide–π interaction identified by ED saddle points. 2fo–fc map is countered at 0.3 sigma; b)

coordinate system of amide–π interaction; c) spatial distribution of aromatic ring center relative to

the carbon atom of the amide plane. Green and yellow indicate edge-on and face-on geometry,

respectively. Illustrations of edge-on and face-on are also provided; (d) and (e) are ellipsoids and

parameters obtained by fitting edge-on and face-on positions, respectively, to the general equation

of an ellipsoid; for (f), (g), and (h), the top part represents the sampling scheme of formamide–

benzene conformation on the ellipsoids, with edge-on conformation sampled in two ways:

perpendicular and parallel; the middle part represents GFN2-xTB level energy landscape, with the

blue arrow pointing to a red star indicating global minimum; bottom part represents the

conformation for global minimum on GFN2-xTB energy landscape and its M06-2x/6-311+G(d,p)

energy calculated by GAMESS; i) energy landscape scan for N- acetyl glycyl glycinamide. The

ellipsoid is with respect to the amide group indicated by the red circle.

To further investigate the interaction geometry for amide–π, we identified the ellipsoids for face-

on and edge-on geometry by fitting the aromatic center positions of the two types of geometry to

the general equation of an ellipsoid (Figure 5d and 5e). Subsequently, we computed the GFN2-

xTB level energy landscape based on the fitted ellipsoids using a formamide–benzene model

system (Figure 5f, 5g, and 5h). For edge-on geometry (i.e., γ=90), the interaction is favored when

benzene approaches the amide plane from the top of C=O perpendicularly (Figure 5f and 5g), with

a minimum interaction energy of -2.65 kcal/mol calculated using M06-2x/6-311+G(d,p). For face-

on geometry (i.e., γ=0), the result of our energy landscape scan is consistent with previous

studies,[16] showing a favored interaction of NH/π and a repulsive interaction of C=O/π, as shown

in Figure 5h. The same approach was also applied to the amide group in a tripeptide to simulate

the situation in the protein (Figure 5i). The computed energy landscape enjoyed a decent match to

the spatial distribution of the observed amide–π interactions extracted from ExptNCI, with 85%

of the latter covered by the former region with energy lower than -1.44 kcal/mol.

In summary, the use of observed ED saddle points for NCI description is demonstrated in this

case through its support for an energy landscape scan.

DISCUSSION

XRD provides an experimental ED map that contains massive amounts of information. Partial

information is effectively interpreted into atom coordinates, and this information is entered in the

PDB. However, in addition to atom coordinates, there is still plenty of information hidden in the

experimental ED maps. For the first time, we extracted NCI signals from the ED maps and used

them to establish the ExptNCI database.

How does the ED-saddle-point-based observation complement and further improve geometry-

rule-based description? In most cases, the rule-based NCI descriptions have the following two

characteristics: 1) they focus on a pair of atoms or groups by simplifying the environment and 2)

they highly rely on the precision of atomic coordinates. Therefore, such description cannot always

appropriately profile the NCIs in practical situations where the polarization environment is

compilated and the model structures are inaccurate or have missing regions. Because ED-saddle-

point-based NCI observation mainly depends on electron density, it is less sensitive to the accuracy

of the coordinates than rule-based descriptions. In addition, because the saddle points are detected

from the experimental electron density, they can reflect the complex polarization environment and

provide more information such as the relative strength of the NCIs and geometrically rare cases to

support the development of new empirical rules. These superiorities are shown in two cases (Fig.

S1 and S2), where the saddle-point-based method is compared with that of the rules-based method

for sulfur-involved NCIs and π-π stacking.[18, 23, 24]

When exploring the ExptNCI database, users should check the three following aspects if some

seemingly unusual NCIs are found: (i) check whether the structure is correctly determined, which

can be judged by checking positive or negative densities around the NCI region of interest in the

Fo–Fc map; (ii) check whether low-resolution causes merging of saddle points. An ED map

becomes less detailed when the resolution is low, and two proximal saddle points may merge into

one in a low-resolution ED map. As shown in Figure 6, just because there is only one saddle point

between C=O and C=O in a 2.7-Å resolution ED map, it does not necessarily indicate the existence

of NCIs between the two sp^2 oxygen atoms. In other words, the case in Figure 6 resulted from the

merging of two saddle points representing two individual classical hydrogen bonds; (iii) check

whether there are any dynamics that can make the interaction more reasonable, e.g., the flip of the

side chain for Gln, Asn, and His.

Figure 6. Merging saddle points in low-resolution ED maps (PDB: 6MA1). a) Experimental ED

2Fo–Fc map at 2.7-Å resolution shown at the counter level of 1.4 sigma. Two classical hydrogen

bonds, indicated by orange dash lines, exist within the red dashed circle, but only one saddle point

is observed; b) GFN2-xTB calculated electron density, showing two saddle points at the same

region. The map is countered at 0.03 e$^-$/Å^3. The empirical QM calculation is conducted using xtb[25].

To further improve the data in terms of quantity and quality, we consider two directions. The

first is to expand the scale of the database by extracting NCIs from the interface of protein–protein

interactions (PPI). This may allow us to achieve a more detailed understanding of the interaction

fingerprint and ultimately benefit peptide/protein design. The second direction is to improve the

accuracy of the data by solving multi-crystal variance, which is a problem caused by the lack of

absolute ED values for macromolecule crystals. Such a challenge could be tackled by converting

the ED values from the sigma-scaled density to the number of electrons. Previous studies that were

aimed at measuring the quality of structures in PDB by analyzing ED can serve as a good starting

point.[26, 27]

Including the experimental saddle point ED intensity as NCI information can also be considered

as a solution to support artificial intelligence-based protein–ligand binding prediction. Although

experimental NCI is not always available as input, because most often, pocket–ligand complexes

are generated by docking or molecular dynamics and thus lack experimental ED, we can build two

machine learning models to first predict NCI from a given protein–ligand complex structure, and

then use the predicted NCI to facilitate ligand binding affinity prediction.

In addition to providing more data resources, describing NCI from the perspective of

crystallography ED also inspired us to consider leveraging crystallography as a solution for

molecular representation for machine learning models. To date, the majority of attempts by

researchers to find molecular representations have been in real space, and many reports have been

made using strings, molecular graphs, molecular matrixes, potential fields, and atom density

fields.[28] However, an ideal representation comprehensively reflecting physical and chemical

information, friendly to mathematics, and supported with plenty of experimental data available for

AI model training is still absent. By applying crystallography theory, we can further expand the

attempt in reciprocal space (i.e., frequency domain) and take a big step forward to realizing the

ideal representation for molecules. Specifically, we apply Fourier transform (FT) on the atomic

coordinates to transfer the information from real space to the frequency domain and then apply

reverse FT on the frequency domain to bring back the information to real space as ED. By varying

the resolution when conducting reverse FT in the frequency domain, we can obtain ED in real

space with different levels of detail, emphasizing scaffold, atom, or even bond properties. Unlike

graphs composed of vertices and edges, such representations fill the space in a continuously

differentiable manner, which is favored by the CNN model. Unlike other 3D molecular

representations, such representations are naturally associated with a large amount of testing data:

the experimental ED deposited in the PDB. We have already tested them on a 3D molecule

generation model and have obtained some promising results that will be reported later.

In summary, a massive amount of information is present in the experimental ED maps deposited

in the PDB. The usage of only part of that information has created our current understanding of

protein structures. We hope that our work can shed some light on leveraging experimental ED

maps to further understand NCIs in the macromolecular system and on combining crystallography

and AI from the perspective of providing reliable data sources and exploring better representation

of molecules.

METHODS

Database construction

Experimental ED map processing and critical point labeling

All coordinates and map coefficients were obtained from PDB-REDO.[29] ED maps covering

ligands and pocket residues within 5 Å of the ligands were synthesized at multiple resolutions

using Phenix.[30] The maps were stored in the xplor format with a 0.15-Å grid interval. The critical

points were labeled using the following procedure:

The ligand in our database is defined using PDBbind (version 2019) as a benchmark.[31]

Ligand/receptor atom pairs with a distance <5 Å were identified, and the midpoint was set as the

origin;

The RDG value of all the grids was calculated within 1 Å of the origin;

The gird point with local minimum RDG was found and marked as a saddle point candidate;

For all the saddle point candidates, the eigenvalue of the Hessian matrix was calculated and

sorted so that $\lambda_3 > \lambda_2 > \lambda_1$. If the eigenvalues did not fulfill the criteria of $\lambda_3 > 0 > \lambda_2 > \lambda_1$, the

candidate was discarded;

If there were two saddle point candidates <0.5 Å from each other, the one with the relatively

weaker intensity was discarded.

Atom property annotation

The topology of ligands from PDB entries was curated by RDKit with isosteric SMILES from

RCSB Ligand-Expo, and other ligands with missing data were curated using OpenBabel. The

Mol2/Sybyl atom types of pockets and ligands in the database were annotated using OpenBabel

and PyBel packages, and the rule-based molecular interactions in the database were analyzed and

classified using the ODDT software package (version 0.7).[18]

Web interface implementation

The database website was developed with a Java backend. The ligand similarity search or

substructure search in the database was developed using RDKit, and NCI information was stored

and queried through MySQL. NGL.js was implemented to display the receptor–ligand complex

and the ED map.

Amide–π interaction model

To avoid including O..N.ar hydrogen bonds, only amide–π systems with ED saddle points

between C/N/O on the protein backbones and C.ar on ligands were subject to our analysis. To

profile the spatial distribution of aromatic ring centers, all the amide groups of interest were

superimposed and placed on the X—Y plane with a uniform orientation (Figure 5b), and all the

aromatic centers of the amide–π systems were plotted in the Z-positive sector, given that the amide

plane is a mirror plane.

Four parameters including angles α, β, γ, and distance d are defined as shown in Figure 5b to

describe amide–π geometry, in which the angle α is used to describe whether the π system is

parallel (α=0 °±30° or 180 °±30°) or perpendicular (α=90±30°) to the amide plane; the angle γ is

used to describe whether the aromatic ring center is facing toward the amide group in a "face-on"

geometry (γ=0±30°), or showing its edge toward the amide group in an "edge-on" geometry

(γ=90±30°).

Ellipsoids for face-on and edge-on geometry were identified by fitting the aromatic center

position for the two types of geometry to the general equation of an ellipsoid (Figure 5d, 5e). Then

the fitted ellipsoids are represented by grids with an interval of 0.1 Å along both X and Y axes. To

scan the interaction energy landscape based on fitted ellipsoids for benzene and formamide

systems, we first determined the zero-point energy (-29.70 kcal/mol) by applying GFN2-xTB

calculation on a benzene–formamide complex with a distance of 50 Å between the two groups.

Afterward, we placed benzene on the face-on grid in the pose where γ equals 0° to obtain a face-

on geometry complex subset (Figure 5h). For the edge-on geometry complex subset, when we

placed a benzene group on the grid of edge-on ellipsoids, there were two types of poses that

fulfilled the requirement of γ=90°. Therefore, we divided the edge-on geometry into perpendicular-

edge-on and parallel-edge-on subtypes. In detail, we used a plane defined by the Z-axis and the

vector connecting the amide carbon to the benzene center to distinguish the two subtypes: if the

norm of benzene was in the above-defined plane, then it was a parallel-edge-on sub-type (Figure

5f); if the norm of benzene was perpendicular to the above-defined plane, then it was a

perpendicular-edge-on sub-type (Figure 5g). The energy landscapes for the geometries of face-on,

parallel-edge-on, and perpendicular-edge-on were synthesized by calculating GFN2-xTB energy

and then subtracting the zero-point energy from it for the complexes on the corresponding grids.

Complexes representing the global minimum of the three energy landscapes were also subjected

to the M06-2x/6-311+G(d,p) calculation using GAMESS to obtain the DFT level energy. The

energy landscape scan for benzene interacting with amide groups in the context of tripeptide was

conducted in a similar way using a GFN2-xTB-optimized N-acetyl glycyl glycinamide as a starting

point. Because the optimized molecule is not subject to mirror symmetry, we scanned the entire

ellipsoid and combined the upper and lower halves by overlapping the grids of the two parts and

using the lower energy on the two overlapped grids as the final value to compose the energy

landscape.

Software for figures and tables

The structure and ED figures were made using Pymol.[32] Statistical analysis was performed using

Pandas[33] and Numpy packages.[34] Scatter plots were constructed using Matplotlib[35] and Inkscape.[36]

AUTHOR INFORMATION

Corresponding Author

Bo Huang (huangbo@stonewise.cn)

Wenbiao Zhou (zhouwenbiao@stonewise.cn)

ORCID:

Bo Huang: 0000-0003-3822-9110

Present Addresses

StoneWise Ltd., 1708, Block B, No. 19 Zhongguancun Street, Haidian District, Beijing, China

Author Contributions

B. H. and Y. Z. conceived the idea. B.H. provided instructions for all experiments. W. Z. provided instructions on AI models. K. D. constructed the database. S. Y. developed the saddle point labeling script. Z. L. supported the saddle point labeling and performed quantum mechanics calculations. S. J. implemented the web interface for the database. Y. Y. designed the web interface.

Funding Sources

All the authors received funding from StoneWise Ltd. B.H. also received funding for cloud computing from Beijing Municipal Science & Technology Commission project Z211100003521001.

Data availability

http://ncidatabase.stonewise.cn/#/nci

Code availability

Available upon request.

Competing Interests

The authors declare no competing interests.

ACKNOWLEDGMENT

ABBREVIATIONS

ED, electron density; NCI, non-covalent interaction; PDB, Protein Data Bank; RDG, reduced

electron density gradient; QM, quantum mechanical; DFT, density functional theory; XRD, X-ray

diffraction; PPI, protein–protein interaction; FT, Fourier transform

REFERENCES

1. Anand, P.; Nagarajan, D.; Mukherjee, S.; Chandra, N., PLIC: Protein-Ligand Interaction

Clusters. *Database (Oxford)* **2014**, *2014*, bau029.

2. Angles, R.; Arenas-Salinas, M.; García, R.; Reyes-Suarez, J. A.; Pohl, E., GSP4PDB: A Web

Tool To Visualize, Search And Explore Protein-Ligand Structural Patterns. *BMC Bioinformatics*

2020, *21*, 85.

3. Gallina, A. M.; Bisignano, P.; Bergamino, M.; Bordo, D., PLI: A Web-Based Tool For The

Comparison of Protein-Ligand Interactions Observed on PDB Structures. *Bioinformatics* **2012**, *29*,

395-397.

4. Inhester, T.; Rarey, M., Protein–ligand Interaction Databases: Advanced Tools To Mine Activity

Data And Interactions On A Structural Level. **2014**, *4*, 562-575.

5. Jurecka, P.; Sponer, J.; Cerný, J.; Hobza, P., Benchmark Database Of Accurate (MP2 And

CCSD(T) Complete Basis Set Limit) Interaction Energies Of Small Model Complexes, DNA Base

Pairs, And Amino Acid Pairs. *Physical chemistry chemical physics : PCCP* **2006**, *8*, 1985-93.

6. Murakami, Y.; Omori, S.; Kinoshita, K., NLDB: A Database For 3D Protein-Ligand Interactions

In Enzymatic Reactions. *Journal of structural and functional genomics* **2016**, *17*, 101-110.

7. Rezac, J., Non-Covalent Interactions Atlas Benchmark Data Sets: Hydrogen Bonding. *J Chem*

Theory Comput **2020**, *16*, 2355-2368.

8. Ferreira de Freitas, R.; Schapira, M., A Systematic Analysis Of Atomic Protein-Ligand

Interactions In The PDB. *Medchemcomm* **2017**, *8*, 1970-1981.

9. Kuhn, B.; Gilberg, E.; Taylor, R.; Cole, J.; Korb, O., How Significant Are Unusual Protein-

Ligand Interactions? Insights from Database Mining. *J Med Chem* **2019**, *62*, 10441-10455.

10. Xu, Z.; Zhang, Q.; Shi, J.; Zhu, W., Underestimated Noncovalent Interactions in Protein Data

Bank. *J Chem Inf Model* **2019**, *59*, 3389-3399.

11. Hobza, P., Calculations On Non-Covalent Interactions And Databases Of Benchmark

Interaction Energies. *Acc Chem Res* **2012**, *45*, 663-72.

12. Kasai, H.; Tolborg, K.; Sist, M.; Zhang, J.; Hathwar, V. R.; Filso, M. O.; Cenedese, S.;

Sugimoto, K.; Overgaard, J.; Nishibori, E.; Iversen, B. B., X-Ray Electron Density Investigation

Of Chemical Bonding In Van Der Waals Materials. *Nat Mater* **2018**, *17*, 249-252.

13. Johnson, E. R.; Keinan, S.; Mori-Sanchez, P.; Contreras-Garcia, J.; Cohen, A. J.; Yang, W.,

Revealing Non-Covalent Interactions. *J Am Chem Soc* **2010**, *132*, 6498-506.

14. Saleh, G.; Gatti, C.; Lo Presti, L., Non-Covalent Interaction Via The Reduced Density Gradient:

Independent Atom Model Vs. Experimental Multipolar Electron Densities. *Computational and*

Theoretical Chemistry **2012**, *998*, 148–163.

15. Saleh, G.; Gatti, C.; Lo Presti, L.; Contreras-Garcia, J., Revealing Non-Covalent Interactions

In Molecular Crystals Through Their Experimental Electron Densities. *Chemistry* **2012**, *18*,

15523-36.

16. Imai, Y. N.; Inoue, Y.; Nakanishi, I.; Kitaura, K., Amide-Pi Interactions Between Formamide

And Benzene. *J Comput Chem* **2009**, *30*, 2267-76.

17. Clark, M.; Cramer III, R. D.; Van Opdenbosch, N., Validation Of The General Purpose Tripos

5.2 Force Field. **1989**, *10*, 982-1012.

18. Wojcikowski, M.; Zielenkiewicz, P.; Siedlecki, P., Open Drug Discovery Toolkit (ODDT): A

New Open-Source Player In The Drug Discovery Field. *J Cheminform* **2015**, *7*, 26.

19. Krone, M. W.; Travis, C. R.; Lee, G. Y.; Eckvahl, H. J.; Houk, K. N.; Waters, M. L., More Than

pi-pi-pi Stacking: Contribution of Amide-pi and CH-pi Interactions to Crotonyllysine Binding by

the AF9 YEATS Domain. *J Am Chem Soc* **2020**, *142*, 17048-17056.

20. DeFrees, K.; Kemp, M. T.; ElHilali-Pollard, X.; Zhang, X.; Mohamed, A.; Chen, Y.; Renslo,

A. R., An Empirical Study of Amide-Heteroarene pi-Stacking Interactions Using Reversible

Inhibitors of a Bacterial Serine Hydrolase. *Org Chem Front* **2019**, *6*, 1749-1756.

21. Bootsma, A. N.; Wheeler, S. E., Stacking Interactions of Heterocyclic Drug Fragments with

Protein Amide Backbones. *ChemMedChem* **2018**, *13*, 835-841.

22. Harder, M.; Kuhn, B.; Diederich, F., Efficient Stacking On Protein Amide Fragments.

ChemMedChem **2013**, *8*, 397-404.

23. Koebel, M. R.; Cooper, A.; Schmadeke, G.; Jeon, S.; Narayan, M.; Sirimulla, S., S...O and

S...N Sulfur Bonding Interactions in Protein-Ligand Complexes: Empirical Considerations and

Scoring Function. *J Chem Inf Model* **2016**, *56*, 2298-2309.

24. Beno, B. R.; Yeung, K. S.; Bartberger, M. D.; Pennington, L. D.; Meanwell, N. A., A Survey

of the Role of Noncovalent Sulfur Interactions in Drug Design. *J Med Chem* **2015**, *58*, 4383-438.

25. Bannwarth, C.; Ehlert, S.; Grimme, S., GFN2-xTB-An Accurate and Broadly Parametrized Self-Consistent Tight-Binding Quantum Chemical Method with Multipole Electrostatics and Density-Dependent Dispersion Contributions. *J Chem Theory Comput* **2019**, *15*, 1652-1671.

26. Lang, P. T.; Holton, J. M.; Fraser, J. S.; Alber, T., Protein Structural Ensembles Are Revealed By Redefining X-Ray Electron Density Noise. *Proc Natl Acad Sci U S A* **2014**, *111*, 237-42.

27. Yao, S.; Moseley, H. N. B., A Chemical Interpretation Of Protein Electron Density Maps In The Worldwide Protein Data Bank. *PLoS One* **2020**, *15*, e0236894.

28. Musil, F.; Grisafi, A.; Bartok, A. P.; Ortner, C.; Csanyi, G.; Ceriotti, M., Physics-Inspired Structural Representations for Molecules and Materials. *Chem Rev* **2021**, *121*, 9759-9815.

29. Joosten, R. P.; Long, F.; Murshudov, G. N.; Perrakis, A., The PDB_REDO Server For Macromolecular Structure Model Optimization. *IUCrJ* **2014**, *1*, 213-20.

30. Liebschner, D.; Afonine, P. V.; Baker, M. L.; Bunkoczi, G.; Chen, V. B.; Croll, T. I.; Hintze, B.; Hung, L. W.; Jain, S.; McCoy, A. J.; Moriarty, N. W.; Oeffner, R. D.; Poon, B. K.; Prisant, M. G.; Read, R. J.; Richardson, J. S.; Richardson, D. C.; Sammito, M. D.; Sobolev, O. V.; Stockwell, D. H.; Terwilliger, T. C.; Urzhumtsev, A. G.; Videau, L. L.; Williams, C. J.; Adams, P. D., Macromolecular Structure Determination Using X-Rays, Neutrons And Electrons: Recent Developments In Phenix. *Acta Crystallogr D Struct Biol* **2019**, *75*, 861-877.

31. Liu, Z.; Su, M.; Han, L.; Liu, J.; Yang, Q.; Li, Y.; Wang, R., Forging the Basis for Developing Protein-Ligand Interaction Scoring Functions. *Acc Chem Res* **2017**, *50*, 302-309.

32. Schrodinger, LLC, In; 2015.

33. McKinney, W., Data Structures for Statistical Computing in Python. *Proceedings of the 9th Python in Science Conference* **2010**, *445*, 51-56.

34. Harris, C. R.; Millman, K. J.; van der Walt, S. J.; Gommers, R.; Virtanen, P.; Cournapeau, D.;

Wieser, E.; Taylor, J.; Berg, S.; Smith, N. J.; Kern, R.; Picus, M.; Hoyer, S.; van Kerkwijk, M. H.;

Brett, M.; Haldane, A.; del Río, J. F.; Wiebe, M.; Peterson, P.; Gérard-Marchant, P.; Sheppard, K.;

Reddy, T.; Weckesser, W.; Abbasi, H.; Gohlke, C.; Oliphant, T. E., Array Programming With

NumPy. *Nature* **2020**, *585*, 357-362.

35. Hunter, J. D., Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering*

2007, *9*, 90-95.

36. InkscapeProject, Inkscape 0.92.5. **2020.**