The temporal and contextual stability of activity levels in hippocampal

CA1 cells

Yuichiro Hayashi, Ko Kobayakawa, and Reiko Kobayakawa

Institute of Biomedical Sciences, Kansai Medical University, 2-5-1 Shin-machi,

Hirakata, Osaka 573-1010, Japan

Correspondence: Yuichiro Hayashi

Institute of Biomedical Sciences, Kansai Medical University, 2-5-1 Shin-machi,

Hirakata, Osaka 573-1010, Japan

+81-72-804-2577

E-mail: hayasyui@hirakata.kmu.ac.jp

21 **Keywords**

22 Hippocampus, Place cell, Calcium imaging

Abstract

Recent long-term optical imaging studies have demonstrated that the activity levels of

hippocampal neurons in a familiar environment change on a daily to weekly basis.

However, it is unclear whether there is any time-invariant property in the cells' neural

representations. In this study, using miniature fluorescence microscopy, we measured

the neural activity of the mouse hippocampus in four different environments every 3

days. Although the activity level of hippocampal neurons fluctuated greatly in each

environment across days, we found a significant correlation between the activity levels

for different days, and the correlation was higher for averaged activity levels across

multiple environments. When the number of environments used for averaging was

increased, a higher activity correlation was observed. Furthermore, the number of

environments in which a cell showed activity was preserved. Cells that showed place

cell activity in many environments had greater spatial information content, and thus

carried a higher amount of information about the current position. In contrast, cells that

were active only in a small number of environments provided sparse representation for

the environment. These results suggest that each cell has not only an inherent activity

level but also play a characteristic role in the coding of space.

Significance Statement

Recent studies have revealed that place cell activity in the hippocampal CA1 cells exhibit instability on a daily to weekly scale. However, it is unclear whether there is any invariant property in the activity of the cells. In this study, we found that, although the activity level of CA1 neurons fluctuated greatly in one environment, the mean activity level across multiple environments was more stable. Furthermore, the number of environments in which a cell showed activity was preserved over time. These results suggest that even though the spatial code changes dynamically, each cell has an inherent activity level and plays a characteristic role in spatial coding.

Abbreviations

AAV, adeno-associated virus; GC, genome copy; ANOVA, analysis of variance; EWTL, electrowetting lens; PV, population vector

Introduction

The hippocampus is a brain region that plays a crucial role in episodic memory formation. To support this, the activity of hippocampal neurons encode place, environment, time, and various task-relevant information (O'Keefe and Dostrovsky, 1971; Muller and Kubie, 1987; Wood et al., 2000; Ferbinteanu and Shapiro, 2003; Pastalkova et al., 2008; MacDonald et al., 2011). In particular, place cell activity, i.e., the activity of hippocampal cells in restricted portions in an environment, has been extensively studied (O'Keefe and Nadel, 1978; Moser et al., 2015). The activity pattern of place cells can change or disappear when animals move to a different environment (O'Keefe and Conway, 1978; Muller and Kubie, 1987; Alme et al., 2014; Lu et al., 2015). This phenomenon is known as remapping, and it seems to provide a mechanism through which the hippocampus can represent different environments.

Recent studies have shown that the place fields and activity levels of hippocampal neurons for familiar environments are not constant; rather, they fluctuate on a daily to weekly basis (Mankin et al., 2012; Ziv et al., 2013; Rubin et al., 2015; Cai et al., 2016; Hainmueller and Bartos, 2018; Kinsky et al., 2018a; Gonzalez et al., 2019; Hayashi, 2019; Wirtshafter and Disterhoft, 2022). Such fluctuations are known as "representational drift" and have also been found in other brain regions, including the

parietal cortex, visual cortex, and olfactory cortex (Driscoll et al., 2017; Deitch et al.,

2021; Schoonover et al., 2021). Several studies, however, have provided evidence that

each cell has a preconfigured activity level and physiological role; for example, the

firing rates of the neurons in the hippocampus and entorhinal cortex are preserved

across different brain states and environments (Hirase et al., 2001; Mizuseki and

Buzsáki, 2013; Liberti et al., 2022). Recording in a very large environment (>40 m

long) revealed that the location of the place fields of the hippocampal neurons

fluctuated on a daily to weekly timescale, but that the number of place fields was

preserved over months (Lee et al., 2020). Another previous study demonstrated that

contextual fear memory engram cells, defined as neurons that excite and express c-fos at

memory encoding and whose reactivation results in memory recall, in the CA1 region

showed higher environmental specificity and lower spatial information content than

non-engram cells in the same area (Tanaka et al., 2018).

In an attempt to reconcile these seemingly incompatible observations on the

long-term dynamics of hippocampal activity, we measured neural activity in the mouse

hippocampal CA1 area in four different environments over several days. If each cell has

a preconfigured activity level and changes its tuning continually, the response to a small

set of stimuli (i.e., a small environment) should fluctuate day by day. Conversely, the

averaged response to a large variety of stimuli (i.e., a very large environment or multiple

environments) will be more stable than that for a single environment. We found that the

activity level of hippocampal neurons fluctuated greatly in one environment, and that

there was a significant correlation between the activity levels for different days. When

the number of environments used for averaging increased, a higher activity correlation

was observed. We also found that the number of environments in which the cells

showed activity tended to be conserved over days.

Materials and Methods

Animal subjects

All experiments complied with the applicable guidelines and regulations. The animal

care and use procedure followed protocols that have been approved by the Animal

Research Committee of Kansai Medical University. All experiments complied with the

ARRIVE guidelines

(https://arriveguidelines.org/arrive-guidelines/experimental-animals).

Nine adult male C57BL/6N mice (Japan SLC, Inc., Shizuoka, Japan) aged 12–18 weeks

were used in the experiments. The mice were housed under a standard 12-h light/dark

cycle and were given access to food and water *ad libitum*.

Viral constructs

We obtained an AAVrh10.hSyn.GCaMP6f.WPRE.SV40 virus from Vector Core at the

University of Pennsylvania at a titer of ~1.3×10^{14} GC/mL. The virus was diluted to ~5

$\times 10^{12}$ GC/mL with phosphate buffered saline before use.

Miniature head-mount microscopy

The miniature microscope design was based on the UCLA miniscope v.3

(miniscope.org). Excitation light was emitted by a blue LED (LXML-PB01-0030;

Lumileds, Aachen, Germany). The light passed through an excitation filter (ET470/40x;

Chroma Technology, Bellows Falls, VT) and was reflected by a dichroic mirror

(T495lpxr; Chroma Technology) onto the tissue through an electrowetting lens (EWTL)

(Varioptic A-25H0; Corning, Corning, NY) and an objective lens (tandem configuration

of achromatic doublets; #84-126 and #49-271, Edmund Optics, Burlington, NJ). The

fluorescent emissions collected by the lenses were passed through the dichroic mirror

and an emission filter (ET525/50m; Chroma Technology). The fluorescence image was

focused by a tube lens (#49-277; Edmund Optics) and captured by a CMOS image

sensor (MT9V032; ON Semiconductor, Phoenix, AZ). The microscope body was

created using a 3D printer (From2, resin type FGPBLK03; Formlabs, Somerville, MA).

The design files are available at https://github.com/yuichirohayashi/Zscope. The driving

voltage for the EWTL (square wave, 2 kHz) was generated using a data acquisition

board (PCIe-6259, National Instruments, Dallas, TX) controlled by custom software

written in LABVIEW 7.1 (National Instruments, Dallas, TX) and amplified with a linear

amplifier (As-904-150B, NF Corporation, Yokohama, Japan).

Surgery

Animals were anesthetized using isoflurane. The skull was exposed, and a small hole

(<0.5 mm) was made over the right hemisphere (1.5 mm lateral to the midline, 2.3 mm

posterior to the bregma). Then, 200 nL of GCaMP6f virus was injected into the CA1

region (1.2 mm ventral from the brain surface). One week after the viral injection, the

animals were anesthetized using isoflurane, and a 2.8-mm diameter craniotomy was

performed. The dura was removed, and the underlying cortex was aspirated. A

stainless-steel cannula (2.76 mm outer diameter, 2.40 mm inner diameter, 1.5 mm

height) covered by a glass coverslip (0.12-mm thick) was inserted over the dorsal CA1.

Dental cement (Shofu, Kyoto, Japan) was used to glue the cannula to the skull.

Four weeks after the cannula implantation, GCaMP-expressing neurons were

imaged using the head-mount microscope. The aluminum baseplate of the microscope

was cemented in a position at which the neurons were visible.

Recording arenas

Calcium activity was recorded in four arenas of different shapes and colors: a tall green

box ($37 \times 25 \times 56$ cm), a short gray box ($37 \times 25 \times 35$ cm), a lime round basket ($39 \times$

26×25 cm), and a white square basket ($36 \times 25 \times 25$ cm). One side of each arena was

painted a different color or decorated with plastic tape (Figure 1A).

Behavioral training

One week after the baseplate surgery, the mice were food-restricted and maintained at

above 85% of their initial weight. Mice were trained to forage for food pellets (14 mg,

#F05684, Bio-Serv, Flemington, NJ) scattered on the floor of the arenas.

Recording sessions

After at least 5 days of behavioral training, mice underwent four 5-min recording

sessions per day in the four different arenas in a random sequential order. The arenas

were cleaned with 70% ethanol prior to each recording session to eliminate any

olfactory cues. During the sessions, food pellets were thrown into the arena

approximately every 1 min. The excitation light intensity was approximately

$0.1–0.5$ mW/mm^2, and fluorescence images were captured at 30 Hz. The focus was

adjusted to match the previous image. Animal behavior was recorded at 30 Hz with an

overhead camera (DMK22BUC03, The Imaging Source, Bremen, Germany).

Image processing

ImageJ 1.52 (National Institutes of Health, MA) and MATLAB 2019a (Mathworks,

Natick, MA) software was used for all analyses. The calcium imaging data recorded on

the same day were concatenated and processed using MiniscopeAnalysis

(https://github.com/etterguillaume/MiniscopeAnalysis). First, the images were

downsampled spatially by factors of two and motion-corrected using NoRMCorre

(Pnevmatikakis and Giovannucci, 2017). The downsampled, motion-corrected videos

were then processed using CNMF-E (Zhou et al., 2018) to extract fluorescence traces

from individual neurons. The fluorescence traces were finally processed using OASIS

software (Pnevmatikakis et al., 2016) for denoizing and deconvolving to estimate the

spiking activity. Behavioral image data were binarized such that black pixels

represented the animal body. The position of the animal at each frame was determined

as the centroid of the black region.

Registration of cells across sessions

For day-to-day longitudinal cell registration, the spatial footprints of the cells extracted

with CNMF-E from each day's fluorescence image stream were processed with the

Cellreg routine (Sheintuch et al., 2017) using the default parameters (maximum distance

<12 μm, spatial correlation >0.74, and P_{same} threshold = 0.5). In brief, the sets of spatial

footprints of cells for each session were aligned with rigid-body translation. The

distribution of spatial footprint similarities between pairs of neighboring cells from

different sessions was modeled to obtain an estimation for the probability of them being

the same cell (P_{same}). Registration quality was assessed using the average P_{same} value for

registered cells in an animal and the register score for registered cells (Sheintuch et al.,

2017). The register score was calculated as the total number of reliably identified cell

pairs out of the total number of cell pairs across all sessions using the following

equation:

$$\text{register score} = \frac{1}{\widehat{N}(N-1)} \sum_{k=1}^{\widehat{N}} \sum_{\substack{m=1 \\ m \neq k}}^{N} \delta(k,m)$$

where N represents the total number of sessions, ~N represents the number of active

sessions for a given cell register. The $\delta(k,m)$ value was 1 if a cell-pair in the cluster was

reliable and 0 if the cell-pair was unreliable.

Place fields

To analyze the place fields, calcium events that occurred when the animal's velocity was

less than 1 cm/s were filtered out to eliminate nonspecific activity at rest. The place

field maps were constructed by finding the spike number at each spatial bin (2×2 cm)

divided by the dwell time in the bin. The maps were smoothed using a Gaussian

function with a standard deviation of 3 cm.

212   For each place field of the cells, the mutual information between calcium

transients and the mouse's location was calculated (Skaggs et al., 1993). To assess the

significance of spatial selectivity, Monte Carlo P values were calculated. A total of

1,000 distinct shuffles of the calcium transient times were performed, and the mutual

information for each shuffle was calculated. The P value was defined as the fraction of

shuffles that exceeded the mutual information of the cell. Cells with a p value of <0.05

were considered place cells.

Rate difference

The rate difference was determined by calculating the unsigned rate difference between

the mean calcium transient rates in the two recording sessions and then dividing the

difference by the sum of the two rates.

Population vector correlation

To compute the population vector (PV) correlation between two sessions, the place

fields of all simultaneously recorded place cells were stacked into a 3D matrix (place

field map × cell identity). At each spatial bin, the firing rate vector of all cells

represented the PV for that spatial bin. To compare two recording sessions, the

Pearson's correlation coefficient was calculated between each pair of PVs at the

corresponding locations, and the correlation coefficients of all spatial bins were

averaged (Leutgeb et al., 2005).

Experimental design and statistical analysis

Calcium activity was recorded from nine mice. Recordings from animals with low

activity (average arena coverage <80%) were discarded. Statistical analysis was

performed using GraphPad Prism 5 (GraphPad Software, La Jolla, CA), MATLAB

2019a, and R version 4.1.1. Data were first tested for normality using the Shapiro–Wilk

test and for homogeneity of variance with Bartlett's test or the *F*-test. Normally

distributed data with equal variances were compared with a one-way analysis of

variance (ANOVA) followed by Bonferroni's post hoc comparisons. If data were not

normally distributed, nonparametric tests (Mann–Whitney U test, Kruskal–Wallis test,

or Friedman test) were used. Statistical significance was set at $p < 0.05$ for all statistical

analyses.

Results

Experimental outline

To investigate the place cell activity for multiple spatial environments in the

hippocampus, neural activity was recorded successively in four arenas with different

shapes and colors (Figure 1A). Before commencing the recording sessions, mice (n = 6)

were trained to find food in these arenas for at least 5 days. Each mouse then underwent

four 5-min recording sessions per day at 3-day intervals (Figure 1A).

253   Neural activity in the dorsal CA1 region of the hippocampus was recorded

using a combination of a genetically encoded calcium indicator, GCaMP6f, and a

miniature microscope (Figure 1B, C). We extracted the calcium traces of putative

neurons from the fluorescence image stream using CNMF-E (Zhou et al., 2018). The

firing rate of each neuron was estimated from its calcium trace using OASIS software

(Pnevmatikakis et al., 2016) (Figure 1D). The firing rate distribution of the recorded

cells showed lognormal-like distribution (Figure 1E), consistent with the findings of a

previous electrophysiological study (Mizuseki and Buzsáki, 2013). With the spatial

footprints of the cells extracted with CNMF-E from each day's fluorescence image

stream, we used Cellreg (Sheintuch et al., 2017) to track cells across days.

Activity of CA1 cells showed a variety of environmental specificity

A total of 6,425 cells were recorded from six mice (602–1,580 cells per animal) in four

different environments. Following a previous study, we observed various cells ranging

from those that showed no activity in any environment to those that showed activity in

all four environments in one day (Figure 2A, B) (Lu et al., 2015). We termed the

number of environments where a cell exhibited activity above the threshold as the

"global activity level index" (gA index). We also classified cells according to the

number of environments where a cell showed statistically significant place cell activity.

This value was termed the "global place cell activity level index" (gPC index). As

shown in Figure 2C, CA1 cells displayed various gPC activity levels. However, a

fraction of cells that showed significant place cell activity did not change between

environments or over days, indicating that these mice were highly familiar with all the

environments (Figure 2D, $Q_{(3,69)} = 1.639$, $p = 0.65$, n = 24 recordings per day in six

animals. $Q_{(3,69)} = 5.812$, $p = 0.12$, n = 24 recordings for each environment in six animals,

Friedman test).

Long-term tracking of neural activity for each cell

For longitudinal analysis of the same cells, the cells need to be reliably registered across

days. Therefore, we checked the register score and the P_{same} value to ensure the

reliability of cell registration (Sheintuch et al., 2017). The average registration score

was 0.86 ± 0.020 (mean ± SEM, n = 6 animals, Figure S1A, left), and the average P_{same}

value was 0.83 ± 0.052 (mean ± SEM, n = 6 animals, Figure S1A, right). These values

were comparable to those in the previous study in the hippocampal CA1 area (Sheintuch

et al., 2017).

Next, we evaluated the reliability of cell registration at the functional level. For

this purpose, the PV correlation (Leutgeb et al., 2005) of the spatial representation for

the four environments between sessions was calculated. Consistent with previous

studies (Rubin et al., 2015; Kinsky et al., 2018b), the PV correlation between sessions in

the same environment was significantly higher than between sessions in different

environments for at least a 6-day session interval (Figure S1B, $F_{(5,174)} = 5.302$, $p <$

0.0001, n = 24 session pairs for same environment, n = 36 session pairs for different

environments, one-way ANOVA followed by Bonferroni's post hoc test). The results

also corroborated that the same neurons were reliably tracked across multiple sessions.

Not only across-day but also within-day registration quality affects the analysis

results. For wide-field imaging data, signals from neighboring cells overlap due to light

scattering and lack of optical sectioning capability (Wilt et al., 2009). Furthermore,

although CNMF-E has been shown to effectively demix signals from overlapping cells

(Zhou et al., 2018), a possible concern remains that the activities of multiple spatially

overlapping cells may be merged into the activity of a single high-gA cell. Therefore, to

test this possibility, we examined the relationship between a cell's gA index and its

spatial footprint size of the region of interest extracted using CNMF-E. If multiple

spatially overlapping cells are merged into a single cell, its spatial footprint is

hypothesized to be larger than that for a genuinely single cell. As shown in Figure S1C,

there was a slight negative correlation between the gA index and the spatial footprint

size. This result suggested that the demix error in the cellular signal extraction was not

more frequent for high-gA cells than for low-gA cells.

Activity levels averaged across multiple environments were more stable than those

for a single environment

Next, we analyzed the long-term dynamics of neural activity for each cell in the four

different environments. A significant correlation between the activity levels in an

environment for adjacent sessions (3 days apart) was observed (Figure 3A, "1 env", R =

0.66, n = 77,100 observations of cells). Notably, the correlation for mean activity level

across multiple environments increased gradually according to the increased number of

environments to be averaged. For calculating the firing rate for multiple environments,

to match the data length, we used 5-min data constructed by concatenating data

fragments randomly chosen from recordings in multiple environments (Figure 3A, "2

env", R = 0.69, n = 115,650 observations of cells, "3 env", R = 0.70, n = 77,100

observations of cells, "4 env", R = 0.71, n = 19,275 observations of cells; Figure 3B, $F_{(3,15)}$

= 24.70, $p < 0.0001$, n = 6 animals, one-way repeated measures ANOVA followed

by Tukey's post hoc test).

The gA and gPC indices were preserved over time

We then tested whether the cells' response properties to environments, such as the gA

and gPC indices, were also preserved. A significant correlation was observed for the gA

index between adjacent sessions (Figure 4A, n = 90 session pairs, R = 0.78, $p < 0.0001$).

Furthermore, although we used a threshold value of 0.01 Hz, a significant correlation

was still found with threshold levels ranging from 0.01 to 0.08 Hz (Figure S2). The gPC

index between adjacent sessions was also correlated (Figure 4B, n = 90 session pairs, R

= 0.72, $p < 0.0001$). These results indicated that not only the mean firing rate but also

the gA and gPC indices (gA/gPC indices) of each cell were preserved over time.

Although the gA/gPC indices were preserved over time, a significant fraction of the

cells changed their environmental preferences (Figure 4C, D).

337   Additionally, on the basis that each cell had preserved activity levels and

gA/gPC indices, we determined whether each cell also had an inherent role in spatial

coding. We investigated the relationship between the gA/gPC indices and the cells'

firing properties. Both the mean firing rates and spatial information rates were

positively correlated with the cell's gPC index (Figure 4E, mean firing rate: R = 0.22,

$F_{(3, 8514)} = 424$, $p = 5.3 \times 10^{-92}$, n = 8,516 observations of neurons over all recording

sessions; 6B, spatial information: R = 0.23, $F_{(3, 8514)} = 467$, $p = 6.9 \times 10^{-101}$, n = 8,516

observations of place cells over all recording sessions, *F*-test). These results indicated

that high gPC index cells convey much information about the current position. In

contrast, cells that were active in a small number of environments constituted a sparse

code for environmental identity. To confirm this, we calculated the firing rate

differences between the environments where a cell showed the highest firing rate and

the firing rate averaged over the other three environments. Because cells did not

necessarily possess position information when encoding the environment, we classified

the cells using the gA index. The firing rate difference was negatively correlated with

the gA index (Figure 4G, $R = 0.24$, $F_{(3,12310)} = 761$, $p = 1.6 \times 10^{-162}$, $n = 12{,}312$

observations of cells over all recording sessions, F-test). This result indicated that cells

with a small gA index show sparse representation for environment.

Discussion

In this study, using a head-mount miniature fluorescence microscope, we measured the

neural activity of hippocampal CA1 cells over time in four different environments

(Figure 1A). We found a significant correlation for each cell's activity level for a single

environment between sessions, and the correlation for mean activity level across

environments increased gradually according to the increased number of environments to

be averaged (Figure 3A, B). This result reconciles previous conflicting observations on

the long-term stability of the neuronal activity in the CA1: the place field location and

activity levels of CA1 cells are not constant but fluctuate on a daily to weekly basis (Ziv

et al., 2013; Rubin et al., 2015; Cai et al., 2016; Kinsky et al., 2018a, 2020; Gonzalez et

al., 2019; Hayashi, 2019). In contradiction to these reports, a recent study reported that

the activity levels of CA1 cells in a very large (~40 m long) environment were stable

over time, though the location of their place field fluctuated (Lee et al., 2020). If each

cell has a preconfigured activity level and changes its tuning continually, the response to

a small set of stimuli will fluctuate day by day. Conversely, the averaged response to a

large variety of stimuli will be stable. Animals receive a larger variety of sensory stimuli

from a large environment or multiple environments than from a single environment.

This is probably the reason why averaged activity across multiple environments is

stable over days (Figure 3A, B).

We also showed that the gA and gPC indices tended to be preserved over time

(Figure 4A, B). This suggests that each CA1 cell possesses not only a preconfigured

activity level but also an inherent role in spatial coding. Cells with a high gPC index

convey a higher amount of spatial information per unit of time; that is, these cells

contribute more to coding their location in an environment than those with a low gPC

index (Figure 4F). In contrast, cells with a low-gA index sparsely code the

environmental identity. Cells with diverse gA/gPC indices in the CA1 cooperatively

code external space with multiresolution. The sparse code for environment allows

simple mechanisms for forming associations with specific environments (Olshausen and

Field, 2004). In the CA1, contextual fear memory engram cells, defined as neurons that

excite and express c-fos at memory encoding and whose reactivation results in memory

recall, show high environmental specificity and low spatial information content (Tanaka

et al., 2018). The activity profile of the memory engram cells resembles that of low-gA

cells in the present study (Figure 4F). Therefore, it would be interesting to examine the

relationship between a cell's c-fos expression during spatial memory encoding and its

gA/gPC indices.

Although each cell's activity level and gA/gPC indices were preserved over

time, we also showed that the environmental tuning of its activity was unstable (Figure

4C, D). However, at the population level, CA1 cells showed distinct representations for

different environments for at least 6 days (Figure S1B). These results suggest that

environmental information was retained as population activity in CA1 cells.

We employed calcium imaging for long-term tracking of individual cell activity,

which cannot be accomplished reliably using electrophysiological techniques. For this

technique, however, the neocortex above the hippocampus needs to be removed for

optical access to the CA1 cells (Figure 1B). Although the basic activity profiles of the

CA1 cells were consistent with those reported in previous electrophysiological studies

(Figure 1E, 2A), an unknown effect of this surgery on their neural activity cannot be

excluded. Validation of our results will be obtained using a less invasive optical method

(Kondo et al., 2017).

To drive GCaMP expression, we used human synapsin I gene promoter, which

promotes transgene expression in both excitatory and inhibitory cells. Recent studies

have reported that a subset of inhibitory cells in the CA1 show spatial and

environmental modulation in activity (Geiller et al., 2020; Schuette et al., 2022).

Therefore, it will be of interest and value to study the relationship between gA/gPC

indices and subtypes of neurons.

The studies of long-term hippocampal activity measurement discussed here

were performed in mice. However, recent application of calcium imaging on species

other than mice have revealed different activity profiles of CA1 cells. For example,

highly stable activity and spatial tuning were detected in bat CA1 cells, and a high

percentage of place cells were identified in rat CA1 cells (Liberti et al., 2022;

Wirtshafter and Disterhoft, 2022). The temporal dynamics of gA/gPC indices may also

be different in these species.

Conflicts of interest statement

The author has no conflicts of interest to declare.

Acknowledgments

This work is supported by JSPS KAKENHI (Grant No. 17K19436, 20K07715) to Y.H.,

JSPS KAKENHI 20H04849 to R.K, and AMED-CREST 21GM1510003 to K.K. The

authors would like to thank Enago (www.enago.jp) for the English language review.

References

Alme CB, Miao C, Jezek K, Treves A, Moser EI, Moser M-B (2014) Place cells in the

hippocampus: Eleven maps for eleven rooms. Proc Natl Acad Sci

111:18428–18435.

Cai DJ et al. (2016) A shared neural ensemble links distinct contextual memories

encoded close in time. Nature 534:115–118.

Deitch D, Rubin A, Ziv Y (2021) Representational drift in the mouse visual cortex. Curr

Biol 31:4327–4339.

Driscoll LN, Pettit NL, Minderer M, Chettih SN, Harvey CD (2017) Dynamic

Reorganization of Neuronal Activity Patterns in Parietal Cortex. Cell 170:986–999.

Ferbinteanu J, Shapiro ML (2003) Prospective and retrospective memory coding in the

hippocampus. Neuron 40:1227–1239.

Geiller T, Vancura B, Terada S, Troullinou E, Chavlis S, Tsagkatakis G, Tsakalides P,

Ócsai K, Poirazi P, Rózsa BJ, Losonczy A (2020) Large-Scale 3D Two-Photon

Imaging of Molecularly Identified CA1 Interneuron Dynamics in Behaving Mice.

Neuron 108:968-983.e9.

Gonzalez WG, Zhang H, Harutyunyan A, Lois C (2019) Persistence of neuronal

representations through time and damage in the hippocampus. Science

825:821–825.

Hainmueller T, Bartos M (2018) Parallel emergence of stable and dynamic memory

engrams in the hippocampus. Nature 558:292–296.

Hayashi Y (2019) NMDA receptor-dependent dynamics of hippocampal place cell

ensembles. J Neurosci 39:5173–5182.

Hirase H, Leinekugel X, Czurko A, Csicsvari J, Buzsaki G (2001) Firing rates of

hippocampal neurons are preserved during subsequent sleep episodes and modified

by novel awake experience. Proc Natl Acad Sci 98:9386–9390.

Kinsky NR, Mau W, Sullivan DW, Levy SJ, Ruesch EA, Hasselmo ME (2020)

Trajectory-modulated hippocampal neurons persist throughout memory-guided

navigation. Nat Commun 11:1–14.

Kinsky NR, Sullivan DW, Mau W, Hasselmo ME, Eichenbaum HB (2018a)

Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long

Timescales. Curr Biol 28:3578–3588.

Kinsky NR, Sullivan DW, Mau W, Hasselmo ME, Eichenbaum HB (2018b)

Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long

Timescales. Curr Biol 28:3578-3588.e6.

Kondo M, Kobayashi K, Ohkura M, Nakai J, Matsuzaki M (2017) Two-Photon calcium

imaging of the medial prefrontal cortex and hippocampus without cortical invasion.

Elife 6.

Lee JS, Briguglio JJ, Cohen JD, Romani S, Lee AK (2020) The Statistical Structure of

the Hippocampal Code for Space as a Function of Time, Context, and Value. Cell

183:620–635.

Leutgeb S, Leutgeb JK, Barnes CA, Moser EI, Mcnaughton BL, Moser M-B (2005)

Independent Codes for Spatial and Episodic Memory in Hippocampal Neuronal

Ensembles. Science 309:619–623.

Liberti WA, Schmid TA, Forli A, Snyder M, Yartsev MM (2022) A stable hippocampal

code in freely flying bats. Nature 604:98–103.

Lu L, Igarashi KM, Witter MP, Moser EI, Moser MB (2015) Topography of Place

Maps along the CA3-to-CA2 Axis of the Hippocampus. Neuron 87:1078–1092.

MacDonald CJ, Lepage KQ, Eden UT, Eichenbaum H (2011) Hippocampal "time cells"

bridge the gap in memory for discontiguous events. Neuron 71:737–749.

Mankin EA, Sparks FT, Slayyeh B, Sutherland RJ, Leutgeb S, Leutgeb JK (2012)

Neuronal code for extended time in the hippocampus. Proc Natl Acad Sci U S A

109:19462–19467.

Mizuseki K, Buzsáki G (2013) Preconfigured, skewed distribution of firing rates in the

hippocampus and entorhinal cortex. Cell Rep 4:1010–1021.

Moser M-B, Rowland DC, Moser EI (2015) Place Cells, Grid Cells, and Memory. Cold

Spring Harb Perspect Biol 7:a021808.

Muller RU, Kubie JL (1987) The effects of changes in the environment on the spatial

firing of hippocampal complex-spike cells. J Neurosci 7:1951–1968.

O'Keefe J, Conway DH (1978) Hippocampal place units in the freely moving rat: why

they fire where they fire. Exp brain Res 31:573–590.

O'Keefe J, Dostrovsky J (1971) The hippocampus as a spatial map. Preliminary

evidence from unit activity in the freely-moving rat. Brain Res 34:171–175.

O'Keefe J, Nadel L (1978) The hippocampus as a cognitive map. Oxford University

Press.

Olshausen BA, Field DJ (2004) Sparse coding of sensory inputs. Curr Opin Neurobiol

14:481–487.

Pastalkova E, Itskov V, Amarasingham A, Buzsaki G (2008) Internally Generated Cell

Assembly Sequences in the Rat Hippocampus. Science 321:1322–1327.

Pnevmatikakis EA, Giovannucci A (2017) NoRMCorre: An online algorithm for

piecewise rigid motion correction of calcium imaging data. J Neurosci Methods

291:83–94.

Pnevmatikakis EAA, Soudry D, Gao Y, Machado TA, Merel J, Pfau D, Reardon T, Mu

Y, Lacefield C, Yang W, Ahrens M, Bruno R, Jessell TM, Peterka DSS, Yuste R,

Paninski L (2016) Simultaneous Denoising, Deconvolution, and Demixing of

Calcium Imaging Data. Neuron 89:299.

Rubin A, Geva N, Sheintuch L, Ziv Y (2015) Hippocampal ensemble dynamics

timestamp events in long-term memory. Elife 4:1–16.

Schoonover CE, Ohashi SN, Axel R, Fink AJP (2021) Representational drift in primary

olfactory cortex. Nature 594:541–546.

Schuette PJ, Ikebara JM, Maesta-Pereira S, Torossian A, Sethi E, Kihara AH, Kao JC,

Reis FMC V., Adhikari A (2022) GABAergic CA1 neurons are more stable

following context changes than glutamatergic cells. Sci Rep 12:10310.

Sheintuch L, Rubin A, Brande-Eilat N, Geva N, Sadeh N, Pinchasof O, Ziv Y (2017)

Tracking the Same Neurons across Multiple Days in Ca2+ Imaging Data. Cell Rep

21:1102–1115.

Skaggs WE, McNaughton BL, Gothard KM, Markus EJ (1993) An

information-theoretic approach to deciphering the hippocampal code (Hanson SJ,

Cowan JD, Giles CI, eds). San Mateo, CA: Morgan Kaufmann.

Tanaka KZ, He H, Tomar A, Niisato K, Huang AJY, McHugh TJ (2018) The

hippocampal engram maps experience but not place. Science 361:392–397.

Wilt BA, Burns LD, Wei Ho ET, Ghosh KK, Mukamel EA, Schnitzer MJ (2009)

Advances in light microscopy for neuroscience. Annu Rev Neurosci 32:435–506.

Wirtshafter HS, Disterhoft JF (2022)   In Vivo Multi-Day Calcium Imaging of CA1

Hippocampus in Freely Moving Rats Reveals a High Preponderance of Place Cells

with Consistent Place Fields . J Neurosci 42:4538–4554.

Wood ER, Dudchenko PA, Robitsek RJ, Eichenbaum H (2000) Hippocampal neurons

encode information about different types of memory episodes occurring in the

same location. Neuron 27:623–633.

Zhou P, Resendez SL, Rodriguez-Romaguera J, Jimenez JC, Neufeld SQ, Stuber GD,

Hen R, Kheirbek MA, Sabatini BL, Kass RE, Paninski L (2018) Efficient and

accurate extraction of in vivo calcium signals from microendoscopic video data.

Elife 7:1–37.

Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, El Gamal A, Schnitzer

MJ (2013) Long-term dynamics of CA1 hippocampal place codes. Nat Neurosci

16:264–266.

Figure Legends

Figure 1. Experimental overview.

(**A**) Mice explored four testing arenas per day at 3-day intervals. (**B**) Left: Schematic

showing a miniature microscope. Right: Schematic representation showing the objective

lens and chronic window implant above the CA1 region. (**C**) Maximum intensity

projection of the fluorescence image stream. (**D**) Upper: An example calcium trace of a

cell. Lower: Spike train estimated with the OASIS software. (**E**) Firing rate distribution

of CA1 cells (n = 6,425 cells). Histogram: data, solid line: lognormal fit.

Figure 2. Hippocampal activity recorded in the four environments.

Place cell activity in four different environments. (**A**) Top row: Top view showing the

four environments. Second row: Typical mouse trajectories. Remaining rows: Place

field maps for five representative cells. Red indicates the maximum firing rate, blue

indicates silence. Regions not visited by the mice are shown in gray. Asterisks represent

activity above the threshold (mean firing rate > 0.01 Hz). (**B**) Distribution of the gA

index with a variable threshold for activity. Data are presented as the mean ± SEM (n =

24 sessions involving 6 mice). (**C**) Distribution of the gPC index. Data are presented as

the mean ± SEM (n = 24 sessions involving 6 mice). (**D**) Place cell ratio on each day

(left) and in each environment (right). Each dot shows an individual recording in a

single environment. Solid lines indicate the mean value.

Figure 3. Firing rate correlation across sessions

(**A**) Change in firing rate between two adjacent sessions. Each dot in the upper panels

shows one cell's firing rate across two adjacent sessions. Heat maps in the lower panels

show the fraction of cells. Left to right: columns show the firing rates in a single (same)

environment, average firing rates in 5-min data randomly gathered from 2, 3, and 4

environments. Pearson's correlation coefficient (R) is displayed in each panel. (**B**) The

correlation coefficients for individual animals. The black line shows mean ± SEM. Each

gray circle shows an individual animal (n = 6 animals). Asterisks indicate significant

differences (*$p < 0.05$, ***$p < 0.0001$).

Figure 4. Session-to-session dynamics of the gA/gPC indices

(**A**) Changes in cell's gA index between two adjacent sessions. Left: Open circles

represent each value. The solid line shows a linear regression fit to the data.

Right: Distribution of gA index at $N+1$st session for cells showing each gA index at Nth

session. Data are presented as mean ± SEM (n = 18 session pairs from six animals).

(**B**) Changes in cell's gPC index between two adjacent sessions. Left: Open circles

represent each value. The solid line shows a linear regression fit to the data.

Right: Distribution of gPC index at $N+1$st session for cells showing each gA index at

Nth session. Data are presented as mean ± SEM (n = 18 session pairs from six animals).

Solid line shows a linear regression fit to the data. Asterisks indicate significant

correlations (***$p < 0.0001$). (**C**) Fraction of "preference changed" cells and

"preference unchanged" cells between sessions out of 19,275 observations of the cells.

Others showed no activity in either or both day ("silent->silent" or "silent->active or

active->silent", respectively). (**D**) Cumulative distribution of environment preference

index for "preference changed" cells and "preference unchanged" cells. (**E**) The gPC

index versus the mean firing rates. The solid line shows a linear regression fit to the data

(log-transformed). (**F**) The gPC index versus the spatial information rates. The solid line

shows a linear regression fit to the data (log-transformed). (**G**) gA index versus the

firing rate difference for the environment in which a cell showed the highest firing rate

and firing rate averaged over the other three environments. The solid line shows a linear

regression fit to the data (log-transformed). Dots represent each value. Horizontal lines

show the mean values, and the whiskers show the standard deviation. Asterisks indicate

significant differences (***$p < 0.0001$).

Figure 1

A

env A   env B   env C   env D

Recording:

Day: 1   4   7   10
Session #: 1   2   3   4

B

Image Sensor

Tube Lens

Emission Filter

Dichroic Mirror

Electrowetting Lens

Base Plate

Excitation Filter

Drum Lens

LED

Objective lens

Glass window

C

D

Fluorescence (AU)

Firing rate (Hz)

Time (s)

E

Cell number

Mean firing rate (Hz)

Figure 2

A

env A env B env C env D

0 env active
Cell ID: 2

1 env active
Cell ID: 11

2 env active
Cell ID: 56

3 env active
Cell ID: 15

4 env active
Cell ID: 1

B

Thr (Hz)
0.01
0.02
0.04
0.08

Fraction

Number of active environments
(gA index)

C

Fraction

Number of environments
where each cell shows PC activity
(gPC index)

D

Place cell ratio

Day

Place cell ratio

Env

Figure 3

A

B

Figure 4