1

The temporal and contextual stability of activity levels in

hippocampal CA1 cells

Yuichiro Hayashi[1]*, Ko Kobayakawa[1], and Reiko Kobayakawa[1]

[1] Institute of Biomedical Sciences, Kansai Medical University, 2-5-1 Shin-machi,

Hirakata, Osaka 573-1010, Japan

*Correspondence: hayasyui@hirakata.kmu.ac.jp

Abstract

Recent studies have shown that the activity levels of hippocampal neurons during

exploration are not constant but change on a daily to weekly scale. However, a few

other studies suggest that their activity levels are preserved over time. To resolve this

contradiction, we repeatedly measured the neural activity of the mouse hippocampus in

four different environments. The activity level of hippocampal neurons fluctuated

greatly in one environment, but the mean activity level across all the environments was

more stable. Each cell's preferred environment frequently changed over time. The

number of environments in which a cell showed activity varied across cells, ranging

from zero to four, and the value also tended to be preserved. Cells that showed place

cell activity in many environments had higher mean firing rates and higher spatial

information content, whereas cells active only in a small number of environments were

suitable for coding environmental identity. These results suggest that although the

activity level in a single environment appears unstable, each cell has not only an

inherent activity level but also play a characteristic role in the coding of space.

Introduction

The hippocampus is known as a brain region that plays a crucial role in episodic

memory formation. To support this, the activity of hippocampal neurons has been found

to encode place, environment, time, and various task-relevant information (1–6). In

particular, place cell activity, i.e., the activity of hippocampal cells in restricted portions

in an environment, has been extensively studied (7,8). The activity pattern of place cells

can change or disappear when animals move to a different environment (5,9–11). This

phenomenon is called remapping, and it seems to provide a mechanism through which

the hippocampus represents different environments.

To store episodic information in a neural circuit, the activity must reliably

respond to the same external stimuli across long timescales. However, recent studies

have shown that the place receptive fields and activity levels of hippocampal neurons

are not constant, but fluctuate on a daily to weekly scale (12–17). Such fluctuations are

called "representational drift" and have also been found in other brain regions, such as

the parietal cortex, visual cortex, and olfactory cortex (18–20). On the contrary, several

studies have reported evidence that each cell has a preconfigured activity level and that

the firing rates of the neurons in the hippocampus and entorhinal cortex are preserved

across different brain states (21,22). Recording in a very large environment (>40 m

long) revealed that the location of the place fields of hippocampal neurons fluctuated on

the day to week timescale, but the number of place fields was preserved over months

(23).

One possible explanation for this discrepancy is that the former studies were

recorded in small spaces (<1 m^2), while the latter was measured in a large environment.

In a larger environment, a greater number of cells show place cell activity and the size

and number of place field for each cell increase (24,25). Considering the presence of

representational drift in the place fields, their activity levels appear to fluctuate in a

small environment, but not in a large environment if the size is enough to average out

the fluctuations. If that is the case, the averaged activity for multiple small environments

should be preserved over time.

In the present study, we measured neural activity in the mouse hippocampal

CA1 area in four different environments with different sizes and colors over time. The

activity level of hippocampal neurons fluctuated greatly in one environment but was

stable when averaged across all environments. We also found that the number of

environments in which the cells showed activity also tended to be conserved.

Results

Experimental outline

To investigate the place cell activity for multiple spatial environments in the

hippocampus, neural activity was recorded successively in four arenas with different

shapes and colors (Figure 1A). Before commencing the recording sessions, mice (n = 6)

were trained to chase food in these arenas for at least five days. Each mouse then

underwent four 5-min recording sessions per day for three-day intervals (Figure 1A).

Neural activity in the dorsal CA1 region of the hippocampus was recorded

using a combination of a genetically encoded calcium indicator GCaMP6f and a

miniature microscope (Figure 1B and C). We extracted calcium traces of putative

neurons from the fluorescence image stream using CNMF-E (26). The firing rate of

each neuron was estimated from its calcium trace using OASIS software (27) (Figure

1D). The firing rate distribution of the recorded cells showed lognormal distribution

(Figure 1E), which was consistent with the findings of the previous electrophysiological

study (21). With the spatial footprints of the cells extracted with CNMF-E from each

day's fluorescence image stream, we used Cellreg (28) to track cells across days (Figure

S2).

Activity of CA1 cells showed a variety of environmental specificity

A total of 6425 cells were recorded from six mice in four different environments. In

accordance with a previous study, we found a variety of cells ranging from those that

showed no activity in any environment to those that showed activity in all the four

environments (Figure 2A and B) (10). We termed the number of environments where a

cell exhibited activity above the threshold as the "global activity level index" (gA index).

Because a fraction of cells in the hippocampus show activity but are not place cells, we

also classified cells according to the number of environments where a cell showed

statistically significant PC activity. The value was termed "global place cell activity

level index" (gPC index). As shown in Figure 2C, CA1 cells showed a variety of gPC

activity levels. The fraction of place cell did not change between environments or over

days, indicating that these mice were highly familiar to all the environments (Figure

S1).

Activity level averaged across multiple environments was more stable than that for

a single environment

For longitudinal analysis of the same neuron, the same cells need to be reliably

registered across days. To ensure reliability of cell registration, we checked Register

score and Psame value (28). The average registration score was 0.86 ± 0.020 (mean $\pm$

SEM, n = 6 animals, Figure S2E, *Left*), and the average Psame value was 0.83 ± 0.052

(mean $\pm$ SEM, n = 6 animals, Figure S2E, *Right*). These values were comparable to the

previous study in the hippocampal CA1 area (28).

104   Then we evaluated the reliability of cell registration at the functional level. For

this purpose, population vector (PV) correlation (29) of spatial representation for the

four environments between sessions was calculated. Consistent with previous studies

(14,30), PV correlation between sessions in the same environment was significantly

higher than between sessions in different environments for at least 6-day session

interval (Figure S3). The result also corroborated that the same neurons were reliably

tracked across multiple sessions.

111   Not only across-day, but also within-day registration quality affects the analysis.

In wide-field imaging data, due to light scattering and lack of optical sectioning

capability, signals from neighboring cells are overlapping (31). Although CNMF-E has

shown to effectively demix signals from overlapping cells (26), one possible concern is

that activities of multiple spatially-overlapping cells may be merged into an activity of

single high gA cell. To test this possibility, we examined the relationship between cell's

gA index and its spatial footprint size of the ROI extracted by CNMF-E. If multiple

spatially overlapping cells are merged into single cell, its spatial footprint would tend to

be larger than that for genuine single cell. As shown in Figure S4, there was no positive

but slight negative correlation between gA index and spatial footprint size, suggesting

that demix error on the cellular signal extraction was not frequent for high gA cells than

for low gA cells.

Then we analyzed the stability of neural activity of each cell in a single

environment and across all four environments. To make a fair comparison, the mean

firing rate for four environments was calculated over 5-min of data randomly extracted

from recordings in the four environments. We plotted the relationship between the

activity levels of two consecutive sessions with an interval of three days. As show in

Figure 3A, the activity correlation for each single environment was lower than that for

the averaged activity across all environments. To quantify the activity correlation, we

calculated the firing rate difference, which refers to the absolute values of differences

between mean firing rates divided by their sum. A significantly higher correlation was

observed for the averaged activity across all environments than for an activity in a

single environment (Figure 3B $U = 3.73 \times 10^9$, $p = 2.02 \times 10^5$, n = 77100 observations

of cells for single environment, n = 19275 observations of cells for all environments,

Mann–Whitney U test). The fluctuating activity for single environment and stable

averaged activity across multiple environments (Figure 3A) suggested that each cell

changed preference for its environment. To test this, the environment in which each cell

showed the highest firing rate across all environments was visualized. As shown in

Figure 3C, each cell's preferred environment (highest firing rate environment) changed

frequently across sessions, and the transition rate of the preferred environment was ~0.6

(Figure 3D).

The gA and gPC indexes were preserved over time

Next, we tested whether the gA and gPC indexes were preserved. A significant

correlation of the gA index between adjacent sessions was found (Figure 4A, n = 90

session pairs, r = 0.780, $p < 0.0001$). We used a threshold value of 0.01 Hz, but a

significant correlation was also found with threshold levels ranging from 0.01 to 0.08

Hz (Figure S5). The gPC index between adjacent sessions was also correlated (Figure

4B, n = 90 session pairs, r = 0.724, $p < 0.0001$). These results indicate that not the only

mean firing rate but also the gA/gPC index of each cell tended to be preserved over

time.

Given that each cell has a preserved activity level and gA/gPC index, we

wanted to determine whether cells have inherent roles. We then investigated the

relationship between gA/gPC index and the cells' firing properties. The cells showing

higher gPC index had a higher mean firing rates and spatial information rates (Figure

5A, mean firing rate: $H_{(4,25545)} = 390.8$, $p < 0.0001$, n = 8516 observations of neurons

over all recording sessions; 5B, spatial information: $H_{(3, 25545)} = 427.4$, $p < 0.0001$, n =

8516 observations of place cells over all recording sessions, Kruskal–Wallis test

followed by Dunn's *post hoc* test). However, the spatial information content per spike

was not significantly different among the groups (Figure 5C, $H_{(3,25545)} = 78.73$, $p <$

0.0001, n = 8516 observations of place cells over all recording sessions, Kruskal–Wallis

test followed by Dunn's *post hoc* test). Therefore, the higher information rate by cells

showing gPC index was attributed to their higher firing rates. These results indicate that

high firing rate and high gPC index cells convey an animal's precise location

information.

On the other hand, cells active in only a few environments can sparsely and

efficiently code environmental identity. To test this, we calculated firing rate differences

between the firing rate for the environment where a cell showed the highest firing rate

and firing rate averaged over the other three environments. Cells do not necessarily

possess position information when encoding the environment. Therefore, we classified

cells with the gA index. The firing rate difference between the maximum firing

rate–environment and the others was highest with the gA index = 1 (Figure 5D, $H_{(3,36933)}$

$= 724.3$, $p < 0.0001$, n $= 12312$ observations of cells over all recording sessions,

Kruskal–Wallis test followed by Dunn's *post hoc* test). This indicates that cells with a

low gA index code the environmental identity rather than the animal's location.

Discussion

In this study, using the head-mount miniature fluorescence microscope, we measured

the neural activity of hippocampal CA1 cells over time in four different environments

(Figure 1). There are contrasting reports on the long-term stability of the activity of

CA1 cells. It has been shown that the place field location and activity levels of CA1

cells are not constant but fluctuate on a daily to weekly scale (12–17,32). In

contradiction to these observations, a recent study reported that the activity levels of

CA1 cells in a very large (~40 m long) environment were stable over time, though the

location of their place field fluctuated (23). The results of this study can resolve this

contradiction. In agreement with former studies, we showed that the activity level of

each CA1 cell in a small environment (~0.1 m^2) varied over days (Figure 3A, *Left*).

However, the temporal variation in the averaged activity level across four environments

was smaller (Figure 3A, *Right*). Previous studies have shown that in a small (~0.4 m^2)

environment, the number of place field of each cell is typically one, whereas in larger

environment, the number of place field for each cell increases (24,25). Therefore, it is

possible that the activity level in a large environment is more stable than in a small

environment if the size of the environment is enough to average out the activity

fluctuation due to representational drift. The activity levels for multiple environments

are also more stable than that for single environment due to the presence of

representational drift.

We also showed the gA and gPC indexes tended to be preserved over time

(Figure 4A and B). This suggests that CA1 cells play an inherent role in spatial coding.

Cells that show activity only in a small number of environments (small gA index)

sparsely and efficiently code environmental identity (Figure 5D), whereas cells that

show PC activity in many environments convey rich information of the location in any

environment. Notably, both gA and gPC indexes are not binary, but are gradually

distributed (Figure 2B and C). Therefore, the possible roles of CA1 cells in spatial

coding are also gradual; there are cells with diverse size of spatial receptive fields from

a portion of an environment to multiple environments.

One of the questions to be addressed next is the relationship between gPC

activity level and c-Fos expression. In the present study, we recorded neural activity in a

familiar environment, but not during a new experience (Figure 1). In contrast to

exploration in familiar environment, exposure to a novel environment or learning a new

task induces an immediate early gene c-Fos in a subset of hippocampal cells, and the

artificial activation of these cells induces memory recall (33). Therefore, these cells are

considered to be involved in a specific memory. Moreover, a recent study reported that

c-Fos expressing place cells in the CA1 have distinct firing properties. Place cells

expressing c-Fos during novel experiences showed higher mean firing rates and less

spatial information content from those c-Fos negative cells. Interestingly, the c-Fos

positive cells also showed a spatial environment-selective firing pattern but the c-Fos

negative cells did not (34). The relationship between c-Fos expression and gA/gPC

index during learning is a topic for future research.

Materials and Methods

Animal subjects

All experiments complied with the applicable guidelines and regulations. Animal care

and use followed protocols approved by the Animal Research Committee of Kansai

Medical University. All experiments complied with the ARRIVE guidelines

(https://arriveguidelines.org/arrive-guidelines/experimental-animals). Nine adult male

C57BL/6N mice aged 12–18 weeks were used. The mice were purchased from Japan

SLC, Inc. (Shizuoka, Japan), housed under a standard 12-h light/dark cycle, and allowed

ad libitum access to food and water.

Viral constructs

We obtained an AAVrh10.hSyn.GCaMP6f.WPRE.SV40 virus from the University of

Pennsylvania Vector Core at a titer of ~1.3×10^{14} GC/mL and diluted it to ~5×10^{12}

GC/mL with phosphate buffered saline.

Miniature head-mount microscope

The miniature microscope design was based on the UCLA miniscope v3

(miniscope.org). Excitation light was emitted by a blue LED (LXML-PB01-0030;

Lumileds, Aachen, Germany). The light passed through an excitation filter (ET470/40x;

Chroma Technology, Bellows Falls, VT) and was reflected by a dichroic mirror

(T495lpxr; Chroma Technology) onto the tissue through an EWTL (Varioptic A-25H0;

Corning, Corning, NY), an objective lens (tandem configuration of achromatic doublets;

#84-126 and #49-271, Edmund Optics, Burlington, NJ). The fluorescent emissions

collected by the lenses were passed through the dichroic mirror and an emission filter

(ET525/50m; Chroma Technology). The fluorescence image was focused by a tube lens

(#49-277; Edmund Optics) and captured by a CMOS image sensor (MT9V032; ON

Semiconductor, Phoenix, AZ). The microscope body was made using a 3D printer

(From2, resin type FGPBLK03; Formlabs, Somerville, MA). The design files are

available at https://github.com/yuichirohayashi/Zscope. The driving voltage for the

EWTL (square wave, 2 kHz) was generated using a data acquisition board (PCIe-6259,

National Instruments, Dallas, TX) controlled by custom software written in LABVIEW

7.1 (National Instruments, Dallas, TX) and amplified with a linear amplifier

(As-904-150B, NF Corporation, Yokohama, Japan).

Surgery

The animals were anesthetized using isoflurane. The skull was exposed, and a small

hole (<0.5 mm) was made over the right hemisphere (1.5 mm lateral to the midline, 2.3

mm posterior to the bregma). Then, 200 nL of GCaMP6f virus was injected into CA1

(1.2 mm ventral from the brain surface). One week after the viral injection, the animals

were anesthetized using isoflurane, and a 2.8 mm-diameter craniotomy was performed.

The dura was removed, and the underlying cortex was aspirated. A stainless-steel

cannula (2.76 mm outer diameter, 2.40 mm inner diameter, 1.5 mm height) covered by a

glass coverslip (0.12 mm thick) was inserted over the dorsal CA1. Dental cement

(Shofu, Kyoto, Japan) was used to glue the cannula to the skull.

Four weeks after the cannula implantation, GCaMP-expressing neurons were

imaged using the head-mount microscope. The aluminum baseplate of the microscope

was cemented in a position at which the neurons were visible.

Recording arenas

Calcium activity was recorded in four arenas of different shapes and colors: a tall green

box ($37 \times 25 \times 56$ cm high), a short gray box ($37 \times 25 \times 35$ cm high), a lime round

basket ($39 \times 26 \times 25$ cm high), and a white square basket ($36 \times 25 \times 25$ cm high). One

side of each arena was painted a different color or decorated with plastic tape (Figure

1A).

Behavioral training

One week after the baseplate surgery, the mice were food-restricted and maintained at

above 85% of their initial weight. Mice were trained to forage for food pellets (14 mg,

#F05684, Bio-Serv, Flemington, NJ) scattered on the floor of the arenas.

Recording sessions

After at least five days of behavioral training, mice underwent four 5-min recording

sessions per day in the four different arenas in a random sequential order. The arenas

were cleaned with 70% ethanol prior to each recording session to eliminate any

olfactory cues. During sessions, food pellets were thrown into the arena about every 1

min. The excitation light intensity was approximately 0.1–0.5 mW/mm^2, and

fluorescence images were captured at 30 Hz. The focus was adjusted to match the

previous one. To record mouse behavior, an overhead camera (DMK22BUC03, The

Imaging source, Bremen, Germany) was used.

Quantification and statistical analyses

All significance values are described in the text. Statistical analyses were performed

using GraphPad Prism 5 (GraphPad Software, La Jolla, CA) and MATLAB 2019a. The

statistical significance was set at $P < 0.05$ for all statistical analyses.

Image processing

ImageJ 1.52 (National Institutes of Health, MA) and MATLAB 2019a (Mathworks,

Natick, MA) were used for all analyses. The calcium imaging data recorded at same day

were concatenated and processed using MiniscopeAnalysis

(https://github.com/etterguillaume/MiniscopeAnalysis). First, the images were

downsampled spatially by factors of two and motion-corrected using NoRMCorre (35).

The downsampled, motion-corrected videos were then processed using CNMF-E (26) to

extract fluorescence traces from individual neurons. The fluorescence traces were

finally processed using OASIS software (27) for denoizing and deconvolving to

estimate spiking activity.

Registration of cells across sessions

For day-to-day longitudinal cell registration, the spatial footprints of the cells extracted

with CNMF-E from each day's fluorescence image stream were processed with Cellreg

routine (28) with default parameters (maximum distance < 12 μm, spatial correlation >

0.74, and Psame threshold = 0.5). Registration quality was assessed using Register score

and Psame value for registered cells (28).

Place fields

To analyze the place fields, calcium events that occurred when the animal's velocity was

less than 1 cm/s were filtered out to eliminate nonspecific activity at rest. The place

field maps were constructed by finding the spike number at each spatial bin (2 cm× 2

cm) divided by the dwell time in the bin. The maps were smoothed using a Gaussian

function with a standard deviation of 3 cm.

For each place field of the cells, the mutual information between calcium

transients and the mouse's location was calculated (36). To assess the significance of

spatial selectivity, Monte Carlo P values were calculated. A total of 1000 distinct

shuffles of the calcium transient times were performed and the mutual information for

each shuffle was calculated. The P value was defined as the fraction of shuffles that

exceeded the mutual information of the cell. Cells with $P < 0.05$ were considered place

cells.

Rate difference

The rate difference in the place field was determined by calculating the unsigned rate

difference between the peak calcium transient rates in the two recording sessions and by

dividing the difference by the sum of the two rates.

Population vector correlation

To compute the population vector (PV) correlation between two sessions, the place

fields of all simultaneously recorded place cells were stacked into a 3D matrix (place

field map × cell identity). At each spatial bin, the firing rate vector of all cells represents

the PV for that spatial bin. To compare two recording sessions, the Pearson correlation

coefficient was calculated between each pair of PVs at the corresponding locations, and

the correlation coefficients of all spatial bins were averaged (29).

Funding Disclosure

This work is supported by JSPS KAKENHI (Grant No. 17K19436, 20K07715) to Y.H.,

JSPS KAKENHI 20H04849 to R.K, and AMED-CREST 21gm1510003 to K.K..

Competing Interests

The authors have declared that no competing interests exist.

Author Contributions

Conceptualization, Y.H.; Methodology, Y.H.; Investigation, Y.H.; Writing –Original

Draft, Y.H.; Writing –Review & Editing, Y.H., K.K. and R.K.; Funding Acquisition, Y.H.

and R.K.; Resources, Y.H., K.K. and R.K.; Supervision, Y.H.

Figure Legends

Figure 1. Experimental overview.

(**A**) Mice explored four testing arenas per day for three-day intervals. (**B**) *Left,* Section

diagram of the miniature microscope. *Right,* Schematic representation of the objective

lens and the chronic window implant above CA1. (**C**) Maximum intensity projection of

the fluorescence image stream. (**D**) *Upper,* an example calcium trace of a cell. *Lower,*

spike train estimated with OASIS software. (**E**) Distribution of the firing rates of CA1

cells (n = 6425 cells). Histogram: data; red line: lognormal fit.

Figure 2. Hippocampal activity recorded in the four environments

Place cell activity in four different environments. (**A**) *Top row,* top view of the four

environments. *Second row,* typical mouse's trajectories. *From third to bottom rows,*

Place field maps for five representative cells. Red indicates the maximum firing rate and

blue indicates silent. Regions not visited by the mouse are shown in gray. Asterisks

represent activity above threshold (mean firing rate > 0.01 Hz). (**B**) Distribution of gA

index with variable threshold for activity. Data represent mean ± SEM (n = 24 sessions

from six mice). (**C**) Distribution of gPC index. Data represent mean ± SEM (n = 24

sessions from six mice).

Figure 3. Session-to-session activity dynamics in multiple environments

(**A**) Change of firing rate between two adjacent sessions. Each dot shows one cell's

firing rates across two adjacent sessions. *Left,* firing rates in a single (same)

environment. *Right,* averaged firing rates over all four environments. (**B**) Firing rate

difference between two adjacent sessions. The box plots show the median and quartiles,

and the whisker caps of the box plots show the minimum and maximum values.

Asterisks indicate significant differences (***$P < 0.0001$). (**C**) Change of environmental

selectivity. Environments in which each cell exhibits the highest firing rate in each

session are displayed. (**D**) Transition rates of environmental selectivity across adjacent

sessions. Each dot shows an individual animal. Solid lines indicate mean value.

Figure 4. Session-to-session dynamics of gA/gPC index

(**A**) Changes in cell's gA index between two adjacent sessions. *Left,* data are shown as

mean ± SEM (n = 18 session pairs from six animals). *Right,* open circles represent each

value. The solid line shows a linear regression fit to the data. (**B**) Changes in cell's gPC

index between two adjacent sessions. *Left,* data are shown as mean ± SEM (n = 18

session pairs from six animals). *Right,* open circles represent each value. The solid line

shows a linear regression fit to the data. Asterisks indicate significant correlation (****P*

< 0.0001).

Figure 5. Relationship between gA/gPC index and cell's firing properties

(**A**) gPC index versus mean firing rates. (**B**) gPC index versus spatial information rates.

(**C**) gPC index versus spatial information rate per spike. (**D**) gA index versus firing rate

difference between the firing rate for the environment in which a cell showed the

highest firing rate and firing rate averaged over the other three environments. The box

plots show the median and quartiles, and the whisker caps of the box plots show the

minimum and maximum values. Asterisks indicate significant differences (****P* <

0.0001).

References

1.   O'Keefe J, Dostrovsky J. The hippocampus as a spatial map. Preliminary

evidence from unit activity in the freely-moving rat. Brain Res [Internet].

1971;34(1):171–5. Available from:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&do

pt=Citation&list_uids=5124915

2.   Ferbinteanu J, Shapiro ML. Prospective and retrospective memory coding in the

hippocampus. Neuron. 2003;40(6):1227–39.

3.   MacDonald CJ, Lepage KQ, Eden UT, Eichenbaum H. Hippocampal "time cells"

bridge the gap in memory for discontiguous events. Neuron [Internet].

2011;71(4):737–49. Available from:

http://dx.doi.org/10.1016/j.neuron.2011.07.012

4.   Pastalkova E, Itskov V, Amarasingham A, Buzsaki G. Internally Generated Cell

Assembly Sequences in the Rat Hippocampus. Science [Internet].

2008;321(5894):1322–7. Available from:

http://www.sciencemag.org/cgi/doi/10.1126/science.1159775

5.   Muller RU, Kubie JL. The effects of changes in the environment on the spatial

firing of hippocampal complex-spike cells. J Neurosci. 1987;7(7):1951–68.

6.   Wood ER, Dudchenko PA, Robitsek RJ, Eichenbaum H. Hippocampal neurons

encode information about different types of memory episodes occurring in the

same location. Neuron [Internet]. 2000 Sep [cited 2021 Oct 18];27(3):623–33.

425   Available from: http://www.ncbi.nlm.nih.gov/pubmed/11055443

426 7. O'Keefe J, Nadel L. The hippocampus as a cognitive map. Oxford University

427   Press; 1978.

428 8. Moser M-B, Rowland DC, Moser EI. Place Cells, Grid Cells, and Memory. Cold

429   Spring Harb Perspect Biol [Internet]. 2015 Feb 2 [cited 2017 Jul

430   21];7(2):a021808. Available from:

431   http://www.ncbi.nlm.nih.gov/pubmed/25646382

432 9. Alme CB, Miao C, Jezek K, Treves A, Moser EI, Moser M-B. Place cells in the

433   hippocampus: Eleven maps for eleven rooms. Proc Natl Acad Sci [Internet].

434   2014;111(52):18428–35. Available from:

435   http://www.pnas.org/lookup/doi/10.1073/pnas.1421056111

436 10. Lu L, Igarashi KM, Witter MP, Moser EI, Moser MB. Topography of Place

437   Maps along the CA3-to-CA2 Axis of the Hippocampus. Neuron.

438   2015;87(5):1078–92.

439 11. O'Keefe J, Conway DH. Hippocampal place units in the freely moving rat: why

440   they fire where they fire. Exp brain Res [Internet]. 1978 Apr 14 [cited 2021 Oct

441   20];31(4):573–90. Available from: http://www.ncbi.nlm.nih.gov/pubmed/658182

442 12. Kinsky NR, Sullivan DW, Mau W, Hasselmo ME, Eichenbaum HB.

Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long

Timescales. Curr Biol [Internet]. 2018;28(22):3578–88. Available from:

https://doi.org/10.1016/j.cub.2018.09.037

13.   Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, et al. Long-term

dynamics of CA1 hippocampal place codes. Nat Neurosci [Internet]. 2013/02/12.

2013 Mar [cited 2017 Jul 11];16(3):264–6. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/23396101

14.   Rubin A, Geva N, Sheintuch L, Ziv Y. Hippocampal ensemble dynamics

timestamp events in long-term memory. Elife. 2015;4(DECEMBER2015):1–16.

15.   Hayashi Y. NMDA receptor-dependent dynamics of hippocampal place cell

ensembles. J Neurosci. 2019;39(26):5173–82.

16.   Gonzalez WG, Zhang H, Harutyunyan A, Lois C. Persistence of neuronal

representations through time and damage in the hippocampus. Science.

2019;825(August):821–5.

17.   Cai DJ, Aharoni D, Shuman T, Shobe J, Biane J, Song W, et al. A shared neural

ensemble links distinct contextual memories encoded close in time. Nature

[Internet]. 2016;534(7605):115–8. Available from:

http://www.nature.com/doifinder/10.1038/nature17955

18.   Driscoll LN, Pettit NL, Minderer M, Chettih SN, Harvey CD. Dynamic

462   Reorganization of Neuronal Activity Patterns in Parietal Cortex. Cell.

463   2017;170(5):986–99.

464   19.   Schoonover CE, Ohashi SN, Axel R, Fink AJP. Representational drift in primary

465   olfactory cortex. Nature [Internet]. 2021 Jun [cited 2021 Oct

466   18];594(7864):541–6. Available from:

467   http://www.ncbi.nlm.nih.gov/pubmed/34108681

468   20.   Deitch D, Rubin A, Ziv Y. Representational drift in the mouse visual cortex. Curr

469   Biol [Internet]. 2021 Oct 11 [cited 2021 Oct 23];31(19):4327–39. Available

470   from: http://www.ncbi.nlm.nih.gov/pubmed/34433077

471   21.   Mizuseki K, Buzsáki G. Preconfigured, skewed distribution of firing rates in the

472   hippocampus and entorhinal cortex. Cell Rep. 2013;4(5):1010–21.

473   22.   Hirase H, Leinekugel X, Czurko A, Csicsvari J, Buzsaki G. Firing rates of

474   hippocampal neurons are preserved during subsequent sleep episodes and

475   modified by novel awake experience. Proc Natl Acad Sci [Internet].

476   2001;98(16):9386–90. Available from:

477   http://www.pnas.org/cgi/doi/10.1073/pnas.161274398

23.   Lee JS, Briguglio JJ, Cohen JD, Romani S, Lee AK. The Statistical Structure of

the Hippocampal Code for Space as a Function of Time, Context, and Value. Cell

[Internet]. 2020 [cited 2021 Oct 18];183(3):620–35. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/33035454

24.   Rich PD, Liaw HP, Lee AK. Large environments reveal the statistical structure

governing hippocampal representations. Science. 2014;345(6198):814–7.

25.   Fenton AA, Kao H-Y, Neymotin SA, Olypher A, Vayntrub Y, Lytton WW, et al.

Unmasking the CA1 Ensemble Place Code by Exposures to Small and Large

Environments: More Place Cells and Multiple, Irregularly Arranged, and

Expanded Place Fields in the Larger Space. J Neurosci [Internet].

2008;28(44):11250–62. Available from:

http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.2862-08.2008

26.   Zhou P, Resendez SL, Rodriguez-Romaguera J, Jimenez JC, Neufeld SQ, Stuber

GD, et al. Efficient and accurate extraction of in vivo calcium signals from

microendoscopic video data. Elife [Internet]. 2018;7:1–37. Available from:

http://arxiv.org/abs/1605.07266

27.   Pnevmatikakis EAA, Soudry D, Gao Y, Machado TA, Merel J, Pfau D, et al.

Simultaneous Denoising, Deconvolution, and Demixing of Calcium Imaging

Data. Neuron [Internet]. 2016 Jan [cited 2016 Dec 21];89(2):299. Available

from: http://linkinghub.elsevier.com/retrieve/pii/S0896627315010843

28.   Sheintuch L, Rubin A, Brande-Eilat N, Geva N, Sadeh N, Pinchasof O, et al.

Tracking the Same Neurons across Multiple Days in Ca2+ Imaging Data. Cell

Rep [Internet]. 2017;21(4):1102–15. Available from:

http://www.cell.com/cell-reports/fulltext/S2211-1247(17)31430-4

29.   Leutgeb S, Leutgeb JK, Barnes CA, Moser EI, Mcnaughton BL, Moser M-B.

Independent Codes for Spatial and Episodic Memory in Hippocampal Neuronal

Ensembles. Science [Internet]. 2005;309(5734):619–23. Available from:

http://www.sciencemag.org/cgi/doi/10.1126/science.1114037

30.   Kinsky NR, Sullivan DW, Mau W, Hasselmo ME, Eichenbaum HB.

Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long

Timescales. Curr Biol. 2018;28(22):3578-3588.e6.

31.   Wilt BA, Burns LD, Wei Ho ET, Ghosh KK, Mukamel EA, Schnitzer MJ.

Advances in light microscopy for neuroscience. Annu Rev Neurosci [Internet].

2009;32:435–506. Available from:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&do

pt=Citation&list_uids=19555292

32.   Kinsky NR, Mau W, Sullivan DW, Levy SJ, Ruesch EA, Hasselmo ME.

Trajectory-modulated hippocampal neurons persist throughout memory-guided

navigation. Nat Commun [Internet]. 2020;11(1):1–14. Available from:

http://dx.doi.org/10.1038/s41467-020-16226-4

33. Liu X, Ramirez S, Pang PT, Puryear CB, Govindarajan A, Deisseroth K, et al.

Optogenetic stimulation of a hippocampal engram activates fear memory recall.

Nature [Internet]. 2012 Mar 22 [cited 2017 Jul 11];484(7394):381–5. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/22441246

34. Tanaka KZ, He H, Tomar A, Niisato K, Huang AJY, McHugh TJ. The

hippocampal engram maps experience but not place. Science.

2018;361(6400):392–7.

35. Pnevmatikakis EA, Giovannucci A. NoRMCorre: An online algorithm for

piecewise rigid motion correction of calcium imaging data. J Neurosci Methods

[Internet]. 2017;291:83–94. Available from:

http://dx.doi.org/10.1016/j.jneumeth.2017.07.031

36. Skaggs WE, McNaughton BL, Gothard KM, Markus EJ. An

information-theoretic approach to deciphering the hippocampal code. Hanson SJ,

Cowan JD, Giles CI, editors. Vol. 5, Advances in neural information processing

systems. San Mateo, CA: Morgan Kaufmann; 1993. 1030–1037 p.

Figure 1

A

env A   env B   env C   env D

Recording:
Day: 1   4   7   10
Session #: 1   2   3   4

B

Image Sensor

Tube Lens
Emission Filter
Dichroic Mirror
Electrowetting Lens
Base Plate

Excitation Filter
Drum Lens
LED

C

D

Fluorescence (AU)

Firing rate (Hz)

Time (s)

E

Cell number

Mean firing rate (Hz)

Figure 2

A

env A   env B   env C   env D

0 env active
Cell ID: 2

1 env active
Cell ID: 11

2 env active
Cell ID: 56

3 env active
Cell ID: 15

4 env active
Cell ID: 1

B

C

Figure 3

Figure 4

Figure 5

Figure S1: Fraction of cells with statistically significant place fields

(A) Place cell ratio in each day. $Q_{(3,69)}$ = 1.639, p = 0.65, n = 24 recordings per day in 6 animals, Friedman test. (B) Place cell ratio in each environment. $Q_{(3,69)}$ = 5.812, p = 0.12, n = 24 recordings for each environment in 6 animals, Friedman test. Each dot shows an individual recording in single environment. Solid lines indicate mean value.

Figure S2: Tracking the same neurons across sessions using Cellreg

Alignment of the imaging data acquired in sessions on different days from a representative animal. (A) The spatial footprints of all identified cells in session 1 (red), 2 (green), and 3 (blue). (B) The centroid position of all identified cells in the all sesisons. (C) Distributions of centroid distances (*left*) and spatial correlation (*right*) between cell-pairs of nearest neighbors and other neighbors across the four sessions. (D) Distributions of centroid distances (*left*) and spatial correlations (*right*) between all neighboring cell-pairs (blue bars) and the modeled distributions of same cells (dashed green curves), different cells (dashed red curves), and their weighted sum (solid black curves). Black dashed lines show the intersection between the two models. Estimated fractions of same cells (green) or different cells (red) above and below these intersections are indicated. (E) Distribution of registration scores (*left*) and average Psame value (*right*) for all animals. Each dot shows an individual animal. Solid lines indicate mean value.

Figure S3: PV correlation versus time

PV correlation for the same environment (blue) and different environment (red) across time. Data represent mean ± SEM. Open circles represent each value (n = 24 session pairs for same environment, n = 36 session pairs for different envitonment). Statistical significance were tested using two-tailed *t* test with Bonferroni correction. Asterisks indicate significant differences (*P < 0.05, ***P < 0.0001, ns: not significant).

Figure S4: gA index versus spatial footprint size

Each cell's gA index versus its spatial footprint size of the spatial components extracted by CNMF-E. Spatial footprint size is defined as the number of pixel whose value is greater than the half-maximal of the cell's spatial components. Dots represent each value. The box plots show the median and quartiles. The solid line shows a linear regression fit to the data. $r = 0.112$, $p < 0.0001$.

Figure S5: Session-to-session dynamics of gA index with variable threshold

(A) Changes in gA index between two adjacent sessions. Activity threshold = 0.02 Hz. n = 90 session pairs, r = 0.934, p < 0.0001. (B) Changes in gA index between two adjacent sessions. Activity threshold = 0.04 Hz. n = 90 session pairs, r = 0.929, p < 0.0001. , (C) Changes in gA index between two adjacent sessions. Activity threshold = 0.08 Hz. n = 90 session pairs, r = 0.772, p < 0.0001.