Ligand-independent modulation of GIPR signaling by splice variants

Kaini Hang[1,2,3,4,†], Lijun Shao[2,3,†], Qingtong Zhou[5], Fenghui Zhao[6], Antao Dai[6], Xiaoqing Cai[6], Raymond C. Stevens[1,2], Dehua Yang[3,6,7,8,*], Ming-Wei Wang[2,3,5,6,7,8,*]

[1]iHuman Institute, ShanghaiTech University, Shanghai 201210, China.

[2]School of Life Science and Technology, ShanghaiTech University, Shanghai 201210, China

[3]University of Chinese Academy of Sciences, Beijing 100049, China.

[4]CAS Center for Excellence in Molecular Cell Science, Shanghai Institute of Biochemistry and Cell Biology, Shanghai 200031, China.

[5]Department of Pharmacology, School of Basic Medical Sciences, Fudan University, Shanghai 200032, China.

[6]The National Center for Drug Screening, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China.

[7]The CAS Key Laboratory of Receptor Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China.

[8]Research Center for Deepsea Bioresources, Sanya, Hainan 572025, China.

†These authors contributed equally to this work.

*Corresponding author. Email: dhyang@simm.ac.cn (D.Y.); mwwang@simm.ac.cn (M.-W.W.)

Abstract

Glucose-dependent insulinotropic polypeptide receptor (GIPR) is a potential drug target for metabolic disorders. It works with glucagon-like peptide-1 receptor (GLP-1R) and glucagon receptor (GCGR) in humans to maintain glucose homeostasis. Unlike the other two receptors, GIPR has at least 7 reported (EMBL-EBI, 2022; NCBI, 2022a, 2022b) splice variants (SVs) with previously undefined functions. To explore their roles in endogenous peptide mediated GIPR signaling, we investigated the outcome of co-expressing each of the four SVs in question with GIPR in terms of ligand binding, cAMP accumulation, G_s activation, β-arrestin recruitment and cell surface localization. The effects of these SVs on intracellular cAMP responses

modulated by receptor activity-modifying proteins (RAMPs) were also studied. It was

found that while SVs of GIPR neither bound to the hormone nor affected its signal

transduction *per se*, they differentially regulated GIPR-mediated cAMP and β-arrestin

responses. Specifically, SV1 and SV4 were preferable to G_s signaling, SV3 was

biased towards β-arrestin recruitment, whereas SV2 was inactive on both pathways. In

the presence of RAMPs, only SV1 and SV4 synergized the repressive action of

RAMP3 on GIP-elicited cAMP production. The results suggest a new form of signal

bias that is constitutive and ligand-independent, thereby expanding our knowledge of

biased signaling beyond pharmacological manipulation (*i.e.*, ligand specific) as well

as constitutive and ligand-dependent (*e.g.*, SV1 of the growth hormone-releasing

hormone receptor).

Introduction

G protein-coupled receptors (GPCRs) represent the largest family of membrane

proteins that are universally expressed in human tissues (Pavlos & Friedman, 2017).

They can recognize a diverse range of extracellular ligands and transduce signals to

intracellular coupling partners, thereby governing crucial physiological functions

(Strange, 2008). GPCR-mediated signaling and pharmacological activities could be

profoundly affected by alternative splicing, leading to functional diversity (Furness,

Wootten, Christopoulos, & Sexton, 2012; Marti-Solano et al., 2020).

Splice variants (SVs) have been observed in many GPCRs, in which sequence

variations may include N terminus truncation or/and substitution, C terminus

truncation or/and substitution, intracellular/extracellular loop differences, severe

truncation leading to variants with less than 7 transmembrane domains (TMDs) or

soluble variants (Markovic & Challiss, 2009). In general, N terminus variations

impair ligand binding properties, such as corticotropin releasing hormone receptor 1

(CRH1R) and 2 (CRHR2) (Evans & Seasholtz, 2009), calcitonin receptor (CTR) (Nag,

Sultana, Kato, & Hirose, 2007) and parathyroid hormone 1 receptor (PTH1R) (Joun et

al., 1997), while C terminus variations show altered signaling or protein interactions,

such as metabotropic glutamate receptors (mGluRs) (Cai, Schools, & Kimelberg,

2000), μ-opioid receptor (MOR) (Lu et al., 2015) and 5-hydroxytryptamine (5-HT)

receptors (Coupar, Desmond, & Irving, 2007). Intracellular loop (ICL) differences

affect G protein coupling preference and extracellular loop (ECL) differences alter

ligand specificity and binding kinetics, as illustrated by pituitary adenylate

cyclase-activating polypeptide (PACAP) type 1 receptor (PAC1R) (McCulloch et al.,

2001) and D3 dopamine receptor (D3R) (Karpa, Lin, Kabbani, & Levenson, 2000;

Richtand, 2006), respectively. Variants with less than 7 TMDs caused by severe N

terminus truncation exhibit negative effect on wild-type (WT) receptor signaling

(Markovic & Challiss, 2009).

Glucose-dependent insulinotropic polypeptide receptor (GIPR) belongs to class

B1 subfamily of GPCRs and is present in pancreatic cells, adipose tissues and

osteoblasts. Upon GIP stimulation, it regulates insulin secretion, fat accumulation and

bone formation by increasing intracellular adenosine 3,5-cyclic monophosphate

(cAMP) levels (Campbell, 2021; Yabe & Seino, 2011; Yue & Lam, 2019). GIPR is

reported to have a truncated SV showing a dominant negative effect on the

translocation of WT GIPR from the endoplasmic reticulum (ER) to the cell surface,

leading to a decreased activity (Harada et al., 2008). However, the functionalities of

GIPR SVs remain to be defined.

In this study, we constructed and expressed four representative GIPR SVs to

elaborate their biological properties on GIPR-mediated cAMP accumulation and

β-arrestin recruitment. They were selected based on expression levels and splicing

modes: SV1, SV2, SV3 and SV4 with residue lengths of 419, 430, 405 and 265,

respectively (Figure 1). SV1 has a truncated C terminus and a 20 amino acid

substitution. SV2 lacks the sequence of residues 58-93 at the N terminus. SV3 has a

replaced N terminus of residues 1-93. SV4 only has 3 TMDs.

Results

Splice variants neither bind nor affect GIP-induced cAMP response

We first expressed GIPR and four SVs separately in HEK293T cells to investigate

their ability to bind GIP_{1-42} and elicit cAMP accumulation and β-arrestin recruitment.

Figure 2 shows that none of the SVs displayed any ligand-binding and signaling

properties, whereas the WT receptor was highly active in each parameter analyzed.

Splice variants differentially modulate GIPR-mediated signaling

Since SVs are usually express in cells and tissues where GIPR are present (GTEx, 2022; Harada et al., 2008), we co-transfected GIPR with each SV in order to study if they influence the signaling profile of the WT receptor. While the binding affinity of GIP_{1-42} to the cognate receptor were significantly reduced by 0.44, 0.54, 0.50 and 0.74 folds for SV1, SV2, SV3 and SV4, respectively (Figure 3A and Table 1), cAMP and β-arrestin responses at GIPR were differentially and negatively modulated. Both SV1 and SV4 decreased cAMP signaling but the effect of SV4 was nearly 5-fold stronger than that of SV1, SV2 and SV3 were inactive (Figure 3B and Table 1). Although SV1, SV3 and SV4 decreased the E_{max} values of β-arrestin 2 recruitment by 0.59, 0.49 and 0.42-folds, respectively, and SV2 remained inactive (Figure 3D and Table 1), none of them influenced on β-arrestin 1 recruitment (Figure 3C).

GIPR and splice variants are co-localized on the membrane

GIPR and SVs could be localized either on the membrane or the cytoplasm of transfected HEK293T cells (Figure 4A and 4B). Figure 4C illustrates that GIPR, SV1 and SV4 were co-expressed mostly on the cell surface, whereas SV2 and SV3 only exhibited a partial membrane co-localization. Upon co-transfection with GIPR, most of SV3 were translocated to the membrane (3rd panel of Figure 4C), but SV2 remained in the cytoplasm along with redistributed GIPR (2nd panel of Figure 4C), consistent with the silent role of SV2 observed.

Synergistic effect exerted by splice variants and RAMP3

Receptor activity-modifying protein 3 (RAMP3) was reported to reduce GIP_{1-42} induced cAMP accumulation at GIPR as opposed to RAMPs 1 and 2 that showed no effect (Harris, Mackie, Pawlak, Carvalho, & Ladds, 2021; Shao et al., 2021). After co-expression of individual SVs with GIPR and each RAMP, no alteration was noted with RAMP1 and RAMP2 (Figure 5A and 5B), but the suppression of GIPR-mediated cAMP production by RAMP3 was moderately augmented by SV1 and SV4 (with EC_{50} increased by 0.54 and 0.96-fold, respectively) (Figure 5C and Table 2).

SV1 and SV4 are repressive on G_s activation

We also studied the effect of SVs on G_s protein coupling by GIPR. G_s activation was

assessed using a split luciferase NanoBiT G protein sensor. Individually expressing

SVs showed no ability to active G_s (Figure 6A), consistent with their lack of cAMP

signaling (Figure 2B). SV1 and SV4 markedly impaired G_s coupling with 4.89- and

2.68-fold increased EC_{50} values, respectively (Figure 6B and Table 1). Although the P

values of pEC_{50} for SV1 and SV4 were greater than 0.05 probably due to inherent

assay variations, the difference between their EC_{50} and that of GIPR alone was

statistically significant (Figure 6C and D), thus in line with their repressive action on

cAMP accumulation (Figure 3B).

Diminished interaction between splice variants and signaling partners

As shown in Figure 7A-C, the helix 8 (H8) of SV1 adopted a distinct conformation

from that of GIPR during molecular dynamics (MD) simulations, bent upwards and

moved away from Gβ, thereby resulting in a reduced receptor-Gβ interface area. Of

note, the specific residue in GIPR H8 (such as R405) stabilized the Gβ binding, which

was absent in SV1, consistent with its role in G_s-mediated signaling. Compared to

GIPR, obvious differences in peptide-binding and β-arrestin 1 interface were observed

for SV3 (Figure 7D-F). By replacing the GIPR extracellular domain (ECD) with a

smaller domain (61 fewer residues), SV3 reorganized its extracellular half including

ECL1 to accommodate peptide binding with a smaller peptide-receptor interface

(from 3,280 $Å^2$ in the last 500 ns MD simulation of GIPR to 2,826 $Å^2$ in that of SV1).

As far as the intracellular half is concerned, β-arrestin 1 inserted deeper to the SV3

core compared with that of GIPR (Figure 7E). These different structural and dynamics

features between SV3 and GIPR highlight their distinct signaling properties.

Discussion

GIP, glucagon-like peptide-1 (GLP-1) and glucagon (GCG) together play a pivotal

role in glucose homeostasis mediated via their respective receptors (Cho, Merchant, &

Kieffer, 2012; Sekar, Singh, Arokiaraj, & Chow, 2016). GCG increases the release of

glucose, while GIP and GLP-1R work synergistically to cause postprandial insulin

secretion, regulate glucagon secretion, stimulate β cell proliferation and protect it

from apoptosis (Alexiadou, Anyiam, & Tan, 2019; Estall & Drucker, 2006; Hansotia

& Drucker, 2005; Seino, Fukushima, & Yabe, 2010; Skow, Bergmann, & Knop,

2016). Of note is that GIP promotes the release of both insulin and glucagon

(Gasbjerg et al., 2018) thereby modulating the action of GLP-1 and GCG on sugar

metabolism, probably involving some SVs of GIPR.

A common feature of the SVs examined is that they neither bind the native ligand,

GIP_{1-42}, nor elicit signal transduction. When co-expressed with WT GIPR, all of them

reduced peptide binding in a similar manner while displaying distinct signaling

profiles (Figure 8). Both SV1 and SV4 decreased GIPR-mediated cAMP and

β-arrestin 2 responses; SV3 selectively suppressed β-arrestin 2 recruitment, and SV2

had no effect on the two signaling events, but diminished GIPR presence in the

membrane. While SV1 (SV4 to less extent) may have the preference for activating the

G_s pathway, SV3 obviously is biased towards β-arrestin 2 recruitment.

Consistent with previous findings showing that SVs are capable of altering

signaling profiles compared to WT receptors (Kochman, 2014; Maggio et al., 2016),

our data suggest a constitutive biased mechanism different from signal bias caused by

various ligands. For example, SVs of the C-X-C chemokine receptor 3 (CXCR3)

could activate different signaling pathways through biased agonism (Berchiche &

Sakmar, 2016), and SV1 of the growth hormone–releasing hormone receptor

(GHRHR) preferentially transduces signals via β-arrestins while GHRHR

predominantly activates G_s proteins (Cong et al., 2021). However, unlike other GPCR

SVs, that of GIPR are incapacitated in terms of ligand-binding and signal transduction

per se, but negatively affect that of the WT receptor in a ligand-independent and

signaling biased manner.

Bidirectional regulation of carbohydrate levels by GIP_{1-42} is essential to the

maintenance of glucose homeostasis, although this hinders the development of

therapeutic agents targeting GIPR (Killion et al., 2018). It seems that such a unique

modulation of gut hormone actions is finely tuned by SVs with differentiated

functionalities: the repression of cAMP response imposed by RAMP3 could be

strengthened by SV1 and SV4, whereas β-arrestin 2 signaling is solely modified by

SV3. Unlike the other three SVs, SV2 appears as a sequester that redistributes the

membrane GIPR to the cytoplasm, evidenced by immunofluorescence staining when
both WT GIPR and SV2 are co-expressed. Whether this constitutes a shutdown
mechanism for GIPR function remains to be explored.

The roles of SVs in GIPR functioning are unique not only because they are of
repressive nature but also due to their synergistic actions with RAMP3 that itself is a
negative regulator of most members of the glucagon receptor subfamily of class B1
GPCRs (Shao et al., 2021). It remains elusive if the above described phenomenon
constitutes a "doubly insured" mechanism for signal modulation in order to fine tune
the action of GIP_{1-42}. Clearly both in-depth structural and biochemical studies are
required to solve the puzzle.

Materials and Methods

Key resource table

Reagent type (species) or resource	Designation	Source or reference	Identifier	Additional information
Transcript ID	WT GIPR	ENSEMBL	ENST00000 590918.6	
Transcript ID	Splice variant 1	ENSEMBL	ENST00000 263281.7	
Transcript ID	Splice variant 2	ENSEMBL	ENST00000 304207.12	
Transcript ID	Splice variant 3	ENSEMBL	ENST00000 652180.1	
Transcript ID	Splice variant 4	ENSEMBL	ENST00000 585889.1	
Cell line (*Homo sapiens*)	HEK293T	ATCC	Cat. No.: CRL-3216	
Cell line (hamster)	CHO-K1	ATCC	Cat. No.: CCL-61	
Recombinant DNA reagent	pcDNA3.1-GIPR (WT and splice variants)	This paper	N/A	
Recombinant DNA reagent	pcDNA3.1-GIPR (WT)-HA	doi: 10.1016/j.bcp.2020.1 14001	N/A	
Recombinant DNA reagent	pcDNA3.1-GIPR (splice variants) -FLAG	This paper	N/A	
Recombinant DNA reagent	pcDNA3.1-GIPR (WT and splice variants)-Rluc8 (Renilla	This paper	N/A	

	luciferase 8)			
Recombinant DNA reagent	Venus-β-arrestins 1 and 2	doi: 10.1016/j.bcp.2020.114001	N/A	
Recombinant DNA reagent	pcDNA3-RAMP3	doi: 10.1016/j.apsb.2021.07.028	N/A	
Peptide, recombinant protein	GIP$_{1-42}$	GenScript	N/A	
Chemical compound, drug	Bovine serum albumin (BSA)	ABCONE	Cat. No.: A23088-100G	
Chemical compound, drug	3-Isobutylene-1-methyl xanthine (IBMX)	ABCONE	Cat. No.: I72182-250MG	
Chemical compound, drug	Fetal bovine serum (FBS)	Gibco	Cat. No.: 10099-141	
Chemical compound, drug	Dulbecco's modified Eagle's medium (DMEM)	Gibco	Cat. No.: 12430-054	
Chemical compound, drug	Ham's F-12 nutrient mix (F12)	Gibco	Cat. No.: C11765500	
Chemical compound, drug	Hanks′ Balanced Salt solution (HBSS)	Gibco	Cat. No.: C14175500	
Chemical compound, drug	HEPES	Gibco	Cat. No.: 15630-080	
Chemical compound, drug	Sodium pyruvate	Gibco	Cat. No.: 11360-0'70	

Chemical compound, drug	Lipofectamine 2000 transfection reagent	Invitrogen	Cat. No.: 11668-019	
Chemical compound, drug	^{125}I-GIP	PerkinElmer	Cat. No.: NEX402010UC	
Chemical compound, drug	Coelenterazine H	Yeasen Biotech	Cat. No.: 40906ES02	
Antibody	Anti-FLAG primary antibody	Sigma-Aldrich	Cat. No.: F3165	
Antibody	Anti-HA primary antibody	Cell Signaling Technology	Cat. No.: 3724S	
Antibody	Anti-mouse Alexa Fluor 647 conjugated secondary antibody	Invitrogen	Cat. No.: A31571	
Antibody	Anti-rabbit Alexa Fluor 488 conjugated secondary antibody	Invitrogen	Cat. No.: A21206	
Commercial assay, kit	LANCE Ultra cAMP kit	PerkinElmer	Cat. No.: 2675984	
Software, algorithm	GraphPad Prism v8.4	GraphPad Software	N/A	https://www.graphpad.com/
Software, algorithm	FreeSASA	doi: 10.12688/f1000research.7931.1	N/A	http://freesasa.github.io/
Software, algorithm	Gromacs 2020.1	doi: 10.1016/j.softx.2015.06.001	N/A	https://manual.gromacs.org/2020.1/download.html
Software, algorithm	Protein Preparation Wizard	Schrödinger	N/A	https://www.schrodinger.com/products

Software, algorithm	CHARMM-GUI Membrane Builder v3.6	doi: 10.1002/jcc.23702	N/A	/protein-preparatio n-wizard
				https://charmm-gui. org/
Software, algorithm	CHARMM36-CAMP	doi: 10.1021/ct200328p	N/A	
Software, algorithm	LINCS algorithm	doi: 10.1021/ct700200b	N/A	
Software, algorithm	Semi-isotropic Parrinello-Rahman barostat	doi: 10.1016/0022-3093(93)90111-A	N/A	

Cell culture

HEK293T cells were maintained in DMEM (Gibco) supplemented with 10% FBS

(Gibco) and 100 mM sodium pyruvate (Gibco). CHO-K1 cells were maintained in F12

(Gibco) supplemented with 10% FBS. All cells were incubated in a humidified

environment at 37°C in 5% CO_2.

Construct

cDNAs were inserted into pcDNA3.1 vector by one-step cloning. Addition of FLAG-

and HA-tags to WT GIPR or SVs was carried out by site-directed mutagenesis. WT

GIPR or SVs were cloned to the backbone of Rluc8 at the C terminus. All constructs

were confirmed by DNA sequencing (GENEWIZ, Suzhou, China). To optimizing the

co-transfection assays, three GIPR *vs*. SV ratios (1:1, 1:3 and 1:6) were tried. Since

the impact of 1:1 on GIPR activity was hard to observe and that of 1:3 and 1.6 was

similar, we selected 1:3 for the entire study.

cAMP accumulation assay

GIP_{1-42} stimulated cAMP accumulation was measured by a LANCE Ultra cAMP kit

(PerkinElmer). Cells were seeded onto 6-well cell culture plates and transiently

transfected with 4 μg plasmid using Lipofectamine 2000 transfection reagent

(Invitrogen). After 24 h culture, the transfected cells were seeded onto 384-well

microtiter plates at a density of 3,000 cells per well in HBSS (Gibco) supplemented with 5 mM HEPES (Gibco), 0.1% (w/v) bovine serum albumin (BSA) and 0.5 mM IBMX (Sigma-Aldrich). The cells were stimulated with different concentrations of GIP_{1-42} for 40 min at room temperature (RT). Eu and Ulight were then diluted by cAMP detection buffer and added to the plates separately to terminate the reaction. Plates were incubated at RT for 40 min and the fluorescence intensity measured at 620 nm and 650 nm by an EnVision multilabel plate reader (PerkinElmer).

Whole-cell binding assay

CHO-K1 cells were seeded at a density of 30,000 cells/well in Isoplate-96 plates (PerkinElmer). The WT GIPR or SVs were transiently transfected using Lipofectamine 2000 transfection reagent. Twenty-four hours after transfection, cells were washed twice, and incubated with blocking buffer (F12 supplemented with 33 mM HEPES and 0.1% BSA, pH 7.4) for 2 h at 37°C. For homogeneous binding, cells were incubated in binding buffer with a constant concentration of ^{125}I-GIP (40 pM, PerkinElmer) and increasing concentrations of unlabeled GIP_{1-42} (3.57 pM to 1 μM) at RT for 3 h. Following incubation, cells were washed three times with ice-cold PBS and lysed by addition of 50 μL lysis buffer (PBS supplemented with 20 mM Tris-HCl, 1% Triton X-100, pH 7.4). Fifty μL of scintillation cocktail (OptiPhase SuperMix, PerkinElmer) was added and the plates were subsequently counted for radioactivity (counts per minute, CPM) in a scintillation counter (MicroBeta2 Plate Counter, PerkinElmer).

β-arrestin 1/2 recruitment

HEK293T cells (3.5×10^6 cells/mL) were grown for 24 h before transfection with 4 μg plasmid containing a GIPR/SV-Rluc8:Venus-β-arrestin1/2 at ratio of 1:9, or a GIPR-Rluc8:SV:Venus-β-arrestin1/2 at a ratio of 1:3:9. Transiently transfected cells were then seeded onto poly-D-lysine coated 96-well culture plates (50,000 cells/well) in DMEM with 10% FBS. Cells were grown overnight before incubation in assay buffer (HBSS supplemented with 10 mM HEPES and 0.1% BSA, pH 7.4) for 30 min at 37°C. Coelentrazine-h (Yeasen Biotech) was added to a final concentration of 5 μM for 5 min before bioluminescence resonance energy transfer (BRET) readings were

made using an EnVision plate reader. BRET baseline measurements were collected

for 15 cycles prior to ligand addition. Following peptide addition, BRET was

measured for 55 cycles. The BRET signal (ratio of 535 nm over 470 nm emission)

was corrected to the baseline and then vehicle-treated condition to determine

ligand-induced changes in BRET response. Concentration-response values were

obtained from the area-under-the-curve (AUC) of the responses elicited by GIP_{1-42}.

Immunofluorescence staining

HEK293T cells were seeded in 6-well plates and transfected with 4 μg plasmid

containing GIPR-HA or/and SV-FLAG. After 24 h, cells were collected and reseeded

in 96-well plates until they reached 50%~70% confluence. Cells were washed with

PBS before fixation with 4% paraformaldehyde for 15 min. Then they were washed

three more times and blocked with 5% BSA plus 0.1% Triton X-100 for 1 h at RT.

Rabbit anti-HA primary antibody (diluted 1:500) or/and mouse anti-FLAG primary

antibody (diluted 1:300) were diluted with incubation buffer (PBS supplemented with

5% BSA) for 1 h followed by 3-time wash. Cells were reacted with 200 μL interaction

buffer containing donkey anti-rabbit Alexa 488-conjugated secondary antibody or/and

donkey anti-mouse Alexa 647-conjugated secondary antibody (diluted 1:1000) at RT

for 1 h in the dark. After final washing, nuclei were stained with Hoechst 33258 for 5

min. Cells were imaged using a high-resolution microscope DeltaVision™ Ultra (GE

Healthcare, Boston, USA).

G protein NanoBiT assay

HEK293T cells (3.5×10^6 cells/mL) were grown for 24 h to reach 70% to 80%

confluence. Then the cells were transiently transfected with GIPR, $G\alpha_s$-LgBiT, Gβ1,

and Gγ2-SmBiT at a 2:1:5:5 mass ratio, or GIPR, SV, $G\alpha_s$-LgBiT, Gβ1, and

Gγ2-SmBiT at a 2:6:1:5:5 mass ratio. Twenty-four hours after transfection, cells were

seeded into poly-D-lysine coated 96-well culture plates at a density of 50,000 cells per

well in DMEM with 10% FBS. Cells were grown overnight before incubation in

HBSS buffer (pH 7.4) supplemented with 0.1% BSA and 10 mM HEPES for 30 mins

at 37°C (no CO_2). They were then reacted with coelenterazine H (5 μM) for 2 h at RT.

Luminescence signals were measured using an EnVision plate reader at 15-s intervals

(25°C). Briefly, following the baseline reading for 3.5 min, GIP_{1-42} was added, and

the reading continued for 13.5 min. Data were corrected to baseline and

vehicle-treated samples. The area-under-the-curve (AUC) across the time-course

response curve was determined and normalized to the WT GIPR, which was set to

100%.

Molecular dynamics simulation

Molecular dynamic simulations were performed by Gromacs 2020.1. The homology

models of SV1 and SV3 were generated using the cryo-EM structure of the full-length

GIPR (PDB code: 7DTY) (Zhao et al., 2021) and the X-ray structure of the GIPR

ECD (PDB code: 2QKH) (Parthier et al., 2007) as templates. All peptide–receptor–G_s

complexes were built based on the cryo-EM structure of the GIP–GIPR–G_s complex

(PDB code: 7DTY) (Zhao et al., 2021) and prepared by the Protein Preparation

Wizard (Schrodinger 2017-4) with the Nb35 nanobody removed. To build the model

of peptide–receptor–β-arrestin 1 complex, the receptor in complex with both peptide

and G_s was aligned to the published β-arrestin 1–bound $β_1AR$ structure (PDB code:

6TKO) (Lee et al., 2020). The receptor chain termini were capped with acetyl and

methylamide. The residues G2 and C3 of $Gα_s$ were *N*-myristoylated and palmitoylated,

respectively (Kato et al., 2019). All missing backbone and side chains were modeled

using Prime (Schrodinger 2017-4) and the titratable residues were left in their

dominant state at pH 7.0. To build MD simulation systems, the complexes were

embedded in a bilayer composed of 254~315 POPC lipids and solvated with 0.15 M

NaCl in explicit TIP3P waters using CHARMM-GUI Membrane Builder v3.6 (Wu et

al., 2014). The CHARMM36-CAMP force filed (Guvench et al., 2011) was adopted

for protein, peptides, lipids and salt ions. The Particle Mesh Ewald (PME) method

was used to treat all electrostatic interactions beyond a cut-off of 12 Å and the bonds

involving hydrogen atoms were constrained using LINCS algorithm (Hess, 2008).

The complex system was first relaxed using the steepest descent energy minimization,

followed by slow heating of the system to 310 K with restraints. The restraints were

reduced gradually over 50 ns. Finally, restrain-free production run was carried out for

each simulation, with a time step of 2 fs in the NPT ensemble at 310 K and 1 bar

using the Nose-Hoover thermostat and the semi-isotropic Parrinello-Rahman barostat (Aoki & Yonezawa, 1992), respectively. The buried interface areas were calculated with FreeSASA using the Sharke-Rupley algorithm with a probe radius of 1.2 Å (Mitternacht, 2016).

Statistical analysis

Statistical analysis was performed using GraphPad Prism 8.4 (GraphPad Software). For signaling assays, data of individual experiments were normalized to the maximum responses in cells expressing only the WT GIPR. Non-linear curve fit was performed using a three-parameter logistic equation (log (agonist *vs*. response)). All data are presented as means ± SEM. Significant differences were determined by one-way ANOVA with Dunnett's test. For co-localization analysis, Pearson's correlation coefficients (r) were performed using the co-localization threshold plugin of ImageJ. Five separate Regions of Interest (ROI) were selected and means ± SEM were determined.

References

Alexiadou, K., Anyiam, O., & Tan, T. (2019). Cracking the combination: Gut hormones for the treatment of obesity and diabetes. *J Neuroendocrinol, 31*(5), e12664. doi:10.1111/jne.12664

Aoki, K. M., & Yonezawa, F. (1992). Constant-pressure molecular-dynamics simulations of the crystal-smectic transition in systems of soft parallel spherocylinders. *Physical Review A, 46*(10), 6541-6549. doi:10.1103/physreva.46.6541

Berchiche, Y. A., & Sakmar, T. P. (2016). CXC Chemokine Receptor 3 Alternative Splice Variants Selectively Activate Different Signaling Pathways. *Mol Pharmacol, 90*(4), 483-495. doi:10.1124/mol.116.105502

Cai, Z., Schools, G. P., & Kimelberg, H. K. (2000). Metabotropic glutamate receptors in acutely isolated hippocampal astrocytes: developmental changes of mGluR5 mRNA and functional expression. *Glia, 29*(1), 70-80. doi:10.1002/(sici)1098-1136(20000101)29:1<70::aid-glia7>3.0.co;2-v

Campbell, J. E. (2021). Targeting the GIPR for obesity: To agonize or antagonize? Potential mechanisms. *Mol Metab, 46*, 101139. doi:10.1016/j.molmet.2020.101139

Cho, Y. M., Merchant, C. E., & Kieffer, T. J. (2012). Targeting the glucagon receptor family for diabetes and obesity therapy. *Pharmacol Ther, 135*(3), 247-278. doi:10.1016/j.pharmthera.2012.05.009

Cong, Z., Zhou, F., Zhang, C., Zou, X., Zhang, H., Wang, Y., Zhou Q., Cai X., Liu Q., Li J., Shao L., Mao C., Wang X., Wu J., Xia T., Zhao L.-H., Jiang H., Zhang Y.,

Xu H. E., Cheng X., Yang D., Wang, M.-W. (2021). Constitutive signal bias mediated by the human GHRHR splice variant 1. *Proc Natl Acad Sci U S A, 118*(40). doi:10.1073/pnas.2106606118

Coupar, I. M., Desmond, P. V., & Irving, H. R. (2007). Human 5-HT(4) and 5-HT(7) receptor splice variants: are they important? *Curr Neuropharmacol, 5*(4), 224-231. doi:10.2174/157015907782793621

EMBL-EBI. (2022). Gene: GIPR ENSG00000010310 Structural variants. Ensembl release 105. Retrieved from https://asia.ensembl.org/Homo_sapiens/Gene/StructuralVariation_Gene?db=core;g=ENSG00000010310;r=19:45668221-45683722

Estall, J. L., & Drucker, D. J. (2006). Glucagon and glucagon-like peptide receptors as drug targets. *Curr Pharm Des, 12*(14), 1731-1750. doi:10.2174/138161206776873671

Evans, R. T., & Seasholtz, A. F. (2009). Soluble corticotropin-releasing hormone receptor 2alpha splice variant is efficiently translated but not trafficked for secretion. *Endocrinology, 150*(9), 4191-4202. doi:10.1210/en.2009-0285

Furness, S. G., Wootten, D., Christopoulos, A., & Sexton, P. M. (2012). Consequences of splice variation on Secretin family G protein-coupled receptor function. *Br J Pharmacol, 166*(1), 98-109. doi:10.1111/j.1476-5381.2011.01571.x

Gasbjerg, L. S., Gabe, M. B. N., Hartmann, B., Christensen, M. B., Knop, F. K., Holst, J. J., & Rosenkilde, M. M. (2018). Glucose-dependent insulinotropic polypeptide (GIP) receptor antagonists as anti-diabetic agents. *Peptides, 100*, 173-181. doi:10.1016/j.peptides.2017.11.021

GTEx. (2022). Isoform Expression of GIPR: ENSG00000010310.8 gastric inhibitory polypeptide receptor [Source:HGNC Symbol;Acc:HGNC:4271]. (dbGaP Accession phs000424.v8.p2). Retrieved from https://gtexportal.org/home/gene/GIPR#gene-transcript-browser-block

Guvench, O., Mallajosyula, S. S., Raman, E. P., Hatcher, E., Vanommeslaeghe, K., Foster, T. J., Jamison F. W. 2nd, Mackerell, A. D., Jr. (2011). CHARMM additive all-atom force field for carbohydrate derivatives and its utility in polysaccharide and carbohydrate-protein modeling. *J Chem Theory Comput, 7*(10), 3162-3180. doi:10.1021/ct200328p

Hansotia, T., & Drucker, D. J. (2005). GIP and GLP-1 as incretin hormones: lessons from single and double incretin receptor knockout mice. *Regul Pept, 128*(2), 125-134. doi:10.1016/j.regpep.2004.07.019

Harada, N., Yamada, Y., Tsukiyama, K., Yamada, C., Nakamura, Y., Mukai, E., Hamasaki A., Liu X., Toyoda K., Seino Y., Inagaki, N. (2008). A novel GIP receptor splice variant influences GIP sensitivity of pancreatic beta-cells in obese mice. *Am J Physiol Endocrinol Metab, 294*(1), E61-68. doi:10.1152/ajpendo.00358.2007

Harris, M., Mackie, D. I., Pawlak, J. B., Carvalho, S., & Ladds, G. (2021). RAMPs regulate signalling bias and internalisation of the GIPR.

Hess, B. (2008). P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation. *J Chem Theory Comput, 4*(1), 116-122. doi:10.1021/ct700200b

Joun, H., Lanske, B., Karperien, M., Qian, F., Defize, L., & Abou-Samra, A. (1997). Tissue-specific transcription start sites and alternative splicing of the parathyroid hormone (PTH)/PTH-related peptide (PTHrP) receptor gene: a new PTH/PTHrP receptor splice variant that lacks the signal peptide. *Endocrinology, 138*(4), 1742-1749. doi:10.1210/endo.138.4.5085

Karpa, K. D., Lin, R., Kabbani, N., & Levenson, R. (2000). The dopamine D3 receptor interacts with itself and the truncated D3 splice variant d3nf: D3-D3nf interaction causes mislocalization of D3 receptors. *Mol Pharmacol, 58*(4), 677-683. doi:10.1124/mol.58.4.677

Kato, H. E., Zhang, Y., Hu, H., Suomivuori, C. M., Kadji, F. M. N., Aoki, J., Krishna Kumar K., Fonseca R., Hilger D., Huang W., Latorraca N. R., Inoue A., Dror R. O., Kobilka B. K., Skiniotis, G. (2019). Conformational transitions of a neurotensin receptor 1-Gi1 complex. *Nature, 572*(7767), 80-85. doi:10.1038/s41586-019-1337-6

Killion, E. A., Wang, J., Yie, J., Shi, S. D., Bates, D., Min, X., Komorowski R., Hager T., Deng L., Atangan L., Lu S.-C, Kurzeja R. J. M, Sivits G., Lin J., Chen Q., Wang Z., Thibault S. A., Abbott C. M., Meng T., Clavette B., Murawsky C. M., Foltz I. N., Rottman J. B., Hale C., Véniant M. M., Lloyd, D. J. (2018). Anti-obesity effects of GIPR antagonists alone and in combination with GLP-1R agonists in preclinical models. *Sci Transl Med, 10*(472). doi:10.1126/scitranslmed.aat3392

Kochman, K. (2014). Superfamily of G-protein coupled receptors (GPCRs)--extraordinary and outstanding success of evolution. *Postepy Hig Med Dosw (Online), 68*, 1225-1237. doi:10.5604/17322693.1127326

Lee, Y., Warne, T., Nehme, R., Pandey, S., Dwivedi-Agnihotri, H., Chaturvedi, M., Edwards P. C., García-Nafría J., Leslie A. G. W., Shukla A. K., Tate, C. G. (2020). Molecular basis of beta-arrestin coupling to formoterol-bound beta1-adrenoceptor. *Nature, 583*(7818), 862-866. doi:10.1038/s41586-020-2419-1

Lu, Z., Xu, J., Rossi, G. C., Majumdar, S., Pasternak, G. W., & Pan, Y. X. (2015). Mediation of opioid analgesia by a truncated 6-transmembrane GPCR. *J Clin Invest, 125*(7), 2626-2630. doi:10.1172/JCI81070

Maggio, R., Fasciani, I., Rossi, M., Di Gregorio, J., Pietrantoni, I., Puca, V., Flati V., Scarselli, M. (2016). Variants of G protein-coupled receptors: a reappraisal of their role in receptor regulation. *Biochem Soc Trans, 44*(2), 589-594. doi:10.1042/BST20150239

Markovic, D., & Challiss, R. A. (2009). Alternative splicing of G protein-coupled receptors: physiology and pathophysiology. *Cell Mol Life Sci, 66*(20), 3337-3352. doi:10.1007/s00018-009-0093-4

Marti-Solano, M., Crilly, S. E., Malinverni, D., Munk, C., Harris, M., Pearce, A., Quon T., Mackenzie A. E., Wang X., Peng J., Tobin A. B/, Ladds G., Milligan G., Gloriam D. E., Puthenveedu M. A., Babu, M. M. (2020). Combinatorial expression of GPCR isoforms affects signalling and drug responses. *Nature, 587*(7835), 650-656. doi:10.1038/s41586-020-2888-2

McCulloch, D. A., Lutz, E. M., Johnson, M. S., Robertson, D. N., MacKenzie, C. J., Holland, P. J., & Mitchell, R. (2001). ADP-ribosylation factor-dependent phospholipase D activation by VPAC receptors and a PAC(1) receptor splice variant. *Mol Pharmacol, 59*(6), 1523-1532. doi:10.1124/mol.59.6.1523

Mitternacht, S. (2016). FreeSASA: An open source C library for solvent accessible surface area calculations. *F1000Res, 5*, 189. doi:10.12688/f1000research.7931.1

Nag, K., Sultana, N., Kato, A., & Hirose, S. (2007). Headless splice variant acting as dominant negative calcitonin receptor. *Biochem Biophys Res Commun, 362*(4), 1037-1043. doi:10.1016/j.bbrc.2007.08.107

NCBI. (2022a). Gastric inhibitory polypeptide receptor isoform X6 [Homo sapiens]. v109.20211119. Retrieved from https://www.ncbi.nlm.nih.gov/protein/XP_016882074.1

NCBI. (2022b). Gastric inhibitory polypeptide receptor isoform X9 [Homo sapiens]. v109.20211119. Retrieved from https://www.ncbi.nlm.nih.gov/protein/XP_011525018.1

Parthier, C., Kleinschmidt, M., Neumann, P., Rudolph, R., Manhart, S., Schlenzig, D., Fanghänel J., Rahfeld J.-U., Demuth H.-U., Stubbs, M. T. (2007). Crystal structure of the incretin-bound extracellular domain of a G protein-coupled receptor. *Proc Natl Acad Sci U S A, 104*(35), 13942-13947. doi:10.1073/pnas.0706404104

Pavlos, N. J., & Friedman, P. A. (2017). GPCR Signaling and Trafficking: The Long and Short of It. *Trends Endocrinol Metab, 28*(3), 213-226. doi:10.1016/j.tem.2016.10.007

Richtand, N. M. (2006). Behavioral sensitization, alternative splicing, and d3 dopamine receptor-mediated inhibitory function. *Neuropsychopharmacology, 31*(11), 2368-2375. doi:10.1038/sj.npp.1301163

Seino, Y., Fukushima, M., & Yabe, D. (2010). GIP and GLP-1, the two incretin hormones: Similarities and differences. *J Diabetes Investig, 1*(1-2), 8-23. doi:10.1111/j.2040-1124.2010.00022.x

Sekar, R., Singh, K., Arokiaraj, A. W., & Chow, B. K. (2016). Pharmacological Actions of Glucagon-Like Peptide-1, Gastric Inhibitory Polypeptide, and Glucagon. *Int Rev Cell Mol Biol, 326*, 279-341. doi:10.1016/bs.ircmb.2016.05.002

Shao, L., Chen, Y., Zhang, S., Zhang, Z., Cao, Y., Yang, D., & Wang, M.-W. (2021). Modulating effects of RAMPs on signaling profiles of the glucagon receptor family. *Acta Pharm Sin B*. doi:10.1016/j.apsb.2021.07.028

Skow, M. A., Bergmann, N. C., & Knop, F. K. (2016). Diabetes and obesity treatment based on dual incretin receptor activation: 'twincretins'. *Diabetes Obes Metab, 18*(9), 847-854. doi:10.1111/dom.12685

Strange, P. G. (2008). Signaling mechanisms of GPCR ligands. *Curr Opin Drug Discov Devel, 11*(2), 196-202.

Wu, E. L., Cheng, X., Jo, S., Rui, H., Song, K. C., Davila-Contreras, E. M., Qi Y., Lee J., Monje-Galvan V., Venable R. M., Klauda J. B., Im, W. (2014).

CHARMM-GUI Membrane Builder toward realistic biological membrane simulations. *J Comput Chem, 35*(27), 1997-2004. doi:10.1002/jcc.23702

Yabe, D., & Seino, Y. (2011). Two incretin hormones GLP-1 and GIP: comparison of their actions in insulin secretion and beta cell preservation. *Prog Biophys Mol Biol, 107*(2), 248-256. doi:10.1016/j.pbiomolbio.2011.07.010

Yue, J. T. Y., & Lam, T. K. T. (2019). Antiobesogenic effects of central GIPR antagonism. *J Clin Invest, 129*(9), 3532-3535. doi:10.1172/JCI130755

Zhao, F., Zhang, C., Zhou, Q., Hang, K., Zou, X., Chen, Y., Wu F., Rao Q., Dai A., Yin W., Shen D.-D., Zhang Y., Xia T., Stevens R. C., Xu H. E., Yang D., Zhao L., Wang, M.-W. (2021). Structural insights into hormone recognition by the human glucose-dependent insulinotropic polypeptide receptor. *Elife, 10*, e68719. doi:10.7554/eLife.68719

Acknowledgments

We are grateful to Dr. Zhaotong Cong, Ms. Yan Chen and Mr. Chao Zhang for technical assistance. This work was partially supported by National Natural Science Foundation of China 81872915 (M.-W.W.), 82073904 (M.-W.W.), 82121005 (D.Y.), 81973373 (D.Y.) and 21704064 (Q.Z.); National Science & Technology Major Project of China–Key New Drug Creation and Manufacturing Program 2018ZX09735–001 (M.-W.W.) and 2018ZX09711002–002–005 (D.Y.); National Science & Technology Major Project of China – Innovation 2030 for Brain Science and Brain-Inspired Technology 2021ZD0203400 (Q.Z.); the National Key Basic Research Program of China 2018YFA0507000 (M.-W.W.); Novo Nordisk-CAS Research Fund grant NNCAS-2017–1-CC (D.Y.) and SA-SIBS Scholarship Program (D.Y.).

Author contributions: D.Y. and M.-W.W. designed research; K.H., L.S., F.Z., A.D., and X.C. performed research; K.H., L.S., Q.Z., R.C.S. and D.Y. analyzed data; and K.H., L.S., and M.-W.W. wrote the paper.

Competing interests: Authors declare that they have no competing interests.

Data availability: All data generated or analyzed during this study are included in the manuscript. Source data files have been provided for Figures 2-6.

Figures

Figure 1. Schematic diagram of GIPR and its splice variant (SV) constructs. **A**,
Construct of the wild-type (WT) GIPR. **B**, Construct of SV1. Residues 400-466 are
replaced by a 20-amino acid sequence (GRDPAAAPALWRQRGVRRRL). **C**,
Construct of SV2. Residues 58-93 are missing compared to that of WT. **D**, Construct
of SV3. Residues 1-93 are replaced by a 32-amino acid sequence
(MNSAHCNFRLPGSSDSPASASREAGITEAGIT). **E**, Construct of SV4. Residues
266-466 are missing and G265 is replaced by aspartic acid (D). **F**, Sequence
alignment of GIPR and the four SVs.

508
Figure 2. Ligand-binding and signaling profiles of GIPR and its splice variants (SVs).
A, Competitive inhibition of ^{125}I-GIP$_{1-42}$ binding to GIPR and SVs by unlabeled
GIP$_{1-42}$. Binding affinity is quantified by reduction of radioactivity (counts per minute,
CPM). **B**, Concentration-response curves of cAMP accumulation elicited by GIP$_{1-42}$ at
GIPR and SVs. **C and D**, β-arrestins 1 (β-arr1) and 2 (β-arr2) recruitment by GIPR
and SVs. Concentration-response characteristics are shown as the
area-under-the-curve (AUC) across the time-course response curve (0 to 10 min) for
each concentration. Data shown are means ± SEM of at least three independent
experiments (n=3-5) performed in quadruplicate (cAMP accumulation) or duplicate
(specific binding and β-arrestin recruitment). Signals were normalized to the
maximum (max) response of the wild-type (WT) GIPR and concentration-response
curves were analyzed using a three-parameter logistic equation.
The following figure supplements are available for figure 2:
Source data 1. Ligand-binding and signaling profiles of GIPR and its splice variants
(SVs).
524

Figure 3. Effects of GIPR splice variants (SVs) on ligand binding and the wild-type (WT) GIPR mediated signal transduction in HEK293T cells co-expressing GIPR and individual SVs. **A**, Effects of SVs on competitive binding of ^{125}I-GIP$_{1-42}$ to GIPR. **B**, Effects of SVs on GIP$_{1-42}$ induced cAMP accumulation at GIPR. **C and D**, Effects of SVs and GIPR on β-arrestins 1 (β-arr1) and 2 (β-arr2) recruitment by GIPR. Cells were co-transfected with GIPR and each SV at a 1:3 ratio. Data shown are means ± SEM of at least three independent experiments (n=3-5) performed in quadruplicate (cAMP accumulation) or duplicate (specific binding and β-arrestin recruitment). Signals were normalized to the maximum (max) response of GIPR and concentration-response curves were analyzed using a three-parameter logistic equation.

The following figure supplements are available for figure 3:

Source data 1. Effects of GIPR splice variants (SVs) on ligand binding and the wild-type (WT) GIPR mediated signal transduction in HEK293T cells co-expressing GIPR and individual SVs.

Figure 4. Co-localization of GIPR and its splice variants (SVs). Immunofluorescence
staining of HEK293T cells transfected with GIPR-HA (**A**) or each SV-FLAG (**B**)
alone. To estimate their co-localization, co-transfection of GIPR and individual SVs
(**C**) was performed at a ratio of 1:3 (green, GIPR-HA; red, SV-FLAG; yellow, merge).
Data show representative results from three independent experiments. Inset
demonstrates the overlapping positions of GIPR and SVs in the cell surface (SV1,
SV3 and SV4) or cytoplasm (SV2). Cells were observed by DeltaVision™ Ultra.
Scale bar = 10 μm.
The following figure supplements are available for figure 4:
Source data 1. Co-localization of GIPR and its splice variants (SVs).

Figure 5. Synergistic effects of splice variants (SVs) and receptor activity-modifying proteins (RAMPs) on GIPR-mediated cAMP signaling in HEK293T cells co-expressing GIPR, each SV and RAMP1 (**A**), RAMP2 (**B**) or RAMP3 (**C**). Signals were normalized to the maximum (max) response of the wild-type (WT) GIPR and data were fitted with a non-linear regression of three-parameter logistic curve. Data shown are means ± SEM of three independent experiments performed in quadruplicate.

The following figure supplements are available for figure 5:

Source data 1. Effects of splice variants (SVs) and receptor activity-modifying proteins (RAMPs) on GIPR-mediated cAMP signaling in HEK293T cells co-expressing GIPR, each SV and RAMP1.

	GIPR alone	GIPR+SV1	GIPR+SV2	GIPR+SV3	GIPR+SV4
pEC$_{50}$	9.66 $\pm$ 0.08	8.89 $\pm$ 0.24	9.39 $\pm$ 0.19	9.67 $\pm$ 0.21	9.10 $\pm$ 0.22
P value of pEC$_{50}$	-	0.0609	0.8366	0.9999	0.1435
ΔpEC$_{50}$	0	0.72 $\pm$ 0.13**	0.24 $\pm$ 0.06	0.05 $\pm$ 0.21	0.51 $\pm$ 0.09*
P value of ΔpEC$_{50}$	-	0.0026	0.5083	0.9956	0.0379

Figure 6. Effects of GIPR splice variants (SVs) on the wild-type (WT) GIPR mediated G$_s$ protein coupling in HEK293T cells co-expressing GIPR and individual SVs. **A**, GIP$_{1-42}$-induced G$_s$ coupling of individually expressed SVs and GIPR. Concentration-response curves are expressed as area-under-the-curve (AUC) across the time-course response curve (0 to 13.5 min) for each concentration and normalized to WT GIPR. **B**, Effects of SVs on GIP$_{1-42}$ induced G$_s$ protein coupling at GIPR. **C**, EC$_{50}$ differences of GIPR mediated G$_s$ protein coupling under the influence of SVs. **D**, G$_s$ protein coupling profiles of GIPR affected by SVs. Cells were co-transfected with GIPR and each SV at a 1:3 ratio. Data shown are means $\pm$ SEM of six independent experiments performed in duplicate. Signals were normalized to the maximum (max) response of GIPR and concentration-response curves were analyzed using a three-parameter logistic equation. *P < 0.05; **P < 0.01.

The following figure supplements are available for figure 6:

Source data 6. Effects of GIPR splice variants (SVs) on the wild-type (WT) GIPR mediated G$_s$ protein coupling in HEK293T cells co-expressing GIPR and individual SVs.

Figure 7. Molecular dynamics (MD) simulations of SV1 and SV3. **A,** Comparison of receptor-G_s conformation between the cryo-EM structure and final simulation snapshots of SV1 and GIPR. **B,** Comparison of the H8–Gβ interface between GIPR and SV1. **C,** Analysis of the MD simulation trajectories in (**A**): top, root mean square deviation (RMSD) of Cα positions of the receptor TMD, where all snapshots were superimposed on the cryo-EM structure of both GIP- and G_s-bound GIPR TMD (PDB code: 7DTY) using the Cα atom; upper middle, the buried surface area between receptor and Gβ, interface areas were calculated using freeSASA; lower middle, minimum distances between the charged atoms of D312^{Gβ} and R164^{ICL2} during MD simulations; bottom, minimum distances between the charged atoms of D312^{Gβ} and the side-chain atoms of R405^{H8} (GIPR) or A405^{H8} (SV1) during MD simulations. **D,** Comparison of the final MD snapshots between GIPR-β-arrestin 1 and SV3-β-arrestin 1. **E,** Different β-arrestin 1 modes between GIPR and SV3 according to MD simulations. **F,** Analysis of the MD simulation trajectories in (**D**): top, Cα RMSD of the receptor TMD, where all snapshots were superimposed on the cryo-EM structure of both GIP- and G_s-bound GIPR TMD (PDB code: 7DTY); upper middle, Cα RMSD calculation for β-arrestin 1, where all snapshots were superimposed on the cryo-EM structure of $β_1$AR-bound β-arrestin 1 (PDB code: 6TKO) using the Cα atom; lower middle, the buried surface area between receptor and peptide during MD simulations; bottom, the buried surface area between receptor and β-arrestin 1 during MD simulations.

	SV1	SV2	SV3	SV4
Binding (Span)	0.56***	0.46***	0.50***	0.26***
cAMP (EC_{50})	1.64***	1.09	0.95	4.79***
G_s (ΔEC_{50})	5.89**	1.88	0.98	3.68*
β-arr1 (E_{max})	1.28	0.90	0.97	0.97
β-arr2 (E_{max})	0.41***	0.96	0.51***	0.58***

▬ Negative modulation

Figure 8. Characterization of the effects of SVs on ligand binding and signaling profiles in HEK293T cells co-expressing GIPR and individual SVs. **A-D**, Schematic diagram representing the effects of SVs on ligand binding (**A**), cAMP signaling and G_s activation (**B**), β-arrestins 1 (β-arr1) (**C**) and 2 (β-arr2) recruitment (**D**). **E**, Ratio of parameters of ligand binding and signaling to GIPR alone. One-way ANOVA were used to determine statistical difference (*P< 0.05, **P< 0.01, ***P< 0.0001).

Table 1. Effects of splice variants (SVs) on ligand binding and GIPR-mediated signal transduction in HEK293T cells co-expressing GIPR and individual SVs.

	Ligand binding		cAMP accumulation		G_s coupling		β-arr1 recruitment		β-arr2 recruitment	
	$pIC_{50} \pm SEM$	Span (%) $\pm$ SEM	$pEC_{50} \pm SEM$	E_{max} (%) $\pm$ SEM	$pEC_{50} \pm SEM$	E_{max} (%) $\pm$ SEM	$pEC_{50} \pm SEM$	E_{max} (%) $\pm$ SEM	$pEC_{50} \pm SEM$	E_{max} (%) $\pm$ SEM
WT GIPR	7.91 ± 0.10	100.00 ± 4.87	11.14 ± 0.02	100.00 ± 0.63	9.66 ± 0.08	99.79 ± 2.20	7.94 ± 0.26	99.03 ± 6.83	8.19 ± 0.06	99.78 ± 1.81
GIPR+SV1	8.09 ± 0.09	54.21 ± 2.27*	10.92 ± 0.02*	100.08 ± 0.50	8.89 ± 0.24	91.36 ± 6.42	7.29 ± 0.52	126.65 ± 16.04	8.32 ± 0.35*	40.99 ± 4.17
GIPR+SV2	7.96 ± 0.11	46.52 ± 2.30*	11.10 ± 0.02	100.79 ± 0.52	9.39 ± 0.19	102.39 ± 5.10	7.67 ± 0.44	88.87 ± 9.90	8.21 ± 0.06	96.02 ± 1.69
GIPR+SV3	7.82 ± 0.13	51.01 ± 3.26*	11.16 ± 0.02	100.38 ± 0.43	9.67 ± 0.21	93.61 ± 5.31	7.70 ± 0.39	96.31 ± 8.45	8.22 ± 0.34*	50.62 ± 5.12
GIPR+SV4	8.15 ± 0.13	25.77 ± 1.59*	10.46 ± 0.02*	100.38 ± 0.68	9.10 ± 0.22	85.77 ± 5.40	8.08 ± 0.60	96.27 ± 10.78	8.13 ± 0.16*	58.17 ± 2.88

cAMP accumulation, G_s activation and β-arrestin 1/2 recruitment assays were performed in HEK293T cells. Whole cell binding assay was performed in CHO-K1 cells. All the measures were fitted to non-linear regression three-parameter logistic curves. pEC_{50} and pIC_{50} are the negative logarithm of the concentration of an agonist that gives a half of the maximum response. E_{max} and span values are the percentage (%) of the maximum response in cells expressing GIPR only. Data shown are means ± SEM of at least three independent experiments. One-way ANOVA was used to determine statistical difference (*P< 0.0001). β-arr1, β-arrestin 1; β-arr2, β-arrestin 2; WT, wild-type.

Table 2. Synergistic effects of splice variants (SVs) and receptor activity-modifying proteins (RAMPs) on GIPR-mediated cAMP signaling in HEK293T cells co-expressing GIPR.

	cAMP accumulation					
	RAMP1		RAMP2		RAMP3	
	$pEC_{50} \pm SEM$	E_{max} (%) $\pm$ SEM	$pEC_{50} \pm SEM$	E_{max} (%) $\pm$ SEM	$pEC_{50} \pm SEM$	E_{max} (%) $\pm$ SEM
GIPR	11.39 ± 0.03	100.00 ± 0.88	11.39 ± 0.03	100.00 ± 0.88	11.76 ± 0.03	100.00 ± 0.78
SV1	11.36 ± 0.04	100.00 ± 0.96	11.36 ± 0.04	100.00 ± 0.96	11.56 ± 0.03[b]	100.47 ± 0.67
SV2	11.42 ± 0.04	100.17 ± 1.11	11.42 ± 0.04	100.17 ± 1.11	11.64 ± 0.03[a]	100.40 ± 0.73
SV3	11.48 ± 0.04	99.89 ± 0.89	11.48 ± 0.04	99.89 ± 0.89	11.70 ± 0.03	100.08 ± 0.62
SV4	10.89 ± 0.04[b]	100.15 ± 1.18	10.89 ± 0.04[b]	100.15 ± 1.18	11.13 ± 0.03[b]	99.48 ± 0.67
GIPR alone	11.27 ± 0.04	98.71 ± 1.09	11.34 ± 0.04	98.64 ± 1.00	10.00 ± 0.03	100.45 ± 0.94
GIPR+SV1	11.27 ± 0.04	99.11 ± 1.04	11.26 ± 0.04	98.61 ± 1.17	9.81 ± 0.03[d]	100.40 ± 1.12
GIPR+SV2	11.31 ± 0.05	99.06 ± 1.28	11.31 ± 0.04	99.01 ± 1.09	9.84 ± 0.03[c]	100.83 ± 1.25
GIPR+SV3	11.38 ± 0.04	99.07 ± 0.98	11.42 ± 0.04	98.72 ± 0.93	9.96 ± 0.03	100.32 ± 0.84
GIPR+SV4	10.66 ± 0.04[e]	98.54 ± 1.27	10.81 ± 0.04[e]	97.94 ± 1.12	9.70 ± 0.02[e]	100.72 ± 0.82

cAMP accumulation was performed in HEK293T cells. Dose-response curves were analyzed using a three-parameter logistic equation to obtain pEC_{50} and E_{max}. Data shown are means $\pm$ SEM of at least three independent experiments. One-way ANOVA was used to determine statistical difference. [a], $P< 0.05$, [b], $P< 0.0001$ compared with GIPR. [c], $P< 0.01$, [d], $P<0.001$ and [e], $P<0.0001$ compared with GIPR alone.