Clinical and molecular correlates of CD39+ CD8 T cells in patients with lung cancer

Andrew Chow[1,2,3], Fathema Z. Uddin[1], Levi Mangarin[2], Hira Rizvi[4], Anton Dobrin[5,6],

Sam Tischfield[7], Alvaro Quintanal-Villalonga[1], Joseph M. Chan[1,3], Nisargbhai Shah[1], Viola Allaj[1],

Parvathy Manoj[1], Marissa Mattar[8], Maximiliano Meneses[8], Rebecca Landau[8], Mariana Ward[8],

Amanda Kulick[8], Charlene Kwong[8], Matthew Wierzbicki[8], Jessica Yavner[8], Shweta S. Chavan[7],

Abigail Farillas[1], Michael Liu[1], Aliya Holland[2], Metamia Ciampricotti[1],

Daniel Hirschhorn[2], Allison L Richards[7], Mark T.A. Donoghue[7], Glenn Heller[9],

Christopher A. Klebanoff[3,5,10,11,12], Matthew D. Hellmann[1,3,4,12,13], Elisa de Stanchina[8],

Triparna Sen[1], Jedd D. Wolchok[2,3,11,12,15], Taha Merghoub[2,3,11,12,15], Charles M. Rudin[1,3,4,14,15*]

[1]Thoracic Oncology Service, Department of Medicine, Memorial Sloan Kettering Cancer Center,

USA

[2]Ludwig Collaborative and Swim Across America Laboratory, Memorial Sloan Kettering Cancer

Center, New York, NY, USA

[3]Department of Medicine, Weill Cornell Medical College, USA

[4]Druckenmiler Center for Lung Cancer Research, Memorial Sloan Kettering Cancer Center, USA

[5]Center for Cell Engineering, Memorial Sloan Kettering Cancer Center, USA

[6]Louis V. Gerstner, Jr. Graduate School of Biomedical Sciences, Memorial Sloan Kettering

Cancer Center, USA

[7]Marie-Josée and Henry R. Kravis Center for Molecular Oncology, Memorial Sloan Kettering

Cancer Center, USA

[8]Antitumor Assessment Core, Memorial Sloan Kettering Cancer Center, USA

[9]Department of Epidemiology & Biostatistics, Memorial Sloan Kettering Cancer Center, USA

[10]Breast Medicine Service, Department of Medicine, Memorial Sloan Kettering Cancer Center,

USA

[11]Human Oncology and Pathogenesis Program, Memorial Sloan Kettering Cancer Center, USA

[12]Parker Institute for Cancer Immunotherapy, Memorial Sloan Kettering Cancer Center, USA

[13]Present Address: Oncology R&D, Astra-Zeneca, USA

[14]Molecular Pharmacology Program, Memorial Sloan Kettering Cancer Center, USA

[15]These senior authors contributed equally to this work.

*Lead Contact

Correspondence: merghout@mskcc.org (TM) and rudinc@mskcc.org (CMR)

HIGHLIGHTS

- Intratumoral CD39$^+$ CD8 T cells from patients with lung cancer demonstrate transcriptional and proteomic features of tumor reactivity.

- CD39 expression on CD8 T cells is poorly correlated with PD-L1, tumor mutation burden, and driver mutation in lung cancer.

- PD-1 axis blockade, but not cytotoxic chemotherapy, increases intratumoral CD39$^+$ CD8 T cells.

- High pre-treatment/early-on-treatment levels are associated with improved clinical outcomes from immune checkpoint blockade.

Background

Recent reports have identified CD39 as a marker of tumor-reactive CD8 T cells. Whether quantification of CD39$^+$ CD8 T cells as a proxy of endogenous anti-tumor reactivity yields information that is non-redundant to tumoral features that currently drive lung cancer management (e.g. PD-L1, tumor mutation burden, driver mutations) is unknown.

Patients and methods

We performed single-cell CITE/RNA/TCRseq on T cells isolated from 4 lung cancer clinical specimens. Cell surface CD39 were compared to transcriptional and proteomic features of these T cells. We subsequently performed flow cytometry on 440 fresh biospecimens from patients with lung cancer and quantified CD39 levels on CD8 T cells. %CD39 on CD8 T cells was compared with PD-L1, tumor mutation burden, histology, driver mutation, and anatomic site for each individual lesion evaluated. CD39 levels on CD8 T cells were also compared by prior chemotherapy or prior immunotherapy. In patients with stage IV metastatic disease from whom clinical samples were obtained pre- or within 3 weeks of treatment, CD39 levels on CD8 T cells was correlated with response rate and progression-free survival.

Results

Single-cell CITE/RNA/TCRseq revealed that CD39$^+$ CD8 T cells in lung cancers demonstrate

transcriptional and proteomic features of exhaustion, tumor reactivity, and clonal expansion. Flow

cytometry of 440 lung cancer specimens revealed that CD39 level on CD8 T cells is only weakly

correlated with tumoral features, such as histology, driver mutation, PD-L1 and tumor mutation

burden. CD39 correlated with PD-1 expression on CD8 T cells and high pre-treatment/early-on-

treatment levels were associated with improved clinical outcomes from immune checkpoint

blockade.

Conclusion

This comprehensive profiling of the clinical, pathological and molecular features highlights the

utility of CD39 as a proxy for tumor-reactive CD8 T cells in human lung cancer.

KEYWORDS

Immune-checkpoint blockade, lung cancer, CD8 T cells

INTRODUCTION

Lung cancer is the leading cause of cancer death in the world. Immune checkpoint blockade (ICB) utilizing antibodies that block programmed cell death protein 1 (PD-1), programmed death-ligand 1 (PD-L1), and cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) has ushered in a new era of hope, with long-term durable disease control in a subset of patients with lung cancer (*1, 2*). However, most patients do not respond to ICB therapy and many of those who initially respond eventually develop recurrent and progressive disease. While PD-L1 on tumor cells and tumor mutation burden (TMB) have been validated as predictors of benefit from ICB in lung cancer (*3, 4*), there is an opportunity to further refine these biomarkers by incorporating features of CD8 T cells, which are critical mediators of ICB efficacy.

Despite early suggestions that measurement of bulk CD8 tumor infiltrating lymphocytes (TILs) would predict benefit from ICB (5), multiple recent studies across different diagnostic modalities have demonstrated that total CD8 T cell content alone is not a reliable predictor of clinical benefit from ICB (6-10). Part of the reason for this lack of predictive benefit is that a large proportion of CD8 T cells in the tumor microenvironment are passive bystanders that lack tumor reactivity (*11, 12*). Recent elegant work has highlighted that CD8 T cells that express CD39 are enriched for features of a clonal, proliferative lymphocyte population that expresses high levels of activation/exhaustion (e.g. PD-1) and cytotoxicity (e.g. granzyme B) markers (*2, 12-15*). Moreover, CD39 was highly expressed on empirically defined neoantigen- and tumor-associated antigen-reactive T cells in lung cancer and melanoma (*16-19*). The largest profiling of CD39 expression on CD8 T cells to date in a single disease was performed in 68 patients with head and neck squamous cell cancer; this paper contained an additional 42 samples across a mixture of seven disease types (13). Herein, we describe the tumoral features that are associated with CD39 levels on CD8 T cells from 440 clinical samples of human lung cancer, validate the utility of CD39

as a surrogate of tumor-reactive CD8 T cells, and report that baseline CD39 on CD8 T cells

predicts enhanced efficacy of ICB in patients with lung cancer.

METHODS

Human biospecimens

Fresh primary tumors, metastatic lesions and pleural/peritoneal/pericardial effusions were

obtained from August 2018 to September 2021 with permission from the MSKCC IRB. Informed

consent was collected from all patients enrolled in this study. Clinical samples were annotated

with tumor histology and driver mutation. Adenocarcinoma, adenosquamous, and NSCLC NOS

tumors were annotated by their molecular driver mutations, if known. The category 'Unknown

Driver' refers to adenocarcinoma, adenosquamous, and NSCLC NOS histology tumors for which

a driver mutation (defined as 'known to be oncogenic' by OncoKB ([20])) was not identified; notably,

this category does not include squamous or small-cell histology biospecimens. With the exception

of a single case in which a tumor sample with squamous histology harbored a *MET* exon 14

mutation and two transformed small-cell lung cancer tumors with *EGFR* mutations, squamous

and small-cell lung cancers were annotated by their histology.

Flow cytometry

Cells were incubated with TruFCX (Biolegend) to block nonspecific binding, and then stained

(15 min, 4 °C) with appropriate dilutions of various combinations of fluorochrome-conjugated anti-

mouse antibodies. The stained cells were acquired on a LSRII Flow Cytometer or Aria 7 cell sorter

and the data were processed using FlowJo software (Treestar). Doublets and dead cells were

excluded based on forward scatter (FSC) and side scatter (SSC) and 4′,6-diamidino-2-

phenylindole staining (DAPI, 1 μg/ml; ThermoFisher). All depicted flow cytometry plots were pre-

gated on non-debris (by FSC and SSC), viable (DAPI⁻) single CD45⁺ CD3⁺ cells, unless otherwise

indicated in the Figure legend. Gating for %CD39 was determined by gating of fluorescence minus

one (FMO). PD-1 mean fluorescence intensity was measured for a subset of samples.

Single-cell transcriptome sequencing

Sorted or dissociated tumor cells were stained with Trypan blue and Countess II Automated Cell

Counter (ThermoFisher) was used to assess both cell number and viability. Following QC, the

single-cell suspension was loaded onto Chromium Chip A or Next GEM Chip K (10X Genomics

PN 230027/1000286) and GEM generation, cDNA synthesis, cDNA amplification, and library

preparation of 700-3,300 cells proceeded using the Chromium Single Cell 5' Reagent Kit or Next

GEM Single Cell 5' Kit v2 (10X Genomics PN 1000006/1000263) according to the manufacturer's

protocol. cDNA amplification included 14-16 cycles and 5.8 ng-20 ng of the material was used to

prepare sequencing libraries with 16 cycles of PCR. Indexed libraries were pooled equimolar and

sequenced on a NovaSeq 6000 in a PE26/91 or PE28/91 run using the NovaSeq 6000 SP or S1

Reagent Kit (100 cycles) (Illumina). An average of 108 million reads were generated per sample.

Single-cell V(D)J analysis from RNA

An aliquot of complementary DNA (cDNA) generated using the methods described above was

used to enrich for V(D)J regions using the Chromium Single Cell V(D)J Enrichment Kit Human T

Cell (10X Genomics PN 1000005) according to the manufacturer's protocol with 10 cycles of PCR

during enrichment and 9 cycles during library preparation. Indexed libraries were pooled

equimolar and sequenced on a NovaSeq 6000 in a PE150 or PE26/91 run using the NovaSeq

6000 SP or S4 Reagent Kit (100 or 300 cycles) (Illumina). An average of 24 million paired reads

was generated per sample.

Cell surface protein feature barcode analysis

Amplification products generated using the methods described above included both cDNA and

feature barcodes tagged with cell barcodes and unique molecular identifiers. Smaller feature

barcode fragments were separated from longer amplified cDNA using a 0.6X cleanup with

aMPure XP beads (Beckman Coulter catalog # A63882). Libraries were constructed using the

Chromium Single Cell 5' Feature Barcode Library Kit (10X Genomics PN 1000080) according to

the manufacturer's protocol with 9 cycles of PCR. Indexed libraries were pooled equimolar and

sequenced on a NovaSeq 6000 in a PE26/91 or PE28/91 run using the NovaSeq 6000 SP or S2

Reagent Kit (100 cycles) (Illumina). An average of 60 million paired reads was generated per

sample.

Single cell CITE/RNA/TCR analysis

Single-cell sequencing data were aligned to the Genome Reference Consortium Human Build 38

(GRCh38) using Cell Ranger (v3.1.0; 10X Genomics) in order to obtain T cell clonotypes, feature

barcoding, CITEseq antibody detection and gene expression profiles associated with individual

single cells. Each data type was matched to create a UMI matrix and cells were filtered out based

on three metrics: (1) cells with fewer than 200 detectable genes; (2) cells with more than 3000

detectable genes; (3) cells that had fewer than 5% percentage of counts related to mitochondrial

genes. Data normalization, Principal Component Analysis and subsequent Uniform Manifold

Approximation and Projections (UMAP) were performed on the dataset using the R package

Seurat v.3.1.1 (https://github.com/satijalab/seurat). The differential expression comparisons were

generated using the DESeq2 package with selected genes (FDR < 0.05). After filtering, we

created subclusters of cells using the 26 Louvain algorithm. Raw counts were normalized by

library size per cell. CD39^{neg} was defined as a normalized value of 0 for adt_CD39. CD39^{int} was

defined as the adt_CD39 level between 0 and 1.0 (non-inclusive) since 1.0 was the mean of all

non-zero values for adt_CD39 among all the cells in the CD8 cluster. CD39hi was defined as

adt_CD39 level greater than or equal to 1.0. Signature scores for exhaustion and T cell reactivity

were calculated by *AddModuleScore* in Seurat. Clonal proportion was calculated as the fractional

representation of all CD8 clones by clonotypes that were categorized by mean adt_CD39

expression with the same cutoffs for CD39^{low}, CD39^{int}, and CD39^{hi} as above.

Multivariate analyses

We used the R package caret (*21*) to implement the glmnet (*22*) algorithm and evaluate the

performance of a lasso regression model using 10-fold cross-validation. The data was center

and scaled during preprocessing and lambda was chosen based on the minimal RMSE

value. Additionally a linear model was evaluated using the R function "lm". In both models, the

response variable, CD39 expression, was log base 2 transformed as well as the predictor

variables TMB and total neoantigens per sample.

Clinical outcomes analyses

RFS assessment was performed on 188 biospecimens obtained from stage I-IIIA lung cancer

who did not receive neoadjuvant ICB and were not lost to follow up after resection. For the PFS

assessment in stage IV patients not treated with ICB, only patients that received at least two

cycles of platinum-based chemotherapy were included in the analysis (n=26). For the PFS

assessment in stage IV patient treated with ICB, only patients that received at least two cycles of

ICB without chemotherapy were included in the analysis (n=22). The cutoffs for %CD8 and

%CD39 of 15.4 and 23.3 were selected from the means of the 440-sample cohort. The PD-1

cutoff was selected to divide the cohort above and below the median for the stage IV cohort.

Response criteria were annotated per RECIST v1.1.

Statistical analysis

Data are expressed and statistical analyses were performed as described in the Figure legend

for each analysis. Statistical significance was determined by two-way ANOVA with Tukey's

multiple comparison test, student's t-test or Mantel-Cox log-rank test using Prism 7 software as

indicated.

Materials availability

This study did not generate new unique reagents.

Data availability

All data generated and supporting the findings of this study are available within the paper. The

single-cell CITE/TCR/RNAseq will be deposited in the Gene Expression Omnibus (GEO)

database. The TCRseq data will be available through ImmuneACCESS on the Adaptive

Biotechnologies website.

RESULTS

Single-cell cellular indexing of transcriptomes and epitopes (CITE)/RNA/T cell receptor

(TCR)seq reveals that CD39^{high} CD8 T cells are enriched for features of exhaustion, tumor

reactivity, and clonal expansion

To characterize CD8 T cells expressing CD39, we sorted CD3$^+$ T cells from four non-small

cell lung cancer (NSCLC) biospecimens and performed droplet-based single-cell

CITE/RNA/TCRseq (*23*). The four NSCLC samples comprised a range of histologies (e.g.

adenocarcinoma and squamous), driver mutations (*KRAS* and *EGFR*), stages (e.g. early and

metastatic), and anatomic sites (e.g. lung, pleural fluid, and brain) (**Table S1**). None of these

patients had received prior ICB at the time of sample collection. After coarse clustering, we

visualized separated clusters of regulatory T cells, CD4 T cells, CD8 T cells, and an additional T

cell NOS (not otherwise specified) cluster (**Fig. 1A, Table S2**). We focused our subsequent

analyses on CD8 T cells since this subset has the greatest known contribution to anti-tumor

immunity. The 896 single CD8 T cells that passed quality control (QC) were divided by CD39

protein expression (assessed by oligo-tagged anti-CD39 antibody) into high (hi), intermediate

(int), and negative (neg) groups. Transcriptional dropout is a well-known limitation of single cell

RNAseq, and oligonucleotide-tagged antibodies to surface molecules (e.g. CITE-seq) represents

a strategy to overcome this hurdle (*24*). Concordantly, CITEseq detected protein expression of

CD39 and other activation markers in many cells in which there was absent transcriptional

expression (**Fig S1A**). Genes that were differentially expressed in CD39^{hi} CD8 T cells included

the exhaustion marker *LAYN*, tissue residence marker *ITGAE* (CD103), and the activation

markers *CXCL13*, *GNLY*, *HLADRA*, and *VCAM1* (**Table S3**). CD39^{hi} CD8 T cells had the highest

transcriptional levels of *ENTPD1* (CD39), *PDCD1* (PD-1), *ITGAE*, *CXCL13*, *TNFRSF4* (OX-40),

HAVCR2 (TIM-3), and *LAG3*, which are features of tumor-reactive CD8 T cells (**Fig. S1B**) (*25*).

Protein level assessment by CITE-seq revealed that CD39^{hi} CD8 T cells across the four samples

consistently expressed the highest protein levels of PD-1, CD103, OX40, 4-1BB, and LAG-3 (**Fig.**

1B). Concordantly, CD39^{hi} CD8 T cells expressed the highest transcriptomic signature score for

T cell exhaustion and tumor reactivity (**Fig. 1C-D, Table S4**).

Two recent reports demonstrated that empirically validated tumor-reactive CD8 T cells

express CD39 (*16, 17*). In these studies, the authors assessed for CD8 tumor reactivity to tumor-

associated or viral-associated antigens and generated gene expression signatures of the reactive

CD8 T cells. Thus, we next evaluated whether the gene expression profiles of CD39^{hi} CD8 T in

our dataset overlapped with the CD8 T cells in these two recent datasets. CD39^{hi} CD8 T cells in

our dataset indeed expressed the highest 'tumor specific' and "MANA-TIL (mutation-associated

neoantigen-tumor infiltrating lymphocyte)' signature scores and lowest 'virus specific' and

'influenza' signatures (**Fig. 1E-F, S1C-D**).

Since CD39^{hi} CD8 T cells were enriched for features of tumor reactivity, we hypothesized

that these cells would display greater clonal expansion. Indeed CD39^{hi} CD8 T cells expressed the

highest proliferation score and both CD39^{int} and CD39^{hi} CD8 T cells comprised a higher clonal

proportion among all CD8 T cells (**Fig 1G-H**). Thus, our single cell CITE/RNA/TCR sequencing

demonstrates that CD39^{hi} CD8 T cells express features of exhaustion, tumor reactivity, and clonal

expansion.

CD39 expression on CD8 T cells is a non-redundant biomarker

We compared CD39, Tim-3, 4-1BB, and PD-1 expression by flow cytometry to compare

their relative staining resolution, as defined by the separation of the positive and negative

population. FACS-based detection of CD39 consistently yielded higher resolution compared to

the other three markers (**Fig. 2A**). The enhanced resolution of CD39 compared to PD-1 is

consistent with our prior report that CD39 has more durable expression in CD8 T cells (*26*). Due

to the staining resolution of the CD39 marker and its association with tumor reactivity, we sought

to utilize CD39 expression on CD8 T cells to estimate the frequency of tumor-reactive CD8 T cells

in lung cancer clinical samples. From August 2018 to September 2021, we evaluated 440 fresh

lung cancer clinical specimens by flow cytometry for CD39 expression on CD8 T cells. These

biospecimens ranged across stages (I-IV) and lung cancer subtypes (lung adenocarcinoma,

squamous cell cancer, and small cell lung cancer) (**Table S5**). The mean CD8 level was 15.4%

(of CD45$^+$) and mean CD39 level was 23.3% (of CD8 T cells), and these were utilized as cutoffs

throughout the rest of the study. On univariate analysis, CD39 expression on CD8 T cells had a

weak correlation with total CD8 T cells, smoking history, TMB, and PD-L1 (**Fig. S2A-D**). Since

TMB $\geq$ 10 mutations/megabase and PD-L1 $\geq$ 50% represent subgroups with favorable clinical

outcomes from ICB therapy (*3, 4*), we assessed %CD39 on these subpopulations. When divided

into quartiles by PD-L1 and TMB expression, tumor material from the two TMB $\geq$ 10

mutations/megabase subgroups (both PD-L1 < 50% and $\geq$ 50%) had nearly two-fold higher

%CD39 among CD8 T cells, relative to the TMB < 10 and PD-L1 < 50% subgroup (**Fig. S2E**). In

contrast, the proportion of CD8 T cells (**Fig. S2F**) showed less variation. There was largely no

association of CD39 expression on CD8 T cells with lung cancer stage with the exception of a

reduction in stage IVA tumors (**Fig. S2G**). This is concordant with reduced CD39 expression on

CD8 T cells in pleural fluid and pleural metastases biospecimens relative to the lung biospecimens

(**Fig. S2H**). Notably, we previously reported that the pleural and peritoneal cavities are

immunosuppressed microenvironments due to the presence of Tim-4$^+$ cavity-resident

macrophages (*26*), and this may contribute to the low %CD39 in these anatomic compartments.

CD39 levels on CD8 T cells showed two patterns of expression among the lung cancer

subtypes. *ALK* fusion, *ROS1* fusion, *MET* exon 14 fusion, *BRAF* V600E, *RET* fusion and *EGFR*

mutant (except exon 20 insertion) lung adenocarcinomas, which are not associated with tobacco

use, tended to have below average CD39 expression on CD8 T cells (**Fig. 2B, Table S6**). The

reduced CD39 expression on *EGFR* mutant lung cancer tumors is in line with a prior report (*12*).

In contrast, CD39 expression on lung adenocarcinomas with *KRAS* G12C, G12D, G12V or

ERBB2 (HER2) mutations were above average. 'Other *KRAS*' mutations, including G12A, G12R,

G12S, G13C, G13D, KRAS Q61H, and Q61R, which are known to be oncogenic per OncoKB

(*20*), were associated with a lower-than-mean CD39 level on CD8 T cells. The proportion of

CD39$^+$ CD8 T cells in squamous and small cell lung cancers averaged 30.3% and 27.3%,

respectively. In contrast, only 6.4% of CD8 T cells in neuroendocrine carcinoid (other

neuroendocrine) tumors expressed CD39. These findings are consistent with *KRAS*, squamous,

and small-cell lung cancers being strongly associated with tobacco use, which is correlated with

a higher number of tumor mutations and neoantigens to which CD8 T cells can react.

Nonetheless, even among the lung cancer subtypes with higher %CD39, there was a wide range

of CD39 levels (**Fig. 2B**), suggesting that even for a given driver mutation, there is substantial

heterogeneity in tumor-reactive CD8 immunosurveillance. Total CD8 T cell infiltration, TMB, and

PD-L1 were overall not associated with consistent differences across lung cancer histological

subtypes (**Fig. 2C, S2I-J**).

Since our data was retrospective and derived from a heterogeneous cohort of patients,

we performed multivariate analyses to determine the clinical and molecular features that best

correlate with CD39 expression on CD8 T cells. We included tissue site, stage, histology, driver

mutation, TMB, smoking history, and PD-L1 as potential covariates. We also included human

leukocyte antigen (HLA) heterozygosity (*27, 28*) in the model and also the number of predicted

neoantigens and strong binding neoantigens from next-generation sequencing of tumor biopsies

by MSK-IMPACT (*29*). A linear model predicted the variance in %CD39 on CD8 T cells at an

adjusted R-squared of 0.24, suggesting that the features in the model poorly predict the variance

observed in CD39 levels (**Table S7**). In the model, only TMB, PD-L1, pericardial metastases, and

prior chemotherapy passed significance thresholds of $P < 0.05$. Since some of the features in the

dataset had few observations and to avoid overfitting, we also applied a lasso regression with 10-

fold cross validation. The lambda was selected based on the root-mean-square error (RMSE) and

the model achieved an R-squared of 0.17 with a RMSE of 1.5 at lambda = 0.1 (**Table S8**). Notably,

at higher lambdas, the algorithm removed all predictors suggesting that there was no linear

combination of any regressed parameters that predicted CD39 levels well. Overall, these results

indicate that CD39 expression on CD8 T cells is a feature of the tumor that is non-redundant to

the tumoral parameters that currently guide therapy in lung cancer (e.g. histology, driver

alterations PD-L1, and TMB).

CD39 is upregulated on CD8 T cells in the tumor microenvironment

We next sought to understand the patterns of tissue-specific expression for CD39 and

thus performed CD39 staining on CD8 T cells from matched tumor and peripheral blood. We

analyzed three paired samples from peripheral blood and matched tumor lesions with varied

anatomic sites (e.g. the brain, lymph node, and lung). In all three cases, high levels of CD39 were

only observed in the tumor tissue and not in the peripheral blood (**Fig. S3A**). This suggests that

CD8 T cell interactions with cancer cells drives the expression of CD39 and is in line with our

recent report that CD39 was expressed at higher levels in regions of a resected tumor with viable

cancer cells compared to regions without viable tumor, normal adjacent regions, and lymph nodes

(*30*). Since there was a small CD39[+] population of intermediate intensity in the peripheral blood,

we evaluated whether this population gave rise to the CD39$^+$ CD8 T cells in the tumor tissue. To

address this question, we sorted CD39$^-$ and CD39$^+$ CD8 T cells from the peripheral blood and

CD39$^-$ or CD39$^+$ from the tumor tissue. We then performed bulk TCR sequencing analysis and

assessed for clonal overlap. As assessed by the Morisita Index, there was a minimal level of

clonal overlap between CD39$^+$ CD8 T cells from the peripheral blood and CD39$^+$ CD8 T cells from

tumor tissue, suggesting that circulating CD39$^+$ CD8 T cells may not be the dominant precursors

for the CD39$^+$ CD8 T cells in the tumor microenvironment (**Fig. S3A-C**).

We have previously shown that CD39 surface expression manifests more durable

expression after TCR stimulation compared to other markers associated with activation, such as

phosphatidylserine, PD-1, and CD44 (*26*). Due the greater durability of CD39 even after removal

of TCR stimulation, we next assessed the robustness of CD39 levels across spatially and

temporally distinct lesions. We analyzed 12 paired samples in which two anatomically distinct

lesions were simultaneously assessed (e.g. two resected lung lesions, lymph node and primary

lung lesions, or pleural metastasis and fluid). We noted reasonable concordance across spatially

distinct lesions (**Fig. S3D**), as exemplified by the low CD39 level on CD8 T cells from the paired

pleural metastasis and fluid from MSK 1266 (**Fig. S3E**). There was one notable case of divergent

CD39 levels. MSK 1372a and b were simultaneous resections of two right-sided separate primary

lung cancer lesions with marked differences in CD39 expression (59.0% vs. 92.5%; **Fig. S3F**).

Consistent with being separate primary lesions, these two tumors were comprised of two different

KRAS driver mutations with differential levels of TMB, which may underly the divergence of CD39

expression. We also followed CD39 levels on CD8 T cells from the same anatomic site multiple

times (e.g. recurrent pleural effusions, initial lung biopsy followed by resection, and serial

resections for recurrent disease in the brain). The CD39 levels over time across nine patients

were largely durable, including for three brain lesions that were resected over the course of nine

months (MSK 1265; **Fig. S3G-H**). These findings highlight that CD39 is upregulated on CD8 T

cells in proximity to the tumor and that CD39 levels on CD8 T cells are relatively preserved across

spatially and temporally distinct lesions in the same patient.

PD-1 axis blockade increases the frequency of CD39$^+$ CD8 T cells

Since our data and prior reports (*12, 16*) demonstrate that tumor antigen-reactive CD8 T

cells express CD39 in lung cancer, we sought to leverage CD39 as a proxy for CD8 anti-tumor

immunosurveillance. It has been proposed that chemotherapy can result in 'immunogenic cell

death' that can prime an anti-tumor CD8 T cell response (*31*). We reasoned that CD39 levels

would be increased in patients after cytotoxic chemotherapy if such immune priming occurred.

Analogously, ICB therapy has been described to expand the pool of antigen-specific CD8 T cells

that infiltrate the tumor (*32*), which we reasoned might be reflected in increased CD39$^+$ CD8 T

cells in the tumor. Thus, we sought to evaluate in our dataset whether we found increases of

CD39$^+$ CD8 T cells with either cytotoxic chemotherapy or ICB therapy.

We first investigated the 218 patients in our cohort with stage IV lung cancer who were

treated with or without chemotherapy and/or ICB in the prior three or six months. Both total CD8

infiltration and CD39 levels on CD8 T cells were unchanged in patients receiving cytotoxic

chemotherapy in the preceding three or six months (**Fig. 3A-D**). Total CD8 infiltration was also

unchanged in patients receiving preceding ICB therapy (**Fig. 3A,C**). In contrast, CD8 T cells from

patients that received ICB in the prior three or six months expressed higher levels of CD39 (**Fig.**

3B,D). In patients that received both cytotoxic chemotherapy and ICB therapy in prior three or six

months, the level of CD39 on CD8 T cells was unchanged compared to patients that had not

received chemotherapy or ICB in the prior 3 or 6 months, which is consistent with chemotherapy

and ICB having opposing effects on CD39 expression (**Fig. 3B,D**).

Since the stage IV lung cancer cohort is quite heterogeneous, we subsequently examined

208 early-stage (stage I-III) NSCLC biospecimens from patients that did or did not receive

neoadjuvant therapy prior to sample collection. Among early-stage patients with NSCLC, we

again did not observe differences in total CD8 infiltration with preceding neoadjuvant

chemotherapy; however, neoadjuvant immunotherapy did increase total CD8 infiltration (**Fig. 3E**).

CD8 T cells from patients who underwent neoadjuvant chemotherapy had reduced CD39 levels

compared to patients without neoadjuvant therapy; in contrast, in patients who underwent

neoadjuvant ICB, we observed an increase in CD39 expression on CD8 T cells (**Fig. 3F**). Thus,

across two clinical cohorts, we did not find evidence for 'immunogenic cell death' with standard

cytotoxic chemotherapy utilized for lung cancer; however, we did discover that ICB exposure was

associated with an enhanced infiltration of CD39$^+$ CD8 T cells. This is consistent with recent

appreciation that ICB results in clonal expansion and infiltration of tumor-reactive CD8 T cells

(*32*).

CD39 and PD-1 expression on CD8 T cells portends benefit from ICB therapy

We next sought to evaluate whether CD39 on CD8 T cells had prognostic significance in

our dataset. We first assessed whether CD39 levels at the time of resection in 188 patients with

stage I-IIIA lung cancer resulted in differential recurrence-free survival (RFS). Since CD39 levels

are modulated by prior ICB (**Fig 3F**), we excluded patients who had previously received

neoadjuvant ICB in this analysis. Across this early-stage resection cohort, we did not observe

differences in RFS based on total CD8 T cells or %CD39 among CD8 T cells utilizing mean cutoff

values from the total 440 patient dataset (**Fig. 4A-C, Fig. S4A-C**). The data for CD8 T cells is in

line with a recently published prospective cohort of early-stage NSCLC resection specimens

which showed no differences in clinical outcomes based on total CD8 abundance (*33*). Similarly,

neither total CD8 T cells or %CD39 among CD8 T cells resulted in differential progression-free

survival (PFS) in the 26 stage IV lung cancer patients undergoing cytotoxic chemotherapy without

ICB (**Fig. 4D, S4D**). Hence, CD39 expression on CD8 T cells was not prognostic for lung cancer

in our dataset.

We next asked whether we could find evidence that CD39 levels on CD8 T cells was

associated with response to ICB. There were 22 patients in our cohort from whom we had

obtained clinical biospecimens prior to or within three weeks of commencing ICB therapy (**Fig.**

4E). Although the tumoral PD-L1, TMB, and %CD8 among DAPI⁻ cells were not different between

non-responders (stable disease/progression of disease, SD/POD) and responders (partial

response/complete response, PR/CR) to ICB, we observed that responders had a higher level of

PD-1 and CD39 on CD8 T cells (**Fig. 4F-J**). While total CD8 infiltration did not differentiate PFS

in these patients (**Fig. 4K**), patients with higher levels of PD-1 or CD39 on CD8 T cells had an

improved PFS (p=0.0145 for PD-1 and p=0.0063 for CD39, **Fig. 4L-M**). Thus, consistent with prior

reports in lung cancer (*2, 34, 35*), levels of baseline exhausted CD8 T cells are associated with

clinical benefit from ICB therapy.

DISCUSSION

Lung cancer therapy is currently guided by tumor cell-intrinsic features, including

histology, driver alterations, PD-L1 and TMB. Integration of tumor cell-extrinsic features such as

the immune infiltrate can potentially lead to improved therapeutic options for patients with lung

cancer. This was exemplified by the recent report of a 'lung cancer activation module', consisting

of activated T cells, plasma cells, and macrophages, that served as a predictor of response to

immunotherapy (*36*). Since CD8 T cells are the critical effectors in ICB therapy, features of CD8

T cells could also potentially serve as a non-redundant biomarker of response to ICB. Although

methods to empirically verify tumor reactivity have improved substantially in sensitivity and

throughput (e.g. MANIFEST (*37*) and 4-1BB assay (17)), there remains a need in certain contexts

for simple and reliable methods to distinguish and quantify the tumor-reactive CD8 sub-population

from bystanders without cognate TCR reactivity. We observed from single-cell profiling of

heterogeneous lesions that CD8 T cells expressing high levels of CD39 are enriched for features

of exhaustion, tumor reactivity and clonal expansion. Motivated by this finding and recent reports

that empirically demonstrated that tumor-reactive CD8 T cells express CD39 (*2, 12, 13, 16, 17,*

25, 38), we profiled CD39 on 440 lung cancer biospecimens obtained at MSKCC. We observed

that CD39 on CD8 T cells is expressed more highly in tobacco-associated lung cancer genotypes

and poorly correlates with tumor TMB and PD-L1. Hence, CD39 is a biomarker that is non-

redundant to tumoral features of lung cancer. CD39 is likely dependent on other variables not

captured in our dataset, including T cell repertoire, HLA subtype, peptide binding, and likely the

intersection of these three highly diverse attributes. CD39 is upregulated on CD8 T cells in the

tumor microenvironment relative to the periphery and is generally stable across space and time.

These data nominate CD39-guided strategies for TCR-based therapeutics against these targets

(*39*). Finally, we demonstrate that although the level of CD39 on CD8 T cells was not prognostic

in any of the patient cohorts that we examined, it was associated with clinical benefit from ICB

therapy.

There have been conflicting data regarding whether the presence of CD8 T cells

expressing markers of exhaustion can predict clinical benefit from ICB. Single-cell RNAseq of

tumors from metastatic melanoma revealed that a CD8 T cell state called CD8_B, which is

enriched in *ENTPD1* (CD39), *PDCD1* (PD-1), *CD38*, *HAVCR2* (TIM-3), and *LAG3*, is

preferentially expressed in non-responders to ICB (9). Also, the use of multiplex

immunofluorescence staining in lung cancer suggests that an 'effector burnt out' (EBO) CD8 T

effector population preferentially expressing PD-1 and LAG-3 was enriched in patients who did

not derive durable benefit from ICB and that these patients had reduced overall survival (*40*).

Notably, CD39 was not a selective feature of the EBO CD8 T cells. On the other hand, lung cancer

patients with high pre-treatment levels of PD-1 by flow cytometry or immunofluorescence

demonstrated a greater response rate and PFS (*2, 34*). Moreover, high levels of intratumoral

CD39$^+$ CD8 T cells is associated with response to ICB therapy in lung cancer (*35*). These findings

dovetail with other studies that revealed that high levels of CD8 T cells with features of exhaustion

prior to ICB are associated with improved clinical benefit in colorectal cancer and ER$^+$ breast

cancer (*41, 42*). Our data confirm that the PD-1 and CD39 on CD8 T cells are both predictive of

improved outcomes from ICB therapy.

Two recent reports highlight that the predictive significance of CD39 for immunotherapy is

context dependent. Caushi et. al. demonstrated that although CD8 T cells that recognize

neoantigen mutations express CD39, neoantigen-reactive CD8 T cells that expressed higher

levels of CD39 were associated with inferior pathological responses (16). Krishna et. al. similarly

showed that neoantigen-reactive CD8 T cells in *ex vivo* expanded TILs prior to infusion

preferentially expressed CD39 and CD69, but improved clinical outcomes were observed in

patients whose pre-infusion products harbored a high number of CD39$^-$ CD69$^-$ TILs (*43*). One

explanation for this apparent paradox may be that the relative timing of the evaluation of

exhaustion likely influences the predictive capability of this biomarker since our data demonstrates

that CD39 increases after ICB. Notably, in the two cases highlighted above, the correlation of

CD39 with worse outcomes was read out after immune based intervention (i.e. ICB or *ex vivo*

expansion followed by adoptive cell transfer). It is plausible that pre-treatment/early on-treatment

exhaustion is reflective of a mobilizable pool of tumor-reactive T cells that portends a favorable

response to ICB; conversely, exhaustion after a course of ICB might be indicative of a persistent

hypofunctional state that is associated with a worse clinical outcome.

There are several limitations of this study that we acknowledge. First, the data are derived

from patients treated at a single center. Second, while we profiled CD39 on CD8 T cells, our data

does not capture the likely contribution/modulation of other immune cell types such as B cells,

innate lymphoid cells, and myeloid cells. Lastly, our cohort to assess the predictive significance

of CD39 expression on CD8 T cells in stage IV lung cancer treated with ICB monotherapy is

limited to 22 patients despite >3 years of fresh tissue collection. Part of this limited sample size is

attributable to the routine incorporation of cytotoxic chemotherapy to ICB for frontline treatment

of the majority of patients with lung cancer. Despite these limitations, the data presented here

contribute substantially to our understanding of CD39 on CD8 T cells in the context of lung cancer,

including factors that modulates its pattern of expression. Our study identifies CD39 on CD8 T

cells as a biomarker that is non-redundant to TMB and PD-L1, and which can be readily captured

on clinical samples to serve as a proxy of the tumor-reactive CD8 T cell population. Subsequent

studies can also leverage CD39 to enrich for TCR candidates to evaluate for TCR-based

immunotherapies.

ACKNOWLEDGEMENTS

We are grateful for experimental support from the MSKCC Molecular Cytology Core Facility, Flow

Cytometry Core Facility, and Integrated Genomics Operation Core (funded by the NCI Cancer

Center Support Grant (CCSG, P30 CA08748), Cycle for Survival, and the Marie-Josée and Henry

R. Kravis Center for Molecular Oncology). We are grateful for manuscript editing provided by Dr.

Clare Wilhelm and Reeja Thomas.

Funding

This research was funded in part through the NIH NCI CCSG P30 CA008748, NCI R01

CA056821, U24 CA213274, P01 CA129243, R01 CA197936, R37 CA259177 (CAK), Cancer

Research Institute CRI3176 (CAK), Stony-Wold Herbert Fund, International Association of Lung

Cancer Research (IASLC)/ International Lung Cancer Foundation (ILCF), the MSKCC Society

Grant, the Ludwig Collaborative and Swim Across America Laboratory, the Emerald Foundation,

the Parker Institute for Cancer Immunotherapy at MSKCC, the Department of Medicine at

MSKCC, and Stand Up To Cancer (SU2C)-American Cancer Society Lung Cancer Dream Team

Translational research grant (SU2C-AACR-DT17-15). AC was supported by an MSKCC

Investigational Cancer Therapeutics Training Program fellowship (T32 CA-009207) and Clinical

Investigator Award from National Cancer Institute (K08 CA-248723).

Disclosure

CAK received research grant support from Kite/Gilead; is on the Scientific and/or Clinical Advisory

Boards of Achilles Therapeutics, Aleta BioTherapeutics, Bellicum Pharmaceuticals, Catamaran

Bio, Obsidian Therapeutics, and T-knife, and has performed consulting services for Bristol Myers

Squibb, PACT Pharma, and Roche/Genentech. CAK is a co-inventor on patent applications

related to TCRs targeting public neoantigens unrelated to the current work. MDH received

research grant from BMS; personal fees from Achilles, Arcus, AstraZeneca, Blueprint, BMS,

Genentech/Roche, Genzyme, Immunai, Instil Bio, Janssen, Merck, Mirati, Natera, Nektar, Pact

Pharma, Regeneron, Shattuck Labs, Syndax, as well as equity options from Arcus, Factorial,

Immunai, and Shattuck Labs. A patent filed by MSKCC related to the use of tumor mutational

burden to predict response to immunotherapy (PCT/US2015/062208) is pending and licensed by

PGDx. JDW is a consultant for Adaptive Biotech, Amgen, Apricity, Ascentage Pharma, Arsenal

IO, Astellas, AstraZeneca, Bayer, Beigene, Boehringer Ingelheim, Bristol Myers Squibb, Celgene,

Chugai, Daiichi Sankyo, Dragonfly, Eli Lilly, Elucida, F Star, Georgiamune, Idera, Imvaq, Kyowa

Hakko Kirin, Linneaus, Maverick Therapeutics, Merck, Neon Therapeutics, Polynoma, Psioxus,

Recepta, Takara Bio, Trieza, Truvax, Trishula, Sellas, Serametrix, Surface Oncology, Syndax,

Syntalogic, and Werewolf Therapeutics. JDW has received grant/research support from Bristol

Myers Squibb and Sephora. JDW has equity in Tizona Pharmaceuticals, Adaptive

Biotechnologies, Imvaq, Beigene, Linneaus, Apricity, Arsenal IO, and Georgiamune. JDW is a co-

inventor on patent applications related to heteroclitic cancer vaccines and recombinant poxviruses

for cancer immunotherapy. JDW and TM are co-inventors on patent applications related to CD40

and in situ vaccination (PCT/US2016/045970). TM is a consultant for Immunos Therapeutics and

Pfizer. TM is a cofounder of and equity holder in IMVAQ Therapeutics. TM receives research

funding from Bristol-Myers Squibb, Surface Oncology, Kyn Therapeutics, Infinity

Pharmaceuticals, Peregrine Pharmaceuticals, Adaptive Biotechnologies, Leap Therapeutics, and

Aprea Therapeutics. TM is an inventor on patent applications related to work on oncolytic viral

therapy, alpha virus–based vaccine, neoantigen modeling, CD40, GITR, OX40, PD-1, and CTLA-

4. C.M.R. has consulted regarding oncology drug development with AbbVie, Amgen, Ascentage,

AstraZeneca, BMS, Celgene, Daiichi Sankyo, Genentech/Roche, Ipsen, Loxo and PharmaMar

and is on the scientific advisory boards of Elucida, Bridge and Harpoon.

TABLE LEGENDS

Table S1. Clinical features of four lung cancer clinical biospecimens that underwent single cell

CITE/TCR/RNA sequencing.

Table S2. Differentially expressed genes for CD8, CD4, T cell NOS, and Treg clusters from single

cell CITE/TCR/RNA sequencing.

Table S3. Differentially expressed genes for CD39^{hi} CD8 T cells compared to all other CD8 T

cells from single cell CITE/TCR/RNA sequencing.

Table S4. List of gene signatures used for scoring.

Table S5. Clinical and molecular features of 440 patient lung cancer cohort.

Table S6. Median and mean values for %CD39 on CD8 T cells among lung cancer subtypes.

Table S7. Linear regression multivariate analysis of %CD39 on CD8 T cells.

Table S8. Lasso regression multivariate analysis of %CD39 on CD8 T cells.

REFERENCES

1.   A. Ribas, J. D. Wolchok, Cancer immunotherapy using checkpoint blockade. *Science* **359**,
1350-1355 (2018).
2.   D. S. Thommen *et al.*, A transcriptionally and functionally distinct PD-1(+) CD8(+) T cell
pool with predictive potential in non-small-cell lung cancer treated with PD-1 blockade.
Nat Med **24**, 994-1004 (2018).
3.   M. D. Hellmann *et al.*, Nivolumab plus Ipilimumab in Lung Cancer with a High Tumor
Mutational Burden. *N Engl J Med* **378**, 2093-2104 (2018).
4.   M. Reck *et al.*, Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell
Lung Cancer. *N Engl J Med* **375**, 1823-1833 (2016).
5.   P. C. Tumeh *et al.*, PD-1 blockade induces responses by inhibiting adaptive immune
resistance. *Nature* **515**, 568-571 (2014).
6.   N. Riaz *et al.*, Tumor and Microenvironment Evolution during Immunotherapy with
Nivolumab. *Cell* **171**, 934-949 e916 (2017).
7.   V. Verma *et al.*, PD-1 blockade in subprimed CD8 cells induces dysfunctional PD-
1(+)CD38(hi) cells and anti-PD-1 resistance. *Nat Immunol* **20**, 1231-1243 (2019).
8.   B. A. Helmink *et al.*, B cells and tertiary lymphoid structures promote immunotherapy
response. *Nature* **577**, 549-555 (2020).
9.   M. Sade-Feldman *et al.*, Defining T Cell States Associated with Response to Checkpoint
Immunotherapy in Melanoma. *Cell* **175**, 998-1013 e1020 (2018).
10.   J. C. Osorio *et al.*, Lesion-Level Response Dynamics to Programmed Cell Death Protein
(PD-1) Blockade. *J Clin Oncol* **37**, 3546-3555 (2019).
11.   W. Scheper *et al.*, Low and variable tumor reactivity of the intratumoral TCR repertoire
in human cancers. *Nat Med* **25**, 89-94 (2019).
12.   Y. Simoni *et al.*, Bystander CD8(+) T cells are abundant and phenotypically distinct in
human tumour infiltrates. *Nature* **557**, 575-579 (2018).
13.   T. Duhen *et al.*, Co-expression of CD39 and CD103 identifies tumor-reactive CD8 T cells
in human solid tumors. *Nat Commun* **9**, 2724 (2018).
14.   H. Li *et al.*, Dysfunctional CD8 T Cells Form a Proliferative, Dynamically Regulated
Compartment within Human Melanoma. *Cell*, (2018).
15.   K. E. Yost *et al.*, Clonal replacement of tumor-specific T cells following PD-1 blockade.
Nat Med **25**, 1251-1259 (2019).
16.   J. X. Caushi *et al.*, Transcriptional programs of neoantigen-specific TIL in anti-PD-1-
treated lung cancers. *Nature*, (2021).
17.   G. Oliveira *et al.*, Phenotype, specificity and avidity of antitumour CD8(+) T cells in
melanoma. *Nature* **596**, 119-125 (2021).
18.   F. J. Lowery *et al.*, Molecular signatures of antitumor neoantigen-reactive T cells from
metastatic human cancers. *Science*, eabl5447 (2022).
19.   B. Liu *et al.*, Glioma Stem Cells Upregulate CD39 Expression to Escape Immune Response
through SOX2 Modulation. *Cancers (Basel)* **14**, (2022).
20.   D. Chakravarty *et al.*, OncoKB: A Precision Oncology Knowledge Base. *JCO Precis Oncol*
2017, (2017).

21.   M. Kuhn, Building Predictive Models in R Using the caret Package. *Journal of Statistical*
596   *Software* **28**, 1 - 26 (2008).

597   22.   J. Friedman, T. Hastie, R. Tibshirani, Regularization Paths for Generalized Linear Models
598   via Coordinate Descent. *J Stat Softw* **33**, 1-22 (2010).

599   23.   M. Stoeckius *et al.*, Simultaneous epitope and transcriptome measurement in single
600   cells. *Nat Methods* **14**, 865-868 (2017).

601   24.   S. H. Gohil, J. B. Iorgulescu, D. A. Braun, D. B. Keskin, K. J. Livak, Applying high-
602   dimensional single-cell technologies to the analysis of cancer immunotherapy. *Nat Rev*
603   *Clin Oncol* **18**, 244-256 (2021).

604   25.   A. M. van der Leun, D. S. Thommen, T. N. Schumacher, CD8(+) T cell states in human
605   cancer: insights from single-cell analysis. *Nat Rev Cancer* **20**, 218-232 (2020).

606   26.   A. Chow *et al.*, Tim-4(+) cavity-resident macrophages impair anti-tumor CD8(+) T cell
607   immunity. *Cancer Cell* **39**, 973-988 e979 (2021).

608   27.   D. Chowell *et al.*, Evolutionary divergence of HLA class I genotype impacts efficacy of
609   cancer immunotherapy. *Nat Med* **25**, 1715-1720 (2019).

610   28.   D. Chowell *et al.*, Patient HLA class I genotype influences cancer response to checkpoint
611   blockade immunotherapy. *Science* **359**, 582-587 (2018).

612   29.   A. Zehir *et al.*, Mutational landscape of metastatic cancer revealed from prospective
613   clinical sequencing of 10,000 patients. *Nat Med* **23**, 703-713 (2017).

614   30.   J. A. Pai *et al.*, Regional and clonal T cell dynamics at single cell resolution in immune
615   checkpoint blockade. *bioRxiv*, 2021.2009.2027.461389 (2021).

616   31.   L. Galluzzi, A. Buque, O. Kepp, L. Zitvogel, G. Kroemer, Immunogenic cell death in cancer
617   and infectious disease. *Nat Rev Immunol* **17**, 97-111 (2017).

618   32.   K. E. Yost, H. Y. Chang, A. T. Satpathy, Recruiting T cells in cancer immunotherapy.
619   *Science* **372**, 130-131 (2021).

620   33.   L. Federico *et al.*, Distinct tumor-infiltrating lymphocyte landscapes are associated with
621   clinical outcomes in localized non-small cell lung cancer. *Ann Oncol*, (2021).

622   34.   S. Kumagai *et al.*, The PD-1 expression balance between effector and regulatory T cells
623   predicts the clinical efficacy of PD-1 blockade therapies. *Nat Immunol* **21**, 1346-1358
624   (2020).

625   35.   J. Yeong *et al.*, Intratumoral CD39(+)CD8(+) T Cells Predict Response to Programmed Cell
626   Death Protein-1 or Programmed Death Ligand-1 Blockade in Patients With NSCLC. *J*
627   *Thorac Oncol*, (2021).

628   36.   A. M. Leader *et al.*, Single-cell analysis of human non-small cell lung cancer lesions
629   refines tumor classification and patient stratification. *Cancer Cell* **39**, 1594-1609 e1512
630   (2021).

631   37.   L. Danilova *et al.*, The Mutation-Associated Neoantigen Functional Expansion of Specific
632   T Cells (MANAFEST) Assay: A Sensitive Platform for Monitoring Antitumor Immunity.
633   *Cancer Immunol Res* **6**, 888-899 (2018).

634   38.   K. I. Hanada *et al.*, A phenotypic signature that identifies neoantigen-reactive T cells in
635   fresh human lung cancers. *Cancer Cell*, (2022).

636   39.   S. S. Chandran, C. A. Klebanoff, T cell receptor-based cancer immunotherapy: Emerging
637   efficacy and pathways of resistance. *Immunol Rev* **290**, 127-147 (2019).

638   40.   M. F. Sanmamed *et al.*, A burned-out CD8+ T-cell subset expands in the tumor
639   microenvironment and curbs cancer immunotherapy. *Cancer Discov*, (2021).
640   41.   M. Chalabi *et al.*, Neoadjuvant immunotherapy leads to pathological responses in MMR-
641   proficient and MMR-deficient early-stage colon cancers. *Nat Med* **26**, 566-576 (2020).
642   42.   M. Terranova-Barberio *et al.*, Exhausted T cell signature predicts immunotherapy
643   response in ER-positive breast cancer. *Nat Commun* **11**, 3584 (2020).
644   43.   S. Krishna *et al.*, Stem-like CD8 T cells mediate response of adoptive cell immunotherapy
645   against human cancer. *Science* **370**, 1328-1334 (2020).
646

Figure 1. Single-cell CITE/RNA/TCR-sequencing reveals that CD39ʰⁱ CD8 T cells are enriched for features of exhaustion, tumor-reactivity, and clonal proliferation in human lung cancer. A) UMAP of sorted CD3⁺ T cells from four patients with lung cancer (see Table S1 for patient characteristics). Clusters are annotated on left panel. Surface levels of CD4, CD8, and CD39 as assessed by CITEseq antibody-derived tags (adt) are depicted in right three panels. B) Levels of various proteins (each column) across the four samples (each row) as determined by CITEseq adt levels. C-G) Scaled scores for exhaustion, tumor reactivity, tumor specific, virus specific, and proliferation gene signatures (see Table S4 for full gene list). H) Clonal proportion among CD8 T cells of clonotypes that were categorized by mean CD39 expression. Statistical significance was determined with two-way ANOVA with Tukey's multiple comparisons test and p value is indicated if < 0.05.

Figure 2. CD39 expression among CD8 TILs varies with lung cancer subtype. A) Representative flow cytometry staining of CD39, Tim-3, 4-1BB, and PD-1 on DAPI⁻ CD45⁺ CD3⁺ CD8⁺ T cells from MSK 1105b. Left plot represents fluorescence minus one (FMO) and right plot represents the CD8 T cells stained with the indicated antibody. B-C) Violin plots of %CD39⁺ (among CD8⁺ T cells) and %CD8 (among CD45⁺) for various histological subtypes/driver mutation categories among the 440-sample cohort. Dashed line indicates 23.3% mean CD39 level for entire 440-sample cohort. Red bars indicate histological subtypes/driver mutations with above average %CD39 values.

Figure 3. PD-1 axis blockade increases the frequency of CD39+ CD8 tumor-infiltrating lymphocytes.
A-B) %CD8 (among CD45+) or %CD39 (among CD8 T cells) from 218 biospecimens obtained from patients with stage IV metastatic disease that did or did not receive chemotherapy or ICB therapy in the prior 3 months. Statistical significance was assessed by two-tailed student's t-test. C-D) %CD8 (among CD45+) or %CD39 (among CD8 T cells) from 218 biospecimens obtained from patients with stage IV metastatic disease that did or did not receive chemotherapy or ICB therapy in the prior 6 months. Statistical significance assessed by two-tailed student's t-test. E-F) %CD8 (among CD45+) or %CD39 (among CD8 T cells) from 208 biospecimens obtained from patients with early stage (stage I-III) NSCLC. Statistical significance was assessed by two-tailed student's t-test. *p<0.05, ***p<0.001.

Figure 4. CD39 expression among CD8 tumor-infiltrating lymphocytes is not prognostic, but portends a beneficial outcome to immune checkpoint blockade in lung cancer. A-C) Kaplan-Meier curve of recurrence-free survival after resection with above or below mean %CD39 (among CD8 T cells) for stage I, stage II, and stage IIIA lung cancer patients who did not receive ICB. Statistical significance determined by Mantel-Cox test. D) Kaplan-Meier curve of progression-free survival after resection with above or below mean %CD39 (among CD8 T cells) for stage IV lung cancer patients who did not receive ICB. Statistical significance determined by Mantel-Cox test. E) Cohort of stage IV lung cancer patients who received ICB monotherapy. F-J) Tumor proportion score for PD-L1, tumor mutation burden, %CD8 (among CD45+), PD-1 mean fluorescence intensity (MFI) and % CD39 (among CD8 T cells) among non-responders (stable disease/progression of disease, SD/POD) or responders (partial response/complete response, PR/CR) to ICB in the cohort described in E). K-M) Kaplan-Meier survival curve of progression-free survival patients described in cohort E based on stratification for CD8, PD-1, and CD39. Statistical significance was determined by Mantel-Cox test.

Figure S1. Single-cell CITE/TCR/RNA-sequencing reveals that CD39hi CD8 T cells are enriched for features of tumor, but not viral, reactivity.

A) Correlation of transcriptional expression (y axis) with protein expression (x axis) for indicated genes. B) Pooled transcript levels (scaled counts) of indicated genes in CD39hi, CD39int, and CD39neg CD8 T cells as classified by adt_CD39 levels. C-D) Scaled scores for MANA-TIL and influenza- reactivity gene signatures (see Table S4 for full gene list). Statistical significance was determined by two-way ANOVA with Tukey's multiple comparisons test and p value is indicated if < 0.05.

Figure S2. Clinical and molecular correlates of CD39 expression on CD8 tumor-infiltrating lymphocytes.
A-D) Correlation of %CD39 on CD8 T cells with %CD8 (among CD45+), tumor mutation burden, PD-L1 tumor proportion score, and smoking history. E-F) %CD39+ (among CD8 T cells) and %CD8 (among CD45+) on four subgroups based on cutoffs for TMB and PDL-1 in the 440-sample cohort. Statistical significance was assessed by two-tailed student's t-test. G) Correlation of %CD39 on CD8 T by lung cancer stage. H) Correlation of %CD39 on CD8 T by anatomic site of biospecimen. I-J) Correlation of tumor mutation burden and PD-L1 tumor proportion score with various histological subtypes/driver mutation categories among 440-sample cohort. Red bars indicate histological subtypes/driver mutations with above average %CD39 values.

Figure S3. CD39 is upregulated in the tumor microenvironment and is consistent across space and time. A-C) Top row depicts flow cytometry histogram plots of CD39 levels on DAPI⁻ CD45⁺ CD3⁺ CD8⁺ T cells from peripheral blood or matched tumor lesion from three patients with lung cancer. Bottom row depicts Morisita indices depicting TCR-β overlap among sorted CD39⁻ and CD39⁺ fractions of blood and tumor CD8 T cells. D) %CD39 among CD8 T cells from two spatially distinct lesions that were sampled concurrently in the same patient (n=12). Blue line indicates MSK 1266 whose CD39 expression is depicted in E. Purple line indicates MSK 1372 depicted in F. E) Flow cytometry plots of CD39 expression on CD8 T cells on two concurrent pleural lesions in MSK 1266. F) Flow cytometry plots of CD39 expression on CD8 T cells in two concurrent spatially distinct lung lesions in MSK 1372. G) %CD39 among CD8 T cells from two lesions from the same anatomic site that were sampled at temporally distinct times in the same patient (n=9). Orange line indicates MSK 1265 whose CD39 expression is depicted in H. H) Flow cytometry plots of CD39 expression on CD8 T cells over the course of three brain resection lesions in MSK 1265.

Figure S4. Total CD8 tumor-infiltrating lymphocytes is not prognostic for survival in lung cancer. A-C) Kaplan-Meier curve of recurrence-free survival stratified by %CD8 (among CD45+) for stage I, stage II, and stage IIIA lung cancer patients who did not receive ICB. Statistical significance was determined by Mantel-Cox test. D) Kaplan-Meier curve of progression-free survival stratified by %CD8 (among CD45+) for stage IV lung cancer patients who did not receive ICB. Statistical significance was determined by Mantel-Cox test.