Repressing ABCB7 Potentiates Cisplatin Response in Pediatric Group 3 Medulloblastoma by Triggering Ferroptosis

Ranjana K. Kanchan[1], Naveenkumar Perumal[1], Parvez Khan[1], David Doss[1,3], Prakadeeshwari Gopalakrishnan[1], Ramakanth Chirravuri Venkata[1], Ishwor Thapa[4], Raghupathy Vengoji[1], Jyoti B. Kaushal[1], Jawed A. Siddiqui[1], Mohd Wasim Nasser[1], Surinder K. Batra[1], and Sidharth Mahapatra [1,5]

[1] Department of Biochemistry, University of Nebraska Medical Center, Omaha, NE 68198, U.S.A
[2] School of Interdisciplinary Informatics, University of Nebraska at Omaha, Omaha, NE.
[3] School of Medicine, Creighton University, Omaha, NE 68178, USA
[4] School of Interdisciplinary Informatics, University of Nebraska at Omaha, Omaha, NE
[5] Department of Pediatrics, Children's Hospital and Medical Center, Omaha, NE 68114, U.S.A.

Corresponding author: Sidharth Mahapatra, MD, PhD, Department of Pediatrics, University of Nebraska Medical Center, 601 South Saddle Creek Road, Omaha, Nebraska, email: sidharth.mahapatra@unmc.edu

Author contributions: RKK and SM conceptualized and designed the study. RKK, JBK, NP, DD, and PG performed all experiments. PK developed ABCB7 knockout clone for the study. DD, RCV, and IT provided bioinformatics support for the study. SM coordinated and supervised data analysis. RKK wrote the first draft of the manuscript. NP, PK, DD, PG, RCV, JAS, RV, MWN SKB and SM provided scientific feedback, critical reviews, and helped in revision of the manuscript. All authors have read and approved the final manuscript.

Funding Information: This work was supported by the National Institutes of Health (NICHD K12HD047349); the Edna Ittner Pediatric Research Support Fund; and the Team Jack Brain Tumor Foundation.

Conflicts of Interest: Authors declare that they have no competing interests**.**

Abbreviations: ABC, ATP binding cassette; ANOVA, analysis of variance; BLC-2, B-cell lymphoma 2; BRCA1, breast cancer gene 1;CB, cerebellum; CD276, cluster of differentiation 276 (B7-H3); CDK6, cyclin-dependent kinase 6; c-Myc, c-myelocytomatosis oncogene; DDR, DNA damage repair; DCFDA, 2′,7′-Dichlorofluorescin diacetate; DFO, deferoxamine; FACs, fluorescence-activated cell sorting; FISH, fluorescence *in situ* hybridization; GPX4, glutathione peroxidase 4; GSH, glutathione; GSS, glutathione synthetase; IC_{50}, 50% inhibitory concentration; IHC, immunohistochemistry; i(17q), isochromosome 17q; KEGG, Kyoto encyclopedia of genes and genomes; KO, knock-out; LIP, labile iron pool; LPO, lipid peroxidation; MB, medulloblastoma; MDR, multiple drug resistance; miR, microRNA; miR-1253, microRNA 1253; MnTBAP, Mn(III)tetrakis(4-benzoic acid)porphyrin; mtROS, mitochondrial reactive oxygen species; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; NC, negative control; NHA, normal human astrocytes; Non-SHH/WNT, non-Sonic Hedgehog/non-Wingless; PARP, poly ADP ribose polymerase; PCR, polymerase chain reaction; Ped, pediatric; PNET, primitive neuro-ectodermal tumor; ROS, reactive oxygen species; RSL3, RAS-selective lethal; SHH, Sonic Hedgehog; WNT, Wingless; XIAP, X-linked inhibitor of apoptosis; XTT, sodium 3′-[1-[(phenylamino)-carbony]-3,4-tetrazolium]-bis(4-methoxy-6-nitro)benzene-sulfonic acid hydrate

Abstract

Medulloblastomas (MB) are the most common malignant pediatric brain tumor and a leading

cause of childhood mortality. Aggressive tumors belonging to group 3 (G3MB) are distinguished

by a marked reduction in programmed cell death by ferroptosis. These aggressive tumors also

enrich iron transport and glutathione metabolism. A highly enriched pathway in these tumors is

iron-sulfur (Fe-S) cluster binding corresponding to significant upregulation of ABCB7, a poor

prognostic feature linked with accelerated mortality. This study elucidated whether repressing

ABCB7 activates ferroptosis to mitigate G3MB aggressiveness and whether this pathway is

pharmacologically targetable. *In silico* and *in vitro* analyses confirmed upregulation of ABCB7

and GPX4, the central regulator of ferroptosis, in G3MB cell lines and tumors. Repressing

ABCB7 (miR-1253^{OE}, siABCB7, sh-ABCB7) induced iron overload, elicited oxidative stress,

and triggered lipid peroxidation, leading to an abrogation of medullosphere formation and cell

death by ferroptosis. Intriguingly, fractionation studies revealed that ABCB7 repression

abrogated GPX4 expression, most likely by GSH depletion. Repressing ABCB7 induced

mitochondrial dysfunction and reduced oxidative phosphorylation. Cisplatin, a chemotherapeutic

mainstay of G3MB, induces cell death by DNA crosslinking. In ABCB7 repressed cell lines, the

IC$_{50}$ of cisplatin was halved, resulting in augmented oxidative stress and lipid peroxidation,

culminating in a higher index of ferroptosis. Artesunate, an anti-malarial drug capable of

triggering ferroptosis, was shown to synergize with cisplatin, reducing tumor burden and

significantly prolonging survival. Taken together, the current study illustrates how targeting iron

transport can augment ferroptosis in G3MBs. It further identifies an FDA-approved drug capable

of recapitulating these effects and potentiating mainstay chemotherapy.

Key words: ABCB7; ferroptosis; GPX4; medulloblastoma; miR-1253; ROS

Introduction

Of the primitive neuro-ectodermal tumors (PNET) afflicting the posterior fossa, medulloblastoma (MB) is the most common malignant tumor of childhood, accounting for 85% of PNETs and 20% of all posterior fossa tumors.[1-3] Based on high-throughput gene profiling studies, MBs are stratified into four major molecular subgroups: WNT, SHH, group 3 and group 4.[4] Among these, patients with group 3 tumors (G3MB) suffer the worst prognosis (5-year overall survival <50%) due to a highly aggressive phenotype punctuated by c-Myc amplification, haploinsufficiency of chromosome 17p, presence of metastatic lesions at diagnosis, and high rates of recurrence.[4-10] Recurrence can further reduce 5-year overall survival to <10% and is partially fueled by the inability of young patients to tolerate effective chemo- and/or radiation therapy.[4,11-13] For instance, systemic toxicity is an oft-cited justification for limitations in cisplatin dosing, especially for group 3 disease, leading to <20% of treated patients receiving an effective dose.[14-16] These facts underscore the urgent need to develop new strategies for high-risk disease.

Recently, combining microRNAs with chemotherapy has garnered support in enhancing therapeutic efficacy via targeting critical regulators of DNA damage repair (DDR), apoptosis, or cell cycle.[17] In breast cancer, for example, miR-9 and miR-218 improved cisplatin responsiveness by targeting BRCA1 and impeding DNA damage repair.[18,19] The miR-15 family sensitized cisplatin-resistant cancer cells to apoptosis by targeting the G_2/M cell cycle checkpoint.[20] In gastric and lung cancer cell lines, the miRNA cluster miR-200bc/429 sensitized resistant cell lines to both vincristine and cisplatin by targeting BCL2 and XIAP.[21] In a gemcitabine-resistant pancreatic xenograft model, miR-205 delivered with gemcitabine in conjugated micelles resulted in significant tumor growth inhibition.[22] Most notably, miR-34a

replacement therapy (MRX34) has reached a phase 1 clinical trial (NCT01829971) in patients

with primary unresectable or metastatic liver cancer.[23]

Group 3 tumors have amongst the highest frequency of cytogenetic aberrations targeting the

short arm of chromosome 17 compared to the other subgroups, with high incidence reported on

locus 17p13.3.[24-27] We recently revealed strong tumor suppressive properties for miR-212 and

miR-1253, which reside on locus 17p13.3.[27,28] We showed that miR-1253 targets the cell cycle

checkpoint protein, CDK6, and CD276 (B7-H3), an immune checkpoint molecule implicated in

tumor aggressiveness. [27,29-31] Aside from reducing tumor cell viability, migration/invasion, and

colony formation, miR-1253 arrested cells at the G_0/G_1 phase, activated apoptotic pathways, and

triggered oxidative stress.[27] Numerous studies have recapitulated these strong tumor suppressive

properties in various cancers, including non-small cell lung carcinoma[32], osteosarcoma[33],

pancreatic cancer[34], male breast cancer[35], and colon cancer[36].

Amongst the putative targets of miR-1253 are the ATP-binding cassette (ABC) transporters

that confer multi-drug resistance to various tumors, including MB.[37,38] In metastatic breast

cancer, for example, miR-1253 was shown to inhibit the drug efflux pump, ABCB1, thereby

potentiating the cytotoxicity of doxorubicin.[39] However, this particular ABC transporter is not

amongst the deregulated ABC transporters reported in G3MB.[40] Another putative target of miR-

1253 is ABCB7, an iron transporter residing on the inner mitochondrial membrane involved in

iron homeostasis and Fe-S cluster biogenesis.[41] Deregulation of ABCB7 has been shown to help

cancer cells withstand apoptosis and ferroptosis.[42] Ferroptosis is a form of cell death triggered by

iron-mediated oxidative stress leading to lethal lipid peroxidation. This process is tightly

controlled by glutathione peroxidase 4 (GPX4), whose primary substrate is the anti-oxidant,

glutathione (GSH).[43-45] Recent studies have identified ferroptosis mitigation in tumor progression

and drug resistance.[46,47]

Of the standard chemotherapeutic agents used in the treatment of MB, cisplatin, a platinum-

based alkylating agent that induces DNA damage, has been shown to trigger cell death via

oxidative stress and ferroptosis.[46,48] Whether deregulation of ABCB7 leads to cisplatin resistance

in group 3 tumors or if miR-1253 can sensitize cisplatin response through inhibition of ABCB7

remains unstudied. Here, we show that targeting ABCB7 can potentiate cisplatin cytotoxicity in

G3MB by inducing iron overload, oxidative stress, and lipid peroxidation, which together trigger

ferroptosis.

Materials and Methods

Patient Samples

Formalin-fixed paraffin embedded tissue blocks and frozen tissues of normal cerebellum

(pediatric=12, adult=5) and pediatric MB specimens (WNT=1, SHH=9, grp 3=11, grp 4=16,

unknown=7) were obtained from the Children's Hospital and Medical Center, Omaha and the

University of Nebraska Medical Center after Institutional Review Board approval. Informed

consent was not required since the status of the study was exempted. For expression profiles of

ABCB7 and GPX4, we cross-analyzed two primary MB datasets (Kanchan *et al.*, GSE148390

and Weishaupt *et al.*, GSE124814).[49-51] For Spearman correlation, we used GSE148390. For

Kaplan-Meier Survival Analysis, we used the R2 database (Cavalli *et al.*, GSE85217).[9]

Cell Lines and Cell Culture

D283 and D341 were purchased from ATCC (Manassas, Virginia); D425 and D556 were kind

gifts from Darell Binger (Duke University Medical Center, Durham, NC); HDMB03 cells were a

kind gift from Till Milde (Hopp Children's Tumor Center, Heidelberg, Germany). Cell line

genotyping was verified using short tandem repeat (STR) DNA profiling (UNMC). D283, D341,

D425 and D556 cell line were maintained in DMEM supplemented with 10%-20% FBS and

100µg/ml penicillin/streptomycin. Normal human astrocytes (NHA) were purchased from Lonza

Bioscience (Walkersville, MD) and grown in ABM basal medium supplemented with growth

factors (Lonza Biosciences). All cell lines were maintained in 95% humidity, 37°C, 5% CO_2.

Transient Transfections

Cells at a density of 0.5×10^6 were seeded in 6-well plates for 24 h and subsequently serum

starved for 4 h prior to transfection. Cells were transfected with miR-1253 mimic (miRVanaTM

miRNA mimic, ThermoFisher, 100 nM) or scramble negative control (100 nM) with

Lipofectamine 2000 (Invitrogen) for 24 h.

CRISPR/Cas9 Knockouts

Lentiviral particles were prepared by transfection of plasmid expressing Cas9 or sgRNA of

ABCB7 (Addgene) co-transfected with pCMV-dR8.2 dvpr (Addgene) and pCMV-VSV-g

(Addgene) lentiviral packaging plasmids into HEK293T cells using polyethyleneimine (PEI)

transfection reagent. Virus-containing supernatant was collected and filtered 48 h after

transfection. HDMB03 cells in 6-well plates were infected with Cas9 viral supernatant

containing 4 μg/mL polybrene. Following blasticidin selection (10 μg/ml), the expression of

Cas9 was confirmed by Western blotting. Stable HDMB03 cells expressing Cas9 expression

were infected with the ABCB7 sgRNA viral supernatant containing 4 μg/ml polybrene. After 24

h of infection, cells were selected with 0.5 μg/ml puromycin. Single cell clones with Cas9

expression and ABCB7 knockout were amplified and used for subsequent experiments.

Cell Viability

After transfection, HDMB03 cells (5×10^3 cells/well) were re-seeded into 96-well plates and

treated with defined concentrations of cisplatin (1-50 μM) for 24-72 h. Subsequently, MTT (5

mg/mL) or XTT (0.3 mg/mL) was added to each well and incubated for 2 or 6 h, respectively, at

37 °C. MTT absorbance was measured at 570 nm; XTT absorbance was measured at 440 nm.

Data were analyzed using the SOFTMAX PRO software (Molecular Devices Corp., Sunnyvale,

CA, USA).

Colony Formation

163   After transfection, HDMB03 cells (1×10^3 cells/well) were re-seeded in 6-well plates. Cells were

164   treated with cisplatin (2 μM) and grown in complete medium for 9 days. Colonies were stained

165   with 0.25% crystal violet (dissolved in 50% methanol) for 30 min. Crystal violet was dissolved

166   in 10% acetic acid and absorbance read at 590 nm.

Luciferase Assay

Luciferase assay was performed as previously described.[27] Primers ABCB7 Forward: 5'-

TAAGCCTGACATAACGAGGA-3'; ABCB7 Reverse: 5'-GCATCTCAGTATTAACTCTAGC

-3') were purchased from Eurofins. 3'UTR Wild and 3'UTR-Mutant were incorporated into

XbaI restriction site of PGL3-control vector (Promega) expressing firefly luciferase. Dual

luciferase assay was done in HDMB03 cells (3×10^5 cells/well) in 12-well plates. Luciferase

activity was the measured using Dual-Luciferase Reporter Assay System (Promega) with a

Luminometer (Biotek).

Calcein AM Staining

Calcein-acetoxymethyl ester (Calcein AM) is a membrane permeable, non-fluorescent dye which

emits green fluorescence once internalized into the live cells and cleaved by cytoplasmic

esterase. Although calcein fluorescence is stable, it can be quenched by divalent metal ions such

as iron and cobalt. To estimate cytosolic labile iron pool (LIP), HDMB03 and D425 cells were

either transfected with scramble or miR-1253 or had ABCB7 knocked down. Cells were reseeded

on to glass coverslips treated with or without the iron chelator deferoxamine (DFO) for 6 h. Cells

were washed twice with 0.5 ml of PBS and incubated with 0.05 μM Calcein AM for 15 min at 37

°C. Cells were analyzed under confocal microscope. Ex/Em = 488 nm/525 nm.

Detection of Cytosolic and Mitochondrial Fe^{2+}

Scramble vs. miR-1253-transfected cells or wild-type vs. ABCB7^{KO} cells were seeded onto glass

coverslips and were treated with or without DFO (100 μM) for 6 h. For cytosolic LIP, 1 μmol/L

FerroOrange (Dojindo, Japan) and 100 nM MitoTraker^{TM} Deep Red FM were added to each

well; for mitochondrial LIP, 1 μM/L Mito-FerroGreen (Dojindo, Japan) and 100 nM MitoTraker

Deep Red FM were added to each well. Cells were incubated in a 37^{o} C incubator equilibrated

with 95% air and 5% CO_2 for 30 min. After incubation cells were washed and counter-stained

with 4′,6-diamidino-2-phenylindole (DAPI). Cells were observed under a confocal fluorescence

microscope, Ex/Em$_{cytosolicLIP}$ = 561 nm/570-620 nm, Ex/Em$_{mitochondrialLIP}$ = 488 nm/500-550 nm.

Measurement of Intracellular Oxidative Stress

Intracellular ROS (H_2O_2) was measured using oxidation-sensitive fluorescent probe 2′,7′-

Dichlorofluorescin diacetate (DCFDA) (Sigma Aldrich, USA); mitochondrial $O_2^{\bullet-}$ was measured

using MitoSOX™ Red (Thermofisher, USA). Scramble vs. miR-1253-transfected cells or wild-

type vs. ABCB7^{KO} cells were treated with cisplatin or cisplatin and Ferrostatin-1 (Apexbio,

USA) for 24 h. Cells were incubated with 10 μM DCFDA or 5 μM MitoSOX™ Red for 30 min.

Oxidized DCFDA and MitoSOX™ Red were measured using at Ex/Em ~485/528 nm and

Ex/Em ~510/580, respectively.[52] Images were captured using EVOS FL Auto Imaging System

(EVOS FL Auto, Life Technologies). Oxidized DCFDA and MitoSOXRed were measured using

multimode plate reader at Ex/Em ~485/528 nm and Ex/Em ~510/580, respectively

Measurement of Lipid Peroxidation

The Image-iT® LPO kit was used to measure lipid ROS through oxidation of the C-11-

BODIPY® 581/591 sensor according to the manufacturer's instructions. Briefly, scramble vs.

miR-1253-transfected cells or wild-type vs. ABCB7^{KO} cells were treated with or without

cisplatin or a combination of cisplatin and Ferrostatin-1 for 24 h. Cells were then stained with

Image-iT® Lipid Peroxidation Sensor (10 μM) for 30 min and counter-stained with DAPI.

Images were captured by confocal microscopy.

Ferroptosis Assessment by Flow Cytometry

Scramble and miR-1253-transfected cells were treated with or without cisplatin or a combination

of cisplatin and Ferrostatin-1 for 24 h. After completion of treatment, cells were incubated with 5

μM of MitoSOX™ Red for 30 min. Cells were washed and stained with Annexin-V/Cy™5(BD

Biosciences). Cell populations were sorted and measured by flow cytometry.

Glutathione Estimation

Glutathione estimation was performed according to the manufacturer's instructions using GSH-

Glo™ Glutathione Assay kit (Promega, USA). Briefly, scramble vs. miR-1253-transfected cells

or wild-type vs. ABCB7^{KO} cells were seeded (5×10^3 cells/well) in the 96-well plates. Cells were

treated with or without cisplatin or in combination with cisplatin and Ferrostatin-1 for 24 h.

Plates were incubated in the dark for 30 min on a plate shaker with 100 µl of prepared 1x GSH-

Glo™ Reagent. Then, 100 µl of reconstituted Luciferin Detection Reagent was added to each

well and incubated in the dark for another 15 min. A standard curve was prepared using GSH

standard solution to facilitate the conversion of luminescence to GSH concentration.

Immunofluorescence Imaging

In cultured cells: scramble vs. miR-1253-transfected cells or wild-type vs. ABCB7^{KO} cells were seeded onto coverslips, rinsed, and fixed using 4% paraformaldehyde (PFA) in PBS for 10 min at room temperature. Cells were washed, incubated with 0.25% Triton X-100, and blocked with 3% BSA at room temperature. Cells were then incubated with primary antibody ABCB7 or COXIV overnight. Following day, cells were incubated with Alexafluor 488 and Alexaflour 547 conjugated antibodies for 1 h. Cells were washed and counter stained with DAPI. Images were captured at 63X using Carl Zeiss microscope (LSM 800 META).

In tissue: immunofluorescence was performed on surgically-resected, formalin-fixed, paraffin-embedded sections of group 3 medulloblastomas. Deparaffinized tissue sections were blocked with 5% BSA after heat induced epitope retrieval with citrate buffer (pH 6.0) and then incubated with primary antibodies, ABCB7 rabbit monoclonal antibody (1:200), GPX4 rabbit monoclonal antibody, or COXIV mouse monoclonal antibody (1:200). Following overnight incubation with primary antibody, sections were incubated for 1 h with Alexa-488 or Alexa-547-conjugated, mouse and rabbit secondary antibodies (1:200). Sections were counter-stained with DAPI. Micrographs were captured at 20X using Carl Zeiss microscope (LSM 800 META).

PCR Primers

Total RNA extraction and quantitative PCR (qPCR) were performed according to manufacturer protocol. The forward (F) and reverse (R) primer sequences were as follows: ABCB7 (F): 5'-AAGATGTGAGCCTGGAAAGC-3' (R): 5'- AGAGGACAGCATCCTGAGGT-3'; GPx4 (F): 5'-ACAAGAACGGCTGCGTGGTGAA-3' (R): 5'-GCCACACACTTGTGGAGCTAGA-3'; β-Actin (F): 5'-CACCATTGGCAATGAGCGGTTC-3'(R): 5'-AGGTCTTTGCGGATGTCCACGT-3'.

 Statistical Analysis

 Data are presented as mean ± SD. All experiments were conducted at least in duplicates with 3-6

 replicates per experiment. Statistical analyses were performed using Prism 9.2 (GraphPad

 Software). Differences between groups were compared using Student's t-test or one-way analysis

 of variance (ANOVA). Statistical significance was established at *p <0.05; **p <0.01; ***p

 <0.001; ****p <0.0001. Statistical analyses of high-throughput sequencing data were performed

 using R Statistical Software v4.1.1 (R Core Team), expression values were compared using a

 Mann-Whitney U test.

 Results

 Ferroptosis evasion characterizes G3MBs.

 Iron is vital to cellular processes, such as DNA synthesis, proliferation, cell cycle regulation,

 electron transport and catalysis, and oxygen transport and delivery.[53,54] Rapidly dividing cancer

 cells can reprogram iron metabolism by upregulating essential iron regulatory and transport

 genes to support their invasive and rapid growth.[54] Ferroptosis occurs when free ferrous iron

 oxidizes lipid membranes.[55] Ferroptosis evasion, in turn, would result from mitigating iron

 imbalance and oxidative stress.

 Using single sample gene set enrichment (ssGSEA) in a large MB dataset (GSE85217)[9], we

 first observed that ferroptosis was the most enriched programmed cell death (PCD) pathway in

 MB tumors compared to normal **(Supplementar Figure 1A)**. We then compared the survival of

 patients in this cohort based on their ferroptosis scores, a measure of how active this pathway is.

 We observed that patients with lower ferroptosis scores (suggestive of lower activation) had

 lower survival **(Supplementary Figure 1B)**. Amongst the 4 MB subgroups, groups 3 and 4 have

 significantly lower survival than WNT and SHH tumors.[56] And between these two subgroups,

G3MBs had significantly lower ferroptosis scores than G4MB **(Supplementary Figure 1C)**. Our

data suggest that lower ferroptosis activation (ferroptosis evasion) may be associated with more

aggressive tumors.

When comparing G3MBs and normal cerebellum, we discovered enrichment of iron

homeostasis (IREB2, STEAP3), iron transport (TFRC, ABCB6-8, SLC40A1, SLC25A37),

glutathione metabolism (GCLC, GSS, TRIT1, SEPSECS), and ferroptosis regulatory genes that

inhibitor ferroptosis. GPX4, the central regulator of ferroptosis[57], was also enriched in our

analysis **(Supplementary Figure 1D)**. We then queried the Molecular Signature Database

(MSigDB, version 7.4) for upregulated pathways associated with the isolated genes. We

discovered the enrichment of iron-sulfur (Fe-S) cluster binding and downregulation of ROS

pathways **(Supplementary Figure 1E)**. Together, our data suggest that G3MBs, like many

aggressive cancers, enrich iron homeostasis genes and redox pathways.[57,58] This gene enrichment

signature may enable G3MB tumors to counter iron imbalance and withstand oxidative stress

and ferroptosis during their unchecked growth, enhancing their proliferative and aggressive

phenotype.

ABCB7, a novel target of miR-1253, is enriched in group 3 medulloblastomas.

In our previous study, we identified miR-1253 as a novel tumor suppressor gene in MB,

along with its multiple oncogenic targets.[27] Upon restoration of miR-1253, using transient

overexpression in HDMB03 cells, we observed potent downregulation of the ABC transporter

superfamily by KEGG pathways analysis **(Supplementary Figures 2A)**. We wanted to then

isolate specific targets of miR-1253 relevant to G3MB aggressiveness. We began by identifying

ABC transporters enriched in G3MB **(Supplementary Figure 2B, column 1)**. From this list, we

isolated transporters whose high expression conferred a poor prognosis **(Supplementary Figure**

2B, column 2, red). From this cohort, we selected transporters whose expression was inhibited

by miR-1253 overexpression in HDMB03 cells (**Supplementary Figure 2B, column 3, red**).

This comparative analysis revealed ABCB7.

ABCB7 resides on the inner mitochondrial membrane and plays a key role in Fe-S cluster

biosynthesis and export.[59-61] Notably, our transcriptomic analysis isolated Fe-S cluster binding as

a top deregulated pathway and ABCB7 as a top enriched gene in G3MB tumors. In multiple MB

cohorts, ABCB7 was shown to be upregulated (**Supplementary Figure 2C).** We then compared

subgroup-specific ABCB7 expression in 3 MB cohorts and identified consistent enrichment of

ABCB7 in G3MB (**Figure 1A; Supplementary Figures 2D and 2E**). High ABCB7 expression

in MB tumors was associated with dramatically poorer overall survival (**Supplementary Figure**

2F). When the same survival analysis was run in a subgroup-specific manner, we learned that

ABCB7 upregulation is associated with poor survival only in G3MB tumors (**Figure 1B**). These

findings strengthened our premise to study ABCB7 enrichment as a possible mechanism for

G3MB aggressiveness.

Now focusing on ABCB7, we first confirmed by confocal microscopy that ABCB7 co-

localizes to the mitochondria, as evidenced by co-staining with COXIV, an inner mitochondrial

membrane protein critical to electron transport (**Figure 1C**). Additionally, we confirmed robust

expression of ABCB7 in a panel of aggressive group 3 MB cell lines compared to normal human

astrocytes (**Figure 1D**).

To demonstrate that ABCB7 is inhibited by miR-1253, we first showed translational

repression via Western blotting in 2 group 3 MB cell lines, D425 (**Figure 1E**) and HDMB03

(**Figure 1F**). In the same cell lines, we showed co-localization of ABCB7 with COXIV to the

mitochondria and visualized the effect of miR-1253 on ABCB7 expression, as illustrated by a

dramatically reduced green fluorescence by confocal microscopy **(Figure 1G)**. Finally, we

showed direct binding of miR-1253 to ABCB7 via a dual-luciferase reporter assay **(Figures 1H)**.

Together, these data confirmed that i) ABCB7 expression is enriched in group 3 tumors, ii)

ABCB7 enrichment is a poor prognostic sign in group 3 patients, and iii) miR-1253 effectively

inhibits ABCB7 expression.

ABCB7 repression attenuates G3MB cancer cell aggressiveness.

While the pathogenic role of ABCB7 has yet to be elucidated in MB, in glioma cells, ABCB7

inhibition triggered both apoptotic and non-apoptotic cell death through iron overload and

ROS.[42] Given the promiscuous nature of miRs, we adopted additional strategies to repress

ABCB7 expression in G3MB cancer cells by using small-interfering RNA (si-ABCB7) or dox-

inducible knockdown **(Supplementary Figure 2G)**. Using these strategies, we observed potent

inhibition of primary medullosphere formation **(Figure 2A)** and reduced cancer cell viability by

MTT assay **(Figure 2B)**, all evidencing that ABCB7 repression attenuates aggressiveness of

HDMB03 cells.

ABCB7 repression triggers iron overload.

Given its primary role as a Fe-S cluster exporter, we studied the impact of repressing

ABCB7 expression on iron balance in HDMB03 cells using ferric ammonium citrate (FAC) to

mimic iron overload[62] as our positive control. We observed an accumulation of free iron in the

mitochondria (**MitoFerroGreen, Figure 3A and Supplementary Figure 3A**) and the cytoplasm

(**FerroOrange, Figures 3B, 3C, and Supplementary Figure 3B**) in HDMB03 cells with

repressed ABCB7 expression. Iron chelation using deferoxamine (DFO) reverted the phenotype

to wild-type. Calcein AM is a membrane-permeable dye that produces green fluorescence when

internalized and can be rapidly quenched by divalent metals like iron. Repression of ABCB7 by

miR-1253 led to quenching of Calcein AM in D425 **(Supplementary Figure 3C)** and HDMB03

cells **(Supplementary Figure 3D)**, further strengthening our observations of iron overload

induced by ABCB7 inhibition.

ABCB7 repression enhances cellular ROS and induces lipid peroxidation resulting in

ferroptosis.

Free ferrous iron (Fe^{2+}) can convert to toxic ferric iron (Fe^{3+}), via the Fenton reaction, which

can disrupt the lipid bilayer to generate lethal lipid hydroperoxides, culminating in programmed

cell death (PCD) by ferroptosis.[55,57] Thus, after showing iron overload via ABCB7 repression,

we subsequently quantified mitochondrial and total ROS in HDMB03 cells using hydrogen

peroxide and erastin (ERA), a known inducer of ferroptosis[63], as our positive controls. We

observed elevated mitochondrial ROS (**Mitosox**, **Figure 3D and Supplementary Figure 4A**)

and total ROS (**DCFDA**, **Figures 3E and Supplementary Figure 4B**) in ABCB7-repressed

HDMB03 cells. We showed the same impact of ABCB7 repression by miR-1253 on oxidative

stress in D425 cells **(Supplementary Figures 4C)**.

We next quantified lipid peroxidation through BODIPY-C11 staining, a membrane-targeted

lipid ROS sensor, using an Image-iT^{TM} lipid peroxidation kit which provides a ratiometric

indication of fatty acid oxidation that can be quantified by microscopy and flow cytometry

(FACS). Repressing ABCB7 elicited lipid peroxidation in HDMB03 **(Figures 3G and 3H)** and

D425 cells **(Supplementary Figure 4D)**.

Moreover, in the setting of mitochondrial iron overload and ROS, we studied the impact of

repressing ABCB7 on mitochondrial membrane potential and oxidative phosphorylation

capacity. Using tetramethylrhodamine methyl ester (TMRM), which fluoresces within

mitochondria with intact membrane potentials, we showed a dramatic reduction in signal with si-

ABCB7, suggesting that ABCB7 repression leads to mitochondrial dysfunction **(Figure 4A)**.[64,65]

ABCB7 repression by miR-1253^{OE} also impaired the oxygen consumption rate (OCR) of

HMDB03 cells, a direct measure of mitochondrial ATP production. It concurrently increased the

extracellular acidification rate (ECAR), a measure of ATP production under anaerobic conditions

by glycolysis **(Figure 4B)**.

Ferroptosis is defined as an iron-dependent form of regulated cell death resulting from lipid

peroxidation.[43] Ferrostain-1 (FER) scavenges ROS and lipid hyperoxides and prevents

ferroptosis.[66] Given our observations of iron accumulation, ROS generation, and lipid

peroxidation, we wanted to assess whether ABCB7 repression can, in fact, trigger ferroptosis.

D425 and HDMB03 cells transfected with miR-1253 were stained with MitoSOX™ Red (for

$O_2^{\bullet-}$) and Annexin-V/Cy™5 (marker of apoptotic cell death). Flow cytometry analysis revealed a

significant (~3-fold) rise in dual-stained cells (Q2), indicating oxidative stress-mediated cell

death **(Supplementary Figures 4E and 4F)**. Moreover, ferrostatin-1 rescued ABCB7-repressed

cells from both oxidative stress **(Figures 3D, 3E, and Supplementary Figures 4A and 4B)** and

lipid peroxidation **(Figure 3G and 3H)**. These are the <u>first data</u> showing that ABCB7 repression

can trigger events that culminate in ferroptosis.

ABCB7 repression inhibits G3MB tumor growth.

We next examined the impact of ABCB7 repression on tumor burden in a validated G3MB

mouse model[28] using dox-inducible ABCB7 knockdown. We generated tumors in NOD *scid*

gamma (NSG) mice by orthotopically injecting wildtype or ABCB7^{KD} HDMB03 cells (1x10^5)

into the cerebellum. We observed smaller tumors sizes in dox-induced ABCB7^{KD} mice at day 28

and day 30 **(Figures 5A and 5B)**. This was concurrent with significantly lower

immunohistochemical staining for ABCB7 and GPX4, as well as higher staining for 4HNE, a

marker of lipid peroxidation (**Figure 5C**). These data reveal that ABCB7[KD] attenuates G3MB

tumor burden.

ABCB7 repression inhibits GPX4, the primary regulator of ferroptosis.

Cancer cells have the inherent ability to activate redox buffering systems to thrive in a highly

oxidative environment.[67] Glutathione (GSH) is a key player in this response and a critical

cofactor for glutathione peroxidase 4 (GPX4). GPX4 uses GSH to reduce toxic lipid

hydroperoxides to non-toxic lipid alcohols and is thus a central regulator of ferroptosis, found in

the nucleus, cytoplasm, and mitochondria (**Supplementary Figure 5A**).[57] In G3MB, ROS

pathways are downregulated and concurrent with elevated expression of genes involved in

glutathione metabolism, including GSS, GCLC, GPX4 (**Supplementary Figures 1D and 1E**).

GPX4 overexpression elevates aggressiveness in multiple cancers (breast, liver, GI,

pancreatic).[68] In MB, we noted significant enrichment of GPX4 in all MB tumor subtypes, with

highest expression in G3MB (**Supplementary Figures 5B and 5C**). By RNA Sequencing

analysis, critical mediators of GSH biosynthesis (GSS, GCLC) and GPX function (SEPSECS,

TIRT1) were also upregulated in G3MB (**Supplementary Figure 5D**). Immunofluorescence

studies revealed high expression of GPX4 in G3MB and co-localization of GPX4 to the

mitochondria (**Supplementary Figure 5E**). In classic G3MB cell lines, GPX4 expression was

significantly elevated compared to normal human astrocytes (**Supplementary Figure 5F**).

Given the pivotal role of GPX4 in ferroptosis mitigation, we studied the impact of ABCB7

repression on GPX4, expecting an upregulation to counter the oxidative stress induced by iron

overload. Instead, we observed the opposite. While miR-1253[OE] reduced GPX4 transcript and

protein levels in HDMB03 (**Figure 6A**) and D425 (**Figure 6B**), ABCB7[KO] (**Figure 6C**) and

ABCB7[KD] (**Figure 6D**) abrogated expression. In support, we noted a positive Spearman

correlation (R=0.72, $p<0.001$) between GPX4 and ABCB7 **(Figure 6E)**. We further explored

this relationship using fractionation studies. While miR-1253[OE] reduced GPX4 levels in the

cytoplasm and mitochondria **(Figure 6F)**, ABCB7[KO] **(Figure 6G)** and si-ABCB7 **(Figure 6H)**

again showed near abrogation of GPX4 expression from the cell. GSH, a main substrate of

GPX4, was concurrently reduced in miR-1253-overexpressing D425 **(Supplementary Figure**

6A) and HDMB03 cells **(Supplementary Figure 6B**) as well as in si-ABCB7-treated HDMB03

cells **(Supplementary Figure 6C**). This is the <u>first time</u> a link has been proposed between

ABCB7 and GPX4, providing us preliminary insight into the impact of the miR-1253/ABCB7

axis on GPX4 and the cellular response to ferroptosis.

ABCB7 repression potentiates cisplatin cytotoxicity in G3MB cells.

Cisplatin, a platinum-containing chemotherapeutic agent, induces DNA damage via various

mechanisms, including (i) crosslinking DNA purine bases, (ii) inducing oxidative stress, and (iii)

interfering with DNA repair machinery, that can eventually lead to cancer cell death.[69] Recent

data purport that it can also inhibit GSH and GPX4.[46,48,70,71] In HDMB03 cells, we observed

cisplatin to inhibit both ABCB7 and GPX4 expression **(Figure 7A)**. In fact, we observed that

cisplatin depleted GSH in HDMB03 cells with the addition of ferrostatin having some recovery

effects. However, cisplatin added to cells with ABCB7 inhibition had the strongest effect in

depleting GSH; ferrotstatin-1 catalyzed the recovery of reduced glutathione levels in these cells

(Supplementary Figures 7D and 7E).

In HDMB03 cells with repressed ABCB7 expression, cisplatin elicited higher cytotoxicity

compared to controls **(Figure 7B)**. Moreover, in D425 **(Figure 7Ci)** and HDMB03 **(Figure**

7Cii) cells, ABCB7 repression via miR-1253[OE] resulted in a ~2-fold reduction in the IC_{50} of

cisplatin, while si-ABCB7 showed the same effect in HDMB03 cells **(Figure 7Ciii)**. Colony

formation of HDMB03 cells was nearly abrogated in ABCB7-repressed HDMB03 cells treated

with cisplatin. **(Figure 7D)**.

We next investigated whether cisplatin treatment enhanced total cellular and mitochondrial

ROS in G3MB cells. We studied ABCB7 repression by miR-1253^{OE}-transfected cells subjected

to 10 µM cisplatin in D425 and 2 µM cisplatin in HDMB03 cells at 24 h; we concurrently

examined cisplatin treatment (2 µM) of ABCB7^{KO} HDMB03 cells. As prior, ABCB7 repression

by these methods induced significantly higher levels of total and mitochondrial ROS compared

to control. Moreover, in all cases, ABCB7 repression potentiated cisplatin-mediated ROS

(**Figures 8A-C**). Concurrent treatment with ferrostatin-1 returned ROS levels to that of untreated

miR-1253^{OE} or si-ABCB7 (**Figure 8D and 8E)**. In ABCB7KO HDMB03 cells, these effects

were similarly recapitulated **(Supplementary Figures 8A and 8B).** These data provide evidence

for cisplatin-induced cytotoxicity in G3MB cells by ferroptosis.

ABCB7 repression potentiates cisplatin toxicity through ferroptosis in G3MB cells.

We next examined the impact of cisplatin on lipid peroxidation to determine if cisplatin

potentiated ferroptosis in ABCB7 repressed G3MB cells. First, cisplatin elicited a high degree of

lipid peroxidation compared to erastin **(Figure 9A)**. When combined with ABCB7 repression,

cisplatin treatment potentiated lipid peroxidation. Treatment with ferrostatin-1 significantly

reduced lipid peroxidation in cisplatin-treated ABCB7-repressed HDMB03 cells **(Figure 9B and**

9C).

Finally, examining O$_2$-mediated cell death (as a proxy for ferroptosis) at 24 h, cisplatin

showed a small rise in ROS-mediated cell death and apoptosis, while miR-1253 led to a larger

rise in both. Treating miR-1253-transfected HDMB03 cells with cisplatin elicited a dramatic

increase in both ROS-mediated and apoptotic cell death; ferrostatin-1 treatment rescued the

former, strongly implicating the important contribution of ferroptosis to the mechanism of miR-

1253-induced cell death **(Figure 9E)**. Together these data substantiate the cardinal role of

ABCB7 repression in inducing ferroptosis and its potentiating effects on cisplatin cytotoxicity

through ferroptosis.

Artesunate synergizes with cisplatin to kill G3MB cancer cells.

Artesunate is an anti-malarial drug that kills *Plasmodium falciparum* by interacting with free

iron to generate ROS.[72] Its anti-neoplastic action has been demonstrated in head and neck

cancers, where it reduced GSH levels, increased lipid ROS, and triggered ferroptosis.[73]

Moreover, tumor susceptibility to artesunate was enhanced by iron overload, especially in high

TFRC-expressing tumors.[74] In the same study, artesunate was shown to inhibit ABCB7 in human

leukemia and breast cancer cells.[74] These features make artesunate an ideal drug to test in

G3MBs, which are also high TFRC-expressing tumors.

We studied the cytotoxicity of artesunate in HDMB03 cells. Our dose-response curves

revealed a declining IC_{50} over 72h, suggestive of a cumulative effect over time **(Figure 10A)**.

Of note, these IC_{50} values are far lower than what has been determined for artesunate in glioma

cell lines.[75] Then, in orthotopic G3MB tumors developed in NSG mice, artesunate monotherapy

was shown to effectively inhibit tumor growth at 27 days post-implantation and even marginally

enhance survival **(Figure 10B)**.

Given our prior findings of ABCB7 repression potentiating cisplatin cytotoxicity, we tested

whether artesunate can bolster cisplatin cytotoxicity by potentiating iron overload and

ferroptosis. By performing a drug synergy experiment, we discovered that artesunate synergizes

with cisplatin (highest synergy score 15.2) **(Figure 10C)**. Combination therapy was superior to

monotherapy in inducing cell death in HDMB03 cells **(Figure 10D)** and abrogated colony

formation **(Figure 10E)**. Finally, we observed, similar to cisplatin, that artesunate (5 μM)

inhibited both ABCB7 and GPX4 in G3MB cells. However, combination therapy with cisplatin

(2.5 μM) compounded GPX4 inhibition **(Figure 10F)**. These data suggest that artesunate

potentiates cisplatin cytotoxicity in G3MB cells and may serve as an innovative treatment

strategy for G3MB tumors.

Artesunate synergizes with cisplatin to mitigate G3MB tumor burden.

We then tested the impact of combination therapy in our orthotopic G3MB mouse model

generated in NSG mice from HDMB03 cells. We divided our groups into negative control

(PBS), cisplatin i.p. (2.5 mg/kg), artesunate i.p. (100 mg/kg)[76], and artesunate+cisplatin. Mice

were randomized and treatment was started at day 10 post implantation **(Figure 11A)**. At

termination of the experiment, at day 22 when all wildtype mice had died, mice in the

combination arm had significantly smaller tumors and higher survival compared to other groups

(Figure 11B).

Discussion

Amongst the most devastating diagnoses in a pediatric patient is a tumor of the central

nervous system, with medulloblastoma being the most common malignant tumor.[3,4,77,78] Poor

survival (5-year OS <50%) in G3MB patients is attributable to a combination of young age at

diagnosis (peak age 3-5), metastases at diagnosis (up to 50%), and c-Myc amplification.[9,28,79-82]

Current treatment regimens have yet to impact the dismal prognosis with stagnant survival rates

seen over the last decade.[83] An inability to tolerate mainstay of therapy especially for young

patients fuels high relapse rates (~30%) which are universally fatal. For example, in children

under the age of 4 years relapse was noted in ~60% of patients who did not receive upfront

craniospinal irradiation. Moreover, relapse rates were highest in group 3 MB tumors and those

with i17q and c-Myc amplification.[84] In those that do manage to survive, irreversible damage to

the hypothalamic-pituitary axis is sustained from cytotoxic treatment regimens resulting in short

stature, cognitive impairments, and emotional lability.[85,86] Thus, the need for novel therapeutic

strategies that mitigate drug-related cytotoxicity yet accomplish widespread tumor abstraction for

this subgroup is dire.

Elevated recurrence rates have often been conjectured to be linked to mechanisms for drug

resistance in group 3 MB, which may include high expression of multi-drug resistance (MDR)

genes belonging to the ABC transporter family.[37,38,40,87,88] Cytotoxic drugs, in combination with

tumor suppressive miRNAs, have the potential for a more complex and profound effect on

tumorigenesis and may possess the ability to address drug resistance patterns, especially if they

can target MDR genes.[89] Here, we demonstrated a strong negative enrichment of the ABC

transporter family with miR-1253 expression restoration in group 3 MB cells. Of the multiple

transporters identified via our bioinformatics approach, we isolated ABCB7, whose deregulated

expression profile in group 3 tumors was strongly associated with poor prognosis, as a target of

miR-1253.

ABCB7 is an iron transporter residing on the inner mitochondrial membrane and involved in

Fe-S cluster biogenesis.[59,90] ABCB7 deficiency can lead to decreased expression of electron

transport chain (ETC) complex proteins I, II, IV and V, which can trigger impaired oxidative

phosphorylation and mitochondrial membrane integrity resulting in oxidative stress.[41,91-93] In

glioma cells, inhibition of ABCB7 resulted in disruption of iron transport and ROS generation

triggering apoptotic and non-apoptotic cell death.[42] Iron accumulation can trigger oxidative stress

and vice versa[94,95], resulting in lethal lipid peroxidation activating ferroptosis, which is distinct

from apoptosis, necrosis, autophagy, and pyroptosis.[96]

With a strong premise for exploring iron imbalance through ABCB7 repression, our

subsequent studies substantiated a role for this strategy in activating ferroptosis in G3MB. First,

miR-1253 induced intracellular iron accumulation in D425 and HDMB03 cells, effectively

abrogated by pre-treatment with an iron chelator, DFO. ABCB7^{KO} resulted in a similar

phenotype. This resulted in elevated mitochondrial ROS ($O_2^{\bullet-}$) and cytosolic ROS (H_2O_2). We

also recorded a high lipid oxidation profile in these cell lines concurrent with miR-1253

expression. Finally, using AnnexinV and MitoSOX™ Red staining, we revealed a significantly

elevated dual staining population of cells in miR-1253-expressing cells. Taken together, these

results are highly indictive of ferroptosis induction by miR-1253 in group 3 MB cells.

Ferroptosis can also be induced by disruption of glutathione synthesis or inhibition of

glutathione peroxidase 4.[97] Cancer cells have the capacity to activate redox buffering systems to

survive in a highly oxidative environment resulting from deregulated cellular functions.[67]

Glutathione is a key player in this response and a critical cofactor for glutathione peroxidase 4

(GPX4). GPX4, a central regulator of ferroptosis upregulated in various cancers, uses glutathione

to reduce ROS and lipid hydroperoxide levels thus facilitating tumor cell survival in an

environment with high oxidative stress.[98] Additionally, GSH binds Fe^{2+} to prevent iron oxidation

and is thus a critical component controlling the labile iron pool.[99] Of note, ABCB7 harbors a

GSH binding pocket, and GSH is a required substrate for cytosolic and nuclear Fe-S protein

biogenesis and iron homeostasis.[61,100] In group 3 tumors, we not only showed deregulated GPX4

expression but also a strong positive correlation with ABCB7. Resultantly, miR-1253 expression

or ABCB7^{KO} strongly inhibited GPX4 expression and reduced glutathione levels. These data

demonstrate the contribution of disrupting GPX4 and glutathione metabolism in miR-1253-

mediated oxidative stress mechanisms resulting in ferroptosis.

MiRNA mimics have been shown to possess the capability of restoring the sensitivity of

cancer cells for chemotherapeutic agents and to thus subsequently enhance their effectiveness.

For example, miR-429, miR-383, miR-101-3p, miR-195, miR-634, and miR-1294 elicited a 2-5

fold reduction in the EC_{50} and IC_{50} values in combination with gemcitabine, temozolomide, and

paclitaxel.[89,101-106] Of the standard chemotherapies for medulloblastoma tumors, only cisplatin

has been shown to trigger both apoptosis and ferroptosis, via oxidative stress, GSH depletion,

and GPX4 inactivation.[46,48,70,71] Moreover, a microarray-based study of the IC_{50} of cisplatin in 60

NCI cell lines identified multiple ABC transporters, including 3 iron transporters, i.e. ABCB6,

ABCB7 and ABCB10, in conferring cisplatin drug resistance.[107]

Given the ferroptotic mechanism we elucidated for miR-1253, we studied whether miR-1253

expression can potentiate cisplatin cytotoxicity in group 3 MB cells. We report a 2-fold reduction

in the IC_{50} value of cisplatin with miR-1253 induction in D425 and HDMB03 cells. Combination

treatment had dramatic effects on the clonal potential of HDMB03 cells. We then demonstrated

the highest induction of mitochondrial and cytosolic ROS, lipid peroxidation, and GSH depletion

in miR-1253-expressing HDMB03 and ABCB7^{KO} HDMB03 cells treated with cisplatin.

Consequently, combination therapy resulted in the highest degree of ferroptosis in these cell

lines. These effects were reversed by ferrostatin-1, a potent inhibitor of ferroptosis, lending

further validity to the central role of ferroptosis in the mechanism of miR-1253-mediated effects

and cisplatin potentiation.

Overall, our study has identified novel tumor suppressive properties for miR-1253. First,

miR-1253 directly inhibits ABCB7 expression, thus inducing labile iron pool within MB cancer

cells and stimulating ROS production. MiR-1253 can concurrently downregulate the expression

of GPX4 and deplete GSH, further exacerbating ROS. Together, the generation of lipid

hyperoxides progresses unabated, leading to cancer cell death via ferroptosis. We also leveraged

these properties by showing potentiation of cisplatin cytotoxicity and thus enhanced therapeutic

efficacy in group 3 MB cells. Together, our findings provide proof-of-concept for further

exploration of tumor suppressive microRNAs as therapeutic adjuncts to standard chemotherapy.

Such a strategy may mitigate the current limitations to treatment regimens in our youngest high

risk patients.

References

1. Taylor RE, Donachie PH, Weston CL, et al. Impact of radiotherapy parameters on outcome for patients with supratentorial primitive neuro-ectodermal tumours entered into the SIOP/UKCCSG PNET 3 study. *Radiother Oncol.* 2009; 92(1):83-88.
2. Udaka YT, Packer RJ. Pediatric Brain Tumors. *Neurol Clin.* 2018; 36(3):533-556.
3. Ostrom QT, Patil N, Cioffi G, Waite K, Kruchko C, Barnholtz-Sloan JS. CBTRUS Statistical Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2013-2017. *Neuro-oncology.* 2020; 22(12 Suppl 2):iv1-iv96.
4. Taylor MD, Northcott PA, Korshunov A, et al. Molecular subgroups of medulloblastoma: the current consensus. *Acta neuropathologica.* 2012; 123(4):465-472.
5. Zapotocky M, Mata-Mbemba D, Sumerauer D, et al. Differential patterns of metastatic dissemination across medulloblastoma subgroups. *J Neurosurg Pediatr.* 2018; 21(2):145-152.
6. Northcott PA, Korshunov A, Witt H, et al. Medulloblastoma comprises four distinct molecular variants. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology.* 2011; 29(11):1408-1414.
7. Ellison DW, Kocak M, Dalton J, et al. Definition of disease-risk stratification groups in childhood medulloblastoma using combined clinical, pathologic, and molecular variables. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology.* 2011; 29(11):1400-1407.
8. Kool M, Koster J, Bunt J, et al. Integrated genomics identifies five medulloblastoma subtypes with distinct genetic profiles, pathway signatures and clinicopathological features. *PloS one.* 2008; 3(8):e3088.
9. Cavalli FMG, Remke M, Rampasek L, et al. Intertumoral Heterogeneity within Medulloblastoma Subgroups. *Cancer cell.* 2017; 31(6):737-754 e736.
10. Ramaswamy V, Remke M, Bouffet E, et al. Recurrence patterns across medulloblastoma subgroups: an integrated clinical and molecular analysis. *Lancet Oncol.* 2013; 14(12):1200-1207.
11. Rieken S, Mohr A, Habermehl D, et al. Outcome and prognostic factors of radiation therapy for medulloblastoma. *Int J Radiat Oncol Biol Phys.* 2011; 81(3):e7-e13.
12. Packer RJ, Goldwein J, Nicholson HS, et al. Treatment of children with medulloblastomas with reduced-dose craniospinal radiation therapy and adjuvant chemotherapy: A Children's Cancer Group Study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology.* 1999; 17(7):2127-2136.
13. Gajjar A, Chintagumpala M, Ashley D, et al. Risk-adapted craniospinal radiotherapy followed by high-dose chemotherapy and stem-cell rescue in children with newly diagnosed medulloblastoma (St Jude Medulloblastoma-96): long-term results from a prospective, multicentre trial. *The Lancet. Oncology.* 2006; 7(10):813-820.
14. Martin AM, Raabe E, Eberhart C, Cohen KJ. Management of pediatric and adult patients with medulloblastoma. *Curr Treat Options Oncol.* 2014; 15(4):581-594.
15. Enriquez Perez J, Fritzell S, Kopecky J, Visse E, Darabi A, Siesjo P. The effect of locally delivered cisplatin is dependent on an intact immune function in an experimental glioma model. *Sci Rep.* 2019; 9(1):5632.
16. Nageswara Rao AA, Wallace DJ, Billups C, Boyett JM, Gajjar A, Packer RJ. Cumulative cisplatin dose is not associated with event-free or overall survival in children with newly diagnosed average-risk medulloblastoma treated with cisplatin based adjuvant chemotherapy: report from the Children's Oncology Group. *Pediatr Blood Cancer.* 2014; 61(1):102-106.
17. Mognato M, Celotti L. MicroRNAs Used in Combination with Anti-Cancer Treatments Can Enhance Therapy Efficacy. *Mini Rev Med Chem.* 2015; 15(13):1052-1062.

18. He X, Xiao X, Dong L, et al. MiR-218 regulates cisplatin chemosensitivity in breast cancer by targeting BRCA1. *Tumour Biol.* 2015; 36(3):2065-2075.

19. Sun C, Li N, Yang Z, et al. miR-9 regulation of BRCA1 and ovarian cancer sensitivity to cisplatin and PARP inhibition. *J Natl Cancer Inst.* 2013; 105(22):1750-1758.

20. Pouliot LM, Chen YC, Bai J, et al. Cisplatin sensitivity mediated by WEE1 and CHK1 is mediated by miR-155 and the miR-15 family. *Cancer Res.* 2012; 72(22):5945-5955.

21. Zhu W, Xu H, Zhu D, et al. miR-200bc/429 cluster modulates multidrug resistance of human cancer cell lines by targeting BCL2 and XIAP. *Cancer Chemother Pharmacol.* 2012; 69(3):723-731.

22. Mittal A, Chitkara D, Behrman SW, Mahato RI. Efficacy of gemcitabine conjugated and miRNA-205 complexed micelles for treatment of advanced pancreatic cancer. *Biomaterials.* 2014; 35(25):7077-7087.

23. Hong DS, Kang YK, Borad M, et al. Phase 1 study of MRX34, a liposomal miR-34a mimic, in patients with advanced solid tumours. *British journal of cancer.* 2020; 122(11):1630-1637.

24. Hoff C, Seranski P, Mollenhauer J, et al. Physical and transcriptional mapping of the 17p13.3 region that is frequently deleted in human cancer. *Genomics.* 2000; 70(1):26-33.

25. McDonald JD, Daneshvar L, Willert JR, Matsumura K, Waldman F, Cogen PH. Physical mapping of chromosome 17p13.3 in the region of a putative tumor suppressor gene important in medulloblastoma. *Genomics.* 1994; 23(1):229-232.

26. Steichen-Gersdorf E, Baumgartner M, Kreczy A, Maier H, Fink FM. Deletion mapping on chromosome 17p in medulloblastoma. *British journal of cancer.* 1997; 76(10):1284-1287.

27. Kanchan RK, Perumal N, Atri P, et al. MiR-1253 exerts tumor-suppressive effects in medulloblastoma via inhibition of CDK6 and CD276 (B7-H3). *Brain Pathol.* 2020; 30(4):732-745.

28. Perumal N, Kanchan RK, Doss D, et al. MiR-212-3p functions as a tumor suppressor gene in group 3 medulloblastoma via targeting nuclear factor I/B (NFIB). *Acta neuropathologica communications.* 2021; 9(1).

29. Mao Y, Li W, Chen K, et al. B7-H1 and B7-H3 are independent predictors of poor prognosis in patients with non-small cell lung cancer. *Oncotarget.* 2015; 6(5):3452-3461.

30. Kang FB, Wang L, Jia HC, et al. B7-H3 promotes aggression and invasion of hepatocellular carcinoma by targeting epithelial-to-mesenchymal transition via JAK2/STAT3/Slug signaling pathway. *Cancer Cell Int.* 2015; 15:45.

31. Kanchan RK, Doss D, Khan P, Nasser MW, Mahapatra S. To kill a cancer: Targeting the immune inhibitory checkpoint molecule, B7-H3. *Biochim Biophys Acta Rev Cancer.* 2022:188783.

32. Liu M, Zhang Y, Zhang J, et al. MicroRNA-1253 suppresses cell proliferation and invasion of non-small-cell lung carcinoma by targeting WNT5A. *Cell death & disease.* 2018; 9(2):189.

33. Mo J, Zheng T, Lei L, et al. MicroRNA-1253 Suppresses Cell Proliferation Migration and Invasion of Osteosarcoma by Targeting MMP9. *Technol Cancer Res Treat.* 2021; 20:1533033821995278.

34. Xu Y, Yao Y, Gao P, Cui Y. Upregulated circular RNA circ_0030235 predicts unfavorable prognosis in pancreatic ductal adenocarcinoma and facilitates cell progression by sponging miR-1253 and miR-1294. *Biochemical and biophysical research communications.* 2019; 509(1):138-142.

35. Merino MJ, Gil S, Macias CG, Lara K. The Unknown microRNA Expression of Male Breast Cancer. Similarities and Differences with Female Ductal Carcinoma. Their Role as Tumor Biomarker. *Journal of Cancer.* 2018; 9(3):450-459.

36. Yang D, Zhang D. miR-1253, a novel tumor suppressor gene in colon cancer, is associated with poor patients prognosis. *Clin Exp Med.* 2021.

37. Szakacs G, Paterson JK, Ludwig JA, Booth-Genthe C, Gottesman MM. Targeting multidrug resistance in cancer. *Nature reviews. Drug discovery.* 2006; 5(3):219-234.

38. Choi YH, Yu AM. ABC transporters in multidrug resistance and pharmacokinetics, and strategies for drug development. *Current pharmaceutical design.* 2014; 20(5):793-807.

39.   Bao L, Hazari S, Mehra S, Kaushal D, Moroz K, Dash S. Increased expression of P-glycoprotein and
doxorubicin chemoresistance of metastatic breast cancer is regulated by miR-298. *The American
journal of pathology.* 2012; 180(6):2490-2503.
40.   Morfouace M, Cheepala S, Jackson S, et al. ABCG2 Transporter Expression Impacts Group 3
Medulloblastoma Response to Chemotherapy. *Cancer research.* 2015; 75(18):3879-3889.
41.   Maio N, Kim KS, Holmes-Hampton G, Singh A, Rouault TA. Dimeric ferrochelatase bridges ABCB7
and ABCB10 homodimers in an architecturally defined molecular complex required for heme
biosynthesis. *Haematologica.* 2019; 104(9):1756-1767.
42.   Kim JY, Kim JK, Kim H. ABCB7 simultaneously regulates apoptotic and non-apoptotic cell death
by modulating mitochondrial ROS and HIF1alpha-driven NFkappaB signaling. *Oncogene.* 2020;
39(9):1969-1982.
43.   Stockwell BR, Friedmann Angeli JP, Bayir H, et al. Ferroptosis: A Regulated Cell Death Nexus
Linking Metabolism, Redox Biology, and Disease. *Cell.* 2017; 171(2):273-285.
44.   Friedmann Angeli JP, Schneider M, Proneth B, et al. Inactivation of the ferroptosis regulator
Gpx4 triggers acute renal failure in mice. *Nat Cell Biol.* 2014; 16(12):1180-1191.
45.   Hirschhorn T, Stockwell BR. The development of the concept of ferroptosis. *Free Radic Biol Med.*
2019; 133:130-143.
46.   Guo J, Xu B, Han Q, et al. Ferroptosis: A Novel Anti-tumor Action for Cisplatin. *Cancer Res Treat.*
2018; 50(2):445-460.
47.   Jiang L, Kon N, Li T, et al. Ferroptosis as a p53-mediated activity during tumour suppression.
Nature. 2015; 520(7545):57-62.
48.   Zhang X, Sui S, Wang L, et al. Inhibition of tumor propellant glutathione peroxidase 4 induces
ferroptosis in cancer cells and enhances anticancer effect of cisplatin. *J Cell Physiol.* 2020;
235(4):3425-3437.
49.   Drusco A. A set of differentially expressed cerebro-spinal fluid (CSF) miRNAs can diagnose CNS
malignancies. 2015.
50.   Kanchan RK, Perumal N, Atri P, et al. MiR-1253 exerts tumor-suppressive effects in
medulloblastoma via inhibition of CDK6 and CD276 (B7-H3). *Brain pathology (Zurich,
Switzerland).* 2020.
51.   Weishaupt H, Johansson P, Sundstrom A, et al. Batch-normalization of cerebellar and
medulloblastoma gene expression datasets utilizing empirically defined negative control genes.
Bioinformatics. 2019; 35(18):3357-3364.
52.   Kaushal JB, Bhatia R, Kanchan RK, et al. Repurposing Niclosamide for Targeting Pancreatic
Cancer by Inhibiting Hh/Gli Non-Canonical Axis of Gsk3beta. *Cancers (Basel).* 2021; 13(13).
53.   Anderson GJ, Vulpe CD. Mammalian iron transport. *Cell Mol Life Sci.* 2009; 66(20):3241-3261.
54.   Torti SV, Torti FM. Iron and cancer: more ore to be mined. *Nature reviews. Cancer.* 2013;
13(5):342-355.
55.   Battaglia AM, Chirillo R, Aversa I, Sacco A, Costanzo F, Biamonte F. Ferroptosis and Cancer:
Mitochondria Meet the "Iron Maiden" Cell Death. *Cells.* 2020; 9(6).
56.   Ray S, Chaturvedi NK, Bhakat KK, Rizzino A, Mahapatra S. Subgroup-Specific Diagnostic,
Prognostic, and Predictive Markers Influencing Pediatric Medulloblastoma Treatment.
Diagnostics. 2021; 12(1).
57.   Forcina GC, Dixon SJ. GPX4 at the Crossroads of Lipid Homeostasis and Ferroptosis. *Proteomics.*
2019; 19(18):e1800311.
58.   Rouault TA, Tong WH. Iron-sulphur cluster biogenesis and mitochondrial iron homeostasis. *Nat
Rev Mol Cell Biol.* 2005; 6(4):345-351.

59. Pondarre C, Antiochos BB, Campagna DR, et al. The mitochondrial ATP-binding cassette
transporter Abcb7 is essential in mice and participates in cytosolic iron-sulfur cluster biogenesis.
Hum Mol Genet. 2006; 15(6):953-964.
60. Seguin A, Ward DM. Mitochondrial ABC Transporters and Iron Metabolism. *Journal of Clinical &*
Experimental Pathology. 2018; 08(01).
61. Srinivasan V, Pierik AJ, Lill R. Crystal structures of nucleotide-free and glutathione-bound
mitochondrial ABC transporter Atm1. *Science (New York, N.Y.).* 2014; 343(6175):1137-1140.
62. Li SW, Liu CM, Guo J, et al. Iron overload induced by ferric ammonium citrate triggers reactive
oxygen species-mediated apoptosis via both extrinsic and intrinsic pathways in human hepatic
cells. *Hum Exp Toxicol.* 2016; 35(6):598-607.
63. Zhao Y, Li Y, Zhang R, Wang F, Wang T, Jiao Y. The Role of Erastin in Ferroptosis and Its Prospects
in Cancer Therapy. *Onco Targets Ther.* 2020; 13:5429-5441.
64. Creed S, McKenzie M. Measurement of Mitochondrial Membrane Potential with the Fluorescent
Dye Tetramethylrhodamine Methyl Ester (TMRM). *Methods in molecular biology (Clifton, N.J.).*
2019; 1928:69-76.
65. Favaro G, Romanello V, Varanita T, et al. DRP1-mediated mitochondrial shape controls calcium
homeostasis and muscle mass. *Nature communications.* 2019; 10(1):2576.
66. Zilka O, Shah R, Li B, et al. On the Mechanism of Cytoprotection by Ferrostatin-1 and
Liproxstatin-1 and the Role of Lipid Peroxidation in Ferroptotic Cell Death. *ACS Cent Sci.* 2017;
3(3):232-243.
67. Aboelella NS, Brandle C, Kim T, Ding ZC, Zhou G. Oxidative Stress in the Tumor
Microenvironment and Its Relevance to Cancer Immunotherapy. *Cancers (Basel).* 2021; 13(5).
68. Li FJ, Long HZ, Zhou ZW, Luo HY, Xu SG, Gao LC. System X(c) (-)/GSH/GPX4 axis: An important
antioxidant system for the ferroptosis in drug-resistant solid tumor therapy. *Front Pharmacol.*
2022; 13:910292.
69. Dasari S, Tchounwou PB. Cisplatin in cancer therapy: molecular mechanisms of action. *Eur J*
Pharmacol. 2014; 740:364-378.
70. Zhou Y, Shen Y, Chen C, et al. The crosstalk between autophagy and ferroptosis: what can we
learn to target drug resistance in cancer? *Cancer Biol Med.* 2019; 16(4):630-646.
71. Berndtsson M, Hagg M, Panaretakis T, Havelka AM, Shoshan MC, Linder S. Acute apoptosis by
cisplatin requires induction of reactive oxygen species but is not associated with damage to
nuclear DNA. *International journal of cancer. Journal international du cancer.* 2007; 120(1):175-
180.
72. Meshnick SR. Artemisinin: mechanisms of action, resistance and toxicity. *Int J Parasitol.* 2002;
32(13):1655-1660.
73. Roh JL, Kim EH, Jang H, Shin D. Nrf2 inhibition reverses the resistance of cisplatin-resistant head
and neck cancer cells to artesunate-induced ferroptosis. *Redox Biol.* 2017; 11:254-262.
74. Kelter G, Steinbach D, Konkimalla VB, et al. Role of transferrin receptor and the ABC transporters
ABCB6 and ABCB7 for resistance and differentiation of tumor cells towards artesunate. *PloS one.*
2007; 2(8):e798.
75. Hooft van Huijsduijnen R, Guy RK, Chibale K, et al. Anticancer properties of distinct antimalarial
drug classes. *PloS one.* 2013; 8(12):e82962.
76. Berte N, Lokan S, Eich M, Kim E, Kaina B. Artesunate enhances the therapeutic response of
glioma cells to temozolomide by inhibition of homologous recombination and senescence.
Oncotarget. 2016; 7(41):67235-67250.
77. Gilbertson R, Wickramasinghe C, Hernan R, et al. Clinical and molecular stratification of disease
risk in medulloblastoma. *British journal of cancer.* 2001; 85(5):705-712.

78.   Gottardo NG, Hansford JR, McGlade JP, et al. Medulloblastoma Down Under 2013: a report from
764   the third annual meeting of the International Medulloblastoma Working Group. *Acta*
765   *neuropathologica.* 2014; 127(2):189-201.

766   **79.**   Cho YJ, Tsherniak A, Tamayo P, et al. Integrative genomic analysis of medulloblastoma identifies
767   a molecular subgroup that drives poor clinical outcome. *Journal of clinical oncology : official*
768   *journal of the American Society of Clinical Oncology.* 2011; 29(11):1424-1430.

769   **80.**   Tamayo P, Cho YJ, Tsherniak A, et al. Predicting relapse in patients with medulloblastoma by
770   integrating evidence from clinical and genomic features. *Journal of clinical oncology : official*
771   *journal of the American Society of Clinical Oncology.* 2011; 29(11):1415-1423.

772   **81.**   Thompson EM, Hielscher T, Bouffet E, et al. Prognostic value of medulloblastoma extent of
773   resection after accounting for molecular subgroup: a retrospective integrated clinical and
774   molecular analysis. *The Lancet Oncology.* 2016; 17(4):484-495.

775   **82.**   Menyhart O, Giangaspero F, Gyorffy B. Molecular markers and potential therapeutic targets in
776   non-WNT/non-SHH (group 3 and group 4) medulloblastomas. *J Hematol Oncol.* 2019; 12(1):29.

777   **83.**   Ward E, DeSantis C, Robbins A, Kohler B, Jemal A. Childhood and adolescent cancer statistics,
778   2014. *CA Cancer J Clin.* 2014; 64(2):83-103.

779   **84.**   Hill RM, Richardson S, Schwalbe EC, et al. Time, pattern, and outcome of medulloblastoma
780   relapse and their association with tumour biology at diagnosis and therapy: a multicentre cohort
781   study. *The Lancet Child & Adolescent Health.* 2020; 4(12):865-874.

782   **85.**   Hoff KV, Hinkes B, Gerber NU, et al. Long-term outcome and clinical prognostic factors in
783   children with medulloblastoma treated in the prospective randomised multicentre trial HIT'91.
784   *Eur J Cancer.* 2009; 45(7):1209-1217.

785   **86.**   Ribi K, Relly C, Landolt MA, Alber FD, Boltshauser E, Grotzer MA. Outcome of medulloblastoma
786   in children: long-term complications and quality of life. *Neuropediatrics.* 2005; 36(6):357-365.

787   **87.**   Othman RT, Kimishi I, Bradshaw TD, et al. Overcoming multiple drug resistance mechanisms in
788   medulloblastoma. *Acta Neuropathol Commun.* 2014; 2:57.

789   **88.**   Chou PM, Reyes-Mugica M, Barquin N, Yasuda T, Tan X, Tomita T. Multidrug resistance gene
790   expression in childhood medulloblastoma: correlation with clinical outcome and DNA ploidy in
791   29 patients. *Pediatric neurosurgery.* 1995; 23(6):283-291; discussion 291-282.

792   **89.**   Miroshnichenko S, Patutina O. Enhanced Inhibition of Tumorigenesis Using Combinations of
793   miRNA-Targeted Therapeutics. *Front Pharmacol.* 2019; 10:488.

794   **90.**   Cavadini P, Biasiotto G, Poli M, et al. RNA silencing of the mitochondrial ABCB7 transporter in
795   HeLa cells causes an iron-deficient phenotype with mitochondrial iron overload. *Blood.* 2007;
796   109(8):3552-3559.

797   **91.**   Miyake A, Higashijima S, Kobayashi D, et al. Mutation in the abcb7 gene causes abnormal iron
798   and fatty acid metabolism in developing medaka fish. *Dev Growth Differ.* 2008; 50(9):703-716.

799   **92.**   Gonzalez-Cabo P, Bolinches-Amoros A, Cabello J, et al. Disruption of the ATP-binding cassette B7
800   (ABTM-1/ABCB7) induces oxidative stress and premature cell death in Caenorhabditis elegans.
801   *The Journal of biological chemistry.* 2011; 286(24):21304-21314.

802   **93.**   Kumar V, A AK, Sanawar R, Jaleel A, Santhosh Kumar TR, Kartha CC. Chronic Pressure Overload
803   Results in Deficiency of Mitochondrial Membrane Transporter ABCB7 Which Contributes to Iron
804   Overload, Mitochondrial Dysfunction, Metabolic Shift and Worsens Cardiac Function. *Sci Rep.*
805   2019; 9(1):13170.

806   **94.**   Nunez MT, Nunez-Millacura C, Tapia V, et al. Iron-activated iron uptake: a positive feedback loop
807   mediated by iron regulatory protein 1. *Biometals.* 2003; 16(1):83-90.

808   **95.**   Dixon SJ, Stockwell BR. The role of iron and reactive oxygen species in cell death. *Nat Chem Biol.*
809   2014; 10(1):9-17.

96. Mou Y, Wang J, Wu J, et al. Ferroptosis, a new form of cell death: opportunities and challenges
in cancer. *J Hematol Oncol.* 2019; 12(1):34.
97. Abdalkader M, Lampinen R, Kanninen KM, Malm TM, Liddell JR. Targeting Nrf2 to Suppress
Ferroptosis and Mitochondrial Dysfunction in Neurodegeneration. *Front Neurosci.* 2018; 12:466.
98. Yang WS, SriRamaratnam R, Welsch ME, et al. Regulation of ferroptotic cancer cell death by
GPX4. *Cell.* 2014; 156(1-2):317-331.
99. Hider RC, Kong XL. Glutathione: a key component of the cytoplasmic labile iron pool. *Biometals.*
2011; 24(6):1179-1187.
100. Gaschler MM, Andia AA, Liu H, et al. FINO2 initiates ferroptosis through GPX4 inactivation and
iron oxidation. *Nat Chem Biol.* 2018; 14(5):507-515.
101. Fan P, Liu L, Yin Y, et al. MicroRNA-101-3p reverses gemcitabine resistance by inhibition of
ribonucleotide reductase M1 in pancreatic cancer. *Cancer Lett.* 2016; 373(1):130-137.
102. Yu G, Jia B, Cheng Y, et al. MicroRNA-429 sensitizes pancreatic cancer cells to gemcitabine
through regulation of PDCD4. *Am J Transl Res.* 2017; 9(11):5048-5055.
103. Chen H, Liu L, Li X, Shi Y, Liu N. MicroRNA-1294 inhibits the proliferation and enhances the
chemosensitivity of glioma to temozolomide via the direct targeting of TPX2. *Am J Cancer Res.*
2018; 8(2):291-301.
104. Jiang J, Xie C, Liu Y, Shi Q, Chen Y. Up-regulation of miR-383-5p suppresses proliferation and
enhances chemosensitivity in ovarian cancer cells by targeting TRIM27. *Biomed Pharmacother.*
2019; 109:595-601.
105. Tu MJ, Ho PY, Zhang QY, et al. Bioengineered miRNA-1291 prodrug therapy in pancreatic cancer
cells and patient-derived xenograft mouse models. *Cancer Lett.* 2019; 442:82-90.
106. Wang M, Qiu R, Yu S, et al. Paclitaxelresistant gastric cancer MGC803 cells promote
epithelialtomesenchymal transition and chemoresistance in paclitaxelsensitive cells via
exosomal delivery of miR1555p. *Int J Oncol.* 2019; 54(1):326-338.
107. Konkimalla VB, Kaina B, Efferth T. Role of transporter genes in cisplatin resistance. *In Vivo.* 2008;
22(3):279-283.

Figure Legends

Figure 1. ABCB7 expression upregulation in G3MB portends a poor prognosis. (A)

Subgroup-specific ABCB7 expression assessment by RNA sequencing ($\log_2$ transcripts per

million) of a local medulloblastoma patient cohort (Kanchan *et al.*, GSE) showing specific

upregulation in G3MB. *CB*, pediatric cerebellum (n=10); *SHH*, sonic hedgehog (n=6); *G3*, group

3 (n=7); *G4*, group 4 (n=12). **(B)** Poor prognostic profile demonstrable in high-expressing G3MB

patients (Cavali *et al*. GSE85217). **(C)** Confocal microscopic images confirming high ABCB7

expression in G3MB tumors (n=6) compared to pediatric cerebellum (n=6) and colocalization to

the mitochondria based on COXIV fluorescence. Images captured at 10X magnification. **(D)**

Western blotting analysis showing high ABCB7 expression in classic MB cell lines (G3: D341,

D425, HDMB03, D556; G3/4: D283) compared to normal human astrocytes (NHA).

Downregulation of ABCB7 with miR-1253 overexpression in **(E)** D425 and **(F)** HDMB03 cells.

(G) Confocal microscopic images in D425 and HDMB03 cells showing co-localization of

ABCB7 to the mitochondria and downregulation with miR-1253 overexpression. Images

captured at 63X magnification. **(H)** Dual-luciferase assay confirming direct binding of miR-1253

to ABCB7 in HDMB03 cells. Data presented as mean $\pm$ SD, n=3; *ns*, not significant, ***p

<0.001. *NC*, negative control; *Scr miR*, scramble micro RNA.

Figure 2. ABCB7 repression attenuates G3MB cancer cell aggressiveness. Attenuation of

medullosphere formation **(A)** and reduction in viability (by MTT) of HDMB03 cells treated with

miR-1253^{OE} (100 nM) and si-ABCB7 (50 nM). **(C)** Dox-inducible ABCB7^{KD} (1 ug/mL)

demonstrating similarly reduced viability of HDMB03 cells. Data presented as mean $\pm$ SD, n=3;

**p <0.01, ****p <0.0001. *Dox*, doxycycline; *NC*, negative control.

Figure 3. ABCB7 repression triggers iron imbalance, oxidative stress, and lipid

peroxidation in G3MB cells. Iron overload demonstrable in the mitochondria **(A)** and

cytoplasm **(B)** of HDMB03 transfected with si-ABCB7 (100 nM) as measured by

MitoFerroGreen and FerroOrange, respectively. Deferoxamine (DFO) rescues this phenotype.

(C) Cytoplasmic iron overload demonstrable with dox-inducible $ABCB7^{KD}$ (1 ug/mL) in

HDMB03 cells. Elevated mitochondrial **(D)** and total cellular **(E)** oxidative stress in HDMB03

cells transfected with si-ABCB7 (100 nM) by mitochondrial $O_2^{\bullet-}$ (MitoSOX™ Red) and cellular

H_2O_2 (DCFDA), respectively. Ferrostatin-1 (100 nM) rescues this phenotype. **(F)** Elevated

mitochondrial oxidative stress demonstrable with dox-inducible $ABCB7^{KD}$ (1 ug/mL) in

HDMB03 cells. **(G)** Elevated lipid peroxidation in ABCB7-repressed HDMB03 cells as

measured by BODIPY-C11 staining. **(H)** Lipid peroxidation quantified by FACS. Mean ±SD,

n=3; *ns*, not significant, $**p<0.01$, $***p<0.001$ $****p<0.0001$. Fluorescent images for ROS

obtained at 20X; quantification done by averaging 4 fields. Confocal images for lipid ROS

obtained at 40X. Quantification of lipid ROS conducted via FACS, examining ratio of oxidized-

to-reduced BODIPY-C11. *DFO*, deferoxamine; *ERA*, erastin; *FAC*, ferric ammonium citrate;

FER, ferrostatin-1; H_2O_2, hydrogen peroxide; O_2^-, superoxide; s*i*, si-ABCB7.

Figure 4. ABCB7 repression induces mitochondrial dysfunction. (A) Disruption of

mitochondrial membrane potential as measured by TMRM fluorescence in ABCB7-repressed

(si-ABCB7, 100 nM) HDMB03 cells, rescued by ferrostatin-1 (100nM). **(B)** Seahorse assay

further showing disruption of oxidative phosphorylation in ABCB7-repressed (miR-1253 100

nM and si-ABCB7 100 nM) HDMB03 cells. Confocal images obtained at 63X magnification.

Quantification done by taking average of 4 fields. Mean ±SD, n=3; *ns*, not significant,

***$p<0.001$, ****$p<0.0001$ compared to NC. *ERA*, erastin; *FER*, ferrostatin-1; *NC*, negative

control; *si-*, si-ABCB7.

Figure 5. ABCB7 repression reduces G3MB tumor burden. NSG mice were orthotopically

injected with luciferase-tagged wild-type vs. dox-inducible ABCB7^{KD} HDMB03 cells. Dox

induction was started at day 10 post-implantation. **(A and B)** Reduction in tumor burden noted in

dox-induced ABCB7^{KD} group at day 28 and day 30 post-implantation compared to control. **(C)**

Attenuated expression of ABCB7 and GPX4 and enhanced expression of 4HNE, a stable end-

product of ferroptosis, in dox-induced ABCB7^{KD} group by immunohistochemical staining. Mean

±SD, n=3-5 mice per arm; **$p<0.01$ compared to -Dox mice. *4-HNE*, 4-hydroxynonenol; *Dox*,

doxycycline; *GPX4*, glutathione peroxidase 4.

Figure 6. ABCB7 repression inhibits GPX4 expression, the central regulator of ferroptosis.

Reduced GPX4 expression by miR-1253^{OE} in HDMB03 **(A)** and D425 **(B)**, by ABCB7^{KO} **(C)**,

and ABCB7^{KD} **(D)**. **(E)** Spearman correlation showing positive correlation between ABCB7 and

GPX4 expression. (Kanchan *et al.*, GSE148390). Fractionation studies revealing that ABCB7

repression by miR-1253^{OE} **(F)**, ABCB7^{KO} **(G)**, and si-ABCB7 **(H)** abrogates GPX4 expression

from the cytoplasm and mitochondria. Mean ±SD, n=3; ****$p<0.0001$. *KO*, ABCB7^{KO}; *miR*,

miR-1253 overexpression; *NC*, normal control; *si*, si-ABCB7; *WT*, wildtype HDMB03 cells.

Figure 7. ABCB7 repression potentiates cisplatin toxicity in G3MB cells. (A) Cisplatin (2.5

μM) induced repression of ABCB7 and GPX4 expression in HDMB03 cells. **(B)** Potentiation of

cisplatin toxicity in ABCB7-repressed HDMB03 cells by MTT. **(C)** A 2-fold reduction in IC$_{50}$ of

cisplatin in D425 **(i)** and HDMB03 **(ii)** cells transfected with scramble vs. miR-1253 for 24-72h

and in HDMB03 cells transfected with si-ABCB7 (50 uM) for 24h **(iii)**. Tabulated results

presented as fold-change in cisplatin IC$_{50}$ between scramble and miR-1253 in D425 and

HDMB03 cells at different time points. **(D and E)** Colonogenic assay demonstrating inhibitory

effect of cisplatin vs. ABCB7 repression vs. combination on colony formation in HDMB03 cells.

Mean ±SD, n=3; *$p<0.05$, ****$p<0.0001$ compared to NC; ##$p<0.01$ compared to miR;

####$p<0.001$ compared to si. *CIS*, cisplatin; *FER*, ferrostatin-1; *miR*, miR-1253; *NC*, negative

control; *si*, si-ABCB7.

Figure 8. ABCB7 repression potentiates oxidative stress in cisplatin-treated G3MB cells.

MiR-1253-transfected D425 **(A)** and HDMB03 **(B)**, and ABCB7^{KO} HDMB03 cells **(C)** treated

with cisplatin (D425 24-h IC$_{25}$ ~10 µM; HDMB03 24-h IC$_{25}$ ~2 µM) demonstrating potentiated

mitochondrial ROS (MitoSOX™ Red, $O_2^{\bullet-}$) and total ROS (DCFDA, H_2O_2). **(D** and **E)** Cisplatin

potentiating ROS in HDMB03 cells with repressed ABCB7 expression; ferrostatin-1 rescues this

effect. Representative images and graphs showing the potentiating effect of combining miR-1253

with cisplatin. Images captured at 20X magnification. Scale bar 400 µm. Mean ± SD, n=3; *ns*,

not significant, *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$. *CIS*, cisplatin; *FER*, ferrostatin-

1; *miR*, miR-1253; *NC*, negative control; *si*, si-ABCB7.

Figure 9. ABCB7 repression potentiates cisplatin cytotoxicity in G3MB cancer cells by

triggering ferroptosis. Lipid peroxidation was visualized by BODIPY-C11 staining and

quantified by FACs using the ratio of oxidized-to-reduced lipids in HDMB03 cells. **(A)** Cisplatin

(5 uM) induced lipid peroxidation compared to erastin, a known inducer of ferroptosis, with

partial rescue by ferrostatin-1. **(B)** Potentiation of lipid peroxidation in ABCB7-repressed cells

treated with cisplatin (5 uM), with partial rescue by ferrostatin-1. **(C)** Visualization of lipid

peroxidation as measured by BODIPY-C11 staining by confocal microscopy. Images obtained at

63X. **(D and E)** Analysis of cell death by flow cytometry showing significantly higher $O_2^{\bullet-}$

mediated cell death (representing ferroptosis) in ABCB7-repressed HDMB03 cells. This is

demonstrable by quantifying cells staining for both Annexin V-Cy5 (apoptosis) and for $O_2^{\bullet-}$

(Mitosox) (Q2). ABCB7 repression by si- **(D)** and miR-1253 **(E)** potentiated cisplatin-induced

ferroptosis, rescue by ferrostatin-1. Mean ± SD, n=3; *p <0.05, **p <0.01, ***p <0.001, ****p

<0.0001. *CIS*, cisplatin; *ERA*, erastin; *FER*, ferrostatin-1; *miR*, miR-1253; *NC*, negative control;

si, si-ABCB7.

Figure 10. Artesunate synergizes with cisplatin to kill G3MB cancer cells. (A) Dose-response

curve for artesunate in HDMB03 cells. **(B)** Reduction in G3MB tumor burden and prolonged

survival with artesunate (100 mg/kg i.p. daily) in an orthotopic G3MB tumor model. n=3-5 mice

per group. **(C)** Drug synergy experiment between cisplatin and artesunate showing synergy

between both drugs in HDMB03 cells. Potentiated cell death **(D)** and abrogated colony formation

(E) in with combined artesunate + cisplatin. **(F)** Potent inhibition of GPX4 by artesunate and

cisplatin treatments. Mean ±SD, nb=3; ****p<0.0001 compared to VC, $\int\int$$p$<0.0001 compared

to CIS, $^{\$\$\$\$}p$<0.0001 compared to ART. *ART*, artesunate; *CIS*, cisplatin; *FER*, ferrostatin-1;

GPX4, glutathione peroxidase 4; *VC*, vehicle control.

Figure 11. Artesunate synergizes with cisplatin to reduce G3MB tumor burden. (A) NSG

mice were orthotopically injected with luciferase-tagged wild-type HDMB03 cells. Mice were

randomized into four treatment arms (CON: negative control; CIS: cisplatin only at 2.5 mg/kg

i.p.; ART: artesunate only at 100 mg/kg i.p.; ART+CIS: combination). Treatment was started at

day 10 post-implantation. Experiment was terminated at day 22 when all wildtype mice died. **(B)**

Mice in the combination arm had significantly smaller tumors and higher survival. Mean ± SD,

n=3-5 mice per arm; *p<0.05 compared to PBS-treated negative controls. *ART*, artesunate;

ART+CIS, combination arm; *CIS*, cisplatin; *CON*, control.

Supplementary Figure 1. Iron homeostasis and glutathione metabolism are deregulated in

G3MB. (A) Genes associated with programmed cell death highlighting ferroptosis as the top

deregulated pathway in MB patients (GSE82517). **(B)** Survival disadvantage for MB patients

with lower ferroptosis scores. **(C)** G3MB tumors have lower ferroptosis scores than G4MBs. **(D)**

Key genes regulating iron homeostasis, glutathione metabolism, and ferroptosis that are enriched

in G3MB (GSE148389, $p<0.05$ for all presented gene comparisons between normal and G3MB).

(E) Enrichment of iron-sulfur cluster binding and downregulation of ROS pathways in G3MB

tumors.

Supplementary Figure 2. ABCB7 is enriched in MB and portends a poor prognosis. (A)

Enrichment plot by KEGG pathways analysis demonstrating a strong negative enrichment score

for the ABC transporter family with miR-1253 overexpression in HDMB03 cells. **(B)**

Deregulated ABC transporters in G3MB (column 1), effect of deregulated expression on survival

(column 2), and transporters significantly downregulated by miR-1253 expression (column 3).

This analysis isolated ABCB7 as the best putative target for miR-1253. **(C)** Expression profile

for ABCB7 in multiple MB cohorts. *CB*, normal cerebellum, Roth *et al.* 2008 (n=9, GSE3526);

MB 1, Gilbertson *et al.* 2012 (n=76, GSE37418); *MB 2*, Pfister *et al.* 2017 (n=223); *MB 3*,

Delattre *et al.* 2012 (n=57); *MB 4*, Kool *et al.* 2009 (n=62, GSE10327). Subgroup-specific

ABCB7 expression in two separate medulloblastoma patient cohorts, i.e. **(D)** Weishaupt *et al.*,

GSE124814; *CB*, normal cerebellum (n=291); *WNT*, wingless (n=118); *SHH*, sonic hedgehog

(n=405); *G3*, group 3 (n=233); *G4*, group 4 (n=530); and **(E)** Luo *et al.*, GSE164677; *CB*,

normal cerebellum (n=4); *WNT*, wingless (n=6); *SHH*, sonic hedgehog (n=20); *G3*, group 3

(n=14); *G4*, group 4 (n=19); Data normalized via RUV method **(D)** or DeSeq2 median of ratios

(MoR) method **(E)** and both analyzed using Mann-Whitney U (*p <0.05, **p <0.01, ***p

<0.001, ****p <0.0001). **(F)** Poor prognostic profile demonstrable in high ABCB7-expressing

medulloblastoma patients (Cavali *et al.* GSE85217).

Supplementary Figure 3. ABCB7 repression triggers iron overload. (A and B) Escalation in

both mitochondrial and cytoplasmic free Fe^{2+} demonstrable with miR-1253 expression, rescued

by iron chelation with DFO. Images captured at 63X. Confocal images showing calcein AM dye

quenching in miR-1253-transfected D425 **(A)** and HDMB03 **(B)** cells indicating high cytosolic

labile iron compared to scramble transfected cells. *DFO*, deferoxamine; *NC*, negative control.

Supplementary Figure 4. ABCB7 repression by MiR-1253 triggers oxidative stress and

lipid peroxidation leading to cell death in G3MB cells. Confocal images showing elevated

mitochondrial $O_2^{\bullet-}$ (MitoSOX™ Red, red) **(A)** and cytosolic H_2O_2 (DCFDA, green) **(B)**

following miR-1253 expression in HDMB03 cells, which could be rescued by ferrostatin. **(C)** A

similar trend noted in D425 cells. **(D)** Higher lipid peroxidation (by BODIPY11 staining) in

miR-1253 transfected D425 cells. Flow cytometry analysis showing significantly higher $O_2^{\bullet-}$

mediated cell death (representing ferroptosis) in miR-1253-expressing **(E)** D425 and **(F)**

HDMB03 cells demonstrable by quantifying cells staining for both Annexin V-Cy5 (apoptosis)

and for $O_2^{\bullet-}$ (Mitosox) (Q2). Images captured either at 20X **(A, B, C)** or 63X **(D)** magnification.

Scale bar 200 µm. Mean ± SD, n=3; *ns*, not significant, *p<0.05, **p<0.01, ***p<0.001,

****p<0.0001. *ERA*, erastin; *FER*, ferrostatin-1; *miR*, miR-1253 overexpression, *NC*, negative

control.

Supplementary Figure 5. GPX4 is a central regulator of ferroptosis that is upregulated in

MB. (A) Schematic showing the important role of GPX4 in mitigating ferroptosis. Image created

with Biorender.com. **(B)** Subgroup-specific GPX4 expression assessment by RNA sequencing

($\log_2$ transcripts per million) of a local medulloblastoma patient cohort (Kanchan *et al.*,

GSE148390) showing deregulation in MB tumors, highest in G3MB. *CB*, pediatric cerebellum

(n=10); *SHH*, sonic hedgehog (n=6); *G3*, group 3 (n=7); *G4*, group 4 (n=12). **(C)** Subgroup-

specific GPX4 expression data from Weishaupt *et al.*, GSE124814; *CB*, normal cerebellum

(n=291); *WNT*, wingless (n=118); *SHH*, sonic hedgehog (n=405); *G3*, group 3 (n=233); *G4*,

group 4 (n=530); Data normalized via RUV method and analyzed using Mann-Whitney U (**p*

<0.05, ***p* <0.01, ****p* <0.001, *****p* <0.0001). **(D)** Enrichment of key proteins involved in

glutathione (GSS, GCLC) and GPX4 (SEPSECS, TRIT1) biosynthesis enriched in G3MB. **(E)**

Confocal microscopic images confirming high GPX4 expression in group 3 MB tumors (n=6)

compared to pediatric cerebellum (n=6) and colocalization to the mitochondria based on COXIV

fluorescence. Images captured at 10X magnification. **(F)** Western blotting analysis showing high

in vitro GPX4 expression in classic MB cell lines (G3: D341, D425, HDMB03; G3/4: D283)

compared to normal human astrocytes (NHA).

Supplementary Figure 6. ABCB7 repression deplete GSH. Inhibitory effect of ABCB7

repression by miR-1253 on total glutathione levels in D425 **(A)** and HDMB03 **(B)** and by si-

ABCB7 (100 nM) **(C)**. Evaluation of oxidized glutathione (GSH) showing punctuated effects in

combination treatment groups (miR + Cis) with rescue in the presence of ferrostatin in miR-

1253-transfected HDMB03 cells **(D)** and with ABCB7^{KO} **(E)**.

Supplementary Figure 7. ABCB7 knockout potentiates cisplatin cytotoxicity in G3MB cells

by ferroptosis. Oxidative stress measured by quantifying **(A)** mitochondrial $O_2^{\bullet-}$ (MitoSOX™

Red, **red**) and **(B)** cytosolic H_2O_2 (DCFDA, **green**) in ABCB7^{KO} HDMB03 cells showing

potentiating effect of cisplatin, and inhibited by ferrostatin-1. Images captured at 20X

magnification. Scale bar 200 µm. **(C)** As measured by Image-iT® Lipid Peroxidation Kit,

confocal images showing the highest lipid peroxidation in combination treatment groups

(ABCB7^{KO} + Cis), again rescued by ferrostatin-1, in ABCB7^{KO} HDMB03 cells. Images captured

at 63X magnification. Mean ± SD, n=3; *p <0.05, **p <0.01, ***p <0.001, ****p <0.0001.

Figure 1

Figure 2

Figure 3

Figure 4

A

NC   ERA

si-ABCB7   si+FER

TMRM

Mean Intensity

20000
15000
10000
5000
0

NC   ERA   si-ABCB7   si+FER
      ****   ***   ns

B

OCR (pmol/min)

Oligo   FCCP   Rotenone

0   10   20   30   40   50   60   70
Time (minutes)

— NC
— miR-1253

****
****

ECAR (mpH/min)

150
100
50
0

0   20   40   60   80
Time (minutes)

◆ NC
● miR-1253

***

Figure 5

Figure 6

A HDMB03
GPX4 Transcript
NC   miR

GPX4
Actin
NC   miR

B D425
Actin
NC   miR

C HDMB03
GPX4 Transcript
WT   KO

GPX4
Actin
WT   KO

D HDMB03
ABCB7 Transcript
-Dox +Dox   -Dox +Dox
24h   96h
****   ****

GPX4 Transcript
-Dox +Dox   -Dox +Dox
24h   96h
****   ****

E Gp3 MB Spearman Correlation
R=0.71, p 1.6e-05
GPX4
ABCB7

F Cytoplasm   Mitochondria
NC miR   NC miR
GPX4
COXIV
Actin

G Cytoplasm   Mitochondria
NC KO   NC KO
GPX4
Actin

H Mitochondria
NC miR si
GPX4
TOMM20

Figure 7

A

ABCB7
GPX4
Actin

VC   Cis

B % Viability (24h)

CIS (0 µM) / CIS (5 µM) / CIS (10 µM)

NC   miR-1253   si-ABCB7

C

(i) D425 — 24h NC IC₅₀ = 16, 24h CIS IC₅₀ = 10, 48h NC IC₅₀ = 13, 48h CIS IC₅₀ = 6.1, 72h NC IC₅₀ = 4.6, 72h CIS IC₅₀ = 2.7

(ii) HDMB03 — 24h NC IC₅₀ = 13, 24h CIS IC₅₀ = 9.2, 48h NC IC₅₀ = 8.8, 48h CIS IC₅₀ = 5.6, 72h NC IC₅₀ = 0.60, 72h CIS IC₅₀ = 0.40

Log Cisplatin (µM)

Fold Reduction	D425	HDMB03
24h	1.6	1.8
48h	2.1	1.5
72h	1.7	1.5

(iii) — BCR IC₅₀ = 13 µM, miR-1253 IC₅₀ = 9 µM, si-ABCB7 IC₅₀ = 7 µM

% Viability at 24h   Cisplatin µM

D VC   CIS

NC
miR-253

Crystal Violet Abs

NC   CIS   miR   miR CIS

E VC   CIS

si-ABCB7

Crystal Violet Abs

NC   CIS   si   si CIS

Figure 8

Figure 9

Figure 10

Figure 11

A

CON: PBS
CIS: 2.5 mg/kg
ART: 100 mg/kg
ART+CIS

D22

7   14   21

B

	Day6	Day22
CON		
CIS		
ART		
ART+CIS		

Day6

Total Flux (p/s)

CON   CIS   ART   ART+CIS

Day22

Total Flux (p/s)

CON   CIS   ART   ART+CIS

Survival

CON
ART
CIS
ART+CIS

Randomization
p = 0.01

Days post-implantation