3D Visualisation of trans-syncytial nanopores provides a pathway for paracellular

diffusion across the human placental syncytiotrophoblast

Short title: Trans-syncytial nanopores

Rohan M Lewis*[1,3], Harikesan Baskaran[4], Jools Green[1], Stanimir Tashev[2], Eleni Paleologou[1],

Emma M Lofthouse[1], Jane K Cleal[1,3], Anton Page[2], David S Chatelet[2], Patricia Goggin[2], Bram G

Sengers[3,4]

[1] University of Southampton, Faculty of Medicine

[2] University of Southampton, Faculty of Medicine, Biomedical Imaging Unit

[3] University of Southampton, Institute for Life Sciences

[4] University of Southampton, Faculty of Engineering and Physical Sciences

*Corresponding Author: Rohan M Lewis, MP 887, IDS Building University of Southampton,

Faculty of Medicine, Southampton General Hospital, rohan.lewis@soton.ac.uk

Abstract

17
The placental syncytiotrophoblast, a syncytium without cell-cell junctions, is the primary
barrier between the mother and the fetus. Despite no apparent anatomical pathway for
paracellular diffusion of solutes across the syncytiotrophoblast size-dependent paracellular
diffusion is observed. Here we report data demonstrating that the syncytiotrophoblast is
punctuated by trans-syncytial nanopores (TSNs). These membrane-bound TSNs directly
connect the maternal and fetal facing sides of the syncytiotrophoblast, providing a pathway
for paracellular diffusion between the mother and fetus. Mathematical modelling of TSN
permeability based on their 3D geometry suggests that 10-37 million TSNs per cm^3 of
placental tissue could explain experimentally observed placental paracellular diffusion. TSNs
may mediate physiological hydrostatic and osmotic pressure homeostasis between the
maternal and fetal circulations but also expose the fetus to pharmaceuticals, environmental
pollutants and nanoparticles.

Introduction

The placenta was once viewed as a perfect barrier, but as the thalidomide tragedy demonstrated, this is not the case [1]. It is now clear that potentially harmful molecules and particulates can cross the placenta and adversely affect fetal development. However, the mechanism by which these molecules and particulates cross the placenta is not always clear [2, 3]. Understanding how these substances cross the placenta is necessary to identify the risks and prevent long-term consequences of these exposures, which may adversely affect fetal and postnatal health [4].

The primary placental barrier is the syncytiotrophoblast, a continuous syncytial monolayer covering the villi at the maternal-fetal interface. As there are no cell-cell junctions in a syncytium, there is no obvious pathway by which paracellular diffusion can occur. Despite the absence of an anatomical pathway for diffusion, there is physiological evidence for size-dependent paracellular diffusion of solutes [5, 6]. Trans-syncytial channels, or nanopores, have been proposed as mediators of trans-syncytial diffusion, however continuous full-width nanopores have not been previously demonstrated in the human placenta [7]. An alternative hypothesis to explain paracellular diffusion is that it occurs through regions of syncytial damage [8]. These hypotheses are not mutually exclusive but establishing mechanisms of fetal exposure is necessary to understand the likely risks of different compounds and develop strategies to mitigate this.

Selective placental transfer of nutrients, IgG, wastes and exogenous toxins is facilitated by membrane transporters and endocytosis [9-11]. However it is not clear how exogenous drugs and toxins reach the fetus as, with a few exceptions such as the apically located exchanger OATP4A1 [10], drug transporters in the placenta mediate efflux from the fetus to the mother [12]. Nanoparticle transfer across the placenta has been observed, but the mechanism is unclear [13]. A more extensive understanding of how metabolites, pharmaceuticals and toxins reach the fetal circulation is necessary to protect fetal health.

Estimates of placental permeability surface area products for hydrophilic solutes have been determined *in vivo* and show size-dependent permeability of solutes, which decreases with increasing molecular radius [5, 6]. In other species with haemochorial placentas, permeability has also been shown to be size-selective with some species having higher or lower overall permeability compared to humans [3, 14]. This data suggests a size-selective permeability of the placenta through low diameter channels [15].

Using serial block-face scanning electron microscopy (SBF SEM) to reconstruct placental ultrastructure in three dimensions, this study demonstrates the presence of full-width trans-syncytial nanopores (TSNs) in the human placenta.

Methods

Term placental tissue was collected after delivery from uncomplicated pregnancies with written informed consent and ethical approval from the Southampton and Southwest Hampshire Local Ethics Committee (11/SC/0529). Villous samples from 8 placentas were processed and imaged using TEM or SBFSEM as described previously [27]. Regions containing TSNs were manually segmented in Avizo v2019.4 (ThermoFisher, Eindhoven).

Modelling of nanopore diffusive transfer capacity

The nanopore diffusive transfer capacity for a particular molecule is determined by its diffusivity in free solution as well as the geometry of the pore, specifically the pore length and cross sectional area. Here we will calculate this geometric contribution in the form of the effective area-over-length ratio A_{pore}/L_{pore} [m], which is a measure for the effect of pore geometry on diffusive transfer, independent of the solute studied. This parameter can then be used to calculate nanopore diffusive transfer using the concentration gradient and diffusion coefficient for any specific solute.

Briefly, the magnitude of the diffusive flux J_{pore} [mol/s] through a pore of uniform cross section is given by:

$$J_{pore} = DA_{pore}\frac{|\Delta C|}{L_{pore}} \qquad\qquad \text{Eq. 1}$$

where D [m^2/s] is the diffusion coefficient, A_{pore} [m^2] the pore cross sectional area, ΔC [mol/m^3] the concentration difference over the pore and L_{pore} [m] the pore length. This can also be expressed in terms of the permeability surface area product PS_{pore} [m^3/s]:

$$J_{pore} = PS_{pore}\Delta C \qquad \text{, where } PS_{pore} = \frac{DA_{pore}}{L_{pore}} \qquad \text{Eq. 2}$$

Since in reality the pore cross sectional area is not uniform, 3D image-based modelling was used to calculate the effective area-over-length ratio A_{pore}/L_{pore} [m] based on Eq. 1 by prescribing the concentration difference and diffusivity and then monitoring the resulting flux in the simulations. Segmented image stacks were imported in Simpleware ScanIP (P-2019.09; Synopsys, Inc., Mountain View, USA). Voxels were resampled isotropically so that the x and y pixel size matched the z axis spacing. Pores were meshed using linear tetrahedral elements. Different coarseness settings were evaluated to establish convergence, with the final number of elements used for the different pores ranging from 1.4–38×10^5.

Steady state diffusion simulations were performed in COMSOL Multiphysics (v5.5; COMSOL AB, Stockholm, Sweden). Since A_{pore}/L_{pore} does not depend on the choice of parameters, $D = 1$ m^2/s was used and a fixed concentration gradient $\Delta C = 1$ mol/m^3 was imposed by constant concentration boundary conditions on the inlet and outlet surface of the pore , while the remaining external surface was subject to no-flux conditions. After simulations, the magnitude of the diffusive flux J_{pore} given by the integral of the solute flux in the direction normal to the pore inlet or outlet taken over the inlet/outlet cross sectional area, was then used to calculate the effective pore area-over-length ratio A_{pore}/L_{pore} based on Eq. 1.

For a specific solute, the permeability surface area product for a single pore PS_{pore} was obtained by multiplying the average A_{pore}/L_{pore} value by the relevant diffusion coefficient (Eq. 2). The number of pores n_{pores} [g^{-1}] per gram of placental tissue was then estimated based on Eq. 3:

$$n_{pores} = \frac{PS_{placenta}}{PS_{pore}} \qquad \text{Eq. 3}$$

where $PS_{placenta}$ [$m^3 \; s^{-1} \; g^{-1}$] is the overall experimentally observed placental permeability surface area product for the corresponding solute from literature.

Results

Trans-syncytial nanopores

This study manually inspected 14 SBFSEM image stacks from five different placentas to identify trans-syncytial nanopores (TSNs) defined as membrane lined pores connecting the apical and basal plasma membranes of the syncytiotrophoblast. These stacks consisted of 7487 SBFSEM images, representing a total volume of 0.00000017 cm^3.

Inspection of serial sections allowed identification of TSNs crossing the syncytiotrophoblast. Continuous TSNs with a clear connection between the apical and basal membranes were identified. Near-continuous TSNs were also identified which had connections to both the apical microvillous and basal plasma membranes but contained discontinuities along the length of the nanopore. Unilateral nanopores, ultrastructurally similar to TSNs but opened from either the apical or basal plasma membrane without connecting to the opposite membrane were identified. An illustration of these nanopore types as we defined then is shown in (figure 1). Regions containing TSNs were manually segmented in Avizo v2019.4 (ThermoFisher, Eindhoven). Cross section from one of these TSNs are shown in figure 2. The lumens of TSNs typically had low electron density, consistent with a primarily fluid filled pore (figure 2a, c and d). However, some TSNs and regions of TSNs were observed with higher electron density indicating diffuse contents (figure 2e). The nanopore shown in figure 2 can be seen reconstructed in 3D (Supplemental movie 1) and a movie showing the nanopore lumen highlighted in sequential sections (supplemental movie 2) can be seen in the supplemental material.

Ten continuous TSNs were identified from three different placentas and these did not have a uniform structure (figure 3). While the least complex TSNs were direct tubes connecting the apical and basal membranes (figure 3c) the most complex full width TSN was a double pore with two bifurcated apical openings (4 in total) and two basal openings (figure 3a). Blind ends are present in the TSNs shown in figure 2d, i and g.

In addition, 20 near-continuous TSNs were identified, with examples in all five placentas studied. In two cases, these near-continuous TSNs were highly complicated with branches, multiple dilations, and blind ends (figure 4b). Finally, 25 unilateral nanopores were identified arising from either the apical (n = 17) or basal (n = 8) plasma membrane of the syncytiotrophoblast but not connecting to the opposing plasma membrane.

On the basal membrane, continuous and near-continuous TSN were observed opening adjacent to basal lamina in 16 cases and regions adjacent to cytotrophoblast in 14 cases.

Of the ten continuous TSNs, eight contained membrane-bound inclusions which, surrounded by pore membrane, formed double-membrane structures (figure 4). Inclusions were also observed in the majority of near-continuous and unilateral nanopores. In three cases, the inclusions appeared to be trophoblastic in origin. In the first case, the inclusion on the apical side appeared to be connected to the syncytiotrophoblast by at least one thin stalk of cytoplasmic material (figure 4a-c). At the available resolution, it was not clear whether other inclusions were also engulfed syncytiotrophoblast. In two cases, an inclusion on the

basal side of the placenta was a clear-cut protrusion from an underlying cytotrophoblast cell
(figure 4d-f).
Another less commonly observed feature associated with TSNs were thin nanopores
within the syncytiotrophoblast, where two closely adjacent membranes were joined by
desmosome like adhesions (figure 5). These were observed in two TEM images and one
SBFSEM image stack. In one case a desmosome associated nanopore was seen almost
crossing the placenta from the basal membrane to near the microvillous membrane (figure
5a). Desmosome associated nanopores were observed in one SBF SEM image stack where
they were found to be ribbon or sheet-like structures 5-16 nm wide and estimated at 200-
800 nm deep (figure 5b). Desmosome containing nanopores could be seen appearing and
disappearing within the same 2D section, which is consistent with a pore rather than a cell-
cell junction. Topologically a cell-cell junction would need to either interact with two cell or
image boundaries or form a circular feature (figure 5c).
164
Modelling diffusion through TSNs
Depending on their geometry, individual nanopores display a large variation in
estimated effective A_{pore}/L_{pore} from 0.14 x 10^{-8} m to 2.08 x 10^{-8} m (figure 2k). The effective
A_{pore}/L_{pore} ratio based on the results of the computational simulations of the ten pores, as
depicted in figure 2, was in the order of 10 nm (8.5 ± 7.2 × 10^{-9} m, mean ± SD). By multiplying
the mean effective A_{pore}/L_{pore} ratio with the diffusion coefficient for particular solutes using
Eq. 2, the average permeability surface area products for a single pore could then be
calculated (table 1). Dividing the experimentally observed placental permeability surface area
products (from the literature) by the average permeability surface area product for a single
pore, resulted in placental pore number estimates between 13 and 37 million pores per gram.
175
Discussion
The demonstration of nanoscale pores punctuating the syncytiotrophoblast
challenges our understanding of the placenta as a barrier. It suggests that rather than being
a continuous physical barrier between mother and fetus, the syncytiotrophoblast is a
molecular sieve facilitating diffusion of small solutes via active maintenance of TSNs.
Computational modelling diffusion though these channels suggests that these solutes may
include nutrients and toxins, so this transfer route may have profound implications for the
fetus.
TSNs were structurally heterogeneous, including simple tubes, tubes with dilated
regions and branched structures with blind ends. Many TSNs also contained inclusions of
membrane-bound material, in three cases trophoblastic in origin but in most cases the origin
was unclear. Some TSNs had multiple openings to the syncytiotrophoblast apical microvillous
or basal membranes. The near-continuous and unilateral nanopores are likely in the process
of formation, degradation or remodelling. It is also possible that near-continuous TSNs are
continuous but that the connections were too thin to observe. The relatively large number of

near-continuous and unilateral nanopores is consistent with TSNs being dynamically
remodelled.

There was a wide range in calculated permeability parameters for the modelled TSNs,
with diffusion through the most permeable being 10 x greater than the least permeable. As
might be expected, shorter, wider TSNs had the greatest permeability while the longer TSNs
had the lowest permeability (excluding the TSN with multiple apical and basal openings).

The time scale for diffusion $T = \frac{L_{pore}^2}{D}$ through the TSNs is < 60 ms for the compounds
studied. As such TSNs will operate under quasi steady state conditions, given that the
timescale of diffusion through the TSN is very fast compared to any changes in concentration
in the much larger volumes of the maternal intervillous space and fetal capillary to either side.

There are differences in the placental permeability of different compounds from
literature, which could not be fully explained based on diffusivity and thus affected the
number of pores estimated, with higher estimated pore numbers for smaller molecules. One
explanation for this is that there may be more thinner pores available that only allow passage
of smaller molecules. However, we do not think that this is the explanation for what is
observed here. Paracellular diffusion across the placenta is size-selective, but the TSNs are
likely to be too wide to impose this size selectivity on smaller molecules. Physiological
estimates of placental pore sizes in rodents are in the order of 17 nm, which is consistent with
the thinnest regions of the nanopores described here and with the desmosome associated
channels we observed [15]. For comparison glucose is 1.5 nm long so would be expected to
diffuse freely though pores of this size. If the TSNs do not impose size selectivity, other
structures such as the trophoblast basal lamina or the capillary endothelium might, and
further research is required to understand this better [16].

Molecules such as IgG or nanoparticles such as ultrafine diesel exhaust could diffuse
into most regions of the TSNs, but the diameter would sterically hinder their transfer. TSN
diameter might be a significant barrier to the transfer of large molecules unless an additional
biological mechanism facilitates this transfer.

On the basal membrane TSNs opened adjacent to both basal lamina and
cytotrophoblast cells. TSNs opening to the basal lamina would provide the most direct
pathway to the fetal circulation. Substances diffusing through TSNs opening adjacent to
cytotrophoblast would have a less direct path to reach the fetus but these solutes could
interact with the cytotrophoblast themselves.

Further research is required to determine TSN densities in placental tissue. An initial
approximation for the TSN density was obtained by dividing the number of observed TSNs (10
continuous or 30 including continuous and near-continuous TSNs) by the imaged volume of
0.00000017 cm^3. This initial approximation for TSN density is 58 million TSNs per cm^3, and
including near-continuous TSNs is 162 million/cm^3. This estimate is in line with modelled
estimate of the number of TSNs required to mediate transfer of known paracellular markers
such as inulin and creatinine.

The 50 nm z-resolution is a potential technical limitation in this study. Firstly, it may
lead us to classify some continuous TSNs as near continuous and secondly could also affect

the accuracy of the geometric representation in the computational models affecting the
calculated permeability parameters. The use of focused ion beam scanning electron
microscopy could reduce z depth, but this approach has a smaller field of view making finding
TSNs more difficult.

The physiological role of TSNs is likely to include the maintenance of ionic and osmotic
homeostasis between mother and fetus. Diffusion of ions and bulk fluid flow would balance
osmotic and pressure gradients that could build up across an impermeable placenta. TSNs
may also facilitate the placental transfer of nutrients, such as glucose, with maternal to fetal
gradients to the fetus [17]. However, the physiological roles of TSNs may come at the cost of
allowing non-selective transfer of potentially harmful substances.

The presence of membrane-bound inclusions within the TSNs suggests their role could
be more complex than simply being pores for diffusion. The nanopore inclusions and
surrounding TSN membrane create a double membrane structure which show ultrastructural
similarities to autophagosomes [18]. Nanopore inclusions can be observed in a previous study
that sought to identify TSNs using lanthanum perfusion [7].

While the origin and nature of the inclusions remains unclear, in some cases they
appeared trophoblastic in origin. One inclusion appeared to be connected to the
syncytiotrophoblast by a stalk of cytoplasmic material which is consistent with
syncytiotrophoblast being pinched off into a nanopore to form an inclusion. If this
interpretation is correct this could potentially represent the initiation of autophagy or
material that will be shed from the placenta. In other cases, inclusions on the basal side of
the placenta were observed to be protrusions from underlying cytotrophoblast cells.

A rare feature associated with TSNs were desmosome associated nanopores which
could be observed arising from basal membrane folds [19], and within the cytoplasm but not
observed connecting to the microvillous membrane. The connect to dilations that appear
similar to, or are, the TSNs that are the main focus of this paper. It is possible that the TSNs
represent formation or breakdown of TSNs or that they are a separate class of nanopore that
intersect with TSNs. The desmosome associated nanopores have been observed previously in
2D TEM images where it was suggested that they are remnants of cytotrophoblast-
syncytiotrophoblast fusion. We cannot exclude this trophoblast fusion explanation, but in the
limited number of 3D examples we identified the structures were more like ribbon than a cell-
cell junctions [20]. Desmosome associated nanopores are very thin and so may not be visible
when the block is in the wrong orientation and so could be more common than they appear.

Cells that cross or penetrate epithelial barriers have been observed to express tight
junction proteins to facilitate movement through cell-cell junctions, and the desmosome
containing nanopores in the syncytiotrophoblast could provide a route and a mechanism by
which maternal or fetal cells could cross the syncytiotrophoblast [21, 22]. Images of
erythrocytes protruding through the syncytiotrophoblast are consistent with distensible
channels that could allow cell transfer [23].

There is considerable diversity in placental structures across species and given the
possibility that the syncytiotrophoblast has evolved independently in different branches of

the evolutionary tree, the diversity of TSNs (or similar structures) is of interest. To date, full
width trans-syncytial channels have only been described in the degu which has both short
direct channels in thin regions of syncytiotrophoblast and more complex channels connecting
infoldings of basal and apical surfaces [24]. The human TSNs had a distinct appearance but
were most similar to the more complex Degu channels, although longer and thinner. The
distribution of TSNs across species may inform their biological role and origins.

Identifying the molecular processes underlying TSN formation will be key to
understanding these structures. While TSNs may form by a unique mechanism, they are more
likely to co-opt at least some of the known molecular mechanisms that mediate endocytosis.
There are multiple endocytic processes, including clathrin and caveolin, micropinosomes or
CLIC/GEEC, which form tubular structures [25]. The CLIC/GEEC may be of particular interest
as these form tube-like structures [26].

In conclusion, this study has demonstrated the existence of TSNs providing a non-
selective diffusion pathway across the placenta. These TSNs may allow pharmacological
drugs, environmental toxins and even particulate pollutants across the syncytiotrophoblast.
Accurately determining the density of TSNs in healthy pregnancy is necessary to confirm the
capacity of TSNs to mediate transfer. Furthermore, establishing the density of TSNs in disease
may provide novel insights into disease processes.

Acknowledgements: This work was supported by BBSRC project grant BB/R002762/1.
Equipment in the Biomedical Imaging Unit was supported by MR/L012626/1 Southampton
Imaging under MRC UKRMP Funding.

Competing interests: The authors have no competing interests to declare

References

1. Dally A. Thalidomide: was the tragedy preventable? Lancet. 1998;351(9110):1197-9. Epub 1998/06/27. doi: 10.1016/S0140-6736(97)09038-7. PubMed PMID: 9643709.

2. Lofthouse EM, Torrens C, Manousopoulou A, Nahar M, Cleal JK, O'Kelly IM, et al. Ursodeoxycholic acid inhibits uptake and vasoconstrictor effects of taurocholate in human placenta. FASEB journal : official publication of the Federation of American Societies for Experimental Biology. 2019;33(7):8211-20. Epub 2019/03/30. doi: 10.1096/fj.201900015RR. PubMed PMID: 30922127; PubMed Central PMCID: PMCPMC6593889.

3. Bove H, Bongaerts E, Slenders E, Bijnens EM, Saenen ND, Gyselaers W, et al. Ambient black carbon particles reach the fetal side of human placenta. Nat Commun. 2019;10(1):3866. Epub 2019/09/19. doi: 10.1038/s41467-019-11654-3. PubMed PMID: 31530803; PubMed Central PMCID: PMCPMC6748955.

4. Lewis RM, Demmelmair H, Gaillard R, Godfrey KM, Hauguel-de Mouzon S, Huppertz B, et al. The Placental Exposome: Placental Determinants of Fetal Adiposity and Postnatal Body Composition. Annals of Nutrition and Metabolism. 2013;63(3):208-15. doi: 10.1159/000355222. PubMed PMID: WOS:000329057400005.

5. Thornburg KL, Burry KJ, Adams AK, Kirk EP, Faber JJ. Permeability of placenta to inulin. American journal of obstetrics and gynecology. 1988;158(5):1165-9. Epub 1988/05/01. doi: 10.1016/0002-9378(88)90246-3. PubMed PMID: 3369499.

6. Bain MD, Copas DK, Taylor A, Landon MJ, Stacey TE. Permeability of the human placenta in vivo to four non-metabolized hydrophilic molecules. The Journal of physiology. 1990;431:505-13. PubMed PMID: 2129229; PubMed Central PMCID: PMCPMC1181787.

7. Kertschanska S, Kosanke G, Kaufmann P. Pressure dependence of so-called transtrophoblastic channels during fetal perfusion of human placental villi. Microscopy research and technique. 1997;38(1-2):52-62. doi: 10.1002/(SICI)1097-0029(19970701/15)38:1/2<52::AID-JEMT7>3.0.CO;2-W. PubMed PMID: 9260837.

8. Brownbill P, Mahendran D, Owen D, Swanson P, Thornburg KL, Nelson DM, et al. Denudations as paracellular routes for alphafetoprotein and creatinine across the human syncytiotrophoblast. American journal of physiology Regulatory, integrative and comparative physiology. 2000;278(3):R677-83. PubMed PMID: 10712288.

9. Cleal JK, Glazier JD, Ntani G, Crozier SR, Day PE, Harvey NC, et al. Facilitated transporters mediate net efflux of amino acids to the fetus across the basal membrane of the placental syncytiotrophoblast. The Journal of physiology. 2011;589(4):987-97. doi: 10.1113/jphysiol.2010.198549. PubMed PMID: WOS:000287743400026.

10. Lofthouse EM, Cleal JK, O'Kelly IM, Sengers BG, Lewis RM. Estrone sulphate uptake by the microvillous membrane of placental syncytiotrophoblast is coupled to glutamate efflux. Biochem Biophys Res Commun. 2018;506(1):237-42. Epub 2018/10/23. doi: 10.1016/j.bbrc.2018.10.074. PubMed PMID: 30343886.

11. Firan M, Bawdon R, Radu C, Ober RJ, Eaken D, Antohe F, et al. The MHC class I-related receptor, FcRn, plays an essential role in the maternofetal transfer of gamma-globulin in humans. Int Immunol. 2001;13(8):993-1002. Epub 2001/07/27. doi: 10.1093/intimm/13.8.993. PubMed PMID: 11470769.

12. Iqbal M, Audette MC, Petropoulos S, Gibb W, Matthews SG. Placental drug transporters and their role in fetal protection. Placenta. 2012;33(3):137-42. Epub 2012/01/24. doi: 10.1016/j.placenta.2012.01.008. PubMed PMID: 22265855.

13. Gruber MM, Hirschmugl B, Berger N, Holter M, Radulovic S, Leitinger G, et al. Plasma proteins facilitates placental transfer of polystyrene particles. J Nanobiotechnology.

2020;18(1):128. Epub 2020/09/11. doi: 10.1186/s12951-020-00676-5. PubMed PMID:
32907583; PubMed Central PMCID: PMCPMC7487953.

14. Sibley CP. Understanding placental nutrient transfer--why bother? New biomarkers of
fetal growth. The Journal of physiology. 2009;587(Pt 14):3431-40. doi:
10.1113/jphysiol.2009.172403. PubMed PMID: 19417095; PubMed Central PMCID:
PMC2742272.

15. Stulc J. Extracellular transport pathways in the haemochorial placenta. Placenta.
1989;10(1):113-9. Epub 1989/01/01. PubMed PMID: 2654915.

16. Harris SE, Matthews KS, Palaiologou E, Tashev SA, Lofthouse EM, Pearson-Farr J, et al.
Pericytes on placental capillaries in terminal villi preferentially cover endothelial junctions in
regions furthest away from the trophoblast. Placenta. 2021;104:1-7. Epub 2020/11/16. doi:
10.1016/j.placenta.2020.10.032. PubMed PMID: 33190063; PubMed Central PMCID:
PMCPMC7921774.

17. Day PE, Cleal JK, Lofthouse EM, Hanson MA, Lewis RM. What factors determine
placental glucose transfer kinetics? Placenta. 2013;34(10):953-8. doi:
10.1016/j.placenta.2013.07.001. PubMed PMID: WOS:000330331500016.

18. Mizushima N, Yoshimori T, Levine B. Methods in mammalian autophagy research. Cell.
2010;140(3):313-26. Epub 2010/02/11. doi: 10.1016/j.cell.2010.01.028. PubMed PMID:
20144757; PubMed Central PMCID: PMCPMC2852113.

19. Tashev SA, Parsons D, Hillman C, Harris S, Lofthouse EM, Goggin P, et al. Folding of the
syncytiotrophoblast basal plasma membrane increases the surface area available for
exchange in human placenta. Placenta. 2022;117:57-63. Epub 20211104. doi:
10.1016/j.placenta.2021.11.002. PubMed PMID: 34768170.

20. Jones CJ, Fox H. Ultrastructure of the normal human placenta. Electron Microsc Rev.
1991;4(1):129-78. PubMed PMID: 1873486.

21. Rescigno M, Urbano M, Valzasina B, Francolini M, Rotta G, Bonasio R, et al. Dendritic
cells express tight junction proteins and penetrate gut epithelial monolayers to sample
bacteria. Nat Immunol. 2001;2(4):361-7. Epub 2001/03/29. doi: 10.1038/86373. PubMed
PMID: 11276208.

22. Murrieta-Coxca JM, Aengenheister L, Schmidt A, Markert UR, Buerki-Thurnherr T,
Morales-Prieto DM. Addressing microchimerism in pregnancy by ex vivo human placenta
perfusion. Placenta. 2021;117:78-86. Epub 2021/11/14. doi: 10.1016/j.placenta.2021.10.004.
PubMed PMID: 34773744.

23. Palaiologou E, Goggin P, Chatelet DS, Lofthouse EM, Torrens C, Sengers BG, et al. Serial
block-face scanning electron microscopy of erythrocytes protruding through the human
placental syncytiotrophoblast. Journal of anatomy. 2017;231(4):634-7. Epub 2017/07/18. doi:
10.1111/joa.12658. PubMed PMID: 28714092; PubMed Central PMCID: PMCPMC5603844.

24. King BF. Ultrastructural evidence for transtrophoblastic channels in the
hemomonochorial placenta of the degu (Octodon degus). Placenta. 1992;13(1):35-41. Epub
1992/01/01. PubMed PMID: 1502135.

25. Cooke LDF, Tumbarello DA, Harvey NC, Sethi JK, Lewis RM, Cleal JK. Endocytosis in the
placenta: An undervalued mediator of placental transfer. Placenta. 2021;113:67-73. Epub
2021/05/18. doi: 10.1016/j.placenta.2021.04.014. PubMed PMID: 33994009.

26. Ferreira APA, Boucrot E. Mechanisms of Carrier Formation during Clathrin-
Independent Endocytosis. Trends Cell Biol. 2018;28(3):188-200. Epub 2017/12/16. doi:
10.1016/j.tcb.2017.11.004. PubMed PMID: 29241687.

27.   Palaiologou E, Etter O, Goggin P, Chatelet DS, Johnston DA, Lofthouse EM, et al.
Human placental villi contain stromal macrovesicles associated with networks of stellate cells.
Journal of anatomy. 2020;236(1):132-41. Epub 2019/09/13. doi: 10.1111/joa.13082. PubMed
PMID: 31512233.
28.   Atkinson DE, Robinson NR, Sibley CP. Development of passive permeability
characteristics of rat placenta during the last third of gestation. The American journal of
physiology.   1991;261(6   Pt   2):R1461-4.   Epub   1991/12/01.   doi:
10.1152/ajpregu.1991.261.6.R1461. PubMed PMID: 1750569.
29.   Cleal JK, Brownbill P, Godfrey KM, Jackson JM, Jackson AA, Sibley CP, et al.
Modification of fetal plasma amino acid composition by placental amino acid exchangers in
vitro. The Journal of physiology. 2007;582(Pt 2):871-82. doi: 10.1113/jphysiol.2007.130690.
PubMed PMID: 17478537; PubMed Central PMCID: PMCPMC2075319.
30.   Edwards D, Jones CJ, Sibley CP, Nelson DM. Paracellular permeability pathways in the
human placenta: a quantitative and morphological study of maternal-fetal transfer of
horseradish peroxidase. Placenta. 1993;14(1):63-73. doi: 10.1016/s0143-4004(05)80249-8.
PubMed PMID: 8456090.
31.   Brownbill P, Edwards D, Jones C, Mahendran D, Owen D, Sibley C, et al. Mechanisms
of alphafetoprotein transfer in the perfused human placental cotyledon from uncomplicated
pregnancy. The Journal of clinical investigation. 1995;96(5):2220-6. doi: 10.1172/JCI118277.
PubMed PMID: 7593608; PubMed Central PMCID: PMCPMC185872.
32.   Collins MC, Ramirez WF. Transport through polymeric membranes. J Phys Chem
1979;83:2294–301. doi: https://doi.org/10.1021/j100480a022.
412

413   **Figures and tables**

415   *Table 1: Estimated number of pores per gram required to explain the experimentally*

416   *observed placental permeability surface area products.*

	MW (da)	D (m² s⁻¹)	PS_{pore} (m³ s⁻¹)	$PS_{placenta}$ (m³ s⁻¹ g⁻¹)	number of pores (g⁻¹)
Inulin*	6179	2.6×10^{-10}	2.20×10^{-18}	0.28×10^{-10}	13×10^{6}
CrEDTA*	344	7.0×10^{-10}	5.92×10^{-18}	1.06×10^{-10}	18×10^{6}
Mannitol*	182	9.9×10^{-10}	8.37×10^{-18}	2.50×10^{-10}	30×10^{6}
Creatinine**	113	12.9×10^{-10}	10.9×10^{-18}	4.00×10^{-10}	37×10^{6}

*Placental permeability surface area product data taken from Bain *et al.* (1990) [6] and diffusion coefficients
from Atkinson *et al.* (1991) [28]. ** Placental permeability surface area product based on the average of values
from Cleal *et al.* (2007)[29], Edwards et al 1993 [30] & Brownbill et al 1995 [31] and diffusion coefficient from
Collins *et al.* (1979) [32].

Figure 1, Classification of nanopores found in human placental syncytiotrophoblast. A)
continuous TSNs connecting the apical and basal membrane without any breaks. **B)** near-
continuous TSNs with sections connected to the apical and basal plasma membranes but
contained discontinuities where no clear connection could be observed but the ends of the
discontinuous sections were adjacent. **C & D)** Unilateral nanopores were ultrastructurally
similar to TSNs but opened from either the apical or basal plasma membrane but without any
apparent connection to the opposing membrane. **E)** Desmosome associated nanopores were
also observed arising from the basal membrane but none of these could be observed
connecting to the apical membrane.

Figure 2, SBF SEM serial sectioning allows identification of TSNs crossing the human placental syncytiotrophoblast. A) An SBF SEM image showing a cross section of the syncytiotrophoblast (STB) from the maternal intervillous space (IVS) to the underlying cytotrophoblast (CTB). Within this image a cross section of a nanopore can be seen (white arrow) illustrating how difficult it would be to identify these structures from individual 2D images. The inset shows the nanopore reconstructed in 3D from 44 consecutive sections with the white arrow indicating where the section comes from and modelled solute concentration indicated by colour from high (red) to low (blue). **B)** the apical opening of the TSN with the inset showing this highlighted in green. **C)** A thin region of nanopore with the inset showing this highlighted in green. **D)** a dilated region of the nanopore with the inset showing this highlighted in green. In B, C and D the nanopore has a low electron density consistent with a fluid filled pore **E)** A region of nanopore where the lumen has a higher electron density than in other regions. The inset showing this cross section of the nanopore highlighted in green. This region is close to the end of the nanopore and the cytotrophoblast boundary can be seen bottom left. * Indicates inclusions in other TSNs in the section. A movie showing the nanopore reconstructed in 3D and another movie with the nanopore lumen highlighted in sequential sections can be seen in the supplemental material.

Figure 3, **Modelled solute concentration gradients within the continuous TSNs.** High concentrations (red) were applied to the maternal facing TSN opening, and concentration gradients are shown for diffusion through the channel. The molecular flux associated with these gradients was used to calculate the effective cross sectional surface area used to calculate permeability. **A-J)** show the nanopores presented in order of calculated permeability from highest to lowest. The apical opening is presented on top right and basal openings on the bottom or left. The TSN shown in A is a double pore with four apical and basal openings and a connection on the basal side which accounts for its higher permeability despite its longer length. Blind ends are present in D, I and G. TSN A has two openings to the microvillous membrane and two openings to the basal membrane of the syncytiotrophoblast. **J)** shows a scatterplot showing the distribution of permeabilities with the line indicating the median value.

483
Figure 4, **Examples of TSN inclusions formed from engulfed syncytiotrophoblast or villous cytotrophoblast. A-C)** An
example of an apical nanopore inclusion (highlighted by an *) which in 3D can be seen to be budding off the
syncytiotrophoblast within the nanopore. **A)** shows a SBF SEM image containing a cross section of the TSN inclusion. **B)** shows
the pink inclusion reconstructed in 3D within the TSN formed from green syncytiotrophoblast. This inclusion lies within a
complex near-continuous TSN which, apart from regions around the inclusion, is shown in red. This TSN has discontinuities
indicated by white arrows but gives the appearance of being part of a trans-syncytial network. A black arrow indicates the
apical opening of this TSN. **C)** shows a higher magnification image of the pink inclusion and the connection to the green
syncytiotrophoblast is indicted by a white arrow. **D-F)** An example of a TSN inclusion, which in 3D is seen to be derived from
underlying villous cytotrophoblast. In panels D-F an * indicates the inclusion, white arrows indicate the lumen of the TSN and
black arrow indicates the apical opening of the TSN on the microvillous membrane. **D)** shows a 2D SBF SEM image of the
syncytiotrophoblast containing a cross section of a TSN and an inclusion containing an inclusion. **E)** shows the 3D structure
of this region with the villous cytotrophoblast in yellow and the syncytiotrophoblast in green. **F)** shows a higher magnification
image of the structure of the villous cytotrophoblast derived inclusion and the TSN channel. BM and MVM are the basal
membrane and apical microvillous membranes of the syncytiotrophoblast.

Figure 5, Desmosome associated nanopores. A) a TEM image of a desmosome associated
nanopore traversing most of the width of the syncytiotrophoblast (white dotted line). The
desmosomes are indicated by black arrows. This type of nanopore was associated with
dilations typical of the TSNs described in the previous figures (indicated by *). **B)** an SBF SEM
image of a desmosome associated nanopore connecting basal membrane folds to a TSN like
opening (indicated by *). **C)** A TEM image of a desmosome associated nanopore which begins
and ends in the same field of view (red arrows). This demonstrates that these structures are
not a cell-cell junction, as if this were the case it would need to intersect 2 sides of the image
or be a circle. This channel contains an empty TSN like dilation (indicated by the top*) and
one which contains inclusion material (indicated by the bottom*). EC = endothelial cell, VS =
villous stroma, BL = basal lamina, STB = syncytiotrophoblast, BMF = basal membrane folds.
IVS = intervillous space.