Title: Inactivity and Ca^{2+} signaling regulate synaptic compensation in motoneurons

following hibernation in American bullfrogs

Authors: Tanya Zubov, Lara do Amaral-Silva, Joseph M. Santin*

*Email for correspondence: jmsantin@uncg.edu

Affiliations: [1]The University of North Carolina at Greensboro, Department of Biology

Keywords: homeostatic plasticity, synaptic scaling, AMPA receptor, compensation, control of breathing

Abstract

Neural networks tune synaptic and cellular properties to produce stable activity.

One form of homeostatic regulation involves scaling the strength of synapses up or

down in a global and multiplicative manner to oppose activity disturbances. In American

bullfrogs, excitatory synapses scale up to regulate breathing motor function after

inactivity in hibernation, connecting homeostatic compensation to motor behavior. In

traditional models of homeostatic synaptic plasticity, inactivity is thought to increase

synaptic strength *via* mechanisms that involve reduced Ca^{2+} influx through voltage-

gated channels. Therefore, we tested whether pharmacological inactivity and inhibition

of voltage-gated Ca^{2+} channels are sufficient to drive synaptic compensation in this

system. For this, we chronically exposed *in vitro* brainstem preparations containing the

intact respiratory network to tetrodotoxin (TTX) to stop activity and nimodipine to block

L-type Ca^{2+} channels. We show that hibernation and TTX similarly increased

motoneuron synaptic strength and that hibernation occluded the response to TTX. In

contrast, block of L-type Ca^{2+} channels did not alter mean synaptic strength but

disrupted the multiplicative scaling of synaptic compensation typically observed in

response to hibernation. Thus, inactivity associated with hibernation drives up synaptic

strength through mechanisms that do not rely on reduced L-type channel function, while

Ca^{2+} signaling independently regulates the balance of synaptic weights. Altogether,

these results point to multiple feedback signals for shaping synaptic compensation that

gives rise to proper network function during environmental challenges *in vivo*.

Introduction

Neural circuits must produce correct output patterns to generate reliable behaviors. However, animals face constantly changing internal and external environments, posing a threat to stable neuronal function [1]. To maintain stability, the nervous system is thought to use a wide range of homeostatic mechanisms that sense activity perturbations which then modify cellular and synaptic function to regulate neuronal output. Although diverse homeostatic mechanisms exist [2,3], the most well-studied involves compensatory changes in excitatory synaptic strength through a mechanism termed synaptic scaling. Synaptic scaling is interpreted to involve the up or downregulation of all a neuron's synaptic inputs by the same relative amount to oppose activity disturbances [4]. Mounting evidence implicates synaptic scaling in association with diverse physiological conditions such as development [5,6], sleep [7], and learning [8] in mammals, while inappropriate function of homeostatic mechanisms may lead to various neurological disorders [9-11]. Although a large body of work supports cell-wide multiplicative synaptic compensation [4,5,8,12-15], how these processes are regulated and integrated *in vivo* to generate adaptive behaviors is less clear.

Animals that hibernate must enter and exit states of long-term dormancy, making excellent models to connect activity-dependent neural compensation to behavior [16]. Motor circuits that produce breathing in American bullfrogs, *Lithobates catesbeianus*, are especially intriguing in this regard. The respiratory network typically generates stable rhythmic motor activity throughout life. However, during under water hibernation, gas exchange demands are met through cutaneous respiration, and the motor circuits that produce lung breathing stop entirely [17-19]. After months of inactivity, frogs emerge in

the spring and quickly display normal respiratory motor performance [17,20]. We previously

found that motoneurons scale up excitatory synapses in response to hibernation, which

acts to maintain respiratory motor outflow immediately after the animal restarts

breathing [15]. Therefore, this system allows the study of synaptic compensation in

relation to motor behavior and points to conserved mechanisms across mammals and

amphibians in the natural environment.

In mammalian systems, synaptic compensation is thought to be induced by

variables related to inactivity, for example, decreases in firing rate or postsynaptic

receptor activation [21-23]. From there, decreases in intracellular calcium represents a

message that is proportional to activity, with the capacity to influence Ca^{2+}-dependent

signaling that drives compensation [24]. Reduced Ca^{2+} influx through L-type calcium

channels in particular plays a critical role in eliciting synaptic compensation in diverse

neuronal types [22,25,26]. Given that hibernation causes a large and chronic reduction in

neuronal output, inactivity and subsequent reductions in Ca^{2+} influx may trigger synaptic

compensation that serves to regulate motor performance [15,16]. However, aquatic

hibernation involves diverse physiological changes to the animal [27], raising questions

about whether inactivity or other aspects of the organismal environment, such as

temperature, feedback to control synaptic compensation. To begin to approach this

question, we tested the hypothesis that inactivity and reduced voltage-gated Ca^{2+}

channel function are sufficient to drive up synaptic strength in frog respiratory

motoneurons. The amphibian respiratory network is well-suited to address this question,

as "fictive breathing" motor patterns persist in dissected brainstem preparations for >24

hours *in vitro* [28]. This allowed us to perform chronic pharmacological manipulations to

the native network without the need for cell culture conditions. To this end, we isolated

the respiratory network *in vitro*, silenced activity with tetrodotoxin (TTX) or blocked L-

type Ca^{2+} channels with nimodipine. We then compared synaptic responses to these

manipulations with that seen during hibernation and then tested whether hibernation

occludes the synaptic response induced by these drugs. This allowed us to infer

whether network inactivity and Ca^{2+} channel blockade overlap with pathways associated

with hibernation, a similar "sufficiency/occlusion" approach used to relate long-term

potentiation of synaptic strength to learning at hippocampal synapses [29].

Results

Amphibians inflate the lungs with positive pressure, whereby muscles of the buccal floor and glottis form a pump-valve system for lung ventilation. Laryngeal motoneurons that innervate the glottal dilator, the value muscle, (Fig. 1A) undergo compensatory upregulation of AMPA-glutamate receptor currents in response to ~2 months of aquatic hibernation [15]. To simulate the overwintering environment in this study, frogs were cooled from 20°C to 2°C over one week and then held at 2°C for either two or four weeks before assessing synaptic strength in brain slices. Consistent with our previous report with inactivity at longer time scales, synaptic strength, as assessed by the amplitude of spontaneous excitatory postsynaptic currents (sEPSC), increased in frogs exposed to the simulated hibernation environment at both two and four weeks (Fig. 1B) with no change in sEPSC frequency (Fig. 1C). These results confirm postsynaptic compensation in response to hibernation and demonstrate that upregulation of excitatory synaptic strength plateaus at a new steady-state value relatively early into aquatic submergence.

Variables associated with inactivity (*i.e.*, loss of postsynaptic firing or synaptic transmission) can drive homeostatic increases in synaptic strength [21,22]. Our rationale was that if synaptic compensation during winter is responding to inactivity, pharmacological inactivity may produce a similar degree of synaptic compensation as hibernation, thus demonstrating inactivity as a sufficient signal to generate the hibernation response. To test this, we silenced intact respiratory networks in the brainstem-spinal cord preparation with 100 nM TTX for 16 hours (Fig. 2A). While the time course of hibernation (one month) and shorter-term TTX (16 hours) are different,

synaptic scaling has been observed to manifest after 4 hours of inactivity *in vitro* [22]. We

compared average miniature excitatory postsynaptic current amplitude in slices

(mESPCs) following network activity induced by TTX to Time Controls (control

preparations with rhythmic activity present for 16 hrs at 22°C) and Post-Hibernation

preparations (preparations recovered following hibernation for 16 hrs at 22°C; Fig 2A).

TTX and hibernation groups exhibited similar mean increases in cell-averaged mEPSC

amplitude relative to Time Control preparations. These mean increases were not

different from each other (Fig. 2B-C). Therefore, pharmacological inactivity produces a

similar average amount of synaptic compensation as physiological inactivity in

hibernation.

The previous experiment demonstrated that inactivity *via* TTX is sufficient to drive

synaptic compensation in this system. Next, we determined if hibernation occludes the

synaptic response to TTX. If hibernation prevents further increases by TTX, this may

reflect overlapping signaling processes. However, if TTX and hibernation produce

additive responses, this may suggest divergent signaling pathways for compensation.

When we exposed post-hibernation preparations to TTX for 16 hr, we did not observe

further increases in synaptic strength by TTX compared to hibernation alone (Fig. 3A-

B). Membrane properties, mEPSC frequency, and mEPSC decay time constant were

not different across control, post-hibernation, TTX, and post-hibernation+TTX groups

(Fig. 4). Therefore, these data suggest that hibernation occludes the synaptic

compensation response to TTX.

To upregulate synaptic strength in an inactivity-dependent manner, we

hypothesized that reduced Ca^{2+} influx through voltage-gated Ca^{2+} channels may play a

role. Thus, we exposed rhythmic circuit preparations to 10 µM nimodipine, a blocker of

L-type Ca^{2+} channels in motoneurons of this system [30]. Exposure of warm-acclimated

preparations to nimodipine did not affect mean mEPSC amplitude or the distribution

relative to warm-acclimated time controls (Fig. 5A-B). Therefore, block of L-type Ca^{2+}

does not mimic the synaptic response to TTX and hibernation. Post-hibernation

preparations exposed to nimodipine still resulted in upregulation of cell-averaged

mEPSCs compared to time controls (Fig 5A-B), just as we observed with TTX,

hibernation, and hibernation+TTX. When comparing the post-hibernation+nimodipine

group to post-hibernation alone, mEPSC amplitude were not different (p=0.470, Fig.

3C). Overall, reduced L-type channel function does not drive compensation in this

system.

Given that the entire mEPSC amplitude distribution is thought to be regulated by

a multiplicative scaling factor during inactivity [24], we tested whether the increase in

postsynaptic strength in each of our experimental conditions increases according to a

multiplicative scaling rule. We used the rank-order method to test for synaptic scaling of

mESPC amplitudes in populations of neurons [4,8,31]. To do this, mEPSCs from control

and treatment groups (TTX, hibernation, hibernation+TTX, hibernation+nimodipine)

were rank-ordered from the smallest to largest amplitude, plotted against each other

(ranked control vs. ranked treatment), and fit with a linear regression. The treatment

distribution for each form of inactivity was then mathematically downscaled by the

equation of the line, and the scaled distribution was compared to control. Positive

evidence for multiplicative scaling is usually taken as a statistically indistinguishable

matching between the scaled and control distributions of mEPSC amplitudes [12,32].

As expected, all treatments strongly right-shifted the amplitude distribution of

mEPSCs relative to controls, as expected based on the mean upregulated synaptic

strength for all conditions (Fig. 6A-E; K-S test; time control vs. all treatment conditions;

$p < 0.0001$). When we performed the downscaling transformation on each treatment

distribution, interestingly, not all showed evidence of multiplicative synaptic scaling.

Hibernation and hibernation+TTX, led to multiplicative scaling (Fig. 6B-C), but TTX

alone (Fig. 6A) and hibernation in the presence of nimodipine (Fig. 6D-E) did not. These

results indicate that hibernation upregulates synaptic drive while maintaining the

balance of synaptic weights according to scaling rule as we previously showed [15].

However, the scaling organization appears to be regulated or maintained by active Ca^{2+}

signaling through L-type channels, as blocking voltage-gated Ca^{2+} channels

disorganized the scaling relationship typical of hibernation.

To infer the role of these synaptic modifications on network function, we

assessed circuit activity of the intact respiratory network. The respiratory network is a

central pattern generating network which permits the recording of these motor neurons

as a receive their normal presynaptic inputs *ex vivo* (Fig. 7A). In this preparation,

extracellular population motor burst frequency of the cut nerve root provides information

regarding the function of the rhythm generating networks [33], and integrated burst shape

relates to motoneuron firing and/or recruitment [34]. Thus, we were able to observe

physiological motor output from motoneurons under conditions with no changes in

synaptic strength (control and warm-nimodipine), upregulated and scaled synaptic

strength (post-hibernation), and upregulated but not scaled synaptic strength (post-

hibernation+nimodipine). Across controls, post-hibernation, and post-

hibernation+nimodipine, no differences existed in frequency of the respiratory rhythm at

13 hrs *in vitro* (one-way ANOVA, p=0.3215). However, preparations treated with

nimodipine after hibernation had shorter motor burst widths relative to all other groups

after 13 hrs (Fig. 7 B-C). Burst width narrowing appeared to be a chronic effect induced

by nimodipine following hibernation, as acute application did not affect burst width

before and after winter, and all preparations without nimodipine (control or post-

hibernation) did not change with time (Fig. 7C). In addition, membrane potential, input

resistance, and action potential firing frequency were not different across groups,

indicating chronic nimodipine does not induce changes in intrinsic membrane properties

(Fig. 7D-F). Overall, these experiments suggest that inactivity scales up synaptic

strength in response to hibernation and proper scaling of the mESPC distribution

appears to be important for normal network function associated with breathing after

hibernation.

Discussion

Neural circuits can compensate for activity perturbations with the apparent goal

of regulating neuronal output. American bullfrogs must produce appropriate motor drive

for breathing after potentially months of network inactivity during hibernation. Our

previous results showed that scaling up the strength of motoneuron synapses helps

promote motor output after hibernation to generate breathing. Data here support the

idea that inactivity during hibernation plays a key role in this response. However,

upregulation of synaptic strength did not occur *via* the reduction of Ca^{2+}-dependent

signaling through L-type Ca^{2+} channels as occurs in other models of synaptic

compensation. On the other hand, the precise distribution of synaptic weights is

maintained by Ca^{2+}-sensitive feedback after hibernation to support motor performance.

Overall, these results suggest that multiple signals interact to regulate synaptic

compensation in response to environmental challenges *in vivo*.

Inactivity drives up synaptic strength in response to hibernation

Decades of work support the view that neurons upregulate synaptic strength in

response to inactivity. Given that cold temperatures silence the respiratory network [17,18],

we hypothesized that inactivity may trigger scaling up of synaptic strength in response

to hibernation. Previously, we showed that synaptic strength is increased at ~2 months

of hibernation. Here we demonstrate that increases in synaptic drive are present at 2

weeks into aquatic hibernation, which then saturate a new steady-state level (Fig. 1B).

These responses were comparable to inactivity by TTX (Fig. 2C) and are therefore

consistent with upregulation of synaptic strength in response to a signal related to

inactivity. Further supporting this conclusion, TTX did not cause greater increases in

synaptic strength when applied after hibernation, which suggests the mechanisms

involved in enhancing synaptic strength were already engaged by hibernation. In this

regard, inactivity may upregulate synaptic strength through mechanisms that involve the

loss of spiking, reduced AMPA receptor activation, and reduced neuromodulation

[21,35,36]. That said, synaptic compensation does not seem to strictly follow negative

feedback regulation as predicted for activity-dependent homeostatic plasticity. First,

inactivity over vastly different time scales (16 hours to 1 month) produced similar

degrees of mean synaptic strengthening. This is inconsistent with compensation that

follows the accumulation of an error signal over time while the neuron remains below its

set point [4,37]. Additionally, synaptic strength remained elevated despite the return of

activity for at least 16 hrs after hibernation and did not differ when activity was present

or absent during the post-hibernation period. Thus, our results indicate that inactivity

likely drives up synaptic strength but a more complicated regulation scheme appears to

influence the maintenance of this plasticity.

The previous results indicated that inactivity plays a role in driving up synaptic

strength. L-type Ca^{2+} channels are attractive effectors of this response as they

communicate changes in intracellular Ca^{2+} to the nucleus and other intracellular

signaling pathways that elicit synaptic compensation during inactivity [22,25]. Thus, we

inhibited L-type Ca^{2+} channels to understand if the "inactivity signal" during hibernation

and TTX involved reduced L-type Ca^{2+} channel activity. In contrast to this hypothesis,

chronic block of L-type Ca^{2+} channels did not affect mEPSC amplitude. Although our

data rule out the possibly for a role by nimodipine-sensitive L-type Ca^{2+} channels in

compensation, other Ca^{2+} channels may contribute. For example, in respiratory motor

neurons of this species, nimodipine blocks the high-voltage activated Ca^{2+} current by

~30-40% [30], indicating the expression of other somatodendritic calcium channels on

respiratory motoneurons. Nevertheless, L-type Ca^{2+} channels do not appear to drive

synaptic compensation, differing from traditional models in neuronal cultures [22,25,26,38].

Future work should address the signaling processes that read out inactivity to initiate

and maintain synaptic compensation throughout hibernation.

L-type Ca^{2+} channels maintain multiplicative scaling after hibernation

As a compensation response, synaptic inputs are often upregulated across the

entire distribution by the same multiplicative factor [4]. An interesting finding of this study

is hibernation, but not TTX, led to multiplicative scaling when we performed the

standard rank-order test [4,8,14]. In line with our results, TTX applied to the hippocampus

in vivo drives up synaptic strength without evidence of scaling as determined by the

rank-order test [39]. Additionally, a recent study also suggested that TTX may not lead to

uniform scaling in different mammalian neuronal types when assessed using the rank-

order method [40], although there is significant evidence to the contrary [12,22,41,42]. While

we used the standard method to infer the presence or absence of scaling, we note

recent studies suggesting complexity in this interpretation. Specifically, synaptic spine

density and AMPAR expression across individual synapses do not scale uniformly

within the same neuron even though multiplicative scaling appears to occur at the

population level [43,44]. A non-zero y-intercept in the linear model used to test for scaling

may also suggest that additive processes work alongside multiplicative mechanisms [41];

however, the y-intercept has been interpreted to account for mEPSCs that fall below the

detection cutoff to align control and scaled distributions for the assessment of

multiplicative scaling [4,32]. In support of the later possibility, the y-intercept values we

obtained in the two hibernation groups that exhibited evidence for multiplicative scaling

compensated for our 7pA cutoff within <0.1pA. Therefore, our data support the idea that

excitatory synapses scale up in response to hibernation, but the multiplicative

organization appears to be controlled by mechanisms associated with the winter state

rather than inactivity *per se*.

What signals coordinate the multiplicative scaling in response to hibernation? An

interesting finding of this study was that blocking L-type Ca^{2+} channels *following*

hibernation disrupted multiplicative scaling across the mEPSC distribution without

influencing the average synaptic compensation. These results indicate that Ca^{2+}

feedback is a signal for maintenance of the scaled synaptic amplitude distribution after

hibernation. Given that we observed evidence for multiplicative scaling after hibernation

in the presence of TTX, the sum of our data suggests that the maintenance of synaptic

compensation in the "scaled state" is, therefore, Ca^{2+}-dependent but not activity-

dependent. As intracellular calcium transients tend to follow patterned activity, these

results may appear contradictory. However, spontaneous neurotransmitter release in

silent neurons can activate L-type channels at subthreshold voltages to induce neuronal

plasticity [45,46], suggesting a spike-independent role for Ca^{2+} to maintain the scaling

organization. Results in the present study pertain to the maintenance of scaling soon

after hibernation; yet, as cold temperature increases membrane input resistance [47,48],

spontaneous neurotransmitter release may cause larger postsynaptic depolarizations

compared to warm animals, generating significant Ca^{2+} influx through L-type channels to

orchestrate scaling even during the winter. Altogether, our results seem to diverge from

classic models of synaptic scaling, whereby a multiplicative scaling factor drives up

synaptic strength during inactivity. Instead, our data suggest that inactivity upregulates

synaptic strength through mechanisms that do not involve multiplicative scaling (Fig.

6A), and a separate regulatory system involving L-type Ca^{2+} channels is engaged by the

"winter state" to organize the balance of the mEPSC distribution (Fig. 6D-E). Given that

we can effectively disrupt the scaling with nimodipine, but leave overall compensation

intact following hibernation, provides evidence to this assertion. The mechanisms that

control these parallel signaling pathways for compensation and the scaling organization

warrant further investigation and may be relevant to complex physiological perturbations

in other systems.

Synaptic compensation and regulation of respiratory motor output

Our previous study identified that upregulation of synaptic strength acts to

maintain motor output associated with breathing following hibernation [15]. In this study

we had the unique opportunity to study this network in states where motoneurons had

compensated with (hibernation) and without (hibernation+nimodipine) scaled synapses.

Results here argue for the role of precise multiplicative scaling in the regulation of motor

function. We draw this conclusion because population motor output for breathing was

reduced when synaptic strength of motoneurons was upregulated but not correctly

scaled after hibernation (Fig. 7). We recognize the challenge of attributing changes in

network-level properties to subtle shifts in the distribution of synaptic weights on

motoneurons, and we made several attempts to address other variables that could

explain these results. First, rhythmic burst frequency of the network was the same

across all groups *ex vivo*; thus, overall activity differences and altered rhythm generator

function is unlikely to explain shorter motor bursts. Second, several intrinsic membrane

properties were not different across scaled and "unscaled" conditions (Fig. 7D-F),

suggesting that changes in excitability did not weaken the motor outflow. Third,

exposing warm-acclimated control preparations to nimodipine before hibernation did not

affect the amplitude distribution of mEPSCs, motor burst width, or any other obvious

electrophysiological property, arguing against other actions of nimodipine on different

parts of the network (Fig. 6-7). Finally, acute exposure to nimodipine did not affect burst

duration in any group (Fig. 7C), suggesting that slow disruption of Ca^{2+} feedback after

hibernation alters synaptic organization that shapes burst duration rather than acute

effects of calcium channel blockade on network excitability. All of this said, we cannot

state with certainty if narrower respiratory bursts after hibernation in the presence of

nimodipine reflect improperly scaled synapses on motoneurons (Fig. 6D-E) or other

actions of nimodipine on premotor neurons that regulate motor outflow in a hibernation-

dependent manner. Nevertheless, these results, in concert with our previous findings [15],

suggest that synaptic scaling is important for regulating motor performance shortly after

inactivity in hibernation. Future work must aim to understand how a failure to accurately

scale motoneurons synapses translates to altered respiratory-driven firing responses at

the single neuron level.

Conclusion

Compensation *via* postsynaptic AMPA receptor currents is a well-documented

response to reduced activity across diverse neurons and systems. Proper synaptic

compensation has clear adaptive value following hibernation in the frog, as it ensures

adequate motor drive for lung ventilation (15; present study Fig. 5). More broadly, our

results suggest that homeostatic synaptic plasticity could represent a more complicated

regulation scheme when studied in the context of living animals in response to natural

perturbations. Synaptic compensation is often viewed as a conceptually simple

feedback process, whereby neurons sense their activity state and elicit appropriate

compensation in attempt to regulate firing or network output. In contrast, our data

suggest that different aspects of the same compensation response, for example the

mean synaptic drive and the scaled nature of the synaptic amplitude distribution, may

be controlled by different variables. Along these lines, many neurological disorders are

thought to involve a failure of homeostatic synaptic and intrinsic mechanisms [9], all of

which occur in complex organisms with diverse types of environmental feedback onto

the nervous system. Our data suggest confronting how the environment shapes

compensation is necessary to understand the expression and control of homeostatic

compensation. Learning from animals that use these mechanisms to survive in their

natural environments, therefore, has the potential to provide new insight into how

neuronal function in regulated in both health and disease states.

Materials and Methods

Animals

Bullfrogs of either sex weighing approximately 100 g were purchased from Rana Ranch (Twin Falls, ID, USA). Control frogs were maintained at 20 ºC in plastic tubs and were provided aerated water with access to wet and dry areas. Frogs were fed pellets provided by Rana Ranch once per week. Hibernation conditions were simulated in temperature controlled incubators. The temperature of water was lowered from 20ºC to 2ºC over the course of a week, dropping about 2-3 degrees per day. On the seventh day, a plastic screen was placed at the top of the water to block to prevent access to the surface. Animals do not eat at cold temperatures; therefore, food was also withheld while frogs remained at 2°C. All frogs in each condition were kept on a 12 hr:12 hr light: dark cycle. Experiments were only performed during the light cycle.

Dissection and Tissue Preparation

Brainstem-spinal cord dissections were performed as previously described [15]. Briefly, frogs were deeply anesthetized with approximately 1 ml of isoflurane per liter and rapidly decapitated. The head was immersed in chilled bullfrog artificial cerebrospinal fluid (aCSF) and the brainstem-spinal cord was removed, keeping the motor rootlets of the vagus intact (aCSF; concentrations in [mM]: 104 NaCl, 4 KCl, 1.4 $MgCl_2$, 7.5 glucose, 40 $NaHCO_3$, 2.5 $CaCl_2$ and 1 NaH_2PO_4, and gased with 98.5% O_2, 1.3% CO_2; pH = 7.85). Following the dissection, the brain was superfused with aCSF with a flow rate of ~6 mL per minute.

To selectively label laryngeal branch motoneurons that innervate the glottal

dilator, the 4th branch of the vagus nerve was isolated and then backfilled

tetramethylrhodamine dextran 3000 MW (Life Technologies Corporation, Eugene, OR

USA). Approximately 1 uL of 10% dextran was loaded into the tip of a glass pipette that

fit snuggly around the cut 4th root of the vagus, as this nerve contains axons that mainly

innervate the glottal dilator see references provided in [15]. Backfills lasted two hours and

produced robustly labeled neurons for electrophysiological recording. Following the dye

loading process, tissue was sliced with a vibratome at 300 µM. Sections approximately

300-600 µm rostral to the obex were used for electrophysiological recording. Tissue

slices were left to recover for an hour before electrophysiological recording.

Whole-cell patch clamp recordings

Slices containing labeled motoneurons were transferred to the recording

chamber, stabilized with a nylon grid, and exposed to aCSF at a flow rate of ~1-2 mL

per min supplied via a gravity-fed system. The region containing labeled neurons was

imaged at 4x, and then individual neurons were located within the slice at 60x

(excitation, 540 nm: emission, 605 nm). Imaging was performed with a Hamamatsu

ORCA Flash 4.0LT sCMOS Camera (Hamamatsu Photonics, Japan). Identified neurons

were approached with the whole-cell patch pipette controlled by a Sutter Instruments

micromanipulator system, which was equipped with a MP-285 micromanipulator and a

MPC-200 controller (Sutter Instruments, CA, USA). A P87 horizontal pipette puller

(Sutter Instruments, Novato, CA, USA) was used to pull patch pipette, which had

resistances of 2-3 MΩ when filled with solution containing the follow composition (in

mM): 110 K-gluconate, 2 $MgCl_2$, 10 HEPES, 1 Na_2-ATP, 0.1 Na_2-GTP, 2.5 EGTA. The

cell of interest was approached with gentle positive pressure on the pipette. When an

indentation could be seen on the cell in the bright field with IR-DIC optics, positive

pressure was removed and a >1 GΩ was formed by applying slight negative pressure

through the pipette by mouth. Whole-cell access was obtained by rapid negative

pressure applied by mouth to rupture the GΩ seal. For measurements of excitatory

postsynaptic currents, cells were voltage clamped at -80 mV to record synaptic currents

mediated by AMPA receptors [15]. When spontaneous excitatory synaptic currents

(sEPSCs) were recorded in aCSF, no TTX was included during the recording. When

miniature excitatory postsynaptic currents (mESPCs) were recorded, 500 nM TTX was

added the aCSF. A +10 mV step was periodically performed to estimate series

resistance (peak of the transient current) and input resistance (steady-state current)

using Ohm's law. Measurements of membrane potential were made in current clamp,

and firing rate was assessed at 3X rheobase in some current clamp prior to the

application of TTX to measure mEPSCs. We often recorded more than one neuron per

slice; however, TTX does not have a fast washout time in this preparation; therefore,

firing properties in current clamp were not assessed for every cell. Previous work in

these neurons suggests that firing response to current injection does not change in

these neurons following hibernation [15], and our results in this study, despite a smaller

sample size, support this previous finding.

Experimental Treatments

Three main series of experiments were performed in this study: (1) those that varied the

duration of cold acclimation and then assessed synaptic function immediately after

removing the animal from the hibernation environment (Supplemental Fig. 1), (2) those

that used either warm-acclimated control animals or 1 month hibernation animals and

then exposed these brainstem preparations to TTX for 16 hrs in vitro (Fig 1-2

Supplemental Fig. 2-4), and (3) those that used either warm-acclimated control animals

or 1 month hibernation animals and exposed each to 10 µM nimodipine for 16 hrs (Fig

3-4). For series 1 experiments control frogs (20°C) or cold-acclimated (2°C for 2 or 4

weeks) were used, and then synaptic function was assessed after animals were

removed from these environments to roughly determine the time course of synaptic

compensation. In series two experiments our main goal was to compare to post-

hibernation synaptic function to inactivity with 100 nM TTX. In addition, we assessed

synaptic function in post-hibernation animals after silencing rhythmic motor output with

TTX. TTX silenced all preparations for the entire 16 hr period (data not shown). In series

three experiments, we exposed warm-acclimated control and post-hibernation rhythmic

preparations to 10 µM nimodipine for 16 hrs to block L-type Ca^{2+} channels in frog

respiratory motoneurons [30], and a separate time control group was run in parallel.

Following these treatments all synaptic recordings were performed in standard aCSF

with 500 nM TTX. Extracellular output from the cut vagus nerve via a bipolar suction

electrode connected to an A-M Systems Model 1700 amplifier (bandpass filtered

between 10-1000 Hz, 100X gain). Activity of this nerve is primarily generated by the 4th

branch of the vagus nerve (the same neurons we studied in slices), and provides the

main drive to the glottal dilator, which gates airflow into and out of the lung in anuran

amphibians [49].

Data Analysis

All data analysis was performed using the peak analysis module on LabChart 8

(ADInstruments, Sydney, Australia). sEPSC and mESPCs events were recorded for 30

seconds and averaged to estimate excitatory synaptic drive due to AMPAR

transmission. Cutoff amplitude was set at 7 pA. All analyses were checked by eye to

verify EPSCs were correctly identified. To infer synaptic scaling, we used the standard

rank-order method as originally described by Turrigiano et al. [4]. Briefly, a series of 50

mESPCs were chosen for each cell in the group and the entire distribution was ranked

from lowest to highest for control and treatment conditions. In some cases, control and

treatment conditions had different numbers of cells. In these situations, cells were

randomly removed for the scaling analysis by assigning a number to each cell and then

using a random number generator chose which cell(s) would be removed. This allowed

equal number of mESPCs to be ranked and scaled in each group. Ranked data were

plotted and then fit by a linear regression to obtain the slope of the line and intercept for

the entire population. ~5-10 mEPSC events in each group showed extreme values that

deviated from the line. These events may be caused by activation of subsaturated

AMPA receptors, as suggested previously (23), and were therefore removed for the

fitting of the ranked data. Each mESPC event in the treatment distribution was then

mathematically downscaled by the equation of the fit from the linear regression and

plotted as cumulative distributions. Input and series resistances were calculated by

making a 10 mV step, from -80 mV to -70 mV. Series resistance did not vary across

groups, and cells with series resistance of >25MΩ were discarded.

Statistics:

Data were analyzed and plotted using GraphPad Prism. Comparisons between

two groups were made using a two-tailed unpaired t test. When standard deviations

were different between groups, a Welch's test was applied. Analyses involving three or

more groups were performed using a one-way ANOVA and followed by Holm-Sidak's

multiple comparisons test. When comparing two cumulative distributions, the

Kolmogorov Smirnov test was run. When comparing two cumulative distributions, the

Kolmogorov Smirnov test was run. Significance was accepted at $p < 0.05$.

Acknowledgements: This research was funded in part by the Department of Defense

(W911NF2010275), the National Institutes of Health (1R01NS114514 and

1R15NS112920), and lab startup funds from UNC-Greensboro to J.M.S.

Competing Interests: The authors declare no competing interests.

References:

1 Marder, E. & Goaillard, J.-M. Variability, compensation and homeostasis in
neuron and network function. *Nature Reviews Neuroscience* **7**, 563-574 (2006).
2 Davis, G. W. Homeostatic signaling and the stabilization of neural function.
Neuron **80**, 718-728 (2013).
3 Turrigiano, G. Too many cooks? Intrinsic and synaptic homeostatic mechanisms
in cortical circuit refinement. *Annual review of neuroscience* **34**, 89-103 (2011).
4 Turrigiano, G. G., Leslie, K. R., Desai, N. S., Rutherford, L. C. & Nelson, S. B.
Activity-dependent scaling of quantal amplitude in neocortical neurons. *Nature*
391, 892-896 (1998).
5 Desai, N. S., Cudmore, R. H., Nelson, S. B. & Turrigiano, G. G. Critical periods
for experience-dependent synaptic scaling in visual cortex. *Nature neuroscience*
5, 783-789 (2002).
6 Pratt, K. G. & Aizenman, C. D. Homeostatic regulation of intrinsic excitability and
synaptic transmission in a developing visual circuit. *Journal of Neuroscience* **27**,
8268-8277 (2007).
7 Diering, G. H. *et al.* Homer1a drives homeostatic scaling-down of excitatory
synapses during sleep. *Science* **355**, 511-515 (2017).
8 Wu, C.-H., Ramos, R., Katz, D. B. & Turrigiano, G. G. Homeostatic synaptic
scaling establishes the specificity of an associative memory. *Current Biology*
(2021).
9 Dickman, D. & Wondolowski, J. Emerging links between homeostatic synaptic
plasticity and neurological disease. *Frontiers in cellular neuroscience* **7**, 223
(2013).
10 Styr, B. & Slutsky, I. Imbalance between firing homeostasis and synaptic
plasticity drives early-phase Alzheimer's disease. *Nature neuroscience*, 1 (2018).
11 Wolfart, J. & Laker, D. Homeostasis or channelopathy? Acquired cell type-
specific ion channel changes in temporal lobe epilepsy and their antiepileptic
potential. *Frontiers in physiology* **6**, 168 (2015).
12 Gainey, M. A., Hurvitz-Wolff, J. R., Lambo, M. E. & Turrigiano, G. G. Synaptic
scaling requires the GluR2 subunit of the AMPA receptor. *Journal of*
Neuroscience **29**, 6479-6489 (2009).
13 Gainey, M. A., Tatavarty, V., Nahmani, M., Lin, H. & Turrigiano, G. G. Activity-
dependent synaptic GRIP1 accumulation drives synaptic scaling up in response
to action potential blockade. *Proceedings of the National Academy of Sciences*
112, E3590-E3599 (2015).
14 Lambo, M. E. & Turrigiano, G. G. Synaptic and intrinsic homeostatic mechanisms
cooperate to increase L2/3 pyramidal neuron excitability during a late phase of
critical period plasticity. *Journal of Neuroscience* **33**, 8810-8819 (2013).
15 Santin, J. M., Vallejo, M. & Hartzler, L. K. Synaptic up-scaling preserves motor
circuit output after chronic, natural inactivity. *eLife* **6** (2017).
16 Santin, J. M. Motor inactivity in hibernating frogs: Linking plasticity that stabilizes
neuronal function to behavior in the natural environment. *Developmental*
neurobiology (2019).

17 Santin, J. M. & Hartzler, L. K. Activation of respiratory muscles does not occur
during cold-submergence in bullfrogs, Lithobates catesbeianus. *Journal of*
Experimental Biology **220**, 1181-1186 (2017).
18 Morales, R. D. & Hedrick, M. S. Temperature and pH/CO_2 modulate respiratory
activity in the isolated brainstem of the bullfrog *Rana catesbeiana*. *Comp*
Biochem Physiol- A Mol Int Physiol **132**, 477-487 (2002).
19 Vallejo, M., Santin, J. M. & Hartzler, L. K. Noradrenergic modulation determines
respiratory network activity during temperature changes in the in vitro brainstem
of bullfrogs. *Respiratory physiology & neurobiology* **258**, 25-31 (2018).
20 Santin, J. M. & Hartzler, L. K. Control of lung ventilation following overwintering
conditions in bullfrogs, Lithobates catesbeianus. *Journal of Experimental Biology*
In press, doi:10.1242/jeb.136259, doi:10.1242/jeb.136259 (2016).
21 Fong, M.-f., Newman, J. P., Potter, S. M. & Wenner, P. Upward synaptic scaling
is dependent on neurotransmission rather than spiking. *Nature communications* **6**
(2015).
22 Ibata, K., Sun, Q. & Turrigiano, G. G. Rapid synaptic scaling induced by changes
in postsynaptic firing. *Neuron* **57**, 819-826 (2008).
23 Mahamed, S., Strey, K. A., Mitchell, G. S. & Baker-Herman, T. L. Reduced
respiratory neural activity elicits phrenic motor facilitation. *Respiratory physiology*
& neurobiology **175**, 303-309 (2011).
24 Turrigiano, G. Homeostatic synaptic plasticity: local and global mechanisms for
stabilizing neuronal function. *Cold Spring Harbor perspectives in biology* **4**,
a005736 (2012).
25 Wang, H.-L., Zhang, Z., Hintze, M. & Chen, L. Decrease in calcium concentration
triggers neuronal retinoic acid synthesis during homeostatic synaptic plasticity.
Journal of Neuroscience **31**, 17764-17771 (2011).
26 Thiagarajan, T. C., Lindskog, M. & Tsien, R. W. Adaptation to synaptic inactivity
in hippocampal neurons. *Neuron* **47**, 725-737 (2005).
27 Tattersall, G. J. & Ultsch, G. R. Physiological ecology of aquatic overwintering in
ranid frogs. *Biological Reviews* **83**, 119-140 (2008).
28 McLean, H., Kimura, N., Kogo, N., Perry, S. F. & Remmers, J. E. Fictive
respiratory rhythm in the isolated brainstem of frogs. *Journal of Comparative*
Physiology A **176**, 703-713 (1995).
29 Whitlock, J. R., Heynen, A. J., Shuler, M. G. & Bear, M. F. Learning induces long-
term potentiation in the hippocampus. *science* **313**, 1093-1097 (2006).
30 Bueschke, N., Amaral-Silva, L. d., Hu, M. & Santin, J. M. Lactate ions induce
synaptic plasticity to enhance output from the central respiratory network. *The*
Journal of physiology.
31 Pacheco, A. T., Bottorff, J., Gao, Y. & Turrigiano, G. G. Sleep promotes
downward firing rate homeostasis. *Neuron* **109**, 530-544. e536 (2021).
32 Blackman, M. P., Djukic, B., Nelson, S. B. & Turrigiano, G. G. A critical and cell-
autonomous role for MeCP2 in synaptic scaling up. *Journal of Neuroscience* **32**,
13529-13536 (2012).
33 Greer, J., Smith, J. & Feldman, J. Role of excitatory amino acids in the
generation and transmission of respiratory drive in neonatal rat. *The Journal of*
physiology **437**, 727-749 (1991).

34 Johnson, S. M., Wilkerson, J. E., Wenninger, M. R., Henderson, D. R. & Mitchell,
G. S. Role of synaptic inhibition in turtle respiratory rhythm generation. *The*
Journal of physiology **544**, 253-265 (2002).
35 Santin, J. M. & Schulz, D. J. Membrane voltage is a direct feedback signal that
determines ion channel expression patterns in neurons. *Current Biology* **in press**
(2019).
36 Khorkova, O. & Golowasch, J. Neuromodulators, not activity, control coordinated
expression of ionic currents. *Journal of Neuroscience* **27**, 8709-8718 (2007).
37 Wierenga, C. J., Ibata, K. & Turrigiano, G. G. Postsynaptic expression of
homeostatic plasticity at neocortical synapses. *Journal of Neuroscience* **25**,
2895-2905 (2005).
38 Joseph, A. & Turrigiano, G. G. All for one but not one for all: Excitatory synaptic
scaling and intrinsic excitability are coregulated by CaMKIV, while inhibitory
synaptic scaling is under independent control. *Journal of Neuroscience*, 0618-
0617 (2017).
39 Echegoyen, J., Neu, A., Graber, K. D. & Soltesz, I. Homeostatic plasticity studied
using in vivo hippocampal activity-blockade: synaptic scaling, intrinsic plasticity
and age-dependence. *PloS one* **2**, e700 (2007).
40 Hanes, A. L. *et al.* Divergent synaptic scaling of miniature epscs following activity
blockade in dissociated neuronal cultures. *Journal of Neuroscience* **40**, 4090-
4102 (2020).
41 Kim, J., Tsien, R. W. & Alger, B. E. An improved test for detecting multiplicative
homeostatic synaptic scaling. *PloS one* **7**, e37364 (2012).
42 Louros, S. R., Hooks, B. M., Litvina, L., Carvalho, A. L. & Chen, C. A role for
stargazin in experience-dependent plasticity. *Cell reports* **7**, 1614-1625 (2014).
43 Wang, G., Zhong, J., Guttieres, D. & Man, H.-Y. Non-scaling regulation of AMPA
receptors in homeostatic synaptic plasticity. *Neuropharmacology* **158**, 107700
(2019).
44 Barnes, S. J. *et al.* Deprivation-induced homeostatic spine scaling in vivo is
localized to dendritic branches that have undergone recent spine loss. *Neuron*
96, 871-882. e875 (2017).
45 Li, Y., Gorassini, M. A. & Bennett, D. J. Role of persistent sodium and calcium
currents in motoneuron firing and spasticity in chronic spinal rats. *Journal of*
neurophysiology **91**, 767-783 (2004).
46 Li, B. *et al.* Neuronal Inactivity Co-opts LTP Machinery to Drive Potassium
Channel Splicing and Homeostatic Spike Widening. *Cell* (2020).
47 Santin, J. & Hartzler, L. Temperature influences CO_2/pH-sensitivity in locus
coeruleus neurons of the bullfrog, *Lithobates catesbeianus*. *FASEB J* **27** (2013).
48 Santin, J. & Hartzler, L. Activation state of the hyperpolarization-activated current
(Ih) modulates temperature-sensitivity of firing in locus coeruleus neurons from
bullfrogs. *American journal of physiology. Regulatory, integrative and*
comparative physiology **308**, R1045-R1061 (2015).
49 Sakakibara, Y. The pattern of respiratory nerve activity in the bullfrog. *The*
Japanese journal of physiology **34**, 269-282 (1983).

 Figures and Legends

Figure 1 sEPSC amplitude, but not frequency, increases by two weeks of cold-acclimation. (A) Schematic of the experimental process to label respiratory motoneurons. The 4th root of the vagus nerve was backfilled with fluorescent dextran, and neurons were identified in slices through fluorescence imaging microscopy. Animals were either warm-acclimated (control; 22°C; n=17), or cold-acclimated for two (n=20) or four weeks (n=19). (B) sEPSC amplitude (One-way ANOVA; p=0.0013) but not frequency (One-way ANOVA; (C) was increased by 2 weeks and remained elevated at four weeks. Circles represent cell-averaged mEPSCs for each neuron. Error bars are S.E.M. Horizontal line represents the mean. **p<0.01. Error bars are S.E.M.

Figure 2 Hibernation and TTX both induce increases in mEPSC amplitude to the same extent. (A) Example respiratory-related motor output from the cut vagus nerve root (CN X) highlight activity for the entire incubation period. TTX silenced activity in all preparations (B) Example mEPSC traces recorded in whole-cell voltage clamp in laryngeal motoneurons in time control, TTX, and following hibernation. (C) Mean data showing enhanced mEPSC amplitude following both hibernation and TTX exposure relative to time control (One-way ANOVA, Holm-Sidak Multiple Comparisons Test, control vs. TTX, p=0.004, control vs. post-hibernation, p=0.0139, TTX vs. post-hibernation, p=0.4405). Circles represent cell-averaged mEPSCs for each neuron. Error bars are S.E.M. Horizontal line represents the mean. Control (n=18 from 5 animals), TTX (n=19 from 5 animals), post-hibernation (n=17 from 4 animals). *p<0.05, **p<0.01.

Figure 3 Hibernation occludes the mEPSC amplitude increase by TTX. (A) shows
the mean mEPSC amplitude after hibernation and hibernation in the presence of 16 hr
TTX (unpaired t test) and (B) shows the cumulative distribution (Kolmogorov-Smirnov
Test). Circles represent cell-averaged mEPSCs for each neuron. Error bars are S.E.M.
Horizontal line represents the mean.

Figure 4 Membrane and other mEPSC properties are not affected following hibernation with or without activity in the post-hibernation period. (A-E) All variables were compared using a one-way ANOVA. No statistical difference was detected among treatments. Control (n=18), TTX (n=19), post-hibernation (n=17), post-hibernation+TTX (n=16). Box and whisker plots show the interquartile range. Whiskers extend to maximum and minimum values.

Figure 5 Block of L-type Ca²⁺ channels does not increase mEPSC amplitude. (A)
Example mEPSC traces recorded in whole-cell voltage clamp in time control (n=22 from
6 animals), 10 µM nimodipine (n=20 from 5 animals), and post-hibernation+nimodipine
(n=19 from 5 animals). (B') Cell-averaged mEPSC amplitudes and (B'') cumulative
distribution of mEPSC amplitudes. Nimodipine exposure does not increase mean
synaptic strength or alter the distribution of mEPSC amplitudes. After hibernation,
exposure to nimodipine still results in enhanced synaptic strength (time control vs post-
hibernation+nimodpine, p=0.0163; nimodipine vs. post-hibernation+nimodipine,
p=0.0367). (C) Post-hibernation (n=17 from 5 animals) and post-hibernation+
nimodipine (n=19) have similar group means (unpaired t test). Circles represent cell-
averaged mEPSCs for each neuron. Error bars are S.E.M. Horizontal line represents
the mean. * indicates p< 0.05

Figure 6 Hibernation, but not TTX, scales synapses, and multiplicative scaling is maintained by Ca^{2+} feedback . Cumulative distributions of mEPSC amplitudes for time control, treatment (TTX, post-hibernation, post-hibernation+TTX, post-hibernation+nimodipine), and mathematically downscaled treatment distributions. TTX does not lead to precise scaling of mEPSC amplitudes (A). Upregulated TTX mEPSCs were downscaled by the linear equation y=1.47x-2.21 (B) Hibernation and hibernation+TTX increases mEPSCs through multiplicative scaling (C). Upregulated hibernation mEPSCs were downscaled by the linear equation y=1.53x-3.58 and upregulated mEPSC after hibernation+TTX were downscaled by the equation y=1.61x-4.14. (D-E) After hibernation in the presence of nimodipine, mEPSC amplitudes were no longer increased according to multiplicative scaling. (D) uses time control as the control group and (E) uses the warm-acclimated nimodipine as the control. When using the time control as control, upregulated mEPSC amplitudes after hibernation+nimodipine were downscaled according to the equation, y=1.45x-3.76. When using nimodipine from warm acclimated animals as the control, upregulated mEPSCs were downscaled according to the equation y=1.52-4.75. All comparisons between control and treatment yielded p values <0.0001 (Kolmogorov-Smirnov Test.). P values shown in the figures compare control vs. scaled distributions (Kolmogorov-Smirnov Test).

**Figure 7 A failure to precisely scale motoneuron synaptic strength is associated
with reduced respiratory motor outflow.** (A) Simplified schematic of the brainstem
preparation, highlighting integrated respiratory-related motor output recorded from the
vagus nerve. The four groups shown are time control, post-hibernation (associated with
scaled compensation), nimodipine in warm-acclimated control (no compensation, no
scaling), and post-hibernation+nimodipine (unscaled compensation) for 13 hrs *in vitro*.
Traces are compressed or expanded and overlayed to highlight differences across
groups. (B) Nimodipine applied following hibernation resulted in narrower respiratory
bursts from the vagus nerve (One-way ANOVA with Holm-Sidak's Multiple Comparisons
Test; control vs. post-hibernation nimo, p=0.049, post-hibernation vs. posthibernation
nimo, p=0.007, nimo vs posthibernation nimo=0.041). (C) Nimodipine resulted in a slow,
reduction of burst width in the post-hibernation group, as acute exposure to nimodipine
did not affect burst amplitude in any group (Two-way ANOVA followed by Holm-Sidak's
multiple comparisons test). However, burst width decreases with longer exposure of
nimodipine only after hibernation (p<0.01 relative to all other groups). n= 6,6,6, and 5
brainstem preparations for control, post-hibernation, nimodipine, and post-hibernation
nimodipine, respectively. Intrinsic properties were not different across groups.
Membrane potential (C), input resistance (D), firing frequency (at 3X rheobase; E). For
(A), n= 6,6,6, and 5 brainstem preparations for control, post-hibernation, nimodipine,

and post-hibernation nimodipine, respectively. For (C &D), n= 20, 17, 20, and 19
neurons for control, post-hibernation, nimodipine, and post-hibernation+nimodpine,
respectively. For assessment of firing rate (E), n= 7,6,6,7 neurons were used in control,
post-hibernation, nimodipine, and post-hibernation+nimodpine, respectively. Box and
whisker plots show the interquartile range. Whiskers extend to maximum and minimum
values *p<0.05, **p<0.01.

34