1  **Autolysis affects the iron cargo of ferritins in neurons and glial cells at**
2  **different rates in the human brain**
3

4  **Sowmya Sunkara[1], Snježana Radulović[1]\*, Saška Lipovšek[1,2,3,4], Christoph Birkl[5],**
5  **Stefan Eggenreich[6], Anna Maria Birkl-Toeglhofer[7], Maximilian Schinagl[1,8,10], Daniel**
6  **Funk[1], Michael Stöger-Pollach[9], Johannes Haybaeck[7,10], Walter Goessler[11], Stefan**
7  **Ropele[6], Gerd Leitinger[1]**
8

[1]Gottfried Schatz Research Center, Division of Cell Biology, Histology and Embryology, Research Unit
Electron Microscopic Techniques, Medical University of Graz, Graz, Austria
[2]Faculty of Medicine, University of Maribor, Maribor, Slovenia
[3]Department of Biology, Faculty of Natural Sciences and Mathematics, University of Maribor, Maribor,
Slovenia
[4]Faculty of Chemistry and Chemical Engineering, University of Maribor, Maribor, Slovenia
[5]University Clinic for Neuroradiology, Medical University of Innsbruck, Innsbruck, Austria
[6]Neuroimaging Research Unit, Department of Neurology, Medical University of Graz, Austria
[7]Institute for Pathology, Neuropathology and Molecular Pathology, Medical University of Innsbruck,
Innsbruck, Austria
[8]Institute of Chemical Technologies and Analytics, Technische Universität Wien, Vienna, Austria
[9]University Service Centre for Transmission Electron Microscopy (USTEM), Technische Universität Wien,
Vienna, Austria
[10]Diagnostic & Research Center for Molecular BioMedicine, Institute of Pathology, Medical University of
Graz, Graz, Austria
[11]Institute for Chemistry, University of Graz, Graz
25

26 \*Correspondence: snjezana.radulovic@medunigraz.at

27
28

**Abstract**

Iron is known to accumulate in neurological disorders, so a careful balance of the
iron concentration is essential for healthy brain functioning. An imbalance in iron
homeostasis could arise due to the dysfunction of the proteins involved in iron
homeostasis. Here, we focus on ferritin – the primary iron storage protein of the

brain. Though it is known that glial cells and neurons differ in their concentration
of ferritin, the change in the number of iron-filled ferritin cores or their distribution
between different cell types during autolysis has not been revealed yet. Here, we
show the cellular and region-wide distribution of ferritin in the human brain using
state-of-the-art analytical electron microscopy. We validated the concentration of
iron-filled ferritin cores to the absolute iron concentration measured by
quantitative MRI and inductively coupled plasma mass spectrometry. We show
that ferritins lost iron from their cores with progressing autolysis whereas the
overall iron concentrations were unaffected. Though the highest concentration of
ferritins was found in glial cells, we found that as the total ferritin concentration
increased in a patient, ferritin accumulated more in neurons than in glial cells.
Collectively our findings point out the unique behaviour of neurons in storing iron
during autolysis and explain the differences between the absolute iron
concentrations and iron-filled ferritin in a cell-type-dependent fashion in the
human brain.

49
50  **Keywords**
51

56
57  **Significance statement**
58
59   Balance of the iron load of the brain is crucial to preventing neurodegenerative
60   disorders. Our study establishes a relation between autolysis, iron, and ferritin in
61   the human brain with emphasis on the role of different cells in ferritin storage. We

demonstrate that the iron load of ferritins does not correlate with mean iron concentrations during autolysis. Neurons retain more iron-loaded ferritin than glial cells with increasing ferritin count, which may make neurons more susceptible and exacerbate neuronal loss during iron overload. Neurons are also depleted of iron-loaded ferritin cores faster than glial cells during autolysis, demonstrating their unique role in iron storage. This paves the way to understanding the respective roles of neurons and glial cells in preventing or promoting neurodegeneration.

**Introduction**

Non-heme iron accumulates in the normal ageing brain until the fourth decade (Hallgren and Sourander, 1958). The rate of accumulation and the absolute iron concentration vary strongly across brain regions with the highest concentrations in the basal ganglia, including the putamen and the globus pallidus (GP) (Hallgren and Sourander, 1958; Krebs et al., 2014). The cause of iron accumulation is still unclear, in particular since much more iron is stored than metabolically relevant.

Basal levels of iron are essential in an array of functions in the brain: e.g. myelin synthesis, as cofactors for enzymatic reactions (Codazzi et al., 2015), and synaptic plasticity (Muñoz et al., 2011). Labile iron ions act as free radicals, induce reactive oxygen species (Halliwell, 2006) and can lead to neuroferritinopathy followed by neurodegeneration (Friedman et al., 2011). In a normal functioning brain, excess iron that is unused is stored in its primary storage protein, ferritin (Wang and Pantopoulos, 2011; (Finazzi and Arosio, 2014)), shielding the brain from the potential iron toxicity.

Ferritin is at the heart of the iron regulation. Ferritin acts as a buffer that stores and
releases iron upon cellular demand, which is crucial in the prevention of ferroptosis
(apoptosis induced by iron overload) (Jiang et al., 2021) and other detrimental
effects caused by free iron. Ferritin contains between 500 to 5000 ferric iron atoms
in its inner core (Harrison and Arosio, 1996; Iancu, 2011; Jian et al., 2016). The
size of the ferritin core in electron micrographs ranges from 5.3 to 11 nm (Iancu,
2011) depending on various factors such as iron load, contrasting agent, hydration,
or the focus of the micrograph. Despite the difference in the concentrations of total
iron and ferritin-bound iron between brain areas, the size of ferritin cores remains
largely uniform (Friedman et al., 2011).
98
Ferritin takes a unique position amongst the iron-binding proteins, owing to its size
and cargo capacity. Other iron-containing proteins like transferrin and
haemoglobin contain only 2 or 4 iron atoms respectively (Giometto et al., 1993;
Wagner et al., 2003; Marengo-Rowe, 2006), and hence are below the detection
threshold for energy-filtered transmission electron microscopy (EFTEM), our
method of choice for ferritin detection. Conversely, hemosiderin is composed of
conglomerates of clumped ferritin particles, denatured proteins, and lipids with a
size of only 1 - 2 μm (Wagner et al., 2003). In view of that, hemosiderin is easily
distinguished by size from ferritin but is rarely detected in the normal brain.

*Post-mortem* studies have the advantage that tissue is available for a variety of
measurements and procedures, including destructive ones. But autolysis sets off
immediately after death, causing a breakdown of cells and tissues, changing
structural components, and making it difficult to identify them (Sele et al., 2019),
reviewed in (Lewis et al., 2019).
113

The total iron concentration in freeze-dried tissue samples can be determined precisely by inductively coupled plasma mass spectrometry (ICP-MS), e.g. in (Krebs et al., 2014). However, this method is destructive. A non-invasive and thus non-destructive way to measure spatial iron concentrations is quantitative Magnetic Resonance Imaging (qMRI). qMRI allows mapping the local transverse relaxation rate $R_2^*$, which has been validated as a measure for iron content in brain tissue using ICP-MS (Langkammer et al., 2010). However, neither method provides information on the amount of iron stored in ferritins or its cellular distribution. We thus established a workflow using EFTEM for localising and quantifying the iron–filled cores of ferritins. We combined the three methodological approaches in our study to obtain complementary insights into the effects of autolysis on iron storage within ferritin cores.

For this, we investigated the concentrations of iron and ferritin cores filled with iron in the human brain samples *post-mortem* at the region and cellular level. We observed that autolysis influences the loss of iron-filled ferritin cores in a time-dependent manner. Moreover, our study reveals different rates of loss of iron–filled cores between neurons and glial cells during autolysis.

**Materials and methods**

**Human brain samples**

The samples used here are collected from six deceased human subjects who had to undergo routine brain autopsy at the Diagnostic and Research Institute of Pathology of the University Hospital Graz (approved by the Ethics Committee of the Medical University of Graz, votum number 28-549 ex 15/16). The subjects had

not suffered from a known neurological disorder. The sex of the subjects was not considered for this study. To ensure that the analyses were not confounded by age, the specimens were collected from patients with ages ranging from 61- 86. At this age saturation in iron accumulation has already been reached (Hallgren and Sourander, 1958). The age and *post-mortem* intervals (PMI) are given in Table 1. One-half of the brain was preserved for quantitative magnetic resonance imaging (qMRI) while four brain areas were dissected out of the other half and further divided into samples for EFTEM and ICP-MS. The brain regions selected included two regions with the lowest iron concentrations - the frontal gray matter (FGM), the superficial frontal white matter (FWM), and two regions with higher concentrations of iron - the putamen, and the globus pallidus (GP) (Bilgic et al., 2012; Ward et al., 2014; Ramos et al., 2014). Once the tissue was cleared for research purposes by the pathologists, samples were dissected out and either immersed in fixative for EM, or placed into a plastic tube, weighed, and frozen for ICP-MS. The brain hemisphere for qMRI remained unfixed.

Table 1. Patient information

| Patient code | Age [years] | PMI [h] |
|:---:|:---:|:---:|
| 1 | 73 | 06:30 |
| 2 | 72 | 9 |
| 3 | 61 | 16 |
| 4 | 69 | 18 |
| 5 | 86 | 20 |
| 6 | 81 | 24 |

**Ferritin Isolation**

Ferritin was isolated from human *post-mortem* brain samples in a series of
purification steps. In brief, the brain sample weighing approximately 10g
immersed in 10mM Tris-HCL, 150mM NaCl containing protease inhibitor was
homogenized using Turrax homogenizer IKA T -10. Followed by centrifugation at
10000g for 30-60 min at 4°C and sonication for 2 min at 4°C using a Branson
Sonifier 250. The sample was centrifuged again at 10000 g for 30-60 min at 4°C
retaining the supernatant for further purification. The supernatant was heated up to
70-75°C for 10 mins under constant stirring and immediately cooled. 75%
saturated ammonium sulfate was used to precipitate the proteins in the supernatant
overnight at 4 °C. The precipitated proteins were collected the following day by
centrifugation at 10000 g for 60 min at 4 °C. This was followed by size exclusion
chromatography and density gradient centrifugation. Isolated ferritin was loaded
onto a carbon-coated nickel grid and stained with 1% uranyl acetate to negatively
contrast the sample before visualization by EM.
175
176  **Electron Microscopy**
177
Following the autopsy, 6-10 sub-samples of approximate wet weight 0.1-0.2 g
(maximum dimensions 1x1x1 mm³) were cut out of the samples and immersed in
chemical fixative containing 2% formaldehyde, 2% glutaraldehyde in 0.1 M
sodium cacodylate buffer. After fixation, the samples were rinsed in the same
buffer, postfixed in 1% osmium tetroxide solution in the same buffer, dehydrated
in a graded series of alcohol, immersed in propylene oxide, and embedded in
TAAB embedding resin (TAAB, Aldermaston, UK). After curing for 3 days at 60
°C, thin sections were cut using a Leica UC6 or a Leica UC7 ultramicrotome at a
thickness of either 60 or 70 nm. The sections were contrasted with platinum blue
and lead citrate solutions. One block was randomly selected from each brain area
for imaging and elemental mapping.

EM was performed with a Thermo Fisher Tecnai G2 20 electron microscope
operated at 200 keV. An Ametek Gatan Quantum GIF energy filter with Ametek
Gatan Ultrascan 1000 XP camera was used for EFTEM, and a bottom-mounted
Ametek Gatan Ultrascan 1000 camera for bright field imaging. For elemental
mapping, three windows – Pre-edge 1, Pre-edge 2, and Post-edge – were made at
80000x magnification, a binning of 2, a slot width of 40 eV, and an exposure time
of 30s. The energy losses were: 718 eV for the iron L – post-edge window, and
633 eV and 673 eV for the two pre-edge windows. Drift correction was done
manually, and both an iron L – elemental map and an iron L – jump-ratio were
computed on Digital Micrograph (Gatan, Inc) using respectively the three-window
method and the two-window method. The iron cores inside the ferritins are visible
as bright spots using EFTEM.

An unbiased sampling protocol described earlier (; Wernitznig et al., 2019)
automatically selects 60 random locations on each sample determined using a
random number generator. The elemental maps were accompanied by overview
micrographs taken from the same locations using the bright field mode at 3500x
magnification.

**Ferritin quantification and cell type assessment**

Four independent researchers (SS, SnR, SL, and GL) each quantified the number
of iron-loaded ferritins and assessed the cell types. Ferritins were only counted if
they were clearly visible as a bright spot on both the iron L – jump-ratio and the
iron L – elemental maps and if at least three of the four researchers agreed on their
number. In case of disagreement, the images were analysed again simultaneously.
If the number of ferritins exceeded 20 in any map, the mean value of the counts

from four researchers was considered as the final count. Cell types were identified
on the bright-field images and only noted if three of the four researchers agreed.
For cell identification, the criteria described by (Nahirney and Tremblay, 2021)
and the Boston University web atlas https://www.bu.edu/agingbrain/ (download
2022-01-14) were used. A total of 1440 iron L – elemental maps and 1440 iron L
– jump ratios were generated and used for this study.
223
224 **Quantitative Magnetic Resonance Imaging**
225
MR imaging of the extracted brain hemisphere was performed using a 3.0 Tesla
scanner (MAGNETOM PRISMA, Siemens Healthineers, Erlangen, Germany). A
phased array head coil with 20 elements was used.  For acquiring $R_2^*$ relaxation
data, a two-dimensional radiofrequency-spoiled multi-echo gradient-echo
sequence with 6 equally spaced echoes was applied (repetition time = 1150 ms,
first echo time = 4.92 ms, echo spacing = 5.42 ms, flip angle = 15°; field of view
= 256 x 256 mm² ; in-plane resolution = 0.75 x 0.75 mm$^2$ ; 2.4-mm-thick sections
covering the entire hemisphere). Generalized auto-calibrating partially parallel
acquisition (GRAPPA) was performed with an acceleration factor of 2.
235
236 To reduce phase dispersion effects and to increase B0 homogeneity, second-order
237 shimming was applied. $R_2^*$ maps were calculated with the relaxometry
238 toolbox(2016).
239
**Inductively Coupled Plasma Mass spectrometry**
All solutions were prepared with ultrapure water (18.2 MΩ*cm, Merck Millipore,
Bedford, USA). Nitric acid (>65 % p.a.) was further purified via sub-boiling. For

quantification, single elements standards (ROTI®Star, Carl Roth GmbH + Co.
KG, Karlsruhe, Germany) were used.

Brain tissue was freeze-dried to constant mass with a Gamma 1-16 LSC freeze-
dryer (Martin Christ GmbH, Osterode am Harz, Germany) and the wet/dry mass
ratio was determined. Freeze-dried samples were weighed to 0.1 mg into the 12
mL quartz vessels of a microwave-heated autoclave, UltraCLAVE IV (EMLS,
Leutkirch, Germany). After the addition of 3 mL subboiled nitric acid and 2 mL of
water the quartz vessels were placed in the 40 positions rack of the autoclave and
the autoclave was pressurized with Argon to a pressure of $4*10^6$ Pa. The following
microwave heating program was applied: from room temperature to 80°C in 5
minutes, from 80°C to 150°C in 15 minutes, from 150°C to 250°C in 20 minutes,
and 250°C for 30 minutes. After cooling to 80°C, the pressure was released and
the samples were transferred to 50 mL polypropylene tubes (CELLSTAR®, blue
screw cap, Greiner Bio-One GmbH, Kremsmuenster, Austria) and filled to the
mark.

For the quantification of iron (Fecalibration curves were prepared from 0.0100 to
10.0 (Mg, Ca, Fe). Germanium (Ge) at a concentration of 200 µg/L was added as
an internal standard online before the nebulizer. The samples were pumped
through a peristaltic pump tubing with an i.d. of 1.05 mm and the internal standard
with a tubing of 0.19 mm. The elemental iron was quantified with an inductively
coupled plasma mass spectrometer (ICP-MS, Agilent 7700x, Agilent
Technologies, Waldbronn, Germany) at a mass-to-charge ratio of Fe m/z = 56. All
isotopes were measured in the collision mode using He at a flow rate of 5.0 ml/min.
The determined concentrations were calculated for fresh tissue.

The accuracy of the determined concentrations was confirmed with the certified
material bovine muscle (RM 8414; NIST; now BOVM-1 NRCC).

**Statistical Analysis**

Statistical analysis was performed using GraphPad Prism 9.2.0. One-way ANOVA
with a Tukey's multiple comparisons test was used for statistical analysis with * $p$
$< 0.05$; *** $p < 0.001$; **** $p < 0.0001$ for figures 2,3,5 and 8. For all comparison
plots of data from three methodological approaches in figures 6, 7, and 9, a simple
linear correlation analysis was performed.

**Results**

**EFTEM elemental mapping allows visualizing the ferritins' cores in the**
**human brain**

Ferritins consist of a shell and an iron-filled core, and the core is visible as an
electron-dense body in negative contrast electron micrographs of isolated ferritins
(Fig. 1B). The same core is visible as a bright spot in iron L – elemental maps (Fig.
1C).

To visualize the ferritin cores in electron micrographs, we generated iron L –
elemental maps and iron L – jump-ratios of brain samples for four different brain
areas in each of six different deceased subjects. The maps consistently exhibited
ferritins as bright spots of 5-10 nm in diameter in all four examined brain regions
(Fig. 2A-D). The size of the spots corresponded to the published sizes of ferritin
cores (Iancu, 2011), indicating that iron-filled ferritin cores could be visualized
with EFTEM.

**Ferritin cores were clustered in distinct cells**

We next counted the number of ferritin cores in samples taken from FGM, FWM,
putamen, and GP of each patient. The counts were made within 60 micrographs of

556 x 556 nm² dimensions and showed that the ferritins were not evenly distributed but clustered in certain micrographs (Fig. 3). We observed that the majority of ferritin accumulated as clusters as shown in figure 1, inside the cells. At a closer examination, it became apparent that certain cells accumulated ferritins, whereas others, often neighbouring cells, did not accumulate ferritins (Fig. 2). None of the ferritins appeared to be extracellular. We were able to assess the cell type (neuron, oligodendrocyte, astrocyte, or other unidentified glial cells) in 2,195 (62 %) of the 3,537 ferritin cores that we had identified in five out of six samples. 471 additional ferritin cores were identified in the sixth sample with a *post-mortem* interval (PMI) of 24 h, but in this sample, autolysis had progressed considerably, preventing us from faithfully identifying cell types. Accordingly, this sample was excluded from cell type identification experiments.

**Ferritins accumulate in the putamen and globus pallidus**

We next calculated the number of ferritins per μm³ from the micrographs' 3D dimensions. Plotting ferritin concentration in each brain region for each patient revealed that either the putamen, GP, or both, had a significantly higher concentration of ferritins than either the FGM or FWM in five out of six patients (Fig. 4). Further, between the putamen and GP (4 out of 6 patients) significant differences were found (Fig. 4). The GP had a significantly higher concentration compared to the former in three out of six patients and the putamen had a significantly higher ferritin concentration than the GP in one out of six patients (Fig. 4).

**qMRI and ICP-MS data are on par with the EFTEM data**

The ferritin concentrations counted from elemental maps were validated with the mean iron concentrations. The latter was determined by both mass spectrometry in adjacent tissue samples and by the proton transverse relaxation rate $R_2^*$ – a qMRI

measure proportional to the iron concentration – measured in the contralateral
hemisphere (Fig. 5). Each of the three methodological approaches provided higher
mean values (of all the six samples) in the putamen and GP than in FGM and FWM
(Fig. 6A–C)). qMRI revealed a significant $R_2^*$ increase in putamen and GP
compared to FGM and FWM (Fig. 6B). Significant differences were found in the
iron concentrations between the putamen and FGM using mass spectrometry (Fig.
6C).

To compare the three different methodological approaches, the mean values of the
measured brain regions from all six patients were correlated with each other. There
was a high correlation between qMRI and ICP-MS results, each measuring the
absolute iron concentration. The higher the $R_2^*$ value, the higher the iron
concentration (Fig. 7A, slope differs significantly from 0). However, we observed
that the mean ferritin count obtained using EFTEM does not correlate with the iron
concentration obtained with either qMRI (Fig. 7B) or ICP-MS (Fig. 7C). This
apparent discrepancy prompted us to check whether the autolysis had an influence
on the ferritin count or its cellular distribution.

**Autolysis enhances the unloading of the iron cargo from ferritin cores**

We next verified whether putative autolysis that progresses with increasing PMI
influenced the iron concentration, the storage of iron in ferritins' core, or the
cellular distribution of the ferritins.

Our analysis showed that the mean ferritin count was inversely correlated to the
PMI (Fig. 8A). In contrast, neither the $R_2^*$ value (Fig. 8B) nor the ICP-MS
determined iron concentration correlated with the PMI (Fig. 8C). In other words,
ferritin loses iron from its core with progressing autolysis while the absolute iron
concentration remains stable in each patient. This indicates that the longer the *post-*

*mortem* time, the more the ongoing autolysis progresses causing the unloading of
iron from its storage protein.

**Cell type identification reveals that ferritins are highly concentrated in glial**
**cells, especially oligodendrocytes**

The counted ferritin was assigned to different cell types by four independent
researchers. The cellular ferritin distribution showed that most of the 2195 ferritin
cores that could be assigned to a specific cell type were found within glial cells.
Oligodendrocytes represented the major cell type within this fraction (59 %). 17
% were found in neurons, 8% in astrocytes, and 1.7% in microglial cells (Fig. 9).

**Ferritin retention in neurons varies with both autolysis and the total ferritin**
**count**

Next, we studied the influence of autolysis on the iron load within ferritins in
neurons. We found that the percentage of iron-filled ferritin cores found using
EFTEM in neurons was inversely proportional to the PMI (Fig. 10A), indicating
that unloading of ferritins during autolysis is faster in neurons than in glial cells
(Fig. 10B).

To evaluate whether the cell-specific accumulation of ferritin is dependent on the
total ferritin amount, we plotted the percentage of ferritin found in neurons against
the total ferritin count in each sample. We observed that the percentage of ferritins
stored in neurons was proportional to the total ferritin count. In other words, when
the ferritin concentration increased in a patient, the probability of ferritins
accumulating in neurons increased. This result indicates that, although the majority

of ferritins was situated in glial cells, total ferritin load scaled with the proportion
of ferritins found in neurons (Fig. 10C) and vice-versa for glial cells (Fig. 10D).
370

**Discussion**

Our studies implemented analytical electron microscopy to determine the
concentration of ferritin cores filled with iron and used ICP-MS and $R_2^*$ relaxation
rate from qMRI to determine the absolute iron concentration in the human brain.
Samples from four brain regions of six deceased subjects were examined. The
subjects had not suffered from a known neurological disorder and the samples were
taken during routine autopsies.

Due to the electron density of the ferritins' iron cores, they are directly visible in
classical electron micrographs (reviewed in (Iancu, 2011)), but they can become
obscured by the contrasting agent in routine contrasted sections (Iancu, 2011).
Accordingly, we chose an energy filter to create iron maps for trustworthy ferritin
core identification. The size (Iancu, 2011) and detectability of these clusters
indicate that they correspond to the iron-loaded cores of ferritins. After identifying
the ferritin cores, we compared the elemental maps with a classical electron
micrograph of the same area for cell-type identification and ferritin localization.
We were able to demonstrate that the putamen and the GP store more ferritin iron
clusters than the FGM and the FWM, confirming a trend also seen in data obtained
with both qMRI and ICP-MS.

The EFTEM method is based on visualizing iron clusters inside ferritin using
analytical electron microscopy. Ferritins are known to vary in their iron load
(Harrison and Arosio, 1996; Iancu, 2011; Jian et al., 2016) and those ferritin
proteins that contain a low amount of iron remain undetectable using EFTEM.
Instead of using a fixed detection threshold, this study relied on four independent

researchers agreeing on the ferritin count from both iron L – elemental maps and
iron L – jump-ratios. There is high variability in the data obtained using the
EFTEM method due to the small sampling size. Hence, the mean concentrations
of ferritin both from each patient and from each region were used for this study.

We were able to demonstrate that the putamen and GP had a higher mean
concentration of detectable (iron-loaded) ferritins than the FGM and FWM. The
basal ganglia as the storage site for iron have long been known (Hallgren and
Sourander, 1958) and have since been confirmed in several different studies that
used a variety of different methods (Maeda et al., 1997; Pfefferbaum et al., 2009;
Krebs et al., 2014; Ramos et al., 2014). To validate our findings, we measured the
iron concentration in adjacent pieces of tissue taken from the same patients with
ICP-MS, and measured the proton relaxation rate $R_2^*$, using qMRI in the
contralateral region of the same patients. Our study on iron concentration validates
and confirms earlier studies in that the basal ganglia have both higher iron
concentrations and a higher $R_2^*$ relaxation rate than the frontal gray and white
matter (Langkammer et al., 2010).

Our study found a high correlation between the $R_2^*$ relaxation rate (qMRI) and the
absolute iron concentration (ICP-MS), confirming an earlier, comparative study of
both methods [16, 25]. In contrast, the mean ferritin concentration of all the brain
areas did not correlate with the mean iron concentration, indicating that the number
of iron-loaded ferritins was not directly dependent on the total iron concentration
in each patient. It has to be borne in mind that the EFTEM method measures only
those ferritin cores that are filled with a minimum threshold amount of iron (cargo),
whereas both qMRI and ICMPS measure the total elemental iron load of the brain
sample. We thus investigated the factors influencing the iron cargo load of ferritins
and found that this load is highly dependent on the post-mortem interval: the longer

the PMI, the fewer the iron-loaded ferritin cores found in a sample. This indicates
that autolysis progresses very rapidly during the post-mortem time and leads to the
unloading of iron from the ferritins' core. Two factors may contribute to unloading:
either the ferritin shell shielding the iron core is damaged during autolysis causing
the leakage of its cargo, or the ferritin holoprotein is decayed dissipating the iron.
Unloading of ferritin cores explains why the absolute iron concentrations remained
independent of the post-mortem time, whereas the number of iron-loaded ferritin
cores decreased with increasing post-mortem time. We thus can only speculate as
to the cellular mechanisms that govern the dispersion or decaying of ferritin during
the post-mortem interval. Further studies will be necessary to detect possible
changes in the oxidation state during the unloading or decay of ferritins.

Our second aim was to reveal the role of different cell types in preserving ferritin
during autolysis. We were able to determine the cell type (neuron,
oligodendrocyte, astrocyte, or other glial cells) that contained most (86 %) of the
ferritins we had detected in five out of the six patients. We found that most of the
iron in the human brain is stored in oligodendrocytes, but iron is also found in
other or unidentifiable glial cells and neurons. Owing to the fact that
oligodendrocytes are the myelinating cells, it is apparent that more iron is stored
in oligodendrocytes since it is used as a co-factor in lipid biosynthesis (Connor
and Menzies, 1996; Todorich et al., 2009). Earlier studies support our results using
a variety of different techniques (Connor et al., 1990; Quintana et al., 2004;
Quintana et al., 2006; Meguro et al., 2008).

In examining the cell types that are involved in iron-filled ferritin loss during
autolysis, we found that the loss occurs at a higher rate in neurons than in glial
cells, as schematically depicted in Fig. 11. Interestingly, we found that the
probability of the ferritins being stored in neurons rather than glial cells scales

positively with total ferritin concentration in each patient. To our knowledge, this
is the first study that shows that the iron–filled ferritin increases faster in neurons
than in glial cells with increasing total ferritin concentration.

Neurons were shown to have a higher ferritin H isoform distribution compared
with glial cells(Han et al., 2002; Muñoz et al., 2011). Previous studies have shown
that iron mediates the upregulation of ferritin H isoform in neurons(Nahirney and
Tremblay, 2021). Thus, we speculate that iron overload might lead to increased
production of ferritin H in neurons. This could also be a potential explanation for
enhanced ferritin load increase in neurons when a higher concentration of iron is
present in the brain.

In summary, we found high involvement of neurons in the storage of ferritins.
Moreover, iron-loaded ferritins are lost post-mortem, in a yet unknown process,
and loss of iron from ferritin cores occurs at a higher rate in neurons than in glial
cells.

**Footnotes**

• This work was funded by the Austrian Science Foundation [FWF], grant P
29370, and approved by the Ethics Committee of the Medical University of
Graz, votum number  28-549 ex 15/16.
464
• We are grateful to Dr. Sigurd Lax, for advice and for contributing material,
Dr. Peter Simmons (Newcastle University, UK), for proofreading and
language corrections, Mariella Sele, for processing the samples and for
initially evaluating most of the samples, Elisabeth Bock, Elisabeth Pritz, Luca
Schmid, Susanne Sumerauer and Nina Schlögl for expert technical assistance.
We especially thank Manuel Hündler for ferritin isolation.

• Figures 1 and 11 are created with BioRender.com

• The authors declare no competing financial interests.

• Correspondence should be addressed to Snježana Radulović at
snjezana.radulovic@medunigraz.at

**Abbreviations**

EFTEM – energy-filtered transmission electron microscopy

EM – electron microscopy

FGM – frontal gray matter

FWM – frontal white matter

GP – globus pallidus

ICP-MS – inductively coupled plasma mass spectrometry

MR – magnetic resonance

PMI – *post-mortem* interval

qMRI – quantitative magnetic resonance imaging

**Author contributions** SS, SnR, and GL planned and performed the experiments

at the electron microscope, and computed elemental maps and jump ratios. SS,

SnR, GL, and SL analyzed and quantified the raw data from elemental maps. JH, AMTB provided and assessed the brain tissue. StR, SE, and CB performed all qMRI experiments and analysis. WG performed ICP-MS experiments and analysis. PS proofread the manuscript. GL designed the project and study plan. All authors revised, read, and approved the manuscript.

## References

Bilgic B, Pfefferbaum A, Rohlfing T, Sullivan EV, Adalsteinsson E (2012) MRI estimates of brain iron concentration in normal aging using quantitative susceptibility mapping. Neuroimage 59:2625–2635.

Codazzi F, Pelizzoni I, Zacchetti D, Grohovaz F (2015) Iron entry in neurons and astrocytes: a link with synaptic activity. Front Mol Neurosci 8:18.

Connor JR, Menzies SL (1996) Relationship of iron to oligondendrocytes and myelination. Glia 17:83–93.

Connor JR, Menzies SL, St Martin SM, Mufson EJ (1990) Cellular distribution of transferrin, ferritin, and iron in normal and aged human brains. J Neurosci Res 27:595–611.

(2016) ESMRMB 2016, 33rd Annual Scientific Meeting, Vienna, AT, September 29 - October 1: ePoster / Paper Poster / Clinical Review Poster / Software Exhibits. MAGMA 29 Suppl 1:401–475.

Finazzi D, Arosio P (2014) Biology of ferritin in mammals: an update on iron storage, oxidative damage and neurodegeneration. Arch Toxicol 88:1787–1802.

Friedman A, Arosio P, Finazzi D, Koziorowski D, Galazka-Friedman J (2011)
Ferritin as an important player in neurodegeneration. Parkinsonism Relat
Disord 17:423–430.
Giometto B, Gallo P, Tavolato B (1993) Transferrin Receptors in the Central
Nervous System. In: Methods in neurosciences (Conn PM, ed), pp 122–134.
Elsevier.
Hallgren B, Sourander P (1958) The effect of age on the non-haemin iron in the
human brain. J Neurochem 3:41–51.
Halliwell B (2006) Oxidative stress and neurodegeneration: where are we now? J
Neurochem 97:1634–1658.
Han J, Day JR, Connor JR, Beard JL (2002) H and L ferritin subunit mRNA
expression differs in brains of control and iron-deficient rats. J Nutr 132:2769–
2774.
Harrison PM, Arosio P (1996) The ferritins: molecular properties, iron storage
function and cellular regulation. Biochimica et Biophysica Acta (BBA) -
Bioenergetics 1275:161–203.
Iancu TC (2011) Ultrastructural aspects of iron storage, transport and metabolism.
J Neural Transm (Vienna) 118:329–335.
Jian N, Dowle M, Horniblow RD, Tselepis C, Palmer RE (2016) Morphology of
the ferritin iron core by aberration corrected scanning transmission electron
microscopy. Nanotechnology 27:46LT02.
Jiang X, Stockwell BR, Conrad M (2021) Ferroptosis: mechanisms, biology and
role in disease. Nat Rev Mol Cell Biol 22:266–282.
JoVE Video Dataset.
Krebs N, Langkammer C, Goessler W, Ropele S, Fazekas F, Yen K, Scheurer E
(2014) Assessment of trace elements in human brain using inductively coupled
plasma mass spectrometry. J Trace Elem Med Biol 28:1–7.

Langkammer C, Krebs N, Goessler W, Scheurer E, Ebner F, Yen K, Fazekas F,
Ropele S (2010) Quantitative MR imaging of brain iron: a postmortem
validation study. Radiology 257:455–462.
Lewis AJ, Genoud C, Pont M, van de Berg WD, Frank S, Stahlberg H,
Shahmoradian SH, Al-Amoudi A (2019) Imaging of post-mortem human brain
tissue using electron and X-ray microscopy. Curr Opin Struct Biol 58:138–
148.
Maeda H, Sato M, Yoshikawa A, Kimura M, Sonomura T, Terada M, Kishi K
(1997) Brain MR imaging in patients with hepatic cirrhosis: relationship
between high intensity signal in basal ganglia on T1-weighted images and
elemental concentrations in brain. Neuroradiology 39:546–550.
Marengo-Rowe AJ (2006) Structure-function relations of human hemoglobins.
Proc (Bayl Univ Med Cent) 19:239–245.
Meguro R, Asano Y, Odagiri S, Li C, Shoumura K (2008) Cellular and subcellular
localizations of nonheme ferric and ferrous iron in the rat brain: a light and
electron microscopic study by the perfusion-Perls and -Turnbull methods.
Arch Histol Cytol 71:205–222.
Muñoz P, Humeres A, Elgueta C, Kirkwood A, Hidalgo C, Núñez MT (2011) Iron
mediates N-methyl-D-aspartate receptor-dependent stimulation of calcium-
induced pathways and hippocampal synaptic plasticity. J Biol Chem
286:13382–13392.
Nahirney PC, Tremblay M-E (2021) Brain Ultrastructure: Putting the Pieces
Together. Front Cell Dev Biol 9:629503.
Pfefferbaum A, Adalsteinsson E, Rohlfing T, Sullivan EV (2009) MRI estimates
of brain iron concentration in normal aging: comparison of field-dependent
(FDRI) and phase (SWI) methods. Neuroimage 47:493–500.

Quintana C, Bellefqih S, Laval JY, Guerquin-Kern JL, Wu TD, Avila J, Ferrer I,
Arranz R, Patiño C (2006) Study of the localization of iron, ferritin, and
hemosiderin in Alzheimer's disease hippocampus by analytical microscopy at
the subcellular level. J Struct Biol 153:42–54.
Quintana C, Cowley JM, Marhic C (2004) Electron nanodiffraction and high-
resolution electron microscopy studies of the structure and composition of
physiological and pathological ferritin. J Struct Biol 147:166–178.
Ramos P, Santos A, Pinto NR, Mendes R, Magalhães T, Almeida A (2014) Iron
levels in the human brain: a post-mortem study of anatomical region
differences and age-related changes. J Trace Elem Med Biol 28:13–17.
Sele M, Wernitznig S, Lipovšek S, Radulović S, Haybaeck J, Birkl-Toeglhofer
AM, Wodlej C, Kleinegger F, Sygulla S, Leoni M, Ropele S, Leitinger G
(2019) Optimization of ultrastructural preservation of human brain for
transmission electron microscopy after long post-mortem intervals. Acta
Neuropathol Commun 7:144.
Todorich B, Pasquini JM, Garcia CI, Paez PM, Connor JR (2009)
Oligodendrocytes and myelination: the role of iron. Glia 57:467–478.
Wagner KR, Sharp FR, Ardizzone TD, Lu A, Clark JF (2003) Heme and iron
metabolism: role in cerebral hemorrhage. J Cereb Blood Flow Metab 23:629–
652.
Wang J, Pantopoulos K (2011) Regulation of cellular iron metabolism. Biochem
J 434:365–381.
Ward RJ, Zucca FA, Duyn JH, Crichton RR, Zecca L (2014) The role of iron in
brain ageing and neurodegenerative disorders. The Lancet Neurology
13:1045–1060.
Wernitznig S, Reichmann F, Sele M, Birkl C, Haybäck J, Kleinegger F, Birkl-
Töglhofer A, Krassnig S, Wodlej C, Holzer P, Kummer D, Bock E, Leitinger

G (2019) An Unbiased Approach of Sampling TEM Sections in Neuroscience.
J Vis Exp. doi:10.3791/58745.

**Table and figures**

Table 1. Patient information

Figure 1. Visualization of the iron-core of ferritin. **A,** Schematic diagram of the
molecular architecture of ferritin. **B**, Negative contrasted electron micrograph of
isolated ferritin from the human brain. The ferritin shell appears as white rings
shielding the electron-dense iron core (inset). **C,** Iron L – elemental map displaying
the iron cores as bright spots.

Figure 2. EFTEM mapping of iron in different regions in post-mortem human brain
sample. **A**, Bright-field electron micrographs of overview images of the FGM
region, the FWM region, the putamen region, and the GP region. **B**, Inverted post-
edge micrographs. **C**, iron L – jump-ratios. **D**, corresponding iron L - elemental
maps of the same region as B. Ferritin filled with iron core appear as bright spots.
Examples are marked by red arrows. Corresponding iron L – jump-ratios and iron
L - elemental maps (C, D) show ferritin present in clusters in the cells.

Figure 3. The ferritin concentration in different brain regions varies between
patients 1-6. **A**, FGM. **B**, FWM. **C**, Putamen. **D**, GP. Data are shown as bar graphs
with whiskers representing the mean $\pm$ SEM. Each dot represents the ferritin
concentration measured in one iron L - elemental map, ($n = 60$ for each patient in
each brain region). One-way ANOVA with Tukey's multiple comparisons test was
used.

Figure 4. The ferritin concentration is often, but not always, higher in the basal ganglia than in the frontal gray or frontal white matter. Graphs show the ferritin concentration in the FGM, FWM, putamen, and GP in **A**, patient 1, **B**, patient 2, **C**, patient 3, **D**, patient 4, **E**, patient 5, and **F**, patient 6. Data are shown as bar graphs with whiskers representing the mean $\pm$ SEM. Each dot represents the ferritin concentration measured in one iron L - elemental map, respectively (n = 60 for each brain region and each patient). * $p < 0.05$; ** $p < 0.01$; **** $p < 0.0001$, One-way ANOVA with a Tukey's multiple comparisons test.

Figure 5. MRI scan showing an axial T1-weighted image of **A**, one cerebral hemisphere, and **B**, the corresponding $R_2^*$ map. The bright areas in (B) are the putamen and the globus pallidus with higher $R_2^*$ rates as a consequence of age-related iron accumulation.

Figure 6. The differences in ferritin concentration determined by EFTEM between brain areas follow the same trend as the sample from the counter brain half in qMRI data and adjacent samples in ICP-MS data. **A**, Ferritin concentration in the FGM, FWM, putamen, and GP measured by EFTEM. **B**, Iron concentration in the FGM, FWM, putamen, and GP measured by ICP-MS. **C**, Proton transverse relaxation rate ($R_2^*$) in the FGM, FWM, putamen, and GP measured by qMRI. Data are shown as bar graphs with whiskers representing the mean $\pm$ SEM. Each dot represents the mean ferritin concentration measured in one patient, respectively (n = 6). * $p < 0.05$; ** $p < 0.01$; **** $p < 0.0001$, One-way ANOVA with a Tukey's multiple comparisons test.

Figure 7. Correlation between mean concentrations of ferritin and iron measured via EFTEM, qMRI, and ICP-MS respectively. **A**, Proton transverse relaxation rate (qMRI) plotted over the iron concentration (ICP-MS) for each patient. The mean proton transverse relaxation rate of each patient correlates significantly with the

mean iron concentration, $r^2 = 0.8560$, p = 0.00082. **B**, Mean ferritin concentration
(EFTEM) plotted over the proton transverse relaxation rate ($R_2$*) (qMRI). **C**, Mean
ferritin concentration (EFTEM) plotted over the iron concentration (ICP-MS). In
contrast to (A), the mean concentration of ferritin particles does not correlate with
proton relaxation rate (B), $r^2 = 0.0230$, p = 0.7742 nor with the iron concentration
(C), $r^2 = 0.0043$, p= 0.9015. Each dot represents the mean ferritin concentration or
the mean iron concentration measured in one patient, respectively (n = 6). All the
plots are based on mean values of the FGM, FWM, the putamen, and the GP. * p
< 0.05; ** p < 0.01; **** p < 0.0001, Linear correlation analysis.

Figure 8. Ferritins disintegrate and their concentration is reduced over post-mortem
time. **A**, Mean values of the ferritin concentration (EFTEM) in all the brain regions
plotted against PMI, $r^2 = 0.6421$, p = 0.0553. **B**, Mean values of the proton
transverse relaxation rate (qMRI) in all the brain regions plotted against PMI, $r^2 =$
0.1323, p = 0.4785. **C**, Mean values of the iron concentration (ICP-MS) in all the
brain regions plotted against PMI, $r^2 = 0.1215$, p = 0.4983. The ferritin
concentration decreases with PMI in regions of the brain (A) whereas the total iron
load in the four regions of the brain does not seem to be influenced by the PMI (B,
C). Each dot represents the mean ferritin concentration or the mean iron
concentration measured in one patient, respectively (n = 6). All the plots are based
on mean values of the FGM, FWM, the putamen, and the GP. * p < 0.05, Linear
correlation analysis.

Figure 9. Ferritin is predominately stored in glial cells (82%) and especially in
oligodendrocytes (59%); followed by 8% in astrocytes and 1.7% in microglial cells
of the identified glial cells. 17 % is stored in neurons. The graph shows those
ferritins in the FGM, FWM, putamen, and GP of patients 1-5 whose cell type could

be determined. One-way ANOVA with Tukey's multiple comparisons test was
used for statistical analysis.

Figure 10. The percentage of ferritin in neurons varies with post-mortem time and
with total ferritin concentration. **A**, Percentage of ferritins in neurons from patients
1 to 5 plotted against PMI. Mean values of the percentage of ferritin concentration
in neurons (EFTEM) reduce over time with increasing PMI, $r^2 = 0.8338$, $p =$
0.0303. **B**, Percentage of ferritins in glial cells from patients 1 to 5 plotted against
PMI. Mean values of the percentage of ferritin concentration in glial cells
(EFTEM) reduce over time with increasing PMI, $r^2 = 0.8338$, $p = 0.0303$. **C**, Mean
values of the percentage of ferritins found in neurons in all brain regions from
patients 1 to 5 were plotted against total ferritin count. The probability of ferritins
being found in neurons rises with the total ferritin concentration, $r^2 = 0.8101$, $p =$
0.0374. **D**, the probability of ferritins being found in glial cells decreases with the
total ferritin concentration, $r^2 = 0.8101$, $p = 0.0374$. Each dot represents the mean
percentage of ferritin found in neurons in all brain regions measured in patients 1-
5, respectively (n = 5). * $p < 0.05$, Linear correlation analysis.

Figure 11. The rate of loss of the iron-filled ferritin cores during autolysis is higher
in neurons than in glial cells.

**A** Ferritin shell    Iron core

Ferritin protein

**B** Ferritin shell

Isolated ferritin
electron micrograph

**C** Iron core

Iron L- elemental map

# Figure 2

# Figure 3

# Figure 4

A

## Patient 1

Ferritin count (number/μm³)

FGM  FWM  Putamen  GP

B

## Patient 2

Ferritin count (number/μm³)

FGM  FWM  Putamen  GP

C

## Patient 3

Ferritin count (number/μm³)

FGM  FWM  Putamen  GP

D

## Patient 4

Ferritin count (number/μm³)

FGM  FWM  Putamen  GP

E

## Patient 5

Ferritin count (number/μm³)

FGM  FWM  Putamen  GP

F

## Patient 6

Ferritin count (number/μm³)

FGM  FWM  Putamen  GP

# Figure 6

# Figure 7

# Figure 8

# Figure 9

# Figure 10

Ferritin

Neuron

Progressing
autolysis

Ferritin

Glial cell

Progressing
autolysis