1 **Transcription factor FOXM1 specifies the loading of chromatin DNA to extracellular**

2 **vesicles**

Yunsheng Zhang[1,2], Nana Ding[1], Yizhen Li[1], Xiaoyun Zhao[1], Shiping Yang[1], Ping Fu[1], Yousong Peng[1],

Haojie Cheng[1], Min Ouyang[1], Ruiping Wang[1], Yu Wang[1], Hanyun Liu[1], Yan Chen[1], Xiaoqin Huang[1], Li

Yu[1], and Yongjun Tan[1*]

[1] State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Biology, Hunan Engineering

Research Center for Anticancer Targeted Protein Pharmaceuticals, Hunan University, Changsha, Hunan

410082, PR China.

[2] The Second Affiliated Hospital, University of South China, Hengyang, Hunan 421001, PR China.

*** Correspondence to:*** Dr. Yongjun Tan, E-mail: yjtan@hnu.edu.cn

Abstract

Extracellular vesicle DNAs (evDNAs) possess the important diagnostic value for multiple diseases and

play roles for horizontally transferring genetic materials among cells. In this study, we have found that

transcription factor FOXM1 can mediate the loading of certain chromatin genes or DNA fragments

(named FOXM1-chDNAs) to extracellular vesicles (EVs). FOXM1 interacts with LC3 in nucleus and

FOXM1-chDNAs (such as *DUX4* gene and Telomere DNA) are specified by FOXM1 and translocated

to cytoplasm. These DNAs are released to EVs through the process of an LC3-involved autophagosome-

multivesicular body (MVB) transport. The roles of FOXM1 on loading FOXM1-chDNAs to EVs are

further confirmed by DNA-FISH experiments, tracing the translocation of selected chromatin loci with

the TetO/TetR-GFP method, and PCR analysis of the DNA samples from MVBs and EVs. Furthermore,

disrupting the expression of FOXM1 or the process of autophagosome-MVB transport impairs the

loading of FOXM1-chDNAs to EVs. This discovery suggests that transcription factor FOXM1

contributes the constitution of evDNAs from nuclear chromatin, providing the first example to explain

how chromatin DNA fragments are specified and loaded to EVs. It also provide a foundation to further

explore the roles of evDNAs in biological processes such as the horizontal gene transfer.

Key words

Extracellular vesicle DNA; Transcription factor FOXM1; FOXM1-LC3 interaction; The process of

autophagosome-MVB transport; Extracellular vesicles

Introduction

Extracellular vehicles (EVs) are membranous vesicles released by a variety of cells into the extracellular microenvironment and contain DNAs, RNAs, and proteins acting as intercellular messengers for exchanging variable materials among cells[1,2]. Because extracellular vesicle DNAs (evDNAs) carry the portions of genetic materials from their parent cells under certain physiological or pathological conditions, extensive studies have focused on the diagnostic value of evDNAs for multiple diseases, while their biological functions are not explored as deeply as that of the RNAs or proteins of EVs[3,4]. The evDNAs have been proposed to play a role in so called the horizontal gene transfer (HGT)[5], which allows to exchange functional genes and DNA fragments among cells[6]. Increasing evidences support that EVs contain functional genes or chromosome DNA fragments as evDNAs and transfer them to recipient cells[7]. For example, LINE-1 retrotransposon as an evDNA of EVs is horizontally transferred among human cancer cells[8]. Another example is that Telomere DNA carried by the EVs from antigen-presenting cells (APCs) is horizontally transferred to primarily naïve T cells and central memory T cells[9]. Therefore, it is critical to understand how chromatin DNA fragments are specified and loaded to EVs, not only for improving the application of evDNAs in diagnosis but also for clarifying their functions in biological processes such as HGT.

Recently, chromatin DNA has been found to translocate to cytoplasm through the interaction between laminB1 and LC3 in nucleus during autophagy[10], providing a clue to explore the origin of evDNAs. LC3 is a key member of the Atg8 family proteins that are responsible for the cargo selection of autophagy[11]. Besides existing in cytoplasm, LC3 is present in nucleus and interacts with the chromatin-binding laminB1 (a nuclear lamina protein[12]). During autophagy, a part of regular LC3 (LC3-I) need be

conjugated with phosphatidyl ethanolamine (PE) to form LC3-PE (LC3-II), which is termed as the

lipidation of LC3 and allows the LC3 protein embedding in membranes and then loading cargos to

autophagosomes[13]. Although cytoplasmic autophagosomes fuse with endosomes or lysosomes to degrade

their components[14,15], accumulated evidences prove that a portion of endosomes fused with

autophagosomes can mature to form multivesicular bodies (MVBs, a special kind of late endosomes[16])

for the secretion of their cargos[1,17,18]. MVBs contain LC3-positive intraluminal vesicles (ILVs)[18], which

can be released to extracellular environment as EVs through the fusion of MVBs with cell membrane[2].

It's well known that LC3 specifies the loading of proteins and RNAs to EVs[18,19], confirming that variable

materials can be released by EVs through the process of an LC3-involved autophagosome-MVB

transport. However, because laminB1 possesses no sequence specificity for its DNA binding[20], it is hard

to figure out how certain chromatin DNA fragments are specified and loaded to EVs, if only basing on

the mechanism of laminB1-LC3-mediated chromatin DNA cytoplasmic translocation.

Here, we show a transcription factor-LC3-involved mechanism specifying certain chromatin genes

or DNA fragments to EVs during autophagy. FOXM1, a member of Forkhead Box transcription factor

family[21], participates in regulating cell proliferation[22], DNA damage repair[23], cell stemness[24], and

metastasis[25,26] through stimulating gene transcription in nucleus[27]. FOXM1 can specify its binding

regions on chromatin directly through its DNA-binding consensus sequence[28] or indirectly through

interacting with other transcription factors such as B-Myb, MuvB, and NFY[29]. In this study, we have

confirmed that FOXM1 interacts with LC3 in nucleus and mediates the loading of specific chromatin

DNA fragments (named FOXM1-chDNAs) to EVs during autophagy. The interaction between FOXM1

and LC3 is mediated by the LC3-interacting region motif of FOXM1 (317-320aa) and facilitated by the

lipidation of LC3, while the FOXM1-LC3 interaction does not disrupt the DNA binding ability of

FOXM1. FOXM1-chDNAs, including *DUX4* gene and Telomere DNA, etc., have been identified by

analyzing the sequencing data from FOXM1-specific ChIP-seq, LC3-specific ChIP-seq, and evDNA-

seq. The loading of specific FOXM1-chDNAs to EVs has been confirmed by DNA-FISH experiments,

tracing the translocation of selected chromatin loci with the TetO/TetR-GFP method, and PCR analysis

of the DNA samples from MVBs and EVs. We have confirmed that the expression of FOXM1 or the

process of LC3-involved autophagosome-MVB transport is essential for the loading of FOXM1-chDNAs

to EVs. Together, this study has provided the first example to explain how chromatin DNA fragments are

specified and loaded to EVs and a foundation to further explore the roles of evDNAs in biological

processes such as HGT.

Results

The interaction between FOXM1 and LC3

LC3 mediated the loading of variable materials to EVs through the process of an LC3-involved autophagosome-MVB transport[18,19]. Interestingly, we identified LC3 as a potential partner interacting with transcription factor FOXM1, which directly or indirectly bound to the specific regions on chromatin, from our mass spectrometry analysis of the FOXM1 interactome (data not shown), implicating that FOXM1 might participate in specifying and loading chromatin DNA fragments to EVs. To test this hypothesis, we first confirmed the interaction between FOXM1 and LC3 in nucleus of lung cancer A549 cells. The immunostaining of endogenous FOXM1 and LC3 showed that the two proteins were co-localized in nucleus with the Pearson's Correlation Coefficient value around 0.66 ± 0.08 (**Fig. 1a**). The co-immunoprecipitation (co-IP) experiments revealed that FOXM1 interacted with LC3, especially with the lipidated LC3-II in nucleus (**Fig. 1b and Extended Data Fig. 1a, b**). A bimolecular fluorescence complementation (BiFC) assay[30] further confirmed that the FOXM1-LC3 interaction happened mainly in nucleus (**Fig. 1c**). The co-IP of exogenous Flag-FOXM1 with GFP-LC3 WT or G120A mutant, which disrupted the LC3 lipidation on its G120 residue[10], showed that the deficiency of the LC3 lipidation decreased the FOXM1-LC3 interaction (**Extended Data Fig. 1c, d**), implicating that LC3 embedding in nuclear membranes facilitated its interaction with FOXM1.

Next, we identified the regions of FOXM1 and LC3 to mediate the FOXM1-LC3 interaction. We screened FOXM1 protein and found that FOXM1 DNA-binding domain (FOXM1-DBD) was required for its LC3 binding (**Fig. 1d**) and FOXM1-DBD interacted with LC3 in GST-LC3-pulldown assays (**Fig. 1e**). Moreover, the overexpression of GFP-FOXM1-DBD abolished the endogenous FOXM1-LC3

interaction (**Extended Data Fig. 1e, f**). It was known that so-called LC3-interacting region (LIR) motif

existed in LC3' substrate proteins[31,32]. We screened the protein sequence of FOXM1-DBD and identified

four LIR motifs (263-266, 273-276, 307-310 and 317-320), which were evolutionarily conserved among

FOXM1s of human, mouse, rat, and macaque (**Extended Data Fig. 1g**). We generated four FOXM1-

DBD mutants (Y263A, F273A, F307A, and Y317A, corresponding to the four LIR motifs respectively),

among which only the Y317A mutation on FOXM1-DBD abolished its interaction with LC3 in GST-

LC3-pulldown assays (**Fig. 1f**), suggesting that the LIR motif (317-320) in FOXM1-DBD mediated the

binding of FOXM1 to LC3. On the other hand, because the residues R10, R11, and F52 of LC3 mediated

the binding of LC3 to its protein partners[10,33], we tested whether the LC3 aa1-28 region (containing R10

and R11) or the LC3 aa29-128 region (containing F52) could interact with FOXM1 in GST-pulldown

assays. We found that only the aa29-128 region (containing F52) of LC3 interacted with FOXM1 (**Fig.**

1g), implicating that the residues F52 of LC3 mediated the binding of LC3 to FOXM1. Thus, we

generated an LC3 aa29-128 truncated F52A mutant and found that this mutant could not bind to FOXM1

in GST-pulldown assays (**Fig. 1h**). Rosetta Dock[34] was performed with the known structures of FOXM1-

DBD[35] and LC3[36] to simulate the FOXM1-LC3 interaction and predicted the formation of hydrogen

bonds between the LIR motif (317-320) of FOXM1 and the residue F52 of LC3 (**Fig. 1i**), further

supporting the conclusion of the interaction between FOXM1 and LC3 in biochemical experiments.

To test whether the binding of LC3 to FOXM1-DBD affected the DNA-binding ability of FOXM1,

we performed the electrophoretic mobility shift assays (EMSAs) with recombinant FOXM1 proteins and

a FAM-labeled DNA probe containing putative FOXM1 binding sites. We observed that the DNA probe

formed FOXM1-DNA complexes with FOXM1 alone, and created a supershift of LC3-FOXM1-DNA

complexes by adding recombinant LC3 proteins in the reactions (**Fig. 1j**), suggesting that the FOXM1-

LC3 interaction did not impair the DNA-binding ability of FOXM1 and the LC3-FOXM1-DNA complex

could be formed in cells.

FOXM1-chDNAs are loaded to extracellular vesicles during autophagy

From the results of LC3 or FOXM1 immuno-staining and DNA DAPI-staining of A549 cells, we

observed that DNAs, FOXM1, and LC3 co-localized in cytoplasmic vesicles (**Fig. 2a and Extended**

Data Fig. 2a), whose numbers were increased by starvation-induced autophagy (from 3% in control cells

to 15% in starvation-treated cells) (**Fig. 2b**), suggesting that the formation of LC3-FOXM1-DNA vesicles

normally happened in cytoplasm and were enhanced during autophagy. Similar phenomena were also

observed from multiple human cancer cell lines, such as Hela and HEK293T cells, or the cells treated

with different autophagy-inducing conditions, such as No-amino acid, No-serum, or Rapamycin

treatment (20 μM) (data not shown). The BiFC assays with exogenous VN173-FOXM1 and VC155-LC3

further confirmed that FOXM1 and LC3 co-existed in cytoplasmic DNA-containing vesicles (**Fig. 2c**).

Interestingly, the expression of RFP-FOXM1 ΔDBD, in which FOXM1-DNA binding domain (DBD)

was deleted, could not appear in cytoplasmic DNA-containing vesicles (**Extended Data Fig. 2b**),

confirming that the FOXM1 DNA-binding ability was required for the formation of cytoplasmic LC3-

FOXM1-DNA vesicles during autophagy. Next, we collected EVs (including Large EVs (lEVs) and

Small EVs (sEVs)) from starved A549 cells by serial differential ultracentrifugation[37] (**Extended Data**

Fig. 2c). We showed that both FOXM1 and LC3 existed in the samples of EVs by Western blotting (**Fig.**

2d). Because researchers usually focused on sEVs when studying EVs[38,39] and only sEVs from our

samples possessed EV marker TSG101[40] (**Fig. 2d**), we just collected sEVs as EV samples in following

experiments and characterized their size range around 80-250 nm through nanoparticle tracking analysis

and transmission electron microscopy (**Extended Data Fig. 2d and e**). The DNA samples from EVs

(evDNAs) were purified and analyzed by the vertical agarose gel electrophoresis (**Fig. 2e**), which showed

the size range of evDNAs around 2 to 15 kb (majorly around 12 kb), confirming that DNA, FOXM1,

and LC3 co-existed in the samples of EVs.

To identify the chromatin DNA fragments specified by FOXM1 (FOXM1-chDNAs) that were

finally loaded to EVs, we first performed chromatin immunoprecipitation sequencing (ChIP-seq)

analysis for FOXM1 or LC3 separately with starved A549 cells and obtained 103 specific loci on

chromatin bound by both FOXM1 and LC3 (**Table S**) from the overlapping of FOXM1-ChIP fragments

(n=8, 838) (GSE216672) and LC3-ChIP fragments (n=7, 951) (GSE216672). Then we performed

evDNA-sequencing analysis with the EVs of starved A549 cells and obtained 15, 544 evDNAs

(GSE216672). After analyzing the overlapped sequences of FOXM1-LC3 ChIP loci and evDNAs, we

identified 25 chromatin loci as FOXM1-chDNAs (**Table**), which could be annotated as functional genes

(n=8, including *DUX4* gene), telomere (n=1), centromere (n=8), and nonsense regions (n=8). We chose

three FOXM1-chDNAs (FOXM1-chDNA1 from a telomere locus, FOXM1-chDNA2 from a nonsense

region, and FOXM1-chDNA3 from *DUX4* locus) as examples to perform ChIP-qPCR and confirmed the

binding of FOXM1 and LC3 on these FOXM1-chDNAs at chromatin but not on the control region such

as β-actin promoter (**Extended Data Fig. 2f, g**).

To verify that FOXM1-chDNAs were translocated to cytoplasm and finally loaded to EVs, we first

purified cytoplasmic DNA (cytoDNA) particularly[41] and found the levels of the three FOXM1-chDNAs

were elevated in cytoplasm of starved A549 cells (**Fig. 2f**). A DNA-FISH probe (10 Kb) specific for the

FOXM1-chDNA1-adjacent region was generated and DNA-FISH experiments showed that FOXM1-

chDNA1 appeared in cytoplasm post starvation (**Fig. 2g**), confirming that FOXM1-chDNAs were

translocated to cytoplasm during autophagy. Then, we purified evDNAs from the EVs of starved A549

cells and found the levels of the three FOXM1-chDNAs were elevated in the EVs (**Fig. 2h**), confirming

that FOXM1-chDNAs were finally loaded to EVs during autophagy. In addition, we also treated the EVs

of starved A549 cells with the Plasmid-Safe ATP-dependent DNase (PS), which only digested linear

DNA[42], and observed the decreased levels of the three FOXM1-chDNAs from the PS-treated samples

(**Extended Data Fig. 2h**), identifying that FOXM1-chDNAs existed as linear DNA on outside the lumen

of EVs. This result was consistent with the previous finding that evDNAs mainly localized on the outside

the lumen of EVs[43]. Then, we treated the EVs of starved A549 cells with formaldehyde to cross-link their

DNA and proteins and the lysates of EVs were immune-precipitated with anti-FOXM1 or anti-LC3

antibodies for evDNA purification. The levels of the three FOXM1-chDNAs in the FOXM1 or LC3

specific IP-evDNA samples were significantly elevated (**Fig. 2i**), further confirming that FOXM1-

chDNAs were bound by FOXM1 or LC3 on EVs. We also collected FOXM1-specific EVs or LC3-

specific EVs from GFP-FOXM1 or GFP-LC3-overexpressed A549 cells through the direct

immunoaffinity capture (DIC) approach[1] with GFP-Trap and observed the elevated levels of the three

FOXM1-chDNAs in the DIC samples (**Fig. 2j**), suggesting that both exogenous GFP-FOXM1 and GFP-

LC3 facilitated the loading of FOXM1-chDNAs to EVs. Together, these data suggested that FOXM1 and

LC3 mediated the loading of FOXM1-chDNAs to EVs during autophagy.

The FOXM1-chDNAs knocked-in with the TetO array were loaded to EVs

To confirm the translocation of FOXM1-chDNAs to EVs, we knocked in a TetO array (96x) at the loci

of selected FOXM1-chDNAs by CRISPR/cas9 in A549 cells (**Extended Fig. 3a**). DNAs containing the

TetO array then were visualized directly by exogenously expressed TetR-EGFP in cells[44]. Because

Telomere DNA was proved to load to EVs[9], we first edited the adjacent region of FOXM1-chDNA1 (a

Telomere locus) at chromosome 18 (**Extended Fig. 3b**) and observed the signal of TetO-Telomere in

nucleus followed by its elevated levels in cytoplasm post starvation (**Extended Fig. 3c, d**), confirming

the validity of the TetO/TetR-GFP method. Next, we edited the locus of FOXM1-chDNA3 (*DUX4* locus)

at chromosome 4 (**Extended Fig. 3b**) for following experiments. DUX4 was identified as a transcription

factor playing roles in embryonic development[45] and in promoting immune evasion of cancers[46]. The

signals of TetO array-containing DUX4 DNA (TetO-DUX4) could be visualized in nucleus of A549[*TetO-*

DUX4] cells and its cytoplasmic signals were induced by starvation (**Fig. 3a**), which were further confirmed

by the qPCR results of the cytoDNA of TetO-DUX4 (**Fig. 3b**). The signal of TetO-DUX4 observed in

cytoplasm post starvation was co-localized with the immunofluorescent signal of LC3 (**Fig. 3c**),

confirming that FOXM1-chDNA3 (DUX4 locus) appeared in autophagosomes during autophagy.

Because autophagosomes could fuse with endosomes to mature to MVBs[16] that finally released their

cargos as EVs through fusing with cell membrane[2], we intended to test whether cytoplasmic TetO-DUX4

appeared in MVBs. We used the method of discontinuous sucrose gradient to prepare different fractions

to define MVB compartments[47] from postnuclear supernatants of A549[*TetO-DUX4*] cells. Based on the MVB

specific markers, such as Rab7[48], CD63[49], or TSG101[50], we first confirmed that FOXM1, same as LC3,

existed in the MVB fractions (**Fig. 3d**). The DNA samples were prepared from the harvested fractions

and the results of qPCR specific for TetO-DUX4 DNA showed its significant enrichment in the MVB

fractions (**Fig. 3e**). Furthermore, we observed that the signal of TetO-DUX4 in cytoplasm post starvation

was co-localized with the immunofluorescent signal of CD63 (**Fig. 3f**). We also isolated the samples of

MVBs by immunoprecipitating A549[TetO-DUX4] cell lysates with anti-CD63 antibodies[51] to prepare their

DNA, which showed the elevated levels of TetO-DUX4 DNA in these MVBs compared to the IgG-IP

controls (**Fig. 3g**), suggesting that this FOXM1-chDNA was transported to MVBs during autophagy.

We collected the EVs from A549 cells and A549[TetO-DUX4] cells to prepare DNA samples and the

results of PCR confirmed that TetO-DUX4 DNA appeared in EVs (**Fig. 3h**). The starvation in A549[TetO-

DUX4] cells resulted in the elevated levels of TetO-DUX4 DNA in EVs (**Fig. 3i**). We also collected EVs

from A549[TetO-DUX4] cells through the DIC approach with anti-FOXM1 or anti-LC3 antibodies to prepare

DNA samples and observed the elevated levels of TetO-DUX4 DNA in the samples of FOXM1-specific

EVs or LC3-specific EVs (**Fig. 3j**), suggesting that FOXM1 and LC3 were involved in loading TetO-

DUX4 DNA to EVs. Moreover, after the EVs were treated with PS DNase, the decreased levels of TetO-

DUX4 DNA were observed but not for the mitochondrial DNA controls (**Fig. 3k**), implicating that TetO-

DUX4 DNA existed as linear DNA on the outer membrane of EVs. Together, we confirmed that this

FOXM1-chDNA was loaded to EVs during autophagy.

The loading of FOXM1-chDNAs to EVs relied on FOXM1 and the autophagosome-MVB process

We found that the PCR-amplified DNA samples from *DUX4* locus were able to pulled down FOXM1 or

lipidated LC3-II *in vitro* (**Fig. 4a**), proving the two proteins binding to this FOXM1-chDNA. To confirm

the roles of FOXM1 on the loading of FOXM1-chDNAs to EVs, we generated an A549[FOXM1-/-] cell line

by the CRISPR/cas9 technology targeting *FOXM1* gene in A549 cells (**Extended Data Fig. 4a**). In

A549[FOXM1-/-] cells, the levels of all the tested three FOXM1-chDNAs (FOXM1-chDNA1 from a telomere

locus, FOXM1-chDNA2 from a nonsense region, and FOXM1-chDNA3 from *DUX4* locus) were

decreased in cytoplasm compared to that of wild type control cells post starvation in the qPCR analysis

of purified cytoDNA samples (**Fig. 4b**). Consequently, the EVs from A549$^{FOXM1-/-}$ cells contained much

lower levels of the three FOXM1-chDNAs than that from control cells post starvation (**Fig. 4c**).

Furthermore, we knocked down the expression of FOXM1 in A549$^{TetO-DUX4}$ cells by FOXM1 siRNA

(**Extended Data Fig. 4b**) and observed that the cytoplasmic levels of TetO-DUX4 signals were decreased

in the cells post starvation (**Fig. 4d**). In addition, we constructed a linear DNA containing 72x FOXM1

binding sites (M1-binding DNA) that was bound by recombinant FOXM1 *in vitro* (**Extended Data Fig.**

4c) or FOXM1-DBD-GFP protein *in vivo* when transfected into cells (**Extended Data Fig. 4d**). We

transfected this M1-binding DNA or control DNA at the equal amount into cells and the qPCR analysis

of the isolated evDNA samples showed that the levels of M1-binding DNA in EVs were higher than that

of control DNA (**Fig. 4e**), suggesting that FOXM1 facilitated the FOXM1-recognized DNA fragments

transporting to EVs. We also transfected M1-binding DNA into A549$^{FOXM1-/-}$ cells and found that the

levels of M1-binding DNA in the EVs of A549$^{FOXM1-/-}$ cells were significantly lower than that of wild

type control cells (**Fig. 4f**). Together, the data suggested that the loading of FOXM1-chDNAs to EVs

relied on FOXM1 in cells.

To confirm the formation of autophagosomes contributing the loading of FOXM1-chDNAs to EVs,

we generated an A549$^{Atg5-/-}$ cell line by the CRISPR/cas9 technology targeting *Atg5* gene in A549 cells

(**Extended Data Fig. 4e**). It was well known that the Atg5 knockout resulted in the losing of LC3

lipidation and abolished the formation of autophagosomes[52,53]. In A549$^{Atg5-/-}$ cells, the cytoplasmic and

EVs' levels of the three FOXM1-chDNAs were lower than that of control cells post starvation (**Fig. 4g,**

h), correlated with the decreased levels of FOXM1 and LC3 in the EVs from A549$^{Atg5-/-}$ cells (**Extended**

Data Fig. 4f). In addition, we treated A549[TetO-DUX4] cells with 3-Methyladenine (3-MA), which prevented

the formation of autophagosomes[54], and observed that the cytoplasmic levels of TetO-DUX4 signals were

also decreased in the cells post starvation (**Fig. 4i**). To confirm the formation of MVBs contributing the

loading of FOXM1-chDNAs to EVs, we treated A549 cells with chloroquine (CQ), a lysosome inhibitor

that facilitated the formation of MVBs by inhibiting the fusion of MVBs and lysosomes[16,55]. We observed

that the levels of the three FOXM1-chDNAs were elevated in the EVs of the CQ-treated cells post

starvation (**Fig. 4j**). In addition, we treated A549 cells with a chemical compound GW4869, a neutral

sphingomyelinase inhibitor that prevented the fusion of MVBs with cell membrane and decreased the

secretion of EVs[56]. We collected the EVs from the equal numbers of A549 cells non-treated or treated

with GW4869 and found that the secretion of TSG101 via EVs was decreased by the GW4869 treatment,

confirming the abolished MVB-cell membrane fusion (**Extended Data Fig. 4g**). Consequently, the

secretion of FOXM1 or LC3 via EVs was also decreased by the GW4869 treatment (**Extended Data Fig.**

4g). Through the qPCR analysis of the prepared evDNA samples, we observed that the levels of the three

FOXM1-chDNAs in EVs, similar as that of mtDNA, were decreased by the GW4869 treatment,

suggesting that the secretion of FOXM1-chDNAs to EVs were majorly mediated by MVBs (**Fig. 4k**).

Together, the data suggested that the loading of FOXM1-chDNAs to EVs relied on the autophagosome-

MVB process.

Discussion

In this study, we have discovered that transcription factor FOXM1 mediates the loading of chromatin

DNA fragments to EVs. Specifically, certain FOXM1-bound chromatin DNA fragments (FOXM1-

chDNAs) form LC3-FOXM1-DNA vesicles with FOXM1 and LC3 in cytoplasm during autophagy. The

cytoplasmic LC3-FOXM1-DNAs are released to EVs through the process of an autophagosome-MVB

transport, providing an example of the transcription factor determining the sources of evDNA from

chromatin DNA (**Extended Data Fig. 5**). FOXM1 can theoretically specify its binding regions in

chromatin, providing a clue to determine the sources of evDNA from certain chromatin DNA loci. Based

on this understanding, the accuracy and specificity of disease diagnosis can be improved in the future by

focusing on specific sequences from patient evDNAs.

Although this study focuses on FOXM1, we anticipate that the FOXM1-LC3 interaction is not an

isolated event in nucleus. We have found that other Forkhead Box transcription factors such as FOXA2

and FOXP2 can interact with LC3 (data not shown), implicating that these two transcription factors may

also participate in determining the sources of evDNA from chromatin DNA. In addition, LC3 interacts

with multiple transcription factors that exist in EVs, based on the results of mass spectrometry

analysis[18,37], suggesting that these transcription factors may also play a similar role as FOXM1 for

specifying evDNA from chromatin DNA. Transcription factors are generally considered to bind to

specific DNA regions of chromosome and regulate specific gene expression in nucleus[57]. The findings

of this study expand the understanding of the functions of transcription factors and provide a new

perspective to explore their roles in regulating the composition of EVs' DNA components. Furthermore,

the further enrichment analysis of the specific sequences of patient evDNAs can predict certain

transcription factors that bind to the sequences, so as to establish the association between the transcription

factors and diseases to deduce the pathogenesis or disease progression.

The FOXM1-chDNAs identified in this study can be annotated as functional genes, telomere,

centromere, and nonsense regions, implicating their potential effects in the cells that receive these DNAs.

Interestingly, a recent study confirmed that Telomere DNAs carried by the EVs from antigen-presenting

cells can horizontally transfer to T cells and promote long-term immunological memory by rescuing T

cells from senescence[9]. We have observed the signals of TetO-DUX4 in recipient cells that are treated

with the A549[TetO-DUX4]-derieved EVs and noticed the changes of phenotypes of recipient cells that are

transfected with the PCR-amplified DNA of *DUX4* locus (data not shown), providing another example

that the certain FOXM1-chDNAs of donor cells can perform specific biological functions in recipient

cells after the horizontal gene transfer (HGT) and may regulate the extracellular microenvironment of

the cells. Furthermore, because the EVs of local cells are able to enter the circulatory system, we

speculate that certain FOXM1-chDNAs may possess relevant roles in HGT to distant tissues. These

hypotheses are being investigated in our laboratory.

FOXM1 is generally considered as a transcription factor regulating gene expression in nucleus of

cells[27]. This study extends its function to loading certain chromatin DNA fragments to EVs. One key

question still remaining is how chromatin DNA breaks to specific fragments. So far, the appearance of

chromatin DNA in cytoplasm is mediated mainly by the micronuclei and cytoplasmic chromatin

fragment pathways[58], while both the mechanisms cannot explain the specific breakage of chromatin DNA.

Although nuclear LC3 interacts with LaminB1 to result in the cytoplasmic translocation of

transcriptionally inactive heterochromatin DNA[10], the mechanism of specific DNA fragments breaking

off chromosomes has not been explored. Recently, the complex of Rec114, Mei4 and Mer2 in cells has

been reported to control the precise DNA breakage[59], suggesting that the breakage of chromatin can be

precisely achieved. Future studies of FOXM1's interaction with the controlled DNA breakage machinery

will reveal how relevant chromatin DNA fragments are precisely produced.

Methods

The complete Materials and Methods were described in Supplementary Materials.

ChIP sequencing, evDNA sequencing, Bioinformatics analysis, RT-qPCR. For sequencing, ChIP and evDNA samples were prepared as described previously[60]. The ChIP and evDNA were purified and used for constructing sequencing libraries with a NEBNext Ultra DNA Library Prep Kit for Illumina (New England Biolabs, Ipswich, MA, USA). The library quantifications were assessed on the Agilent Bioanalyzer 2100 system. The library preparations were sequenced on an Illumina Hiseq platform and 50 bp single-end reads or 150 bp paired-end reads were generated.

For bioinformatics analysis, the following pipeline was used for analysis of all sequence data sets. Firstly, fastp (version 0.20.0, parameter: '-q 15 -u 40 -n 6 -l 15 ')[61] was used to trim adaptors and low-quality reads. Then, we evaluated the quality of NGS short reads by searching them on the FastQC (version 0.11.8) program (http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Thirdly, the remaining reads were aligned to human genome hg38 by bwa men (version 0.7.17,parameter: '-t 20')[62], since ChIP-Seq and evDNA-Seq interrupted fragments were usually small and the percentage of unique sequences in the total number of sequences was the focus of attention. Finally, peak calling of the aligned reads was made using macs3 (version 3.0.0a6, parameter: '-f BAM -g hs -p 0.1 -B')[63], and peaks were annotated using the R/Bioconductor package ChIPseeker[64]. Browser views of peaks are shown using Integrated Genomics Viewer (IGV; broadinstitute.org/igv)[65].

For qPCR, the following primers were used for qPCR analyses of chromatin loci. FOXM1-chDNA1 (chr18): forward, 5'-TGA TCA CCC AGG AGA CGG-3'; reverse, 5'-CTT GGG TGA TCA GTG GCG AG-3'. FOXM1-chDNA2 (chr9): forward, 5'-CTG CAA AAT GGA CCA ATC AGC-3'; reverse, 5'-AAG ATG GTG TGT CCG GAA TTT G-3'. FOXM1-chDNA3 (chr4): forward, 5'-TCA CAA GCC CCC

TGT AGG-3'; reverse, 5'-TCC AAC TCT TGC CTG GTC TC-3'.

DNA Fluorescence *in situ* hybridization (FISH). DNA-FISH probes were constructed by Exon

Biotechnology Inc (Guangzhou, China). The probes covered genomic regions (hg38) used in this study

was as follow: chr18: 103010-113570 (chr18_FOXM1-chDNA1_10 Kb). For fluorescence *in situ*

hybridization assay, DNA FISH was performed as described[66]. Briefly, coverslips containing fixed and

permeabilized cells were quenched by 3% H_2O_2, followed by dehydration in 70%, 85%, and 100%

ethanol for 1 min each. FISH probe (chr18_ FOXM1-chDNA1_10 Kb) was added onto a glass slide, and

lower the coverslips onto the slide slowly. Place the slides with samples to be hybridized in a heating

block at 85℃ for 5min, and then 37℃ for 20 hours. The slide was then removed, and the sample was

washed with Wash Buffer (1 X, preheated to 73℃) for 2 min. Added blocking buffer onto the hybridized

samples and incubated at 37℃ for 30 min. Added primary antibody onto coverslip and incubated at 37℃

for 30 min. The sample was washed with Wash Buffer (1 X, preheated to 45℃) for 5 min. Added

secondary antibody onto coverslip and incubated at 37℃ for 30 min. Wash coverslip in Wash Buffer (1x)

three times at 45℃ for 5 min. Re-wash once with PBS. Preparation of TSA solution at TSA: 0.15%H2O2:

TSA amplification buffer = 1:5:500-1:10:1000. Add TSA solution to coverslip and incubate 5-15min at

RT. Wash coverslip in Wash Buffer (1 X) three times for 30 min. Mount coverslip with Anti-fade

Fluorescence Mounting Medium (with DAPI, Abcam, #ab104139).

CRISPR/Cas9-mediated knock-in of a TetO array (96x) for visualization of FOXM1-chDNAs. To

obtain a TetO-knockin cell line, we designed a pair of gRNAs using http://crispr.mit.edu/. The pair of

gRNAs (5'-CAC CTG GGT GAT CAG AGC AA-3' and 5'-TCT AGG CTT TGG CCT ACA GG-3')

bound to FOXM1-chDNA1 or gRNAs (5'-GCA GGC AGA GCG TAA GCA AA-3' and 5'-GGA CAC

ACG TTT AAT CGA GT-3') bound to FOXM1-chDNA3 was cloned into the pX335 vector (Addgene,

42335) before the gRNA scaffold. Furthermore, we designed two donor plasmids that contain the TetO

array (96x) flanked by 500-bp homology arms of the FOXM1-chDNAs in the pSP2-96-merTetO-EFS-

BLaR vector (Addgene, 118713). The primers (5'-CCC TTT CGT CTT CAA GAA TTC CAT CAC CTG

GGT GAT CAG TGT AGA-3', 5'-ACC CAT TCC TAG GGC GAA TTC GCT CTC TGA TCA CCC

AGG TGA T-3'; 5'-GCG CTGC TAG CTT AAG GTA CCA GCG TAG GCC AAA GCC TAG AC-3',

5'-CCC AGA TCT ATC GAT GGT ACC CTT GTC TAG GAT CTG CCT ACA GGG-3') were

synthesized and amplified homology arms of the FOXM1-chDNA1. The primers (5'- CCC TTT CGT

CTT CAA GAA TTC CCT GTA GGC AAG CCT ACA CAA GT-3', 5'-ACC CAT TCC TAG GGC GAA

TTC AGC GCT TAC GCT CTG CCT G-3'; 5'-GCG CTG CTA GCT TAA GGT ACC GAC CGA TTA

AAC GTG TGT CCT TT-3', 5'-CCC AGA TCT ATC GAT GGT ACC GTT TTT TCC TTT AAG ACT

TAT GTA ATG AAT T-3') were synthesized and amplified homology arms of the FOXM1-chDNA3. The

paired gRNAs and donor plasmids were co-transfected into A549 cells. Blasticidin (10 μg/ml) selection

was started 1 day after transfection. Three weeks after blasticidin selection, TetO-knockin cell clones

were transplanted into 24-well plates. Cells were collected from 24-well plates and extracted genomic

DNA (gDNA) for genotyping.

For FOXM1-chDNAs loci visualization, the TetO-knockin cells were transiently expressed with

TetR-EGFP. Cells were fixed in 4% paraformaldehyde and mounted with Mounting Medium with DAPI

(Abcam, ab104139). Images were captured by FV1200 laser scanning microscopes (Olympus, Japan).

For qPCR, the following primers were used for qPCR analyses of knock-in of FOXM1-chDNAs. TetO-

Telomere DNA (FOXM1-chDNA1 locus): forward, 5'-GAA GAC TAC AGC GTC GCC AG-3'; reverse,

5'-CGC GAC GAT ACA AGT CAG GT-3'. TetO-DUX4 DNA (FOXM1-chDNA3 locus): forward, 5'-

TCT GAA GAC TAC AGC GTC GC-3'; reverse, 5'-ACA CAT AAC CAG AGG GCA GC-3'.

Extracellular vesicle isolation. Cells seeded in 150 mm culture dishes at approximately 80% confluence

were incubated with serum-free DMEM for 1 day or 2 days. Cell viability was assessed using trypan blue

and only > 95% viability was used for EVs isolation. The cell-conditioned medium was collected by

sequential ultracentrifugation[1,18]. Briefly, the medium was first subjected to a centrifugation at 400 × g

for 10 min to remove cells. Next, the supernatant was centrifuged at 2,000 × g for 20 min to remove

debris and apoptotic bodies. Then, the supernatant was centrifuged at 15,000 × g for 40 min to obtain

large EVs. The resulting large EVs pellet was resuspended in a large volume of PBS buffer followed by

ultracentrifugation at 15,000 × g for 40 min to wash the sample. To remove any remaining any large EVs,

the media supernatant from the first 15,000 × g step was passed through a 0.22-μm pore PES filter

(Millipore, Bedford, MA, USA). This supernatant (pre-cleared medium) was next subjected to

ultracentrifugation at 120,000 × g for 4 h in a SW 28 Ti Swinging-Bucket Rotor (Beckman Coulter,

Fullerton, CA) to sediment small EVs. The crude small EVs pellet was resuspended in a large volume of

PBS followed by ultracentrifugation at 120,000 × g for 4 h to wash the sample. All following

centrifugation steps were performed at 4℃. Notably, for the comparison of EVs in different conditions,

the results must be corrected by total cell number or total protein concentration to eliminate differences

in cell seeding.

Acknowledgements

We thank Dr. Zhuoxian Rong of Central South University for help with construction of FOXM1 knockout A549 cell line by CRISPR/Cas9 technology; thank Dr. Jianhui Jiang of Hunan University for help with confocal microscopy. This work was supported by the National Natural Science Foundation of China (grant number 81773169 to Y.T.), China Changsha Development and Reform Commission "Mass entrepreneurship and innovation program" (2018-68) and "Innovation platform construction program" (2018-216).

Author contributions

Y.Z. and Y.T. conceived this project. Y.Z., N.D., Y.L., X.Z., S.Y., H.L. and Y.W. performed all of experiments. Y.Z., H.C., P.F., Y.P., L.Y. and T.Y. analyzed ChIP-seq and evDNA-seq. M.O. performed molecular docking simulation. Y.C, L.Y. and X.H. provided administrative and material supports. Y.T. supervised this study. Y.Z. and Y.T. composed the manuscript. All authors reviewed the manuscript and discussed the work.

Competing interests

The authors declare no competing interests.

Supplementary Material

Supplementary methods, figures and tables.

References

| 415 | **References** |

1 Jeppesen, D. K. *et al.* Reassessment of Exosome Composition. *Cell* **177**, 428-445 e418 (2019).

2 van Niel, G., D'Angelo, G. & Raposo, G. Shedding light on the cell biology of extracellular
vesicles. *Nat Rev Mol Cell Biol* **19**, 213-228 (2018).

3 Sina, A. A. *et al.* Methylation dependent gold adsorption behaviour identifies cancer derived
extracellular vesicular DNA. *Nanoscale Horiz* **5**, 1317-1323 (2020).

4 Elzanowska, J., Semira, C. & Costa-Silva, B. DNA in extracellular vesicles: biological and
clinical aspects. *Mol Oncol* **15**, 1701-1714 (2020).

5 Douanne, N. *et al.* Leishmania parasites exchange drug-resistance genes through extracellular
vesicles. *Cell Rep* **40**, 111121 (2022).

6 Soucy, S. M., Huang, J. & Gogarten, J. P. Horizontal gene transfer: building the web of life. *Nat*
Rev Genet **16**, 472-482 (2015).

7 Cai, J., Wu, G., Jose, P. A. & Zeng, C. Functional transferred DNA within extracellular vesicles.
Exp Cell Res **349**, 179-183 (2016).

8 Kawamura, Y., Sanchez Calle, A., Yamamoto, Y., Sato, T. A. & Ochiya, T. Extracellular vesicles
mediate the horizontal transfer of an active LINE-1 retrotransposon. *J Extracell Vesicles* **8**,
1643214 (2019).

9 Lanna, A. *et al.* An intercellular transfer of telomeres rescues T cells from senescence and
promotes long-term immunological memory. *Nat Cell Biol* **24**, 1461-1474 (2022).

10 Dou, Z. *et al.* Autophagy mediates degradation of nuclear lamina. *Nature* **527**, 105-109 (2015).

11 Lee, Y. K. & Lee, J. A. Role of the mammalian ATG8/LC3 family in autophagy: differential and
compensatory roles in the spatiotemporal regulation of autophagy. *BMB Rep* **49**, 424-430 (2016).

12 Oldenburg, A. R. & Collas, P. Mapping Nuclear Lamin-Genome Interactions by Chromatin
Immunoprecipitation of Nuclear Lamins. *Methods Mol Biol* **1411**, 315-324 (2016).

13 Tanida, I., Ueno, T. & Kominami, E. LC3 and Autophagy. *Methods Mol Biol* **445**, 77-88 (2008).

14 Itakura, E., Kishi-Itakura, C. & Mizushima, N. The hairpin-type tail-anchored SNARE syntaxin
17 targets to autophagosomes for fusion with endosomes/lysosomes. *Cell* **151**, 1256-1269
(2012).

15 Diao, J. *et al.* ATG14 promotes membrane tethering and fusion of autophagosomes to
endolysosomes. *Nature* **520**, 563-566 (2015).

16 Fader, C. M. & Colombo, M. I. Autophagy and multivesicular bodies: two closely related
partners. *Cell Death Differ* **16**, 70-78 (2009).

17 Leidal, A. M. & Debnath, J. Emerging roles for the autophagy machinery in extracellular vesicle
biogenesis and secretion. *FASEB Bioadv* **3**, 377-386 (2021).

18 Leidal, A. M. *et al.* The LC3-conjugation machinery specifies the loading of RNA-binding
proteins into extracellular vesicles. *Nat Cell Biol* **22**, 187-199 (2020).

19 Leidal, A. M. & Debnath, J. LC3-dependent extracellular vesicle loading and secretion
(LDELS). *Autophagy* **16**, 1162-1163 (2020).

20 Gruenbaum, Y., Wilson, K. L., Harel, A., Goldberg, M. & Cohen, M. Review: nuclear lamins--
structural proteins with fundamental functions. *J Struct Biol* **129**, 313-323 (2000).

21 Costa, R. H., Kalinichenko, V. V., Holterman, A. X. & Wang, X. Transcription factors in liver

development, differentiation, and regeneration. *Hepatology* **38**, 1331-1347 (2003).

22   Fu, Z. *et al.* Plk1-dependent phosphorylation of FoxM1 regulates a transcriptional programme
required for mitotic progression. *Nat Cell Biol* **10**, 1076-1082 (2008).

23   Tan, Y., Raychaudhuri, P. & Costa, R. H. Chk2 mediates stabilization of the FoxM1 transcription
factor to stimulate expression of DNA repair genes. *Mol Cell Biol* **27**, 1007-1016 (2007).

24   Song, I. S. *et al.* FOXM1-Induced PRX3 Regulates Stemness and Survival of Colon Cancer
Cells via Maintenance of Mitochondrial Function. *Gastroenterology* **149**, 1006-1016 e1009
(2015).

25   Xie, Z. *et al.* Foxm1 transcription factor is required for maintenance of pluripotency of P19
embryonal carcinoma cells. *Nucleic Acids Res* **38**, 8027-8038 (2010).

26   Li, D. *et al.* The critical role of dysregulated FOXM1-PLAUR signaling in human colon cancer
progression and metastasis. *Clin Cancer Res* **19**, 62-72 (2013).

27   Sanders, D. A. *et al.* FOXM1 binds directly to non-consensus sequences in the human genome.
Genome Biol **16**, 130 (2015).

28   Marceau, A. H. *et al.* An order-to-disorder structural switch activates the FoxM1 transcription
factor. *Elife* **8**, e46131 (2019).

29   Liu, C., Barger, C. J. & Karpf, A. R. FOXM1: A Multifunctional Oncoprotein and Emerging
Therapeutic Target in Ovarian Cancer. *Cancers (Basel)* **13**2021).

30   Hu, C. D. & Kerppola, T. K. Simultaneous visualization of multiple protein interactions in living
cells using multicolor fluorescence complementation analysis. *Nat Biotechnol* **21**, 539-545
(2003).

31   Birgisdottir, A. B., Lamark, T. & Johansen, T. The LIR motif - crucial for selective autophagy.
J Cell Sci **126**, 3237-3247 (2013).

32   Johansen, T. & Lamark, T. Selective Autophagy: ATG8 Family Proteins, LIR Motifs and Cargo
Receptors. *J Mol Biol* **432**, 80-103 (2020).

33   Sentelle, R. D. *et al.* Ceramide targets autophagosomes to mitochondria and induces lethal
mitophagy. *Nat Chem Biol* **8**, 831-838 (2012).

34   Chaudhury, S. *et al.* Benchmarking and analysis of protein docking performance in Rosetta v3.2.
PLoS One **6**, e22477 (2011).

35   Littler, D. R. *et al.* Structure of the FoxM1 DNA-recognition domain bound to a promoter
sequence. *Nucleic Acids Res* **38**, 4527-4538 (2010).

36   Kuang, Y. *et al.* Structural basis for the phosphorylation of FUNDC1 LIR as a molecular switch
of mitophagy. *Autophagy* **12**, 2363-2373 (2016).

37   Hoshino, A. *et al.* Extracellular Vesicle and Particle Biomarkers Define Multiple Human
Cancers. *Cell* **182**, 1044-1061 e1018 (2020).

38   Moller, A. & Lobb, R. J. The evolving translational potential of small extracellular vesicles in
cancer. *Nat Rev Cancer* **20**, 697-709 (2020).

39   Gao, Y. *et al.* Small Extracellular Vesicles: A Novel Avenue for Cancer Management. *Front*
Oncol **11**, 638357 (2021).

40   Yoshioka, Y. *et al.* Comparative marker analysis of extracellular vesicles in different human
cancer types. *J Extracell Vesicles* **2**, eCollection 2013 (2013).

41   Abmayr, S. M., Yao, T., Parmely, T. & Workman, J. L. Preparation of nuclear and cytoplasmic

extracts from mammalian cells. *Curr Protoc Mol Biol* **Chapter 12**, Unit 12 11 (2006).

42   Wang, Y. *et al.* eccDNAs are apoptotic products with high innate immunostimulatory activity.
Nature **599**, 308-314 (2021).

43   Liu, H. *et al.* Analysis of extracellular vesicle DNA at the single-vesicle level by nano-flow
cytometry. *J Extracell Vesicles* **11**, e12206 (2022).

44   Tasan, I. *et al.* CRISPR/Cas9-mediated knock-in of an optimized TetO repeat for live cell
imaging of endogenous loci. *Nucleic Acids Res* **46**, e100 (2018).

45   Hendrickson, P. G. *et al.* Conserved roles of mouse DUX and human DUX4 in activating
cleavage-stage genes and MERVL/HERVL retrotransposons. *Nat Genet* **49**, 925-934 (2017).

46   Chew, G. L. *et al.* DUX4 Suppresses MHC Class I to Promote Cancer Immune Evasion and
Resistance to Checkpoint Blockade. *Dev Cell* **50**, 658-671 e657 (2019).

47   de Araujo, M. E., Lamberti, G. & Huber, L. A. Homogenization of Mammalian Cells. *Cold*
Spring Harb Protoc **2015**, 1009-1012 (2015).

48   Shearer, L. J. & Petersen, N. O. Distribution and Co-localization of endosome markers in cells.
Heliyon **5**, e02375 (2019).

49   Piper, R. C. & Katzmann, D. J. Biogenesis and function of multivesicular bodies. *Annu Rev Cell*
Dev Biol **23**, 519-547 (2007).

50   Nabhan, J. F., Hu, R., Oh, R. S., Cohen, S. N. & Lu, Q. Formation and release of arrestin domain-
containing protein 1-mediated microvesicles (ARMMs) at plasma membrane by recruitment of
TSG101 protein. *Proc Natl Acad Sci U S A* **109**, 4146-4151 (2012).

51   Ferreira, J. V. *et al.* LAMP2A regulates the loading of proteins into exosomes. *Sci Adv* **8**,
eabm1140 (2022).

52   Klionsky, D. J. *et al.* Guidelines for the use and interpretation of assays for monitoring
autophagy (4th edition)(1). *Autophagy* **17**, 1-382 (2021).

53   Otomo, C., Metlagel, Z., Takaesu, G. & Otomo, T. Structure of the human ATG12~ATG5
conjugate required for LC3 lipidation in autophagy. *Nat Struct Mol Biol* **20**, 59-66 (2013).

54   Seglen, P. O. & Gordon, P. B. 3-Methyladenine: specific inhibitor of autophagic/lysosomal
protein degradation in isolated rat hepatocytes. *Proc Natl Acad Sci U S A* **79**, 1889-1892 (1982).

55   Mauthe, M. *et al.* Chloroquine inhibits autophagic flux by decreasing autophagosome-lysosome
fusion. *Autophagy* **14**, 1435-1455 (2018).

56   Trajkovic, K. *et al.* Ceramide triggers budding of exosome vesicles into multivesicular
endosomes. *Science* **319**, 1244-1247 (2008).

57   Todeschini, A. L., Georges, A. & Veitia, R. A. Transcription factors: specific DNA binding and
specific gene regulation. *Trends Genet* **30**, 211-219 (2014).

58   Miller, K. N. *et al.* Cytoplasmic DNA: sources, sensing, and role in aging and disease. *Cell* **184**,
5506-5526 (2021).

59   Claeys Bouuaert, C. *et al.* DNA-driven condensation assembles the meiotic DNA break
machinery. *Nature* **592**, 144-149 (2021).

60   Kharchenko, P. V., Tolstorukov, M. Y. & Park, P. J. Design and analysis of ChIP-seq experiments
for DNA-binding proteins. *Nat Biotechnol* **26**, 1351-1359 (2008).

61   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884-i890 (2018).

62   Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*,

1303.3997 (2013).

63   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

64   Yu, G., Wang, L. G. & He, Q. Y. ChIPseeker: an R/Bioconductor package for ChIP peak

annotation, comparison and visualization. *Bioinformatics* **31**, 2382-2383 (2015).

65   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26 (2011).

66   Tsuchiya, K. D. Fluorescence in situ hybridization. *Clin Lab Med* **31**, 525-542, vii-viii (2011).

 Figure Legends

Figure 1. The interaction between FOXM1 and LC3. (a) Representative FOXM1 and LC3 images in

A549 cells that were fixed by 4% paraformaldehyde and immunostained with anti-FOXM1 and anti-LC3

antibodies. Scale bar, 10 μm. The Pearson Correlation Coefficient (PCC) value of FOXM1 and LC3

colocalization was 0.66±0.08. **(b)** Co-immunoprecipitation of endogenous FOXM1 and LC3 in A549

cells. **(c)** The BiFC assays of FOXM1-LC3 interaction. HEK293T cells were transfected with the

indicated combination of split Venus constructs for 24 hrs. Scale bar, 10 μm. **(d)** HEK293T cells (5 x 10^6)

were transfected with Myc-tagged FOXM1 deletion mutant plasmids (5 μg). All cell lysates were pulled

down with bacterially purified GST-LC3 protein (30 μg) and subjected to immunoblotting as indicated.

(e) The GFP-fused FOXM1 truncations were expressed and pulled down with bacterially purified GST-

LC3 as described above. **(f)** The GFP-fused FOXM1-DBD point mutations were expressed and pulled

down with bacterially purified GST-LC3 as described above. **(g)** The Myc-tagged FOXM1 was pulled

down with bacterially purified GST-LC3 truncated mutants as described above. **(h)** The Myc-tagged

FOXM1 was expressed and pulled down with bacterially purified GST-LC3 truncated point mutants as

described above. **(i)** The molecular docking simulation for FOXM1-LC3 interaction by Rosetta Dock. **(j)**

EMSA showed FOXM1, LC3 and DNA formed LC3-FOXM1-DNA complex.

Figure 2. FOXM1-chDNAs are loaded to extracellular vesicles during autophagy. (a)

Representative LC3-FOXM1-DNA vesicles image in cytoplasm of starved A549 cells (6 hrs) that were

immunostained with anti-FOXM1 and anti-LC3 antibodies. The white arrows indicated LC3-FOXM1-

DNA vesicle. Scale bar, 10 μm. **(b)** Quantification of cytoplasmic LC3-FOXM1-DNA vesicle in non-

starved or starved A549 cells (6 hrs). (n = 4 independent experiments). **$P<0.01$; unpaired two-tailed.

(c) The BiFC assays showed that the exogenous VN173-FOXM1, VC155-LC3 and chromatin DNA

vesiculated in cytoplasm of starved A549 cells (6 hrs). Scale bar, 10 μm. **(d)** EVs were collected from

starved A549 cells (48 hrs) and subjected to immunoblotting for TSG101, FOXM1 and LC3. **(e)** evDNAs

were purified from EVs collected from starved A549 cells (48 hrs). The different concentrations of

evDNAs were analyzed by 1% vertical agarose gel electrophoresis. **(f)** The cytoplasmic DNA (cytoDNA)

was extracted from non-starved or starved A549 cells (24 hrs) for qPCR. The mitochondria DNA

(mtDNA, *ND-2* gene) was used as the internal loading control. The Bars, mean±s.e.m.; n=3; *$P<0.05$;

NS, non-significant; unpaired two-tailed. **(g)** Representative DNA-FISH images of FOXM1-chDNA1-

adjacent region (10 Kb probe) in starved A549 cells (6 hrs). The white arrows indicated FOXM1-

chDNA1. **(h)** The evDNA was extracted from non-starved or starved A549 cells (48 hrs) for qPCR. The

data were normalized by mtDNA. **(i)** The qPCR analyses of FOXM1-chDNAs in anti-FOXM1-IP

evDNA or anti-LC3-IP evDNA from EVs collected from starved A549 cells (48 hrs). **(j)** The qPCR

analyses of FOXM1-chDNAs in the DIC-purified GFP-FOXM1-specific EVs or DIC-purified GFP-

LC3-specific EVs collected from starved A549 stably expressing GFP-FOXM1 or GFP-LC3 cells (48

hrs). The Bars, mean±s.e.m.; n=3; *$P<0.05$, #$P<0.05$, **$P<0.01$; unpaired two-tailed.

Figure 3. The FOXM1-chDNAs knocked-in with the TetO array were loaded to EVs. (a)

Representative TetO-DUX4 images in starved A549$^{TetO-DUX4}$ cells (6 hrs) that were transiently expressed

TetR-EGFP. The white arrows indicated TetO-DUX4. Scale bar, 10 μm. **(b)** The A549$^{TetO-DUX4}$ cells were

treated with starvation and cytoplasmic DNA (cytoDNA) was extracted 24 hours later for qPCR. The

mtDNA was internal control. The Bars, mean±s.e.m.; n=3; **$P<0.01$; NS, non-significant; unpaired two-

tailed. **(c)** Immunostaining of the endogenous LC3 and TetO-DUX4 in starved A549$^{TetO\text{-}DUX4}$ cells (6 hrs)

that were exogenously expressed TetR-EGFP. The white arrows indicated TetO-DUX4 that was

translocated into cytoplasmic autophagosome. Scale bar, 10 μm. **(d)** MVB compartments were isolated

from starved A549$^{TetO\text{-}DUX4}$ cells (24 hrs) by discontinuous sucrose gradient and subjected to

immunoblotting. **(e)** The qPCR analyses of TetO-DUX4 in MVB fractions collected from starved

A549$^{TetO\text{-}DUX4}$ cells (24 hrs) by discontinuous sucrose gradient. The Bars, mean±s.e.m.; n=3; **$P<0.01$;

NS, non-significant; unpaired two-tailed. **(f)** Immunostaining of the endogenous CD63 and TetO-DUX4

in starved-A549$^{TetO\text{-}DUX4}$ cells (6 hrs). The white arrows indicated TetO-DUX4 that was translocated into

cytoplasmic MVB. Scale bar, 10 μm. **(g)** The qPCR analyses of TetO-DUX4 in MVBs collected from

starved-A549$^{TetO\text{-}DUX4}$ cells (24 hrs) by CD63-immunoprecipitation. The Bars, mean±s.e.m.; n=3;

$P<0.01$; NS, non-significant; unpaired two-tailed. **(h) PCR was used to amplify the TetO-DUX4,

DUX4, and mitochondria DNAs in the evDNAs collected from starved A549^{WT} or A549$^{TetO\text{-}DUX4}$ cells (48

hrs). **(i)** The qPCR analyses of TetO-DUX4 in the EVs collected from starved-A549$^{TetO\text{-}DUX4}$ cells (48

hrs). **(j)** The qPCR analyses of TetO-DUX4 in the DIC-purified FOXM1-specific EVs or LC3-specific

EVs collected from starved-A549$^{TetO\text{-}DUX4}$ cells (48 hrs). **(k)** The qPCR analyses of TetO-DUX4 in the

Plasmid-Safe ATP-dependent DNase (PS)-digested EVs collected from starved-A549$^{TetO\text{-}DUX4}$ cells. The

Bars, mean±s.e.m.; n=3; *$P<0.05$; #$P<0.05$; **$P<0.01$; ##$P<0.01$; unpaired two-tailed.

Figure 4. The loading of FOXM1-chDNAs to EVs relied on FOXM1 and the autophagosome-MVB

process. (a) A549 cell lysates were incubated with biotinylated DUX4 DNA (500 ng). The bound

proteins were immunoprecipitated with streptavidin microbeads and blotted by anti-FOXM1 or LC3

antibodies. **(b, c)** The qPCR analyses of three FOXM1-chDNAs in the cytoplasm or EVs collected from

starved A549 or A549$^{FOXM1-/-}$ cells. The data were normalized by mtDNA. The Bars, mean±s.e.m.; n=3;

$P<0.01$; unpaired two-tailed. **(d) A549$^{TetO-DUX4}$ cells were transfected with 50 nM *FOXM1* siRNA and

1 μg TetR-EGFP plasmid. After the transfection (48 hrs), cells were starved (6 hrs) and imaged for the

signal of TetO-DUX4 (left). The white arrows indicated TetO-DUX4. Scale bar, 10 μm. The TetO-DUX4

signal in cytoplasm was quantitated (right). n = 3 independent experiments; *$P<0.05$; unpaired two-tailed.

(e) HEK293T cells were transfected with linear M1-binding or control DNA at the equal amount. After

the transfection (24 hrs), all cells were starved (48 hrs) for collecting EVs**.** The qPCR analyses were

performed for the mtDNA and transfected DNA in evDNAs purified from the EVs. **(f)** A549 and

A549$^{FOXM1-/-}$ cells were transfected with linear M1-binding DNA at the equal amount. After the

transfection (24 hrs), all cells were starved (48 hrs) for collecting EVs**.** The qPCR analyses were

performed for the mtDNA and transfected DNA in evDNAs purified from the EVs. The Bars,

mean±s.e.m.; n=3; *$P < 0.05$; **$P<0.01$; NS, non-significant; unpaired two-tailed. **(g, h)** The qPCR

analyses of three FOXM1-chDNAs in the cytoplasm or EVs collected from starved A549 or A549$^{Atg5-/-}$

cells. The data were normalized by mtDNA. The Bars, mean±s.e.m.; n=3; **$P<0.01$; unpaired two-tailed.

(i) A549$^{TetO-DUX4}$ cells were transfected with TetR-EGFP plasmid (1 μg). After the transfection (24 hrs),

cells were treated with 3-MA (5 mM) and starved (6 hrs). Cells were immunostained with anti-LC3

antibodies and imaged for TetO-DUX4. The white arrows indicated the signal of TetO-DUX4. Scale bar,

10 μm. The TetO-DUX4 signal in cytoplasm was compared (right). n = 3 independent experiments;

*$P<0.05$; unpaired two-tailed. **(j)** EVs were collected from A549 cells that were treated with CQ (20 μM)

and starvation (24 hrs). The qPCR analyses were performed for the three FOXM1-chDNAs in evDNAs

purified form the EVs. **(k)** EVs were collected from A549 cells that were treated with GW4869 (20 μM)

and starvation (24 hrs). The qPCR analyses were performed for the three FOXM1-chDNAs in evDNAs

purified form the EVs.

Figure 1

633
634

Figure 2

636
637

638   **Figure 3**

639
640

641 **Figure 4**

642
643

644 **Table. FOXM1-chDNAs.**

Names	Sequence signatures	Chr	geneID
FOXM1-chDNA1	Telomere	18	ENSG00000263006
FOXM1-chDNA2	Nonsense region	9	ENSG00000228522
FOXM1-chDNA3	Coding region (*DUX4*)	4	ENSG00000260596
FOXM1-chDNA4	Coding region (*PGK1*)	X	ENSG00000102144
FOXM1-chDNA5	Coding region (*ITFG2*)	12	ENSG00000111203
FOXM1-chDNA6	Coding region (*FOXM1*)	12	ENSG00000111206
FOXM1-chDNA7	Coding region (*FOXM1*)	12	ENSG00000111206
FOXM1-chDNA8	Coding region (*TBC1D5*)	3	ENSG00000131374
FOXM1-chDNA9	Coding region (*XR_946063.2*)	10	ENSG00000166317
FOXM1-chDNA10	Coding region (*XR_943570.1*)	14	ENSG00000258314
FOXM1-chDNA11	Centromere	5	ENSG00000170571
FOXM1-chDNA12	Centromere	1	ENSG00000224857
FOXM1-chDNA13	Centromere	20	ENSG00000227195
FOXM1-chDNA14	Centromere	3	ENSG00000251727
FOXM1-chDNA15	Nonsense region	16	ENSG00000261239
FOXM1-chDNA16	Nonsense region	16	ENSG00000261239
FOXM1-chDNA17	Nonsense region	16	ENSG00000261239
FOXM1-chDNA18	Nonsense region	16	ENSG00000261239
FOXM1-chDNA19	Nonsense region	16	ENSG00000261239
FOXM1-chDNA20	Centromere	17	ENSG00000263433
FOXM1-chDNA21	Centromere	17	ENSG00000263433
FOXM1-chDNA22	Centromere	17	ENSG00000263433
FOXM1-chDNA23	Centromere	17	ENSG00000264970
FOXM1-chDNA24	Nonsense region	1	ENSG00000276118
FOXM1-chDNA25	Nonsense region	5	ENSG00000284042

645