cir-DNA sequencing revealed the landscape of extrachromosomal circular DNA in articular cartilage and the potential roles in osteoarthritis

Wei Xiang[#1], Tongyi Zhang[#1,2]，Song Li[#3]，Yunquan Gong[1]，Xiaoqing Luo[3], Jing Yuan[4]，Yaran Wu[5]，Xiaojing Yan[4]，Yan Xiong[6], Jiqin Lian[5], Guangyu Zhao[7], Changyue Gao[1], Liang Kuang[3] and Zhenhong Ni[1]

[1] State Key Laboratory of Trauma, Burns and Combined Injury, Department of Rehabilitation Medicine, Daping Hospital, Army Medical University, Chongqing, China;

[2] Department of General practice, Chinese PLA General Hospital of the Central Theater Command，Wuhan, China

[3] Department of Wound Repair and Rehabilitation Medicine, Center of Bone Metabolism and Repair, Laboratory for Prevention and Rehabilitation of Training Injuries, State Key Laboratory of Trauma, Burns and Combined Injury, Trauma Center, Research Institute of Surgery, Daping Hospital, Army Medical University, Chongqing, China;

[4] Department of Biochemistry and Molecular Biology, College of Basic Medical Sciences, Army Medical University, Chongqing, China;

[5] Department of pharmacy and clinical laboratory, Army Medical University, Chongqing, China;

[6] Department of orthopedics, Daping Hospital, Army Medical University, Chongqing, China;

[7] Seventeen Squadron Five Brigade, School of Basic Medical Science, Army Medical University, Chongqing, China;

[#] Contributed equally

Correspondence: Zhenhong Ni (nizhenhong1986@163.com) or Liang Kuang (blue_sky128@163.com) or Changyue Gao (gaochangyue1990@163.com)

Abstract

Objective: Extrachromosomal circular DNA (eccDNA) has been shown to be involved in several physiological and pathological processes including immunity, inflammation, aging and tumor. However, the expression of eccDNA in cartilage has been not reported until now. In this study, we aimed to investigate the landscape of eccDNA in articular cartilage and analyze the potential roles in Osteoarthritis (OA).

Methods: The samples of articular cartilage were obtained from total knee arthroplasty (TKA) donors with OA. The mtDNAs (mitochondrial DNA) and the

linear DNAs from chondrocytes of articular cartilage were removed. Then the eccDNAs were enriched for cir-DNA sequencing. After quality control evaluation, we systematically revealed the identified eccDNA data including size distribution, the size range and sequence pattern. Moreover, we explored and disscussed the potential roles of eccDNA in OA via motif analysis and GO/KEGG pathways analysis.

Results: The chondrocytes from OA cartilage contained an abundance of eccDNAs, which was termed as OC-eccDNAs (OA Cartilage-derived eccDNA). The characteristics of OC-eccDNAs were tissue specific, including the distribution, the size range and sequence pattern. Moreover, the functional analysis indicated that eccDNA may be involved in the homeostasis maintenance of chondrocytes and participated in the process of OA.

Conclutions: Our data firstly showed the landscape of eccDNA in articular cartilage, and preliminarily indicated the potential roles of eccDNA in OA.

Key messages:

● **What is already known about this subject?**

1. The eccDNA can be detected in some normal and pathological tissues.

2. The eccDNA can regulate a number of cellular functions and affect cell phenotypes via different action modles.

3. The formation of eccDNA was tightly related to the abnormal repair of DNA damage, which was often induced by inflammation and aging.

● **What does this study add?**

1. An abundance of eccDNA were detected in articular cartilage, OA-related transcription factors may participate in the formation of OC-eccDNA.

2. The eccDNA may be involved in cartilage degeneration via multiple mechanisms, such asepigenetic regulation.

● **How might this impact on clinical practice or future developments?**

The eccDNA could be a new target for the treatment of cartilage-related diseases, such as OA. The roles and mechanisms of eccDNA in the physiology and pathology of OA cartilage should be further investigated.

Introduction

Osteoarthritis (OA) is a common degenerative disease of musculoskeletal system. The total number of patients with OA disease has exceeded 250 million in the world, and the number of patients is still rising rapidly and continuously[1]. With the aging of our society, the incidence rate of osteoarthritis is increasing, which will become the most important factor leading to the dysfunctions of the elderly and seriously affect the quality of life of the patients[2]. At present, the clinical therapeutic effect of OA is still not ideal. Therefore, it is urgent to strengthen the relevant research so as to provide new theoretical support for the prevention and treatment of OA.

OA is a total joint disease involving cartilage, subchondral bone and synovium[3]. Among them, cartilage injury is the core pathological change of OA and the basic element of OA disease diagnosis. A variety of biological factors such as aging, inflammation and mechanical stimulation will destroy the homeostasis of chondrocyte, and then aggravate OA progression[4]. The articular cartilage consists of chondrocytes and extracellular matrix, which mainly depends on the infiltration of joint fluid to maintain nutrition. As there is no distribution of blood vessels, nerves and lymphatic vessels, the repair and regeneration ability of articular cartilage after injury is very poor[5]. Until now, there are no effective measures to protect OA chondrocytes from degeneration. More studies on the homeostasis maintenance of chondrocytes in the process of OA needed to provide a new potential target for this disease.

Extrachromosomal circular DNA (eccDNA) refers to the circular DNA derived from genomic DNA but free of chromosomes. Recent studies have shown that eccDNA is extensively expressed in different tissues and may be involved in varied physiological and pathological processes including immune, inflammation, aging and tumors[6-9]. However, the expression of eccDNA in cartilage tissue and its pathological significance in OA are still unknown. Here, we extracted eccDNA from total knee arthroplasty (TKA) donors and firstly investigated the characteristics of eccDNA in articular cartilage using cir-DNA sequencing, and discussed the potential roles of eccDNA in OA.

Methods

Human Cartilage Tissue Collection and Isolation of Chondrocytes

Human cartilage was obtained from OA patients (n=10, aged 69.3 ± 6.53 years, four male and six females). Subjects with tumors, diabetes or other severe diseases in the last 5 years were excluded. Human articular cartilage explants were cultured according to the previously described methods[10]. Briefly, freshly obtained germ-free human cartilage was washed with PBS and cut into 4-mm-diameter blocks by the bistoury. Subsequently, the bocks were digested in high-glucose DMEM supplemented with 0.2% type II collagenase for 6h. Then the digestion suspension was collected and centrifuged at 400g for 5min following the filtration with a 40 μm cell strainer. Finally, the sediment (primary chondrocyte) was resuspended with high glucose DMEM containing 10% FBS and cultured in a 5% C02 incubator.

Circle-Seq

The method of Circle-Seq Chondrocyte eccDNA was optimized based on the description of Henrik Devitt Møller et.al, including of multiple steps showed below: 1) Total DNA was extracted using QIAamp DNA Mini Kit (Qiagen); 2) Then, total DNA were alkaline treated to separate chromosomal DNA, lipids, and protein by rapid DNA denaturing–renaturing, followed by column chromatography on an ion exchange membrane column (Plasmid Mini AX; A&A Biotechnology); 3) Remaining linear DNA was removed by exonuclease (Plasmid-Safe ATP-dependent DNase, Epicentre), assisted by rare-cutting endonuclease MssI (only support Homo sapiens) that digested mitochondrial circular DNA (mtDNA, 16 kb) and made additional accessible DNA ends for exonuclease; 4) EccDNA-enriched samples was used as template for phi29 polymerase reactions (REPLI-g Midi Kit) amplifying DNA; 5) Amplified circular DNA was cleaned (AMPure XP beads) and sheared by sonication (Bioruptor) to average fragment size of 200 –300 bp; 6) Libraries for next-generation sequencing were prepared using the NEBNext Ultra DNA Library Kit for Illumina according to the manufacturer's protocol (New England Biolabs), and subjected to sequencing on Illumina Novaseq 6000 using PE150, collecting up to 80 million paired-end

reads/sample.

EccDNA Mapped and identified Method for Circle-Seq Data

The sequencing reads from Circle-Seq were aligned to reference genome sequences

(GRChg38) using the BWA program (Burrows-Wheeler-Alignment Tool,

https://github.com/lh3/bwa) to dig eccDNA originated from the chromosome.

Subsequently, the candidate reads were sorted, labeled, and ranked by Circle-Map to

detect and identify eccDNA refers to the practices of Henrik Devitt Møller.

Results

cir-DNA sequencing revealed the landscape of eccDNA in articular cartilage

from OA patients.

Firstly, we harvested articular cartilage (n=10) from OA patients post-TKA operation

(supplemental Figure 1A) ，and collected eccDNA of chondrocytes for cir-DNA

sequencing (**Figure 1A** and supplemental Figure 1B). The quality of cir-sequencing

output reads was evaluated as excellent (supplemental Figure 1C and D). Then we

used BWA software to perform genomic alignment of reads with GRch38. For

mapped reads, we utilized Circle-Map software for eccDNA identification and

detected 39,342 eccDNA (detected eccDNA) (**Figure 1A**). Next, we developed a filter

standard of split reads> 1 to identify the detected eccDNA, and finally 15121 eccDNA

(identified eccDNA) were obtained (**Figure 1 B**). Subsequently, we analyzed the size

of the identified eccDNA in Figure 1 B. We found that the size distribution of

eccDNA in chondrocytes was between 0.1 kb and 10 kb, which was similar to the size

distribution of eccDNA in muscle tissue but was distinct from that of serum (<1kb)

and of tumor cells(1kb-1mb) (**Figure 1 C**). As to the distribution of eccDNA on

chromosomes, the results uncovered that chondrocyte eccDNA distributed in all other

regions except for Y chromosome and the short arm of chromosome 13,14,15, which

was similar to that in Hela cells, but a bit different from that of glioma cells and yeast

cells (**Figure 1 D**). In addition, previous studies have demonstrated that eccDNA may

derive from repetitive sequences on the genome[11]. Indeed, we found that most of the

reads corresponding to eccDNA of chondrocytes matched the repetitive sequences of

the genome, including long interspersed nuclear elements (LINEs, 19.2%), short

interspersed nuclear elements (SINEs_Alu, 18.8%), long terminal repeat (LTR,

11.3%), and a small number of telomere sequences (**Figure 1 E**), which was partly the

same with muscle tissue, but different from plasma[12]. Excluding the overlapping parts,

8915 genes were annotated, which included mRNA (6468), lncRNA (1075)),

multipletype (697), pseudogene (219), miRNA (1), and others (455) (**Figure 1F**).

These data indicate that articular cartilage contains an abundance of eccDNA, whose

size range and chromosome origin may be relatively specific.

OA-related transcription factors that involved in eccDNA cyclization

Previous studies have suggested that the junction point of eccDNA and its nearby area

played an important role in eccDNA formation. Therefore, we separately extracted the

200bp sequence from upstream and downstream of the junction point start site or the

stop site, combing with HOMER software to blast the extracted sequence with the

known motif and predict the de novo motif (**Figure 2A**). To explore the role of the

transcription factor in eccDNA formation and effects, we performed transcription

factor-based enrichment analysis on mapped known motifs and de novo motifs

respectively, and extracted the TOP20(based on p-value) collection of the two outputs

(**Figure 2B**, supplemental table 1 and 2). Interestingly, most of the transcription

factors with high p-value were highly related to the development and degeneration of

cartilage, such as MYB, STAT3, NKX, NRF2, ZNF215, FOX. Moreover, some

transcription factors including KLF4, NRF2, STAT3 and NF-kB have been proved to

be the key pathogenic factors in OA (**Figure 2B**). These results indicate that

OA-related transcription factors may be involved in the formation of eccDNA in

chondrocytes.

The GO/KEGG pathways analysis for the eccDNA in OA

Next, we performed GO analysis and KEGG analysis of the above genes (**Figure**

3A-B). The data suggests that eccDNA may be involved in the pathophysiological

regulation of degenerated chondrocytes in morphology, cell polarization, cell-ECM

interaction, etc. From the top enriched terms, we found that abundant genes were

involved in anabolism and catabolism, senescence, and inflammation of chondrocyte,

such as ACAN, MMPs, ADAMTSs, IL1B, and so on (**Figure 3C**), indicating that

eccDNA may participate in the regulation of cartilage homeostasis. Moreover, GO

and KEGG analysis both revealed that the target genes of eccDNA were significantly

enriched in nervous system-related pathways, such as axon guidance, synapse

organization, suggesting that a potential role of eccDNA in joint pain of OA. Besides,

we sorted the eccDNA according to circle score to analyze the genes annotated by

TOP20 eccDNA and found that these genes are mainly related to the formation of

nerve synapses (CNTN4, CABLES), cell stress response (SGK1), chromatin stability

and DNA damage repair (NT5DC1, PDS5A, STK11), protein degradation and

reparation (FBXW7, PCMT1, UBXN8), transcription inhibition (CBFA2T2, NCOR2),

cell adhesion and skeletal remodeling (MAGI1, ARHGEF26) (**Figure 3C-D**).

Furthermore, certain eccDNAs that potentially targeted OA-related genes from Figure

1J and 1K were analyzed and displayed (**Figure 3E**).

Discussion

In recent years, more and more studies have shown that eccDNA is closely

involved in the occurrence and development of several diseases such as cancer[9][13].

Turner, K. M. et al. reported that eccDNA was found in nearly half of human cancers

and its frequency varied by tumor type[9][13]. They also revealed that driver oncogenes in

eccDNA were amplified commonly, which contributed to the transcription of

oncogenes out of chromosomes[14]. Wu, S. et al. investigated the structure of eccDNA

by integrating ultrastructural imaging, long-range optical mapping and computational

analysis of whole-genome sequencing, and their data suggested that oncogenes

encoded on eccDNA increased copy number with high transcription levels[15]. In

addition, eccDNA displays significantly enhanced chromatin accessibility than typical

of chromosomes and has more ultra-long-range interactions with active chromatin [15],

indicating it possesses powerful transcriptional capacity potentially. Koche, R. P. and

their colleagues also found that cancer-causing lesions can increase the formation of

functional eccDNA in neuroblastoma, which may play an important role in tumor

progression[16]. Apart from tumor, eccDNA has also been related to aging process[17-19].

The accumulation of CUP1 eccDNA has unique-site specificity in the copper-induced

aging model of yeast[18]. Furthermore, the heterogeneity of circular DNA that offers

flexibility in adaptation is significantly diminished with age[19]. As aging is the main

cause of OA, it is speculated that eccDNA may play an important role in the

maintenance of chondrocyte homeostasis and the process of OA disease. Just like

tumor cells, Gene transcription on eccDNA (not on Chromosomal DNA) may be

actived in aged chondrocytes and be involved in OA process. As eccDNA comes from

genome, it may regulate chondrocyte phenotypes by epigenetic means, such as

affecting the replication and transcription of Chromosomal DNA. More studies are

needed to investigate the roles and mechanisms of eccDNA on chondrocyte.

Synovium is an important part of the joint sliding system, which is composed of

loose connective tissue. Its main functions include secreting synovial fluid, reducing

joint surface friction, preventing joint adhesion, providing nutrients for articular

cartilage, absorbing and swallowing various metabolites and cell fragments in the

joint cavity, etc. When the joint is subjected to stress injury (such as degeneration,

trauma, infection and rheumatism), the synovium will be stimulated to produce

inflammatory reaction and participate in the repair of articular cartilage injury. It is

found that chronic and low-grade inflammation is the main feature of OA synovitis, 6

which is closely involved in the formation of OA cartilage damage repair

microenvironment. Therefore, trying to regulate the state of synovitis to reshape the

microenvironment of OA joint, and then promote the damage and repair of

chondrocytes, is expected to become a feasible strategy for clinical OA treatment.

However, the formation mechanism of OA synovitis has not been very clear until now.

Previous studies have suggested that the fragments from dead chondrocytes and

degraded cartilage matrix can stimulate synovial tissue to produce inflammatory

response and mediate the formation of chronic synovitis in the process of OA[20]. In

addition, our group reported that the exosome-like vesicles from OA chondrocytes

with inflammatory stimulation could increase the mature IL-1β production of synovial

macrophages and aggravated synovitis in osteoarthritis[10]. Recently, Wang, Y. et al.

revealed that eccDNAs are apoptotic products with high innate immunostimulatory activity. Their data indicated that apoptosis inducers can increase eccDNA generation and eccDNAs can function as potent innate immunostimulants in a manner that is dependent on eccDNA circularity and the cytosolic DNA sensor Sting[6]. Therefore, we deduced that eccDNA from the apoptotic or senescent chondrocytes may participate in the formation of OA chronic synovitis via stimulating synovial inflammatory cells.

This study firstly mapped out the landscape of eccDNA in articular cartilage and preliminarily indicated the potential roles of eccDNA in OA, which may open up a new vision on the roles and mechanisms of eccDNA in joint diseases. Based on the biological characteristics of eccDNA, we speculate that it may play a potential role in cartilage-related diseases, such as OA. The roles and mechanisms of eccDNA in joint diseases need further research. In addition, the formation mechanisms of eccDNA in chondrocytes and its changes in the pathological process are also worthy of further attention.

Competing interests The authors declare that there are no personal, organizational or financial conflicts of interests.

Contributorship ZN, LK and YC conceived and designed the experiments. WX, TZ , YG, YW, JY, JY and LS performed experiments. YX, JL and XL provided expert advice. All authors analysed the data. WX, LK, XL and ZN wrote the paper.

Acknowledgements We are grateful to Ms. JiaLi for excellent technical assistance.

Funding This work was supported by National Natural Science Foundation of China (No. 81871817; No. 82002360; No. 81772330).

Ethical approval information All procedures and protocols were approved by the Ethics Committee with informed consent obtained (2018128) and carried out in accordance with standard operative procedures.

Data sharing statement The authors declare that all the data will be shared online after the paper was published.

References:

1. Safiri S, Kolahi AA, Smith E, et al. Global, regional and national burden of osteoarthritis 1990-2017: a
systematic analysis of the Global Burden of Disease Study 2017. *Ann Rheum Dis*
2020;79(6):819-28. doi: 10.1136/annrheumdis-2019-216515 [published Online First:
2020/05/14]
2. Sharma L. Osteoarthritis of the Knee. *N Engl J Med* 2021;384(1):51-59. doi:
10.1056/NEJMcp1903768 [published Online First: 2021/01/07]
3. Astephen Wilson JL, Kobsar D. Osteoarthritis year in review 2020: mechanics. *Osteoarthritis*
Cartilage 2021;29(2):161-69. doi: 10.1016/j.joca.2020.12.009 [published Online First:
2021/01/10]
4. Greene MA, Loeser RF. Aging-related inflammation in osteoarthritis. *Osteoarthritis Cartilage*
2015;23(11):1966-71. doi: 10.1016/j.joca.2015.01.008 [published Online First: 2015/11/03]
5. Huey DJ, Hu JC, Athanasiou KA. Unlike bone, cartilage regeneration remains elusive. *Science*
2012;338(6109):917-21. doi: 10.1126/science.1222454 [published Online First: 2012/11/20]
6. Wang Y, Wang M, Djekidel MN, et al. eccDNAs are apoptotic products with high innate
immunostimulatory activity. *Nature* 2021;599(7884):308-14. doi:
10.1038/s41586-021-04009-w [published Online First: 2021/10/22]
7. Wu SH, Turner KM, Nguyen N, et al. Circular ecDNA promotes accessible chromatin and high
oncogene expression. *Nature* 2019;575(7784):699-+. doi: 10.1038/s41586-019-1763-5
8. Prada-Luengo I, Moller HD, Henriksen RA, et al. Replicative aging is associated with loss of genetic
heterogeneity from extrachromosomal circular DNA in Saccharomyces cerevisiae. *Nucleic*
Acids Research 2020;48(14):7883-98. doi: 10.1093/nar/gkaa545
9. Henriksen RA, Jenjaroenpun P, Sjostrom IB, et al. Circular DNA in the human germline and its
association with recombination. *Mol Cell* 2022;82(1):209-17 e7. doi:
10.1016/j.molcel.2021.11.027 [published Online First: 2021/12/25]
10. Ni Z, Kuang L, Chen H, et al. The exosome-like vesicles from osteoarthritic chondrocyte enhanced
mature IL-1beta production of macrophages and aggravated synovitis in osteoarthritis. *Cell*
Death Dis 2019;10(7):522. doi: 10.1038/s41419-019-1739-2 [published Online First:
2019/07/10]
11. Paulsen T, Kumar P, Koseoglu MM, et al. Discoveries of Extrachromosomal Circles of DNA in Normal
and Tumor Cells. *Trends Genet* 2018;34(4):270-78. doi: 10.1016/j.tig.2017.12.010
12. Sin STK, Jiang P, Deng J, et al. Identification and characterization of extrachromosomal circular DNA
in maternal plasma. *Proc Natl Acad Sci U S A* 2020;117(3):1658-65. doi:
10.1073/pnas.1914949117 [published Online First: 2020/01/05]
13. Wu S, Bafna V, Mischel PS. Extrachromosomal DNA (ecDNA) in cancer pathogenesis. *Curr Opin*
Genet Dev 2021;66:78-82. doi: 10.1016/j.gde.2021.01.001 [published Online First:
2021/01/22]
14. Turner KM, Deshpande V, Beyter D, et al. Extrachromosomal oncogene amplification drives tumour
evolution and genetic heterogeneity. *Nature* 2017;543(7643):122-25. doi:
10.1038/nature21356 [published Online First: 2017/02/09]
15. Wu S, Turner KM, Nguyen N, et al. Circular ecDNA promotes accessible chromatin and high
oncogene expression. *Nature* 2019;575(7784):699-703. doi: 10.1038/s41586-019-1763-5
[published Online First: 2019/11/22]
16. Koche RP, Rodriguez-Fos E, Helmsauer K, et al. Extrachromosomal circular DNA drives oncogenic
genome remodeling in neuroblastoma. *Nat Genet* 2020;52(1):29-34. doi:

10.1038/s41588-019-0547-z [published Online First: 2019/12/18]

17. Ain Q, Schmeer C, Wengerodt D, et al. Extrachromosomal Circular DNA: Current Knowledge and
Implications for CNS Aging and Neurodegeneration. *Int J Mol Sci* 2020;21(7) doi:
10.3390/ijms21072477 [published Online First: 2020/04/08]

18. Hull RM, King M, Pizza G, et al. Transcription-induced formation of extrachromosomal DNA during
yeast ageing. *PLoS Biol* 2019;17(12):e3000471. doi: 10.1371/journal.pbio.3000471 [published
Online First: 2019/12/04]

19. Prada-Luengo I, Moller HD, Henriksen RA, et al. Replicative aging is associated with loss of genetic
heterogeneity from extrachromosomal circular DNA in Saccharomyces cerevisiae. *Nucleic*
Acids Res 2020;48(14):7883-98. doi: 10.1093/nar/gkaa545 [published Online First:
2020/07/02]

20. Mathiessen A, Conaghan PG. Synovitis in osteoarthritis: current understanding with therapeutic
implications. *Arthritis Res Ther* 2017;19(1):18. doi: 10.1186/s13075-017-1229-9 [published
Online First: 2017/02/06]

Figure 1

Figure 2

Figure 3