1   **Role of posterodorsal medial amygdala urocortin-3 in pubertal timing in female mice**

Deyana Ivanova[1], Xiao-Feng Li[1], Yali Liu[2], Caitlin McIntyre[1], Cathy Fernandes[3,4], Geffen

Lass[1], Lingsi Kong[1] and Kevin T O'Byrne[1]

[1]Department of Women and Children's Health, Faculty of Life Science and Medicine, King's

College London, UK

[2]Department of Assisted Reproduction, Shanghai Ninth People's Hospital, Shanghai Jiaotong

University School of Medicine, Shanghai, People's Republic of China.

[3]Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology &

Neuroscience, King's College London, UK

[4] MRC Centre for Neurodevelopmental Disorders, King's College London, UK

Corresponding author's contact details: Deyana Ivanova (deyana.ivanova@kcl.ac.uk;

https://orcid.org/0000-0003-1508-166X) and Kevin T O'Byrne (kevin.obyrne@kcl.ac.uk;

https://orcid.org/0000-0002-2548-4182), Department of Women and Children's Health,

School of Life Course and Population Sciences, Faculty of Life Science and Medicine, King's

College London, 2.92W Hodgkin Building, Guy's Campus, London SE1 1UL, UK.

Reprint requests: Deyana Ivanova, Department of Women and Children's Health, School of

Life Course Sciences, Faculty of Life Science and Medicine, King's College London, 2.92W

Hodgkin Building, Guy's Campus, London SE1 1UL, UK.

Abstract

Post-traumatic stress disorder impedes pubertal development and disrupts pulsatile LH secretion in humans and rodents. The posterodorsal sub-nucleus of the medial amygdala (MePD) is an upstream modulator of the hypothalamic gonadotropin-releasing hormone (GnRH) pulse generator, pubertal timing, as well as emotional processing and anxiety. Psychosocial stress exposure alters neuronal activity within the MePD increasing the expression of Urocortin3 (Ucn3) and its receptor corticotropin-releasing factor type-2 receptor (CRFR2) while enhancing the inhibitory output from the MePD to key hypothalamic reproductive centres. We test the hypothesis that psychosocial stress, processed by the MePD, is relayed to the hypothalamic GnRH pulse generator to delay puberty in female mice. We exposed C57Bl6/J female mice to the predator odor, 2,4,5-Trimethylthiazole (TMT), during pubertal transition and examined the effect on pubertal timing, pre-pubertal LH pulses and anxiety-like behaviour. Subsequently, we virally infected Ucn3-cre-tdTomato female mice with stimulatory DREADDs targeting MePD Ucn3 neurons and determined the effect on pubertal timing and pre-pubertal LH pulse frequency. Exposure to TMT during pubertal development delayed puberty, suppressed pre-pubertal LH pulsatility and enhanced anxiety-like behaviour, while activation of MePD Ucn3 neurons reduced LH pulse frequency and delayed puberty. Early psychosocial stress exposure decreases GnRH pulse generator frequency delaying puberty while inducing anxiety-behaviour in female mice, an effect potentially involving Ucn3 neurons in the MePD.

Introduction

In humans post-traumatic stress disorder (PTSD) is associated with altered pubertal timing and the development of anxiety disorders (1,2). Predator odor exposure is a classic model for PTSD in rodents (3) and has been shown to delay puberty (4), suppress luteinizing hormone (LH) pulse frequency (5) and inhibit the pre-ovulatory LH surge (6). The gonadotropin-releasing hormone (GnRH) pulse generator that controls the hypothalamus-pituitary-gonadal (HPG) axis and pubertal development is restrained in the juvenile, but reactivated at a critical time point with increased frequency leading to the initiation of puberty (7). However, the neural mechanisms controlling pubertal timing are not fully established.

Kisspeptin (kiss1) is known to be a major gatekeeper of pubertal onset. An absence of the *kiss1* gene and kiss1 receptor (kiss1r) in mice and humans leads to a loss of gonadal maturation and a lack of pubertal onset (8,9), and treatment with kisspeptin can reactivate the HPG axis during the juvenile hiatus in monkeys and rats (10,11). Kiss1 neurons are located in the anteroventral periventricular nucleus (AVPV) and arcuate nucleus (ARC) of the hypothalamus in mice (12). The ARC kiss1 neurons, known as KNDy because they co-express neurokinin B (NKB) and dynorphin A (Dyn), are a critical component of the GnRH pulse generator (12–14). The activation of KNDy neurons induces pulsatile release of kiss1 acting on GnRH dendrons in the median eminence to stimulate pulsatile GnRH secretion (15). The AVPV kiss1 neurons project to GnRH cell bodies and proximal dendrites and are known to control the preovulatory LH surge and ovulation in rodents (16). In the juvenile, a neurobiological break exerted on the ARC GnRH pulse generator suppresses pulsatile kiss1 release in the median eminence, however the nature of this upstream inhibition of the GnRH pulse generator, which is released at the end of the juvenile phase to trigger puberty is unknown (7).

The amygdala, a stress-sensitive part of the limbic brain, is involved in processing anxiety and fear as well as modulating pubertal timing and the HPG axis. Lesioning of the amygdala

advances menarche in female rhesus macaques (17), whereas electrical stimulation of this

region delays puberty in rats (18). Specific lesioning of the posterodorsal sub-nucleus of the

medial amygdala (MePD) advances puberty in rats (19), suggesting this region may play a key

role in exerting an inhibitory break on pubertal timing. The MePD sends GABAergic

projections to various hypothalamic reproductive centres (20,21) and has been shown to project

directly to ARC KNDy neurons, although their neurochemical phenotype is undetermined

(22,23). Interestingly, the MePD contains a kiss1 neuronal population (24) and antagonism of

MePD kiss1 delays puberty, disrupts estrous cyclicity and the LH surge in rats (25). Moreover,

selective optogenetic stimulation of MePD kiss1 neurons increases LH pulse frequency in mice

(26). The MePD is highly responsive to psychosocial stress, exhibiting a distinct firing pattern

in response to predator odor (27) and recently cat odor was found to delay puberty, disrupt

estrous cyclicity and induce a fear response in rats (4). The MePD may be a central hub

involved in the integration of external olfactory and anxiogenic signals with the GnRH pulse

generator.

Central administration of the stress neuropeptide corticotrophin releasing factor (CRF) dose-

dependently delays puberty, while CRF-receptor antagonism advances puberty in rats (28).

Urocortin-3 (Ucn3), a member of the CRF family and an endogenous ligand for CRF type 2

receptors (CRFR2) is found in abundance in the MePD (29). Psychosocial stressors, including

social defeat and restraint, increase cfos expression in CRFR2 positive neurons and increase

Ucn3 mRNA expression in the MePD of rodents (30). Moreover, we have recently shown that

MePD Ucn3 mediates predator odor and restraint stress-induced suppression of pulsatile LH

secretion in mice (5).

In this study, we aimed to determine whether chronic predator odor stress exposure suppresses

pre-pubertal LH pulse frequency and delays pubertal timing in female mice. Additionally, we

assessed social, anxiety and fear-like behaviour as measures of psychosocial stress in predator

odor-exposed pre-pubertal female mice. Finally, we aimed to determine whether MePD Ucn3

signalling is involved in modulating pre-pubertal LH pulse frequency and pubertal timing in

female mice.

Materials and Methods

Mice

Breeding pairs of C57Bl6/J mice were purchased from Charles River Laboratories

International, Inc. For Ucn3-cre mice, cryopreserved sperm of strain (Tg(Ucn3-

cre)KF43Gsat/Mmucd; congenic on C57BL/6 background) was acquired from MMRRC

GENSAT. Breeding pairs of heterozygous transgenic Ucn3-cre mice were recovered by

insemination of female C57Bl6/J mice at King's College London. Genotyping of Ucn3-cre

mice was performed using PCR to detect heterozygosity. Heterozygous Ucn3-cre mice were

bred with cre-activated tdTomato reporter mice (strain B6.Cg-Gt(ROSA)26Sortm9(CAG-

tdTomato)Hze/J; congenic on C57BL/6 background obtained from The Jackson Laboratory,

Bar Harbor, ME, USA) to obtain Ucn3-cre-tdTomato mice, as previously described (5,29).

Litter size was reduced to 6-8 pups 2-3 days after birth (on pnd 2-3) to standardise body weight,

which can alter pubertal development. Mice aged between 21 to 45 days were group housed

in individually ventilated cages equipped with wood-chip bedding and nesting material with

food and water ad libitum and sealed with a HEPA-filter at 25 ± 1 °C in a 12:12 h light/dark

cycle, lights on at 07:00 h. All procedures were carried out following the United Kingdom

Home Office Regulations and approved by the Animal Welfare and Ethical Review Body

Committee at King's College London.

Stereotaxic adeno-associated-virus injection

All surgical procedures were carried out with aseptic conditions and under general anaesthesia using ketamine (Vetalar, 100 mg/kg, i.p.; Pfizer, Sandwich, UK) and xylazine (Rompun, 10 mg/kg, i.p.; Bayer, Leverkusen, Germany). The mouse brain atlas of Paxinos and Franklin (31) was used to obtain target coordinates for the MePD (2.15 mm lateral, -1.25 mm from bregma, at a depth of -5.30 mm below the skull surface) of pre-pubertal mice. Mice at pnd 14 were temporarily taken from their mother and secured in a David Kopf stereotaxic frame (Kopf Instruments), a small skin incision was made to reveal the skull and two small holes were drilled above the location of the MePD. Bilateral stereotaxic viral injections of the stimulatory adeno-associated-virus (AAV) carrying the DIO-hM3D-mCitrine, DREADD, construct (AAV-hSyn-DIO-HA-hM3D(Gq)-IRES-mCitrine, 3×10^{11} GC/ml, Serotype:5; Addgene) was administered intra-MePD using the robot stereotaxic system (Neurostar, Tubingen, Germany), performed for the targeted expression of DIO-hM3D-mCitrine in MePD Ucn3 neurons in Ucn3-cre-tdTomato mice. AAV-hSyn-DIO-HA-hM3D(Gq)-IRES-mCitrine (150 nl) was bilaterally injected into the MePD using a 2-μl Hamilton micro syringe (Esslab, Essex, UK) over 10 min and the needle was left in position for a further 5 min then slowly lifted over 2 min. Mice that were cre-positive received the AAV-hM3D injection (test mice) or a control virus AAV-YFP (Addgene) (control mice) where the control virus does not contain the DIO-hM3D-mCitrine construct. The mice were placed back with the mother for a further 7 days until wean day (pnd 21). Nine Ucn3-cre-tdTomato mice received the AAV-hM3D injection, 6 Ucn3-cre-tdTomato mice received the control AAV-YFP, 4 Ucn3-cre-negative mice without surgery received only CNO and 3 Ucn3-cre-tdTomato mice without surgery did not receive CNO.

Chronic pre-pubertal predator-odor exposure and puberty evaluation

Female pups were weaned on pnd 21 and separated into control and test groups. To investigate the effects of chronic pre-pubertal psychosocial stress mice were removed from their home cages, singly housed in new cages and left to habituate for 10 min. After the habituation period, the mice were exposed to 12 μl of 2,4,5-Trimethylthiazole (TMT; synthetic extract of fox urine; ≥98% purity; Sigma-Aldrich, UK) pipetted on a small circular piece of filter paper in a petri dish placed in the centre of the cage for 20 min. The mice were exposed to TMT daily, at random time points, for 14 days (from pnd 21 to 35). Control mice were exposed to filter paper soaked with 12 μl of ddH$_2$O water. To rule out the possibility that physiological and behavioural responses in TMT-exposed mice resulted from novelty of scent we repeated the experiment with 85 mM of ethyl vanillin (≥98% purity; Sigma-Aldrich UK) as a control scent in a separate cohort of mice. Mice were monitored from pnd 24 for vaginal opening (VO) and first estrous (FE) indicated by epithelial cell cornification. Once the occurrence of VO was detected, vaginal smears were taken to determine the exact day of FE.

Pre-pubertal blood sampling

The effect of TMT exposure on pre-pubertal LH pulses was measured, on pnd 26 and 29. Mice were handled twice daily for 10 min to habituate to tail-tip blood collection for at least 7 days prior to blood sampling, as described previously (5). For LH measurement, 3 μl of blood was collected every 5 min for 80 min. Blood sampling was performed on pnd 26 and 29 during the period of daily TMT exposure (pnd 21-35). Blood samples were collected at least 2 h before TMT-exposure on these days. Blood collection was performed between 09:00-12:00 h.

For Ucn3 DREADDs experiments, blood sampling was performed on pnd 29 during the period of daily CNO administration (pnd 21-35), as described above. Blood collection was performed between 09:00-12:00 h.

Behavioural tests

Light Dark Box

The same cohort of C57Bl6/J female mice were transported to a new room (lights off) and left to habituate for 30 min. Mice were placed in the centre of the light compartment facing towards the dark compartment. The time spent in each compartment (entry with all 4 limbs) was recorded manually over a period of 5 min by two independent observers. The LDB test was carried out on pnd 19, 27 and 40. Behaviour testing performed during the period of daily TMT exposure (pnd 21-35) and was measured before the exposure to TMT on that day. This applies to all behaviour tests performed during the TMT exposure period. All behaviour tests were performed between 11:00-13:00 h.

Social interaction with familiar conspecific

The same cohort of C57Bl6/J female mice were transported to a new dimly lit room and left to habituate for 30 min. Experimental mice and a same sex familiar conspecific (similar age, weight and strain) were placed simultaneously into the test arena (a new cage with clean wood-chip bedding). The time spent following, sniffing, grooming and mounting the conspecific was monitored and recorded manually over a period of 5 min by two independent observers. The social interaction test was carried out on pnd 19, 27 and 40.

Elevated Plus maze

The same cohort of C57Bl6/J female mice were transported to a new room (lights on) and left to habituate for 30 min. Mice were placed in the centre of the maze facing towards the closed arms. The time spent in each part of the maze (open, centre and closed; entry with all 4 limbs) was recorded manually over a period of 5 min by two independent observers. The EPM test was carried out on pnd 20, 28 and 41.

Chronic DREADD activation of Ucn3 neurons in the MePD

For DREADDs experiments, stock solution of Clozapine-N-oxide (CNO) (Tocris Bio-techne,

Abingdon, UK) was made by dissolving 5 mg of CNO in 1 ml of 0.9% sterile saline solution

and stored at 4 °C. CNO was made fresh daily and administered via drinking water at a final

concentration of 0.5 mg CNO/ kg, as described previously (32) for 14 days from pnd 21-35.

Mice were monitored from pnd 22 for vaginal opening (VO) and first estrous (FE), as described

above.

Validation of AAV injection

On pnd 50, Ucn3-cre-tdTomato mice were anaesthetised with a lethal dose of ketamine.

Transcardial perfusion was performed with heparinised saline for 5 min followed by ice-cold

4% paraformaldehyde (PFA) in phosphate buffer (pH 7.4) for 15 min with a pump (Minipuls,

Gilson, Villiers Le Bel, France). Brains were immediately collected and fixed in 15% sucrose

in 4% PFA at 4 °C and left to sink. Brains were then transferred to 30% sucrose in phosphate-

buffered saline (PBS) and left to sink. Brains were snap-frozen in isopropanol on dry ice and

stored in -80°C. Every third coronal brain section, 30-μm/section, was collected using a

cryostat (Bright Instrument Co., Luton, UK) through-out the MePD region corresponding to -

1.34 mm to -2.70 mm from bregma. Brain sections were mounted on microscope slides, air

dried and covered with ProLong Antifade mounting medium (Molecular Probes, Inc. OR,

USA). The number of tdTomato labelled and AAV-hM3D-mCitrine infected Ucn3 neurons

per side per slice in the MePD was quantified from 4 slices. For AAV-hM3D-mCitrine injected

Ucn3-cre-tdTomato mice we determined whether Ucn3 neurons were infected in the MePD

region by merging td-Tomato fluorescence of Ucn3 neurons with mCitrine fluorescence in the

MePD. Images were taken using Axioskop 2 Plus microscope (Carl Zeiss) equipped with

axiovision, version 4.7 (Carl Zeiss). Only data from animals with correct AAV injection were

analysed.

LH pulse detection and analysis

Blood samples were processed with a LH ELISA, as previously reported (33). Capture

antibody (monoclonal antibody, anti-bovine LHβ subunit, AB_2665514) was purchased from

Department of Animal Science at the University of California, Davis. Mouse LH standard

(AFP-5306A) and primary antibody (polyclonal antibody, rabbit LH antiserum, AB_2665533)

were obtained from Harbour-UCLA (California, USA). Secondary antibody (Horseradish-

Peroxidase (HRP)-linked donkey anti-rabbit IgG polyclonal antibody, AB_772206) was

purchased from VWR International (Leicestershire, UK). Inter-assay and intra-assay

variations were 4.6% and 10.2%, respectively and the assay sensitivity was 0.0015 ng/mL.

ODs of the standards were plotted against the log of the standard concentrations, non-linear

regression to fit the points and parameters were extracted to calculate the concentration of LH

(ng/ml) in blood samples, as previously described (33). The LH concentration at every time

point of blood collection was plotted as a line and scatter graph using Igor Pro 7, Wavemetrics,

Lake Oswego, OR, USA. DynPeak algorithm was used for the detection of LH pulses (34).

Statistics

For TMT-exposed and DREADD injected female pre-pubertal mice, data obtained for VO and

FE was analysed using RM one-way ANOVA. Data obtained for performance on LDB, SI and

EPM, LH pulse frequency and body weight measure were analysed using RM two-way

ANOVA. Statistics were performed using Igor Pro 7, Wavemetrics, Lake Oswego, OR, USA.

Data was represented as mean ± SEM and $+p<0.05$, $++p<0.001$ and $+++p<0.0001$ were

considered to be significant.

Results

Chronic TMT exposure delays puberty onset

TMT exposure during the pubertal transition period, pnd 21 to 35, delayed FE without altering VO compared water and ethyl vanillin controls in C57Bl6/J female mice (Fig. 1, A and B; Control vs TMT, +++$p<0.0001$; TMT, n=14, water control, n=6, ethyl vanillin control, n=4). Data for water and ethyl vanillin exposed control mice were combined as control since there was no significant difference between the two control groups. Body weight was unaffected between the experimental groups (Fig. 1, C; Control vs TMT). The data from this study shows that predator-odor stress exposure during the pubertal transition delays pubertal onset.

Chronic TMT exposure decreases pre-pubertal LH pulse frequency

TMT exposure during the pubertal transition period, pnd 21 to 35, suppressed pre-pubertal LH pulse frequency on pnd 26 and 29 compared to water and ethyl vanillin controls in C57Bl6/J female mice (Fig. 2, A-E; Control vs TMT, ++$p<0.001$; TMT, n=8, water control, n=6, ethyl vanillin control, n=3). On pnd 29, the average LH pulse frequency for the control group tended to increase compared to the control group on pnd 26 potentially marking an acceleration of GnRH pulse generator activity approaching the onset of puberty (Fig. 2, A, C and E). The results of this experiment are summarised in the figure 2E. Data for water and ethyl vanillin exposed control mice were combined as control since there was no significant difference between the two control groups. The data from this study shows that predator-odor stress exposure during the pubertal transition inhibits pre-pubertal LH pulsatility.

Chronic TMT exposure induced long-lasting anxiety and fear-like behaviour

We investigated anxiety and fear-like behaviour induced by TMT-exposure during the pubertal transition period, pnd 21 to 35 in in C57Bl6/J female mice. Anxiety and fear-behaviour were

assessed using standard anxiety and fear tests, the LDB, SI and EPM, on pnd 19 or 20 (prior to

day of weaning and TMT-exposure), pnd 27 or 28 (during the TMT exposure period) and pnd

40 or 41 (a week after termination of TMT-exposure). The first LDB, SI and EPM trial on pnd

19 or 20, prior to weaning, confirms no group difference in anxiety before the beginning of

TMT-exposure (Fig. 3, A-C). TMT-exposed mice spent significantly less time in the light

compartment of the LDB as well as socialising with a familiar conspecific compared to the

control group on pnd 27, but not on pnd 40 (Fig. 3, A, B; Control vs TMT, +$p<0.05$; TMT,

n=14, water control, n=6, ethyl vanillin control, n=4). TMT-exposed mice spent less time in

the open arms of the EPM on pnd 28 and pnd 41 compared to controls (Fig. 3, C; Control vs

TMT, ++$p<0.001$; TMT, n=14, water control, n=6, ethyl vanillin control, n=4). It is

acknowledged that repeat testing on the EPM assesses phobia/fear (35). On pnd 20, pnd 28

and 41, within-group analysis showed no significant differences in the total time spent in the

open arms of the EPM across the different test days in the control group. However, within-

group analysis of the TMT-exposed group showed a significant difference in time-spent in the

open arm of the EPM on pnd 41 compared to pnd 28 (Fig. 3, C; TMT on pnd 28 vs TMT on

pnd 41, ###$p<0.0001$). TMT-exposure during pubertal transition had a long-lasting effect on

anxiety and fear/phobia.

Selective expression of DREAD(Gq) in MePD Ucn3 neurons

Evaluation of m-Citrine, hM3D, expression in tdTomato labelled neurons from AAV-injected

Ucn3-cre-tdTomato mice revealed that $86 \pm 5\%$ of MePD Ucn3 neurons expressed hM3D and

the number of tdTomato labelled Ucn3 neurons per side per slice in the MePD was counted at

72.90 ± 5.48 (mean $\pm$ SEM) with the number of AAV-hM3D-mCitrine infected neurons being

63.00 ± 6.83 (mean $\pm$ SEM) (n=8). A representative example is shown in Fig 4, A-F.

Selective DREADD activation of Ucn3 neurons in the MePD delays puberty onset

Bilateral DREADD activation of MePD Ucn3 neurons during the pubertal transition period, pnd 21 to 35, of Ucn3-cre-tdTomato mice delayed FE without altering VO compared to non-surgery and AAV-YFP controls (Fig. 5, A and B; Control vs DREADD, $+p<0.05$; DREADD, n=8, control AAV, n=6, cre-negative control, n=4, no CNO, n=3). Data for cre-negative control, no CNO and control AAV-YFP injected mice were combined as control since there was no significant difference between the control groups. Body weight was unaffected between the experimental groups (Fig. 5, C; Control vs DREADD). These data show that activation of MePD Ucn3 during the pubertal transition delays pubertal timing. Mice with misplaced injections (n=1) were excluded from the analysis.

DREADD activation of Ucn3 neurons in the MePD suppresses pre-pubertal LH pulse frequency

Bilateral DREADD activation of MePD Ucn3 neurons during the pubertal transition period, pnd 21 to 35, of Ucn3-cre-tdTomato mice suppressed pre-pubertal LH pulse frequency sampled on pnd 29 compared to controls (Fig. 6, A, B and C; Control vs DREADD, $+p<0.05$; DREADD, n=5, control AAV, n=3, cre-negative control, n=3). Data for cre-negative control and control AAV-YFP injected mice were combined as control since there was no significant difference between the control groups. These data show that activation of MePD Ucn3 during pubertal transition suppresses pre-pubertal pulsatile LH secretion. Mice with misplaced injections (n=1) were excluded from the analysis.

Discussion

The present study shows for the first time that chronic exposure to the psychogenic stressor, predator odor, from pnd 21 for 14 days suppresses the pubertal acceleration of LH pulse

frequency, delaying puberty onset, while concurrently inducing long-lasting anxiety and fear-

like behaviour. Moreover, DREADD activation of MePD Ucn3 neurons during this same

developmental period delays puberty onset while suppressing pre-pubertal LH pulse frequency,

demonstrating that MePD Ucn3 signalling may play an important role in stress-related changes

in pubertal timing.

Hypothalamic kiss1 signalling is pivotal in regulating puberty onset in various species,

including rodents and humans (7,11). Pubertal onset is thought to be timed by an increase in

GnRH pulse generator frequency, however the mechanisms underlying the timing of ARC

kiss1 neuronal activation to trigger pubertal onset are not well established. Nevertheless,

puberty onset has been associated with increased ARC Tac2 (encoding NKB) and kiss1 mRNA

expression in mice (36,37), which are critical components of the GnRH pulse generator and

there is an increase in ARC kiss1 promoter activity during pubertal development (38). The

amygdala is known to influence pubertal timing in primates (17), and more specifically, the

MePD has been shown to regulate pubertal timing and reproductive function in rodents (4,19),

vis-à-vis MePD kiss1 receptor antagonism delays puberty, reduces the occurrence of

preovulatory LH surges in rats (25) and MePD kiss1 neuronal activation heightens sexual

partner preference in male mice (39). Recently, we have shown optogenetic stimulation of

MePD kiss1 neurons increases GnRH pulse generator frequency in adult female mice (26) and

preliminary data from our lab shows that activation of MePD kiss1 neurons during pubertal

transition advances the onset of first estrous, thus MePD kiss1 neurons may be an upstream

regulator of pubertal timing (40).

We explored the effect of chronic psychological stress exposure on pubertal timing, pre-

pubertal GnRH pulse generator activity and anxiety-like behaviour. Stress exposure during the

juvenile period is known to alter pubertal timing and induce anxiety behaviour in mammals,

including rodents, primates and humans (1,2,4,41). Predator odor exposure in rodents is a

model for PTSD; increasing fear and anxiety behaviour as well as reducing social interaction

(3). Previously, we have shown that acute TMT exposure suppresses LH pulsatility (5) and

restraint stress, another psychological stressor, inhibits LH pulse frequency while reducing

ARC kiss1 cfos expression in adult mice (42). Moreover, we have shown that cat odor delays

puberty and induces a fear response in rats (4). In the present study we show that the predator

odor, TMT, reduces LH pulsatility and delays the first estrous in female mice. In mice, VO

and FE do not coincide, wherein VO is not tightly coupled to the first ovulation, but rather a

variable time gap exists between VO and FE; the latter associated with first ovulation and

considered the unequivocal marker of puberty in this species (43). Additionally, pre-pubertal

LH pulse frequency in the control group tended to increase potentially marking the acceleration

of the GnRH pulse generator as the time of puberty onset approaches. Contrastingly, pre-

pubertal LH pulse frequency in the TMT-exposed group was significantly lower compared to

controls and we did not observe an increase in GnRH pulse generator activity between pnd 26

and 29.

Puberty is a sensitive period where enhanced neuroplasticity provides a context for the impact

of stress on the development of long-term anxiety disorders in humans (44). Early-life stress

increases anxiety-like behaviour in juvenile and adult mice linked to hyper-connectivity

between fronto-limbic circuits, involving the amygdala (45). The medial amygdala is activated

in response to external stressors and electrophysiological recordings reveal that predator urine

robustly activates the MePD, which is associated with delaying pubertal onset (4) and

supressing LH pulsatility in rodents (5,46). We found that predator odor-induced pubertal

delay was accompanied by increased anxiety-like behaviour, with mice spending significantly

less time in the light compartment of the LDB and in the open arms of the EPM during stress

exposure. Moreover, we observed that mice exposed to TMT exhibited long lasting fear-like

behaviour where they spent significantly less time in the open arm of the EPM compared to

controls on pnd 41; 6 days after termination of stress exposure. This is consistent with our

previous studies showing rats exposed to cat odor during pubertal development display

increased long-lasting fear-like behaviour on the EPM even after the termination of stress

exposure (4). Moreover, exposure to fox odor induces long-lasting anxiety on the EPM in rats

(47), while mice exposed to foot-shock stress during adolescence exhibit increased startle

reflexes into adulthood indicative of lasting anxiety-like behaviour (48). Additionally, TMT-

exposed mice showed significantly reduced social interaction with familiar con-specifics

during stress exposure whereas sociability tended to increase along with reproductive

maturation in the control group, which is consistent with our previous observations where

exposure to cat odor during puberty decreased sociability in rats (4).

Central administration of CRF inhibits kiss1 expression in the ARC and POA in rats (49). In

women with functional hypothalamic amenorrhea there is a correlation between decreased

serum kisspeptin and increased CRF concentrations (50). Moreover, an endogenous CRF tone

regulates pubertal timing where central administration of CRF in pnd 28 rats delays puberty

onset and administration of astressin-B, a CRFR1 antagonist, advances puberty (28). Central

antagonism of CRFR2 blocks the suppressive effect of stress on LH pulsatility in rodents (51).

These data indicate stress neuropeptides may provide an inhibitory tone on the timing of

puberty. Predator odor activates CRFR2 positive neurons in the medial amygdala of rats (52)

and we have recently shown that TMT-induced suppression of LH pulsatility is mediated by

MePD Ucn3 and CRFR2 signalling in mice (5). In the present study, selective activation of

MePD Ucn3 neurons during pubertal development delayed puberty and suppressed pre-

pubertal LH pulse frequency, demonstrating that MePD Ucn3 signalling plays a key role in

regulating pubertal timing.

The MePD has a major GABAergic output to key hypothalamic reproductive nuclei (21,53)

and we know the amygdala exerts an inhibitory brake on pubertal timing (17,19). Ucn3

neurons in the medial amygdala have an interneuron-like appearance (54) where Ucn3 fibres

overlap with CRFR2 expression (55,56) and we have shown that antagonism of MePD CRFR2

blocks the suppressive effect of TMT on LH pulse frequency in adult mice (5), thus Ucn3

neurons possibly connect to and signal via CRFR2 within the MePD. Moreover, the majority

of MePD CRFR2 neurons co-express GAD65 and 67 (29) indicating they are GABAergic and

may be involved in mediating stress-induced suppression of the GnRH pulse generator (22,53).

Therefore, MePD Ucn3 neurone activation may modulate pubertal timing possibly via

enhancing GABAergic output to inhibit GnRH pulse generator activity (5,21,23).

Our findings show for the first time that early life exposure to a psychosocial stressor disrupts

GnRH pulse generator frequency to delay puberty, and Ucn3 signalling in the MePD may be

involved in mediating this response. These findings provide novel insight into the key

interactions between the emotional stress centres and reproductive centres in the brain that

control pubertal onset in response to the external environment. Alterations in the timed onset

of puberty has major implications later in life with increased risk of diverse adverse outcomes,

including cancer, gynaecologic/obstetric, cardio-metabolic and neuro-cognitive categories in

humans (57). Understanding the mechanisms involved in controlling pubertal timing and the

impact of stress is crucial to ultimately aid future development of more effective treatments for

stress-related disorders of puberty.

Acknowledgements

The authors gratefully acknowledge the financial support from UKRI: BBSRC

(BB/S000550/1) and MRC (MR/N022637/1). DI is a PhD student funded by MRC-DTP

studentship at King's College London.

Funding

Grants supporting paper: Financial support from UKRI: BBSRC (BB/S000550/1) and MRC (MR/N022637/1). DI is a PhD student funded by MRC-DTP studentship at King's College London.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Data Availability Statement

The original contributions presented in the study are included in the article/supplementary material. Further inquiries can be directed to the corresponding author.

Ethics Statement

The animal study was reviewed and approved by the Animal Welfare and Ethical Review Body Committee at King's College London.

Bibliography

1. Ponnapakkam A, Gensure R. Effects of stress after Hurricanes Katrina and Rita on pubertal disorders in children. Ochsner J. 2008;8(3):129–33.

2. Gur RE, Moore TM, Rosen AF, Barzilay R, Roalf DR, Calkins ME, Ruparel K, Scott JC, Almasy L, Satterthwaite TD, Shinohara RT . Burden of Environmental Adversity Associated with Psychopathology, Maturation, and Brain Behavior Parameters in Youths. JAMA Psychiatry. 2019;76(9):966–75.

3. Zoladz PR, Diamond DM . Predator-based psychosocial stress animal model of PTSD:

Preclinical assessment of traumatic stress at cognitive, hormonal, pharmacological,

cardiovascular and epigenetic levels of analysis. Exp Neurol. 2016;284(Pt B):211–9.

4.   Li XF, Adekunbi DA, Alobaid HM, Li S, Pilot M, Lightman SL, O'Byrne KT . Role

of the posterodorsal medial amygdala in predator odour stress-induced puberty delay

in female rats. J Neuroendocrinol. 2019;31(6):e12719.

5.   Ivanova D, Li XF, McIntyre C, Liu Y, Kong L, O'Byrne KT . Urocortin3 in the

Posterodorsal Medial Amygdala Mediates Stress-induced Suppression of LH

Pulsatility in Female Mice. Endocrinology. 2021;162(12):bqab206.

6.   Wagenmaker ER, Moenter SM. Exposure to acute psychosocial stress disrupts the

luteinizing hormone surge independent of estrous cycle alterations in female mice.

Endocrinology. 2017;158(8):2593–602.

7.   Terasawa E, Guerriero KA, Plant TM. Kisspeptin and Puberty in Mammals. Adv Exp

Med Biol. 2013;784:253–73.

8.   De Roux N, Genin E, Carel JC, Matsuda F, Chaussain JL, Milgrom E.

Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived peptide

receptor GPR54. Proc Natl Acad Sci U S A. 2003;100(19):10972–6.

9.   Seminara SB, Messager S, Chatzidaki EE, Thresher RR, Acierno Jr JS, Shagoury JK,

Bo-Abbas Y, Kuohung W, Schwinof KM, Hendrick AG, Zahn D . The GPR54 gene as

a regulator of puberty. N Engl J Med. 2003;349(17):1614–27.

10.   Plant TM, Ramaswamy S, DiPietro MJ. Repetitive activation of hypothalamic G

protein-coupled receptor 54 with intravenous pulses of kisspeptin in the juvenile

monkey (Macaca mulatta) elicits a sustained train of gonadotropin-releasing hormone

discharges. Endocrinology. 2006;147(2):1007–13.

11.   Navarro VM, Fernandez-Fernandez R, Castellano JM, Roa J, Mayen A, Barreiro ML,

Gaytan F, Aguilar E, Pinilla L, Dieguez C, Tena-Sempere M . Advanced vaginal

opening and precocious activation of the reproductive axis by KiSS-1 peptide, the

endogenous ligand of GPR54. J Physiol. 2004;561(2):379–86.

12.   Clarkson J, Han SY, Piet R, McLennan T, Kane GM, Ng J, Porteous RW, Kim JS,

Colledge WH, Iremonger KJ, Herbison AE. Definition of the hypothalamic GnRH

pulse generator in mice. Proc Natl Acad Sci U S A. 2017;114(47):e10216–23.

13.   Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, Rønnekleiv OK .

High-frequency stimulation-induced peptide release synchronizes arcuate kisspeptin

neurons and excites GnRH neurons. Elife. 2016;5:e16246.

14.   Voliotis M, Li XF, De Burgh R, Lass G, Lightman SL, O'Byrne KT, Tsaneva-

Atanasova K . The Origin of GnRH pulse generation: An integrative mathematical-

experimental approach. J Neurosci. 2019;39(49):9738–47.

15.   Liu X, Yeo SH, McQuillan HJ, Herde MK, Hessler S, Cheong I, Porteous R, Herbison

AE . Highly redundant neuropeptide volume co-transmission underlying episodic

activation of the gnrh neuron dendron. Elife. 2021;10:e62455.

16.   Wang L, Moenter SM. Differential Roles of Hypothalamic AVPV and Arcuate

Kisspeptin Neurons in Estradiol Feedback Regulation of Female Reproduction.

Neuroendocrinology. 2020;110(3–4):172–84.

17.   Stephens SBZ, Raper J, Bachevalier J, Wallen K. Neonatal amygdala lesions advance

pubertal timing in female rhesus macaques. Psychoneuroendocrinology. 2015;51:307–

17.

18.   Bar-Sela M, Critchlow V. Delayed puberty following electrical stimulation of

amygdala in female rats. Am J Physiol. 1966;211(5):1103–7.

19.   Li XF, Hu MH, Hanley BP, Lin YS, Poston L, Lightman SL, O'Byrne KT . The

posterodorsal medial amygdala regulates the timing of puberty onset in female rats.

Endocrinology. 2015;156(10):3725–36.

20.   Choi GB, Dong HW, Murphy AJ, Valenzuela DM, Yancopoulos GD, Swanson LW,

Anderson DJ . Lhx6 delineates a pathway mediating innate reproductive behaviors

from the amygdala to the hypothalamus. Neuron. 2005;46(4):647–60.

21.   Keshavarzi S, Sullivan RKP, Ianno DJ, Sah P. Functional properties and projections of

neurons in the medial amygdala. J Neurosci. 2014;34(26):8699–715.

22.   Moore AM, Coolen LM, Lehman MN. Kisspeptin/Neurokinin B/Dynorphin (KNDy)

cells as integrators of diverse internal and external cues: evidence from viral-based

monosynaptic tract-tracing in mice. Sci Rep. 2019;9(1):1–5.

23.   Yeo SH, Kyle V, Blouet C, Jones S, Colledge WH. Mapping neuronal inputs to Kiss1

neurons in the arcuate nucleus of the mouse. PLoS One. 2019;14(3):e0213927.

24.   Kim J, Semaan SJ, Clifton DK, Steiner RA, Dhamija S, Kauffman AS. Regulation of

Kiss1 expression by sex steroids in the amygdala of the rat and mouse. Endocrinology.

2011;152(5):2020–30.

25.   Adekunbi DA, Li XF, Li S, Adegoke OA, Iranloye BO, Morakinyo AO, Lightman SL,

Taylor PD, Poston L, O'Byrne KT . Role of amygdala kisspeptin in pubertal timing in

female rats. PLoS One. 2017;12(8):e0183596.

26.   Lass G, Li XF, de Burgh RA, He W, Kang Y, Hwa-Yeo S, Sinnett-Smith LC,

Manchishi SM, Colledge WH, Lightman SL, O'Byrne KT . Optogenetic stimulation of

kisspeptin neurones within the posterodorsal medial amygdala increases luteinising

hormone pulse frequency in female mice. J Neuroendocrinol. 2020;32(2):e12823.

27.   Govic A, Paolini AG. In vivo electrophysiological recordings in amygdala subnuclei

reveal selective and distinct responses to a behaviorally identified predator odor. J

Neurophysiol. 2015;113(5):1423–36.

28.   Kinsey-Jones JS, Li XF, Knox AM, Lin YS, Milligan SR, Lightman SL, O'Byrne KT .

Corticotrophin-releasing factor alters the timing of puberty in the female rat. J

Neuroendocrinol. 2010;22(2):102–9.

29.   Shemesh Y, Forkosh O, Mahn M, Anpilov S, Sztainberg Y, Manashirov S,

Shlapobersky T, Elliott E, Tabouy L, Ezra G, Adler ES . Ucn3 and CRF-R2 in the

medial amygdala regulate complex social dynamics. Nat Neurosci. 2016;19(11):1489–

96.

30.   Jamieson PM, Li C, Kukura C, Vaughan J, Vale W. Urocortin 3 modulates the

neuroendocrine stress response and is regulated in rat amygdala and hypothalamus by

stress and glucocorticoids. Endocrinology. 2006;147(10):4578–88.

31.   Paxinos G, Franklin KB . Mouse Brain in Stereotaxic Coordinates. Acad Press.

2019;5:246.

32.   Zhan J, Komal R, Keenan WT, Hattar S, Fernandez DC . Non-invasive strategies for

chronic manipulation of dreadd-controlled neuronal activity. J Vis Exp.

2019;2019(150):e59439.

33.   Steyn FJ, Wan Y, Clarkson J, Veldhuis JD, Herbison AE, Chen C. Development of a

methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile

and adult male mice. Endocrinology. 2013;154(12):4939–45.

34.   Vidal A, Zhang Q, Médigue C, Fabre S, Clément F. Dynpeak: An algorithm for pulse

detection and frequency analysis in hormonal time series. PLoS One.

2012;7(7):e39001.

35.   Fernandes C, File SE. The influence of open arm ledges and maze experience in the

elevated plus-maze. Pharmacol Biochem Behav. 1996;54(1):31–40.

36.   Gill JC, Navarro VM, Kwong C, Noel SD, Martin C, Xu S, Clifton DK, Carroll RS,

Steiner RA, Kaiser UB . Increased Neurokinin B (Tac2) Expression in the Mouse

Arcuate Nucleus Is an Early Marker of Pubertal Onset with Differential Sensitivity to

Sex Steroid-Negative Feedback than Kiss1. Endocrinology. 2012;153(10):4883.

37.   Simavli S, Thompson IR, Maguire CA, Gill JC, Carroll RS, Wolfe A, Kaiser UB,

Navarro VM . Substance P Regulates Puberty Onset and Fertility in the Female

Mouse. Endocrinology. 2015;156(6):2313–22.

38.   Wright H, Aylwin CF, Toro CA, Ojeda SR, Lomniczi A . Polycomb represses a gene

network controlling puberty via modulation of histone demethylase Kdm6b

expression. Sci Rep. 2021;11(1):1–7.

39.   Adekunbi DA, Li XF, Lass G, Shetty K, Adegoke OA, Yeo SH, Colledge WH,

Lightman SL, O'Byrne KT. Kisspeptin neurones in the posterodorsal medial amygdala

modulate sexual partner preference and anxiety in male mice. J Neuroendocrinol.

2018;30(3):e12572.

40.   Ivanova D, Li XF, McIntyre C, O'Byrne KT . Chronic Exposure to Predator Odour

Stress Disrupts LH Pulsatility and Delays Puberty While Activation of Amygdala

Kisspeptin Advances Puberty. Endocr Soc. 2020;Supplement:SUN-LB49.

41.   Pincus M, Godfrey JR, Feczko E, Earl E, Miranda-Dominguez O, Fair D, Wilson ME,

Sanchez MM, Kelly C . Chronic psychosocial stress and experimental pubertal delay

affect socioemotional behavior and amygdala functional connectivity in adolescent

female rhesus macaques. Psychoneuroendocrinology. 2021;127:105154.

42.   Yang JA, Song CI, Hughes JK, Kreisman MJ, Parra RA, Haisenleder DJ, Kauffman

AS, Breen KM . Acute psychosocial stress inhibits LH pulsatility and kiss1 neuronal

activation in female mice. Endocrinology. 2017;158(11):3716–23.

43.   Gaytan F, Morales C, Leon S, Heras V, Barroso A, Avendaño MS, Vazquez MJ,

Castellano JM, Roa J, Tena-Sempere M . Development and validation of a method for

precise dating of female puberty in laboratory rodents: The puberty ovarian maturation

score (Pub-Score). Sci Rep. 2017;7(1):1–11.

44.   Marshall AD . Developmental timing of trauma exposure relative to puberty and the

nature of psychopathology among adolescent girls. J Am Acad Child Adolesc

Psychiatry. 2016;55(1):25–32.

45. Johnson FK, Delpech JC, Thompson GJ, Wei L, Hao J, Herman P, Hyder F, Kaffman

A . Amygdala hyper-connectivity in a mouse model of unpredictable early life stress.

Transl Psychiatry. 2018;8(1):1–14.

46. Lin Y, Li XF, Lupi M, Kinsey-Jones JS, Shao B, Lightman SL, O'Byrne KT . The role

of the medial and central amygdala in stress-induced suppression of pulsatile LH

secretion in female rats. Endocrinology. 2011;152(2):545–55.

47. Dopfel D, Perez PD, Verbitsky A, Bravo-Rivera H, Ma Y, Quirk GJ, Zhang N .

Individual variability in behavior and functional networks predicts vulnerability using

an animal model of PTSD. Nat Commun. 2019;10(1):1–2.

48. Chester JA, Barrenha GD, Hughes ML, Keuneke KJ . Age- and sex-dependent effects

of footshock stress on subsequent alcohol drinking and acoustic startle behavior in

mice selectively bred for high-alcohol preference. Alcohol Clin Exp Res.

2008;32(10):1782–94.

49. Kinsey-Jones JS, Li XF, Knox AM, Wilkinson ES, Zhu XL, Chaudhary AA, Milligan

SR, Lightman SL, O'byrne KT . Down-regulation of hypothalamic kisspeptin and its

receptor, Kiss1r, mRNA expression is associated with stress-induced suppression of

luteinising hormone secretion in the female rat. J Neuroendocrinol. 2009;21(1):20–9.

50. Agnieszka Podfigurna, Anna Szeliga BM. Serum kisspeptin and corticotropin-

releasing hormone levels in patients with functional hypothalamic amenorrhea.

Gynecol Reprod Endocrinol Metab. 2020;1(1):37–42.

51. Li XF, Bowe JE, Lightman SL, O'Byrne KT . Role of corticotropin-releasing factor

receptor-2 in stress-induced suppression of pulsatile luteinizing hormone secretion in

the rat. Endocrinology. 2005;146(1):318–22.

52.   Fekete ÉM, Zhao Y, Li C, Sabino V, Vale WW, Zorrilla EP. Social defeat stress

564   activates medial amygdala cells that express type 2 corticotropin-releasing factor

565   receptor mRNA. Neuroscience. 2009;162(1):5–13.

566   53.   Pardo-Bellver C, Cádiz-Moretti B, Novejarque A, Martínez-García F, Lanuza E.

567   Differential efferent projections of the anterior, posteroventral, and posterodorsal

568   subdivisions of the medial amygdala in mice. Front Neuroanat. 2012;6(33):1–26.

569   54.   Deussing JM, Breu J, Kühne C, Kallnik M, Bunck M, Glasl L, Yen YC, Schmidt MV,

570   Zurmühlen R, Vogl AM, Gailus-Durner V . Urocortin 3 modulates social

571   discrimination abilities via corticotropin-releasing hormone receptor type 2. J

572   Neurosci. 2010;30(27):9103–16.

573   55.   Cavalcante JC, Sita LV, Mascaro MB, Bittencourt JC, Elias CF. Distribution of

574   urocortin 3 neurons innervating the ventral premammillary nucleus in the rat brain.

575   Brain Res. 2006;1089(1):116–25.

576   56.   Li C, Vaughan J, Sawchenko PE, Vale WW. Urocortin III-immunoreactive projections

577   in rat brain: Partial overlap with sites of type 2 corticotrophin-releasing factor receptor

578   expression. J Neurosci. 2002;22(3):991–1001.

579   57.   Day FR, Elks CE, Murray A, Ong KK, Perry JRB. Puberty timing associated with

580   diabetes, cardiovascular disease and also diverse health outcomes in men and women:

581   the UK Biobank study. Sci Rep. 2015;5(1):1–12.

583 **Figure 1.** Female C57Bl6/J mice chronically exposed to 2,4,5-Trimethylthiazole (TMT) from

584 post-natal day (pnd) 21 for 14 days showed delayed first estrous (FE) without affecting vaginal

585 opening (VO) and body weight (BW). Effect of chronic TMT exposure on day of VO (A), FE

586 (B) and (C) BW weight gain between pnd 21 and 45. +++$p < 0.0001$ Control group (combined

587 water control: n=6; ethyl vanillin control: n=4) vs TMT group (n=14).

Figure 2. Female C57Bl6/J mice chronically exposed to 2,4,5-Trimethylthiazole (TMT) from post-natal day (pnd) 21 for 14 days showed suppressed pre-pubertal pulsatile luteinising hormone (LH) secretion on pnd 26 and 29. Representative LH pulse profile with (A) control mouse on pnd 26, (B) TMT-exposed mouse on pnd 26, (C) control mouse on pnd 29 and (D) TMT-exposed mouse on pnd 29. (E), Summary of LH pulse frequency on pnd 26 and 29 of control and TMT-exposed group during the 80 min blood sampling period. LH pulses detected by the DynePeak algorithm are indicated with an asterisk located above each pulse on the representative LH pulse profiles. ++$p<0.001$ control group (combined water control: n=6; ethyl vanillin control: n=3) vs. TMT-exposed group (n=8) on pnd 26 and 29.

Figure 3. Female C57Bl6/J mice chronically exposed to 2,4,5-Trimethylthiazole (TMT) showed increased anxiety-like behavior on the light-dark box (LDB) and elevated plus maze (EPM) and decreased social interaction with familiar conspecifics on the social interaction (SI) test. (A) Summary of time spent in the light compartment of the LDB on post-natal day (pnd) 19 (before beginning of TMT-exposure), 27 (during TMT exposure) and 40 (after termination of TMT-exposure), (B) time spent socially interacting with familiar conspecifics on pnd 19, 27 and 40, and (C) time spent in the open arm of the EPM on pnd 20, 28 and 41. +$p<0.05$ time spent on LDB, SI and EPM of control group (combined water control: n=6; ethyl vanillin control: n=4) vs TMT-exposed group (n=14) on pnd 27 or 28 (during TMT-exposure); ###$p<0.0001$ time spent in the open arm of the EPM in the TMT-exposed group on pnd 28 vs pnd 41.

Figure 4. Expression of AAV5-hSyn-DIO-HA-hM3D(Gq)-IRES-mCitrine in posterodorsal medial amygdala (MePD) Urocortin 3 (Ucn3) neurons. (A-F) Representative dual fluorescence photomicrographs of the MePD from a Ucn3-cre-tdTomato female mouse injected with AAV5-hSyn-DIO-HA-hM3D(Gq)-IRES-mCitrine. Ucn3 neurons labelled with m-Citrine (A)

and tdTomato (C) appear yellow/orange (E). (B), (D) and (F) are a higher power view of (A),

(C) and (E) respectively. Scale bars represent A, C, E 100 μm and B, D, F 25 μm; OT, Optic

tract (blue line).

Figure 5. DREADD (hM3D) activation of urocortin 3 (Ucn3) neurons in the posterodorsal

sub-nucleus of the medial amygdala (MePD) of Ucn3-cre-tdTomato female mice from post-

natal day (pnd) 21 for 14 days delays first estrous (FE) without affecting vaginal opening (VO)

and body weight (BW) in urocortin3-cre-tdTomato female mice. Effect of MePD Ucn3

activation on day of VO (A), FE (B) and (C) BW weight gain between pnd 21 and 45. $+p<0.05$

Control group (combined control AAV: n=6; cre-negative control: n=4; no CNO n=3) vs

DREADD group (n=8).

Figure 6. DREADD (hM3D) activation of urocortin 3 (Ucn3) neurons in the posterodorsal

sub-nucleus of the medial amygdala (MePD) in Ucn3-cre-tdTomato female mice from post-

natal day (pnd) 21 for 14 days suppresses pre-pubertal LH pulse frequency sampled on pnd 29.

Representative LH pulse profile from Ucn3-cre-tdTomato female mice on pnd 29 with (A)

control mouse, (B) DREADD (hM3D) activated mouse, (C) Summary of LH pulse frequency

on pnd 29 of control and DREADD group during the 80 min blood sampling period. LH pulses

detected by the DynePeak algorithm are indicated with an asterisk located above each pulse on

the representative LH pulse profiles. $+p<0.05$ Control group (combined control AAV: n=3;

cre-negative control: n=3) vs DREADD group (n=5).

Figure 1

A Vaginal Opening

B First Estrous

C

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6