Title: Sporadic Diurnal Fluctuations of Cyanobacterial Populations in

Oligotrophic Temperate Systems Can Prevent Accurate Characterization of Change

and Risk in Aquatic Systems

Authors: Ellen S. Cameron[a], Anjali Krishna[a], Monica B. Emelko[b], Kirsten M. Müller[a,*]

[a] Department of Biology, University of Waterloo, 200 University Ave. W, Waterloo,

Ontario, Canada, N2L 3G1

[b] Department of Civil & Environmental Engineering, University of Waterloo, 200

University Ave. W, Waterloo, Ontario, Canada, N2L 3G1

* Corresponding author. Tel.: +1 519 888 4567x32224

E-mail address: kirsten.muller@uwaterloo.ca (K.M. Müller).

Abstract

Continental-scale increases in aquatic system eutrophication are linked with increased

cyanobacteria threats to recreational water use and drinking water resources

globally. Increasing evidence suggests that diurnal vertical migration of cyanobacteria are

key factors that must be considered in cyanobacterial bloom risk management. While this

has been discussed in marine and eutrophic freshwater contexts, reports of diurnal

vertical migration of cyanobacteria in oligotrophic freshwater lakes are scant. Typical

monitoring protocols do not reflect these dynamics and frequently focus only on surface

water sampling approaches, and either ignore sampling time or recommend large midday

timeframes (e.g., 10AM-3PM), thereby preventing accurate characterization of

cyanobacterial community dynamics. To evaluate the impact of diurnal migrations and

water column stratification on cyanobacterial abundance and composition, communities

were characterized in a shallow well-mixed lake interconnected to a thermally stratified

lake in the Turkey Lakes Watershed (Ontario, Canada) using amplicon sequencing of the

16S rRNA gene across a multi-time point sampling series in 2018 and 2022. This work

showed that cyanobacteria are present in oligotrophic lakes and their community structure

varies (i) diurnally, (ii) across the depth of the water column, (iii) interannually within the

same lake and (iv) between different lakes that are closely interconnected within the same

watershed. It underscored the need for integrating multi-timepoint, multi-depth discrete

sampling guidance into lake and reservoir monitoring programs to describe cyanobacteria

community dynamics and signal change to inform risk management associated with the

potential for cyanotoxin production. Ignoring variability in cyanobacterial community

dynamics (such as that reported herein) and reducing sample numbers can lead to a false

sense of security and missed opportunities to identify and mitigate changes in trophic

status and associated risks such as toxin or taste and odor production, especially in

sensitive, oligotrophic systems.

Graphical Abstract

Keywords

Amplicon sequencing, diel cycles, vertical migration, buoyancy, algae, eutrophication

Highlights

- Cyanobacterial populations fluctuate sporadically across diurnal cycles

- Cyanobacterial communities can vary significantly between interconnected lakes

- Significant annual shifts in communities signal higher risk and need for monitoring

- Cyanobacteria monitoring for risk management should incorporate time and depth

1. Introduction

Cyanobacteria are recognized as a threat to surface water quality as they can form dense blooms and produce secondary metabolites including potent cyanotoxins (Harke et al., 2016; Huisman et al., 2018; Paerl, 2014; Vu et al., 2020) and, compounds associated with taste and odor problems (e.g., geosmin, 2-methyl isoborneol). As a result of anthropogenic activities, cyanobacterial bloom frequency and intensity have been increasing globally (Huisman et al., 2018) and climate change-exacerbated landscape disturbances (e.g., wildfires) further promote their proliferation (Emelko et al., 2016; Silins et al., 2014). These organisms can challenge water treatment processes beyond their response capacities (Emelko et al., 2011) and exacerbate disinfection by-product formation potential after landscape disturbances such as wildfires (Tsai et al., 2017). Accordingly, continued increases in toxic bloom occurrence globally (Huisman et al., 2018) underscore the need to advance source water protection and drinking water safety plan development guidance. Improved accuracy in characterization of cyanobacterial communities and their distributions which are underpinned by diurnal migrations will accelerate that advancement and associated risk management.

The diurnal migration of cyanobacterial populations is driven by cellular characteristics (Naselli-Flores et al., 2021) and water column stability (Walsby et al., 1997), as summarized in Table S1. Diurnal migration rates (e.g., flotation and/or sinking rates) and vertical distribution of populations may be distinctive to individual taxa arising from characteristic differences in cell sizes or the presence of specialized cellular structures impacting cellular density (Naselli-Flores et al., 2021; Reynolds et al., 1987). However, these characteristic distributions are further impacted by water column stability

(Walsby et al., 1997) with the potential for oscillatory diurnal variation in the distribution

of cyanobacterial populations within systems of differing water column stability (Hunter

et al., 2008). Rapid changes caused by oscillatory diurnal variation in the distribution of

cyanobacterial populations can challenge detection if community structure and system

dynamics are not well characterized.

Climate variability and landscape change-associated impacts on freshwater systems

are key drivers of the need to better describe cyanobacterial growth and behavior in

freshwater systems. Warming temperatures alter thermal stratification and mixing

regimes in lakes (Woolway and Merchant, 2019) and increases in extreme precipitation

and associated runoff increase nutrient delivery and eutrophication in receiving waters

(Scholz and Brahney, 2022; Stoddard et al., 2016) and contribute to increases in

cyanobacterial blooms, even in oligotrophic systems (Reinl et al., 2021; Sterner et al.,

2020). Improved community characterization that reflects contemporary understanding of

the diversity of cyanobacterial populations and their adaptations is therefore essential to

managing risks attributable to the presence of these potentially toxic organisms in water

supplies and better informing pre-emptive mitigation of potential health impacts and

drinking water treatment challenges.

Key factors that must be considered in cyanobacterial bloom risk management

include diurnal vertical migration of cyanobacteria, which has been reported in marine

(Olli, 1999) and eutrophic freshwater (Hunter et al., 2008; von Orgies-Rutenberg et al.,

2018) systems. More recently, advanced technologies including remote sensing

techniques have been used to monitor cyanobacterial blooms evolution in eutrophic

systems and has enabled high-resolution exploration of diurnal variability of visible

bloom biomass (Li et al., 2023; Wang et al., 2023; Xue et al., 2023). In contrast, reports

of diurnal vertical migration of cyanobacteria in oligotrophic freshwater lakes are scant

and have been limited to spectrophotometric analysis of chlorophyll-*a* and microscopic

cell enumeration over periods of 24 hours or less (Frempong, 1981; Ganf, 1974; Gilbert

et al., 2010; Ibelings et al., 1991; Olli, 1999; Shahraki et al., 2020). Despite next-

generation sequencing and metagenomic approaches being used to characterize

cyanobacterial communities in freshwater lakes (Valadez-Cano et al., 2022) and in the

context of water treatment (Jalili et al., 2021; Le et al., 2022). Studies using next-

generation sequencing technology to characterize diurnal variation of cyanobacteria

remain limited in number and scope, however (Gilbert et al., 2010; Shahraki et al., 2020).

While the use of multi-time point sampling series provides insights into the variability

observed in community composition over periods of hours, investigations that use them

do not typically encompass the natural dynamic variability that aquatic systems may

experience day-to-day.

Here, we provide guidance for the development of cyanobacterial monitoring

programs for risk management in oligotrophic freshwater systems, the impact of

sampling time and depth on cyanobacterial community composition were evaluated using

multi-time point sampling over consecutive days and amplicon sequencing of the V4

region in the 16S rRNA gene. Taxonomic composition and community diversity analyses

were used to examine diurnal trends in cyanobacterial distribution and the associated

impact of sampling time. Fluctuations in cyanobacterial community composition were

evaluated (i) temporally over a 48-hour multi-time point sampling period and (ii)

spatially within the water column of a shallow well-mixed lake and an interconnected,

thermally stratified lake. Temporal differences were further characterized in the thermally

stratified lake during a second aquatic growing season.

2. Methods

2.1 Study Site: Turkey Lakes Watershed

The Turkey Lakes Watershed (TLW) Study was established in 1980 to investigate

ecosystem effects of acidic atmospheric deposition; Jeffries et al. (1988) provide a

comprehensive description of the physical characteristics of the watershed. In brief, it is

approximately 50 km north of Sault Ste Marie, Ontario on the Canadian Shield in an

uneven-aged tolerant hardwood and mixed conifer forest (Jeffries et al., 1988). This

watershed consists of four interconnected lakes fed by both first order streams and

groundwater: Batchawana Lake, Wishart Lake, Little Turkey Lake, and Big Turkey Lake

(Jeffries et al., 1988; Figure S1). The shallow depth of Wishart Lake often results in

complete wind-induced mixing that prevents water column stratification whereas Little

Turkey Lake, Big Turkey Lake and Batchawana Lake are deeper and regularly undergo

thermal stratification (Figure S2). These lakes are classified as oligotrophic to

mesotrophic, and cyanobacteria have been reported as the dominant members of

phytoplankton communities and previously reported as major contributors to peaks in

primary production during summer periods (Jeffries et al., 1988).

2.2 Sample Collection

Water samples were collected from the deepest point in Little Turkey

(47°02'37.2"N 84°24'24.4"W) and Wishart (47°03'00.0"N 84°23'58.3"W) Lakes during

a two-day period across three time points on both days (2018a/b, 2022a/b): morning (8-9

a.m.), midday (12-1 p.m.) and afternoon (4-5 p.m.) during two sampling events (August

2018 and August 2022). Samples were collected from Secchi depth, the water depth at

which light penetration is approximately 1% of surface illumination and is considered the

maximum depth at which there is sufficient light for photosynthesis (Bukata et al., 1988).

In Wishart Lake, these samples were collected from near the bottom (4 m) due to high

water clarity. Secchi depths ranged from 4 to 5.25 m in Little Turkey Lake. Water

samples were also collected at the surface (0 m) in Little Turkey Lake on the first

sampling day in 2018 (2018a) and both sampling days in 2022 (2022a, 2022b) to evaluate

potential temporal correlations in cyanobacterial communities between depths.

Water samples collected using a Masterflex E/S portable sampler peristaltic pump

were vacuum filtered through a 47 mm GF/C filter (Whatman plc, Buckinghamshire,

United Kingdom). For samples collected in August 2018, 250 mL of vacuum filtered

water were filtered a second time through a 0.22 µm Sterivex™ filter to collect additional

microbes. All filters were stored at -20°C prior to DNA extraction. Sampling details are

provided in Table S2.

2.3 DNA Extraction, V4 16S rRNA Gene Amplicon Sequencing

DNA was extracted using the DNeasy PowerSoil Kit (QIAGEN Inc., Venlo,

Netherlands) following the manufacturer's protocol. Following this, elution buffer was

added to spin columns for 15 minutes prior to elution. DNA was quantified using a

NanoDrop spectrophotometer (Table S2); absolute values were only accurate at DNA

concentrations of more than 10ng/µl. The DNA extracts were submitted to a commercial

laboratory (Metagenom Bio Inc.,Waterloo, ON) for amplicon sequencing. Primers

designed to target the 16S rRNA gene V4 region [515FB

(GTGYCAGCMGCCGCGGTAA) and 806RB (GGACTACNVGGGTWTCTAAT)]

(Walters et al., 2015) were used for PCR amplification. Polymerase chain reactions

(PCR) were conducted in triplicate for each sample (25 µL). Reaction mixtures were

composed of 2.5 µL of 10X *Taq* buffer, 0.5µL of 10mM dNTP, 0.25 µL of BSA

(20mg/mL), 5.0 µL of 1 µM forward primer, 5.0 µL of 1 µM of reverse primer, 5.0 µL of

sample DNA, 0.2 µL *Taq* DNA polymerase (5u/µL) and 6.55 µL of PCR water. DNA

was initially denatured at 95°C for 5 minutes, followed by 35 cycles of: 1) 95°C for 30

seconds, 2) 30°C for 30 seconds, 3) 72°C for 50 seconds. A final extension was

performed at 72°C for 10 minutes. Triplicate PCR products were pooled, gel purified and

quantified using Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific Inc.). Library

DNA was prepared and sequenced with MiSeq Reagent Kit v2 (2x 250 cycles) and

sequenced with the Illumina MiSeq platform (Illumina Inc., San Diego, United States).

2.4 Sequence Processing & Library Size Normalization

The program QIIME2 (v. 2019.10; Bolyen et al., 2019) was used for

bioinformatic processing. Demultiplexed paired-end sequences were trimmed and

denoised, including the removal of chimeric sequences and singleton sequence variants,

using *DADA2* (Callahan et al., 2016). Amplicon sequence variants across the sample sets

were clustered at 99% identity to account for single nucleotide discrepancies in

sequenced amplicons using *vsearch* (Rognes et al., 2016) to generate a representative set

of operational taxonomic units (OTUs). Taxonomic classification was performed using a

Naïve-Bayes taxonomic classifier trained using the SILVA138 database (Quast et al.,

2013; Yilmaz et al., 2014). Sequences classified as cyanobacteria at the phylum level

were aligned to reference 16S rRNA cyanobacterial sequences in the Cyanoseq database

(Lefler et al., 2023) using *MAFFT* (v7.490; Katoh and Standley, 2013). A phylogenetic

tree was generated using *iqtree* (v. 2.2.0; Nguyen et al., 2015) using an unrestricted

model (UNREST+FO+R10) selected by the ModelFinder Bayesian Information Criterion

and visualized with Interactive Tree of Life (iTOL; Letunic and Bork, 2021; Figure S3).

Confirmed taxonomic classifications for OTUs classified as cyanobacteria at the phylum

level were manually curated to reflect taxonomic levels above the genus level according

to taxonomic labels in the Cyanoseq database based on placements in the phylogenetic

tree.

Files from QIIME2 were imported into R (v. 4.0.1; R Core Team, 2020) for

downstream analyses using *qiime2R* (v. 0.99.23; Bisanz, 2018). Initial sequence libraries

were filtered to exclude OTUs that were taxonomically classified as mitochondria or

chloroplast sequences using *phyloseq* (v. 1.32.0; McMurdie and Holmes, 2012). For

cyanobacterial community analysis, OTUs classified as cyanobacteria at the phylum level

were selected to create libraries consisting of only cyanobacterial sequences.

2.5 Cyanobacterial Communities - Taxonomic Composition & Diversity Analyses

Bacterial community composition was assessed at the phylum level and relative

abundances were visualized using a stacked bar plot produced with *mirlyn* (Cameron and

Tremblay, 2020). Relative abundances were randomized across bacterial phyla within

samples to identify significantly dominant groups within the bacterial community in

relation to sampling time and adjusted with a Bonferroni correction using the

randomseqsig function in *mirlyn*. Cyanobacterial community composition was assessed

through the selection of a subset of sequences classified as "cyanobacteria" at the phylum

level and relative abundances of cyanobacterial families were also visualized using a

stacked bar plot. For bacterial and cyanobacterial community composition bar plots,

groups with an average relative abundance lower than 1% across samples were collapsed

into a low abundance category for visualization purposes.

As amplicon sequencing only indirectly and partially represents source diversity

and clustering, amplification, and the lack of certainty about how many zeros should be

included in the data compromise statistical inference about source diversity (Schmidt et

al., 2021), community diversity analyses were performed using relative abundances

instead of rarefying. While rarefying can be utilized to compare libraries in a statistically

meaningful manner (Cameron et al., 2021), library sizes between samples collected in

2018 and 2022 differed by orders of magnitude creating extreme data loss of sequences

or entire samples. OTUs present at >1% in the bacterial community were utilized for

diversity analyses to reduce the risk of accidental non-detection. Alpha diversity was

assessed using direct counts of unique OTUs and the Shannon Index (Shannon, 1948) to

identify trends in sample diversity as a function of time. Bray-Curtis dissimilarity metric

was calculated on compositional community data using *vegdist* in *vegan* (Oksanen et al.,

2022) and used for permutational analysis of variance with *adonis2*. Bray-Curtis

distances were visualized using ComplexHeatMap (Gu, 2022). To identify individual

OTUs driving compositional changes, highly abundant (>10% of cyanobacterial

community) were selected and normalized using Z-scores to identify enrichment or

under-enrichment of individual abundant OTUs as a function of time.

2.6 Data Availability

Sequence data analyzed in this study is available in the European Nucleotide

Archive (ENA) under study accession ERP134980.

3. Results

3.1 Diurnal Conservation of Bacterial & Cyanobacterial Communities

Cyanobacteria classified sequences were found to comprise a variable proportion of the bacterial community composition across sampling events in 2018 and 2022 (Figure 1A). Relative abundances of cyanobacteria were more similar during sampling events in Little Turkey Lake than Wishart Lake in 2018 (Figure 1A; Figure 2A). Cyanobacteria did not contribute to a significant portion of the bacterial community in all sampling events of 2018 ($p = 1$; Table S4). In contrast, cyanobacterial sequences comprised a significant portion of the bacterial community in Little Turkey Lake in 2022. Specifically, cyanobacteria comprised a significant portion of the bacterial community (i) at all sampling times at Secchi depth ($p < 0.005$) on both sampling days (2022a; 2022b) and (ii) in morning and midday surface samples on the second sampling day (2022b; $p < 0.003$).

Cyanobacterial communities were frequently dominated by sequences classified to the family Prochlorococcaceae with minimal diurnal fluctuations (Figure 1B; Figure 2B). Cyanobacterial communities observed in Little Turkey Lake in 2018 included large proportions of taxonomic families not observed in Wishart Lake or during the 2022 sampling events. A higher abundance of Cyanothrichaceae, Microcystaceae, and Trichocoleusaceae classified sequences were observed in the 2018 samples from Little Turkey Lake. Relative abundances of Cyanothrichaceae decreased from morning (11.1%) to afternoon (1.4%) at surface depths, possibly due to downward vertical migration. At Secchi depth on the first sampling day (2018a), Trichocoleusaceae were found in similar relative abundances (48.6%) in the morning sampling event but experienced a drastic

decrease by midday (10.1%) before a second increase at the final sampling time (19.8%).

On the second sampling day (2018b) in Little Turkey Lake, Trichocoleusaceae relative

abundances were relatively consistent from morning (19.0%) to midday (19.1%) and then

increased in the afternoon (25%). Notably, the relative abundances observed in the

morning sampling point at Secchi depth of the 2018a sampling event had higher

similarity to those at the water surface than those at Secchi depth during subsequent

sampling events. Fluctuations in Prochlorococcaceae relative abundances were negatively

correlated with Trichocoleusaceae and Microcystaceae abundances due to the

compositional constraint of amplicon sequencing data.

Similarity between cyanobacterial communities across sampling events was

analyzed using the Bray-Curtis dissimilarity metric (Figure 3). Sampling time did not

significantly impact cyanobacterial community composition in either of the study lakes (p

= 0.97; Table S5) suggesting diurnal conservation of cyanobacterial community

composition. However, significant compositional differences were observed between the

two lakes sites (p = 0.002). The insignificance of sampling time and significance of

sampling site were also observed for bacterial community composition (p = 0.99; p =

0.001; Table S6).

Within Little Turkey Lake, interannual variability between sampling events had a

significant impact on cyanobacterial communities (p = 0.001). Sampling depth of the

stratified water column in Little Turkey did not significantly impact the community

structure (p = 0.154) and high similarity was still observed between samples collected at

the same sampling depth. Additionally, similarity was observed within lakes across the

sampling days of each year, except in the unique case of the morning of the first sampling

day of 2018 (2018a). Cyanobacterial communities at Secchi depth in the morning of the

2018a sampling event were highly similar to surface communities observed in Little

Turkey Lake, a phenomenon that was not observed during other sampling events (Figure

3).

Figure 1: Stacked bar plots depicting the taxonomic composition of bacterial

communities in terms of the relative abundance of major bacterial phyla (A) and

cyanobacterial communities in terms of relative abundance of major cyanobacterial

families (B) measured across a multi-time point sampling series over two consecutive

sampling days (2018a/2018b, 2022a/2022b) in a thermally stratified lake (Little Turkey

Lake).

Figure 2: Stacked bar plots depicting the taxonomic composition of bacterial

communities in terms of the relative abundance of major bacterial phyla (A) and

cyanobacterial communities in terms of relative abundance of major cyanobacterial

families (B) measured across a multi-time point sampling series over two consecutive

sampling days (2018a/2018b) in a shallow, non-stratified lake (Wishart Lake).

Figure 3: Compositional similarity of cyanobacterial communities across a multi-time

point sampling series conducted over two consecutive sampling days (2018a/2018b,

2022a/2022b). Bray-Curtis dissimilarity metric (0 = completely identical, 1 = completely

dissimilar) characterizes the differences present in the OTU composition of

cyanobacterial communities. Overall proportion of cyanobacteria in bacterial

communities displayed on the right-hand side highlights and contextualizes the

interannual variability observed within Little Turkey Lake bacterial communities.

3.2 Sporadic Diurnal Variability in Relative Abundances of OTUs

Cyanobacterial community diversity across sampling times was assessed using

(i) counts of unique OTUs (>1% abundance) and (ii) the Shannon Index (Figure 3A).

Little Turkey Lake exhibited different trends in sample diversity dependent on the

sampling event. Within samples collected during the 2018 sampling events, a higher

number of cyanobacterial OTUs were detected in comparison to the 2022 sampling

series. However, there was no consistent trend of diurnal fluctuations, revealing the

sporadic fluctuations which may occur throughout a diurnal period. For example, 2018

sampling events at Secchi depth revealed opposing trends at midday where diversity

scores decrease during the 2018a sampling event but increase during the 2018b sampling

event. Similar results are also observed during the 2022 sampling events. The 2022b

sampling event also reveals a consistent OTU count across sampling times, but the

Shannon Index suggests changes in proportionality of these detected OTUs across time.

Although results from Wishart Lake are limited to the 2018 sampling events, contrary to

the sporadic diurnal fluctuations of diversity observed in Little Turkey Lake, Wishart

Lake was shown to have corresponding spikes in diversity at the midday sampling point

on both sampling days.

Diurnal fluctuations of highly abundant cyanobacterial OTUs (>10%) were

identified using Z-scores; time points associated with a higher Z-score reflect greater

deviation from the mean (Figure 3B). Additionally, temporal distribution of

cyanobacterial OTUs was visualized to identify ubiquitous OTUs within cyanobacterial

communities (Figure 4). Specifically, OTU5755 (Prochlorococcaceae) was found in all

samples independent of sampling site, sampling year or sampling time. Diurnal

fluctuations in the occurrence of OTU5755 could not be discerned based on the Z scores;

both enrichment and de-enrichment were observed. Absence of diurnal linked

fluctuations was also observed for other highly independent OTUs including those from

families which may be associated with gas vacuolate taxa (e.g., Trichocoleusaceae –

OTU5691). Distribution of OTUs across sampling events indicated that the predominant

populations varied temporally (Figure 4). While OTU5691 and OTU5695, both classified

to the family Trichocoleusaceae and comprised a significant portion of cyanobacterial

communities in 2018, they were rarely detected and detected at substantially lower

relative abundances in the 2022 sampling events. 2022 sampling events were found to

have a higher count of unique Prochlorococcaceae OTUs than those detected in 2018.

Figure 3: Temporal Fluctuations in alpha-diversity metrics of cyanobacterial

communities driven by changes in OTU detection (OTU counts; A) and composition

(Shannon Index; B). Z-score normalized relative abundances for highly abundant

operational taxonomic units (>10% of cyanobacterial community) for sampling event to

demonstrate diurnal trends of specific cyanobacterial populations (C).

Figure 4: Temporal distribution of individual OTUs of cyanobacterial families present in

>1% average relative abundances coloured based on family level taxonomic

classification. Sizes of points represent relative abundance of OTU in cyanobacterial

community.

4. Discussion

Diurnal migrations of cyanobacterial populations have been observed in marine

and eutrophic freshwater systems and attributed to both cellular characteristics (Naselli-

Flores et al., 2021) and water column stability (Walsby et al., 1997). Oscillatory diurnal

variation in the distribution of cyanobacterial populations would thus be expected across

depth in the water column. In contrast to trends frequently observed in marine and

eutrophic freshwater systems, we demonstrate that in highly oligotrophic systems, diurnal

fluctuations of cyanobacterial populations are sporadic, but overall bacterial community

composition is conserved temporally. These observations contradict the patterns

commonly described for diurnal migrations of gas vacuolate taxa. Consequently, these

contrasting results—especially the unpredictable nature of cyanobacterial population

distributions—have significant implications for the development of water quality

monitoring programs and associated risk management strategies, including source water

protection and drinking water safety plans.

Evidence of diurnal migrations of colonial and filamentous cyanobacteria has

been reported previously; larger colonies have been shown to migrate across greater

depths more rapidly than smaller ones (Reynolds et al., 1987; Westwood and Ganf,

2004); they also experience faster sinking rates (Ganf, 1974). In both Little Turkey and

Wishart Lakes, cyanobacterial communities were predominantly composed of OTUs

classified to the family Prochlorococcaceae. Notably, the family Prochlorococcaceae

includes small unicellular and colonial taxa that may facilitate their ability to maintain

water column position (Śliwińska-Wilczewska et al., 2018). Furthermore, smaller

colonies also experience more random spatial movement (Chien et al., 2013). The

dominance of small unicellular and colonial taxa of the Prochlorococcaceae family

concurrent with the absence of visible cyanobacterial biomass may thus help to explain

the absence of discernible diurnal trends in the studied cyanobacterial OTUs.

Previous studies have demonstrated that thermal stratification of the water column

plays a significant role in shaping the distribution of cyanobacteria and other

phytoplankton through the creation of discrete zones in the water column (*i.e.*, nutrient

depleted, light rich surface waters *vs.* nutrient rich, light limited deep waters; Reynolds et

al., 1987; Spigel and Imberger, 1987). To investigate the impact of thermal stratification

on cyanobacterial communities in an oligotrophic lake (*i.e*., Little Turkey Lake), samples

were collected in both the epilimnion (water surface) and metalimnion (Secchi depth)

zones (Figure S2). During both sampling events this analysis revealed common OTUs

over time and across depth, thereby suggesting a broader distribution of cyanobacterial

species across the water column rather than restriction to specific zones. In 2022, the

composition of the cyanobacterial communities was highly similar regardless of spatio-

temporal differences. In contrast, in 2018, cyanobacterial communities at the water

surface and Secchi depth were distinguishable, consistent with previous spatial

distribution research conducted in Little Turkey Lake (Cameron et al., 2022). On both

occasions, sampling time did not significantly impact the cyanobacterial community

composition, further supporting the hypothesis that small unicellular and colonial

cyanobacterial populations will exhibit random fluctuations.

Notably, cyanobacterial communities of Little Turkey Lake in 2018 and 2022 did

not exhibit similarity. This result is consistent with seasonal studies conducted in

consecutive years (*i.e*., 2018-2020) in Little Turkey Lake and that reported (i) similar

cyanobacterial community composition between sampling years, and (ii) variability in the

relative abundance of cyanobacteria within the broader bacterial community (Cameron et

al., 2022). Here, cyanobacteria were a statistically significant portion of the bacterial

community in 2022, but not 2018, underscoring the potential for significant interannual

variability in bacterial community structure and the need for further investigation into

environmental drivers of these differences in oligotrophic systems.

Critically, the increased availability and rapidly decreasing costs of next-

generation sequencing tools have eliminated most of the barriers to their wider use in

resource management and the drinking water industry. The insights enabled by these

approaches allow for proactive rather than reactive management of drinking water

supplies (Chapman, 2010). Although amplicon sequencing has revolutionized our ability

to study microbial communities, these data do not quantify abundance of community

members. Rather, they only indicate community composition and relative abundance

(Gloor et al., 2017, 2016) which introduces challenges in the interpretation of changes in

community structure. For example, an observed decrease in relative abundances of

cyanobacteria may only be an artifact of increased relative abundances of sequences

classified to other taxonomic groups and not reflective of an actual decrease in

cyanobacterial populations. Thus, to best inform cyanobacterial community dynamics for

the purposes of resource and risk management, traditional cell enumeration techniques

(e.g., flow cytometry; Patel et al., 2019) or other molecular approaches (e.g., qPCR; Chiu

et al., 2017) that provide community density information are needed in conjunction with

amplicon sequencing.

The sporadic diurnal variations in cyanobacterial populations and the differences

between interconnected oligotrophic lakes that were reported herein emphasize the

challenge in developing lake and reservoir monitoring protocols in absence of sufficient

baseline knowledge of ecological community structures in the specific systems of

interest. Here, cyanobacterial relative abundances fluctuated substantially between the

hours of 9 a.m. and 4 p.m., demonstrating that large, generalized time frames are

insufficient for characterizing cyanobacteria occurrence and dynamics in aquatic systems

and associated threats to water quality and treatability. This work specifically

demonstrated that following typical guidance that limits sample collection to a discrete

depth in stratified lakes or a single timepoint may result in vast underestimation of

cyanobacterial relative abundances. While cyanobacteria sampling protocols and

guidelines do occasionally mention that time and depth of sample collection can affect

outcomes, they are often vague and refer to collection of "surface waters" at times "later

in the day" or from "10 a.m. to 3 p.m." (Table S7), and do not address the potential for or

implications of random non-diurnal migrations of smaller taxa. The present investigation

fills a critical knowledge gap by reporting these insights for oligotrophic systems.

The design of sample collection schemes is closely linked to the utility of the

information that is delivered using these approaches. The analysis provided herein has

delivered clear evidence to demonstrate the need for integrating multi-timepoint, multi-

depth discrete sampling guidance into lake and reservoir monitoring programs to describe

cyanobacterial community dynamics and signal change to inform risk management

associated with the potential for cyanotoxin production. This work further emphasized

that cyanobacteria are present in oligotrophic lakes and their community structure varies

(i) diurnally, (ii) across the depth of the water column, (iii) interannually within the same

lake and (iv) between different lakes that are closely interconnected within the same

watershed. Ignoring this variability and reducing sample numbers can lead to a false

sense of security and missed opportunities to identify and mitigate changes in trophic

status and associated risks such as toxin or taste and odor production, especially in

sensitive, oligotrophic systems.

5. Conclusions

- The potential for diurnal migration should be reflected in cyanobacterial monitoring programs through the inclusion of multiple-sampling times or conducting sampling at an ecologically significant time of day.

- Sampling of lakes should not be restricted to the surface and should include discrete sampling and multiple depths to reflect spatial heterogeneity of cyanobacterial communities present in the water column.

- Cyanobacterial communities are highly dynamic and may vary significantly in composition, prevalence and occurrence times between sampling years at the same sampling site.

- Cyanobacterial monitoring may be enhanced through the incorporation of system characteristics (e.g., thermal stratification) and characterization of communities for design of sampling protocols that are system specific.

Acknowledgements

We acknowledge the support of the *for*Water NSERC Network for Forested Drinking Water Source Protection Technologies [NETGP-494312-16]. We are also grateful for the continued support of Natural Resources Canada and Environment and Climate Change Canada in sample collection at Turkey Lakes Watershed Research Station.

6, References

Bisanz, J.E., 2018. qiime2R: Importing QIIME2 artifacts and associated data into R

sessions.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith,

G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz,

J.E., Bittinger, K., Brejnrod, A., Brislawn, C.J., Brown, C.T., Callahan, B.J.,

Caraballo-Rodríguez, A.M., Chase, J., Cope, E.K., da Silva, R., Diener, C.,

Dorrestein, P.C., Douglas, G.M., Durall, D.M., Duvallet, C., Edwardson, C.F.,

Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J.M., Gibbons, S.M., Gibson,

D.L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H.,

Huttenhower, C., Huttley, G.A., Janssen, S., Jarmusch, A.K., Jiang, L.,

Kaehler, B.D., Kang, K. bin, Keefe, C.R., Keim, P., Kelley, S.T., Knights, D.,

Koester, I., Kosciolek, T., Kreps, J., Langille, M.G.I., Lee, J., Ley, R., Liu,

Y.X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B.D.,

McDonald, D., McIver, L.J., Melnik, A. v., Metcalf, J.L., Morgan, S.C.,

Morton, J.T., Naimey, A.T., Navas-Molina, J.A., Nothias, L.F., Orchanian,

S.B., Pearson, T., Peoples, S.L., Petras, D., Preuss, M.L., Pruesse, E.,

Rasmussen, L.B., Rivers, A., Robeson, M.S., Rosenthal, P., Segata, N., Shaffer,

M., Shiffer, A., Sinha, R., Song, S.J., Spear, J.R., Swafford, A.D., Thompson,

L.R., Torres, P.J., Trinh, P., Tripathi, A., Turnbaugh, P.J., Ul-Hasan, S., van der

Hooft, J.J.J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel, M.,

Walters, W., Wan, Y., Wang, M., Warren, J., Weber, K.C., Williamson,

C.H.D., Willis, A.D., Xu, Z.Z., Zaneveld, J.R., Zhang, Y., Zhu, Q., Knight, R.,

Caporaso, J.G., 2019. Reproducible, interactive, scalable and extensible

484   microbiome data science using QIIME 2. Nat Biotechnol 37, 852–857.

485   https://doi.org/10.1038/s41587-019-0209-9

486   Bukata, R.P., Jerome, J.H., Bruton, J.E., 1988. Relationships Among Secchi Disk

487   Depth, Beam Attenuation Coefficient, and Irradiance Attenuation Coefficient

488   for Great Lakes Waters. J Great Lakes Res 14, 347–355.

489   https://doi.org/10.1016/S0380-1330(88)71564-6

490   Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes,

491   S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon

492   data. Nat Methods 13, 581–583. https://doi.org/10.1038/nmeth.3869

493   Cameron, E.S., Müller, K.M., Stone, M., Buttle, J., Leach, J., Webster, K., Emelko,

494   M.B., 2022. Early Seasonal Increases and Persistence in Relative Abundance of

495   Potentially Toxic Cyanobacteria: Concerning Impacts of Extended Ice-Free

496   Periods in Northern Temperate Lakes. bioRxiv 2022.12.20.521158.

497   https://doi.org/10.1101/2022.12.20.521158

498   Cameron, E.S., Schmidt, P.J., Tremblay, B.J.M., Emelko, M.B., Müller, K.M., 2021.

499   Enhancing diversity analysis by repeatedly rarefying next generation

500   sequencing data describing microbial communities. Sci Rep 11.

501   https://doi.org/10.1038/s41598-021-01636-1

502   Chapman, A.D., 2010. Cyanobacteria, in: Algae Source to Treatment - Manual of

503   Water Supply Practices M57. American Water Works Association, pp. 125–

504   146.

Chien, Y.C., Wu, S.C., Chen, W.C., Chou, C.C., 2013. Model simulation of diurnal

vertical migration patterns of different-sized colonies of microcystis employing

a particle trajectory approach. Environ Eng Sci 30, 179–186.

https://doi.org/10.1089/ees.2012.0318

Chiu, Y.T., Chen, Y.H., Wang, T.S., Yen, H.K., Lin, T.F., 2017. A qPCR-based tool

to diagnose the presence of harmful cyanobacteria and cyanotoxins in drinking

water sources. Int J Environ Res Public Health 14.

https://doi.org/10.3390/ijerph14050547

Emelko, M.B., Silins, U., Bladon, K.D., Stone, M., 2011. Implications of land

disturbance on drinking water treatability in a changing climate: Demonstrating

the need for " source water supply and protection" strategies. Water Res 45,

461–472. https://doi.org/10.1016/j.watres.2010.08.051

Emelko, M.B., Stone, M., Silins, U., Allin, D., Collins, A.L., Williams, C.H.S.,

Martens, A.M., Bladon, K.D., 2016. Sediment-phosphorus dynamics can shift

aquatic ecology and cause downstream legacy effects after wildfire in large

river systems. Glob Chang Biol 22, 1168–1184.

https://doi.org/10.1111/gcb.13073

Frempong, E., 1981. Diel Variation in the Abundance, Vertical Distribution, and

Species Composition of Phytoplankton in a Eutrophic English Lake. J Ecol 69,

919. https://doi.org/10.2307/2259645

Ganf, G.G., 1974. Diurnal Mixing and the Vertical Distribution of Phytoplankton in

526   a Shallow Equatorial Lake (Lake George , Uganda). Journal of Ecology 62,

527   611–629.

528   Gilbert, J.A., Field, D., Swift, P., Thomas, S., Cummings, D., Temperton, B.,

529   Weynberg, K., Huse, S., Hughes, M., Joint, I., Somerfield, P.J., Mühling, M.,

530   2010. The taxonomic and functional diversity of microbes at a temperate

531   coastal site: A "multi-omic" study of seasonal and diel temporal variation.

532   PLoS One 5. https://doi.org/10.1371/journal.pone.0015545

533   Gloor, G.B., Macklaim, J.M., Pawlowsky-Glahn, V., Egozcue, J.J., 2017.

534   Microbiome datasets are compositional: And this is not optional. Front

535   Microbiol 8, 1–6. https://doi.org/10.3389/fmicb.2017.02224

536   Gloor, G.B., Macklaim, J.M., Vu, M., Fernandes, A.D., 2016. Compositional

537   uncertainty should not be ignored in high-throughput sequencing data analysis.

538   Austrian Journal of Statistics 45, 73–87. https://doi.org/10.17713/ajs.v45i4.122

539   Gu, Z., 2022. Complex heatmap visualization. iMeta 1.

540   https://doi.org/10.1002/imt2.43

541   Harke, M.J., Steffen, M.M., Gobler, C.J., Otten, T.G., Wilhelm, S.W., Wood, S.A.,

542   Paerl, H.W., 2016. A review of the global ecology, genomics, and

543   biogeography of the toxic cyanobacterium, Microcystis spp. Harmful Algae 54,

544   4–20. https://doi.org/10.1016/j.hal.2015.12.007

545   Huisman, J., Codd, G.A., Paerl, H.W., Ibelings, B.W., Verspagen, J.M.H., Visser,

546   P.M., 2018. Cyanobacterial blooms. Nat Rev Microbiol 16, 471–483.

547   Hunter, P.D., Tyler, A.N., Willby, N.J., Gilvear, D.J., 2008. The spatial dynamics of

548   vertical migration by Microcystis aeruginosa in a eutrophic shallow lake: A

549   case study using high spatial resolution time-series airborne remote sensing.

550   Limnol Oceanogr 53, 2391–2406. https://doi.org/10.4319/lo.2008.53.6.2391

551   Ibelings, B.W., Mur, L.R., Walsby, A.E., 1991. Diurnal changes in buoyancy and

552   vertical distribution in populations of Microcystisin two shallow lakes. J

553   Plankton Res 13, 419–436. https://doi.org/10.1093/plankt/13.2.419

554   Jalili, F., Trigui, H., Guerra Maldonado, J.F., Dorner, S., Zamyadi, A., Shapiro, B.J.,

555   Terrat, Y., Fortin, N., Sauvé, S., Prévost, M., 2021. Can Cyanobacterial

556   Diversity in the Source Predict the Diversity in Sludge and the Risk of Toxin

557   Release in a Drinking Water Treatment Plant? Toxins (Basel) 13.

558   https://doi.org/10.3390/TOXINS13010025

559   Jeffries, D.S., Kelso, J.R.M., Morrison, I.K., 1988. Physical, Chemical and

560   Biological Characteristics of the Turkey Lakes Watershed, Central Ontario,

561   Canada. Canadian Journal of Fisheries and Aquatic Sciences 45, 3–13.

562   Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software

563   version 7: Improvements in performance and usability. Mol Biol Evol 30, 772–

564   780. https://doi.org/10.1093/molbev/mst010

565   Le, K.T.N., Maldonado, J.F.G., Goitom, E., Trigui, H., Terrat, Y., Nguyen, T.L.,

566   Husk, B., Shapiro, B.J., Sauvé, S., Prévost, M., Dorner, S., 2022. Shotgun

567   Metagenomic Sequencing to Assess Cyanobacterial Community Composition

568   following Coagulation of Cyanobacterial Blooms. Toxins (Basel) 14.

569   https://doi.org/10.3390/toxins14100688

570   Lefler, F.W., Berthold, D.E., Laughinghouse, H.D., 2023. Cyanoseq: A database of

571   cyanobacterial 16S rRNA gene sequences with curated taxonomy. J Phycol.

572   https://doi.org/10.1111/jpy.13335

573   Li, H., Qin, C., He, W., Sun, F., Du, P., 2023. Learning and inferring the diurnal

574   variability of cyanobacterial blooms from high-frequency time-series satellite-

575   based observations. Harmful Algae 123.

576   https://doi.org/10.1016/j.hal.2023.102383

577   McMurdie, P.J., Holmes, S., 2012. Phyloseq: A bioconductor package for handling

578   and analysis of high-throughput phylogenetic sequence data, in: Pacific

579   Symposium on Biocomputing.

580   Naselli-Flores, L., Zohary, T., Padisák, J., 2021. Life in suspension and its impact on

581   phytoplankton morphology: an homage to Colin S. Reynolds. Hydrobiologia

582   848, 7–30. https://doi.org/10.1007/s10750-020-04217-x

583   Nguyen, L.T., Schmidt, H.A., Von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: A fast

584   and effective stochastic algorithm for estimating maximum-likelihood

585   phylogenies. Mol Biol Evol 32, 268–274.

586   https://doi.org/10.1093/molbev/msu300

587   Olli, K., 1999. Diel vertical migration of phytoplankton and heterotrophic flagellates

588   in the Gulf of Riga. Journal of Marine Systems 23, 145–163.

589   https://doi.org/10.1016/S0924-7963(99)00055-X

Paerl, H.W., 2014. Mitigating harmful cyanobacterial blooms in a human- and

climatically-impacted world. Life 4, 988–1012.

https://doi.org/10.3390/life4040988

Patel, R., de Oliveira, A., Newby, R., Chu, T., 2019. Flow cytometric analysis of

freshwater cyanobacteria: A case study. Water (Switzerland) 11.

https://doi.org/10.3390/w11071422

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J.,

Glöckner, F.O., 2013. The SILVA ribosomal RNA gene database project:

Improved data processing and web-based tools. Nucleic Acids Res 41, 590–

596. https://doi.org/10.1093/nar/gks1219

Reinl, K.L., Brookes, J.D., Carey, C.C., Harris, T.D., Ibelings, B.W., Morales-

Williams, A.M., De Senerpont Domis, L.N., Atkins, K.S., Isles, P.D.F.,

Mesman, J.P., North, R.L., Rudstam, L.G., Stelzer, J.A.A., Venkiteswaran, J.J.,

Yokota, K., Zhan, Q., 2021. Cyanobacterial blooms in oligotrophic lakes:

Shifting the high-nutrient paradigm. Freshw Biol.

https://doi.org/10.1111/fwb.13791

Reynolds, C.S., Oliver, R.L., Walsby, A.E., 1987. Cyanobacterial dominance: The

role of buoyancy regulation in dynamic lake environments. N Z J Mar

Freshwater Res 21, 379–390. https://doi.org/10.1080/00288330.1987.9516234

Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: A

versatile open source tool for metagenomics. PeerJ 2016.

https://doi.org/10.7717/peerj.2584

Schmidt, P.J., Cameron, E.S., Müller, K.M., Emelko, M.B., 2021. Ensuring that

fundamentals of quantitative microbiology are reflected in microbial diversity

analyses based on next-generation sequencing. bioRxiv.

https://doi.org/https://doi.org/10.1101/2021.06.19.449110

Scholz, J., Brahney, J., 2022. Evidence for multiple potential drivers of increased

phosphorus in high-elevation lakes. Science of the Total Environment 825.

https://doi.org/10.1016/j.scitotenv.2022.153939

Shahraki, A.H., Chaganti, S.R., Heath, D.D., 2020. Diel Dynamics of Freshwater

Bacterial Communities at Beaches in Lake Erie and Lake St. Clair, Windsor,

Ontario. Microb Ecol 81, 1–13.

Shannon, C.E., 1948. A Mathematical Theory of Communication. The Bell System

Technical Journal 27, 623–656.

Silins, U., Bladon, K.D., Kelly, E.N., Esch, E., Spence, J.R., Stone, M., Emelko,

M.B., Boon, S., Wagner, M.J., Williams, C.H.S., Tichkowsky, I., 2014. Five-

year legacy of wildfire and salvage logging impacts on nutrient runoff and

aquatic plant, invertebrate, and fish productivity. Ecohydrology 7, 1508–1523.

https://doi.org/10.1002/eco.1474

Śliwińska-Wilczewska, S., Maculewicz, J., Felpeto, A.B., Latała, A., 2018.

Allelopathic and bloom-forming picocyanobacteria in a changing world. Toxins

(Basel) 10, 1–20. https://doi.org/10.3390/toxins10010048

Spigel, R.H., Imberger, J., 1987. Mixing processes relevant to phytoplankton

dynamics in lakes. N Z J Mar Freshwater Res 21, 361–377.

https://doi.org/10.1080/00288330.1987.9516233

Sterner, R.W., Reinl, K.L., Lafrancois, B.M., Brovold, S., Miller, T.R., 2020. A first

assessment of cyanobacterial blooms in oligotrophic Lake Superior. Limnol

Oceanogr 65, 2984–2998. https://doi.org/10.1002/lno.11569

Stoddard, J.L., van Sickle, J., Herlihy, A.T., Brahney, J., Paulsen, S., Peck, D. v.,

Mitchell, R., Pollard, A.I., 2016. Continental-Scale Increase in Lake and

Stream Phosphorus: Are Oligotrophic Systems Disappearing in the United

States? Environ Sci Technol 50, 3409–3415.

https://doi.org/10.1021/acs.est.5b05950

Tsai, K.P., Uzun, H., Karanfil, T., Chow, A.T., 2017. Dynamic Changes of

Disinfection Byproduct Precursors following Exposures of Microcystis

aeruginosa to Wildfire Ash Solutions. Environ Sci Technol 51, 8272–8282.

https://doi.org/10.1021/acs.est.7b01541

Valadez-Cano, C., Hawkes, K., Calvaruso, R., Reyes-Prieto, A., Lawrence, J., 2022.

Amplicon-based and metagenomic approaches provide insights into toxigenic

potential in understudied Atlantic Canadian lakes. Facets 7, 194–214.

https://doi.org/10.1139/facets-2021-0109

von Orgies-Rutenberg, M., Rolfes, C., Eckel, T., Quiroz, A., Skalbeck, J., Riley, D.,

Sander, H., 2018. Diurnal vertical migration of cyanobacteria and chlorophyta

in eutrophied shallow freshwater lakes. Fundamental and Applied Limnology

191. https://doi.org/10.1127/fal/2017/1021

Vu, H.P., Nguyen, L.N., Zdarta, J., Nga, T.T.V., Nghiem, L.D., 2020. Blue-Green

Algae in Surface Water: Problems and Opportunities. Curr Pollut Rep 6, 105–

122. https://doi.org/10.1007/s40726-020-00140-w

Walsby, A.E., Hayes, P.K., Boje, R., Stal, L.J., 1997. The selective advantage of

buoyancy provided by gas vesicles for planktonic cyanobacteria in the Baltic

Sea. New Phytologist 136, 407–417. https://doi.org/10.1046/j.1469-

8137.1997.00754.x

Walters, W., Hyde, E.R., Berg-Lyons, D., Ackermann, G., Humphrey, G., Parada,

A., Gilbert, J.A., Jansson, J.K., Caporaso, J.G., Fuhrman, J.A., Apprill, A.,

Knight, R., 2015. Improved Bacterial 16S rRNA Gene (V4 and V4-5) and

Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial

Community Surveys. mSystems 1, e0009-15.

https://doi.org/10.1128/mSystems.00009-15.Editor

Wang, S., Zhang, X., Wang, C., Chen, N., 2023. Temporal continuous monitoring of

cyanobacterial blooms in Lake Taihu at an hourly scale using machine learning.

Science of the Total Environment 857.

https://doi.org/10.1016/j.scitotenv.2022.159480

Westwood, K.J., Ganf, G.G., 2004. Effect of mixing patterns and light dose on

growth of Anabaena circinalis in a turbid, lowland river. River Res Appl 20,

115–126. https://doi.org/10.1002/rra.725

Woolway, R.I., Merchant, C.J., 2019. Worldwide alteration of lake mixing regimes

in response to climate change. Nat Geosci 12, 271–276.

https://doi.org/10.1038/s41561-019-0322-x

Xue, K., Ma, R., Shen, M., Wu, J., Hu, M., Guo, Y., Cao, Z., Xiong, J., 2023.

Horizontal and vertical migration of cyanobacterial blooms in two eutrophic

lakes observed from the GOCI satellite. Water Res 240, 120099.

https://doi.org/10.1016/j.watres.2023.120099

Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T.,

Peplies, J., Ludwig, W., Glöckner, F.O., 2014. The SILVA and "all-species

Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids Res 42,

D643–D648. https://doi.org/10.1093/nar/gkt1209