Title - Life history speed, population disappearances and noise-induced ratchet effects

Authors:

Christopher J. Greyson-Gaito[*1], Gabriel Gellner[1], Kevin S. McCann[1]

Affiliations:

1. Department of Integrative Biology, University of Guelph, Guelph, Ontario, Canada

*Corresponding Author – Email: christopher@greyson-gaito.com (CJGG)

ORCID:

CJGG – 0000-0001-8716-0290

GG - 0000-0001-8170-1463

KSM – 000-0001-6031-7913

Abstract

Nature is replete with variation in the body sizes, reproductive output, and generation times of species that produce life history responses known to vary from small and fast to large and slow. Although researchers recognize that life history speed likely dictates fundamental processes in consumer-resource interactions like productivity and stability, theoretical work remains incomplete in this critical area. Here, we examine the role of life history speed on consumer-resource interactions by using a well used mathematical approach that manipulates the speed of the consumer's growth rate in a consumer-resource interaction. Importantly, this approach holds the isocline geometry intact allowing us to assess the impacts of altered life history speed on stability (coefficient of variation, CV) without changing the underlying qualitative dynamics. Although slowing life history can be initially stabilizing, we find that in stochastic settings slowing ultimately drives highly destabilizing population disappearances, especially under reddened noise. Our results suggest that human-driven reddening of noise may decrease species stability because the autocorrelation of red noise enlarges the period and magnitude of perturbations, overwhelming a species' natural compensatory responses via a ratchet-like effect. This ratchet-like effect then pushes species' population dynamics far away from equilibria that can lead to precipitous local extinction.

Keywords

Slow-fast, life history, stability, consumer-resource, red noise, stochasticity

Introduction

Life history is a fundamental axis of variation within nature. Researchers have argued cogently that this variation tends to follow a "slow-fast" continuum where slow life histories have small population growth rates (low r), large body sizes, and long generation times while fast life histories have high population growth rates, small body sizes, and short generation times [1–4]. Indeed, empirical work has found remarkably consistent relationships that suggest body size is a key attribute of life history speed (Fig. 1a; [5]). These covarying traits along the "slow-fast" continuum can impact many ecological processes and properties including population growth, ecosystem productivity and stability [1,6]. Given the changing nature of abiotic variation under climate change [7,8] and the increasing propensity for fast life histories under global change [9,10], understanding how life history speed governs the ability for organisms and whole communities to persist (i.e., retain densities well above zero densities) is critical.

Theory has begun to unpack how population life history "speed" may tie into stability, with initial findings of fast and slow life histories amplifying or muting noise respectively. Specifically, research has found that because of the high population growth rates, faster life history organisms tend to produce over-compensatory dynamics (i.e., overshoot the equilibrium) and instability compared to slower organisms [11,12]. In contrast to fast organisms, slow organisms, with longer lives, buffer noise and thus maintain high stability. Consistent with this, empirical research has found negative correlations between population variability (one measure of stability) and both body size and generation time, and positive correlations between population variability and growth rates across multiple taxa and kingdoms [13–16]. Taken altogether, larger and slower organisms appear capable of buffering noise better than small and fast organisms.

Nonetheless, while large slow-growth organisms may be able to buffer perturbations, this same potentially stabilizing slow growth response can be destabilizing when perturbations are large enough to push population densities far away from the equilibrium [17,18]. Once a slow-growth species is pushed away from the equilibrium they become subject to the vagaries of different nonlinear dynamics (hereafter referred to as non-local nonlinear dynamics, see Box 1) that can fundamentally alter the outcome of the system including taking the dynamics close to zero or local extinction [17,18]. On the other hand, small fast-growing species may be less likely to be pushed far away from equilibrium because their rapid growth responses can keep them closer to equilibrium and local dynamic properties. These results suggest complex slow-fast life history stability responses such that some aspects of life history speed are beneficial (e.g., large and slow can buffer a perturbation) but also produces costs (e.g., large and slow growth can be detrimental once the population is pushed away from local equilibrium dynamics). Consequently, more research is required to understand how stability manifests along the slow-fast life history continuum.

Another related factor that will impact stability along the slow-fast continuum is the color of the stochastic noise (see Box 1 for definition). Climate change is known to be reddening environmental noise by increasing the spatial and temporal autocorrelation of climate

variables [7,8] which then can act to lengthen periods of suboptimal conditions for organisms
[19]. Under extended suboptimal conditions, populations can be driven to smaller and smaller
numbers thus often hastening local extinction [19]. In a sense, these extended suboptimal
conditions are akin to large perturbations (although each individual noise event is small). By
pushing an organism's dynamics far from equilibria, red noise is then also likely to cause non-
local nonlinear dynamic outcomes. In this latter case, although not well explored, these
autocorrelated perturbations could even lead beneficial perturbations to drive strong nonlinear
effects that eventually produce local extinction (e.g., consumer increases in a consumer-
resource interaction can lead to overshoot and dangerously low densities). These collective
theoretical results suggest a complex range of dynamical responses that demand further
understanding.

Towards understanding the complex responses of stochastic models, Higgins et al. [20]
pointed out that the dynamics of an underlying deterministic model are critical to
understanding the full stochastically forced model. Through elegantly decomposing a
stochastic population model of Dungeness crab into a deterministic model (the deterministic
skeleton, see Box 1) and a stochastic process, Higgins et al. [20] found that the novel
responses produced from the interaction of the density-dependent dynamics and stochasticity
matched empirical observations. Similarly, qualitatively different dynamics of the deterministic
consumer-resource (C-R) model are known to yield qualitatively different stochastic
dynamics. For example, weakly interacting deterministic C-R models are known to produce
non-excitable monotonic dynamics (i.e., the deterministic equilibrium is stable with real
eigenvalues) that when perturbed tend to simply produce noisy equilibrium dynamics [21,22].
As the stochastic dynamics suggest, the monotonic deterministic skeleton is aptly named
non-excitable. In contrast, as the C-R interactions are strengthened, the deterministic
equilibrium begins to show excitable dynamics with decaying cycles to the equilibrium (i.e.,
the deterministic equilibrium is stable with complex eigenvalues). Here, stochastic
perturbations now tend to excite the underlying density dependent frequencies of the
deterministic C-R skeleton producing stochastic cycles or quasi-cycles [22]. Again, the
dynamics of the underlying deterministic skeleton is critical to understanding the full
stochastic model.

Here, towards synthesizing slow-fast life history stability responses to perturbations, we
examine how stability (i.e., defined as per many empirical studies as the coefficient of
variation, CV) is influenced by life history speed along the slow-fast continuum. Specifically,
we examine the C-R interaction – a fundamental building block of whole communities –
because it has been well described by allometric arguments that are conducive to slow-fast
theory [23,24]. We alter the consumer's life history speed using a common mathematical
scaler (ε) employed for the analysis of slow-fast mathematical systems [25]. The parameter, ε,
allows us to mimic known variation in growth rates (Fig. 1a,b) and to alter the life history
speed while maintaining the underlying qualitative dynamical conditions (i.e. the isocline
geometry of the C-R model is preserved when changing ε alone). We employ this model over
a range of white to red noise perturbations (Fig. 1c). Finally, we vary the C-R interaction
strength (i.e. energy flux via consumer conversion efficiency, e) to explore both the non-
excitable and the excitable deterministic skeletons (Fig. 1c). In doing so, we cover a broad
range in the underlying deterministic skeleton and seek a general answer to the role slow-fast

life histories play in stability. Overall, we show that life history speed drives a range of stability
responses to noise with slow life histories especially sensitive to non-local nonlinear dynamics
and population disappearances when perturbed by reddened noise.

Figure 1 a) Empirical relationship between body size and population growth rate (r_{max}) show
the existence of slow-fast life histories (e.g., Savage et al. [1]). b) In our consumer- resource
model, we use the parameter ε to scale the consumer's growth rate to replicate the empirical
relationship between body size and growth rate. ε scales the consumer equation such that the
consumer growth rate decreases with increasing $1/\varepsilon$ producing a fast to slow life history
continuum. This method simultaneously holds the isocline arrangement constant and so is an
experiment that changes "life history speed" in and of itself. c) To explore the impacts of life
history speed, we will examine how slowing the consumer's life history through $1/\varepsilon$ impacts
the stability of the C-R interaction (using coefficient of variation). Stochastic perturbations,
from white to reddened, will be added to the consumer. Because we know that the underlying
deterministic skeleton interacts with noise in different ways [21,22], we will set the efficiency
parameter (e) of the C-R model to produce the non-excitable (i.e., real eigenvalues,
monotonic dynamics) and excitable (i.e., complex eigenvalues, oscillatory decay dynamics)
deterministic skeletons.

Box 1 Key terms and definitions

Deterministic skeleton: The deterministic skeleton mathematically describes the processes of interest in the system being modeled. These are usually a dynamical system (e.g., ordinary differential equation or difference equations) and are effectively the model without stochastic processes [26].

Noise/Stochasticity: In ecology, noise is considered to be whatever we do not understand in the system [26,27]. Modelers add stochastic (random) noise to the deterministic skeleton. The interaction between the deterministic skeleton and noise can generate radically different dynamical responses, providing key insights into ecological phenomena (see review by Coulson et al. [27]).

Red & white noise: Noise from time points close together that are similar to each other (positive autocorrelation) is classified as red noise. Red noise is the opposite of blue noise where noise from time points close together are completely different (negatively autocorrelated). White noise lies between red and blue noise, that is, noise from time points close together are neither always similar nor always different (i.e. neither positively nor negatively autocorrelated).

Stability: Here, we define stability in terms of persistence where increasing the lower limit of population density away from zero increases stability/persistence (general stability in McCann [28]). Notably, the lower a population's density is the greater the likelihood that a stochastic process can push it to local extinction. The coefficient of variation (CV) is a common theoretical and empirical metric that normalizes standard deviation relative to the mean and so is a good metric for general stability (see Figure 1c). A high SD relative to a low mean means that the population necessarily attains a very low density where risk of collapse is high.

Non-local nonlinear dynamics: Nonlinear dynamics occur in systems that react disproportionately to initial conditions or a small perturbation. These dynamics can include chaos and limit cycles. In this study, we differentiate between local and non-local nonlinear dynamics. Local nonlinear dynamics are dynamics when system trajectories are close to the equilibria in phase space. Non-local nonlinear dynamics are dynamics when system trajectories are far from equilibria.

Quasi-cycle: Quasi-cycles are a type of dynamic behaviour that occur when stochasticity resonates with damped oscillations surrounding an interior equilibrium [21,26,29]. Frequencies in the stochastic noise that most closely resemble the period of the damped oscillations are amplified. Thus, a power spectrum would show all frequencies with the frequency of the damped oscillations having the highest power [21].

Quasi-canard: Quasi-canards are another type of dynamic behaviour that occur when stochasticity induces a pattern similar to a deterministic canard. A deterministic canard occurs when a system's solution follows an attracting state space area (manifold), passes over a critical point along this manifold, and then follows a repelling manifold [30]. In the C-R model,

the canard solution slowly follows the resource isocline (the attracting manifold) until the maximum point of the isocline is reached (the critical point), then the solution quickly jumps to the consumer axis before slowly following the consumer axis (the repelling manifold). Finally, the solution quickly jumps back to the resource isocline and repeats the canard cycle. When stochasticity is introduced, trajectories combine both small oscillations around the equilibria and large relaxation oscillations qualitatively similar to deterministic canards. Generally, these patterns are called mixed mode oscillations [30]. However, we use the term quasi-canards in this study to evoke the importance of stochastic noise similar to the importance of stochastic noise in quasi-cycles.

Methods

Model

In this study, we used the classic Rosenzweig-MacArthur consumer-resource (C-R) model

$$\frac{dR}{dt} = r R \left(1 - \frac{R}{k}\right) - \frac{a R C}{1 + a h R} \tag{1}$$

$$\frac{dC}{dt} = \left(\frac{e a R C}{1 + a h R} - m C\right) \tag{2}$$

where r is the intrinsic growth rate of the resource (R), k is the carrying capacity of the resource, a is the attack rate of the consumer (C), e is the efficiency or conversion rate of consumed resources into new consumers, h is the handling time, and m is the consumer mortality.

Previous exploration has illustrated the multiple bifurcations and different dynamics of this model [14,31]. The bifurcations are dictated by the energy entering the consumer versus exiting the consumer (the energy flux of the consumer) or as an equation:

$$\frac{K a e}{m} \tag{3}$$

As the energy entering the consumer increases relative to the energy exiting the consumer (the numerator of (3) increases relative to the denominator), the first bifurcation is a transcritical bifurcation (switching of stable points) creating an interior stable equilibrium (see Figure 1c). At first, this equilibrium is monotonic but then exhibits damped oscillations. The next bifurcation is a Hopf bifurcation (the stable interior equilibrium becomes unstable) leading to period cycling. The mathematics underlying the shift from monotonic to dampened oscillations for the stable interior equilibrium is a shift in eigenvalues from real (no imaginary part) to complex (with an imaginary part). Gellner et al. [22] termed these two regions the non-excitable region and the excitable region respectively (see Figure 1c).

156 Now, to manipulate the life history of the consumer, we added the parameter ε to the
157 consumer's equation. The consumer is fast when ε is one and slow when ε is tiny.

158
$$\frac{dC}{dt} = \varepsilon \left(\frac{e\,a\,R\,C}{1+a\,h\,R} - m\,C \right) \qquad (4)$$

Because ε scales the consumer's intrinsic growth rate ($e \cdot a_{max} - m$ where a_{max} is $1/h$, see S.I.
Section "Scaling of the consumer's growth rate by ε") but preserves the isocline geometry of
the C-R model, we can manipulate the life history of the consumer along a slow-fast
continuum (relative to the resource) (Figure 1a) while controlling the underlying deterministic
skeleton.

For all analyses, r = 2.0, k = 3.0, a = 1.1, h = 0.8, and m = 0.4.

Stability along the slow-fast continuum

We used the coefficient of variation to measure stability as the consumer's life history was
varied from fast to slow ($1/\varepsilon$ was varied from 1.0 to 1000). We used a flow-kick approach to
add stochasticity (versus a stochastic ordinary differential equation approach) [32]. In this
flow-kick approach, we kick the consumer variable every unit of time and let the model flow
(integrate) without noise until the next kick. The kick is generated from normally distributed
noise (with mean 0.0 and standard deviation 0.01). For each value of $1/\varepsilon$, we ran 50
simulations of 24,000 time units (package DifferentialEquations.jl v6.20.0, Algorithms: Vern7 &
Rodas4 with automatic stiffness detection). From these simulations for each value of $1/\varepsilon$, we
calculated the average coefficient of variation (CV) after removing the first 2000 time units.
We did this for the efficiency values (e) of 0.5 (non-excitable deterministic dynamics with real
eigenvalues) and 0.71 (excitable deterministic dynamics with complex eigenvalues). Note,
any simulations that landed on the axial solution (where the resource isocline intersects the
Consumer = 0 axis) were removed from the average coefficient of variation calculation.

To unpack the CV result as the consumer's life history is slowed, we examined the dynamical
behaviours that drove the change in CV. Three general dynamical behaviors occur depending
on whether the C-R interaction is excitable (complex eigenvalues) or non-excitable
(completely real eigenvalues) (see Figure 2a). The first dynamical behaviour is quasi-cycling
and occurs only when the C-R interaction is excitable (see Box 1 and Figure 2b). Quasi-
cycling is when stochastic perturbations resonate with the excitability of the C-R model to
extend the range of cyclic dynamics (see Box 1 and [21,22]). Compared to the deterministic
limit cycles, these quasi-cycles generally do not threaten persistence because quasi-cycles
exhibit smaller variation. Nevertheless, because slowing the consumer's life history reduces
the excitability of the C-R interaction (see S.I. Section "Slowing the consumer decreases
excitability"), we may see an impact of slowing the consumer's life history on the stability (CV)
of the C-R interaction. The second dynamical behavior is quasi-canards which can happen for
both excitable and non-excitable C-R interactions (see Box 1 and Figure 2b). A quasi-canard
is similar to a deterministic canard. A deterministic canard occurs after the Hopf bifurcation
(i.e. to the left of the resource isocline maximum) and does not require stochastic noise. In
contrast, a quasi-canard occurs before the Hopf bifurcation (i.e. to the right of the resource
isocline maximum) and requires stochastic noise (see Box 1). Because quasi-canard

trajectories are quite large, stability will decrease markedly (i.e. CV will increase). The third
dynamical behaviour is what we term stretched wandering and occurs for non-excitable C-R
interactions only (see Figure 2b). The dynamics are stretched along the resource isocline with
no obvious cycling.

Figure 2 a) Table of dynamical behaviours when the C-R interaction is non-excitable and
excitable and when the consumer's life history is fast or slow. b) Phase diagrams and time
series of quasi-cycles (yellow), quasi-canards (purple), and stretched wandering (green).
Dashed vertical lines in the phase diagrams denote where the deterministic Hopf bifurcation
occurs. Note, the standard deviation of the noise was increased to 0.05 to help emphasize
what quasi-cycles, quasi-canards, and stretched wandering look like.

Quasi-cycles

To examine how slowing the consumer's life history impacted quasi-cycles, we ran 100
simulations each of the C-R model with $1/\varepsilon$ values of 1 and 10 and where, again using the
flow-kick approach, the consumer variable was perturbed every time unit with normally

distributed noise (with mean 0.0 and standard deviation 0.01) (package
DifferentialEquations.jl v6.20.0, Algorithms: Vern7 & Rodas4 with automatic stiffness
detection). To examine the initial decrease in CV as the consumer's life history was slowed,
we chose $1/\varepsilon$ values of 1 and 10. We calculated the autocorrelation of the last 1000 time units
of each simulation for lags between 0 and 40 (function autocor, package StatsBase v0.33.13).
We then calculated the average autocorrelation value for each lag across the 100 simulations
for each simulation set with a different $1/\varepsilon$ value. We did this for the efficiency values (*e*) of 0.5
(non-excitable) and 0.71 (excitable). As suggested by Pineda-Krch et al. [21], we used
autocorrelation to identify the occurrence of quasi-cycles. An autocorrelation function (ACF)
from quasi-cycles would show pronounced amplitude oscillations that decrease in amplitude
with increasing lags. No oscillations in the ACF indicate that quasi-cycles are not occurring
and constant ACF oscillations indicate noisy period cycling. An ACF with smaller damped
oscillations indicates that the periodicity of the quasi-cycles is being removed. Note, that in
our ACF figures, the mean of each time series was removed from each time point value. In
the ACF figures in Pineda-Krch et al. [21], the mean of each time series was not removed
from each time point value. Thus, our figures look different (our ACF scale goes from -1 to 1
whereas Pineda-Krch et al.'s [21] scale goes from 0.95 to 1). Our method makes spotting the
damped oscillations easier.

Quasi-canards

To examine how slowing the consumer's life history impacted quasi-canards, we examined
the prevalence of quasi-canards in stochastically perturbed simulations of the C-R model with
varying ε and efficiency values. First, we examined the prevalence of quasi-canards under
white noise stochasticity. We ran 1000 simulations with 24,000 time units for each value of $1/\varepsilon$
varied from 6.667 to 1000, where, using the flow-kick approach, the consumer variable was
perturbed every time unit with normally distributed noise (with mean 0.0 and standard
deviation 0.01) (package DifferentialEquations.jl v6.20.0, Algorithms: Vern7 & Rodas4 with
automatic stiffness detection). Our lower bound for $1/\varepsilon$ was set at 6.667 because no quasi-
canards can be found below this point. This was done for efficiency values of 0.5 (non-
excitable) and 0.71 (excitable). For each combination of ε and efficiency value, we calculated
the proportion of simulations that exhibited at least one quasi-canard. To calculate the
proportion of simulations with quasi-canards, we created an algorithm to check whether a
time series contained a quasi-canard. More details of this algorithm can be found in the
Supporting Information (S.I. section "Explanation of quasi-canard finder algorithm"), but in
short the algorithm includes a return map at the maximum point of the resource isocline which
canards and quasi-canards must pass through. The algorithm also includes boxes along the
attracting and repelling manifolds (the right side of the resource isocline and the consumer
axis respectively) through which a quasi-canard should pass.

Second, we examined how reddened noise impacts the prevalence of quasi-canards. For two
values of $1/\varepsilon$ (12.66 and 250.0) and for two values of efficiency (0.5 and 0.71), we varied the
stochasticity from white to red noise and measured the proportion of simulations that
exhibited at least one quasi-canards. Note, we also measured the proportion of simulations
that exhibited no quasi-canards but landed on the axial solution (where the resource isocline
intersects the Consumer = 0 axis) because landing on the axial solution readily occurs under

red noise. We ran 1000 simulations with 6,000 time units for each value of $1/\varepsilon$, efficiency, and
noise autocorrelation (package DifferentialEquations.jl v6.20.0, Algorithms: Vern7 & Rodas4
with automatic stiffness detection). We used an autoregressive model of order 1 (AR_1) to
create reddened noise with autocorrelation varying from 0 (white noise) to 1 (red noise). We
scaled the variance of the red noise to the variance from the original white noise using the
technique in Wichmann et al. [33] where the ratio of white noise to red noise variances scales
individual noise values in the red noise sequence.

All analyses were done using julia version 1.7.0. [34]

Results

Stability along the slow-fast continuum

When the C-R model is non-excitable (e = 0.5), slowing the consumer's life history increases
the coefficient of variation (Fig. 3). When the C-R is highly excitable (e = 0.71), slowing the
consumer's life history initially decreases then increases the coefficient of variation (Fig. 3).

Figure 3 Average coefficient of variation for time series of Rosenzweig-MacArthur C-R model
with efficiency of either 0.5 (non-excitable) or 0.71 (excitable) along a continuum of $1/\varepsilon$ from 1
to 1000 (50 simulations per value of $1/\varepsilon$ and efficiency). Note, reducing life history speed is
first modestly stabilizing, in effect removing the oscillatory potentials of the underlying
excitable model before life history slowing ultimate generates large increases in CV that likely
produce population disappearances (R~0 during trajectories) due to quasi-canards.

Quasi-cycles

When the C-R model is non-excitable (e = 0.5), quasi-cycles are never found regardless of
how slow the consumer's life history is (Fig. 4 a, c, & e). In contrast, when the C-R model is
highly excitable (e = 0.71) quasi-cycles are found when $1/\varepsilon$ = 1.0 (Fig. 4b & d). With a slowing
of the consumer's life history (now $1/\varepsilon$ = 10) the average autocorrelation function (ACF) line
flattens out indicating reduced manifestation of quasi-cycles (Fig. 4b & f).

Figure 4 a) & b) Average ACF for each lag value for 100 simulations of the C-R model
perturbed each time unit by normally distributed noise (with mean 0.0 and standard deviation
0.01) with $1/\varepsilon$ values of 1 and 10 and with efficiency values of 0.5 (non-excitable) and 0.71
(excitable) respectively. c) & d) Time series of the consumer with $1/\varepsilon$ value of 1 and efficiency
values of 0.5 and 0.71 respectively. e) & f). Time series of the consumer with $1/\varepsilon$ value of 10
and efficiency values of 0.5 and 0.71 respectively.

Quasi-canards

When the C-R model is non-excitable (e = 0.5), quasi-canards can be found under white noise but require a very slow consumer life history (Fig. 5a). When the C-R model is highly excitable (e = 0.71), quasi-canards can be easily found along the gradient of fast to slow consumer life histories (Fig. 5b). Reddening the noise increases the likelihood of finding both quasi-canards and the axial solution regardless of the consumer's life history (Fig. 5 c, d, e, f). Note for a highly excitable C-R interaction, landing on the axial solution is less likely than for a non-excitable C-R interaction because there is more phase space that the dynamics must traverse through. Slowing the consumer's life history down increases the likelihood of the highly excitable C-R interaction landing on the axial solution.

Figure 5 a) & b) Proportion of 1000 simulations under white noise per value of 1/ε that exhibited quasi-canards with constant efficiency values of 0.5 (non-excitable) & 0.71 (excitable) respectively. Vertical dashed lines correspond to log10(1/ε) values used in c), d), e), & f). c), d) Proportion of 1000 simulations per value of log10(1/ε) = 1.1 that exhibited quasi-canards or axial solution or neither with constant efficiency values of 0.5 & 0.71 respectively and with noise correlation (AR1 process) varied from 0.0 to 0.9. e), & f) Proportion of 1000 simulations per value of log10(1/ε) = 2.4 that exhibited quasi-canards or

305 axial solution or neither with constant efficiency values of 0.5 & 0.71 respectively and with
306 noise correlation (AR_1 process) varied from 0.0 to 0.9.

Discussion

Using the simple technique of changing the time scales within the Rosenzweig-MacArthur
consumer-resource (C-R) model, we set up a biologically motivated mathematical experiment
exploring the stability of faster and slower life histories under white to red stochastic noise.
Here, we experimentally manipulated the consumer's intrinsic growth rate while maintaining
the underlying qualitative dynamical conditions. When the energy flux is high, we find that
initially slowing the consumer's life history increased stability through reduced manifestation
of quasi-cycles (alternatively, when energy flux is low, slowing the consumer's life history
always decreased stability). However, further slowing the consumer's life history decreased
stability through increasing the likelihood of quasi-canards and population disappearances.
Furthermore, we found that increasing the autocorrelation of noise tended to increase the
likelihood of quasi-canards and population disappearances. Finally, Yodzis & Innes's [23]
biologically plausible C-R model similarly exhibits quasi-canards (see S.I. Section "Biologically
plausible parameters" & S.I. Figure 5). Consequently, our results are general and suggest that
such instability ought to occur under increasingly reddened perturbations arising from climate
change.

Our results illuminate a gradient in stability along the slow-fast life history continuum that is
dependent on energy flux. If the C-R interaction is highly excitable, initial slowing of the
consumer's life history can increase stability. Stability increases because slowing the
consumer reduces the excitability of the C-R interaction which reduces the manifestation of
the quasi-cycles (see S.I. Section "Slowing the consumer decreases excitability"). If the C-R is
non-excitable or moderately excitable, initial slowing of the consumer's life history decreases
stability (although slowing the consumer's life history does reduce quasi-cycles for the
moderately excitable C-R interaction). Here, perturbations push a slow growth consumer
away from the equilibrium and thus the coefficient of variation continually increases. For all
types of C-R interaction, when the consumer's life history is sufficiently slow, additional
perturbations multiply the effect of each perturbation resulting in a consumer biomass that is
"far from equilibrium" and prone to the distant nonlinear effects of the C-R dynamics (i.e, the
quasi-canard). Overall, there appears to be a gradient in stability along the slow-fast life
history continuum that is mediated by energy flux.

The combination of life history and red noise significantly increased the opportunities for non-
local nonlinear dynamics to be expressed. Reddening the noise in our C-R model increased
the likelihood of quasi-canards (and landing on the axial solution) for both fast and slow
organisms. However, the onset of quasi-canards occurred with less autocorrelation for faster
organisms. To understand this pattern, we must examine the relative time scales of the
autocorrelation and the system population processes [26]. Slowing the life history is
effectively increasing the time scale of the population response processes, and thus the
system will stay for longer in the phase space region the system was pushed into after a
perturbation. Increasing the autocorrelation increases the time scale of the perturbations, and
thus the perturbations effectively are magnified over time.

To unpack the idea of the magnification of the perturbations, we can use similar research on
how multiple discrete disturbances can kick dynamics out of basins of attraction to produce
different dynamical outcomes (flow-kick dynamics: Meyer et al. [32]). In the flow-kick
framework, models are kicked at discrete time points and are allowed to flow (integrate)
without kicks in between the discrete time points. Meyer et al. [32] found that rare but large
disturbances can have the same effect as frequent and small disturbances. Reddened noise
is technically lots of frequent and small disturbances (the kicks in flow-kick dynamics). But
because of the autocorrelation pushing dynamics far from their attracting equilibria, reddened
noise has the same effect as a large disturbance. Furthermore, slowing life histories reduces
the relative time available for organisms to respond to the disturbances (the flow in flow-kick
dynamics), thus pushing dynamics far from their attracting equilibria.

Overall, we can use the analogy of a rusty ratchet to understand how slow life histories and
reddened noise interact. White noise is akin to a ratchet that can spin in any direction (without
a pawl), and red noise is akin to a ratchet with a pawl that can spin in only one direction for a
period of time (Figure 6). Because the reddened noise has a tendency to produce similar
values for a period of time, reddened noise consistently pushes the dynamics of the C-R
model far from the local area around the stable equilibria. Whereas the fast life history is akin
to oil in the ratchet, the slow life history is akin to rust in the ratchet which slows the spinning
speed and in the system increases the time required for trajectories to return to the local area
around the equilibria. To describe this process, we use the term noise-induced ratchet effects
inspired by the rate-induced critical transition literature [35,36]. In a rate-induced critical
transition, canards can be produced when the equilibrium shifts at a slow rate. The dynamics
enter non-local non-linear dynamics because the equilibrium has been pulled from under the
dynamics. For the quasi-canard, the dynamics enter non-local non-linear dynamics because
the dynamics have been pushed by the noise (thus quasi-canards are noise-induced critical
transitions). When slow life histories are combined with red noise, we get noise-induced
ratchet effects.

Figure 6 Illustration of simplified trajectories with white noise (top) and red noise (bottom) together with the consumer and resource isoclines and vector field when the consumer's life history is slow (1/ε is large). We use the analogy of a rusty ratchet to illustrate the interaction of slow life histories with reddened noise. White noise is similar to a ratchet wheel without the pawl (can spin in any direction) and reddened noise is similar to a ratchet with the pawl (can spin in only one direction for a period of time). Slow life history is akin to rust in the ratchet which slows the spinning speed.

Intriguingly, our examination of slow life histories with stochasticity exhibiting sudden population disappearances is a further example of how stochasticity is immensely useful in ecological research to understand the full nonlinear dynamics of ecosystems [20,26,29]. First, stochasticity can act to uncover underlying processes (Boettiger [26] coined the phrase *noise the informer* for this phenomenon). Similar to quasi-cycles where stochastic resonance is visible in advance of a Hopf bifurcation, the quasi-canards also occur in advance of the Hopf bifurcation after which deterministic canards occur. Although noise has the same effect of uncovering an imminent Hopf bifurcation leading to either deterministic cycles or canards, the mechanism producing the quasi-cycles and quasi-canards are different. The quasi-cycles are stochasticity interacting with local nonlinear dynamics (excitability). The quasi-canards are slow life histories plus reddened noise pushing dynamics towards non-local nonlinear dynamics.

Second, stochasticity can be used to reveal differences in stability from what our normal linear stability analysis would find and indeed linear stability analysis would not have predicted the highly destabilizing behaviour for slow consumers. One method to reveal these differences in stability is through using the mathematical tools of the potential function or quasi-potential where potential functions and quasi-potentials can be simplified conceptually to the ball and cup analogy [37]. In this analogy, the state of a system is the position of the ball rolling around a surface (the cup) with minimums being attractors. The potential and quasi-potential are the

cup. The potential function can be found for any system that exhibits solely fixed point
attractors (gradient systems). Quasi-potentials can be found for gradient and non-gradient
systems (the C-R model is a non-gradient system because it exhibits cycling). The quasi-
potentials for our model show the stretching of the quasi-potential along the resource isocline
due to stochasticity with low efficiency and a fast consumer life history, then the quasi-canard
shape when efficiency is large enough, and finally the flattening of the quasi-potential quasi-
canard shape with a slow consumer life history (see S.I. Figure 6). Linear stability analysis
would solely focus on the tiny region around the intersection of the consumer and resource
isoclines. In contrast, the flattened quasi-potentials reveal other possible dynamics (quasi-
canards) in addition to the stable interior equilibria. Overall, stochasticity can both produce
and reveal dynamical outcomes not predicted by normal linear stability analysis.

Conclusion

By varying growth rates along the slow-fast life history continuum within the classic
Rosenzweig-MacArthur model, we have shown that life history has interesting stability
consequences for the system. Slowing the life history can initially increase stability up to a
point in the face of many tiny perturbations, but further slowing can dramatically decrease
stability and increase the potential for highly variable dynamics and population
disappearances. Noise-induced ratchet effects occur when positively correlated noise is
added to an already slowed consumer. This noise-induced ratchet effect is another
mechanism that selects against slow life histories leading to greater proportions of species
with fast life histories. Our study examines a single C-R interaction, which is a fundamental
component of food web theory. Moving forward, the interaction of many more organisms with
varying life histories is required for a more comprehensive understanding of life history and
stability. Furthermore, other examples of non-local nonlinear dynamics should be explored,
especially in the context of slow life histories and reddened noise. Taken together, we have
shown how life history along the slow-fast life history continuum can impact the stability of
systems and shown how human-caused reddened noise will disproportionately impact slow
living organisms through noise-induced ratchet effects and population disappearances.

Acknowledgments

429 We wish to thank members of the McCann laboratory for thoughts and feedback.

Funding

431 CJGG was supported by a Natural Sciences and Engineering Research Council of Canada
432 (NSERC) CGS-D and a Ontario Graduate Scholarship. Financial support was provided by an
433 NSERC Discovery Grant (400353) and a CFREF Food from Thought grant (499075) to K. S.
434 McCann.

Author Contributions

436 All authors contributed to the idea generation and analysis of the model. CJGG wrote the first
437 draft and all authors contributed to editing the manuscript.

Code accessibility

All code to reproduce the above analyses and figures are publicly available on GitHub and have been archived on Zenodo (version 2.0).

References

1. Savage VM, Gillooly JF, Brown JH, West GB, Charnov EL. 2004 Effects of body size and temperature on population growth. *Am. Nat.* **163**, 429–441. (doi:10.1086/381872)

2. Gaillard J -M., Yoccoz NG, Lebreton J -D., Bonenfant C, Devillard S, Loison A, Pontier D, Allaine D. 2005 Generation time: A reliable metric to measure life-history variation among mammalian populations. *Am. Nat.* **166**, 119–123. (doi:10.1086/430330)

3. Réale D, Garant D, Humphries MM, Bergeron P, Careau V, Montiglio P-O. 2010 Personality and the emergence of the pace-of-life syndrome concept at the population level. *Phil. Trans. R. Soc. B* **365**, 4051–4063. (doi:10.1098/rstb.2010.0208)

4. Healy K, Ezard THG, Jones OR, Salguero-Gómez R, Buckley YM. 2019 Animal life history is shaped by the pace of life and the distribution of age-specific mortality and reproduction. *Nat. Ecol. Evol.* **3**, 1217–1224. (doi:10.1038/s41559-019-0938-7)

5. Peters RH. 1984 *The ecological implications of body size.* Cambridge, United Kingdom: Cambridge University Press.

6. Brown JH, Gillooly JF, Allen AP, Savage VM, West GB. 2004 Toward a metabolic theory of ecology. *Ecology* **85**, 1771–1789.

7. Lenton TM, Dakos V, Bathiany S, Scheffer M. 2017 Observed trends in the magnitude and persistence of monthly temperature variability. *Sci. Rep.* **7**, 5940. (doi:10.1038/s41598-017-06382-x)

8. Di Cecco GJ, Gouhier TC. 2018 Increased spatial and temporal autocorrelation of temperature under climate change. *Sci. Rep.* **8**, 14850. (doi:10.1038/s41598-018-33217-0)

9. Audzijonyte A *et al.* 2016 Trends and management implications of human-influenced life-history changes in marine ectotherms. *Fish Fish* **17**, 1005–1028. (doi:10.1111/faf.12156)

10. Couet J, Marjakangas E-L, Santangeli A, Kålås JA, Lindström Å, Lehikoinen A. 2022 Short-lived species move uphill faster under climate change. *Oecologia* **198**, 877–888. (doi:10.1007/s00442-021-05094-4)

11. Stone L. 1993 Period-doubling reversals and chaos in simple ecological models. *Nature* **365**, 617–620. (doi:10.1038/365617a0)

12. Gellner G, McCann KS. 2016 Consistent role of weak and strong interactions in high- and low-diversity trophic food webs. *Nat. Commun.* **7**, 11180. (doi:10.1038/ncomms11180)

13. Gaston KJ, Lawton JH. 1988 Patterns in body size, population dynamics, and regional distribution of bracken herbivores. *Am. Nat.* **132**, 662–680. (doi:10.1086/284881)

14. Rip JMK, McCann KS. 2011 Cross-ecosystem differences in stability and the principle of energy flux: Cross-ecosystem differences in stability. *Ecol. Lett.* **14**, 733–740. (doi:10.1111/j.1461-0248.2011.01636.x)

15. Májeková M, de Bello F, Doležal J, Lepš J. 2014 Plant functional traits as determinants of population stability. *Ecology* **95**, 2369–2374. (doi:10.1890/13-1880.1)

16. Röpke C, Pires THS, Zuanon J, Freitas CEC, Hernandes MC, Souza F, Amadio S. 2021 Growth–reproduction trade-off and fecundity regulate population stability in Amazon floodplain fishes. *Freshwater Biol.* , fwb.13702. (doi:10.1111/fwb.13702)

17. Sæther B-E *et al.* 2013 How life history influences population dynamics in fluctuating environments. *Am. Nat.* **182**, 743–759. (doi:10.1086/673497)

18. Gamelon M, Gimenez O, Baubet E, Coulson T, Tuljapurkar S, Gaillard J-M. 2014 Influence of life-history tactics on transient dynamics: A comparative analysis across mammalian populations. *Am. Nat.* **184**, 673–683. (doi:10.1086/677929)

19. Schwager M, Johst K, Jeltsch F. 2006 Does red noise increase or decrease extinction risk? Single extreme events versus series of unfavorable conditions. *Am. Nat.* **167**, 879–888. (doi:10.1086/503609)

20. Higgins K, Hastings A, Sarvela JN, Botsford LW. 1997 Stochastic dynamics and deterministic skeletons: Population behavior of Dungeness Crab. *Science* **276**, 1431–1435. (doi:10.1126/science.276.5317.1431)

21. Pineda-Krch M, J. Blok H, Dieckmann U, Doebeli M. 2007 A tale of two cycles - distinguishing quasi-cycles and limit cycles in finite predator-prey populations. *Oikos* **116**, 53–64. (doi:10.1111/j.2006.0030-1299.14940.x)

22. Gellner G, McCann KS, Hastings A. 2016 The duality of stability: towards a stochastic theory of species interactions. *Theor. Ecol.* **9**, 477–485. (doi:10.1007/s12080-016-0303-2)

23. Yodzis P, Innes S. 1992 Body size and consumer-resource dynamics. *Am. Nat.* **139**, 1151–1175. (doi:10.2307/2462335)

24. Brose U, Williams RJ, Martinez ND. 2006 Allometric scaling enhances stability in complex food webs. *Ecol. Lett.* **9**, 1228–1236. (doi:10.1111/j.1461-0248.2006.00978.x)

25. Poggiale J-C, Aldebert C, Girardot B, Kooi BW. 2020 Analysis of a predator–prey model with specific time scales: a geometrical approach proving the occurrence of canard solutions. *J. Math. Biol.* **80**, 39–60. (doi:10.1007/s00285-019-01337-4)

26. Boettiger C. 2018 From noise to knowledge: how randomness generates novel phenomena and reveals information. *Ecol. Lett.* **21**, 1255–1267. (doi:10.1111/ele.13085)

27. Coulson T, Rohani P, Pascual M. 2004 Skeletons, noise and population growth: the end of an old debate? *Trends in Ecology & Evolution* **19**, 359–364. (doi:10.1016/j.tree.2004.05.008)

28. McCann KS. 2000 The diversity-stability debate. *Nature* **405**, 228–233. (doi:10.1038/35012234)

29. Hastings A, Abbott KC, Cuddington K, Francis TB, Lai Y-C, Morozov A, Petrovskii S, Zeeman ML. 2021 Effects of stochasticity on the length and behaviour of ecological transients. *J. R. Soc. Interface.* **18**, 20210257. (doi:10.1098/rsif.2021.0257)

30. Touboul J, Krupa M, Desroches M. 2015 Noise-induced canard and mixed-mode oscillations in large-scale stochastic networks. *SIAM J. Appl. Math.* **75**, 2024–2049.

31. Rosenzweig ML. 1971 Paradox of enrichment: destabilization of exploitation ecosystems in ecological time. *Science* **171**, 385–387. (doi:10.2307/1731139)

32. Meyer K, Hoyer-Leitzel A, Iams S, Klasky I, Lee V, Ligtenberg S, Bussmann E, Zeeman ML. 2018 Quantifying resilience to recurrent ecosystem disturbances using flow–kick dynamics. *Nat Sustain* **1**, 671–678. (doi:10.1038/s41893-018-0168-z)

33. Wichmann MC, Johst K, Schwager M, Blasius B, Jeltsch F. 2005 Extinction risk, coloured noise and the scaling of variance. *Theor. Pop. Biol.* **68**, 29–40. (doi:10.1016/j.tpb.2005.03.001)

34. Bezanson J, Edelman A, Karpinski S, Shah VB. 2017 Julia: A fresh approach to numerical computing. *SIAM Rev.* **59**, 65–98. (doi:10.1137/141000671)

35. Siteur K, Eppinga MB, Doelman A, Siero E, Rietkerk M. 2016 Ecosystems off track: rate-induced critical transitions in ecological models. *Oikos* **125**, 1689–1699. (doi:10.1111/oik.03112)

36. Vanselow A, Wieczorek S, Feudel U. 2019 When very slow is too fast - collapse of a predator-prey system. *Journal of Theoretical Biology* **479**, 64–72. (doi:10.1016/j.jtbi.2019.07.008)

37. Nolting BC, Abbott KC. 2016 Balls, cups, and quasi-potentials: quantifying stability in stochastic systems. *Ecology* **97**, 850–564. (doi:10.1890/15-1047.1)

Authors:

Christopher J. Greyson-Gaito[*1], Gabriel Gellner[1], Kevin S. McCann[1]

Affiliations:

1. Department of Integrative Biology, University of Guelph, Guelph, Ontario, Canada

*Corresponding Author – Email: christopher@greyson-gaito.com (CJGG)

ORCID:

CJGG – 0000-0001-8716-0290

GG - 0000-0001-8170-1463

KSM – 000-0001-6031-7913

Scaling of the consumer's growth rate by ε

To see the impact of ε on the consumer's growth rate let's rearrange the parameters in the
model to get the functional response into a monod equation

$$\frac{dR}{dt} = r R \left(1 - \frac{R}{k}\right) - \frac{a R C}{1 + a h R}$$

$$\frac{dC}{dt} = \varepsilon \left(\frac{e a R C}{1 + a h R} - m C\right)$$

becomes

$$\frac{dR}{dt} = r R \left(1 - \frac{R}{k}\right) - \frac{a_{max} R C}{R_0 + R}$$

$$\frac{dC}{dt} = \varepsilon \left(\frac{e a_{max} R C}{R_0 + R} - m C\right)$$

where $a_{max} = \frac{1}{h}$ and $R_0 = \frac{1}{ah}$.

From this version of the model, we can see that ε scales $e \cdot a_{max} - m$ (the consumer's growth rate).

Isoclines of Rosenzweig-MacArthur consumer-resource model

Figure S.I. 1 a), b), c) resource and consumer isoclines with efficiency values of 0.5, 0.6, 0.71
respectively. Dashed lines depicts where the Hopf birfurcation occurs

Moderately excitable consumer-resource interaction

Figure S.I. 2 a) Average coefficient of variation for time series of Rosenzweig-MacArthur C-R resource model with efficiency of 0.6 (moderately excitable) along a continuum of 1/ε from 1 to 1000 (50 simulations per value of 1/ε and efficiency). b) Average ACF for each lag value for 100 simulations of the C-R model perturbed each time unit by normally distributed noise (with mean 0.0 and standard deviation 0.01) with 1/ε values of 1 and 10 and with an efficiency value of 0.6 (moderately excitable). c) Proportion of 1000 simulations under white noise per value of 1/ε that exhibited quasi-canards with an efficiency value of 0.6 (moderately excitable). Vertical dashed lines correspond to 1/ε values used in d), & e). d), e) Proportion of 1000 simulations per value of log10(1/ε) = 1.1 and log10(1/ε) = 2.4 respectively that exhibited quasi-canards or axial solution or neither with an efficiency value of 0.6 and with noise correlation (AR1 process) varied from 0.0 to 0.9.

Slowing the consumer decreases excitability

Below, we numerically and analytically prove that slowing the consumer's life history (increasing $1/\varepsilon$) decreases the excitability of the C-R interaction. By reducing the excitability, we mean that the divide between real and complex eigenvalues (from the linear stability analysis of the interior equilibrium) moves towards the Hopf bifurcation.

Numerical proof

We numerically calculated the real to complex divide for each value of $1/\varepsilon$ between 0.01 and 10 — with a step size of 0.001 in ε (package LinearAlgebra.jl). For each value of $1/\varepsilon$, we found the efficiency value, $e_{R/C}$, where the real/complex divide occurs (i.e. after $1/\varepsilon$ is set, the efficiency value is increased from the efficiency value that produces the transcritical bifurcation until the eigenvalues switch from real to complex, akin to finding the value at which the tip of the checkmark occurs in Gellner and McCann [12]). This efficiency value was then used to calculate the proportion of efficiency "parameter space" that produces real eigenvalues (Proportion Real) by subtracting the real/complex divide efficiency value ($e_{R/C}$) from the efficiency value at the Hopf bifurcation (e_{Hopf}) and then dividing this value by the efficiency parameter distance between the deterministic Hopf (e_{Hopf}) and transcritical ($e_{transcritical}$) bifurcation efficiency values (note ε does not change where the Hopf and transcritical bifurcation occur):

$$Proportion\,Real = \frac{e_{Hopf} - e_{R/C}}{e_{Hopf} - e_{transcritical}}$$

Increases in $1/\varepsilon$ move the real-complex divide towards the Hopf bifurcation and thus, increase the proportion of efficiency "parameter space" that produces real eigenvalues (Figure SI1). In other words, increasing $1/\varepsilon$, reduces the excitability of the system.

503 Figure S.I. 3 Proportion of efficiency "parameter space" that produces completely real
504 eigenvalues for each 1/ε value where the full efficiency "parameter space" corresponds to the
505 distance between the efficiency values that produce the transcritical and Hopf bifurcation.

Analytical Proof

We also proved, using the non-dimensional type I version of the Rosenzweig-MacArthur C-R
model, that increasing 1/ε, increases the efficiency value where the real/complex divide
occurs and thus decreases excitability.

With change of variables

$X = x k, Y = \dfrac{y r}{a}, t = \dfrac{t}{r}$

and with non-dimensional parameters $\alpha = \dfrac{k a e}{r}, \beta = \dfrac{m}{r}$

the non-dimensionalized form of the type I functional response C-R model is:

$\dfrac{d x}{d t} = x(1 - x) - x y$

$\dfrac{dy}{dt} = \epsilon(\alpha x y - \beta y)$

Equilibria exist at

$\widehat{x}_1 = 0, \widehat{y}_1 = 0$

$\widehat{x}_2 = \dfrac{\beta}{\alpha}, \widehat{y}_2 = 1 - \dfrac{\beta}{\alpha}$

The jacobian of the model is

$\begin{pmatrix} 1 - 2x - y & -x \\ \epsilon \alpha y & \epsilon(\alpha x - \beta) \end{pmatrix}$

Inputting the interior equilibrium, $\widehat{x}_2, \widehat{y}_2$, into the jacobian returns

$\begin{pmatrix} \dfrac{-\beta}{\alpha} & \dfrac{-\beta}{\alpha} \\ \epsilon(\alpha - \beta) & 0 \end{pmatrix}$

Using the trace and determinant of this jacobian matrix we can get the characteristic
polynomial and the quadratic equation to solve for the eigenvalues:

$Trace = \dfrac{-\beta}{\alpha}$

$Determinant = \dfrac{\beta \epsilon(\alpha - \beta)}{\alpha}$

$Characteristic\ polynomial = \lambda^2 + \dfrac{\beta}{\alpha} \lambda + \dfrac{\beta \epsilon(\alpha - \beta)}{\alpha}$

$\lambda = \dfrac{\dfrac{-\beta}{\alpha} \pm \sqrt{\left(\dfrac{\beta}{\alpha}\right)^2 - 4 \dfrac{\beta \epsilon(\alpha - \beta)}{\alpha}}}{2}$

We are determining the boundary of real to complex eigenvalues, thus we must examine what
is inside the square root of the quadratic equation:

When $\left(\dfrac{\beta}{\alpha}\right)^2 - 4 \dfrac{\beta \epsilon(\alpha - \beta)}{\alpha} < 0$ the eigenvalues are complex

We can solve for α to find what parameter values produce α at the real/complex divide

$$a = \frac{\beta\epsilon \pm \sqrt{\beta\epsilon(\beta\epsilon+1)}}{2\epsilon}$$

We can ignore the minus square root part (because $\beta\epsilon < \sqrt{\beta\epsilon(\beta\epsilon+1)}$ always and we get a
negative alpha value which is impossible biologically).

Thus, we concentrate on

$$\alpha = \frac{\beta\epsilon + \sqrt{\beta\epsilon(\beta\epsilon+1)}}{2\epsilon}$$

We can differentiate the above equation with respect to ε to find out how the α value (at which
the real/complex divide occurs) changes.

$$\frac{d\alpha}{d\epsilon} = \frac{-\beta}{4\epsilon\sqrt{\beta\epsilon(\beta\epsilon+1)}}$$

which is always negative when β and ε are positive (biologically they have to be).

Therefore, if we decrease ε (slowing the consumer by increasing $1/\varepsilon$), the α value — at which
the real/complex divide occurs — increases. Converting α back into its original dimensional
parameters, we see that if k and a are kept constant, e must increase to increase the non-
dimensional α parameter.

Explanation of quasi-canard finder algorithm

The algorithm checks that the trajectory has the characteristics of a quasi-canard. Thus, the
algorithm includes a return map at the maximum point of the resource isocline where canards
and quasi-canards must pass through. The algorithm also includes boxes along the attracting
and repelling manifolds (the right side of the resource isocline and the consumer axis
respectively) through which a quasi-canard should pass. The quasi-canard passes through
these checks in a particular order and so the algorithm ensures the order is correct. Below are
the six steps that the quasi-canard finder algorithm goes through. The full code can be found
in slowfast_canardfinder.jl of the Github repository.

1. The algorithm finds all the points in the time series where the next sequential point
creates a vector that intersects with a line that sits at the Hopf bifurcation point on the
Resource isocline (the maximum of the Resource isocline). The line has a length of 5%
of the Hopf bifurcation point above and below the Hopf bifurcation point. If no points
are found, the algorithm does step six. If points are found, the points are collated and
passed to the next step.

2. The algorithm then takes all of these points and moves along the time series after
these points to identify the first point within a box that sits between the Hopf bifurcation

563	point and where the Resource isocline intersects with the Consumer axis. The box has
564	a width of 0.1. If no points are found, the algorithm does step six. If points are found,
565	the points are collated and passed to the next step.

3. The algorithm then takes all of these points and moves along the time series after
these points to identify the first point within a box that sits between the 0 consumers
and 80% of where the Resource isocline intersects with the Consumer axis. The box
has a width of 0.1. If no points are found, the algorithm does step six. If points are
found, the points are collated and passed to the next step.

4. The algorithm then takes all of these points and moves along the time series after
these points to identify the first point that sits close to the resource isocline. If no points
are found, the algorithm does step six. If points are found, the points are collated and
passed to the next step.

5. The algorithm then takes all of these points and repeats step 1 to ensure a full cycle of
the quasi-canard. If the return map check is passed for the second time, the algorithm
returns "quasi-canard", otherwise the algorithm does step six.

6. The algorithm checks whether the final point in the time series is 0.0 consumers and
3.0 resources (where the axial solution exists). If so, the algorithm returns "axial",
otherwise the algorithm returns "nothing".

Note, the sensitivity of this algorithm to find quasi-canards can be changed by varying the top
of the box in step three (changing the percentage of where the Resource isocline intersects
with the Consumer axis).

Figure S.I. 4 C-R phase plot with a quasi-canard, the resource isocline and the consumer
isocline in purple, blue, and orange respectively. The six steps of the algorithm outlined above
are depicted with green lines, boxes, and circles.

Biologically Plausible Parameters

We used Yodzis & Innes' [23] biologically plausible parameterization of the C-R model to test
whether our sudden population disappearance results are general to other parameter sets.

$$\frac{dR}{dt} = R\left(1 - \frac{R}{K}\right) - \frac{\frac{x\,y}{(1-\delta)f_e}C\,R}{R+R_0}$$

$$\frac{dC}{dt} = C\,x\left(-1 + \frac{y\,R}{R+R_0}\right)$$

where $x = \left(\frac{a_T}{f_r a_r}\right)\left(\frac{m_R}{m_C}\right)^{0.25}$

$y = \frac{f_J a_J}{a_T}$

Similar to Yodzis & Innes [23], we expressed the resource body mass in terms of the mass of
an equivalent endotherm operating at its physiological limit. We set our consumer as an
herbivorous endotherm.

Thus, $a_T=54.9, a_r=34.3, \delta=0.55, f_J=0.99, a_J=89.2, K=3.0$

To slow the consumer relative to the resource, we multiplied the resource/consumer body
mass ratio by ε (ε B = ε m_{ER}/m_C). To maintain the same biomass loss from the resource (i.e.
when using the original B = m_{ER}/m_C), we set

$$f_e=\frac{\left(\varepsilon B\right)^{0.25}}{B^{0.25}}$$

To ensure the full model was feasible (between the feasibility and Hopf boundaries in Yodzis
& Innes [23]), we set B = 10^{-6} and we restricted R_0 to [0.7,1.82].

All other methods are the same as in the main Methods section.

Figure S.I. 5 In the Yodzis & Innes (1992) model, quasi-canards can be found for a variety of
efficiency values within certain boundaries of ε parameter space. a), b), c), d) resource and
consumer isoclines with R_0 values of 1.2, 1.0, 0.8 respectively. e), f), g), h) proportion of 1000

simulations per value of $1/\varepsilon$ that exhibited quasi-canards with constant R_0 values of 1.2, 1.0,
0.8 respectively. Bold and thin dashed lines correspond to 6,000 and 24,000 time units
respectively.

Quasi-potentials

The quasi-potentials depicted below were created using the Rosenzweig-MacArthur
consumer -resource model with the same parameter values as the model in the main article.
We used the QPot package (version 1.2) in R to calculate the quasi-potentials (Moore *et al.*
2016). We maintained the overall intensity of noise but had different relative noise intensities
between the resource and the consumer (specifically 1:4, see Moore *et al.* (2016) for how to
specify different relative noise intensities).

Figure S.I. 6 Quasi-potentials for the C-R models. The top row (a) & b)) had a $1/\varepsilon$ value of
12.7 and the bottom row (c) & d)) had a $1/\varepsilon$ value of 250. Each column of plots had efficiency
values of 0.5 and 0.7 respectively. Resource and consumer isoclines are the blue and orange
lines respectively.

Christopher Moore, Christopher Stieha, Ben Nolting, Maria Cameron and Karen Abbott
(2016). QPot: Quasi-Potential Analysis for Stochastic Differential Equations. R package
version 1.2. https://github.com/bmarkslash7/QPot.