Title: STK25 inhibits PKA signaling by phosphorylating PRKAR1A

Authors: Xiaokan Zhang[1], Bryan Z. Wang[1,2], Michael Kim[1], Trevor R. Nash[1,2], Bohao Liu[1,2], Jenny

Rao[1], Roberta Lock[2], Manuel Tamargo[2], Rajesh Kumar Soni[3], John Belov[1], Eric Li[1], Gordana

Vunjak-Novakovic[1,2] and Barry Fine[1*]

Affiliations

[1] Department of Medicine, Division of Cardiology, Columbia University Irving Medical Center, New

York, NY, USA

[2] Department of Biomedical Engineering, Columbia University, New York, NY, USA

[3] Proteomics and Macromolecular Crystallography Shared Resource, Herbert Irving

Comprehensive Cancer Center, Columbia University Irving Medical Center, New York, NY, USA

***Corresponding Author**

Barry Fine MD PhD

Department of Medicine, Division of Cardiology

622 West 168th Street

PH8-405B

New York, NY 10032

Preprint Server: bioRxiv, 2022.01.27.478084
21
22
Keywords: STK25, PRKAR1A, Protein Kinase A, induced pluripotent stem cell

Summary

In the heart, Protein Kinase A (PKA) is critical for activating calcium handling and sarcomeric

proteins in response to beta adrenergic stimulation leading to increased myocardial contractility

and performance. The catalytic activity of PKA is tightly regulated by regulatory subunits which

inhibit the catalytic subunit until released by cAMP binding. Phosphorylation of Type II regulatory

subunits promotes PKA activation, however the role of phosphorylation in Type I regulatory

subunits remain uncertain. Here we utilized human induced pluripotent stem cell cardiomyocytes

(iPSC-CM) to identify STK25 as a kinase of the Type Iα regulatory subunit PRKAR1A.

Phosphorylation of PRKAR1A led to inhibition of PKA kinase activity and increased binding to the

catalytic subunit in the presence of cAMP. *Stk25* knockout in mice diminished Prkar1a

phosphorylation, increased Pka activity and augmented contractile response to beta adrenergic

stimulation. Together, these data support STK25 as a negative regulator of PKA signaling

through phosphorylation of PRKAR1A.

Introduction

Second messengers are key relays in the transduction of external cues into intracellular

signaling cascades. cAMP is a prototypical second messenger generated downstream from G

protein coupled receptors and is involved in integrating a diverse set of signaling pathways

(Gancedo, 2013). The classic and most widely studied effector of cAMP is PKA and it has been

broadly implicated in a diverse set of cellular processes including the cell cycle, proliferation,

cytoskeletal dynamics, ion flux and beta adrenergic signaling (Tasken and Aandahl, 2004; Taylor

et al., 2012). Specificity of signaling in a system with such a broad set of substrates relies on

careful regulation of both cAMP metabolism as well as compartmentalization of PKA through a

series of A kinase anchoring proteins (AKAP) (Colledge and Scott, 1999). These AKAPs bind to

the regulatory subunits of PKA and direct its activity to a discrete set of effectors and substrates,

thereby allowing for enhanced spatiotemporal control over PKA and the pathways it activates.

The PKA holoenzyme is comprised of a regulatory subunit dimer and two catalytic units

(Kim et al., 2007). The main function of the regulatory subunit is to inactivate kinase activity in

the absence of cAMP. There are two classes of regulatory subunits, Type I and Type II, each

with an alpha and beta isoform. Binding of cAMP to the C terminal tandem cAMP binding domains

of the regulatory subunits releases the catalytic domain, allowing for kinase activity (Barradeau et

al., 2002). Type II regulatory subunits have a PKA substrate motif in the N -terminus (RRXS),

which is phosphorylated by the catalytic subunit upon binding to cAMP, enhancing its release and

activation (Taylor et al., 1990). They also are the primary target of AKAPs, binding at nanomolar

affinities (Colledge and Scott, 1999). The regulation of Type I subunits however remains less well

understood.   They   contain   a   pseudo-substrate   (RRXxA/G)   sequence   that   cannot   be

phosphorylated by the catalytic subunit. Although phosphorylation of the Type I subunits has

been reported as part of large phosphoproteomic projects, their role in regulation of PKA activity

has not been demonstrated *in vivo* (Daub et al., 2008; Olsen et al., 2006; Olsen et al., 2010;

Oppermann et al., 2009; Zhou et al., 2013). Furthermore, the vast majority AKAP proteins do not

interact with Type I subunits, and the dual specificity AKAP proteins that do interact with Type I subunits do so with significantly lower affinities compared to Type II subunits.

In the heart, PKA mediates beta adrenergic activation of the physiologic flight or fight response, modulating both heart rate and myocardial contractility (Wang et al., 2018). Substrates of PKA in the cardiomyocyte include troponin (TNNT2), myosin binding protein c (MYBPC3), phospholamban (PLN), and ryanodine receptor (RYR2) (Marks, 2013). Acute activation of PKA in hyperadrenergic states leads to improved calcium flux and myocardial performance and is necessary for increases in stroke volume to meet cardiac output demands. In heart failure, chronic beta-adrenergic stimulation from high levels of catecholamines results in compensatory downregulation of beta adrenergic signaling to leading diminished phosphorylation of several PKA substrates (Bristow et al., 1982; Piacentino et al., 2003). Long term activation of Pka was shown to be deleterious in mouse models and is thought to underly the maladaptive remodeling of chronic beta-adrenergic activation in heart failure (Antos et al., 2001).

STK25 is a stress response kinase in the sterile-20 (Ste20) kinase superfamily and has been shown through a series of *in vivo* models to impact glucose utilization, insulin homeostasis, and lipid metabolism (Amrutkar et al., 2015a; Amrutkar et al., 2015b; Amrutkar et al., 2016a; Amrutkar et al., 2016b; Cansby et al., 2013; Chursa et al., 2017; Nerstedt et al., 2012; Nunez-Duran et al., 2017; Sutt et al., 2018). Overexpression of STK25 leads to steatosis and was shown to aggravate atherosclerosis in a PCSK9 gain of function mouse model (Cansby et al., 2018). Recently, STK25 was shown to interact with the Hippo signaling via phosphorylation of LATS1/2 and/or regulation of SAV1-STRIPAK (Bae et al., 2020; Lim et al., 2019). In this study, we use a combination of *in vitro* and *in vivo* cardiomyocytes models to demonstrate that STK25 phosphorylates the Type Iα regulatory subunit of PKA, PRKAR1A. This leads to increased inhibition of PKA signaling, revealing a new regulatory mechanism that can control PKA activity and attenuate cAMP mediated increases in cardiac contractile function.

Methods:

Patient samples

Patients with advanced HF were recruited at Columbia University Medical Center and heart tissue at the time of heart transplant was collected from the left ventricle. Control myocardial samples were obtained from the National Disease Research Interchange (https://ndriresource.org/ats) and were comprised of de-identified specimens collected from non-failing hearts determined to be unusable for cardiac transplantation due to non-cardiac donor issues but without evidence or knowledge of underlying cardiac disease. The study was approved by the Institutional Review Board of Columbia University (IRB# AAAR0055). All patients provided written informed consent before inclusion into the study.

Cell culture

HEK293T cells (ATCC Cat# CRL-3216, RRID:CVCL_0063) were cultured in high glucose (4.5g/L) DMEM supplemented with 10% FBS, penicillin and streptomycin, and grown in a CO_2 incubator maintained at atmospheric oxygen levels and 5% CO_2. Human induced pluripotent stem cells (hiPSC) were obtained through Material Transfer Agreements from Bruce Conklin, Gladstone Institute (WTC cell line). hiPSCs were expanded on growth factor reduced Matrigel-coated plates (Corning) in mTeSR plus medium (Stemcell technologies) containing mTeSR plus supplement (Stemcell technologies), 50U penicillin and 50U streptomycin. The cell culture medium was changed every other day, and cell passaged upon reaching 70% confluence. During the first 24 hrs after passaging, 5 µM Y-27632 dihydrochloride (Tocris, 1254) was supplemented to culture medium.

CRISPR-Cas9

CRISPR-Cas9 Genome editing was used to generate a homozygous knockout of *STK25* in WTC iPSCs following the manufacturer's protocol (ORIGENE) with gRNA vector (KN203215G).

Single-cell clones were expanded and confirmation of homozygous editing was determined by

sanger sequencing of gDNA and protein analysis by western blot.

<u>iPS Cardiomyocyte Differentiation</u>

Cardiac differentiation of iPSC's was initiated in confluent monolayers two days after replating at

a density of $200,000/cm^2$ (24hrs in mTeSR plus medium with 5µM Y-27632 dihydrochloride,

followed by 24hrs in mTeSR plus medium without Y-27632). On day 0 of differentiation, media

was changed to cardiac differentiation medium (CDM, consisting of RPMI-1640, Life

Technologies), 500µg/ml human recombinant albumin (Sigma), 213µg/ml L-ascorbic acid

(Sigma), 50U penicillin and 50U streptomycin) with 3-6 µM CHIR (4423, Tocris). 2 days after

initial CHIR addition, media was changed to CDM with 2 µM Wnt-C59 (5148, Tocris). From day

4 onwards, media was changed fresh CDM every other day until cells start contracting. Media

was then switched to RPMI-B27 (consisting of RPMI-1640, 1x B27 supplement (Life

Technologies), 213µg/mL L-ascorbic acid, 50U penicillin and 50U streptomycin). Cardiomyocytes

were characterized by flow cytometry using the cardiomyocyte-specific marker TNNT2 (BD

Biosciences Cat# 565744, RRID:AB_2739341). Differentiation typically resulted in cell

populations containing 80–90% TNNT2-positive cells at day 12.

<u>PKA activity assay</u>

PKA activity assay was performed using whole cell lysate from iPSC-CMs treated with either PBS

or 10µM forskolin (Sigma-Aldrich, F6886) for 30min at 37°C prior to assay, or immunoprecipitated

type I and II PKA holoenzymes from iPSC-CMs whole cell lysate or whole heart lysates from

Stk25[+/+] or *Stk25*[-/-] mice following the manufacturer's protocol (EIAPKA, Invitrogen). Briefly, PKA

standards or diluted samples and reconstituted ATP were added into wells of PKA substrate plate.

Phospho-PKA substrate antibody and Goat anti-Rabbit IgG HRP conjugate antibody were added

into the wells per manufacturer protocol. After incubation TMB substrate was added into each

well, and the plate was incubated for 30 min at room temperature. Stop solution was added, and

absorbance at 450nm was analyzed in a 96-plate reader.

<u>Global Phosphoproteomics analysis</u>

Sorted $STK25^{+/+}$ and $STK25^{-/-}$ iPSC-CMs were lysed/homogenized by bead-beating in 8 M urea,

1%SDS and 200 mM EPPS (pH 8.5), protease, and phosphatase inhibitors. Lysates were

cleared by centrifugation at 21,000 g for 30 min at 4^0C, and protein concentration was

measured by BCA. Proteins were reduced with 5 mM TCEP, alkylated with 10 mM

iodoacetamide (IAA), and quenched with 10 mM DTT. A total of 500 µg of protein was

chloroform−methanol precipitated. Protein was reconstituted in 200 mM EPPS (pH 8.5) and

digested by Lys-C overnight and trypsin for 6 h, both at a 1:50 protease-to-peptide ratio.

Digested peptides were quantified using a Nanodrop at 280 nm, and 200 µg of peptide from

each sample were labeled with 800 µg TMT reagent using a 10-plex TMT kit (Navarrete-Perea

et al., 2018). TMT labels were checked, 100 ng of each sample was pooled and desalted and

analyzed by short SPS-MS3 method, and using normalization factor samples were bulk mixed

at 1:1 across all channels and desalted using a 200 mg Sep-Pak solid-phase extraction column

and dried using vacuum centrifugation.

Desalted isobaric labeled peptides were enriched for phospho-peptides using a mixture

of MagReSyn Ti-IMAC and Zr-IMAC resins according ReSyn Bioscience instructions. In brief,

1.4 mg of labeled peptide were resuspended in 1 ml of binding buffer (80% Acetonitrile, 1M

glycolic acid and 5% TFA) and incubated with equilibrated 150 µl (75 µl of each Ti-IMAC and Zr-

IMAC) resins at room temperature for 30 min, and the resin was washed 3 three times to

remove the unbound, non-phosphorylated peptides. Phospho-peptides were eluted using 1%

ammonium hydroxide. The enriched phospho-peptides were further fractionated in eight

fractions using Pierce™ High pH Reversed-Phase Peptide Fractionation Kit and each fraction

dried down in a speed-vac.

The isobaric labeled dried phospho-peptides were resuspended in 10 µl of (3%

acetonitrile/ 0.1% formic acid), and analyzed on an Orbitrap Fusion mass spectrometer coupled

to a Dionex Ultimate 3000 (ThermoFisher Scientific) using the MSA-SPS-MS3 and NL SPS-

MS3 method (Jiang et al., 2017). Peptides were separated on an EASY-Spray C18 50cm

column (Thermo Scientific). Peptide elution and separation were achieved at a non-linear flow

rate of 250 nl/min using a gradient of 5-30% of buffer B (0.1% (v/v) formic acid, 100%

acetonitrile) for 110 minutes with a temperature of the column maintained at 50 °C during the

entire experiment. For both methods, MS1 data were collected using the Orbitrap (120,000

resolution; maximum injection time 50 ms; AGC 4×10^5). Determined charge states between 2

and 5 were required for sequencing and a 45 s dynamic exclusion window was used. Data-

dependent top10 MS2 scans were performed in the ion trap with collision-induced dissociation

(CID) fragmentation (Turbo; NCE 35%; maximum injection time 60ms; AGC 5×10^4). MS3

quantification scans were performed using the multi-notch MS3-based TMT method (ten SPS

ions; 50,000 resolution; NCE 65% for MSA-SPS-TMT and 38% for NL-SPS-TMT maximum

injection time 105 ms; AGC $1x10^5$) using the Orbitrap.

Raw mass spectrometric data were analyzed using Proteome Discoverer 2.2 to perform

database search and TMT reporter ions quantification. TMT tags on lysine residues and

peptide N termini (+229.163 Da) and the carbamidomethylation of cysteine residues (+57.021

Da) was set as static modifications, while the oxidation of methionine residues (+15.995 Da),

deamidation (+0.984) on asparagine and glutamine and phosphorylation (+79.966) on serine,

threonine, and tyrosine were set as a variable modification. Data were searched against a

UniProt Human database with peptide-spectrum match (PSMs) and protein-level FDR at 1%

FDR. The signal-to-noise (S/N) measurements of each protein normalized so that the sum of

the signal for all proteins in each channel was equivalent to account for equal protein loading.

Phospho-peptides identification and quantification were imported into Perseus (Tyanova et al.,

2016) for t-test statistical analysis (FDR<0.05) to identify phospho-peptides demonstrating

statistically significant changes in abundance. Pathway analysis were performed using

Ingenuity Pathway Analysis (Qiagen).

<u>Animal studies</u>

Stk25 knockout mice were generated by CRISPR/Cas9 mediated genome engineering in

C57BL/6J background (Jackson Laboratory, Bar Harbor, Maine). Exon 3-5 of Stk25 gene were

deleted with two CRISPR guides from Synthego (sgRNA-stk25-7367 and sgRNA-stk25-10150).

Both guides were mixed with IDT Cas9V3 protein to form RNP, which was further injected into

the pronuclei of fertilized C57BL/6J eggs to generate knockout founders. Genotyping was

performed to show the *Stk25* knockout allele. Further breeding generated homozygous *Stk25$^{-/-}$*

mice and protein immunoblotting was performed to confirm an absence of Stk25. The protocol for

all mouse experiments (AABC1503) was approved by the Columbia University Institutional Animal

Care and Use Committee.

Cardiac function of 20 week and 52 week old mice was assessed by echocardiography

by the Columbia University Mouse Imaging Core Facility (imaged by Visulasonics VEVO 3100

High Frequency Ultrasound imaging system and analyzed by Vevo LAB software). For 20 week

old mice, cardiac function was recorded at baseline and 3 minutes after administration of the β-

adrenergic receptor agonist isoproterenol (0.2ug/g, i.p.). Systolic function parameters including

ejection fraction (EF, %), fractional shortening (FS, %), left ventricular end systolic diameter

(LVESD, mm) and left ventricular end diastolic diameter (LVEDD, mm) were measured in the two-

dimensional parasternal short-axis imaging plane of M-mode tracings close to the papillary

muscle level.

<u>iPS Cardiomyocyte sorting</u>

Cardiomyocytes were pretreated with 10μM of Y-27632 in B27 culture medium for 6 hrs, then

digested with cell dissociation buffer (containing Hank's buffered saline solution, 100 Units/ml

collagenase, 10μM Y-27632) at room temperature overnight. Cells were collected and

centrifuged at 300g for 5 min, and then resuspended in cell sorting buffer (CSB, containing

Hank's buffered saline solution, 20mM HEPES, 5% FBS, 10 units/ml Turbo DNase). The cell

pellet was resuspended with anti-CD172a/b (SIRPα/β, 423107, Biolegend) and anti-CD90 (Thy-

1, 11-0909-42, Invitrogen) antibodies in CSB to stain for 20 min at 4°C in the dark. Cells were

washed 3 times with CSB and DAPI staining was added. Cardiomyocytes were sorted from the

SIRPA+ and CD90- population and then plated on matrigel coated plates in B27 medium with

5μM Y-27632. The following day, medium was replaced with fresh B27, and cardiomyocytes

generally started to beat in 2-5 days.

RNA Isolation, Sequencing, and Analysis.

RNA was extracted using the RNeasy Mini kit (Qiagen #74004) along with on-column DNAse

digestion (Qiagen #79254). cDNA libraries were generated using the Clontech Ultra Low v4 kit

followed by NextaraXT DNA Library Prep Kit, then sequenced on an Illumina NovaSeq 6000.

Paired-end 100-bp sequenced reads were analyzed as follows: RTA (Illumina) software was

used for base calling and bcl2fastq2 (version 2.19) for converting BCL to fastq format, coupled

with adaptor trimming. A pseudoalignment to a kallisto index created from transcriptomes (build

GRCh38) using kallisto (0.44.0). After pseudoalignment the R package DESeq2 (1.28.1) was

used to normalize the count matrix and calculate differentially expressed genes. Geneset

enrichment analysis was performed using GSEA software (4.10.0) per documentation (Mootha

et al., 2003; Subramanian et al., 2005).

siRNA and vector transfection

Cells were transfected using lipofectamine 3000 (Invitrogen) following the manufacturer's

protocol with siRNA targeting STK25 (On-TargetPlus siRNA, Horizon Discovery, L-004873-00-

0050) or siRNA targeting PRKAR1A (EHU071341, Sigma) for protein knockdown or non-

targeting control siRNA (ON-TARGETplus non-targeting pool, Horizon Discovery D-001810-10-

50) as scramble control.

For heterologous expression studies, cells were transfected using lipofectamine 3000

(Invitrogen, L3000015) following the manufacturer's protocol. Vectors used include Flag-WT-

STK25 vector (EX-M0142-M46, GeneCopoeia), Flag-K49R/T147A-STK25 vector (Vector

Builder), Flag- empty vector (EX-NEG-M46, GeneCopoeia), V5-PRKAR1A (Vector Builder,

VB200124-1141aes), V5-S77A/S83A-PRKAR1A (Vector Builder, VB200124-1126jtp), and V5-

S77E/S83E-PRKAR1A (Vector Builder, VB200124-1127pst).

qRT-PCR assays

Equivalent amounts (2µg) of purified RNA were used as a template to synthesize cDNA using

oligo-d(T) primers and SuperScript™ III First-Strand Synthesis SuperMix (Invitrogen,

18080400). *STK25* was quantified by real-time PCR using Fast SYBR Green mixture (Life

Technologies, 4385612) and was carried out on Applied Biosystems Step One Plus. Relative

levels were calculated using $\Delta\Delta C_T$ method. Data analysis was carried out using the fold change

normalized to GAPDH gene expression.

Gene	Forward Primer 5'-3'	Reverse Primer 5'-3'
STK25	GCTCCTACCTAAAGAGCACCA	TGGCAATGTATGTCTCCTCCAG
GAPDH	GGACTCATGACCACAGTCCATG	CAGGGATGATGTTCTGGAGAGC

Immunoprecipitation and immunoblot analysis

300 µg of total protein from whole cell lysate were used in immunoprecipitation experiments.

The extract was incubated with anti-Flag affinity gel (Sigma-Aldrich Cat# A2220,

RRID:AB_10063035), anti-V5 affinity gel (Sigma-Aldrich Cat# A7345, RRID:AB_10062721),

anti-PRKAR1A (Abcam Cat# ab139695, RRID:AB_2893184) or anti-PRKAR2A (ProteinTech

Cat# 10142-2-AP) overnight at 4°C. The affinity gel beads were centrifuged at 8000xg for 1min,

washed three times in TBS buffer. The affinity gel beads were added with 4x Laemmli SDS

sample buffer (NuPAGE), denatured at for 5min at 95°C and analyzed by SDS-PAGE and

immunoblotting. For Western blotting antibodies include: HRP conjugated anti-GAPDH (Cell

Signaling Technology Cat# 3683, RRID:AB_1642205), anti-STK25 (Abcam Cat# ab157188,

RRID:AB_2725788), anti-Flag (Sigma-Aldrich Cat# F3165, RRID:AB_259529), anti-GM130

(Cell Signaling Technology Cat# 12480, RRID:AB_2797933), anti- PRKAR1A (Abcam Cat#

ab 139695, RRID:AB_2893184), anti-pS77 PRKAR1A (Abcam Cat#ab139682,

RRID:AB_2904566), anti-pS83 PRKAR1A (Abcam Cat#ab154851, RRID:AB_2904567), anti-

PRKAR2A (ProteinTech Cat# 10142-2-AP), anti-phospholamban (Cell Signaling Technology

Cat# 14562, RRID:AB_2798511), anti-phospho-phospholamban -pS16/T17 (Cell Signaling

Technology Cat# 8496, RRID:AB_10949102), anti-Ryanodine receptor 2 (Abcam

Cat#ab196355, RRID:AB_2904568), anti-p- Ryanodine receptor 2-pS2808 (Abcam Cat#

ab59225, RRID:AB_946327), anti-TnI (Cell Signaling Cat# 4002), anti-TnI-pS23/S24 (Cell

Signaling Cat# 4004), anti-V5 (Sigma-Aldrich Cat# V8137, RRID:AB_261889), anti-PKA

Catalytic subunit (Abcam Cat# ab26322, RRID:AB_2170049), HRP-conjugated anti-mouse (Cell

Signaling Technology Cat# 7076, RRID:AB_330924) and HRP-conjugated anti-rabbit (Cell

Signaling Technology Cat# 7074, RRID:AB_2099233) were used for detection.

In vitro kinase assay

Recombinant protein STK25 (0.5µM, TP303215, OriGene) and PRKAR1A (1µM, TP303828

Origene) were mixed with *in vitro* kinase buffer (25mM Tris-HCl, 10mM beta-glycerophosphate,

10mM MgCl$_2$, 20mM NaF, 2mM DTT, 1mM sodium orthovanadate, 1x Protease Inhibitor cocktail

(Roche)) containing ATP (500uM) and incubated at 37°C for 30 min. In vitro kinase assay was

terminated by adding Laemmli SDS sample buffer. Phosphorylation levels of PRKAR1A at S77

and S83 were analyzed by SDS-PAGE and immunoblotting.

Real Time Glo cell viability assay

Cells were seeded into 96-well plates at a density of 3000 cells per well 24h before

measurement. RealTime-Glo (Promega; G9712) reagent was added to each well per

manufacturer protocol and measurement of luminescence was performed at selected time

points on a BioTek Synergy JHTX with Gen5 data analysis software.

Statistical Analysis:

Statistical analyses were performed using Prism 8 (Graphpad Software). Results are presented

as mean ± standard deviation. For comparisons between two groups, a two tailed unpaired t-

test was used unless otherwise specific. Welch's correction was utilized for two groups of

unequal sizes. For multiple group comparisons, either one way or two-way (depending on the

number of variables) ANOVA followed by multiple comparison post-hoc testing was performed

as indicated using Prism 8. Notation in the text is as follows *$p<0.05$, **$p<0.01$, ***$p<0.001$ and

****$p<0.0001$

16
Results:

STK25 inhibits PKA signaling.

As part of an kinome knockout discovery project, CRISPR-Cas9 was used to generate a

homozygous knockout of *STK25* in the wild type iPSC line WTC11 (Bruce Conklin, Gladstone

Institute) (Hayashi et al., 2016), and cardiomyocytes were differentiated and characterized using

an established protocol (Burridge et al., 2014). *STK25* expression in iPSC derived

cardiomyocytes was found to be similar to *Stk25* expression to primary cardiomyocytes from adult

mice (**Figure S1A and S1B**). To identify potential substrates of STK25 in cardiomyocytes, mass

spectrometry based phosphoproteomics was performed with *STK25$^{+/+}$* and *STK25$^{-/-}$*

cardiomyocytes (**Figure 1A**). Signaling pathways impacted by *STK25* were identified using

Ingenuity Pathway Analysis (Qiagen). A decrease in Hippo signaling was observed in response

to loss of STK25 confirming a prior study demonstrating STK25 activates the Hippo pathway

(**Figure 1B**) (Lim et al., 2019). The most significant signaling change in response to loss of *STK25*

was an upregulation of the PKA pathway. Several downstream substrates of PKA displayed

increased phosphorylation including MYBPC3, TNNT2, RYR2, CACNA1C and GSK3α (**Table 1**).

STK25$^{-/-}$ cardiomyocytes exhibited increased PKA kinase activity when stimulated with the

adenylate cyclase agonist forskolin (**Figure 1C**). In heterologous overexpression studies in

HEK293T cells, STK25 decreased forskolin induced PKA activity. This inhibition was dependent

on the kinase domain of STK25 as overexpression of the kinase-dead mutant *STK25$^{K49R/T174A}$*

(Preisinger et al., 2004) did not inhibit PKA activity in response to forskolin (**Figure 1D)**.

STK25 phosphorylates PRKAR1A

Two of the most downregulated phosphorylation sites in *STK25$^{-/-}$* iPSC-CM were S77 and

S83 of PRKAR1A raising the possibility that these sites were directly phosphorylated by STK25

(**Table 1**). PRKAR1A is the R1α member of the regulatory subunit isoforms of the PKA

holoenzyme which binds to and inhibits the catalytic subunit of PKA (PRKACA) and has been

shown to disengage in response to cAMP, leading to PKA activation (Bossis and Stratakis, 2004).

We first confirmed the phosphoproteomic data by demonstrating decreased phosphorylation of

S77 and S83 in *STK25$^{-/-}$* cardiomyocytes by immunoblot protein analysis (**Figure 2A and S2A**).

In order to determine if this observation was dependent on the kinase activity of STK25,

STK25$^{K49R/T174A}$ was expressed in *STK25$^{-/-}$* iPSC-CMs and led to significantly decreased

phosphorylation of S77 and S83 compared to overexpression of wild type STK25 (**Figure 2B and**

S2B). Stimulation of PKA with forskolin led to diminished phosphorylation at both S77 and S83

in *STK25$^{+/+}$* iPSC-CM but did not have significant effect on *STK25$^{-/-}$* iPSC-CMs (**Figure 2C and**

S2C).

Co-immunoprecipitations experiments demonstrated STK25 and PRKAR1A binding using

epitope tagged protein in HEK293T cells (**Figure 2D**). Though forskolin inhibited binding between

PRKAR1A and the catalytic subunit, PRKACA, it did not alter the association between STK25 and

PRKAR1A (**Figure 2E**). Mutation of the kinase domain of STK25 also did not affect binding to

PRKAR1A by immunoprecipitation (**Figure S2D**). Using purified protein in an *in vitro* kinase

assay, STK25 was able to phosphorylate PRKAR1A at S77 and S83(**Figure 2F**).

Phosphorylation of S77/S83 inhibits PKA activity

Phosphorylation at S77 and S83 on PRKAR1A have been described as part of large scale global

phosphoproteomic experiments (Bian et al., 2014; Daub et al., 2008; Olsen et al., 2006;

Oppermann et al., 2009) but their influence on PKA activity has not been characterized. In order

to investigate the effect of pS77 and pS83 on the ability of PRKAR1A to inhibit PKA activity,

phosphomimetic (S77E/S83E) and non-phosphomimetic (S77A/S83A) mutations were generated

in PRKAR1A expression vectors. In HEK293T co-immunoprecipitation experiments, all three

isoforms bind PRKACA similarly (**Figure S3A and S3B**). However, upon stimulation with forskolin

the S77E/S83E mutant exhibited elevated binding to the catalytic subunit compared to the

S77A/S83A and WT PRKAR1A, indicating that positive charges at this residue may interfere with

release of PKA in response to cAMP (**Figure 3A and S3C**). We then further tested this hypothesis

by assessing PKA kinase activity *in vitro* in response to overexpression of these constructs in

HEK293T cells. Though both WT and S77A/S83A PRKAR1A exhibited attenuated increases in

induced PKA activity, expression of the S77E/S83E mutant exhibited no increase in PKA activity

in response to forskolin (**Figure 3B**). Similarly, S77E/S83E PRKAR1A was able to inhibit

metabolic activity of HEK293T in response to forskolin (**Figure 3C)** and in normal growth

conditions (**Figure S3D)** to a significantly greater degree than either PRKAR1A or STK25

overexpression alone. In order to confirm that PRKAR1A is downstream of STK25, STK25 was

overexpressed simultaneously with knock-down of PRKAR1A resulting in rescue of STK25

inhibition of PKA activity (**Figure 3D and Figure S3E**). Together these data demonstrate a new

regulatory event where STK25 phosphorylation increases the ability of PRKAR1A to bind the

catalytic subunit and inhibit PKA activity in response to cAMP **(Figure 3E)**.

Phosphorylation of PRKAR2A was also detected in our phosphoproteomic analysis

however phosphorylation levels increased with the loss of STK25 **(Table 1).** Despite this, given

its role in PKA regulation in cardiomyocytes, we evaluated PKA holoenzyme activity from

PRKAR1A and PRKAR2A immunoprecipitates. Loss of STK25 increased PKA activity from

PRKAR1A complexes but not PRKAR2A complexes (**Figure S4A and S4B**). Furthermore,

immunoprecipitation experiments did not demonstrate binding of PRKAR2A and STK25 (**Figure**

2D). Together these data do not support PRKAR2A as a downstream effector of STK25.

Stk25* loss increases PKA activity *in vivo

In order to validate the STK25-PKA relationship *in vivo*, we generated an $Stk25^{-/-}$ mouse and

investigated the impact of *Stk25* loss on the Pka pathway *in vivo*. Protein analysis demonstrated

that loss of *Stk25* was accompanied by an increase in Prkar1a levels and a significant decrease

in the ratio of pS77 and pS83 to total Prkar1a (**Figure 4A and S5A**). Whole heart lysates of

$Stk25^{-/-}$ mice displayed increased levels of Pka kinase activity compared to wildtype littermates

(**Figure 4B**). Hearts histologically were similar without any fibrosis (**Figure S5B)** and heart weight

to body weight ratio were not different between genotypes (**Figure S5C**).

Since Pka mediates beta adrenergic stimulation of cardiomyocyte contraction function *in*

vivo, we assessed the impact of *Stk25* loss on cardiac function in response to the β1 agonist

isoproterenol using echocardiography. At baseline, there were no significant differences in either

contractile function or chamber dimensions between genotypes; however $Stk25^{-/-}$ mice displayed

statistically greater individual increases in ejection fraction (EF) and fractional shortening (FS)

and trends towards decreased chamber size in response to isoproterenol (**Figure 4C, 4D and**

S5D). Phosphorylation levels of downstream Pka effectors were also evaluated in response to

isoproterenol and loss of *Stk25* was associated with significant increases in isoproterenol induced

phosphorylation of Pln, TnI, and Ryr2 while CamKII remained unchanged (**Figure S6A and S6B**).

Because prolonged Pka activation has been associated with heart failure (Antos et al.,

2001; Kushnir et al., 2010), a survival cohort of mice at 52 weeks were evaluated by

echocardiography and no significant differences in either heart function or left ventricular size

were found (**Figure S7**). We then investigated human heart failure samples and found that *STK25*

expression was broadly *increased* in myocardial samples from patients with end stage heart

failure when compared to normal heart samples (**Figure 4E and Table S1)**.   Using available

protein samples from this heart failure cohort, we analyzed the phosphorylation status of

PRKAR1A and observed a concordant increase in STK25, PRKAR1A and phosphorylated

PRKAR1A. This raises the possibility that upregulation of STK25 and PRKAR1A has a role in

heart failure and may be involved in the compensatory downregulation of beta adrenergic

signaling.

Discussion

Kinases are integral components of signal transduction pathways, playing key roles in

most cellular processes that are initiated by external cues. Therapeutic modulation of kinases

has demonstrated success in many branches of medicine, highlighted especially by those that

leverage inhibition of kinase cascades implicated in oncogenic processes. Within cardiovascular

biology, kinases regulate ion handling, contractility, and metabolism, and their dysregulation is

involved in many cardiovascular diseases (Clerk et al., 2007; Heineke and Molkentin, 2006;

Vlahos et al., 2003). Several kinases including PKA, Ca2+ calmodulin dependent kinase, protein

kinase C and PIP3 kinase have all been implicated in both physiologic and pathologic signaling

(Dhalla and Muller, 2010). Despite this progress, kinase directed therapy has not been developed

successfully for cardiovascular diseases, and identifying new regulators of kinase pathways is an

area of unrealized therapeutic potential.

Here we present evidence that STK25 is an inhibitor of the PKA pathway, whose

phosphorylation of a regulatory subunit drives inhibition of the catalytic subunit's kinase activity.

Though phosphorylation of the Type II regulatory subunits has been described in detail, little is

known about phosphorylation of Type I regulatory subunits. Phosphorylation at S103 *in vitro* has

been shown to be mediated by protein kinase G and interferes potentially with binding to PRKACA

(Haushalter et al., 2018). Phosphorylation at S83 has been shown to modulate PRKAR1A's

association with the replication factor c complex *in vitro*, the consequence of which is unknown

(Gupte et al., 2006). In the model we propose, phosphorylation at S77 and S83 increases affinity

of PRKAR1A for the catalytic subunit and inhibits cAMP mediated PKA activity. This stands in

contrast to the regulation of Type II subunits whose phosphorylation promotes release of the

catalytic subunit and thus enhances PKA activity.

Though chronic PKA activation has been associated with progression of heart failure in

animal models (Marks, 2013), knockout of *Stk25* resulted in an increase in PKA activity without

evidence of heart failure. This discrepancy merits discussion. First it should be noted that heart

failure mouse models overexpressing the catalytic PKA subunit generate up to 8-fold increases

in PKA activity while in our model, mice experience approximately a 1.6-fold increase in PKA

activity. This is in range of prior work demonstrating no heart failure phenotype with *Prkar1a*

heterozygosity despite an increase in PKA activity by approximately 1.4-fold (Liu et al., 2020).

Furthermore, data from human heart failure samples demonstrate *decreased* phosphorylation of

PKA targets (Najafi et al., 2016) as chronic hyperstimulation of the beta-adrenergic receptor leads

to desensitization and what is thought to be decreased PKA substrate activation (Ungerer et al.,

1993). We observe *increased* levels of STK25 and phosphorylated PRKAR1A in heart failure

samples. This would translate into suppression of PKA stimulation yet it remains to be seen

whether these changes are protective (e.g. as part of a downregulation response to

hyperadrenergic state of heart failure) versus maladaptive (e.g. higher levels of STK25 lead to

worsening cardiac performance) or both. While, differences in either heart function or mortality

are not observed with loss of *Stk25*, further longitudinal studies with overlayed chronic heart failure models are needed to determine if sustained STK25 loss is a potential therapeutic to improve cardiac function safely.

Limitations

There are several important limitations to this study. How exactly phosphorylation of PRKAR1A alters its ability to inhibit the catalytic subunits remains unclear. We observe increased binding of S77E/S83E to the catalytic subunits in forskolin stimulated cells which implies there is either increased affinity between PRKAR1A and the catalytic subunit versus decreased affinity for cAMP by phosphorylated PRKAR1A. Another possibility is a change in PRKAR1A's affinity for an AKAP. Though Type I regulatory subunits generally have weak affinities for AKAP proteins and display a diffuse cytoplasmic localization, the impact that phosphorylation on the assembly of a larger macromolecular regulatory complex remains to be investigated. This is particularly relevant since forskolin did not inhibit PRKAR1A and STK25 binding, yet PRKAR1A phosphorylation was diminished. This would imply that there is either inhibition of STK25 kinase activity by cAMP or a phosphatase that acts on PRKAR1A that is stimulated by cAMP. Given that we observe residual phosphorylation of S77 and S83 in both iPSC-CM and mouse knockout studies, it is likely that there are other kinases that phosphorylate these sites and regulate PKA activity. Further studies are needed to elucidate kinases and phosphatases modulate the phosphorylation status of PRKAR1A what their physiological role is in PKA activity in the heart.

In summary, loss of *STK25* leads to increased PKA activity *in vitro* and *in vivo*. We identify PRKAR1A as a new substrate for STK25 and demonstrate that phosphorylation of PRKAR1A at S77 and S83 inhibits PKA signaling. Loss of Stk25 leads to increased physiologic response to beta adrenergic stimulation of cardiac function without evidence of heart failure. Both STK25 and

1   the phosphorylation of PRKAR1A are upregulated in heart failure and potentially represent a new

2   pathway for downregulation of beta adrenergic signaling in the heart.

3
4
5   **Acknowledgements**
6

We would like to thank the Qing Li for his surgical expertise, Erin Bush at the Columbia Genome

Center for RNA-seq help, Christopher Damoci at the Columbia Mouse Imaging Core Facility and

the labs of Veli Topkara, Elain Wan and Emily Tsai for reagents, technical expertise and tissue

samples.   B.F is supported by a grant from the NHLBI (K08HL140201), the Gerstner Foundation

and the Schwartz Foundation.   G.V.N is supported by grants from the NIH (UH3EB025765,

P41EB027062, and R01HL076485) and NSF (NSF16478).   B.W. T.N and B.L are supported by

the MSTP Training Program (T32GM007367).   B.L. is also supported by the NIH (F30HL145921).

R.K.S is supported by 2P30 CA013696-45 Cancer Center Support Grant.   M.K. is supported by

a Glorney Raisbeck Fellowship Award in Cardiovascular Disease.

Competing Interests: None

Data Availability.

RNA-seq data has been deposited in the Gene Expression Omnibus, NCBI: GSE195514

Proteomics data has been deposited in the PRIDE ProteomeXchange: PXD031367

Original Data can be found at:

https://data.mendeley.com/datasets/t2z9bkm78h/draft?a=a7e3489f-7112-4504-94a4-

33f48aa43867

Material Availability

All unique reagents generated in this study are available from the lead contact with a completed

materials transfer agreement.

Figure 1. *STK25 inhibits PKA activity.* A) Differential phosphoproteomic spectra of *STK25*^{+/+} and *STK25*^{-/-} cardiomyocytes. Members of the PKA signaling pathway are highlighted. B) Ingenuity phosphoprotein pathway analysis. Orange indicates pathways upregulated in *STK25*^{-/-} cardiomyocytes while blue indicates upregulation in *STK25*^{+/+}. C) PKA activity in response to 10μM forskolin treatment for 30 minutes in *STK25*^{+/+} and *STK25*^{-/-} cardiomyocytes. n=3 for each condition. D) PKA activity in response to 10μM forskolin treatment for 30 minutes in HEK293T cells overexpressing either empty vector, wild type STK25 or kinase-dead K49R/T174A. n=3 for each condition. Bar graph data is represented as mean +/- SD, *p<0.05 **p<0.01, ****p<0.0001 using ANOVA and Tukey's adjustment for multiple comparisons.

Figure 2. *STK25 binds to and phosphorylates PRKAR1A.* A) Immunoblot of PRKAR1A phospho-S77 and S83 in *STK25$^{+/+}$* and *STK25$^{-/-}$* cardiomyocyte protein lysates. B) Immunoblots of phosphor-S77 and S83 of PRKAR1A in *STK25$^{-/-}$* cardiomyocytes transfected with empty vector (EV), wild type STK25 and kinase dead K49R/T174A STK25. C) Forskolin (10μM, 30 minutes) stimulated *STK25$^{+/+}$* and *STK25$^{-/-}$* cardiomyocytes immunoblotted for phosphorylation of PRKAR1A. D) Immunoprecipitation of Flag-STK25 expressed in HEK293T cells and immunoblotted for PRKAR1A, PRKA2A and GM130 (positive control binding partner). E) Co-immunoprecipitation of PRKAR1A-V5 with STK25 and PRKACA in HEK293T cells treated with forskolin (10μM, 30 minutes). F) *In vitro* kinase assay of purified STK25 and PRKAR1A, immunoblotted for phosphorylation of S77 and S83 of PRKAR1A.

Figure 3. *Phosphorylation of PRKAR1A inhibits PKA activity.* A) Co-immunoprecipitations of V5 tagged PRKAR1A, S77A/S83A PRKAR1A or S77E/S83 PRKAR1A and immunoblotting for PRKARCA in HEK293T cells stimulated with forskolin (10μM, 30 minutes). B) PKA activity in HEK293T cells stimulated with forskolin (10μM, 30 minutes) and transfected with empty vector, wild type PRKAR1A, S77A/S83A PRKAR1A mutant or S77E/S83E PRKAR1A mutant as indicated. C) HEK293T cells transfected with the indicated vectors and assessed for growth by Real Time Glo for 5 hours after stimulation with forskolin (10μM). D) PKA activity in HEK293T cells stimulated with forskolin (10μM, 30 minutes) and either overexpressing STK25 and/or the siRNA of PRKAR1A. E) Model of STK25 downregulation of the PKA pathway through phosphorylation of PRKAR1A. For all graphs in this figure, n=3 for each condition, data presented as mean +/- SD *p<0.05, ***p<0.001, and ****p<0.0001 by ANOVA with Tukey's adjustment for multiple comparisons.

Figure 4. *Stk25 loss increases response to adrenergic stimulation in vivo.* A) Immmunoblot of Stk25, Prkaca, Gapdh, phospho-S77, phospho-S83 and total Prkar1a in *Stk25^{+/+}* and *Stk25^{-/-}* whole heart lysates. B) *Stk25^{+/+}* and *Stk25^{-/-}* mouse heart lysates were assessed for PKA activity *in vitro*, n=6 for each condition. C) Representative m-mode images of *Stk25^{+/+}* and *Stk25^{-/-}* mouse hearts stimulated with either control or isoproterenol. D) Echocardiographic measurements of ejection fraction (EF) and fractional shortening (FS) at unstimulated baseline and in response to isoproterenol, n=5 for *Stk25^{+/+}* and n=6 for *Stk25^{-/-}*. E) RT-PCR (left) from left ventricular myocardium of normal hearts (n=6) or heart failure (n=17) expressed as a ratio of the threshold cycle curve (Ct) of *STK25* to *GAPDH*. Immunoblot (right) of STK25 and PRKAR1A expression and phosphorylation in protein lysates from left ventricular myocardium of normal hearts or failing hearts. Bar graphs presented as mean +/- SD, *p<0.05, **p<0.01 by student t-test in 4B, repeated measures 2 way ANOVA with Sidak's correction for multiple comparisons in 4D and Welch's t-test in 4E.

Function			Protein Descriptions	Fold Change	Phosphorylation Sites	Protein Accessions
PKA	**Regulatory subunits**	**PRKAR1A**	cAMP-dependent protein kinase type I-alpha regulatory subunit	0.17	S77, S83	P10644
		PRKAR2A	cAMP-dependent protein kinase type II-alpha regulatory subunit	3.03	S78	P13861
	Catalytic Subunit	**PRKACA**	cAMP-dependent protein kinase catalytic subunit alpha	0.79	S339	P17612
PKA targets	**Contractility Ca2+ handling**	**MYBPC**	Myosin-binding protein C, cardiac-type	100.03	T498	Q14896
		Troponin	Troponin T, cardiac muscle	6.18	S132	P45379
				5.75	S285	
				5.02	S189	
		RYR2	Isoform 2 of Ryanodine receptor 2	2.51	S2808	Q92736
				2.14	S4368	
		CACNA1C	Isoform 2 of Voltage-dependent L-type calcium channel subunit alpha-1C	1.65	S1756	Q13936
	Energy Metabolism	**Phosphorylase Kinase**	Phosphorylase b kinase gamma catalytic chain	3.48	T325	P15735
		Glycogen Synthase	Glycogen synthase kinase-3 alpha	2.50	S21	P49840
	Gene Expression	**CREB**	Cyclic AMP-responsive element-binding protein 1	0.79	S271	P16220

Table 1: PKA pathway phosphoproteomic changes in STK25$^{-/-}$ compared to wild type. Fold change represents a ratio of STK25$^{-/-}$/STK25$^{+/+}$. Phosphorylation sites, description and accession numbers for each uniprot ID are listed. All changes met a significance threshold of an FDR corrected q value <0.05.

References

Amrutkar, M., Cansby, E., Chursa, U., Nunez-Duran, E., Chanclon, B., Stahlman, M., Friden, V., Manneras-Holm, L., Wickman, A., Smith, U.*, et al.* (2015a). Genetic Disruption of Protein Kinase STK25 Ameliorates Metabolic Defects in a Diet-Induced Type 2 Diabetes Model. Diabetes *64*, 2791-2804.

Amrutkar, M., Cansby, E., Nunez-Duran, E., Pirazzi, C., Stahlman, M., Stenfeldt, E., Smith, U., Boren, J., and Mahlapuu, M. (2015b). Protein kinase STK25 regulates hepatic lipid partitioning and progression of liver steatosis and NASH. FASEB J *29*, 1564-1576.

Amrutkar, M., Chursa, U., Kern, M., Nunez-Duran, E., Stahlman, M., Sutt, S., Boren, J., Johansson, B.R., Marschall, H.U., Bluher, M.*, et al.* (2016a). STK25 is a critical determinant in nonalcoholic steatohepatitis. FASEB J *30*, 3628-3643.

Amrutkar, M., Kern, M., Nunez-Duran, E., Stahlman, M., Cansby, E., Chursa, U., Stenfeldt, E., Boren, J., Bluher, M., and Mahlapuu, M. (2016b). Protein kinase STK25 controls lipid partitioning in hepatocytes and correlates with liver fat content in humans. Diabetologia *59*, 341-353.

Antos, C.L., Frey, N., Marx, S.O., Reiken, S., Gaburjakova, M., Richardson, J.A., Marks, A.R., and Olson, E.N. (2001). Dilated cardiomyopathy and sudden death resulting from constitutive activation of protein kinase a. Circ Res *89*, 997-1004.

Bae, S.J., Ni, L., and Luo, X. (2020). STK25 suppresses Hippo signaling by regulating SAV1-STRIPAK antagonism. Elife *9*.

Barradeau, S., Imaizumi-Scherrer, T., Weiss, M.C., and Faust, D.M. (2002). Intracellular targeting of the type-I alpha regulatory subunit of cAMP-dependent protein kinase. Trends Cardiovasc Med *12*, 235-241.

Bian, Y., Song, C., Cheng, K., Dong, M., Wang, F., Huang, J., Sun, D., Wang, L., Ye, M., and Zou, H. (2014). An enzyme assisted RP-RPLC approach for in-depth analysis of human liver phosphoproteome. J Proteomics *96*, 253-262.

Bossis, I., and Stratakis, C.A. (2004). Minireview: PRKAR1A: normal and abnormal functions. Endocrinology *145*, 5452-5458.

Bristow, M.R., Ginsburg, R., Minobe, W., Cubicciotti, R.S., Sageman, W.S., Lurie, K., Billingham, M.E., Harrison, D.C., and Stinson, E.B. (1982). Decreased catecholamine sensitivity and beta-adrenergic-receptor density in failing human hearts. N Engl J Med *307*, 205-211.

Burridge, P.W., Matsa, E., Shukla, P., Lin, Z.C., Churko, J.M., Ebert, A.D., Lan, F., Diecke, S., Huber, B., Mordwinkin, N.M.*, et al.* (2014). Chemically defined generation of human cardiomyocytes. Nat Methods *11*, 855-860.

Cansby, E., Amrutkar, M., Manneras Holm, L., Nerstedt, A., Reyahi, A., Stenfeldt, E., Boren, J., Carlsson, P., Smith, U., Zierath, J.R.*, et al.* (2013). Increased expression of STK25 leads to impaired glucose utilization and insulin sensitivity in mice challenged with a high-fat diet. FASEB J *27*, 3660-3671.

Cansby, E., Magnusson, E., Nunez-Duran, E., Amrutkar, M., Pedrelli, M., Parini, P., Hoffmann, J., Stahlman, M., Howell, B.W., Marschall, H.U.*, et al.* (2018). STK25 Regulates Cardiovascular Disease Progression in a Mouse Model of Hypercholesterolemia. Arterioscler Thromb Vasc Biol *38*, 1723-1737.

Chursa, U., Nunez-Duran, E., Cansby, E., Amrutkar, M., Sutt, S., Stahlman, M., Olsson, B.M., Boren, J., Johansson, M.E., Backhed, F.*, et al.* (2017). Overexpression of protein kinase STK25 in mice exacerbates ectopic lipid accumulation, mitochondrial dysfunction and insulin resistance in skeletal muscle. Diabetologia *60*, 553-567.

Clerk, A., Cullingford, T.E., Fuller, S.J., Giraldo, A., Markou, T., Pikkarainen, S., and Sugden, P.H. (2007). Signaling pathways mediating cardiac myocyte gene expression in physiological and stress responses. J Cell Physiol *212*, 311-322.

Colledge, M., and Scott, J.D. (1999). AKAPs: from structure to function. Trends Cell Biol *9*, 216-221.

Daub, H., Olsen, J.V., Bairlein, M., Gnad, F., Oppermann, F.S., Korner, R., Greff, Z., Keri, G., Stemmann, O., and Mann, M. (2008). Kinase-selective enrichment enables quantitative phosphoproteomics of the kinome across the cell cycle. Mol Cell *31*, 438-448.

Dhalla, N.S., and Muller, A.L. (2010). Protein Kinases as Drug Development Targets for Heart Disease Therapy. Pharmaceuticals (Basel) *3*, 2111-2145.

Gancedo, J.M. (2013). Biological roles of cAMP: variations on a theme in the different kingdoms of life. Biol Rev Camb Philos Soc *88*, 645-668.

Gupte, R.S., Traganos, F., Darzynkiewicz, Z., and Lee, M.Y. (2006). Phosphorylation of RIalpha by cyclin-dependent kinase CDK 2/cyclin E modulates the dissociation of the RIalpha-RFC40 complex. Cell Cycle *5*, 653-660.

Haushalter, K.J., Casteel, D.E., Raffeiner, A., Stefan, E., Patel, H.H., and Taylor, S.S. (2018). Phosphorylation of protein kinase A (PKA) regulatory subunit RIalpha by protein kinase G (PKG) primes PKA for catalytic activity in cells. J Biol Chem *293*, 4411-4421.

Hayashi, Y., Hsiao, E.C., Sami, S., Lancero, M., Schlieve, C.R., Nguyen, T., Yano, K., Nagahashi, A., Ikeya, M., Matsumoto, Y.*, et al.* (2016). BMP-SMAD-ID promotes reprogramming to pluripotency by inhibiting p16/INK4A-dependent senescence. Proc Natl Acad Sci U S A *113*, 13057-13062.

Heineke, J., and Molkentin, J.D. (2006). Regulation of cardiac hypertrophy by intracellular signalling pathways. Nat Rev Mol Cell Biol *7*, 589-600.

Jiang, X., Bomgarden, R., Brown, J., Drew, D.L., Robitaille, A.M., Viner, R., and Huhmer, A.R. (2017). Sensitive and Accurate Quantitation of Phosphopeptides Using TMT Isobaric Labeling Technique. J Proteome Res *16*, 4244-4252.

Kim, C., Cheng, C.Y., Saldanha, S.A., and Taylor, S.S. (2007). PKA-I holoenzyme structure reveals a mechanism for cAMP-dependent activation. Cell *130*, 1032-1043.

Kushnir, A., Betzenhauser, M.J., and Marks, A.R. (2010). Ryanodine receptor studies using genetically engineered mice. FEBS Lett *584*, 1956-1965.

Lim, S., Hermance, N., Mudianto, T., Mustaly, H.M., Mauricio, I.P.M., Vittoria, M.A., Quinton, R.J., Howell, B.W., Cornils, H., Manning, A.L.*, et al.* (2019). Identification of the kinase STK25 as an upstream activator of LATS signaling. Nat Commun *10*, 1547.

Liu, Y., Xia, P., Chen, J., Bandettini, W.P., Kirschner, L.S., Stratakis, C.A., and Cheng, Z. (2020). PRKAR1A deficiency impedes hypertrophy and reduces heart size. Physiol Rep *8*, e14405.

Marks, A.R. (2013). Calcium cycling proteins and heart failure: mechanisms and therapeutics. J Clin Invest *123*, 46-52.

Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstrale, M., Laurila, E.*, et al.* (2003). PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat Genet *34*, 267-273.

Najafi, A., Sequeira, V., Kuster, D.W., and van der Velden, J. (2016). beta-adrenergic receptor signalling and its functional consequences in the diseased heart. Eur J Clin Invest *46*, 362-374.

Navarrete-Perea, J., Yu, Q., Gygi, S.P., and Paulo, J.A. (2018). Streamlined Tandem Mass Tag (SL-TMT) Protocol: An Efficient Strategy for Quantitative (Phospho)proteome Profiling Using Tandem Mass Tag-Synchronous Precursor Selection-MS3. J Proteome Res *17*, 2226-2236.

Nerstedt, A., Cansby, E., Andersson, C.X., Laakso, M., Stancakova, A., Bluher, M., Smith, U., and Mahlapuu, M. (2012). Serine/threonine protein kinase 25 (STK25): a novel negative regulator of lipid and glucose metabolism in rodent and human skeletal muscle. Diabetologia *55*, 1797-1807.

Nunez-Duran, E., Chanclon, B., Sutt, S., Real, J., Marschall, H.U., Wernstedt Asterholm, I., Cansby, E., and Mahlapuu, M. (2017). Protein kinase STK25 aggravates the severity of non-alcoholic fatty pancreas disease in mice. J Endocrinol *234*, 15-27.

Olsen, J.V., Blagoev, B., Gnad, F., Macek, B., Kumar, C., Mortensen, P., and Mann, M. (2006). Global, in vivo, and site-specific phosphorylation dynamics in signaling networks. Cell *127*, 635-648.

Olsen, J.V., Vermeulen, M., Santamaria, A., Kumar, C., Miller, M.L., Jensen, L.J., Gnad, F., Cox, J., Jensen, T.S., Nigg, E.A.*, et al.* (2010). Quantitative phosphoproteomics reveals widespread full phosphorylation site occupancy during mitosis. Sci Signal *3*, ra3.

Oppermann, F.S., Gnad, F., Olsen, J.V., Hornberger, R., Greff, Z., Keri, G., Mann, M., and Daub, H. (2009). Large-scale proteomics analysis of the human kinome. Mol Cell Proteomics *8*, 1751-1764.

Piacentino, V., 3rd, Weber, C.R., Chen, X., Weisser-Thomas, J., Margulies, K.B., Bers, D.M., and Houser, S.R. (2003). Cellular basis of abnormal calcium transients of failing human ventricular myocytes. Circ Res *92*, 651-658.

Preisinger, C., Short, B., De Corte, V., Bruyneel, E., Haas, A., Kopajtich, R., Gettemans, J., and Barr, F.A. (2004). YSK1 is activated by the Golgi matrix protein GM130 and plays a role in cell migration through its substrate 14-3-3zeta. J Cell Biol *164*, 1009-1020.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S.*, et al.* (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A *102*, 15545-15550.

Sutt, S., Cansby, E., Paul, A., Amrutkar, M., Nunez-Duran, E., Kulkarni, N.M., Stahlman, M., Boren, J., Laurencikiene, J., Howell, B.W.*, et al.* (2018). STK25 regulates oxidative capacity and metabolic efficiency in adipose tissue. J Endocrinol *238*, 187-202.

Tasken, K., and Aandahl, E.M. (2004). Localized effects of cAMP mediated by distinct routes of protein kinase A. Physiol Rev *84*, 137-167.

Taylor, S.S., Buechler, J.A., and Yonemoto, W. (1990). cAMP-dependent protein kinase: framework for a diverse family of regulatory enzymes. Annu Rev Biochem *59*, 971-1005.

Taylor, S.S., Ilouz, R., Zhang, P., and Kornev, A.P. (2012). Assembly of allosteric macromolecular switches: lessons from PKA. Nat Rev Mol Cell Biol *13*, 646-658.

Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., Mann, M., and Cox, J. (2016). The Perseus computational platform for comprehensive analysis of (prote)omics data. Nat Methods *13*, 731-740.

Ungerer, M., Bohm, M., Elce, J.S., Erdmann, E., and Lohse, M.J. (1993). Altered expression of beta-adrenergic receptor kinase and beta 1-adrenergic receptors in the failing human heart. Circulation *87*, 454-463.

Vlahos, C.J., McDowell, S.A., and Clerk, A. (2003). Kinases as therapeutic targets for heart failure. Nat Rev Drug Discov *2*, 99-113.

Wang, J., Gareri, C., and Rockman, H.A. (2018). G-Protein-Coupled Receptors in Heart Disease. Circ Res *123*, 716-735.

Zhou, H., Di Palma, S., Preisinger, C., Peng, M., Polat, A.N., Heck, A.J., and Mohammed, S. (2013). Toward a comprehensive characterization of a human cancer cell phosphoproteome. J Proteome Res *12*, 260-271.

Title: STK25 inhibits PKA signaling by phosphorylating PRKAR1A

Authors: Xiaokan Zhang[1], Bryan Z. Wang[1,2], Michael Kim[1], Trevor R. Nash[1,2], Bohao Liu[1,2], Jenny Rao[1], Roberta Lock[2], Manuel Tamargo[2], Rajesh Kumar Soni[3], John Belov[1], Eric Li[1], Gordana Vunjak-Novakovic[1,2] and Barry Fine[1*]

Affiliations

[1] Department of Medicine, Division of Cardiology, Columbia University Medical Center, New York, NY, USA

[2] Department of Biomedical Engineering, Columbia University, New York, NY, USA

[3] Proteomics and Macromolecular Crystallography Shared Resource, Herbert Irving Comprehensive Cancer Center, Columbia University Irving Medical Center, New York, NY, USA

*Corresponding Author

Barry Fine MD PhD

Department of Medicine, Division of Cardiology

622 West 168th Street

PH8-405B

New York, NY 10032

Supplementary Materials:

Figures S1-*S7*

Table S1: *Patient characteristics of end stage heart failure samples*

Figure S1. Comparison of STK25 expression in adult mice cardiomyocytes and in iPSC-CMs. A) Immunoblot and B) densitometry of STK25 and GAPDH expression in primary cardiomyocytes isolated from adult mouse heart and in iPSC-CMs. n=3 for each condition. Data presented as mean +/- SD, statistical significance was tested by student t-test in S1B.

Figure **S2**. A) Densitometry analysis of phospho-S77 and S83 relative to total PRKAR1A in immunoblot shown in Figure 2A. B) Densitometry analysis of phospho-S77 and S83 relative to total PRKAR1A in immunoblot shown in Figure 2B. C) Densitometry analysis of phospho-S77 and S83 relative to total PRKAR1A in immunoblot shown in Figure 2C. D) Co-immunoprecipitation experiments of either wild type or kinase dead (K49R/T147A) mutant of STK25 in HEK293T cells with overexpressed flag tagged STK25 protein and endogenous PRKAR1A. For all bar graphs in this figure, n=3 for each condition, Data presented as mean +/- SD, *p<0.05, **p<0.01, and ****p<0.0001 by student's t-test in S2A and S2B and two way ANOVA with Tukey's adjustment for multiple comparisons in S2C.

Figure S3. A) Co-immunoprecipitation of PRKAR1A-V5 with STK25 and PRKACA in HEK293T cells without forskolin treatment. B) Densitometry analysis of PRKACA relative to PRKAR1A in immunoblot shown in Figure S2A. C) Densitometry analysis of PRKACA relative to PRKAR1A in immunoblot shown in Figure 3A. D) HEK293T cells transfected with the indicated vectors and assessed for growth by Real Time Glo (Promega) over three days. E) Immunoblot demonstrating both overexpression of STK25 and knockdown of PRKAR1A in HEK293T cells shown in Figure 3D. For all graphs in this figure, n=3 for each condition or time point, Data presented as mean +/- SD, **p<0.01, ***p<0.001 and ****p<0.0001 one way ANOVA in S3A and S3C and two way ANOVA, all with Tukey's adjustment for multiple comparisons.

Figure S4: *Comparison of Type I versus Type II holoenzyme activity in response to STK25 loss.* A) Immunoprecipitations of PRKAR1A or PRKAR2A in STK25$^{+/+}$ and STK25$^{-/-}$ iPSC-CMs and immunoblotting for PRKAR1A or PRKAR2A, PRKACA and GAPDH. B) PKA activity of type I and II PKA holoenzymes precipitated from cell lysates of STK25$^{+/+}$ and STK25$^{-/-}$ iPSC-CMs. N=3 for each condition, mean +/- SD, ****p<0.0001 two way ANOVA with Sidak's adjustment for multiple comparisons.

Figure S5. A) Densitometry analysis phospho-S77 and S83 relative to total Prkar1a in immunoblot shown in Figure 4A. B). Representative pentachrome staining of mid ventricle sections of 20-week old *Stk25*$^{+/+}$ and *Stk25*$^{-/-}$ mice. C) Heart weight (HW) to body weight (BW) ratio of mouse hearts at 20 weeks. D) Chamber measurements LVEDD (mm) and LVESD (mm) of mice at baseline and treated with isoproterenol. n=5 for *Stk25*$^{+/+}$and n=6 for *Stk25*$^{-/}$ mice^{-}. Data presented as mean +/- SD, statistical significance was tested with student's t-test in S5A, Welch's t-test in S5C, and repeated measures 2-way ANOVA with Sidak's correction for multiple comparisons in S5D.

Figure S6: *Stk25 knockout and beta-adrenergic response*. A) Immunoblot of phospho-S23/S24 and total TnI, phospho-S16/T17 and total Pln, phospho-S2808 and total Ryr2, phospho-T286 and total CamkII in *Stk25*$^{+/+}$ and *Stk25*$^{-/-}$ whole heart lysates with and without isoproterenol injection. B) Densitometry analysis of phospho-S23/S24 relative to total TnI, phospho-S16/T17 relative to total Pln, phospho-S2808 relative to total Ryr2, phospho-T286 relative to total CamkII in immunoblot shown in Figure S6C, n=3 for each condition. Data presented as mean +/- SD, *p<0.05, **p<0.01, statistical significance was tested with two way ANOVA with Tukey's correction for multiple comparisons.

Figure S7: *Cardiac function at one year in Stk25-/-mice.* Echocardiographic measurements of LV function and size in a survival cohort at 52 weeks for Stk25+/+ (n=8) and Stk25-/-(n=10). Data presented as mean +/- SD, statistical significance was tested with a Welch's t-test.

Patient Characteristics (n=17)		
	Age (SD)	49.47 (12.8)
	Gender (Male%)	82.4
	BMI (SD)	24.4 (3.3)
Cardiomyopathy		
	Non-ischemic (%)	70.6
	Ischemic (%)	29.4
Comorbidities		
	Hypertension (%)	17.6
	Hyperlipidemia (%)	29.4
	Diabetes(%)	23.5
Heart Function		
	Ejection Fraction (%)	17.9
	LVAD	88.23
	Inotrope	35.3
Medical Therapy		
	ACEi/ARB/ARNI	29.4
	Beta Blocker	76.5
	MRA	52.9
	Anticoagulation	88.23
	Antiarrhythmic	41.2

Table S1: *Patient characteristics of end stage heart failure samples*

Abbreviations: SD, standard deviation; BMI, body mass index; LVAD, left ventricular assist device; ACEi, angiotensin converting enzyme inhibitor; ARB, angiotensin II receptor blocker; ARNI angiotensin II receptor blocker and neprisilyn inhibitor; MRA, mineralcorticoid receptor antagonist.