Macrophages regulate gastrointestinal motility through complement component 1q

Mihir Pendse[1], Yun Li[1], Cristine N. Salinas[1], Gabriella Quinn[1], Nguyen Vo[1], Daniel C. Propheter[1],

Chaitanya Dende[1], Alexander A. Crofts[1], Eugene Koo[1], Brian Hassell[1], Kelly A. Ruhn[1], Prithvi Raj[1],

Yuuki Obata[1*], Lora V. Hooper[1,2*]

[1]Department of Immunology, The University of Texas Southwestern Medical Center, Dallas, TX 75390

[2]The Howard Hughes Medical Institute, The University of Texas Southwestern Medical Center, Dallas, TX 75390

*For correspondence: yuuki.obata@utsouthwestern (YO), lora.hooper@utsouthwestern.edu (LVH)

ABSTRACT

Peristaltic movement of the intestine propels food down the length of the gastrointestinal tract to promote nutrient absorption. Interactions between intestinal macrophages and the enteric nervous system regulate gastrointestinal motility, yet we have an incomplete understanding of the molecular mediators of this crosstalk. Here we identify complement component 1q (C1q) as a macrophage product that regulates gut motility. Macrophages were the predominant source of C1q in the mouse intestine and most extraintestinal tissues. Although C1q mediates complement-mediated killing of bacteria in the bloodstream, we found that C1q was not essential for immune defense of the intestine. Instead, C1q-expressing macrophages were localized to the intestinal submucosal plexus where they were closely associated with enteric neurons and expressed surface markers characteristic of nerve-adjacent macrophages in other tissues. Mice with a macrophage-specific deletion of *C1qa* showed changes in enteric neuronal gene expression, increased peristaltic activity, and accelerated intestinal transit. Our findings identify C1q as a key regulator of gastrointestinal motility and provide enhanced insight into the crosstalk between macrophages and the enteric nervous system.

INTRODUCTION

Peristalsis is the physical force that propels food through the intestine, promoting digestion and nutrient absorption. The gastrointestinal motility that underlies peristalsis is a complex process that requires coordination of the activity of smooth muscle cells by enteric neurons (Rao and Gershon, 2016). Recent studies have revealed that intestinal macrophages impact gastrointestinal motility by regulating the functions of enteric neurons and facilitating their interactions with smooth muscle cells (Muller et al., 2014; Matheis et al., 2020).

Macrophages carry out diverse functions in the intestine that vary according to their anatomic location. For example, macrophages that localize to the tissue located directly underneath the gut epithelium — known as the lamina propria — contribute to immune defense against pathogenic bacteria

(Gabanyi et al., 2016). A distinct group of macrophages localizes to the tissues located beneath the lamina

propria, between the circular and longitudinal muscle layers in the tissue region known as the muscularis

externa. These muscularis macrophages express genes that are distinct from lamina propria macrophages

(Gabanyi et al., 2016). They directly regulate the activity of smooth muscle cells (Luo et al., 2018) and

secrete soluble factors such as bone morphogenetic protein 2 (BMP2) that interact with the enteric neurons

that control smooth muscle activity (Muller et al., 2014). Muscularis macrophages thus play a key role in

regulating gut motility. However, our understanding of the molecular mechanisms by which these

macrophages regulate intestinal neuromuscular activity and gut motility remains limited.

C1q is a member of the defense collagen family that has distinct functions in immune defense and

nervous system development and function (Bossi et al., 2014; Casals et al., 2019; Shah et al., 2015; Thielens

et al., 2017). C1q is composed of six molecules each of C1qA, C1qB, and C1qC, forming a 410 kDa

oligomer. C1q circulates in the bloodstream, where it participates in immune defense against infection by

recognizing antibodies bound to invading bacteria. This binding interaction initiates the classical

complement pathway, which entails the recruitment and proteolytic processing of other complement

components that rupture the bacterial membrane and recruit phagocytic cells (Kishore and Reid, 2000;

Noris and Remuzzi, 2013). C1q is also produced by microglia (brain-resident macrophage-like cells) in the

brain where it promotes the pruning of neuronal synapses through an unclear mechanism (Hammond et al.,

2020; Hong et al., 2016). Consequently, C1q deficiency results in heightened synaptic connectivity in the

central nervous system that can lead to epilepsy (Chu et al., 2010).

C1q is also produced at barrier sites, such as the intestine, where encounters with commensal and

pathogenic microbes are frequent. However, little is known about the physiological role of C1q in barrier

tissues. Liver immune cells, including macrophages and dendritic cells, produce serum C1q; however the

cellular source of C1q in barrier tissues including the intestine remains unclear (Petry et al., 2001). Here,

we show that C1q is selectively produced by macrophages that inhabit the submucosal plexus of the mouse

intestine. These C1q-expressing macrophages are located close to enteric neurons that have a known role

in controlling gut motility. Consistent with their nerve-adjacent localization, mice lacking macrophage C1q

exhibit altered expression of enteric neuronal genes, increased peristaltic activity, and accelerated

gastrointestinal motility. These findings identify C1q as a key mediator of a neuroimmune interaction that

regulates gut motility.

RESULTS

C1q is expressed in intestinal macrophages

Soluble defense collagens are an ancient, evolutionarily conserved family of antimicrobial proteins

with shared structural features including a *C*-terminal globular head and a collagen-like region (Casals et

al., 2019). Little is known about the function of defense collagens at mucosal barrier sites, where microbial

encounter is frequent. Our initial goal in this study was to identify soluble defense collagens that are

expressed by the mouse intestine and to assess their role in host defense. We therefore measured the

expression of 18 defense collagen genes in the mouse small intestine and colon by RNA sequencing (RNA-

seq). This analysis revealed that the most abundant soluble defense collagen transcripts in the small intestine

and colon were those encoding C1qA, C1qB, and C1qC (***Figure 1A; Figure 1 – figure supplement 1***).

Serum C1q is produced by liver dendritic cells, monocytes, and macrophages (El-Shamy et al.,

2018). However, the cellular source(s) of C1q in peripheral tissues including the intestine is unknown.

Quantitative PCR (qPCR) analysis of fluorescence-activated cell sorting (FACS)-sorted cell suspensions

recovered from the small intestines of wild-type C57BL/6 mice revealed that *C1qa*, *C1qb*, and *C1qc*

transcripts were most abundant in $CD45^+$ cells, which include all immune cells, as compared to $CD45^-$ cells,

which encompass non-immune cells such as epithelial cells (***Figure 1B***). Further, C1q transcripts and

protein were most abundant in $CD45^+$ cells recovered from the subepithelial compartment, which includes

both the lamina propria and muscularis, as compared to $CD45^+$ cells recovered from the intraepithelial

compartment of the small intestine (***Figure 1C and D***). Thus, C1q is expressed by immune cells located in

the subepithelial compartment of the intestine and is largely absent from epithelial cells and intraepithelial

immune cells.

To identify the intestinal immune cell population that expresses C1q, we further analyzed the

subepithelial CD45$^+$ cell population by flow cytometry. Expression of C1q transcripts and protein was

highest in CD11b$^+$MHCII$^+$F4/80^{hi} macrophages and was mostly absent from all non-macrophage immune

cells (***Figure 1E-H***). Thus, C1q is expressed by macrophages in the mouse small intestine.

Figure 1. C1q is expressed by macrophages in the mouse small intestine.
(A) RNA-seq analysis of soluble defense collagen expression in the small intestines (ileum) of C57BL/6 mice. Data
were adapted from a previously published RNA-seq analysis (Gattu et al., 2019). Data are available in the Gene
Expression Omnibus repository under accession number GSE122471. Each column represents one mouse.
(B) qPCR measurement of *C1qa*, *C1qb*, and *C1qc* transcript abundance in CD45$^+$ and CD45$^-$ cells purified from mouse
small intestines by flow cytometry. Each data point represents one mouse, and results are representative of two
independent experiments.
(C) qPCR measurement of *C1qa*, *C1qb*, and *C1qc* transcript abundance in subepithelial and intraepithelial cells
recovered from mouse small intestines. Each data point represents one mouse, and results are representative of three
independent experiments.
(D) Representative immunoblot of subepithelial and intraepithelial cells recovered from mouse small intestines, with
detection of C1q and actin (control). Each lane represents cells from one mouse and immunoblot is representative of
three independent experiments.
(E) Flow cytometry gating strategy for analysis of mouse small intestinal cell suspensions in panels F, G, and H. Cells
were pre-gated as live CD45$^+$ cells. SSC, side-scatter; MHCII, major histocompatibility complex II.
(F) qPCR measurement of *C1qa*, *C1qb*, and *C1qc* transcript abundance in cells isolated by flow cytometry from mouse
small intestines as indicated in (E). Each data point represents cells pooled from three mice, and results are
representative of three independent experiments.
(G) Flow cytometry analysis of intracellular C1q in small intestinal subepithelial cells identified as indicated in (E).
(H) Quantitation of flow cytometry analysis in (G). Each data point represents one mouse, and results are
representative of two independent experiments.

105
106 Sm. int., mouse small intestine; Error bars represent SEM. **p<0.01; ***p<0.001; ****p<0.0001; ns, not significant
107 by one way ANOVA (A,F) or two-tailed Student's t-test (B,C,H).
108
109 **Figure supplement 1.** C1q is expressed in the mouse colon.
110

Macrophages are the primary source of C1q in the mouse gastrointestinal tract

We next assessed whether macrophages are the primary source of C1q in the intestine by analyzing

two mouse models. First, we depleted macrophages by injecting neutralizing antibodies directed against

the receptor for colony stimulating factor 1 (CSF1R)(***Figure 2A***), which is required for the development of

a subset of lamina propria macrophages (Bogunovic et al., 2009) and all muscularis macrophages (Muller

et al., 2014). Antibody injection led to a >2-fold reduction in the number of macrophages recovered from

the small intestine (***Figure 2B***), and a corresponding >2-fold reduction in small intestinal C1q gene

expression (***Figure 2C***), suggesting that macrophages are the primary source of intestinal C1q.

Second, we constructed a genetic model of C1q deficiency by crossing $C1qa^{fl/fl}$ mice (Fonseca et

al., 2017) to mice carrying the $Lyz2$-Cre transgene (LysM-Cre mice), which is selectively expressed in

myeloid cells including macrophages (***Figure 2D***). These mice, hereafter designated as $C1qa^{\Delta M\phi}$ mice,

lacked C1q expression in intestinal macrophages (***Figure 2E and F***). Importantly, $C1qa^{\Delta M\phi}$ mice had

markedly lower overall C1q expression in both the small intestine and colon (***Figure 2G***), indicating that

macrophages are the main source of C1q in the intestine. Unexpectedly, the $C1qa^{\Delta M\phi}$ mice also lost $C1q$

gene expression in the lung, skin, kidney, and liver (but not the brain), and C1q protein was undetectable

in the serum (***Figure 2 – figure supplement 1***). These findings indicate that macrophages are the primary

source of C1q in the intestine and suggest that $LysM^{+}$ macrophages or macrophage-like cells are also the

main source of C1q in most extraintestinal tissues and the bloodstream.

Figure 2. Macrophages are the primary source of C1q in the mouse gastrointestinal tract.

(A) Graphic showing selective depletion of macrophages in C57BL/6 mice by intraperitoneal injection of anti-CSF1R antibody. Control mice were injected with isotype-matched non-specific antibody. Mice were analyzed 72 hours after antibody injection. Panel was generated at Biorender.com.

(B) Representative flow cytometry analysis of mouse small intestines after intraperitoneal injection of anti-CSF1R or isotype control antibody. All cells were gated as live CD45$^+$. Macrophages were MHCII$^+$ F4/80^{hi}; B cells were CD19$^+$; T cells were CD3$^+$. Total small intestinal cell yields were $1.5 \times 10^6 \pm 4.9 \times 10^5$ cells.

(C) qPCR measurement of *C1qa*, *C1qb*, and *C1qc* transcript abundance in mouse small intestines after intraperitoneal injection of anti-CSF1R or rat IgG2a (isotype control). Each data point represents one mouse and results are pooled from two independent experiments.

(D) *C1qa*$^{fl/fl}$ mice were crossed with mice harboring a LysM-Cre transgene to generate mice having a macrophage-selective deletion of *C1qa* (*C1qa*$^{\Delta M\varphi}$ mice). Panel was generated at Biorender.com.

(E) Representative flow cytometry analysis of intracellular C1q expression in small intestinal macrophages from *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\varphi}$ mice. Mice were littermates from heterozygous crosses that remained co-housed. Cells were gated on live CD45$^+$CD11b$^+$MHCII$^+$.

(F) Quantitation of the flow cytometry analysis in (E). Each data point represents one mouse. Results are representative of two independent experiments.

(G) qPCR measurement of *C1qa* transcript abundance in the small intestines (sm. int.) and colons of *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\varphi}$ littermates. Each data point represents one mouse.

Error bars represent SEM. **p<0.01; ***p<0.001; ****p<0.0001; ns, not significant by two-tailed Student's t-test.

Figure supplement 1. C1q expression is lost systemically but preserved in the central nervous system of *C1qa*$^{\Delta M\varphi}$ mice.

$C1qa^{\Delta M\varphi}$ mice do not show altered microbiota composition, barrier function, or resistance to enteric infection

The classical complement pathway is a well-studied host defense system that protects against systemic pathogenic infection (Warren et al., 2002; Noris and Remuzzi, 2013). Circulating C1q activates the complement pathway by binding to antibody-antigen complexes or to bacterial cell surface molecules, and thus protects against systemic infection. We therefore assessed whether C1q promotes immune defense of the intestine.

We first determined whether C1q exhibits characteristics of known intestinal antimicrobial proteins, including induction by the intestinal microbiota and secretion into the gut lumen. *C1qa* was expressed at similar levels in the small intestines of germ-free and conventionally-raised mice (***Figure 3A***), suggesting that *C1q* expression is not induced by the gut microbiota. This contrasted with *Reg3g*, encoding the antimicrobial protein REG3G (Cash et al., 2006), which was expressed at a >2-fold higher level in conventional as compared to germ-free mice (***Figure 3A***). Additionally, in contrast to REG3G, C1q was not detected in the gut lumen of either conventional or germ-free mice (***Figure 3B***). *C1qa* expression was also not markedly altered by oral infection with the intestinal pathogenic bacterial species *Salmonella* Typhimurium (***Figure 3C***). These data indicate that C1q is not induced by the gut microbiota or by the bacterial pathogen *S.* Typhimurium, in contrast to other antibacterial proteins.

We next assessed whether C1q regulates the composition of the gut microbiota. 16*S* rRNA gene sequencing analysis of the fecal microbiotas of $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ mice showed that the microbiota composition was not appreciably altered in the absence of macrophage C1q (***Figure 3D***). We also challenged $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ mice with dextran sulfate sodium (DSS), which damages the colonic epithelium and exposes underlying tissues to the commensal microbiota. However, the sensitivity of the $C1qa^{\Delta M\phi}$ mice to DSS was similar to that of their $C1qa^{fl/fl}$ littermates as assessed by change in body weight (***Figure 3E***). There was also no change in intestinal paracellular permeability in $C1qa^{\Delta M\phi}$ mice as measured

by oral administration of FITC-dextran (***Figure 3F***). These results suggest that macrophage C1q does not

substantially impact gut microbiota composition or intestinal epithelial barrier function.

To determine whether C1q protects against enteric infection we conducted oral infection

experiments with the enteric pathogen *Citrobacter rodentium*. We chose *C. rodentium* as our model

organism for two reasons. First, *C. rodentium* is a non-disseminating pathogen, allowing us to test

specifically for C1q's role in intestinal infection. Second, *C. rodentium* clearance depends on

immunoglobulins and complement component C3 (Belzer et al., 2011). Because C1q is bactericidal in

concert with C3 and immunoglobulins, we predicted that $C1qa^{\Delta M\phi}$ mice would show enhanced

susceptibility to *C. rodentium* infection. However, $C1qa^{\Delta M\phi}$ mice cleared *C. rodentium* similarly to their

$C1qa^{fl/fl}$ littermates, indicating that C1q is dispensable for defense against *C. rodentium* infection (***Figure***

3G).

We also did not observe marked alterations in immunity in the absence of C1q. Comparison of

cytokine transcript abundance in the small intestines of $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ littermates revealed no

statistically significant differences in cytokine expression (***Figure 3H***). There were also no statistically

significant differences in the percentages or absolute numbers of various T cell subsets, including $T_{helper}1$

(T_H1), T_H2, T_H17, and regulatory T (T_{reg}) cells between $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ mice (***Figure 3I; Figure 3 –***

figure supplement 1). Although total B cell numbers trended lower in $C1qa^{\Delta M\phi}$ mice, the difference was

not statistically significant (***Figure 3I; Figure 3 – figure supplement 2***). There were also no statistically

significant differences in the percentages or absolute numbers of total plasma cells (***Figure 3I; Figure 3 –***

figure supplement 2), IgA$^+$ plasma cells (***Figure 3I; Figure 3 – figure supplement 2***), myeloid cells

(***Figure 3I; Figure 3 – figure supplement 3***), or innate lymphoid cells (***Figure 3I; Figure 3 – figure***

supplement 4) when comparing $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ mice. These results suggest that the absence of

macrophage C1q has little impact on intestinal immunity. Altogether, our findings suggest that C1q does

not participate substantially in intestinal immune defense and thus might have an intestinal function that is

independent of its canonical role in activating the classical complement pathway.

Figure 3. *C1qa*^ΔMφ mice do not show altered microbiota composition, barrier function, or resistance to enteric infection.

(A) Small intestinal *C1qa* expression is not induced by the intestinal microbiota. qPCR measurement of *Reg3g* and *C1qa* transcript abundances in the small intestines of germ free (GF) and conventional (CV) C57BL/6 mice. Each data point represents one mouse and results are representative of two independent experiments.

(B) C1q is not detected in the mouse intestinal lumen or feces. Representative immunoblot of an ammonium sulfate precipitation of intestinal lumenal contents and feces from germ-free and conventional mice with detection of C1q. C1q in small intestinal tissue (sm. int.) is shown for comparison at right. REG3G was detected as a control, as it is secreted into the intestinal lumen of conventional mice (Cash et al., 2006). Each lane represents multiple mice pooled (n = 5 and 9 for germ-free and conventional, respectively) and the immunoblot is representative of three independent experiments.

(C) C1q gene expression is not altered by enteric infection with *S.* Typhimurium. qPCR measurement of *C1qa* transcript abundance in small intestinal tissue after oral inoculation of mice with 10^9 colony forming units of *S.* Typhimurium strain SL1344. Each data point represents one mouse, and results are representative of two independent experiments.

(D) Intestinal microbiota composition is not altered in *C1qa*^ΔMφ mice. Phylogenetic analysis of 16*S* rRNA gene sequences from fecal pellets collected from *C1qa*^fl/fl and *C1qa*^ΔMφ littermates. Operational taxonomic units with an

average of 100 reads and populations greater than or equal to 1% were included in the graphical analysis. Each bar
represents one mouse. Data are available from the Sequence Read Archive under BioProject ID PRJNA793870.
(E) *C1qa*$^{\Delta M\varphi}$ mice do not show altered susceptibility to dextran sulfate sodium (DSS)-induced colitis. Mice were
provided with 3% DSS in drinking water and body weights were monitored for 7 days. n = 4 and 6 for *C1qa*$^{fl/fl}$ and
C1qa$^{\Delta M\varphi}$ littermates, respectively. Error bars represent SEM. Differences at each time point were not significant by
two-tailed Student's t-test.
(F) *C1qa*$^{\Delta M\varphi}$ mice do not show altered intestinal permeability. To measure intestinal permeability, *C1qa*$^{fl/fl}$ and
C1qa$^{\Delta M\varphi}$ littermates were gavaged with FITC-dextran, and serum FITC-dextran levels were determined by
fluorescence microplate assay against a FITC-dextran standard curve. Indomethacin induces intestinal damage in mice
and was used as a positive control.
(G) Time course of fecal *C. rodentium* burden following oral gavage of *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\varphi}$ mice with 5×10^8 colony
forming units (CFU) of *C. rodentium*. n = 5 and 5 for *C1qa*$^{fl/fl}$ and $^{C1qa\Delta M\varphi}$ littermates, respectively. Error bars represent
SEM. Differences at each time point were not significant by two-tailed Student's t-test.
(H) qPCR measurement of cytokine transcript abundances in the small intestines of *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\varphi}$ littermates.
Error bars represent SEM. Statistics were performed with two-tailed Student's t-test.
(I) Flow cytometry analysis of small intestinal immune cell subsets from *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\varphi}$ littermates. Gating
strategies are shown in Figure 3 - figure supplement 1 through 4. ILC, innate lymphoid cell. Total small intestinal cell
yields were $8.8 \times 10^6 \pm 2.9 \times 10^6$ cells.
243
244 Error bars represent SEM. **p<0.01; ns, not significant by two-tailed Student's t-test.
245
Figure supplement 1. Flow cytometry gating strategy for comparison of T cell populations in *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\varphi}$
mice.
Figure supplement 2. Flow cytometry gating strategy for comparison of B cell and plasma cell populations in *C1qa*$^{fl/fl}$
and *C1qa*$^{\Delta M\varphi}$ mice.
Figure supplement 3. Flow cytometry gating strategy for comparison of myeloid cell populations in *C1qa*$^{fl/fl}$ and
C1qa$^{\Delta M\varphi}$ mice.
Figure supplement 4. Flow cytometry gating strategy for comparison of innate lymphoid cell populations in *C1qa*$^{fl/fl}$
and *C1qa*$^{\Delta M\varphi}$ mice.
254
255
256

C1q-expressing gut macrophages are located near enteric neurons

Intestinal macrophages perform distinct functions depending on their anatomical location.

Macrophages in the lamina propria protect against invasion by pathogenic microbes and promote tissue

repair (Grainger et al., 2017). In contrast, macrophages that reside in deeper intestinal tissues, including the

submucosal plexus and muscularis (***Figure 4A***), regulate enteric smooth muscle cells and neurons that

drive gastrointestinal motility (De Schepper, Stakenborg, et al., 2018; De Schepper, Verheijden, et al.,

2018).

Immunofluorescence analysis revealed that C1q was located primarily beneath the lamina propria,

in a region consistent with the location of the submucosal plexus of the small intestine and colon (***Figure***

4A and B). C1q was detected in a diffuse pattern close to macrophages in *C1qa*$^{fl/fl}$ mice and was absent in

C1qa$^{\Delta M\phi}$ mice (***Figure 4B***) despite the presence of similar overall numbers of CD169$^+$ macrophages

(*Figure 4 – figure supplement 1*). Although we did not observe direct colocalization of C1q with CD169,

a macrophage marker that selectively marks nerve-adjacent macrophages (Ural et al., 2020), the pattern

was consistent with prior observations of C1q deposition in the extracellular matrix in the arterial wall (Cao

et al., 2003)(*Figure 4B*).

We next assessed the location of C1q relative to enteric neurons in the mouse small intestine.

Detection of the neuronal marker βIII tubulin (TUBB3) revealed the presence of two distinct layers of

enteric neurons: one just beneath the lamina propria that was consistent with the location of the submucosal

plexus, and one in the muscularis that was consistent with the location of the myenteric plexus (*Figure 4C*).

Immunofluorescence detection of C1q alongside TUBB3 revealed that C1q was located within an average

of ~5 μm of the nearest TUBB3$^+$ enteric neurons in the submucosal plexus and was largely absent from the

myenteric plexus (*Figure 4C*). Thus, C1q-producing gut macrophages are located close to enteric neurons

in the submucosal plexus.

C1q-producing intestinal macrophages exhibited several characteristics of nerve-adjacent

macrophages in other tissues. First, prior studies have shown that microglia in the central nervous system

and nerve-adjacent macrophages in peripheral tissues, such as the lung, are enriched for C1q gene

expression (Fonseca et al., 2017; Ural et al., 2020). This is consistent with our observations of C1q

expression in nerve-adjacent intestinal macrophages located in the submucosal plexus (*Figure 4C*). Second,

nerve-adjacent macrophages depend on a CSF1-CSF1R signaling axis for their maintenance and are

therefore sensitive to anti-CSF1R antibody-mediated depletion (Muller et al., 2014). Thus, our finding that

anti-CSF1R administration depleted C1q-expressing intestinal macrophages is consistent with their location

near enteric neurons (*Figure 2A-C*). Third, C1q-expressing intestinal macrophages showed elevated

expression of Arginase 1, CD169, and TREM2 (triggering receptor expressed on myeloid cells 2)(*Figure

4D*), which are enriched on macrophages with known neuromodulatory functions (Colonna, 2003; Paloneva

et al., 2002; Ural et al., 2020). Thus, C1q-expressing intestinal macrophages are located near enteric neurons

and exhibit phenotypes characteristic of nerve-adjacent macrophages in other tissues.

Figure 4. C1q-expressing macrophages are closely associated with neurons in the mouse intestine.
(A) Graphic depicting the muscularis of the mouse small intestine. The lumen, epithelium (epi), lamina propria (LP), submucosal plexus (SP), and distinct anatomical regions of the muscularis are indicated. Created at Biorender.com
(B) Immunofluorescence detection of C1q (violet) and macrophages marked with CD169 (green) in the small intestine and colon of $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ littermates. Nuclei were detected with 4',6-diamidino-2-phenylindole (DAPI; blue). Detection with isotype control antibodies on $C1qa^{fl/fl}$ small intestines is shown at right. Anti-rat IgG AlexaFluor 488 and streptavidin-Cy5 were used as secondary stains for CD169 and C1q respectively. The intestinal surface is denoted with a red dotted line and the gut lumen, epithelium, and lamina propria are indicated. The approximate region encompassing the submucosal plexus and the muscularis is denoted with two white dotted lines. Examples of C1q⁺ areas are indicated with yellow arrows and examples of CD169⁺ macrophages are indicated with white arrowheads. Note that the violet staining near the bottom of the muscularis is non-specific, as indicated by its presence in the isotype control image. Images are representative of three independent experiments. Scale bars = 50 μm.
(C) Immunofluorescence detection of C1q (violet), macrophages marked with CD169 (green) and neurons marked with TUBB3 (red) in the small intestines of wild-type C57BL/6 mice. Nuclei are detected with DAPI (blue). The epithelium, lamina propria are indicated. The approximate region encompassing the submucosal plexus and the muscularis is denoted with two white dotted lines. The expanded image area delineated by a yellow square shows an example of the close association between C1q and TUBB3⁺ neurons. Images are representative of images captured from three mice. Anti-rat IgG AlexaFluor 488, anti-rabbit IgG AlexaFluor 594, and streptavidin-Cy5 were used as secondary stains for CD169, TUBB3, and C1q respectively, and an isotype control image is shown at upper right. Scale bars = 50 μm. Lower right panel: Distances between C1q⁺ areas and the nearest TUBB3⁺ neurons were measured in images captured from three mice. Each data point represents one measurement and measurements were made on 10 to 15 microscopic fields for each mouse (71, 70, and 57 total measurements, respectively).
(D) Flow cytometry analysis of CD169, Arginase 1, and TREM2 on C1q⁻ and C1q⁺ macrophages recovered from the small intestines of wild-type C57BL/6 mice. Median fluorescence intensities from multiple mice are quantified in the panels at right. Each data point represents one mouse (n=5-6 mice), and results are representative of two independent experiments. Error bars represent SEM. **p<0.01; ****p<0.0001 by two-tailed Student's t-test.
(E) Colonic longitudinal muscle myenteric plexus was acquired from $C1qa^{\Delta M\phi}$ and $C1qa^{fl/fl}$ littermates. RNA-seq was performed, and annotated gene ontology (GO) biological processes were assigned to genes that were differentially expressed in $C1qa^{\Delta M\phi}$ mice when compared to their $C1qa^{fl/fl}$ littermates. GO biological processes associated with neurons are in bold type. The dotted line indicates the cutoff for statistical significance. Five mice per group were analyzed as pooled biological replicates. Data are available from the Sequence Read Archive under BioProject ID PRJNA793870.
(F) Heat map of representative neuron-related genes differentially expressed in the colonic longitudinal muscle myenteric plexus of $C1qa^{\Delta M\phi}$ mice relative to $C1qa^{fl/fl}$ littermates. Five mice per group were analyzed as pooled biological replicates.

Figure supplement 1. Similar numbers of CD169⁺ macrophages are recovered from the small intestines of $C1qa^{\Delta M\phi}$ and $C1qa^{fl/fl}$ littermates.

Figure supplement 2. The colonic muscle myenteric plexus of $C1qa^{\Delta M\phi}$ mice has a transcriptional profile similar to that of mice with a gastrointestinal motility disorder.

$C1qa^{\Delta M\phi}$ mice have altered gastrointestinal motility

Macrophages engage in crosstalk with the enteric nervous system and regulate functions, including gastrointestinal motility, that depend on the enteric nervous system (Muller et al., 2014). This crosstalk involves the exchange of specific proteins such as bone morphogenetic protein 2 (BMP2) (Muller et al., 2014). Given that C1q⁺ macrophages phenotypically resemble peripheral nerve-adjacent macrophages and

reside near enteric neurons, we postulated that macrophage-derived C1q might also regulate enteric nervous

system function.

As an initial test of this idea, we performed RNA-seq on the colonic longitudinal muscle-myenteric

plexus from $C1qa^{\Delta M\phi}$ and $C1qa^{fl/fl}$ littermates and then conducted unbiased Gene Set Enrichment Analysis.

Of the 22 biological pathways that were enriched in $C1qa^{\Delta M\phi}$ mice, 18 were related to neuronal development

or function, including synapse organization, dendrite development, and neurotransmitter secretion (***Figure***

4E and F). 30 genes were differentially expressed when comparing the muscle-myenteric plexus of

$C1qa^{\Delta M\phi}$ and $C1qa^{fl/fl}$ mice. These included genes with known roles in regulating neuronal activity

(*Dusp26*), synaptic transmission (*Rasgrf2*), and neuropeptide signaling (*Tacr2*) (Mao et al., 2017;

Schwechter et al., 2013; Yang et al., 2017). We also compared the list of genes differentially expressed in

the $C1qa^{\Delta M\phi}$ mice to those differentially expressed in the *TashT* mouse line. This line contains an insertional

mutation that leads to a gut motility disorder comparable to Hirschsprung's disease, a human genetic

disorder resulting in incomplete development of the enteric nervous system (Bergeron et al., 2015). There

was a statistically significant overlap in the transcriptional changes in the colonic longitudinal muscle-

myenteric plexus of $C1qa^{\Delta M\phi}$ mice and *TashT* mice (***Figure 4 – figure supplement 2***). Together, these

results suggested that macrophage C1q impacts enteric nervous system gene expression and function.

Efficient coordination of gastrointestinal motility is necessary for proper digestion, nutrient

absorption, and excretion. Given that muscularis macrophages regulate enteric nervous system functions

that govern gastrointestinal motility (Muller et al., 2014), we assessed whether macrophage C1q impacts

gut motility. We first tested this idea by measuring gut transit time using the nonabsorbable dye carmine

red. $C1qa^{\Delta M\phi}$ and $C1qa^{fl/fl}$ littermates were gavaged with the dye and the time to first appearance of the dye

in the feces was recorded. Transit times were decreased in $C1qa^{\Delta M\phi}$ mice relative to their $C1qa^{fl/fl}$

littermates, indicating accelerated gut motility (***Figure 5A***). This was not due to a change in the lengths of

either the small intestine or the colon, which were unaltered in the $C1qa^{\Delta M\phi}$ mice (***Figure 5B***). By contrast,

gut transit time was unaltered in $C3^{-/-}$ mice, suggesting that macrophage C1q impacts gut motility

independent of its canonical function in the classical complement pathway (***Figure 5A***). Accelerated transit

was also observed in the small intestines of $C1qa^{\Delta M\phi}$ mice as assessed by rhodamine dye transit assay

(***Figure 5C***). We then conducted *ex vivo* colonic peristaltic recordings and observed that the colons of

$C1qa^{\Delta M\phi}$ mice exhibited greater neuromuscular activity as compared to $C1qa^{fl/fl}$ littermates (***Figure 5 –***

figure supplement 1). This suggested that the colons of $C1qa^{\Delta M\phi}$ mice maintained increased peristaltic

activity compared to their $C1qa^{fl/fl}$ littermates even in the absence of gut-extrinsic signals. Finally, we

directly assessed colonic motility by bead expulsion assay and found that $C1qa^{\Delta M\phi}$ mice had accelerated

colonic transit *in vivo* (***Figure 5D***). Thus, the absence of macrophage C1q increases peristaltic activity and

accelerates gut transit, indicating that macrophage C1q regulates gastrointestinal motility.

Figure 5. $C1qa^{\Delta M\phi}$ mice have altered gastrointestinal motility.
(A) Measurement of total intestinal transit time in $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ littermates and $C3^{-/-}$ mice. Mice were gavaged
with 100 µl of carmine red (5% (w/v in 1.5% methylcellulose). Fecal pellets were collected every 15 minutes and
transit time was recorded when the dye was first observed in the feces. Each data point represents one mouse and the
results are pooled from five independent experiments.
(B) Intestinal tract length is not altered in $C1qa^{\Delta M\phi}$ mice. Small intestines and colons from $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$
littermates were excised and measured. Each data point represents one mouse.
(C) Transit of rhodamine B-dextran through the intestines of $C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ littermates. Mice were sacrificed
90 minutes after gavage with rhodamine B-dextran. The intestines were divided into 16 segments, the rhodamine B
fluorescence was measured in each segment (top panel), and the geometric center of the fluorescence was determined
for each mouse (bottom panel). Each data point represents one mouse and the results were pooled from four
independent experiments.
(D) Colonic motility was measured by determining the expulsion time of a glass bead inserted intrarectally into
$C1qa^{fl/fl}$ and $C1qa^{\Delta M\phi}$ littermates. Each data point represents one mouse and the results are representative of three
independent experiments.
391

392 sm. int., small intestine. Error bars represent SEM. **p<0.01; ***p<0.001; ns, not significant by two-tailed Student's
393 t-test.
394
395 **Figure supplements 1 and 2.** *Ex vivo* video recordings of colonic peristalsis in *C1qa*$^{\text{fl/fl}}$ and *C1qa*$^{\Delta M\varphi}$ littermates.
396

DISCUSSION

Here, we have identified a role for C1q in regulating gastrointestinal motility. We discovered that

nerve-adjacent macrophages are the primary source of C1q in the mouse intestine and that macrophage C1q

regulates enteric neuronal gene expression and gastrointestinal transit time. Our findings reveal a previously

unappreciated function for C1q in the intestine and help to illuminate the molecular basis for macrophage-

mediated control of gut motility.

Our study identifies macrophages as the main source of C1q in the mouse small intestine and colon.

Both transient antibody-mediated depletion of macrophages and *in vivo* deletion of the *C1qa* gene from

macrophages led to a marked reduction in intestinal *C1q* expression. The *C1qa*$^{\Delta M\phi}$ mice also lacked C1q in

the circulation, indicating that LysM^{+} macrophages or macrophage-like cells are also the source of

circulating C1q in the absence of infection. This enhances findings from prior studies indicating that

monocytes, macrophages, and immature dendritic cells are the main sources of C1q in the bloodstream (El-

Shamy et al., 2018). Importantly, the *C1qa*$^{\Delta M\phi}$ mice retained C1q expression in the brain, allowing us to

analyze the effects of C1q deficiency without possible confounding effects on the central nervous system.

C1q has two known physiological functions that are distinct and vary according to tissue context.

C1q was originally discovered as having a role in the classical complement pathway, which tags and

destroys invading microbes (Noris and Remuzzi, 2013; Schifferli et al., 1986). Circulating C1q binds to

invading microorganisms and recruits additional proteins that assemble into the membrane attack complex

(MAC) (Kishore and Reid, 2000). C1q-mediated MAC formation has been primarily described in the

bloodstream, where the necessary accessory proteins are present at high levels (Davis et al., 1979).

However, C1q is also expressed in the absence of infection in tissues such as the brain, where it regulates

neuronal development and function (Kouser et al., 2015; van Schaarenburg et al., 2016).

Our findings suggest that C1q does not play a central role in immune defense of the intestine. First,

we found that intestinal C1q expression was not induced by gut commensals or pathogens and was not

deposited into the gut lumen. Second, C1q deficiency did not markedly alter gut microbiota composition or

the course of disease after DSS treatment. There were also no major changes in cytokine expression or

numbers and frequencies of intestinal immune cells that would indicate dysregulated interactions with the

microbiota. Third, C1q was not required for clearance of *C. rodentium*, a non-disseminating enteric

pathogen whose clearance requires antigen-specific IgG and complement component 3 (C3) (Belzer et al.,

2011). Although we cannot rule out a role for C1q in immune defense against other intestinal pathogens,

these findings support the idea that C1q is not essential for intestinal immune defense in mice.

Instead, our results indicate that C1q influences enteric nervous system function and regulates

intestinal motility. First, most C1q-expressing macrophages were localized to the intestinal submucosal

plexus and were largely absent from the lamina propria. Second, C1q-expressing intestinal macrophages

were located close to submucosal plexus neurons and expressed cell surface markers like those expressed

by nerve-adjacent C1q-expressing macrophages in the lung (Ural et al., 2020). Third, macrophage-specific

deletion of *C1qa* altered enteric neuronal gene expression. Finally, consistent with the altered neuronal gene

expression, macrophage-specific *C1qa* deletion altered gastrointestinal motility in both the small and large

intestine. Thus, our results suggest that the function of C1q in the intestine is similar to its function in the

brain, where it regulates the development and function of neurons (Kouser et al., 2015; van Schaarenburg

et al., 2016).

A function for macrophage C1q in intestinal motility adds to the growing understanding of how gut

macrophages regulate intestinal peristalsis. Prior work has shown that CSF1R[+] macrophages selectively

localize to the muscularis of the mouse intestine (Muller et al., 2014; Gabanyi et al., 2016). These

macrophages secrete BMP2, which activates enteric neurons that regulate colonic muscle contraction and

thus colonic motility (Muller et al., 2014). We found that depletion of CSF1R[+] macrophages reduces

intestinal C1q expression, and that macrophage-specific deletion of *C1qa* alters enteric neuronal gene

expression and gut motility. Thus, our findings suggest that C1q is a key component of the macrophage-

enteric nervous system axis.

An important remaining question concerns the molecular mechanism by which C1q regulates gut

motility. One possibility is that C1q shapes microbiota composition which, in turn, impacts gut motility.

This idea is suggested by studies in zebrafish showing that a deficiency in intestinal macrophages leads to

altered gut microbiota composition relative to wild-type zebrafish (Earley et al., 2018). Other studies in

zebrafish and mice have shown that severe defects in enteric nervous system development produce changes

in gut microbiota composition that are linked to dysregulated gut motility (Rolig et al., 2017; Johnson et

al., 2018). However, we did not observe marked changes in the composition of the gut microbiota in

$C1qa^{\Delta M\phi}$ mice, arguing against a central role for the microbiota in C1q-mediated regulation of gut motility.

A second possibility is that the absence of C1q leads to immunological defects that alter gut transit time.

This idea is suggested by studies showing that T cell cytokines can influence gastrointestinal motility

(Akiho et al., 2011). However, this seems unlikely given the lack of pronounced immunological

abnormalities in the intestines of $C1qa^{\Delta M\phi}$ mice.

A third possibility is that C1q interacts directly with enteric neurons or glial cells. Like

macrophage-produced BMP2 (Muller et al., 2014), C1q might bind to specific receptors on neurons to

regulate their activity. In support of this idea, mouse enteric neurons express *Adgrb1*, which encodes BAI1

(Obata et al., 2020), a known C1q receptor on human neural stem cells (Benavente et al., 2020). Other

studies have suggested that C1q functions as a signaling effector on immune cells and neurons (Benoit &

Tenner, 2011; Ling et al., 2018). Macrophages have also been reported to restrain neurogenesis in the

enteric nervous system through phagocytosis of apoptotic neurons (Kulkarni et al., 2017). Given that C1q

can opsonize dying host cells (Botto et al., 1998; Korb and Ahearn, 1997), C1q could regulate enteric

nervous system function by tagging apoptotic neurons for elimination. It is also possible that C1q acts

directly on enteric smooth muscle cells that regulate gut motility. Although we cannot rule out this

possibility, our transcriptional profile of the colonic myenteric plexus of $C1qa^{\Delta M\phi}$ mice suggests that most

of the transcriptional changes were associated with neurons.

Our findings on intestinal C1q have implications for the understanding of human intestinal disease.

Dysregulated peristalsis is a characteristic of irritable bowel syndrome (Vrees et al., 2002) and is present in

a subset of inflammatory bowel disease patients (Bassotti et al., 2014). Our finding that macrophage C1q

regulates gut motility could suggest new strategies to prevent or treat these diseases. Additionally, most

humans with C1q deficiency develop systemic lupus erythematosus (SLE). Since C1q can target cellular

debris for phagocytosis, it is thought that the absence of C1q results in increased exposure of self-antigen

to the immune system, thereby reducing immune tolerance and causing autoimmune disease (Macedo and

Isaac, 2016). Furthermore, roughly 42.5% of SLE patients report gastrointestinal symptoms that range from

acute abdominal pain to chronic intestinal obstruction (Fawzy et al., 2016; Tian and Zhang, 2010). The

exact cause of these symptoms is unclear. Given that C1q deficiency is strongly correlated with SLE in

humans and alters gut motility in mice, we suggest that C1q could be a therapeutic target for SLE patients

that present with chronic constipation or other forms of dysregulated intestinal motility.

484
485 **MATERIALS AND METHODS**
486 **Key Resources Table**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background (*Mus musculus*)	$C1qa^{fl/fl}$; B6(SJL)-C1qa$^{tm1c(EUCOMM)Wtsi}$/TennJ	Jackson Laboratory; Fonseca et al., 2017	Stock #031261	
strain, strain background (*Mus musculus*)	LysM-Cre; B6.129P2-Lyz2$^{tm1(cre)Ifo}$/J	Jackson Laboratory; (Clausen et al., 1999)	Stock #004781	
strain, strain background (*Mus musculus*)	$C1qa^{\Delta M\phi}$	this paper		Generated by crossing $C1qa^{fl/fl}$ mice with LysM-Cre mice
strain, strain background (*Mus musculus*)	$C3^{-/-}$; B6.129S4-C3^{tm1Crr}/J	Jackson Laboratory; Wessels et al., 1995	Stock #029661	
strain, strain background (*Mus musculus*)	Germ-free C57BL/6J mice	UT Southwestern Gnotobiotics Core Facility		

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background (*Salmonella enterica*)	*Salmonella enterica* subsp. enterica serovar Typhimurium strain SL1344	Dr. Vanessa Sperandio; Eichelberg and Galan, 1999		
strain, strain background (*Citrobacter rodentium*)	*Citrobacter rodentium* strain DBS100	ATCC	Strain# 51459	
antibody	Anti-Actin HRP (rabbit monoclonal)	Cell Signaling	Clone: 13E5	Immunoblot (1:5000)
antibody	Anti-ARG1 (sheep monoclonal)	R&D Systems	Clone: P05089	Flow (1:100)
antibody	Anti-B220 (rat monoclonal)	ThermoFisher	Clone: RA3-6B2	Flow (1:500)
antibody	Anti-C1q (rat monoclonal)	Cedarlane Laboratories	Clone: RmC7H8	Flow (1:50)
antibody	Anti-C1q (rabbit polyclonal)	ThermoFisher	Cat# PA5-29586	Immunoblot (1:500)
antibody	Anti-C1q-biotin (mouse monoclonal)	Abcam	Clone: JL1	ELISA (1:1000); Immunofluorescence (1:100)
antibody	Anti-CD3 (rat monoclonal)	ThermoFisher	Clone: 17A2	Flow (1:200)
antibody	Anti-CD4 (rat monoclonal)	BioLegend	Clone: GK1.5	Flow (1:500)
antibody	Anti-CD11b (rat monoclonal)	ThermoFisher	Clone: M1/70	Flow (1:200)
antibody	Anti-CD11c (Armenian hamster monoclonal)	ThermoFisher	Clone: N418	Flow (1:500)
antibody	Anti-CD16/32 (rat monoclonal)	BioLegend	Clone: 93	Fc receptor block (1:1000)
antibody	Anti-CD19 (rat monoclonal)	BioLegend	Clone: 1D3	Flow (1:500)
antibody	Anti-CD45 (rat monoclonal)	BioLegend	Clone: 30-F11	Flow (1:500)
antibody	Anti-CD90.2 (rat monoclonal)	BioLegend	Clone: 30-H12	Flow (1:500)
antibody	Anti-CD169 (rat monoclonal)	BioLegend	Clone: 3D6.112	Flow (1:200)
antibody	Anti-CD169 (rat monoclonal)	Abcam	Clone: 3D6.112	Immunofluorescence (1:200)
antibody	Anti-CSF1R (rat monoclonal)	Bio X Cell	Cat# AFS98	Macrophage depletion (dosage)
antibody	Anti-F4/80	BioLegend	Clone: BM8	Flow (1:100)
antibody	Anti-FoxP3 (rat monoclonal)	ThermoFisher	Clone: FJK-16s	Flow (1:50)
antibody	Anti-GATA3 (mouse monoclonal)	BD Biosciences	Clone: L50-823	Flow (1:50)
antibody	Anti-IgA (rat monoclonal)	ThermoFisher	Clone: 11-44-2	Flow (1:50)
antibody	Anti-LY6C	BioLegend	Clone: RB6-8C5	Flow (1:500)
antibody	Anti-MHCII	Thermo	Clone: M5/114.15.2	Flow (1:500)
antibody	Anti-REG3G antiserum (rabbit)	Cash et al., 2006; antiserum generated by Pacific Biosciences		Immunoblot (dilution)
antibody	Anti-RORγt	ThermoFisher	Clone: AFKJS-9	Flow (1:50)
antibody	Anti-T-BET	BioLegend	Clone: 4B10	Flow (1:50)
antibody	Anti-TREM2	R&D Systems	Clone: 237920	Flow (1:200)
antibody	Anti-TUBB3	Abcam	Cat# ab18207	Immunofluorescence (1:200)
antibody	Goat anti-rabbit IgG HRP conjugate	Abcam	Cat# ab6721	Immunoblot (1:5000)
antibody	secondary antibodies - AlexaFluor 488/594/647	Invitrogen		Immunofluorescence (1:400)
antibody	mouse IgG1	Abcam	Cat# ab18443	ELISA (10 μg/ml)
antibody	Rat IgG2a	ThermoFisher	Clone: 2A3	Isotype control for anti-CSF1R macrophage depletion
antibody	Rat IgG1 PE isotype control	Cedarlane Laboratories	Cat# CLCR104	Flow (1:50)
sequence-based reagent	mouse *C1qa* TaqMan assay	ThermoFisher	Assay ID: Mm00432142_m1	

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
sequence-based reagent	mouse *C1qb* TaqMan assay	ThermoFisher	Assay ID: Mm01179619_m1	
sequence-based reagent	mouse *C1qc* TaqMan assay	ThermoFisher	Assay ID: Mm00776126_m1	
sequence-based reagent	mouse *Reg3g* TaqMan assay	ThermoFisher	Assay ID: Mm00441127_m1	
sequence-based reagent	mouse *Ifng* TaqMan assay	ThermoFisher	Assay ID: Mm01168134_m1	
sequence-based reagent	mouse *Il4* TaqMan assay	ThermoFisher	Assay ID: Mm00445259_m1	
sequence-based reagent	mouse *IL5* TaqMan assay	ThermoFisher	Assay ID: Mm00439646_m1	
sequence-based reagent	mouse *Il10* TaqMan assay	ThermoFisher	Assay ID: Mm01288386_m1	
sequence-based reagent	mouse *Il13* TaqMan assay	ThermoFisher	Assay ID: Mm00434204_m1	
sequence-based reagent	mouse *Il17a* TaqMan assay	ThermoFisher	Assay ID: Mm00439618_m1	
sequence-based reagent	mouse *Il17f* TaqMan assay	ThermoFisher	Assay ID: Mm00521423_m1	
sequence-based reagent	mouse 18*S* gene TaqMan assay	ThermoFisher	Assay ID: Mm03928990_g1	
sequence-based reagent	bacterial 16*S* V3 - rRNA gene forward primer	ThermoFisher; (Klindworth et al., 2013)	16*S* rRNA gene sequencing	5'-TCGTCGGCAGCGTC AGATGTGTA TAAGAGACAGCCTAC GGGNGGCWGCAG-3′
sequence-based reagent	bacterial 16*S* v4 - rRNA gene reverse primer	ThermoFisher; Klindworth et al., 2013	16*S* rRNA gene sequencing	5′-GTCTCGTGGGCTCG GAGATGTGTA TAAGAGACAGGACTA CHVGGGTATCTAATC C-3′
peptide, recombinant protein	recombinant mouse C1q	Complementech	Cat# M099	
commercial assay or kit	FOXP3/Transcription Factor Fixation/Permeabilization Buffer Set	ThermoFisher	Cat# 00-5523-00	
commercial assay or kit	MMLV Reverse Transcriptase Kit	ThermoFisher	Cat# 28025-021	
commercial assay or kit	PE300 (Paired end 300 bp) v3 kit	Illumina	Cat# MS-102-3001	
commercial assay or kit	RNeasy Universal Mini Kit	Qiagen	Cat# 73404	
commercial assay or kit	TaqMan Master Mix	ThermoFisher	Cat# 4369542	
commercial assay or kit	TruSeq RNA sample preparation kit	Illumina	Cat# RS-122-2001	
chemical compound, drug	Agencourt AmpureXP beads	Beckman Coulter Genomics	Cat# A63880	
chemical compound, drug	Carmine Red	Sigma	Cat# C1022-25G	
chemical compound, drug	Collagenase IV	Sigma	Cat# C5138-1G	
chemical compound, drug	Borosilicate glass beads (2mm)	Millipore Sigma	Cat# Z273627-1EA	
chemical compound, drug	Dextran sulfate sodium	Thomas Scientific	Cat# 216011090	
chemical compound, drug	DNase I	Sigma	Cat# DN25	
chemical compound, drug	Dispase II	Sigma	Cat# D4693-1G	
chemical compound, drug	FITC-dextran	Sigma	Cat# FD4-1g	
chemical compound, drug	Ghost 710	Tonbo Biosciences	Cat# 13-0871-T100	Flow cytometry viability dye
chemical compound, drug	Methylcellulose	Sigma	Cat# M0262-100G	
chemical compound, drug	Nalidixic acid, sodium salt	Research Products International	Cat# N23100-25.0	
chemical compound, drug	Optimal Cutting Temperature Compound (OCT)	ThermoFisher	Cat# 23-730-571	
chemical compound, drug	Percoll Plus	GE Healthcare	Cat# GE17-0891-09	

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
chemical compound, drug	Protease inhibitors	Millipore Sigma	Cat# 11836153001	
chemical compound, drug	Rhodamine B-dextran	ThermoFisher	Cat# D1841	
chemical compound, drug	Streptavidin-Cy5	ThermoFisher	Cat# 434316	
chemical compound, drug	Streptavidin-HRP conjugate	Abcam	Cat# ab7403	ELISA
chemical compound, drug	VECTASHIELD Antifade Mounting Medium with DAPI	Vector Labs	Cat# H-1200-10	
software, algorithm	clusterProfiler			
software, algorithm	CLC Genomics Workbench			
software, algorithm	CLC Bio microbial genomics module			https://digitalinsights.qiagen.com/plugins/clc-microbial-genomics-module/
software, algorithm	FlowJo			
software, algorithm	GraphPad PRISM	GraphPad Software	Version 7.0; RRID:SCR_002798	
software, algorithm	Igor Pro 9	WaveMetrics		
software, algorithm	Illumina Nextera Protocol	Illumina	Part # 15044223 Rev. B	
software, algorithm	ImageJ	National Institutes of Health		https://imagej.nih.gov/ij/
software, algorithm	NovoExpress	Agilent Technologies		
software, algorithm	PVCAM software	Teledyne Photometrics		
Other	Agilent 2100 Bioanalyzer	Agilent Technologies	G2939A	
Other	Amicon Ultra centrifugal filters	Millipore	Cat #UFC900324	
Other	BioRad ChemiDoc Touch System	BioRad	Cat# 1708370	
Other	CMOS camera	Teledyne Photometrics	MOMENT	
Other	Leica CM1950 (Cryostat)	Leica		
Other	FACSAria	BD Biosciences		
Other	Illumina MiSeq	Illumina	RRID:SCR_016379	
Other	Illumina NextSeq 500	Illumina		
Other	Keyence Fluorescence Microscope	Keyence	BZ-X800	
Other	NovoCyte 3005	Agilent Technologies		
Other	Organ bath chamber	Tokai Hit Co. (Japan)		
Other	Peristaltic pump	Gilson	MINIPULS3	
Other	QuantStudio 7 Flex Real-Time PCR System	Applied Biosystems	Cat #4485701	
Other	SpectraMax M5 plate reader	Molecular Devices		
Other	Zeiss Axio Imager M1 Microscope	Zeiss		

Mice

Wild-type C57BL/6J (Jackson Laboratory) and $C3^{-/-}$ mice (Jackson Laboratory; Wessels et al., 1995) were

bred and maintained in the SPF barrier facility at the University of Texas Southwestern Medical Center.

$C1qa^{\Delta M\phi}$ mice were generated by crossing $C1qa^{fl/fl}$ mice (Jackson Laboratory; Fonseca et al., 2017) with a

mouse expressing Cre recombinase controlled by the macrophage-specific mouse *Lyz2* promoter (LysM-

Cre mice; Jackson Laboratory; Clausen et al., 1999). Mice 8 to 12 weeks of age were used for all

experiments and cohoused littermates were used as controls. Both male and female mice were analyzed.

Germ-free C57BL/6J mice were bred and maintained in isolators at the University of Texas Southwestern

Medical Center. All procedures were performed in accordance with protocols approved by the Institutional

Animal Care and Use Committees (IACUC) of the UT Southwestern Medical Center.

Quantitative polymerase chain reaction (qPCR)

Tissue RNA was isolated using the RNeasy Universal Mini kit (Qiagen, Hilden, Germany). Cellular RNA

was isolated using the RNAqueous Micro kit (ThermoFisher). cDNA was generated from the purified RNA

using the M-MLV Reverse Transcriptase kit (ThermoFisher). qPCR analysis was performed using TaqMan

primer/probe sets and master mix (ThermoFisher) on a Quant-Studio 7 Flex Real-Time PCR System

(Applied Biosystems). Transcript abundances were normalized to 18S rRNA abundance. TaqMan probe

assay IDs are provided in the Key Resources table.

Isolation and analysis of intestinal immune cells

Lamina propria (LP) cells were isolated from the intestine using a published protocol (Yu et al., 2013; Yu

et al., 2014). Briefly, intestines were dissected from mice and Peyer's patches were removed. Intestines

were cut into small pieces and thoroughly washed with ice-cold phosphate-buffered saline (PBS) containing

5% fetal bovine serum (PBS-FBS). Epithelial cells were removed by incubating intestinal tissues in Hank's

buffered salt solution (HBSS) supplemented with 2 mM EDTA, followed by extensive washing with PBS-

FBS. Residual tissues were digested twice with Collagenase IV (Sigma), DNase I (Sigma) and Dispase (BD

Biosciences) for 45 minutes at 37°C with agitation. Cells were filtered through 70 μm cell strainers

(ThermoFisher) and applied onto a 40%:80% Percoll gradient (GE Healthcare). Subepithelial cell

populations were recovered at the interface of the 40% and 80% fractions. For small intestinal cell

suspensions, the epithelial fraction was kept and combined with enzymatically liberated subepithelial cells.

Cells were washed with 2 mM EDTA/3% FBS in PBS and Fc receptors were blocked with anti-CD16/32

(93). Cells were then stained with the viability dye Ghost 710 (Tonbo Biosciences) followed by antibodies

against cell surface markers including anti-CD45 (30-F11), anti-CD11b (M1/70), anti-MHCII

(M5/114.15.2), anti-F4/80 (BM8), anti-CD3 (17A2), anti-CD4 (GK1.5), anti-CD19 (1D3), anti-B220

(RA3-6B2), anti-CD11c (N418), anti-CD169 (3D6.112), anti-TREM2 (237920) and anti-LY6C (RB6-

8C5). Cells were fixed and permeabilized with the eBioscience FOXP3/Transcription Factor

Fixation/Permeabilization buffer set (ThermoFisher) and then subjected to intracellular staining with anti-

C1Q (RmC7H8), anti-FOXP3 (FJK-16s), anti-GATA3 (L50), anti-T-BET (4B10), anti-RORγ (AFKJS-9),

and anti-ARG1 (P05089). Cells were sorted using a FACSAria (BD Biosciences) or analyzed using a

NovoCyte 3005 (Agilent Technologies). Data were processed with FlowJo software (BD Biosciences) or

NovoExpress (Agilent Technologies).

Macrophage depletion

Anti-mouse CSF1R (ThermoFisher; AFS98) and rat IgG2a isotype control (ThermoFisher; 2A3) antibodies

were administered intraperitoneally at a concentration of 100 mg/kg. Mice were sacrificed 72 hours post-

injection and terminal ileum and colon were collected for qPCR analysis.

Protein extraction from intestinal cells and feces

To isolate proteins from intestinal cell suspensions, cell pellets were resuspended in 100 μl of RIPA Lysis

Buffer (ThermoFisher) supplemented with protease inhibitors (Millipore Sigma) and vortexed vigorously

every 5 minutes for 20 minutes. Lysates were cleared of cellular debris by centrifugation at 13000g for 5

minutes. To isolate proteins from the intestinal lumen, the entire gastrointestinal tract (from duodenum to

distal colon) was recovered from five wild-type C57BL/6J mice. The intestines were flushed with a total of

~ 50 ml cold PBS containing protease inhibitors (Millipore Sigma, 11836153001). The flushes and fecal

pellets were homogenized by rotor and stator (TH Tissue Homogenizer; OMNI; TH01) and large particles

were centrifuged at 100g for 10 min at room temperature. The supernatants were carefully decanted and

centrifuged further at 3000*g* for 20 min at room temperature. The clarified supernatants were precipitated

with 40% ammonium sulfate overnight at 4°C. Precipitated protein was centrifuged at 3000*g* for 30 min at

4°C, then resuspended in cold 40% ammonium sulfate and centrifuged again. The pellets were resuspended

in room temperature PBS and allowed to mix for 10 min. Protein concentrations were determined by

Bradford assay (BioRad).

Immunoblot

50 µg of fecal protein or 25 µg of cellular protein was loaded onto a 4-20% gradient SDS-PAGE and

transferred to a PVDF membrane. Membranes were blocked in 5% nonfat dry milk in Tris-buffered saline

(TBS) with 0.1% Tween-20 and then incubated overnight with the following primary antibodies: anti-C1Q

(PA5-29586, ThermoFisher) and anti-actin (13E5, Cell Signaling). REG3G was detected by incubating

membranes with rabbit anti-REG3G antiserum (Cash et al., 2006). After washing, membranes were

incubated with goat anti-rabbit IgG HRP and then visualized with a BioRad ChemiDoc Touch system.

Enzyme-linked immunosorbent assay (ELISA)

Mouse C1q ELISA was performed as previously described (Petry et al., 2001). Briefly, microtiter plates

were coated overnight with mouse IgG1 and were then blocked with 5% BSA in PBS. Serum samples were

diluted 1:50 and plated for 1 hour at room temperature. After washing with 0.05% Tween-20 in PBS, bound

C1q was incubated with biotinylated anti-C1q antibody (JL1, Abcam). Biotinylated anti-C1q was detected

with a streptavidin-HRP conjugate (Abcam). Optical density was measured using a wavelength of 492 nm.

Plates were then analyzed using a SpectraMax M5 microplate reader (Molecular Devices).

Intestinal permeability assay

Intestinal permeability assays were performed by treating mice with fluorescein isothiocyanate dextran

(FITC-dextran) by oral gavage. The non-steroidal anti-inflammatory drug (NSAID) indomethacin was

administered to mice as a positive control. For the experimental group, mice were treated with 190 µl 7%

dimethyl sulfoxide (DMSO) in PBS by oral gavage. For the positive control group, mice were treated with

190 µl indomethacin (1.5 mg/ml in 7% DMSO in PBS) by oral gavage. After 1 hour, all mice were treated

with 190 ul FITC-dextran (80 mg/ml in PBS) by oral gavage. Mice were sacrificed after 4 hours and sera

were collected. Serum samples were centrifuged for 20 minutes at 4°C at 800g and supernatants were

collected. Serum FITC-dextran levels were measured by a fluorescence microplate assay against a standard

curve using a Spectramax plate reader (Molecular Devices).

Dextran sulfate sodium (DSS) treatment

Age and sex-matched mice were provided with 3% dextran sulfate sodium (weight/volume) in autoclaved

drinking water for 7 days. Animal weight and health were monitored in accordance with institutional

IACUC guidelines.

***Salmonella* Typhimurium infection**

To prepare bacteria for infection, *Salmonella enterica* serovar Typhimurium (SL1344) was cultured in

Luria-Bertani (LB) broth containing 50 µg/ml streptomycin in a shaking incubator at 37°C (Eichelberg and

Galan, 1999). The overnight culture was diluted the next day and grown to mid-log phase (OD_{600} = 0.3-

0.5). *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\phi}$ littermates were inoculated intragastrically with 10^9 CFU. All mice were

sacrificed 24 hours later and small intestinal tissues were harvested for analysis.

***Citrobacter rodentium* infection**

To prepare bacteria for infection, an overnight culture of *C. rodentium* (DBS100, ATCC) in LB broth

containing nalidixic acid (100 µg/ml) in a shaking incubator at 37°C. The overnight culture was diluted the

next day and grown to mid-log phase (OD_{600} = 0.4-0.6). Bacteria were pelleted, washed, and resuspended

in PBS. Sex-matched littermates were inoculated intragastrically with 5×10^8 CFU. Fecal pellets were

collected at a fixed time every 48 hours, homogenized in sterile PBS, diluted, and plated on LB agar with

nalidixic acid (100 μg/ml).

Immunofluorescence analysis of mouse intestines

Mouse small intestines and colons were flushed with PBS and embedded with Optimal Cutting Temperature

compound (OCT) (ThermoFisher). Sections were fixed in ice cold acetone, blocked with 1% BSA, 10%

FBS, 1% Triton X-100 in PBS, and then incubated overnight at 4°C with the following antibodies: mouse

anti-C1q biotin (JL-1), rat anti-CD169 (3D6.112), and rabbit anti-TUBB3 (ab18207, Abcam). Slides were

then washed with PBS containing 0.2% Tween-20 (PBS-T) and incubated with donkey anti-rabbit

AlexaFluor 488, donkey anti-rat AlexaFluor 594, and Streptavidin-Cy5 (ThermoFisher) for 1 hour at room

temperature in the dark. Slides were then washed in PBS-T and mounted with DAPI-Fluoromount-G

(Southern Biotech). Mounted slides were cured overnight at 4°C until imaging.

RNA-seq analysis of colonic longitudinal muscle myenteric plexus

The colonic longitudinal muscle-myenteric plexus was collected from five age and sex matched $C1qa^{fl/fl}$

and $C1qa^{\Delta M\phi}$ mice by manual dissection using a 2 mm metal probe (Fisher Scientific). RNA was isolated

using the RNeasy Mini kit according to the manufacturer's protocol (Qiagen). Quantity and quality of RNA

samples was assessed on an Agilent 2100 Bioanalyzer (Agilent Technologies). RNA sequencing libraries

were prepared using the TruSeq RNA sample preparation kit (Illumina) according to the manufacturer's

protocol. Libraries were validated on an Agilent Bionalyzer 2100. Indexed libraries were sequenced on an

Illumina NextSeq550 for single-end 75 bp length reads. CLC Genomics Workbench 7 was used for

bioinformatics and statistical analysis of the sequencing data. The approach used by CLC Genomics

Workbench is based on a method developed previously (Mortazavi et al., 2008). To identify differentially

enriched biological pathways, all genes were ranked based on their $\log_2$ fold change and pathway

enrichment was identified using the R packages "clusterProfiler" and "msigdbr". For analysis of

differentially expressed genes, gene counts were analyzed using DESeq-2 and differentially expressed

genes were defined as having an adjusted p value <0.05. A Fisher's Exact Test was conducted to assess the

overlap between differentially expressed genes in $C1qa^{\Delta M\phi}$ mice and the TashT mouse (Bergeron et al.,

2015).

16S rRNA gene sequencing and analysis

The hypervariable regions V3 and V4 of the bacterial 16S rRNA gene were prepared using the Illumina

Nextera protocol (Part # 15044223 Rev. B). An amplicon of 460 bp was amplified using the 16S Forward

Primer and 16S Reverse Primer as described in the manufacturer's protocol. Primer sequences are given in

the Key Resources Table. The PCR product was purified using Agencourt AmpureXP beads (Beckman

Coulter Genomics). Illumina adapter and barcode sequences were ligated to the amplicon to attach them to

the MiSeqDx flow cell and for multiplexing. Quality and quantity of each sequencing library were assessed

using Bioanalyzer and picogreen measurements, respectively. Libraries were loaded onto a MiSeqDX flow

cell and sequenced using the Paired End 300 (PE300) v3 kit. Raw fastq files were demultiplexed based on

unique barcodes and assessed for quality. Samples with more than 50,000 quality control pass sequencing

reads were used for downstream analysis. Taxonomic classification and operational taxonomic unit analysis

were done using the CLC Microbial Genomics Module. Individual sample reads were annotated with the

Greengene database and taxonomic features were assessed.

Gastrointestinal motility assays

Motility assays were adapted from previous studies (Luo et al., 2018; Maurer, 2016; Muller et al., 2014).

To determine transit time through the entire gastrointestinal tract, mice were fasted overnight and water

was removed 1 hour prior to the start of the experiment. Mice were then singly housed for 1 hour and then

gavaged with 100 μl of carmine red (5% weight/volume; Sigma) in 1.5% methylcellulose. Fecal pellets

were collected every 15 minutes and transit time was recorded when the dye was first observed in the feces.

For small intestinal motility measurements, mice were fasted overnight and then gavaged with 100

μl of rhodamine B-dextran (5 mg/ml; ThermoFisher) in 2% methylcellulose. After 90 minutes, mice were

sacrificed and their stomachs, small intestines, ceca, and colons were collected. Small intestines were cut

into 8 segments of equal length and colons were cut into 5 segments of equal length. Segments were cut

open lengthwise and vortexed in 1 ml PBS to release rhodamine B-dextran. Fluorescence was then

measured on a SpectraMax M5 microplate reader (Molecular Devices). The geometric center of the dye

was calculated as: GC = Σ (% of total fluorescent signal per segment × segment number).

Relative fluorescence per segment was calculated as: (fluorescence signal in segment / total fluorescence

recovered) x 100.

To measure colonic motility, mice were fasted overnight and lightly anesthetized with isoflurane.

A 2 mm glass bead was inserted 2 cm intrarectally using a 2 mm surgical probe. Mice were then returned

to empty cages and time to reappearance of the bead was recorded.

To account for potential circadian differences in gut motility, the time of day for the initiation of

all experiments was held constant.

Ex vivo **peristaltic imaging**

Ex vivo video imaging and analysis of colonic peristalsis was carried out as described previously (Obata et

al., 2020). Colons were dissected, flushed with sterile PBS, and pinned into an organ bath chamber (Tokai

Hit, Japan) filled with high glucose Dulbecco's Modified Eagle Medium (DMEM). DMEM was oxygenated

(95% O2 and 5% CO_2), run through the chamber using a peristaltic pump (MINIPULS 3, Gilson) and kept

at 37°C. Colons were allowed to equilibrate to the organ chamber for 30 minutes before video recording.

Movies of colonic peristalsis were captured with a camera (MOMENT, Teledyne photometrics) using

PVCAM software (500ms time-lapse delay) over 15 minutes.

Statistical analysis

Graphed data are presented as means ± standard error of the mean (SEM). Statistics were determined with GraphPad Prism software. Statistical analyses were performed using two-tailed Student's t test when comparing two groups, one way ANOVA when comparing multiple groups, and Fisher's exact test to assess overlap between groups of differentially expressed genes. The statistical tests used are indicated in the figure legends. $*P \leq 0.05$; $**P \leq 0.01$; $***P \leq 0.001$; and ns, $P > 0.05$.

ADDITIONAL INFORMATION

Acknowledgements

We thank Shai Bel for assistance with immunofluorescence imaging experiments, the UT Southwestern Genomics Core for assistance with RNA sequencing experiments, and the UT Southwestern Flow Cytometry Core for assistance with flow cytometry experiments. *Citrobacter rodentium* strain DBS100 was a gift from Vanessa Sperandio. This work was supported by NIH grants R01 DK070855 (L.V.H.), Welch Foundation Grant I-1874 (L.V.H.), the Walter M. and Helen D. Bader Center for Research on Arthritis and Autoimmune Diseases (L.V.H.), and the Howard Hughes Medical Institute (L.V.H.). M.P. and A.A.C. were supported by NIH T32 AI005284. E.K. was supported by NIH F31 DK126391. Y.O. is the Nancy Cain Marcus and Jeffrey A. Marcus Scholar in Medical Research, in Honor of Dr. Bill S. Vowell.

Competing interests: The authors declare no competing interests.

Author contributions

Mihir Pendse, Conceptualization, Investigation, Data curation, Formal analysis, Methodology, Supervision, Writing – original draft, Writing – review and editing; Yun Li, Investigation; Cristine N. Salinas, Investigation; Gabriella Quinn, Data curation, Formal analysis; Daniel C. Propheter, Investigation, Writing – review and editing; Chaitanya Dende, Investigation, Writing – review and editing; Alexander A. Crofts,

Data curation, Formal analysis; Eugene Koo, Methodology; Brian Hassell, Investigation; Kelly A. Ruhn,

Investigation; Prithvi Raj, Investigation, Data curation, Formal analysis; Yuuki Obata, Investigation,

Methodology, Writing – original draft, Writing – review and editing; Lora V. Hooper, Conceptualization,

Funding acquisition, Supervision, Writing – original draft, Writing – review and editing.

Data availability

16*S* rRNA gene sequencing data (***Figure 3D***) and RNA sequencing data (***Figure 4E and F; Figure 4 –***

figure supplement 2) are available from the Sequence Read Archive under BioProject ID PRJNA793870.

All mouse strains used are available commercially.

705
706
707
708   **REFERENCES**
709
710

711   Akiho, H., Ihara, E., Motomura, Y., Nakamura, K. (2011). Cytokine-induced alterations of gastrointestinal
712   motility in gastrointestinal disorders. *World J Gastrointest Pathophysiol, 2*, 72-81.
713   https://doi.org/10.4291/wjgp.v2.i5.72
714

715   Bassotti, G., Antonelli, E., Villanacci, V., Salemme, M., Coppola, M., Annese, V. (2014). Gastrointestinal
716   motility disorders in inflammatory bowel diseases. *World J Gastroenterol, 20*, 37-44.
717   https://doi.org/10.3748/wjg.v20.i1.37
718

719   Belzer, C., Liu, Q., Carroll, M. C., Bry, L. (2011). The role of specific IgG and complement in combating
720   a primary mucosal infection of the gut epithelium. *Eur J Microbiol Immunol (Bp), 1*, 311-318.
721   https://doi.org/10.1556/EuJMI.1.2011.4.7
722

723   Benavente, F., Piltti, K.M., Hooshmand, M.J., Nava, A.A., Lakatos, A., Feld, B.G., Creasman, D., Gershon,
724   P.D., Anderson, A. (2020). Novel C1q receptor-mediated signaling controls neural stem cell behavior and
725   neurorepair. *eLife, 9*, e55732. https://doi.org/10.7554/eLife.55732
726

727   Benoit, M. E., Tenner, A. J. (2011). Complement protein C1q-mediated neuroprotection is correlated with
728   regulation of neuronal gene and microRNA expression. *J Neurosci, 31*, 3459-3469.
729   https://doi.org/10.1523/JNEUROSCI.3932-10.2011
730

731   Bergeron, K. F., Cardinal, T., Toure, A. M., Beland, M., Raiwet, D. L., Silversides, D. W., Pilon, N. (2015).
732   Male-biased aganglionic megacolon in the TashT mouse line due to perturbation of silencer elements in a
733   large gene desert of chromosome 10. *PLoS Genet, 11*, e1005093.
734   https://doi.org/10.1371/journal.pgen.1005093
735

Bogunovic, M., Ginhoux, F., Helft, J., Shang, L., Hashimoto, D., Greter, M., Liu, K., Jakubzick, C., Ingersoll, M. A., Leboeuf, M., Stanley, E. R., Nussenzweig, M., Lira, S. A., Randolph, G. J., Merad, M. (2009). Origin of the lamina propria dendritic cell network. *Immunity, 31*, 513-525. https://doi.org/10.1016/j.immuni.2009.08.010

Bossi, F., Tripodo, C., Rizzi, L., Bulla, R., Agostinis, C., Guarnotta, C., Munaut, C., Baldassarre, G., Papa, G., Zorzet, S., Ghebrehiwet, B., Ling, G. S., Botto, M., Tedesco, F. (2014). C1q as a unique player in angiogenesis with therapeutic implication in wound healing. *Proc Natl Acad Sci U S A, 111*, 4209-4214. https://doi.org/10.1073/pnas.1311968111

Botto, M., Dell'Agnola, C., Bygrave, A. E., Thompson, E. M., Cook, H. T., Petry, F., Loos, M., Pandolfi, P. P., Walport, M. J. (1998). Homozygous C1q deficiency causes glomerulonephritis associated with multiple apoptotic bodies. *Nat Genet, 19*, 56-59. https://doi.org/10.1038/ng0598-56

Cao, W., Bobryshev, Y. V., Lord, R. S., Oakley, R. E., Lee, S. H., Lu, J. (2003). Dendritic cells in the arterial wall express C1q: potential significance in atherogenesis. *Cardiovasc Res, 60*, 175-186. https://doi.org/10.1016/s0008-6363(03)00345-6

Casals, C., Garcia-Fojeda, B., Minutti, C. M. (2019). Soluble defense collagens: Sweeping up immune threats. *Mol Immunol, 112*, 291-304. https://doi.org/10.1016/j.molimm.2019.06.007

Cash, H. L., Whitham, C. V., Behrendt, C. L., Hooper, L. V. (2006). Symbiotic bacteria direct expression of an intestinal bactericidal lectin. *Science, 313*, 1126-1130. https://doi.org/10.1126/science.1127119

Chu, Y., Jin, X., Parada, I., Pesic, A., Stevens, B., Barres, B., Prince, D. A. (2010). Enhanced synaptic connectivity and epilepsy in C1q knockout mice. *Proc Natl Acad Sci U S A, 107*, 7975-7980. https://doi.org/10.1073/pnas.0913449107

Clausen, B. E., Burkhardt, C., Reith, W., Renkawitz, R., Forster, I. (1999). Conditional gene targeting in macrophages and granulocytes using LysMcre mice. *Transgenic Res, 8*, 265-277. https://doi.org/10.1023/a:1008942828960

Colonna, M. (2003). TREMs in the immune system and beyond. *Nat Rev Immunol, 3*, 445-453. https://doi.org/10.1038/nri1106

Davis, C. A., Vallota, E. H., Forristal, J. (1979). Serum complement levels in infancy: age related changes. *Pediatr Res, 13*, 1043-1046. https://doi.org/10.1203/00006450-197909000-00019

De Schepper, S., Stakenborg, N., Matteoli, G., Verheijden, S., Boeckxstaens, G. E. (2018). Muscularis macrophages: Key players in intestinal homeostasis and disease. *Cell Immunol, 330*, 142-150. https://doi.org/10.1016/j.cellimm.2017.12.009

De Schepper, S., Verheijden, S., Aguilera-Lizarraga, J., Viola, M. F., Boesmans, W., Stakenborg, N., Voytyuk, I., Schmidt, I., Boeckx, B., Dierckx de Casterle, I., Baekelandt, V., Gonzalez Dominguez, E., Mack, M., Depoortere, I., De Strooper, B., Sprangers, B., Himmelreich, U., Soenen, S., Guilliams, M., Vanden Berghe, P., Jones, E., Lambrechts, D., Boeckxstaens, G. (2018, Oct 4). Self-maintaining gut macrophages are essential for intestinal homeostasis. *Cell, 175*, 400-415 e413. https://doi.org/10.1016/j.cell.2018.07.048

785   Earley, A.M., Graves, C.L., Shiau, C.E. (2018). Critical role for a subset of intestinal macrophages in
786   shaping gut microbiota in adult zebrafish. *Cell Rep, 25*, 424-436. https://doi.org/
787   10.1016/j.celrep.2018.09.025
788

789   Eichelberg, K., Galan, J. E. (1999). Differential regulation of Salmonella typhimurium type III secreted
790   proteins by pathogenicity island 1 (SPI-1)-encoded transcriptional activators InvF and hilA. *Infect Immun,*
791   *67*, 4099-4105. https://doi.org/10.1128/IAI.67.8.4099-4105.1999
792

793   El-Shamy, A., Branch, A. D., Schiano, T. D., & Gorevic, P. D. (2018). The complement system and C1q
794   in chronic hepatitis C virus infection and mixed cryoglobulinemia. *Front Immunol, 9*, 1001.
795   https://doi.org/10.3389/fimmu.2018.01001
796

797   Fawzy, M., Edrees, A., Okasha, H., El Ashmaui, A., Ragab, G. (2016). Gastrointestinal manifestations in
798   systemic lupus erythematosus. *Lupus, 25*, 1456-1462. https://doi.org/10.1177/0961203316642308
799

800   Fonseca, M. I., Chu, S. H., Hernandez, M. X., Fang, M. J., Modarresi, L., Selvan, P., MacGregor, G. R.,
801   Tenner, A. J. (2017). Cell-specific deletion of C1qa identifies microglia as the dominant source of C1q in
802   mouse brain. *J Neuroinflammation, 14*, 48. https://doi.org/10.1186/s12974-017-0814-9
803

804   Gabanyi, I., Muller, P. A., Feighery, L., Oliveira, T. Y., Costa-Pinto, F. A., Mucida, D. (2016). Neuro-
805   immune interactions drive tissue programming in intestinal macrophages. *Cell, 164*, 378-391.
806   https://doi.org/10.1016/j.cell.2015.12.023
807

808   Gattu, S., Bang, Y. J., Pendse, M., Dende, C., Chara, A. L., Harris, T. A., Wang, Y., Ruhn, K. A., Kuang,
809   Z., Sockanathan, S., Hooper, L. V. (2019). Epithelial retinoic acid receptor beta regulates serum amyloid A
810   expression and vitamin A-dependent intestinal immunity. *Proc Natl Acad Sci U S A, 116*, 10911-10916.
811   https://doi.org/10.1073/pnas.1812069116
812

813   Grainger, J. R., Konkel, J. E., Zangerle-Murray, T., Shaw, T. N. (2017). Macrophages in gastrointestinal
814   homeostasis and inflammation. *Pflugers Arch, 469*, 527-539. https://doi.org/10.1007/s00424-017-1958-2
815

816   Hammond, J. W., Bellizzi, M. J., Ware, C., Qiu, W. Q., Saminathan, P., Li, H., Luo, S., Ma, S. A., Li, Y.,
817   Gelbard, H. A. (2020). Complement-dependent synapse loss and microgliosis in a mouse model of multiple
818   sclerosis. *Brain Behav Immun, 87*, 739-750. https://doi.org/10.1016/j.bbi.2020.03.004
819

820   Hong, S., Beja-Glasser, V. F., Nfonoyim, B. M., Frouin, A., Li, S., Ramakrishnan, S., Merry, K. M., Shi,
821   Q., Rosenthal, A., Barres, B. A., Lemere, C. A., Selkoe, D. J., Stevens, B. (2016). Complement and
822   microglia mediate early synapse loss in Alzheimer mouse models. *Science, 352*, 712-716.
823   https://doi.org/10.1126/science.aad8373
824

825   Johnson, C.D., Barlow-Anacker, A.J., Pierre, J.F., Touw, K., Erickson, C.S., Furness, J.B., Epstein, M.L.,
826   Gosain, A. (2018). Deletion of choline acetyltransferase in enteric neurons results in postnatal intestinal
827   dysmotility and dysbiosis. *FASEB J, 32*, 4744–4752. https://doi.org/10.1096/fj.201701474RR
828

829   Kishore, U., Reid, K. B. (2000). C1q: structure, function, and receptors. *Immunopharmacology, 49*, 159-
830   170. https://doi.org/10.1016/s0162-3109(00)80301-x
831

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glockner, F. O. (2013).
Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-
based diversity studies. *Nucleic Acids Res, 41*, e1. https://doi.org/10.1093/nar/gks808

Korb, L. C., Ahearn, J. M. (1997). C1q binds directly and specifically to surface blebs of apoptotic human
keratinocytes: complement deficiency and systemic lupus erythematosus revisited. *J Immunol, 158*, 4525-
4528. https://www.ncbi.nlm.nih.gov/pubmed/9144462

Kouser, L., Madhukaran, S. P., Shastri, A., Saraon, A., Ferluga, J., Al-Mozaini, M., Kishore, U. (2015).
Emerging and novel functions of complement protein C1q. *Front Immunol, 6*, 317.
https://doi.org/10.3389/fimmu.2015.00317

Kulkarni, S., Micci, M. A., Leser, J., Shin, C., Tang, S. C., Fu, Y. Y., Liu, L., Li, Q., Saha, M., Li, C.,
Enikolopov, G., Becker, L., Rakhilin, N., Anderson, M., Shen, X., Dong, X., Butte, M. J., Song, H.,
Southard-Smith, E. M., Kapur, R. P., Bogunovic, M., Pasricha, P. J. (2017). Adult enteric nervous system
in health is maintained by a dynamic balance between neuronal apoptosis and neurogenesis. *Proc Natl Acad*
Sci USA, 114, E3709-E3718. https://doi.org/10.1073/pnas.1619406114

Ling, G. S., Crawford, G., Buang, N., Bartok, I., Tian, K., Thielens, N. M., Bally, I., Harker, J. A., Ashton-
Rickardt, P. G., Rutschmann, S., Strid, J., Botto, M. (2018). C1q restrains autoimmunity and viral infection
by regulating CD8+ T cell metabolism. *Science, 360*, 558-563. https://doi.org/10.1126/science.aao4555

Luo, J., Qian, A., Oetjen, L. K., Yu, W., Yang, P., Feng, J., Xie, Z., Liu, S., Yin, S., Dryn, D., Cheng, J.,
Riehl, T. E., Zholos, A. V., Stenson, W. F., Kim, B. S., Hu, H. (2018,). TRPV4 channel signaling in
macrophages promotes gastrointestinal motility via direct effects on smooth muscle cells. *Immunity, 49*,
107-119 e104. https://doi.org/10.1016/j.immuni.2018.04.021

Macedo, A. C., Isaac, L. (2016). Systemic lupus erythematosus and deficiencies of early components of the
complement classical pathway. *Front Immunol, 7*, 55. https://doi.org/10.3389/fimmu.2016.00055

Mao, Y. L., Shen, C. L., Zhou, T., Ma, B. T., Tang, L. Y., Wu, W. T., Zhang, H. X., Lu, H. L., Xu, W. X.,
Wang, Z. G. (2017). Ablation of Tacr2 in mice leads to gastric emptying disturbance. *Neurogastroenterol*
Motil, 29, e13117. https://doi.org/10.1111/nmo.13117

Matheis, F., Muller, P. A., Graves, C. L., Gabanyi, I., Kerner, Z. J., Costa-Borges, D., Ahrends, T.,
Rosenstiel, P., Mucida, D. (2020). Adrenergic signaling in muscularis macrophages limits infection-
induced neuronal loss. *Cell, 180*, 64-78 e16. https://doi.org/10.1016/j.cell.2019.12.002

Maurer, A. H. (2016). Gastrointestinal motility, part 2: small-bowel and colon transit. *J Nucl Med Technol,*
44, 12-18. https://doi.org/10.2967/jnumed.113.134551

Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L., Wold, B. (2008). Mapping and quantifying
mammalian transcriptomes by RNA-Seq. *Nat Methods, 5*, 621-628. https://doi.org/10.1038/nmeth.1226

Muller, P. A., Koscso, B., Rajani, G. M., Stevanovic, K., Berres, M. L., Hashimoto, D., Mortha, A.,
Leboeuf, M., Li, X. M., Mucida, D., Stanley, E. R., Dahan, S., Margolis, K. G., Gershon, M. D., Merad,
M., Bogunovic, M. (2014). Crosstalk between muscularis macrophages and enteric neurons regulates
gastrointestinal motility. *Cell, 158*, 300-313. https://doi.org/10.1016/j.cell.2014.04.050

Noris, M., Remuzzi, G. (2013). Overview of complement activation and regulation. *Semin Nephrol, 33*,
479-492. https://doi.org/10.1016/j.semnephrol.2013.08.001
883
884 Obata, Y., Castaño, A., Boeing, S., Bon-Frauches, A.C., Fung, C., Fallesen, T., de Agüero, M.G., Yilmaz,
885 B., Lopes, R., Huseynova, A., Horswell, S., Maradana, M.R., Boesmans, W., Vanden Berghe, P., Murray,
886 A.J., Stockinger, B., Macpherson, A.J., Pachnis, V. (2020). Neuronal programming by microbiota regulates
887 intestinal physiology. *Nature, 578*, 284-289. https://doi.org/10.1038/s41586-020-1975-8
888
889 Paloneva, J., Manninen, T., Christman, G., Hovanes, K., Mandelin, J., Adolfsson, R., Bianchin, M., Bird,
890 T., Miranda, R., Salmaggi, A., Tranebjaerg, L., Konttinen, Y., Peltonen, L. (2002). Mutations in two genes
891 encoding different subunits of a receptor signaling complex result in an identical disease phenotype. *Am J*
892 *Hum Genet, 71*, 656-662. https://doi.org/10.1086/342259
893
894 Petry, F., Botto, M., Holtappels, R., Walport, M. J., Loos, M. (2001). Reconstitution of the complement
895 function in C1q-deficient (C1qa-/-) mice with wild-type bone marrow cells. *J Immunol, 167*, 4033-4037.
896 https://doi.org/10.4049/jimmunol.167.7.4033
897
898 Rao, M., Gershon, M. D. (2016). The bowel and beyond: the enteric nervous system in neurological
899 disorders. *Nat Rev Gastroenterol Hepatol, 13*, 517-528. https://doi.org/10.1038/nrgastro.2016.107
900
901 Rolig, A.S., Mittge, E.K., Ganz, J., Troll, J.V., Melancon, E., Wiles, T.J., Alligood, K., Stephens, W.Z.,
902 Eisen, J.S., Guillemin, K. (2017). The enteric nervous system promotes intestinal health by constraining
903 microbiota composition. *PLoS Biol*, 15, e2000689. https://doi.org/10.1371/journal.pbio.2000689
904
905 Schifferli, J. A., Ng, Y. C., Peters, D. K. (1986). The role of complement and its receptor in the elimination
906 of immune complexes. *N Engl J Med, 315*, 488-495. https://doi.org/10.1056/NEJM198608213150805
907
908 Schwechter, B., Rosenmund, C., Tolias, K. F. (2013). RasGRF2 Rac-GEF activity couples NMDA receptor
909 calcium flux to enhanced synaptic transmission. *Proc Natl Acad Sci USA, 110*, 14462-14467.
910 https://doi.org/10.1073/pnas.1304340110
911
912 Shah, D., Romero, F., Zhu, Y., Duong, M., Sun, J., Walsh, K., Summer, R. (2015). C1q deficiency promotes
913 pulmonary vascular inflammation and enhances the susceptibility of the lung endothelium to injury. *J Biol*
914 *Chem, 290*, 29642-29651. https://doi.org/10.1074/jbc.M115.690784
915
916 Thielens, N. M., Tedesco, F., Bohlson, S. S., Gaboriaud, C., Tenner, A. J. (2017). C1q: A fresh look upon
917 an old molecule. *Mol Immunol, 89*, 73-83. https://doi.org/10.1016/j.molimm.2017.05.025
918
919 Tian, X. P., Zhang, X. (2010). Gastrointestinal involvement in systemic lupus erythematosus: insight into
920 pathogenesis, diagnosis and treatment. *World J Gastroenterol, 16*, 2971-2977.
921 https://doi.org/10.3748/wjg.v16.i24.2971
922
923 Ural, B. B., Yeung, S. T., Damani-Yokota, P., Devlin, J. C., de Vries, M., Vera-Licona, P., Samji, T., Sawai,
924 C. M., Jang, G., Perez, O. A., Pham, Q., Maher, L., Loke, P., Dittmann, M., Reizis, B., Khanna, K. M.
925 (2020). Identification of a nerve-associated, lung-resident interstitial macrophage subset with distinct
926 localization and immunoregulatory properties. *Sci Immunol, 5*.
927 https://doi.org/10.1126/sciimmunol.aax8756
928

929 van Schaarenburg, R. A., Suurmond, J., Habets, K. L., Brouwer, M. C., Wouters, D., Kurreeman, F. A.,
930 Huizinga, T. W., Toes, R. E., Trouw, L. A. (2016). The production and secretion of complement component
931 C1q by human mast cells. *Mol Immunol, 78*, 164-170. https://doi.org/10.1016/j.molimm.2016.09.001
932
933 Vrees, M. D., Pricolo, V. E., Potenti, F. M., Cao, W. (2002). Abnormal motility in patients with ulcerative
934 colitis: the role of inflammatory cytokines. *Arch Surg, 137*, 439-445.
935 https://doi.org/10.1001/archsurg.137.4.439
936
937 Warren, J., Mastroeni, P., Dougan, G., Noursadeghi, M., Cohen, J., Walport, M. J., Botto, M. (2002).
938 Increased susceptibility of C1q-deficient mice to Salmonella enterica serovar Typhimurium infection. *Infect
939 Immun, 70*, 551-557. https://doi.org/10.1128/IAI.70.2.551-557.2002
940
941 Wessels, M. R., Butko, P., Ma, M., Warren, H. B., Lage, A. L., Carroll, M. C. (1995). Studies of group B
942 streptococcal infection in mice deficient in complement component C3 or C4 demonstrate an essential role
943 for complement in both innate and acquired immunity. *Proc Natl Acad Sci U S A, 92*, 11490-11494.
944 https://doi.org/10.1073/pnas.92.25.11490
945
946 Yang, C. H., Yeh, Y. J., Wang, J. Y., Liu, Y. W., Chen, Y. L., Cheng, H. W., Cheng, C. M., Chuang, Y. J.,
947 Yuh, C. H., Chen, Y. R. (2017). NEAP/DUSP26 suppresses receptor tyrosine kinases and regulates
948 neuronal development in zebrafish. *Sci Rep, 7*, 5241. https://doi.org/10.1038/s41598-017-05584-7
949
950 Yu, X., Rollins, D., Ruhn, K. A., Stubblefield, J. J., Green, C. B., Kashiwada, M., Rothman, P. B.,
951 Takahashi, J. S., Hooper, L. V. (2013). TH17 cell differentiation is regulated by the circadian clock. *Science,
952 342*, 727-730. https://doi.org/10.1126/science.1243884
953
954 Yu, X., Wang, Y., Deng, M., Li, Y., Ruhn, K. A., Zhang, C. C., Hooper, L. V. (2014). The basic leucine
955 zipper transcription factor NFIL3 directs the development of a common innate lymphoid cell precursor.
956 *Elife, 3, e04406*. https://doi.org/10.7554/eLife.04406
957
958
959
960

Figure 1 - figure supplement 1. C1q is expressed in the mouse colon.
RNA-seq analysis of soluble defense collagen expression in the colons of C57BL/6 mice. Data were reanalyzed from Gattu et al., 2019. Each column represents one mouse. Data are available in the Gene Expression Omnibus repository under accession number GSE122471.

Figure 2 - figure supplement 1. C1q expression is lost systemically but preserved in the central nervous system of *C1qa*^{ΔMφ} mice.

(A) qPCR measurement of *C1qa* expression in lung, skin, liver, kidney, and brain. Each data point represents one mouse. Data are representative of two independent experiments.

(B) C1q is absent from the serum of *C1qa*^{ΔMφ} mice. Enzyme-linked immunosorbent assay (ELISA) detection of serum C1q protein from *C1qa*^{fl/fl} and *C1qa*^{ΔMφ} littermates. Data are presented as C1q serum concentration based on a standard curve generated from purified recombinant mouse C1q. Each data point represents one mouse. nd, not detected. Data are representative of three independent experiments.

Error bars represent SEM. ***p<0.001; ****p<0.0001; ns, not significant by two-tailed Student's t-test.

Figure 3 - figure supplement 1. Flow cytometry gating strategy for comparison of T cell populations in *C1qa*^{fl/fl} and *C1qa*^{ΔMφ} mice.

Small intestinal cells were recovered from *C1qa*^{fl/fl} and *C1qa*^{ΔMφ} littermates and analyzed by flow cytometry. T cells were gated as live CD45$^+$ CD3$^+$ CD4$^+$. T cell subsets were further identified by gating into T$_{reg}$ (FoxP3$^+$), T$_H$1 (T-bet$^+$), T$_H$17 (RORγt$^+$), and T$_H$2 (GATA3$^+$). Representative plots from *C1qa*^{fl/fl} mice are presented and comparisons between *C1qa*^{fl/fl} and *C1qa*^{ΔMφ} littermates are shown in Figure 3. FSC, forward scatter.

Figure 3 - figure supplement 2. Flow cytometry gating strategy for comparison of B cell and plasma cell populations in *C1qa*fl/fl and *C1qa*ΔMφ mice.

Small intestinal cells were recovered from *C1qa*fl/fl and *C1qa*ΔMφ littermates and analyzed by flow cytometry. B cells were gated as live CD45[+] CD19[+] B220[+]. Plasma cells were gated as CD19[+] B220[-], and IgA[+] plasma cells were further identified. Representative plots from *C1qa*fl/fl mice are presented and comparisons between *C1qa*fl/fl and *C1qa*ΔMφ littermates are shown in Figure 3. SSC, side scatter.

Figure 3 - figure supplement 3. Flow cytometry gating strategy for comparison of myeloid cell populations in *C1qa*^{fl/fl} and *C1qa*^{ΔMφ} mice.

Small intestinal cells were recovered from *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\varphi}$ littermates and analyzed by flow cytometry. Macrophages were gated as live CD45$^+$ MHCII$^+$ CD11b$^+$ F4/80^{hi}. Monocytes were gated as live CD45$^+$ MHCII$^-$ Ly6C$^{+.}$ Dendritic cells were gated as live CD45$^+$ CD11c$^+$ MHCII$^+$ F4/80^{lo}. Representative plots from *C1qa*$^{fl/fl}$ mice are presented and comparisons between *C1qa*$^{fl/fl}$ and *C1qa*$^{\Delta M\varphi}$ littermates are shown in Figure 3. SSC, side scatter; MHCII, major histocompatibility complex II.

Figure 3 - figure supplement 4. Flow cytometry gating strategy for comparison of innate lymphoid cell populations in *C1qa*^{fl/fl} and *C1qa*^{ΔMφ} mice.

Small intestinal cells were recovered from *C1qa*$^{fl/fl}$ and *C1qa*^{ΔMφ} littermates and analyzed by flow cytometry. Innate lymphoid cells (ILC) were gated as live CD45^{+} Lin^{-} CD90.2^{+} and then further identified as ILC1 (RORγt- GATA3^{-} T-bet^{+}), ILC2 (RORγt^{-} GATA3^{+}) and ILC3 (RORγt^{+} GATA3^{-}). Representative plots from *C1qa*$^{fl/fl}$ mice are presented and comparisons between *C1qa*$^{fl/fl}$ and *C1qa*^{ΔMφ} littermates are shown in Figure 3. SSC, side scatter.

Figure 4 - figure supplement 1. Similar numbers of CD169$^+$ macrophages are present in the small intestines of *C1qa*$^{\Delta M\varphi}$ and *C1qa*$^{fl/fl}$ littermates.

Flow cytometry analysis of CD169$^+$ macrophages was conducted on cells recovered from the small intestines of *C1qa*$^{\Delta M\varphi}$ and *C1qa*$^{fl/fl}$ littermates. CD169$^+$ cells were determined as a percentage of total macrophages (live CD45$^+$ CD11b$^+$ MHCII$^+$ F4/80^{hi} cells).

Figure 4 - figure supplement 2. The colonic longitudinal muscle myenteric plexus of *C1qa*[ΔMφ] mice has a transcriptional profile similar to that of mice with a gastrointestinal motility disorder.

RNA-seq was performed on the colonic longitudinal muscle-myenteric plexus from five *C1qa*[fl/fl] and five *C1qa*[ΔMφ] littermates. Genes that were differentially expressed are represented in a heatmap that depicts $\log_2$(fold change). Genes that also showed altered expression in the TashT mouse line, which is a model of human Hirschsprung's disease (Bergeron et al., 2015), are indicated in red. Statistical significance of overlap between differentially expressed genes in *C1qa*[ΔMφ] and TashT mice was determined by Fisher's exact test (p = 0.0032).

Figure 5 - figure supplements 1 and 2. *Ex vivo* **video recordings of colonic peristalsis in** *C1qa*^{fl/fl} **and** *C1qa*^{ΔMφ}
littermates.
Still images captured from *ex vivo* video recordings of colonic peristalsis to indicate orientation of the colons. Colons were allowed to equilibrate to the organ chamber for 30 minutes before video recording. Movies were captured over 15 minutes. Colons from *C1qa*^{fl/fl} mice are shown in figure supplement 1; colons from *C1qa*^{ΔMφ} are shown in figure supplement 2.