Staphylococcus aureus **prophages are gatekeepers of exogenous lipid utilization**

to control membrane flexibility and bacterial fitness

Byang Zhou[a], Amit Pathania[a], Deepak Pant[a], Philippe Gaudu[a], Patrick Trieu-Cuot[b], Alexandra

Gruss[a] and Karine Gloux[a].

[a]Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, Jouy en Josas, France,

[b] Institut Pasteur, Université de Paris, CNRS UMR 2001, Unité de Biologie des Bactéries

Pathogènes à Gram-Positif, Paris Cedex 15, France.

For correspondence: karine.gloux@inrae.fr

Key words: triglycerides, diglycerides, sphingomyelins, FASII, bacteriophage, Geh lipase, Hlb

hemolysin B, FakB1.

Significance

Staphylococcus aureus is a major human health challenge. Its lipases activities are crucial for

virulence and immune response escape. Prophages inserted in lipases are common in *S. aureus*

but the selective benefits of their maintenance is little understood. We demonstrate that two

prophages, inserted in triglyceride and sphingomyelin lipase genes, interactively modulate fatty

acid (FA) incorporation into bacterial membranes. The triglyceride lipase facilitates elongation of

incoming FAs and impacts membrane composition. Saturated and unsaturated FAs compete for

entry *via* binding protein FakB1. The events dictated by prophage status illustrate how

competitive processes impact membrane flexibility and bacterial fitness in lipid-rich

environments. These findings highlight the importance of assessing prophage status and lipid

environments when evaluating infection risk and therapy efficiency.

Abstract

Lipids are essential components of bacterial membranes and phages are quasi-ubiquitous in

bacteria. Both are crucial mediators of bacteria-host interactions. In the major pathogen

Staphylococcus aureus, two common prophages, ϕ11.1 and ϕ13, are preferentially integrated in

lipases genes: respectively encoding Geh, a triglyceride lipase, and Hlb, a sphingomyelinase. Here

we discovered that ϕ11.1 and ϕ13 prophage statuses interactively determine bacterial growth,

fatty acid (FA) incorporation and elongation *via* their synthesis pathway (FASII), and membrane

composition. Remarkably, FASII-mediated elongation of C18:2 issued from triglycerides was

stimulated by intact Geh, *i.e.*, in non-ϕ11.1 lysogens. Secondly, while *hlb* has no direct role in

triglyceride metabolism, it inhibited C18:2 incorporation from di- and tri-glycerides into bacterial

membrane as observed in non-ϕ13 lysogens. We show that these interactions are based on

competition between Geh-generated fatty acids from tri- and di-glycerides, and Hlb-generated

C16, for binding the fatty acid kinase FakB1 subunit. These findings lead to a mechanistic model

in which prophage regulates the dynamics of lipid utilization, which in turn dictates FA toxicity,

FASII activity and bacterial fitness. We suggest that testing new therapeutics against *S. aureus*

should consider prophage status and environmental lipids for accurate prediction of efficiency.

Introduction

Environmental lipids comprise an integral, but often overlooked, aspect of bacterial pathogenesis. Lipids are crucial for bacterial fitness and are front-line players in modulating interactions with the host (1-4). Lipids range in structure from simple short hydrocarbon chains to complex molecules, including triglycerides (TGs), phospholipids (PLs), sterols and their esters. Fatty acids (FAs), the basic constituents of complex lipids, vary in chain length, side chains, and number and position of double bonds. A vast array of bacteria scavenge FAs, which they incorporate into their membranes (5-11). This quasi-ubiquity suggests a great potential of the host lipid environment to modulate invading pathogen behavior. The environmental lipid composition depends on human genetics and diet and is correlated to infection risk (2, 3, 12, 13). *Staphylococcus* aureus, a major human pathogen, incorporates exogenous FAs, such as those present in plasma lipids (6, 14). Incorporation involves a FA kinase, FakA, whose preference for saturated and unsaturated FA is reportedly respectively conferred by the FA binding proteins FakB1 and FakB2 (15-17). While *S. aureus* does not synthesize unsaturated and polyunsaturated FAs (UFAs and PUFAs), their incorporation has documented effects on bacterial membrane functions and interactions with the host (18-22).

Remarkably, two commonly found *S. aureus* prophages, ɸ11-1 and ɸ13, are inserted in *geh* and *hlb* genes encoding lipases, respectively Geh glycerol ester hydrolase (EC 3.1.1.3) and Hlb neutral sphingomyelinase C (EC 3.1.4.12) (23, 24) (Fig. 1A). About 15% of *S. aureus* isolates carry ɸ11.1 into *geh* and at least 90% of human clinical isolates carry ɸ13 into *hlb* (1, 25). In contrast ɸ13 prophage was quasi-absent in livestock isolates. ɸ11-1 and ɸ13 prophage excision is induced by environmental signals (such as SOS response or biocide/H_2O_2 respectively), leading to reactivation of Geh and Hlb activities with consequent impacts on *in vivo* inflammatory responses

(1, 26-31). In contrast consequences of lipase genes inactivation by φ11.1 and φ13 on *S. aureus*

metabolism and physiology have been little explored (14). Here we show that these two

widespread prophages interactively affect membrane phospholipid composition and growth

response to toxic UFA C18:2. The FA binding protein FakB1 is identified as a control point for

phage-dependent dynamic competition between FAs issued from triglycerides and

sphingomyelins.

Results

φ13 lysogeny increases C18:2 incorporation from serum into *S. aureus* phospholipids. We

analyzed FA profiles from two isogenic Newman strains (WT and Δφ13), which differ by the

presence of φ13 prophage (21). The φ13-lysogenized WT and Δφ13 strains showed identical FA

profiles in control BHI medium,, indicating that FASII-synthesis was unaffected by φ13 (Fig. S1).

In contrast, addition of 10% mouse serum to medium distinguished the two strains by their

incorporation of UFAs. The Δ φ13 strain had approximately two times less UFAs than the WT (6

± 1% *versus* 13 ± 1%), including oleic acids C18:1Δ9, C18:1Δ11, and the PUFA linoleic acid C18:2

(Fig. 1B). Serum was the only UFA source for *S. aureus*, which lacks UFA synthesis and desaturase

functions (32). These results thus suggested an unexpected link between φ13 lysogenic status

and UFA incorporation from a serum source.

φ11.1 lysogeny decreases trilinolein-derived C18:2 incorporation, elongation and toxicity; φ13

effects depend on φ11.1 status. Trilinolein (TNL), a TG with three C18:2, is a substrate for Geh

lipase (33). We asked how the *geh/*φ11.1 status impacts breakdown of TNL to C18:2, and whether

φ13 lysogeny affects this process. For this, we analyzed FA profiles of isogenic Newman strains

lysogenized or not with φ11 (including the three φ11.1 to φ11.3 prophages) and φ13 prophages

(21). In BHI medium, the ф11 prophage status had no detectable effect on FA profiles which

reflect endogenous FA synthesis (Fig. S1). In contrast, TNL addition led to marked differences in

FA profiles and growth: the ф11 lysogenized strains showed markedly less C18:2 incorporation

from TNL and improved growth compared to non- ф11 lysogens (Fig. 1C). This result is in

accordance with *geh* inactivation by ф11.1 and C18:2 toxicity on *S. aureus* (34). The remaining

18:2 incorporation in the ф11 lysogens may indicate residual Geh activity despite the C-terminal

truncation, or C18:2 release by other lipases. Strikingly, ф11 lysogeny correlated with a quasi-

total inhibition of C18:2 elongation by FASII (Fig. 1C) suggesting that Geh facilitates accessibility

of TNL-derived C18:2 to FASII pathway. Furthermore, the ф13 lysogeny had effects depending on

the ф11 status. Indeed when ф11.1 was present, ф13 lysogeny led to higher C18:2 incorporation

from TNL (11 ± 1% in WT *versus* 4 ± 0.5% in Δф13 strain). In contrast, in absence of ф11.1 (intact

geh), it did not impact the C18:2 incorporation (28 ± 4% in Δф11 *versus* 28 ± 1% in Δф11Δф13

strain, Fig. 1C) but increases elongation of both C18:2 (from 50 to 70%) and C20:2 (from 14 to

28%). These observations point to an unexpected interconnection between *S. aureus* ф11 and

ф13 prophages for both incorporation and elongation of the PUFA C18:2 from TNL.

A *geh* mutant shows decreased incorporation of trilinolein-derived C18:2 and its elongation.

We asked whether ф11 prophage family effects were due to *geh* gene interruption by ф11.1. We

analyzed the strain USA300 which lacks ф11.1 (intact *geh*) and its isogenic *geh::tn* mutant. While

TNL-derived C18:2 was incorporated and efficiently elongated in both these strains,

incorporation and particularly elongation were markedly lowered in the USA300 *geh::tn* mutant

(Fig. 1D). Both *geh*::tn mutant and ф11 Newman lysogens encode C-terminal truncations of Geh

(at respective amino acid [aa] positions 517 and 643 from a full length protein of 690 aa).

Furthermore, orlistat, an inhibitor which binds to Geh (35), inhibited both incorporation and

elongation of C18:2 from TNL (Fig. 1D). These results indicate that the Geh state, rather than an

intrinsic property of φ11.1, determines *S. aureus* capacity to both incorporate and elongate TNL-

derived C18:2.

1,3-diglyceride-derived C18:2 is efficiently incorporated and toxic for growth. In TGs, the UFA

position on the glycerol backbone affects its absorption and metabolism by cells (36-38).

Triglyceride lipases display positional selectivity, which can be discriminated using 1,2-

dilinoeoylglycerol (1.2DG) and 1,3-dilinoleoylglycerol (1.3DG) as substrates (39-41). Geh

reportedly cleaves TNL in both positions 1 and 2 whereas most lipases prefer position 1 (40, 41).

Consequently, Geh activity liberates 1.3DG from TNL, in contrast to other lipases. We therefore

examined C18:2 incorporation from 1.2DG and 1.3DG in WT, Δφ11, Δφ13 and Δφ11Δφ13

Newman isogenic strains. Addition of 30 μM 1.2DG led to low-level C18:2 incorporation whatever

the prophage status. A growth retardation was observed in the Geh$^+$ strains (non-φ11.1 lysogens)

where both incorporation and elongation were favored (Fig. 1E). The same concentration of

1.3DG lead to lysis of non-φ11.1 lysogens (data not shown). Lowering 1.3DG addition to 10 μM

avoided lysis. Under this condition, very high incorporation and elongation of C18:2 clearly

occurred only when Geh is intact (in non-φ11.1 lysogens). TG hydrolysis product, particularly

attributed to Geh activity, leads therefore to high PUFA incorporation and elongation in *S. aureus*.

Serum-derived C16 incorporation is increased in Δ φ13 and *hlb*$^+$ strains. We focused on possible

mechanisms involved in φ13 impact on UFAs incorporation. In the presence of 10% mouse serum,

the percentage of C16 and its derived FAs (C18 + C20) was decreased in Newman WT strain

compared to the Δφ13 strain in favor of incorporated serum UFAs (Fig. 2A). Serum

sphingomyelins are dominant circulating Hlb substrates that contain mainly C16 as FA moieties

(42, 43). To further understand how *hlb* affects UFAs utilization, we complemented the USA300

hlb::tn mutant with a plasmid carrying an intact *hlb* gene. Hlb complementation increased C16

and elongated derived FAs whereas it decreased incorporated serum UFAs (Fig. 2B). These results

indicate that *hlb* contributes to C16 liberation and its further incorporation whereas φ13 lysogeny

increases serum UFAs incorporation. Based on these observations, we hypothesized that the φ13

lysogenic/excised status likely creates a dynamic competition between C16 and UFA from serum

TGs for incorporation in membrane phospholipids.

Incorporation of 1.3DG-derived C18:2 requires *fakB1*. The FakA kinase preference for saturated

and unsaturated FA incorporation is reportedly conferred by FakB1 and FakB2, respectively (15-

17). We first confirmed the reported phenotypes in USA300 derivative *fakA*, *fakB1*, and *fakB2*

mutants in keeping with a minor role of FakB1 in UFAs incorporation, including that of free C18:2

PUFA (Fig. S2 and S3)(15, 17)). We then challenged USA300 derivative *fakA*, *fakB1*, and *fakB2*

mutants with TNL. The *fakA* mutant failed to incorporate TNL-derived C18:2 (data not shown).

Unexpectedly, *fakB1* and *fakB2* mutants failed to show selectivity when TNL was used as C18:2

precursor; both mutants displayed the capacity to incorporate C18:2 from TNL (Fig. S4). Thus,

while free C18:2 incorporation was FakB2-dependent (see Fig. S3), incorporation from TNL-

derived C18:2 was permissive. The strong effects of 1.3DG on non-φ11.1 lysogens growth and on

C18:2 incorporation, compared to 1.2DG (Fig. 2), led us to examine incorporation of these

substrates in *fakA*, *fakB1*, and *fakB2* backgrounds. *fakA* was essential for incorporation of both

substrates, reflecting the phosphorylation requirement ((15; data not shown). Incorporation of

1.2DG-derived C18:2 was in accordance with the reported FakB2 binding phenotype (15-17) (Fig.

S5). In contrast, incorporation of 1.3DG-derived C18:2 was remarkably greater in *fakB2* than in

fakB1 mutant (20 ± 7% versus 13 ± 3%) and *fakB1* inactivation results in markedly lower

incorporation compared to the USA300 wild type (13 ± 3 % and 31 ± 5% respectively, Fig. 3A).

To confirm the direct role of *fakB1*, we used RN450 strain, cured of phage lysogens and with a

natural *fakB1* in-frame deletion (483-bp within the 867-bp functional gene, Fig. S6) and the

isogenic RN450-R strain, repaired for *fakB1*. The RN450-R repaired strain incorporated more

1.3DG-derived C18:2 than RN450 strain (35 ± 3% and 27± 3% respectively, Fig. 4A). Furthermore

1.3DG impairs more growth of *fakB1* defective strains (the USA300 *fakB1* mutant and RN450

strain) compared to their respective *fakB1*[+] controls (WT USA300 and RN450-R, Fig. 3; Fig. 4).

These observations identified therefore FakB1 binding as the unconventional preferred pathway

for efficient incorporation of 1.3DG-derived C18:2 into PL and as a contributor of growth

protection.

1,3-diglyceride-derived C18:2 and C16 compete for FakB1. The unexpected role of FakB1 in

C18:2 incorporation from tri- and di-glycerides (Fig. S4A, 3A, 4A) led us to ask whether C16,

favored in *hlb*[+] strains, competes for FakB1. To test this, C16 was also added in experiments

discussed above. We first established that, as reported, free C16 and free C18:2 were poorly

competitive ((17), Fig. S3). In the *fakB1* background (USA300 WT and *fakB2* mutant), addition of

free C16 led to a slight decrease in C18:2 incorporation from either TNL or 1.3DG (Fig. S4, Fig 3).

The 1.3DG-derived C18:2 incorporation is little affected in the WT (22 ± 1 % compared to 31 ±

5%) and unaffected in the *fakB1* mutant (14 ± 4 % compared to 13 ± 3%). In contrast, when only

FakB1 is available (in the *fakB2* mutant), C16 addition led to a quasi-total arrest of C18:2

incorporation (Fig. S4, Fig 3). As example, *fakb2* mutant has more than three times less 1.3DG-

derived C18:2 in the presence of C16 (6 ± 3% compared to 20 ± 7%). In RN450 (naturally fakB1

defective), *fakB2* can compensate for C18:2 incorporation from 1.3DG but C16 had no

competitive effect (Fig. 4). A direct role for *fakB1* was confirmed using RN450-R, repaired for

fakB1. Notably, total incorporation of 1.3DG-derived C18:2 is highly increased in this strain and

quasi-totally inhibited (80 ± 10% decreased) in the presence of free C16 (Fig. 4). FakB1 is

therefore a key determinant responsible for competition between C16 and 1.3DG-derived C18:2.

Discussion

S. aureus metabolism of environmental lipids is modulated by two common prophages, ϕ11.1

and ϕ13, which interrupt the lipases encoding genes *geh* and *hlb*. The present work reveals a

more complex role of these prophages on exogenous lipid utilization than previously supposed.

We identified three interactive factors that determine the bacterial responses: the

environmental lipids offered, which are highly variable according to the biotope; the status of

prophages, which toggle the lipase switches and control FAs integration in the membrane; and

the FA kinase FakB1, which controls competitive entry of PUFA and saturated FAs released from

lipids. Expanded roles for Geh, Hlb and FakB1 in *S. aureus* physiology were thus uncovered. Our

work demonstrates that lysogeny by ϕ11.1 and ϕ13 protects bacteria from lipid toxicity and

expands membrane flexibility, thus conferring a selective benefit for prophage maintenance. We

propose a model for two-level control of FA incorporation and elongation, first by prophages,

and then by FakB1 (Fig. 5).

The first control is determined by ϕ11.1 lysogeny, which protects *S. aureus* from PUFA toxicity.

When TG is available, ϕ11.1 lysogens carrying truncated Geh show decreased PUFA C18:2

incorporation and limited elongation. To our knowledge this is the first report of the impact of

Geh on FASII elongation efficiency. PUFAs provoke membrane permeabilization and lysis (44-49).

Unlike most lipases, Geh cleaves the TG acyl group at both positions on the glycerol backbone

liberating both 1,2- and 1,3-diglycerides (40, 41). Direct 1.3DG addition led to greater C18:2

toxicity than did 1.2DG. By inactivating Geh, φ11.1 lysogeny exerts a protective role on *S. aureus*

by limiting PUFA incorporation in membrane phospholipids, and likely improves bacterial survival

in natural lipid-rich environments. Considering φ13 lysogeny, its prevalence among human

clinical *S. aureus* isolates (25) suggests a functional benefit of preventing Hlb activity. As

expected, *hlb,* intact in the absence of φ13, led to increased saturated FA incorporation from

serum. Remarkably however, intact *hlb*, which is not implicated in PUFA metabolism, also

reduced PUFA incorporation and elongation.

FakB1 defines the second level of control. While FakB1 is reportedly associated with saturated

FA binding (15-17), it also mediated efficient incorporation of C18:2 issued from 1.3DG. Entry of

1.3DG -derived C18:2 contrasts with both free and 1.2DG-derived C18:2, which involved mainly

FakB2. Thus FakB1 binding is more permissive than initially reported, in accordance with recent

studies (15, 17, 50). FakB1 may act as a chaperone that controls free C18:2, toxic on FASII (34),

prior to membrane incorporation. It may also promote FakA-mediated phosphorylation of FAs,

which is required for C18:2 efflux *via* FarE (51). Further investigation is needed to identify the TG

intermediate recognized by FakB1 and Geh contribution to these binding specificities.

The unexpected role of FakB1 in facilitating PUFA incorporation revealed a newly identified role

of *hlb* in an interference with unsaturated FAs incorporation. Hlb activity in φ13 non-lysogens

increases entry of saturated FAs, which bind FakB1, and compete with PUFA. FakB1 is a

gatekeeper that discriminates between PUFA and saturated FA incorporation in membrane

phospholipids. If PUFA-FakB1 binding is out-competed by saturated FAs, *S. aureus* survival could

be compromised due to deficient FakB1 and FarE protecting effects. We propose that the

combined effects of prophages and FakB1 on FA incorporation are relevant to *S. aureus* infection.

Both φ11.1 and φ13 phage may be maintained episomally and reintegrated, implying transient

activation of *geh* and *hlb* expression; these events may help adjust bacterial membrane lipids

during infection (25-30, 52). PUFAs may in turn impact the life cycle of these prophages.

The composition and levels of TGs and SMs vary among individuals and according to dietary

intake. These factors are determinants of infection risk and predisposition to bacterial resistance

(43, 53-57). TGs are major reservoirs of long chain UFAs and PUFAs liberated by Geh activity and

incorporation of these FAs alters bacterial membrane functions and host interactions (6, 18, 19,

22, 58). Cholesterol was recently reported to be a Geh substrate (18); its effects on *S. aureus*

fitness according to prophage status remains to be investigated. Interestingly, intact Geh

contributes to masking *S. aureus* from the immune surveillance by TLR2 receptor (1, 59) and 1,2-

diglycerols moieties are involved in the interaction TLR2 (60). Roles of position 2 hydrolysis and

ϕ11.1 lysogeny in immune escape merit further investigation.

The *in vivo* significance of ϕ13 impacts on PUFAs incorporation and elongation is intriguing.

Specific interference by Hlb in PUFA incorporation is an interesting new avenue to investigate

ϕ13 maintenance in the *hlb* locus specifically in healthy human host (25,61). Indeed

sphingomyelins and their metabolites vary between animals and in disease associated to atypical

ϕ13 integration (62, 63).

In conclusion, ϕ11.1 and ϕ13 are interactive regulators of *S. aureus* membrane phospholipid

composition in response to environmental TGs and SMs. Our findings lead us to suggest that

prophage lysogeny contributes to protecting bacteria from lipid toxicity. Prophage instability also

offers greater membrane flexibility, which contribute to *S. aureus* adaptation in different host

biotopes. In particular, while Hlb (ß toxin) is required for *S. aureus* virulence in humans, we

showed that the associated production of C16 results in an increase of free 1.3DG-derived C18:2

that is toxic for bacteria. Thus, by controlling the cytoplasmic availability of C18:2 and C16,

respectively, φ11.1 and φ13 exert a fine tuning between metabolism and toxicity. These results

provide new mechanistic insights of *S. aureus* lipid physiology and indicate the need to assess

mechanisms of prophage/lipase switches in developing strategies against *S. aureus* infection.

Materials and methods

Bacterial strains. *S. aureus* Newman, USA300 FPR3757 JE2 (called here USA300), and NCTC 8325

derivative strains RN450 and RN450-R, were used in this study. Isogenic Newman strains

differentiated by the presence of prophages were provided by Dr O. Schneewind (Department of

Microbiology, University of Chicago). Newman strain has 4 prophages which are from

Siphoviridae family: φ11.1, φ11.2 and φ11.3 from Sfi 11/Sa6 subfamily and φ13 from Sfi 21/Sa3

subfamily (24). Methicillin-resistant USA300 and the Nebraska library of transposon insertions

(University of Nebraska Medical Center) were generously supplied by BEI resources (64). USA300

derivatives from this library used in this work contained insertions in the following genes:

SAUSA300_0320 (*geh*), SAUSA300_1973 (*hlb*), SAUSA300_1119 (*fakA*), SAUSA300_0733 (*fakB1*),

SAUSA300_1318 (*fakB2*); all mutants were verified by PCR. RN450 is a phage-free derivative of

the NCTC 8325 strain (65). It was repaired for a natural *fakB1* deletion as described for another

strain (66).

Construction of *hlb* complemented strain. A PCR fragment containing the *S. aureus* subsp.

aureus COL *hlb* promoter and open reading frame from SACOL_RS10470 (-355 from ATG start

codon and +40 after TAG stop codon) was cloned in EcoRI and SmaI sites of modified pAW8 (67).

The primers pairs used for the PCR are 5'-TTGCCGGAATTCTGCAACTTAATTATAGCCAGACTTTC-3'

and 5'-CATCAACCCGGGCGTCCTTTTAGAACGAAGCAAG-3'. Genomic co-ordinates of the 1388 bp

cloned segment are 2063371-2064758, as present on plasmid phlb.

Construction of *fakB1*-repaired strain. The *fakB1* gene of *S. aureus* RN450, as the entire NCTC 8325 lineage, displays a 483-bp deletion removing 56% of the 867-bp functional gene (Fig. S6). To repair this deletion, a 1,939-bp DNA fragment containing a functional *fakB1* was amplified as described for other NCTC 8325 derivatives (66). The *fakB1*-repaired strain is designated as RN450-R.

Growth media. Brain heart infusion (BHI) was the base medium for *S. aureus* cultures unless specified. Cultures were grown aerobically at 37°C and the starting OD_{600nm} was 0.02 from a preculture at OD_{600nm} between 1 and 5. Mouse serum (Clinisciences, France) was added at 10% final concentration. Stocks of trilinolein (TNL, 50 mM), 1,2-dilinoleoylglycerol (1.2DG, 50 mM), 1,3-dilinoleoylglycerol (1.3DG, 50 mM), palmitic acid (C16, 100 mM) (Larodan Fine Chemicals, Sweden) and orlistat (38 mM, MedChemExpress) were prepared in DMSO.

Determination of *S. aureus* fatty acid profiles. FA profiles were performed as described (7). For each condition, profiles were determined from three independent cultures. In these experiments culture aliquots were routinely taken under exponential growth at OD_{600} between 0.5 and 2. After sample preparation (7), analyses were performed in a split-splitless injection mode on an AutoSystem XL Gas Chromatograph (Perkin-Elmer) equipped with a ZB-Wax capillary column (30 m x 0.25 mm x 0.25 mm; Phenomenex, France). Data were recorded and analyzed by TotalChrom Workstation (Perkin-Elmer). FA peaks were detected between 12 and 40 min of elution, and identified by comparing to retention times of purified esterified FA standards (Mixture ME100, Larodan, Sweden). Results are shown as representative gas chromatograph profiles, or as percent of specific FAs as calculated from their proportions compared to total peak areas (TotalChrom Workstation; Perkin Elmer).

Statistics analysis. Graphs were prepared using Microsoft Excel software. Means and standard

deviations are presented for culture growth (OD_{600nm}/h) and FA percentages. Statistical

significance was determined by unpaired, nonparametric Mann Whitney tests (U test calculator,

Wilcoxon rank-sum, https://www.statskingdom.com), as recommended for small sample sizes.

Acknowledments: We gratefully acknowledge funding from the following French granting

agencies: French Agence Nationale de la Recherche (StaphEscape project ANR-16-CE15-0013),

and the Fondation pour la Recherche Medicale (DBF20161136769). We thank Myriam Gominet

for the construction of *S. aureus* RN450 *fakB1*-repaired strain.

306
FIG 1 *S. aureus* lysogenic status impacts incorporation and elongation of C18:2 derived from
acylglycerides.
(*A*) Upper, site of φ13 insertion in *hlb*, encoding sphingomyelinase (Hlb). φ13 insertion results in
enzyme N-terminal truncation (position 56 aa of the 330 aa protein). Lower, preferred site of
φ11.1 insertion is *geh*, encoding triacylglycerol lipase (Geh). φ11-1 insertion results in truncation
of the encoded triglyceride Geh (position 643 aa of the 690 aa protein).
(*B*) φ13 lysogen displays greater C18:2 incorporation from mouse serum than non-lysogen. The
Newman strain (wild type, WT), a φ13 lysogen, and an isogenic Δφ13 derivative were grown 2h
in BHI medium containing 10% mouse serum. FAs were extracted and analyzed by gas

chromatography. The y axis represents relative FA abundance (mV response) at indicated peak
positions. Black arrows indicate FAs that can be endogenously produced. Red arrows indicate FAs
not synthesized by *S. aureus* but incorporated from serum. Representative FA profiles from four
determinations are shown.

(*C*) φ13 and φ11 prophages interfere to modulate C18:2 incorporation and elongation from
trilinolein. Four isogenic Newman strains (NM) vary in prophage content, including φ11.1 and
φ13 (21). These isogenic strains were grown 2h in BHI medium supplemented with 30µM
trilinolein (comprising 3 oleic acid moieties). Growth is presented as the absorbance increase in
one hour (OD_{600nm}/h). Proportions of C18:2 and elongated forms (C20:2 and C22:2) were
determined from the FA profiles as in (*A*) and presented as percent of the total FA areas for each
profile (n=5).

(*D*) *geh* truncation decreases both incorporation and elongation of C18:2 from trilinolein. C18:2
incorporation from trilinolein is favored when *geh* is intact. USA300 (*geh*$^+$) and the *geh*::tn, were
grown in media supplemented with TNL. The Geh inhibitor orlistat (orli; 30 µM) was added as
indicated. The confirmed *geh*::tn mutant has a C-terminal truncated Geh (position 517 aa/690aa).
The percent of C18:2 and its elongated forms were determined as in (B). The OD/h of the *geh*::tn
mutant and the wild type were comparable (0.8 ± 0.3 for both cultures), (n=3).

(*E*) Acylated positions in a diglyceride determine C18:2 incorporation into *S. aureus* membrane
phospholipids**.** The four isogenic Newman strains, which differ in φ11.1 and φ13 prophages (21)
were grown 2h in BHI medium supplemented with 30 µM diglycerides comprising linoleic acid
moieties; 1,2-dilinoleoylglycerol (1.2DG) or 10µM 1,3-dilinoleoylglycerol (1.3DG), (n=4). Note
that 1.3DG was more incorporated and more toxic than 1.2DG; 30µM 1.3DG generated cell lysis
and FA profiles of unrepresentative non-lysed subpopulation (not shown). Studies were
therefore with 10µM. Percent of C18:2 and its elongated forms were determined as in (*B*). FA
profiles and percent of C18:2 and elongated forms are described as in (*B*).

p-values were determined by Mann Whitney test on percent total incorporated C18:2, including
elongated forms ('i'), and on percent elongation of C18:2 ('e1'), and C20:2 ('e2'); n.s., not
significant.

FIG 2 Intact *hlb* favors incorporation of saturated FAs and decreases incorporation of UFAs from serum. (*A*) Newman (NM) WT and Δφ13 cultures were grown for 2 hours in BHI medium supplemented with 10% mouse serum. The proportion of UFAs (left) and SFAs (right) are shown. φ13 lysogeny increases unsaturated FAs while lowering incorporation of C16 and elongated forms. (*B*) Plasmid-driven expression of *hlb* increases incorporation of C16 and elongated forms, while lowering incorporation of unsaturated FAs. Cultures of the USA300 (USA) *hlb*::tn pØ strain bearing an empty plasmid or a plasmid carrying intact *hlb* (pHlb, derived from the COL strain) were grown for 2.5 h in BHI supplemented with 10% mouse serum. FA extractions were performed and FA profiles and area of FAs were determined as in Fig. 1. Proportions of saturated FAs (C16, C18and C20) and unsaturated FAs (C18:1Δ9, C18:1Δ11, C18:2 and C20:1) were compared to total FAs (shown as percent of total). Data presented are means ± standard deviations from three independent experiments. p-values were determined by Mann Whitney test.

FIG 3 *fakB1* and C16 modulate 1.3DG-derived C18:2 incorporation and its toxicity on growth. (*A*) Involvement of *S. aureus fakB1* in 1.3DG-mediated C18:2 incorporation. USA300, *fakB1* and *fakB2* strains were grown 2 hours in BHI supplemented with 10μM 1.3DG. (*B*) 1.3DG-derived C18:2 and C16 compete for FakB1. Effects of C16 (20 μM) addition on C18:2 incorporation from 1.3DG were assessed in USA300, *fakB1*, and *fakB2* strains, as in (*A*). Representative FA profiles are shown in (A) and (B), (n=5). (*C*) Growth of WT and *fakB* mutants under conditions described in (*A*) and (*B*), presented as OD_{600nm} per hour. Representative profiles (*A*, *B*) or means (*C*) from four independent experiments are presented. p-values were determined by Mann Whitney test for percent incorporated C18:2, including elongated forms and OD/h.

FIG 4 Competition between 1.3DG-derived C18:2 and C16 at *fakb1* shapes incorporation into phospholipid. (*A*) FA profiles of RN450 (*fakB1*⁻) and RN450-R (*fakB1*⁺) grown 2.5 h in BHI supplemented with 10µM 1.3DG without or with 40µM C16. (*B*) Growth of RN450 and RN450-R cultures from (*A*). Data presented are representative profiles (A) or means (C) from five independent experiments. p-values were determined by the Mann Whitney test for percent incorporated C18:2 (including elongated forms (*A*) and OD/h (*B*).

FIG 5 A mechanistic model for environmental lipid utilization by *S. aureus*; φ11.1, φ13 and *fakB1*
are lipid toxicity and membrane flexibility checkpoints.

φ11.1 lysogeny protects *S. aureus* from toxic C18:2 incorporation from triglyceride. φ11.1 in *geh*
prevents TG hydrolysis, so that *S. aureus* incorporates low amounts of C18:2. In the absence of
φ11.1, or upon stress-induced excision, intact Geh allows TG hydrolysis to 1.3 DG which promotes
C18:2 incorporation. 1.3DG-derived C18:2 (C18:2$^{1.3DG}$) is incorporated via the FakA-FakB kinase
system and the phosphorylated product (C18:2$^{1.3DG}$-P) may then be incorporated in
phospholipids or cycled towards FASII for rounds of elongation before incorporation. The latter
pathway is favored by intact Geh by an unknown mechanism.

FakB1 binds C18:2$^{1.3DG}$ and protects from its toxicity. Exogenous FAs are phosphorylated *via*
FakAB system comprising FakA kinase and FakB1 or FakB2 acyl binding proteins, initially reported
as being specialized in saturated and unsaturated FAs respectively (15, 16). Despite this bias, we
showed in this study that C18:2 from TG or 1.3DG interacts with FakB1 binding protein.

Intact *hlb* interferes with binding of C18:2$^{1.3DG}$ to FakB1, with consequences on toxicity. The
absence of φ13 (or oxidative stress condition) enables Hlb activity and liberation of C16 from
serum SMs. This C16 competes with C18:2$^{1.3DG}$ for FakB1 binding and increases pools of free
C18:2, which may be toxic. In contrast, φ13 lysogeny inactivates *hlb* and prevent sphingomyelins
(SMs) hydrolysis, thus reducing competition. The entry of C18:2$^{1.3DG}$ via FakB1 may prevent
intracellular toxicity and lead to FA elongation via FASII. φ13 lysogeny thus fine-tunes membrane
flexibility via both C18:2$^{1.3DG}$ incorporation and elongation.

C18:2$^{1.3DG}$ represents C18:2 issued from 1.3DG. Impact of φ11.1, green arrows; impact of φ13,
purple arrows; unknown mechanism of fatty acid diffusion, gray arrows. Experimental results of
this study are indicated by the star (*).

Supplementary information titles and legends

Supplementary FIG S1 φ11.1 or φ13 lysogeny has no effect on fatty acid (FA) profiles of *S. aureus* grown in BHI medium. Four isogenic Newman strains, lysogenized or not by φ11.1 and φ13 prophages (1) were grown in BHI medium. FA were extracted at $OD_{600} = 1$ and analyzed by gas chromatography as described in material and methods. Representative FA profiles from three determinations are shown.

Supplementary FIG S2 Validation of free fatty acids (FAs) incorporation in USA300 (WT), and *fakA, fakB1* and *fakB2* mutant derivatives. The indicated strains were grown in BHI in the presence of 0.17 mM each free FAs, C14:0, C16:0, and C18:1. FAs were extracted and analyzed as in Fig. S1. The three exogenous FAs and elongated forms are indicated in red. Data presented are representative profiles from three independent experiments.

Supplementary FIG S3 Incorporation of free C18:2 in the *fakB2* mutant without and with C16 supplementation. The *fakB2* mutant was grown in BHI in the presence 10μM free C18:2 without or with 40 μM C16, and fatty acids (FAs) were extracted and analyzed as in Fig. S1. C16 in upper panel corresponds to endogenous synthesis; the higher C16 peak in lower panel reflects exogenous FA addition. Exogenous FAs or their elongated forms are represented in red. Data presented are representative profiles from three independent experiments.

Supplementary FIG S4 *fakB1* and *fakB2* mutants both mediate C18:2 incorporation from trilinolein, but only the *fakB2* mutant is affected by C16 addition. (*A*) C18:2 incorporation from trilinolein in USA300 wild type and *fakB* mutants. Strains were grown two hours in BHI supplemented with 30μM trilinolein, and fatty acid (FA) profiles were determined as in Fig. S1. The y-axis represents relative fatty acid (FA) abundance (mV response). Exogenous FA incorporated products (C18:2, C20:2, and C22:2) are in red. (B) Cultures were prepared as in (A), except that 20 μM C16 was further added to medium. The *fakB2* mutant, which expresses FakB1, is totally inhibited for C18:2 incorporation. Green and red arrows indicate changes due to C16 addition on peaks of C16, and C18:2 + elongated products, respectively. (C) Growth of WT, *fakB1* and *fakB2* strains in the presence of trilinolein supplemented or not with C16. Samples correspond to those presented in (*A*) and (*B*); results are presented as OD_{600nm} per hour. Data are representative profiles, (*A* and *B*) or means (*C*) from three independent experiments.

Supplementary FIG S5 *fakB2*, and not *fakB1*, affects C18:2 incorporation from 1,2-dilinoleoylglycerol (1.2DG). C18:2 incorporation from 1.2DG was assessed in USA300 WT, *fakB1*, and *fakB2* strains. Strains were grown two hours in BHI supplemented with 30μM 1.2DG, and fatty acids (FA) profiles were determined as in Fig. S1). Exogenous FA incorporated products (C18:2, C20:2, and C22:2) are in red. The *fakB2* mutant shows significant inhibition of the cumulative percentage of C18:2 and elongated forms (10 ± 3 %) compared to wild type (34 ± 5%). Data are representative profiles from three independent experiments.

Supplementary FIG S6 FakB1 defect in NCTC 8325 lineage, comprising RN450. Alignment of FakB1 sequences from Newman (non-NCTC 8325 lineage) and NCTC 8325 strains. The non-deleted region of NCTC 8325 FakB1 shares 99% identity with the synonymous region in Newman. Red arrows indicate amino acids required for FA binding and for FakA-mediated phosphotransfer (1).

References

1. Chen X, Alonzo F, 3rd. 2019. Bacterial lipolysis of immune-activating ligands promotes evasion of innate defenses. Proc Natl Acad Sci USA 116:3764-3773.

2. Sun H, Zhang X, Shi W, Fang B. 2019. Association of soft tissue infection in the extremity with glucose and lipid metabolism and inflammatory factors. Exp Ther Med 17:2535-2540.

3. Trinder M, Boyd JH, Brunham LR. 2019. Molecular regulation of plasma lipid levels during systemic inflammation and sepsis. Curr Opin Lipidol 30:108-116.

4. Gajenthra Kumar N, Contaifer D, Jr., Baker PR, Ekroos K, Jefferson KK, Wijesinghe DS. 2018. Untargeted lipidomic analysis to broadly characterize the effects of pathogenic and non-pathogenic staphylococci on mammalian lipids. PLoS One 13:e0206606.

5. Gloux K, Guillemet M, Soler C, Morvan C, Halpern D, Pourcel C, Vu Thien H, Lamberet G, Gruss A. 2017. Clinical relevance of type II fatty acid synthesis bypass in *Staphylococcus aureus*. Antimicrob Agents Chemother 61: e02515-16.

6. Kenanian G, Morvan C, Weckel A, Pathania A, Anba-Mondoloni J, Halpern D, Gaillard M, Solgadi A, Dupont L, Henry C, Poyart C, Fouet A, Lamberet G, Gloux K, Gruss A. 2019. Permissive fatty fcid Incorporation promotes staphylococcal adaptation to FASII antibiotics in host environments. Cell Rep 29:3974-3982.

7. Morvan C, Halpern D, Kenanian G, Hays C, Anba-Mondoloni J, Brinster S, Kennedy S, Trieu-Cuot P, Poyart C, Lamberet G, Gloux K, Gruss A. 2016. Environmental fatty acids enable emergence of infectious *Staphylococcus aureus* resistant to FASII-targeted antimicrobials. Nat Commun 7:12944.

8. Morvan C, Halpern D, Kenanian G, Pathania A, Anba-Mondoloni J, Lamberet G, Gruss A, Gloux K. 2017. The *Staphylococcus aureus* FASII bypass escape route from FASII inhibitors. Biochimie 141:40-46.

9. Brinster S, Lamberet G, Staels B, Trieu-Cuot P, Gruss A, Poyart C. 2009. Type II fatty acid synthesis is not a suitable antibiotic target for Gram-positive pathogens. Nature 458:83-6.

10. Brinster S, Lamberet G, Staels B, Trieu-Cuot P, Gruss A, Poyart C. 2010. Essentiality of FASII pathway for *Staphylococcus aureus*. Reply. Nature 463:E4-E5.

11. Hays C, Lambert C, Brinster S, Lamberet G, du Merle L, Gloux K, Gruss A, Poyart C, Fouet A. 2021. Type II fatty acid synthesis pathway and cyclopropane ring formation are dispensable during *Enterococcus faecalis* systemic infection. J Bacteriol 203:e0022121.

12. Feingold KR, Grunfeld C. 2000. The effect of inflammation and infection on lipids and lipoproteins. *In* Feingold KR, Anawalt B, Boyce A, Chrousos G, Dungan K, Grossman A, Hershman JM, Kaltsas G, Koch C, Kopp P, Korbonits M, McLachlan R, Morley JE, New M, Perreault L, Purnell J, Rebar R, Singer F, Trence DL, Vinik A, Wilson DP (ed), Endotext, South Dartmouth (MA). https://pubmed.ncbi.nlm.nih.gov/26561701.

13. Kaysen GA, Ye X, Raimann JG, Wang Y, Topping A, Usvyat LA, Stuard S, Canaud B, van der Sande FM, Kooman JP, Kotanko P, Monitoring Dialysis Outcomes I. 2018. Lipid levels are inversely associated with infectious and all-cause mortality: international MONDO study results. J Lipid Res 59:1519-1528.

14. Delekta PC, Shook JC, Lydic TA, Mulks MH, Hammer ND. 2018. *Staphylococcus aureus* utilizes
host-derived lipoprotein particles as sources of fatty acids. J Bacteriol 200:e00728-17.

15. Parsons JB, Broussard TC, Bose JL, Rosch JW, Jackson P, Subramanian C, Rock CO. 2014.
Identification of a two-component fatty acid kinase responsible for host fatty acid
incorporation by *Staphylococcus aureus*. Proc Natl Acad Sci U S A 111:10532-7.

16. Broussard TC, Miller DJ, Jackson P, Nourse A, White SW, Rock CO. 2016. Biochemical roles
for conserved residues in the bacterial fatty acid-binding protein family. J Biol Chem
291:6292-303.

17. Cuypers MG, Subramanian C, Gullett JM, Frank MW, White SW, Rock CO. 2019. Acyl-chain
selectivity and physiological roles of *Staphylococcus aureus* fatty acid-binding proteins. J Biol
Chem 294:38-49.

18. Hines KM, Alvarado G, Chen X, Gatto C, Pokorny A, Alonzo F, 3rd, Wilkinson BJ, Xu L. 2020.
Lipidomic and ultrastructural characterization of the cell envelope of *Staphylococcus aureus*
grown in the presence of human serum. mSphere 5:e00339-20.

19. Nguyen MT, Hanzelmann D, Hartner T, Peschel A, Gotz F. 2016. Skin-specific unsaturated
fatty acids boost the *Staphylococcus aureus* innate immune response. Infect Immun 84:205-
15.

20. Bosch M, Sanchez-Alvarez M, Fajardo A, Kapetanovic R, Steiner B, Dutra F, Moreira L, Lopez
JA, Campo R, Mari M, Morales-Paytuvi F, Tort O, Gubern A, Templin RM, Curson JEB, Martel
N, Catala C, Lozano F, Tebar F, Enrich C, Vazquez J, Del Pozo MA, Sweet MJ, Bozza PT, Gross
SP, Parton RG, Pol A. 2020. Mammalian lipid droplets are innate immune hubs integrating
cell metabolism and host defense. Science 370:eaay8085.

21. Bae T, Baba T, Hiramatsu K, Schneewind O. 2006. Prophages of *Staphylococcus aureus*
Newman and their contribution to virulence. Mol Microbiol 62:1035-47.

22. Arsic B, Zhu Y, Heinrichs DE, McGavin MJ. 2012. Induction of the staphylococcal proteolytic
cascade by antimicrobial fatty acids in community acquired methicillin resistant
Staphylococcus aureus. PLoS One 7:e45952.

23. Lee CY, Iandolo JJ. 1985. Mechanism of bacteriophage conversion of lipase activity in
Staphylococcus aureus. J Bacteriol 164:288-93.

24. Goerke C, Pantucek R, Holtfreter S, Schulte B, Zink M, Grumann D, Broker BM, Doskar J, Wolz
C. 2009. Diversity of prophages in dominant *Staphylococcus aureus* clonal lineages. J
Bacteriol 191:3462-8.

25. Goerke C, Koller J, Wolz C. 2006. Ciprofloxacin and trimethoprim cause phage induction and
virulence modulation in *Staphylococcus aureus*. Antimicrob Agents Chemother 50:171-7.

26. Tran PM, Feiss M, Kinney KJ, Salgado-Pabon W. 2019. phiSa3mw prophage as a molecular
regulatory switch of *Staphylococcus aureus* beta-toxin production. J Bacteriol 201:e00766-
18.

27. Ubeda C, Maiques E, Knecht E, Lasa I, Novick RP, Penades JR. 2005. Antibiotic-induced SOS
response promotes horizontal dissemination of pathogenicity island-encoded virulence
factors in staphylococci. Mol Microbiol 56:836-44.

28. Tang Y, Nielsen LN, Hvitved A, Haaber JK, Wirtz C, Andersen PS, Larsen J, Wolz C, Ingmer H.
2017. Commercial Biocides Induce Transfer of Prophage Phi13 from Human Strains of
Staphylococcus aureus to Livestock CC398. Front Microbiol 8:2418.

29. Salgado-Pabon W, Herrera A, Vu BG, Stach CS, Merriman JA, Spaulding AR, Schlievert PM.
2014. *Staphylococcus aureus* beta-toxin production is common in strains with the beta-toxin
gene inactivated by bacteriophage. J Infect Dis 210:784-92.

30. Goerke C, Wirtz C, Fluckiger U, Wolz C. 2006. Extensive phage dynamics in Staphylococcus
aureus contributes to adaptation to the human host during infection. Mol Microbiol
61:1673-85.

31. Jung P, Abdelbary MM, Kraushaar B, Fetsch A, Geisel J, Herrmann M, Witte W, Cuny C,
Bischoff M. 2017. Impact of bacteriophage Saint3 carriage on the immune evasion capacity
and hemolytic potential of *Staphylococcus aureus* CC398. Vet Microbiol 200:46-51.

32. Parsons JB, Frank MW, Jackson P, Subramanian C, Rock CO. 2014. Incorporation of
extracellular fatty acids by a fatty acid kinase-dependent pathway in *Staphylococcus aureus*.
Mol Microbiol 92:234-45.

33. Cadieux B, Vijayakumaran V, Bernards MA, McGavin MJ, Heinrichs DE. 2014. Role of lipase
from community-associated methicillin-resistant *Staphylococcus aureus* strain USA300 in
hydrolyzing triglycerides into growth-inhibitory free fatty acids. J Bacteriol 196:4044-56.

34. Zheng CJ, Yoo JS, Lee TG, Cho HY, Kim YH, Kim WG. 2005. Fatty acid synthesis is a target for
antibacterial activity of unsaturated fatty acids. FEBS Lett 579:5157-62.

35. Kitadokoro K, Tanaka M, Hikima T, Okuno Y, Yamamoto M, Kamitani S. 2020. Crystal
structure of pathogenic *Staphylococcus aureus* lipase complex with the anti-obesity drug
orlistat. Sci Rep 10:5469.

36. Yoshinaga K, Sasaki K, Watanabe H, Nagao K, Inoue N, Shirouchi B, Yanagita T, Nagai T,
Mizobe H, Kojima K, Beppu F, Gotoh N. 2015. Differential effects of triacylglycerol positional
isomers containing n-3 series highly unsaturated fatty acids on lipid metabolism in C57BL/6J
mice. J Nutr Biochem 26:57-63.

37. Ikeda I, Yoshida H, Tomooka M, Yosef A, Imaizumi K, Tsuji H, Seto A. 1998. Effects of long-
term feeding of marine oils with different positional distribution of eicosapentaenoic and
docosahexaenoic acids on lipid metabolism, eicosanoid production, and platelet aggregation
in hypercholesterolemic rats. Lipids 33:897-904.

38. Schuhmacher M, Grasskamp AT, Barahtjan P, Wagner N, Lombardot B, Schuhmacher JS, Sala
P, Lohmann A, Henry I, Shevchenko A, Coskun U, Walter AM, Nadler A. 2020. Live-cell lipid
biochemistry reveals a role of diacylglycerol side-chain composition for cellular lipid
dynamics and protein affinities. Proc Natl Acad Sci U S A 117:7729-7738.

39. Lyberg AM, Adlercreutz P. 2008. Lipase specificity towards eicosapentaenoic acid and
docosahexaenoic acid depends on substrate structure. Biochim Biophys Acta 1784:343-50.

40. Douchet I, De Haas G, Verger R. 2003. Lipase regio- and stereoselectivities toward three
enantiomeric pairs of didecanoyl-deoxyamino-O methyl glycerol: a kinetic study by the
monomolecular film technique. Chirality 15:220-6. 10.

41. Horchani H, Ben Salem N, Chaari A, Sayari A, Gargouri Y, Verger R. 2010. Staphylococcal
lipases stereoselectively hydrolyse the sn-2 position of monomolecular films of diglyceride
analogs. Application to sn-2 hydrolysis of triolein. J Colloid Interface Sci 347:301-8.

42. Hammad SM, Pierce JS, Soodavar F, Smith KJ, Al Gadban MM, Rembiesa B, Klein RL, Hannun
YA, Bielawski J, Bielawska A. 2010. Blood sphingolipidomics in healthy humans: impact of
sample collection methodology. J Lipid Res 51:3074-87.

43. Hanamatsu H, Ohnishi S, Sakai S, Yuyama K, Mitsutake S, Takeda H, Hashino S, Igarashi Y.
2014. Altered levels of serum sphingomyelin and ceramide containing distinct acyl chains in
young obese adults. Nutr Diabetes 4:e141.

44. Zhang Q, Wu W, Zhang J, Xia X. 2020. Antimicrobial lipids in nano-carriers for antibacterial
delivery. J Drug Target 28:271-281.

45. Butcher GW, King G, Dyke KG. 1976. Sensitivity of *Staphylococcus aureus* to unsaturated fatty
acids. J Gen Microbiol 94:290-6.

46. Knapp HR, Melly MA. 1986. Bactericidal effects of polyunsaturated fatty acids. J Infect Dis
154:84-94.

47. Kelsey JA, Bayles KW, Shafii B, McGuire MA. 2006. Fatty acids and monoacylglycerols inhibit
growth of *Staphylococcus aureus*. Lipids 41:951-61.

48. Kabara JJ, Swieczkowski DM, Conley AJ, Truant JP. 1972. Fatty acids and derivatives as
antimicrobial agents. Antimicrob Agents Chemother 2:23-8.

49. Desbois AP, Smith VJ. 2010. Antibacterial free fatty acids: activities, mechanisms of action
and biotechnological potential. Appl Microbiol Biotechnol 85:1629-42.

50. Krute CN, Ridder MJ, Seawell NA, Bose JL. 2019. Inactivation of the exogenous fatty acid
utilization pathway leads to increased resistance to unsaturated fatty acids in *Staphylococcus
aureus*. Microbiology 165:197-207.

51. Alnaseri H, Kuiack RC, Ferguson KA, Schneider JET, Heinrichs DE, McGavin MJ. 2019. DNA
binding and sensor specificity of FarR, a novel TetR family regulator required for induction of
the fatty scid efflux pump FarE in *Staphylococcus aureus*. J Bacteriol 201 (3):e00602-18.

52. Deutsch DR, Utter B, Verratti KJ, Sichtig H, Tallon LJ, Fischetti VA. 2018. Extra-chromosomal
DNA sequencing reveals episomal prophages capable of impacting virulence factor
expression in *Staphylococcus aureus*. Front Microbiol 9:1406.

53. Astorg P, Bertrais S, Laporte F, Arnault N, Estaquio C, Galan P, Favier A, Hercberg S. 2008.
Plasma n-6 and n-3 polyunsaturated fatty acids as biomarkers of their dietary intakes: a
cross-sectional study within a cohort of middle-aged French men and women. Eur J Clin Nutr
62:1155-61.

54. Hodson L, Skeaff CM, Fielding BA. 2008. Fatty acid composition of adipose tissue and blood
in humans and its use as a biomarker of dietary intake. Prog Lipid Res 47:348-80.

55. Al-Daghri NM, Torretta E, Barbacini P, Asare H, Ricci C, Capitanio D, Guerini FR, Sabico SB,
Alokail MS, Clerici M, Gelfi C. 2019. Sphingolipid serum profiling in vitamin D deficient and
dyslipidemic obese dimorphic adults. Sci Rep 9(1):16664.

56. Trinder M, Walley KR, Boyd JH, Brunham LR. 2020. Causal inference for genetically
determined levels of high-density lipoprotein cholesterol and risk of infectious disease.
Arterioscler Thromb Vasc Biol 40:267-278.

57. Dobner J, Kaser S. 2018. Body mass index and the risk of infection - from underweight to
obesity. Clin Microbiol Infect 24:24-28.

58. Lopez MS, Tan IS, Yan D, Kang J, McCreary M, Modrusan Z, Austin CD, Xu M, Brown EJ. 2017.
Host-derived fatty acids activate type VII secretion in *Staphylococcus aureus*. Proc Natl Acad
Sci U S A 114:11223-11228.

59. Nguyen MT, Luqman A, Bitschar K, Hertlein T, Dick J, Ohlsen K, Broker B, Schittek B, Gotz F.
2018. Staphylococcal (phospho)lipases promote biofilm formation and host cell invasion. Int
J Med Microbiol 308:653-663.

60. Jimenez-Dalmaroni MJ, Radcliffe CM, Harvey DJ, Wormald MR, Verdino P, Ainge GD, Larsen
DS, Painter GF, Ulevitch R, Beutler B, Rudd PM, Dwek RA, Wilson IA. 2015. Soluble human
TLR2 ectodomain binds diacylglycerol from microbial lipopeptides and glycolipids. Innate
Immun 21:175-93.

61. Rohmer C, Wolz C. 2021. The role of *hlb*-converting bacteriophages in *Staphylococcus aureus*
host adaption. Microb Physiol 31(2):109-122.

62. Ferlazzo AM, Bruschetta G, Di Pietro P, Medica P, Notti A, Rotondo E. 2011. Phospholipid
composition of plasma and erythrocyte membranes in animal species by 31P NMR. Vet Res
Commun 35:521-30.

63. Zardini Buzatto A, Abdel Jabar M, Nizami I, Dasouki M, Li L, Abdel Rahman AM. 2021.
Lipidome alterations induced by cystic fibrosis, CFTR mutation, and lung function. J Proteome
Res 20:549-564.

64. Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW. 2013. A genetic
resource for rapid and comprehensive phenotype screening of nonessential *Staphylococcus*
aureus genes. mBio 4:e00537-12.

65. Herbert S, Ziebandt AK, Ohlsen K, Schafer T, Hecker M, Albrecht D, Novick R, Gotz F. 2010.
Repair of global regulators in *Staphylococcus aureus* 8325 and comparative analysis with
other clinical isolates. Infect Immun 78:2877-89.

66. Pathania A, Anba-Mondoloni J, Gominet M, Halpern D, Dairou J, Dupont L, Lamberet G, Trieu-
Cuot P, Gloux K, Gruss A. 2021. (p)ppGpp/GTP and malonyl-CoA modulate *Staphylococcus*
aureus adaptation to FASII antibiotics and provide a basis for synergistic bi-therapy. mBio
12:e03193-20.

67. Katayama Y, Zhang HZ, Hong D, Chambers HF. 2003. Jumping the barrier to beta-lactam
resistance in *Staphylococcus aureus*. J Bacteriol 185:5465-72.