The histone code of the fungal genus *Aspergillus* uncovered by evolutionary and proteomic analyses

Xin Zhang[1,2], Roberta Noberini[3], Tiziana Bonaldi[3,4], Jérôme Collemare[2,*,+], Michael F. Seidl[1,*,+]

Author affiliations: [1]Theoretical Biology & Bioinformatics Group, Department of Biology, Utrecht University, Padualaan 8, 3584 CH Utrecht, the Netherlands; [2]Westerdijk Fungal Biodiversity Institute, Uppsalalaan 8, 3584 CT Utrecht, the Netherlands; [3]Department of Experimental Oncology, IEO, European Institute of Oncology IRCCS, Via Adamello 16, 20139 Milan, Italy; [4]Department of Oncology and Haematology-Oncology, University of Milano, Via Santa Sofia 9/1, 20122 Milano, Italy

[+]These authors contributed equally

[*]Correspondence: J. Collemare (telephone: (+31) 030 21 22 600; email: j.collemare@wi.knaw.nl) and M. F. Seidl (telephone: (+31) 030 253 36 94; email: m.f.seidl@uu.nl)

Abstract

Chemical modifications of DNA and histone proteins impact the organization of chromatin within the nucleus. Changes in these modifications, catalyzed by different chromatin-modifying enzymes, influence chromatin organization, which in turn is thought to impact the spatial and temporal regulation of gene expression. While combinations of different histone modifications, the histone code, have been studied in several model species, we know very little about histone modifications in the fungal genus *Aspergillus*, whose members are generally well-studied due to their importance as models in cell and molecular biology as well as their medical and biotechnological relevance. Here, we used phylogenetic analyses in 94 Aspergilli as well as other fungi to uncover the occurrence and evolutionary trajectories of enzymes and protein complexes with roles in chromatin modifications or regulation. We found that these enzymes and complexes are highly conserved in Aspergilli, pointing towards a complex repertoire of chromatin modifications. Nevertheless, we also observed few recent gene duplications or losses, highlighting *Aspergillus* species to further study the roles of specific chromatin modifications. SET7 (KMT6) and other components of PRC2 (Polycomb Repressive Complex 2), which is responsible for methylation on histone H3 at lysine 27 in many eukaryotes including fungi, are absent in Aspergilli as well as in closely related *Penicillium* species, suggesting that these lost the capacity for this histone modification. We corroborated our computational predictions by performing untargeted mass spectrometry analysis of histone post-translational modifications in *Aspergillus nidulans*. This systematic analysis will pave the way for future research into the complexity of the histone code and its functional implications on genome architecture and gene regulation in fungi.

Keywords

Aspergilli, chromatin, DNA methylation, histone modification, H3K27methylation, mass

spectrometry

Data Summary

The predicted proteomes used in this study are publicly available at the JGI (Joint Genome

Institute) MycoCosm repository (1); the species names and abbreviations are listed in

Supplementary Table 1. To evaluate the completeness of the predicted proteomes and to

obtain a species phylogeny, 758 fungal BUSCO (Benchmarking Universal Single-Copy

Ortholog) genes were used, and their names are listed in Supplementary Table 2. The fasta,

trimmed alignment, and maximum-likelihood phylogenetic tree files can be found in

Supplementary Data 1 and 2 deposited at Zenodo (10.5281/zenodo.6586562). The mass

spectrometry results mentioned in Supplementary Table 6 are deposited in the PRIDE

database with the dataset identifier PXD033478.

Impact Statement

Uncovering how fungi regulate gene expression and genome organization is an important step

to understand how they control biological processes such as growth or biosynthesis of

bioactive molecules. Despite the known importance of chromatin modifications in controlling

a plethora of biological processes across eukaryotes, they remain particularly poorly

understood in filamentous fungi, even in model organisms like Aspergilli. Our systematic

evolutionary analysis provides a robust framework for the functional analysis of chromatin

modifications in *Aspergillus* species and other fungi. Our results do not only implicate

candidate enzymes to play a role in new chromatin modifications, but they also point at species

that have experienced duplications or losses of genes encoding enzymes for specific

chromatin modifications. We therefore expect that this work will set the stage for future

research into the complexity of the histone code and its functional implications on gene

regulation and genome organization in fungi.

Introduction

Inside eukaryotic cells, the membrane-bound nucleus facilitates the storage and replication of the genetic material as well as the regulation of gene expression (2). Within the nucleus, DNA is arranged as a chromatin fiber formed by repeated nucleosome units consisting of histone protein octamers (two copies of each H2A, H2B, H3, and H4 histones) that are wrapped by 145-147 bp of DNA (3,4). The chromatin structure is regulated by chemical modifications such as methylation of DNA on cytosines (5mC) and adenines (6mA), and different post-translational modifications (PTMs) on the histone proteins, including methylation and acetylation (5,6). The combination of these different modifications, the histone code, influences the recruitment of diverse nucleosome-associated proteins (e.g., DNA binding proteins), which can cause dynamic transitions between the condensed and transcriptionally silent heterochromatin and the accessible and transcriptionally active euchromatin (7–9). Chromatin dynamics is therefore key for organisms to adapt gene expression patterns in response to a variety of developmental and environmental clues.

In the past decades, several enzymes involved in orchestrating chromatin dynamics have been identified and functionally characterized in diverse eukaryotes (5,9,10). These enzymes can be broadly classified into three distinct types: (i) writers that place modifications on nucleosomes or linker DNA; (ii) erasers that remove these modifications; and (iii) readers that first recognize specific modifications and subsequently recruit writers, erasers, or other proteins to the correct genomic location (11). Chromatin modifiers that catalyze the same type of chemical modification often harbor functionally and evolutionarily conserved catalytic domains. For instance, DNA methylation (5mC) is catalyzed by different DNA methyltransferases (DNMTs) which contain a DNA methyltransferase domain, while histone mono-, di-, and tri-methylation are nearly exclusively catalyzed by proteins containing a SET (Su (var) 3-9, E(z) and Trithorax) domain (10). Next to these catalytic domains, the enzymes often contain accessory domains, which can contribute to the overall stability of the enzyme or establish interactions with other proteins (12). Furthermore, these multi-domain proteins,

together with other accessory protein subunits, can form multi-protein complexes to find the

correct target region witin the genome and modify the exact amino acid residue, collectively

contributing to the specific functions of these complexes (13,14). For example, the histone

methyltransferase complex PRC2 (Polycomb Repressive Complex 2) is responsible for the

mono-, di-, and tri-methylation of histone protein H3 on the lysine at position 27 (abbreviated

as H3K27me1/2/3; the same abbreviation principle will be applied below for other histone

PTMs) (15–18). Evolutionary analyses have suggested that the key enzymes responsible for

chromatin modifications have evolved and siginificantly expanded early in eukaryotic evolution

(19). An analysis of 94 gene families with roles in chromatin modifications reveals that 87 out

of 94 gene families emerged more than one billion years ago, and 48 were already present in

the last eukaryotic common ancestor (19).

Among eukaryotes, the fungal kingdom is remarkably large and diverse, comprising

an estimated 1,5-5 million species with various morphologies, life cycles, and ecological

niches (20). The most well-studied model fungal species in respect of chromatin modifications

is the unicellular yeast *Saccharomyces cerevisiae*, in which 75 histone PTMs on 43 distinct

amino acid positions have been reported with histone methylation and acetylation being most

abundant (11). In addition to ancient chromatin modifiers, also lineage-specific innovations

such as duplications and losses were found in fungi. For example, Dim-2 and RID (Repeat-

Induced Point Mutation (RIP)-Defective) are considered to be fungal-specific DNMTs, while

DNMT3 that is typically found in animals and plants seems to be absent in fungi (21–23).

However, *S. cerevisiae* is not representative of the complete fungal kingdom, and detailed

studies on PTMs are needed for other, less studied fungal species.

The fungal genus *Aspergillus* comprises approximately 350 species, including

pathogens of humans, animals, and crops, food contaminants, as well as important cell

factories for industrial and medical applications (24). Chromatin dynamics influences gene

expression in several Aspergilli and regulates growth, sexual development, secondary

metabolite biosynthesis, and virulence (11,25–27). Thus far, very few well-known chromatin

modifications and their catalytic protein complexes have been identified and studied in few

Aspergillus species (11). For example, gene knockout of the H3K9 methyltransferase *Clr4*

homolog in *Aspergillus fumigatus* reduced histone H3K9me1/3 results in reduction of radial

growth and conidia production, as well as delayed conidiation (26). Similarly, CclA, a subunit

of the COMPASS (COMplex of Proteins ASsociated with SET1) complex, is essential for

H3K4me2/3 in *A. fumigatus*, and the deletion strain showed increased secondary metabolite

production and decreased growth (28). The SAGA (Spt-Ada-Gcn5 Acetyltransferase) complex,

which is responsible for histone acetylation, has been functionally characterized in *Aspergillus*

nidulans (29). Notably, this complex in *A. nidulans* lacks the Sgf11 and Sus1 subunits that

have been described in *S. cerevisiae*, suggesting that this complexe evolves in distinct

trajectories in different fungi (29). Besides these few examples, we so far lack a complete view

of the occurrence and evolution of chromatin modifiers in the *Aspergillus* genus. Here, we

report the phylogenetic analyses of core enzymes involved in the deposition and removal of

histone PTMs, and assess the conservation of all the subunits belonging to chromatin

regulator complexes in 94 diverse Aspergilli. We also performed proteomics analyses to

corroborate our computational predictions and reveal a broader spectrum of histone

modifications.

Material and Methods

Acquisition of predicted proteomes for 109 fungal species

The predicted proteomes of 109 fungal species were retrieved from JGI (Joint Genome Institute) MycoCosm (https://mycocosm.jgi.doe.gov/mycocosm/home) (1) on 24th January 2020. These species comprise 94 different Aspergilli species, 13 Ascomycota species from other genera, as well as two Basidiomycota species (*Coprinopsis cinerea* and *Dendrothele bispora*) (Supplementary Table 1).

Phylogenetic analyses of chromatin modifiers

Phylogenetic trees for six major chromatin modifiers were generated using their conserved catalytic domains. Catalytic domain HMMs (Hidden Markov Model) of DNA methyltransferase domain (PF00145), histone methylation SET domain (PF00856), acetyltransferase domain (PF00583), histone acetyltransferase MOZ/SAS FAMILY domain (PF01853), histone deacetylase domain (PF00850), and histone demethylase Jumonji C (JmjC) domain (PF02373) were downloaded from the Pfam database (https://pfam.xfam.org). Hmmsearch, which is part of the HMMER package (v3.1b2) (30), with the Gathering Cut-Off threshold (-cut_ga) was used to identify occurrences of these domains in the predicted proteomes of all 109 fungal species considered here. To filter out short and incomplete fragments, we applied a cutoff calculated by 50% length coverage of the query sequence (domain) or the hit. Subsequently, the identified domain sequences were aligned using MAFFT v7.271 (31). TrimAl v1.2 (-gt 0.1) was used to remove positions in the alignments with gaps in more than 90% of the sequences (32). Maximum-likelihood phylogenetic trees were constructed using IQ-TREE v1.6.10 (33,34) using ultrafast bootstrap and Shimodaira–Hasegawa approximate likelihood ratio test (SH-aLRT) (35) with 1,000 pseudo-replicates, and standard model selection to automatically determine the best-fit model. We predicted additional domains in the analyzed proteins using Hmmsearch with all Pfam domain profiles that are present in the Pfam database (v28.0) and visualized them along the phylogenetic trees using iTOL (36). Lastly, additional TBLASTN searches were performed on the NCBI website

(https://blast.ncbi.nlm.nih.gov/Blast.cgi) and checked manually to confirm dubious absences

in the domain trees.

Species tree construction

To construct a species phylogeny of the here analyzed fungi, we used BUSCO v4.0.1

(Benchmarking Universal Single-Copy Ortholog assessment tool) (37) to identify the

occurrence of 758 fungal single-copy genes (Supplementary Table 2) within the 109 predicted

proteomes (Supplementary Table 1). The identified homologs of single-copy BUSCO genes

were subsequently concatenated into a single super-alignment. Then, the maximum-likelihood

species phylogeny was estimated using IQ-TREE v1.6.10 (33,34) by employing partitioned

model selection, which allows selecting a substitution model for each BUSCO gene. Branch

supports were obtained using ultrafast bootstrap and SH-aLRT (35) with 1,000 pseudo-

replicates separately.

Chromatin modifier subunits homology search and gene tree construction

We obtained previously studied chromatin modifier complexes from literature, and the

associated protein sequences were downloaded from NCBI (https://www.ncbi.nlm.nih.gov)

(Supplementary Table 4). For each subunit of the chromatin modifier complexes, homologs in

the analyzed 109 predicted proteomes were detected based on sequence similarity searches.

First, BLASTP (v2.2.31+) (38) searches were performed, and an e-value threshold of 10^{-2} was

applied. Multiple BLAST matches (High-Scoring Pairs: HSPs) of the same query were

concatenated if they were aligned with different non-overlapping regions. Only the non-

redundant region was retained when HSPs of the same query were overlapping. For these

two conditions, the highest e-value of the different matches was used to represent the merged

match. Only matches with the e-value lower than 10^{-5} and covering more than 50% of query

or subject were retained for the following analyses. If we observed less than ten hits for the

BLASTP search after filtering in Aspergilli, we used PSI-BLAST (Position-Specific Initiated

BLAST v2.2.31+) (38) with five iterations (-num_iterations 5) using the best Aspergilli hit from

the BLASTP as the query; if no Aspergilli match was retained, we used the same query as for

the BLASTP searches. The same filter criterion as for the BLASTP selection was applied to

the PSI-BLAST hits, and the results of these two homology search strategies were

subsequently jointed.

To account for apparent protein absences due to erroneous gene annotation, we

tested these potential losses by using TBLASTN (BLAST v2.2.31+) (38) to query protein

sequences against the genome assemblies. Matches with e-value lower than 10^{-5} together

with 4,500 nucleotides upstream and downstream were extracted, and Exonerate v2.2.0 (39)

was used to predict the gene structure (options: protein2genome, extensive search,

showtargetgff, showcigar, and showquerygff options were set as true, and ">%ti(%tcb

- %tce)\n%tcs\n" option was used to output the complete aligned coding sequence). Transeq

v6.6.0.0 (40) was used to translate the coding sequence into protein sequences. Together

with the protein sequences identified from BLASTP and PSI-BLAST, all protein sequences

from each subunit were retrieved to build a phylogenetic tree (MAFFT, TrimAl, and IQ-TREE

settings were the same as for the tree reconstruction using the catalytic domains above).

For each sequence included in the phylogenic trees, the domain organization was

identified and visualized as described above. When the domain organization of a *Aspergillius*

protein differed from its orthologues, the corresponding gene model was manually checked in

the MycoCosm genome browser (1) and corrected. The presence and absence patterns of 83

subunits from 15 different chromatin modifier complexes were visualized using the Seaborn

library in Python (41,42). Lastly, additional TBLASTN searches for dubious absences in the

outgroup species were performed on the NCBI website as absences could be caused by older

and more fragmented assemblies in the JGI database compared to NCBI.

Histone enrichment and mass spectrometry

Spores of 4- or 5-days old sporulating *A. nidulans* (AnWT pabaA1 veA1, kindly provided by

Prof. Joseph Strauss) growing on ME (Malt Extract) agar medium were harvested with 2 mL

sterile water and cultured in 50 mL ME liquid medium at 28 ℃, 200 rpm for 16 hours. The

sample was centrifuged at 8,000 g for 10 minutes to discard supernatant, and germinating

spores were resuspended in 25 mL 0.8 M NaCl and centrifuged again. Ten mL protoplasting

solution (20 mg/mL *Trichoderma* lysing enzyme and 5 mg/mL driselease in 0.8 M NaCl) were

used to resuspend spores, which were then incubated at 30 ℃, 100 rpm for 2-3 hours.

Protoplasts were retrieved by filtering through sterile Miracloth and were concentrated by

centrifugation at 3,000 rpm, 4 ℃ for 5 minutes. Histone proteins were extracted according to

an adapted histone protein extraction protocol (43). Briefly, 10^8 protoplasts were resuspended

in 1 mL Nuclei Isolation Buffer (1×PBS, 0.5 mM PMSF, 5 µM Leupeptin, 5 µM Aprotinin, and

5mM Na-butyrate) with 1% Triton X-100, followed by vigorous pipetting through a 200 µL

pipette several times. Then, centrifuged at 2,300 g for 15 minutes at 4 ℃, discarded the

supernatant, and resuspended the nuclear pellet in 200 µL Nuclei Isolation Buffer with 0.1%

SDS and 250 U Pierce™ Universal Nuclease and incubated at 37 ℃ for 30 minutes to digest

nucleic acids. The protein concentration was measured using the Bradford Protein Assay (44).

The samples were separated on a 17% SDS-PAGE gel and histone bands were

excised, chemically acylated with propionic anhydride and in-gel digested with trypsin,

followed by peptide N-terminal derivatization with phenyl isocyanate (PIC) (45). Chemical

acylation of lysines (which occurs on unmodified or mono-methylated residues) impairs trypsin

cleavage, resulting in proteolityc cleavage at arginine residues only. This treatment allows

obtaining histone peptides of proper length for MS analysis. The samples were then desalted

on handmade StageTips columns (46). Peptide mixtures were separated by reversed-phase

chromatography on an EASY-Spray column (Thermo Fisher Scientic), 25-cm long (inner

diameter 75 µm, PepMap C18, 2 µm particles), which was connected online to a Q Exactive

Plus instrument (Thermo Fisher Scientific) through an EASY-Spray™ Ion Source (Thermo Fisher Scientific), as described (45).

The acquired RAW data were analyzed using the integrated MaxQuant software v.1.6.10. The Uniprot UP000000560 database was used for identification of *A. nidulans* histone peptides. Enzyme specificity was set to Arg-C. The estimated false discovery rate (FDR) was set at a maximum of 1%. The mass tolerance was set to 6 ppm for precursor and fragment ions. Two missed cleavages were allowed, and the minimum peptide length was set to 4 amino acids. Min. score for modified peptides and min. delta score for modified peptides were set to 1. Variable modifications included lysine propionylation, monomethylation + propionylation, dimethylation, trimethylation and acetylation. N-terminal PIC labeling was set as a fixed modification (45). To reduce the search time and the rate of false positives, with increasing the number of variable modifications included in the database search, the raw data were analyzed through multiple parallel MaxQuant jobs (47), setting different combinations of variable modifications. Peptides identified by MaxQuant with Andromeda score higher than 50 and localization probability score higher than 0.75 were quantitated, either manually or by using a version of the EpiProfile 2.0 software (48) adapted to the analysis of histones from *A. nidulans*. Identifications and retention times were used to guide the manual quantification of each modified peptide using QualBrowser version 2.0.7 (Thermo Fisher Scientific). Site assignment was evaluated from MS2 spectra using QualBrowser and MaxQuant Viewer. Extracted ion chromatograms (XICs) were constructed for each doubly charged precursor, based on its *m/z* value, using a mass tolerance of 10 ppm. For each histone modified peptide, the % relative abundance (%RA) was estimated by dividing the area under the curve (AUC) of each modified peptide for the sum of the areas corresponding to all the observed forms of that peptide (49). The AUC values are reported (Supplementary Table 6) and visualized using GraphPad Prism.

Results and discussion

DNA methylation, histone methylation, and histone acetylation are among the most studied chromatin modifications in animals, plants, and fungi (50–52). Protein complexes involved in writing, reading, and erasing these modifications have been reported in few model fungal species to date (16,22), but the evolution and conservation of these complexes in diverse fungal species have not been largely systematically explored. Besides some research on DNA methyltransferases (21,23) and SET domain-containing proteins (17) that have been performed throughout the fungal kingdom, we still lack the evolutionary overview of other chromatin modifiers and in depth analyses for important fungal groups. To start uncovering the occurrence and evolutionary histories of chromatin modifiers involved in these major modifications in the fungal genus *Aspergillus,* we first focused on six conserved catalytic domains reported to be involved in histone modifications: DNA methyltransferase domain for DNMTs, SET domain for histone methyltransferases (HMTs), acetyltransferase domain and MOZ/SAS FAMILY domain for histone acetyltransferases (HATs), histone deacetylase domain for histone deacetylases (HDACs), and JmjC domain for histone demethylases (HDMTs). We identified homologs harboring these conserved domains in 109 predicted proteomes; the completeness of the predicted proteomes was generally high as 97 had more than 97% complete BUSCO predicted proteins (Supplementary Table 1). The 94 Aspergilli species cover 18 recognized sections: *Polypaecilum*, *Cremei*, *Restricti*, *Aspergillus*, *Cervini*, *Clavati*, *Fumigati*, *Candidi*, *Circumdati*, *Janorum*, *Terrei*, *Flavi*, *Ochraceorosei*, *Bispori*, *Cavernicolarum*, *Usti*, *Nidulantes*, and *Nigri* (53–59), and 15 other species were used as outgroups (Supplementary Figure 1). Two of the outgroup species are Basidiomycota and the remaining species cover each class in the Ascomycota phylum, which provides a robust framework for our evolutionary analyses within Aspergilli. The outgroup species include well-studied model species, some of which are important for crop production, food fermentation, and antibiotics production (60,61). We used the identified homologs for each catalytic domain to determine their evolutionary history using maximum-likelihood phylogenies (Figure 1-4 and Supplementary Figure 2 and 3).

Aspergilli harbor conserved DNA methyltransferases for genome defense and methylation maintenance

The conserved DNA methyltransferase domain is the only domain known to catalyze the addition of methyl groups to DNA specifically at cytosines (5mC) (21). Based on previous domain architecture analyses, phylogeny reconstructions, and functional experiments, five distinct protein families containing a DNA methyltransferase domain have been recognized in fungi: DNMT1, DNMT2, and DNMT5 are commonly found in eukaryotes, while Dim-2 and RID are considered to be fungal specific (21). Among these five families, DNMT2 seems to function as a tRNA methyltransferase rather than DNA methyltransferase (62). In line with previous observation (21,63), DNMT2 only occurs in our study in two Basidiomycota and one Ascomycota (*Schizosaccharomyces pombe*), suggesting it has been lost in most Ascomycota.

Three (DNMT1, DNMT5, and Dim-2) of the four remaining families catalyze the maintenance of DNA methylation pattern from parental DNA strand onto synthesized daughter DNA strands, while Dim-2 is also able to catalyze *de novo* methylation (64,65). Based on our species selection and phylogenetic analyses, we here found *DNMT1* and *Dim-2* to be absent in all Aspergilli and only present in four or seven out of 15 outgroup species, respectively, while *DNMT5* is found in most Aspergilli and outgroup species (Figure 1). Only four outgroup species harboring *DNMT1*, the two Basidiomycota species, *Botrytis cinerea*, and *Plectania melastoma*, which corroborates previous findings (21,66). *Dim-2* was identified only in seven Ascomycota outgroup species, and matches in *B. cinerea* and *Zymoseptoria tritici* were included after an additional TBLASTN search in NCBI (Supplementary Table 6). The original *Dim-2* absence in *B. cinerea* by local Hmmsearch is caused by the older version and fragmented assembly at JGI compared to NCBI. The *Z. tritici* IPO323 strain used in this study carries a non-functional *Dim-2* as previously reported (67) and no significant hit was found with Hmmsearch or BLASTP searches, but several fragmented hits with TBLASTN search in

Figure 1. Conservation of DNA methyltransferase families in the *Aspergillus* genus and outgroup species. The maximum-likelihood phylogeny of DNA methyltransferase domain (PF00145) found in 94 Aspergilli and 15 outgroup species was determined using IQ-TREE (37,38). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT values over 80. Leaves from Aspergilli with trustworthy support are collapsed and shown as red triangles. The numbers of hits and species are separated by the pipe in the '#Homologs | #Species' column. The number in parentheses after * indicates the additional hits found by the TBLASTN search on NCBI. Additional conserved domains in DNA methyltransferase domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

NCBI. This loss of *Dim-2* function in *Z. tritici* IPO323 is very recent because other isolates

harbor a functional *Dim-2* gene (67). Its loss leads to increased activity of transposable

elements (TEs) and decreased mutation rate, thus influencing genome defense and genome

evolution (67). In contrast to the *DNMT1* and *Dim-2* losses in Aspergilli, *DNMT5* is identified

in 88 out of 94 Aspergilli and 10 out of 15 outgroup species (Figure 1), with a conserved

domain architecture consisting of a DNA methyltransferase domain, an SNF2 family domain

(PF00176), and a Helicase conserved C-terminal domain (PF00271), in which the latter two

domains are involved in transcription regulation (68). The five outgroup species without

DNMT5 are *S. cerevisiae*, *S pombe*, *Neurospora crassa*, *Fusarium graminearum*, and

Cladonia grayi, among which the two yeast species also lack the other two DNA methylation

maintaining enzymes, and thus accordingly are considered to be incapable of DNA

methylation (69–71). Five out of the six *Aspergillus* species that lost *DNMT5* belong to a single

clade within the *Nidulantes* section (*A. undulatus*, *A. olivicola*, *A. filifera*, *A. venezuelensis*, and

A. stella-maris), and the sixth one is *A. robustus* which is assigned to the *Circumdati* section

(Supplementary Figure 1). In the Basidiomycota fungus *Cryptococcus neoformans*, DNMT5

maintains DNA methylation using the same cofactors as DNMT1 (70,72). Thus, it is most likely

that DNMT5 is similarly involved in DNA methylation maintenance in Aspergilli as in *C.*

neoformans. Moreover, DNA methylation maintenance does not appear to be required for the

survival of fungi as shown with the absence of all three *DNMT1*, *DNMT5*, and *Dim-2* not only

within the *Nidulantes* section but also in *S. cerevisiae* and *S. pombe* (Figure 1). However, the

absence of these genes could have an impact on the regulation of certain genes like

secondary metabolite biosynthetic gene clusters and it would be interesting to compare

species within the *Nidulantes* section in more detail to identify how such loss may have

affected gene expression and genome organization.

RID is a more recently acquired single domain DNMT that specifically occurs in fungi

and is involved in a fungal-specific genome defense mechanism called RIP (Repeat-Induced

Point mutation) (21). RID is a putative *de novo* methyltransferase for TEs in both CG and non-

CG contexts relating to RIP during the sexual stage in *N. crassa***;** however, the capacity of RID

to catalyze 5mC has not yet been demonstrated (73). Here, we identified RID in all analyzed

Aspergilli species as well as in ten outgroup species. A RID-like homolog knock-out mutant in

the model filamentous Ascomycota *Podospora anserina* shows blocked sexual development,

as gametes formation and fertilization can happen but dikaryotic cells cannot develop normally

(74). Interestingly, the knock-out mutant of *DmtA* (homolog of *RID* in *A. flavus*) shows altered

conidiation, sclerotial production, aflatoxin biosynthesis, and virulence (27), suggesting that

RID might have additional functions next to genome defense. Additionally, fungi lost the gene

for *de novo* DNA methyltransferase, *DNMT3,* around 50–150 million years ago (72,75), and

only *DNMT5* and *RID* were consistently found in Aspergilli (Figure 1). Thus, we hypothesize

that Aspergilli can only maintain DNA methylation *via* DNMT5 but do not spur genome-wide

de novo DNA methylation. This hypothesis seems to be corroborated as bisulfite sequencing

in *A. flavus* has uncovered very little genome-wide DNA methylation (76). However,

systematic analyses of DNA methylation patterns and experimental evidence for the roles of

DNMTs in Aspergilli are still largely missing.

Aspergilli lack *SET7* (*KMT6*), but SET-domain proteins are abundant and other histone

methyltransferases are conserved

Histone methylation is typically catalyzed by SET domain-containing proteins (17,77). We

identified a total of 1,548 SET domain-containing proteins and classified them into 15 groups

based on their domain architecture and phylogenetic support that contain species both from

Aspergilli as well as from the outgroup. Among these groups, 107, 111, 109, 102, and 108

sequences represent homologs of the known histone methyltransferases (HMTs) SET9

(KMT5), Dim-5 (KMT1), SET1 (KMT2), Ash1 (KMT2H), and SET2 (KMT3), respectively

(Figure 2). The remaining homologs were divided into nine groups that we broadly classified

as SET domain-containing proteins. Among them, eight groups contain a SET domain but lack

experimentally determined functions, and 117 sequences could be grouped as Ribosomal

lysine N-methyltransferases (Figure 2). Enzymes in this group contain a SET domain together

with a Rubisco LSMT substrate-binding domain (PF09273). While the homolog in plant

transfers a methyl group to the large subunit of the Rubisco holoenzyme complex (78,79), the

400
Figure 2. Conservation of histone methyltransferase families in the *Aspergillus* genus and
outgroup species. The maximum-likelihood phylogeny of SET domains (PF00856) found in 94
Aspergilli and 15 outgroup species was determined using IQ-TREE (37,38). The black dot on the
branch indicates ultrafast bootstrap values over 95 and SH-aLRT values over 80. Leaves from
Aspergilli with trustworthy support are collapsed and shown as red triangles. Leaves with similar
protein architecture but unknown functions covering both Aspergilli and outgroup species are
collapsed and shown as grey triangles. The numbers of hits and species are separated by the pipe
in the '#Homologs | #Species' column. The number in parentheses after * indicates the additional
hits found by the TBLASTN search on NCBI. Additional conserved domains in SET domain-
containing proteins were identified using the PFAM database (https://pfam.xfam.org).

homolog in *S. cerevisiae* has been characterized as a ribosomal lysine N-methyltransferase

(80).

The well-described HMTs, SET1, Dim-5, SET2, Ash1, and SET9, all contain distinct

accessory domains and catalyze methylation on lysine residues at different positions of H3 or

H4 tails (17). SET1 is composed of the SET domain, the N-SET domain (PF11764), as well

as the SET_assoc domain (PF11767), and is responsible for H3K4me1/2/3 in *S. cerevisiae*

(81). SET1 occurs in a single copy in all species (Figure 2), indicating that this enzyme and its

catalyzed PTM play fundamental roles in fungal biology. Dim-5 contains a SET domain and

Pre-SET domain (PF05033) and acts as an H3K9 methyltransferase inducing heterochromatin

formation notably in *N. crassa* (82). Dim-5 was found in all species included in this study

except *S. cerevisiae* and *A. calidoustus*, among which *S. cerevisiae* is reported to lack

H3K9me (82,83). How such a loss has impacted gene expression in *A. calidoustus* in

comparison to its closest relative *A. carlsbadensis* deserves further investigation. Both SET2

and Ash1 catalyze H3K36me in *N. crassa* and share the SET domain and the AWS domain

(PF17907) (Figure 2), yet SET2 harbor one additional SRI domain (PF08236) that interacts

with RNA polymerase II and links H3K36me with transcript elongation (84–86). In *N. crassa,*

Ash1 was shown to contribute to H3K36me at sub-telomeric regions, while SET2 acts primarily

at euchromatic regions (87,88). Considering the common origin of SET2 and Ash1 as well as

their similar activities, an ancient duplication followed by the loss or gain of the SRI domain is

likely to have contributed to their functional diversification (Figure 2). *SET2* was found in all

outgroup species, while *Ash1* was lost in both Basidiomycota and two yeasts. False absences

of *Ash1* in *B. cinerea* and *F. graminearum* were corrected as they could be retrieved by

TBLASTN search at NCBI (Supplementary Table 6). Both enzymes are conserved in all

Aspergillus species except in *A. carbonarius* from the *Nigri* section, suggesting that either

other SET domain-containing proteins exhibit functional redundancy or that H3K36

methylation is not fully required in this species. In *Fusarium fujikuroi*, deletion of *SET2* and

Ash1 strongly affected vegetative growth and conidiation (88). SET9 homologs harbor only a

SET domain, and were shown to catalyze H4K20me in *S. pombe* (89). This group is well

conserved in most of the species except in *S. cerevisiae* (90) and in the two closely related

Aspergilli, *A. novoparasiticus* and *A. arachidicola* that both belong to section *Flavi* (Figure 2

and Supplementary Figure 1).

SET7 is the catalytic subunit of PRC2, and is highly conserved in mammals, plants,

and *N. crassa*, in which it has been well-characterized to catalyze methylation on H3K27, a

PTM that is deposited at facultative heterochromatin (16). Even though PRC2 is considered

to be a key chromatin modifier complex, its catalytic subunit *SET7* was lost in many fungal

species (17). *SET7* has previously been reported to be absent in *A. nidulans* and *A. fumigatus*

(15–17), and we here demonstrate that *SET7* is absent from all analyzed Aspergilli as well as

from *Penicillium* (Figure 2), suggesting that this enzyme was most likely lost at or before the

divergence of the last common ancestor of these two fungal genera. Thus, in line with previous

reports (17,25,91,92), our results indicate that methylation on H3K4, H3K9, H3K36, and

H4K20 are likely catalyzed by conserved enzymes in most Aspergilli, while H3K27 methylation

is absent. The loss of this methylation is usually reported to trigger serious phenotypes in

humans (94) and mice (95). Loss of H3K27me3 in fungi typically leads to gene expression

changes without severe growth defects (16, 93), and thus further investigating how Aspergilli

and other fungi are able to cope with such a loss will likely provide new insights on the histone

code and its evolution in fungi.

Importantly, the occurrence of a high number of uncharacterized SET domain-

containing proteins opens the possibility that these contribute to not yet characterized histone

modifications. Especially, four uncharacterized SET domain-containing groups with 144, 195,

113, and 113 members are conserved in Aspergilli and thus likely play important roles in

different cellular processes.

Histone acetyltransferases are conserved in Aspergilli

Histone acetyltransferases (HATs) have been classified into five families: MYST (MOZ,

Ybf2/Sas3, Sas2, and Tip60)-related HATs, Gcn5-related acetyltransferases (GNATs),

p300/CBP HATs, general transcription factor HATs, and nuclear hormone-related HATs (96–

98). We here only focused on the first two HAT families because these are known to be

involved in the regulation of histone proteins and appear largely conserved in fungi (96).The

MOZ/SAS FAMILY domain is the catalytic domain of MYST-related HATs (99). As previously

reported (100), the three characterized MYST-related HATs, Sas3, Sas2, and Esa1, belong

to distinct monophyletic clades, and these paralogues are nearly fully conserved in the here

analyzed fungi (Figure 3). MOZ/SAS FAMILY domain and zf-MYST domain (PF17772) are

present in Sas3, which can on its own acetylate histone H3 and H4, and H2A weakly *in vitro*

(101). Within the NuA3 (Nucleosomal Acetyltransferase of histone H3) complex, Sas3 histone

acetylation activity is *in vitro* restricted on H3K14 and to a less extent on H3K23, while in *S.*

cerevisiae, only H3K14ac can be detected (102). *Sas3* occurs in most of the species but is

lost in *S. pombe*, as previously observed (103,104), and is duplicated in *A. carbonarius* (Figure

3). In addition to the two domains found in Sas3, Sas2 and Esa1 carry a PHD_4 domain

(PF16866) or Tudor-knot domain (PF11717), respectively (Figure 3). *Esa*1 is present as a

single copy in all species included in this study, and its functions is known to contribute to the

acetylation of K5, K8, K12, K16 on histone H4, as well as on H2AZK14 in *S. cerevisiae* (105–

108). *Sas2* is similarly found in most of the species except for two Basidiomycota as well as

for *A. carbonarius*. Sas2 is involved in chromatin-mediated gene regulation and H4K16

acetylation in both *S. cerevisiae* and *Candida albicans* (105,109,110). Although Esa1 and

Sas2 catalyze H4K16ac, it was shown in *C. albicans* that they are differentially recruited at

different stages of development (105). Because both Esa1 and Sas2 are well-conserved in

the *Aspergillus* genus, we hypothesize that these two HATs may also have distinct roles during

development.

GNATs have a acetyltransferase domain as its catalytic core, and 2,604

acetyltransferase domain-containing proteins were found in our search (Supplementary

Figure 3. Conservation of MOZ/SAS FAMILY domain-containing histone acetyltransferases in the *Aspergillus* genus and outgroup species. The maximum-likelihood phylogeny of MOZ/SAS FAMILY domains (PF01853) found in 94 Aspergilli and 15 outgroup species was determined using IQ-TREE (37,38). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT values over 80. Leaves from Aspergilli with trustworthy support are collapsed and shown as red triangles. The numbers of hits and species are separated by the pipe in the '#Homologs | #Species' column. '*In vitro*' indicates the acetylation activity of this enzyme was only observed on the substrate *in vitro*. Additional conserved domains in MOZ/SAS FAMILY domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

Figure 2). The genes for two well-characterized GNATs, *Gcn5* and *Elp3*, occur in a single copy

in every species included in our study (Supplementary Figure 2). Gcn5 harbors a

Bromodomain (PF00439) in addition to the Acetyltransferase domain and is a crucial subunit of distinct histone acetyltransferase complexes such as the HAT-A2 complex, the SAGA complex, and the ADA complex in *S. cerevisiae* (96,98). All these complexes contain two coactivator proteins, Ada2 and Ada3, that directly bind to Gcn5 (111) and the recruitment of different other subunits determines the location where they catalyze acetylation (96). Elp3 is characterized by the Acetyltransferase domain, Radical SAM (PF04055), and Radical SAM C-terminal domain (PF16199), and has been shown to catalyze H3K14ac and H4K8ac in *S. cerevisiae* (112–114). Elp3 also exhibits a dual enzymatic activity as it catalyzes demethylation in mouse zygotes (115), which is due to the Radical SAM domain that catalyzes the demethylation of methylated lysyl residues on histones (116).

Next to Gcn5 and Elp3, we have identified 2,386 additional acetyltransferase domain-containing proteins. Because their functions remain unknown due to the lack of homology to sequences with characterized functions, they are valuable candidates that may acetylate histones or other proteins. Interestingly, it has been shown that Gcn5 also exhibits succinyltransferase activity in humans (117) and can also function as histone crotonyltransferase to regulate gene transcription *in vitro* (118). It is expected that other less common histone modifications, i.e., histone lysine butyrylation, propionylation, or malonylation, may also be catalyzed by histone acetyltransferases because these histone modifications all use acyl-Coenzyme A to complete the reactions (119–121). Further research is needed to determine whether these less common histone modifications are catalyzed by the known histone acetyltransferases Gcn5 and Elp3, or as of yet unstudied HATs. In conclusion, all five characterized fungal HATs described above are highly conserved in the *Aspergillus* genus, showing their importance in fungal biology. Many other acetyl-transferase domain-containing proteins are encoded in *Aspergillus* genomes and may be responsible for histone acetylation or additional less common histone modifications like butyrylation, succinylation, and malonylation.

Duplication of *RpdA* histone deacetylase gene in the *Flavi* and *Janorum* sections

Five families of histone deacetylases (HDACs) can be grouped into two classes based on sequence similarity: class I (HOS1, HOS2, and Rpd3/RpdA) and class II (HdaA and HOS3) (122,123). They all harbor the Histone deacetylase catalytic domain, and our phylogenetic tree corroborates the HDAC classification into two distinct classes (Figure 4). *HdaA* is found as a single copy in each species included in this study (Figure 4). It harbors the Histone deacetylase domain and C-terminal Arb2 domain (PF09757) that serves as an anchor to target centromeric heterochromatin region (124,125). HdaA deacetylates lysine in histones H3 and H2B, but not H4 nor H2A in *S. cerevisiae* (126,127), and the Δ*hdaA* knock-out strains show increased or reduced production of different secondary metabolites in *A. nidulans* (128), *A. niger* (128), and *A. fumigatus* (129). By contrast, *HOS3* is found in only 80 *Aspergillus* species, as it is lost in nine species (*A. nomiae*, *A. pseudocaelatus*, *A. tamarii*, *A. pseudotamarii*, *A. caelatus*, *A. sergii*, *A. novoparasiticus*, *A. minisclerotigenes*, *A. arachidicola*) in the *Flavi* section, three species (*A. desertorum*, *A. stercorarius*, and *A. multicolor*) in the *Nidulantes* section, and two species (*A. heteromorphus* and *A. carbonarius*) in the *Nigri* section. *HOS3* is well conserved in all outgroup species except *S. pombe* (Figure 4) (130). HOS3 has been shown to catalyze deacetylation with distinct specificity for histone H2AK7, H2BK11, H3K14 and H3K23, as well as H4K5 and H4K8 in *S. cerevisiae* cell extracts (131). Notably, *HOS1* was only detected in three outgroup species, *S. cerevisiae* and both Basidiomycota (Figure 4). The specificity of Hos1 in catalyzing deacetylation on histone proteins is not known yet (132), but it can catalyze the deacetylation of the Smc3 subunit of Cohesin and influence the regulation of chromosome segregation during mitosis in *S. cerevisiae* (133). *HOS2* is found in 15 outgroup species and most of Aspergilli except *A. oryzae* (Figure 4). It is known to deacetylate lysine residues in H3 and H4 histone tails (134). The conservation of *HOS2* in Aspergilli suggests that it plays a key role, which is supported by the deletion of *HOS2* in *A. niger* where the deletion strains displays severe reduction in growth, sporulation, SM biosynthesis, and stress resistance (128).

Figure 4. Conservation of histone deacetylase domain-containing proteins in the *Aspergillus* genus and outgroup species. The maximum-likelihood phylogeny of Histone deacetylase domains (PF00850) found in 94 Aspergilli and 15 outgroup species was determined using IQ-TREE (37,38). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT values over 80. Leaves from Aspergilli with trustworthy support are collapsed and shown as red triangles. The numbers of hits and species are separated by the pipe in the '#Homologs | #Species' column. '*In vitro*' indicates the deacetylation activity of this enzyme was only observed on the substrate *in vitro*. Additional conserved domains in histone deacetylase domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

Within class I HDACs, the Rpd3/RpdA clade is divided into two sub-clades (Figure 4).

One is nearly fully conserved (except in *A. oryzae*) and comprises the well-characterized

Rpd3/RpdA HDACs, while the other one only comprises sequences from the *Flavi* and

Janorum sections (Figure 4). Our phylogenetic reconstruction suggests a duplication event

that has occurred in the ancestor of all Aspergilli giving rise to a paralog that was retained in

two sections only and could explain the *A. oryzae* loss in the conserved sub-clade. The two

paralogs from the same species vary in length as they share a conserved N-terminal catalytic

domain but have a different C-terminus (Figure 4). Follow-up studies are needed to investigate

the function of both paralogues in *Aspergillus* species. Rpd3 is a crucial deacetylase in

eukaryotes, as it is required for deacetylation in most locations on core histone proteins,

except H4K16 *in vitro* (135) that is catalyzed by HOS2 (134). *S. cerevisiae* lacking *Rpd3* is

sensitive to high osmolarity and shows compromised expression of osmotic stress genes

(136). Deletion of *RpdA* in *A. nidulans* is lethal (137), while only heterokaryon disruptants could

be obtained in *A. oryzae* (138), indicating its crucial role in the fungal genus *Aspergillus*. In

summary, class I and class II deacetylases are well conserved in Aspergilli, and duplication

events like the one observed for *RpdA* may confer new activities. While only one paralog was

retained in *A. oryzae*, the presence of two paralogs in sections *Flavi* and *Janorum* may provide

functional redundancy or diversification, which remains to be determined.

Lastly, we reconstructed the phylogeny of histone demethylases harboring the Jumonji

C (JmjC) domain (PF02373) (Supplementary Figure 3). Six groups are distinguished by their

domain distribution: KDM4A/JHDM3/JMJD2, KDM5/JARID, DMM-1, JMJD6/PKDM11,

KDM2/Epe1, and KDM6B/JMJD3, which also agrees with a previous study (91). The first five

groups occur in most of the outgroup and *Aspergillus* species, but the *KDM6B/JMJD3* is only

present in several outgroup species but not in any of the *Aspergillus* species. Certain JmjC

domain-containing proteins were shown to catalyze the demethylation of histone proteins, e.g.,

KDM5/JARID for H3K4me2/3 and KDM4A/JHDM3/JMJD2 for H3K9me2/3 and H3K36me2/3

(139). The histone demethylase activity has not been detected for DMM-1 (140) or Epe1 (141)

in *N. crassa*, while KDM2 can demethylate H3K36me1/2 in mouse (142) and JMJD6/PKDM11

is an arginine demethylase in *Arabidopsis* (143). Interestingly, the human KDM6B/JMJD3 is

responsible for the demethylation of H3K27me2/3 (144). As we did not find *SET7* and thus

conclude that H3K27me3 is absent, it would be tempting to speculate that H3K27 demethylase

should be absent in *Aspergillus* too. However, based on the fact that other enzymes, e.g.,

UTX can also catalyze the demethylation of H3K27 (144,145), we cannot unambiguously

correlate the absence of *KDMB6B/JMJD3* to *SET7* loss. More studies are needed to

investigate the regulation mechanism and interplay of proteins for H3K27 methylation.

Presence-absence patterns of subunits that compose histone modifier complexes

Histone PTMs are catalyzed by the conserved domains found in HMTs, HATs, and HDACs

(Figure 2-4, Supplementary Figure 2), but these chromatin modifying enzymes are often not

acting alone. Instead, they are part of protein complexes that comprise accessory subunits,

which are involved in the recognition and induction of specific histone modifications at the

correct locations (13,14). To further investigate the presence-absence patterns (PAPs) of

known chromatin modifier complexes in Aspergilli, we focused on 15 chromatin modifier

complexes with 83 subunits previously identified and studied in *S. cerevisiae* or *N. crassa*

(Supplementary Table 4). Forty-four subunits are present in all *Aspergillus* and outgroup

species. Most of the accessory subunits are either fully conserved or absent in Aspergilli

(Figure 5 and Supplementary Table 4). Ten subunits show PAPs in both Aspergilli and

outgroup species, while nine subunits appear specific to *S. cerevisiae* as they are not found

in any other species and are likely to play a specific role in this species. Seventeen subunits

show PAPs in the outgroup species, while eight are absent and nine are present in the

Aspergilli. Lastly, three subunits are present in all outgroup species but show PAPs in

Aspergilli.

        Five histone methyltransferase complexes, COMPASS, DCDC (Dim-5/-7/-

9/CUL4/DDB1 Complex), SET2 Complex, PRC2, and Dot1 (disruptor of telomeric silencing 1)

(KMT4) were included in our analyses (Figure 5). Five COMPASS subunits (*SET1*, *CclA*, *Swd1*, *Swd2*, and *Swd3*) are fully conserved, suggesting an important function for H3K4 methylation. Consistent with a PAP in most outgroup species and the loss in Aspergilli, Spp1p is not essential for H3K4 methylation (146). As *Sdc1* shows a PAP in Aspergilli and outgroup species, it is likely that this subunit also does not play a key role in H3K4 methylation. In the DCDC complex, while Dim-7 and Dim-9 seem to be dispensable for H3K9 methylation, Dim-5, Cul4, and Dim-8 are highly conserved, and we thus expect these enzymes to play key roles in the activity of this complex (Figure 5 and Supplementary Table 4). However, it was shown that Dim-7 and Dim-9 are responsible for heterochromatin recognition, which leads to subsequent recruitment of the complete DCDC to induce H3K9 methylation in *N. crassa* (147). Thus, it is very likely that other accessory subunits are involved in recruiting DCDC at specific chromatin locations in Aspergilli. Furthermore, the unique absence of *Dim-8* in *A. indicus* may suggest altered H3K9 methylation in this species. Consistent with the absence of *SET7* (Figure 2), all components of the PRC2 complex are lost in Aspergilli (Figure 5). Lastly, the interaction and crosstalk of multiple chromatin modifications in *Aspergillus* species should be further explored. For example, heterochromatin formation and regulation are done by H3K9me3, DNA methylation, and H3K27me3 in *N. crassa*. In this species, H3K9me3 is recognized by an adapter protein, HP1, to recruit DNA methyltransferase Dim-2 for DNA methylation (73), and cause the redistribution of H3K27me3 (15). In *Aspergillus* species, although HP1 and the writer for H3K9me3, DCDC complex, are well conserved (Figure 5), the inability for genome-wide *de novo* DNA methylation and H3K27me3 suggest different mechanisms of heterochromatin formation and regulation in Aspergilli which remain to be determined.

We included eight histone acetyltransferase complexes that differ by their distinct catalytic subunits (Figure 5). The HAT-A2, ADA, SAGA, and Elongator complexes use an acetyltransferase domain-containing protein as catalytic subunit (29,148). The SAS (Something About Silencing), NuA4 (Nucleosomal Acetyltransferase of histone H4), and NuA3

Figure 5. Presence-absence patterns of subunits that constitute characterized histone
modification complexes. The top row with distinct color boxes indicates outgroup species and
Aspergilli sections, the order of the species from right to left is the same as in the species tree from
top to bottom (Supplementary Figure 1) without *Fusarium oxysporum* and *Aspergillus kawachii*
since they were excluded in the later analysis. Each row represents a subunit in the histone
modification complexes, the order agrees with the summary list (Supplementary Table 5). Each
row shows the presence/absence of a specific complex subunit. For each complex, the first row
represents the catalytic subunit. Each box in the matrix indicates the presence (dark color) or
absence (light color) of a subunit (Supplementary Table 4). The brown boxes indicate additional
hits found with TBLASTN search on NCBI (Supplementary Table 6).

complexes use a MOZ/SAS FAMILY domain-containing protein (149). The NuB4 (Nuclear

Hat1p-containing type B histone acetyltransferase) complex uses the histone

acetyltransferase HAT1 domain-containing protein (150). Gcn5, Ada2, and Ada3 are fully

conserved in all fungi and constitute the core enzymatic module of the HAT-A2, ADA, and

SAGA complexes. This catalytic module contributes to catalyzing histone acetylation, and

other subunits in these complexes guide them to specific locations and link them to the

transcriptional machinery (151). Besides the catalytic module, the ADA complex also harbors

the Ahc1 subunit that is found in *S. cerevisiae* only (148,152). Twelve additional subunits of

the SAGA complex are well conserved in Aspergilli and outgroup species (Figure 5). The

SAGA complex also contains a deubiquitylase (DUB) module, for which the composition

differs between Aspergilli and other species. *Sgf11* is unique to *S. cerevisiae* and *Sus1* was

likely lost in the ancestor of Pezizomycotina because we can only identify *Sus1* orthologs in

both Basidiomycota and the yeasts (Figure 5). The Elongator complex appears more variable

as four subunits are mostly conserved, but *Elp6* is specific to *S. cerevisiae*, and *Hpa2* and

Hpa3 show PAPs in both Aspergilli and outgroup species.

The NuA4, SAS, and NuB4 complexes are well-conserved in Aspergilli as well as in

the outgroup species (Figure 5). These complexes appear slightly different in *S. cerevisiae*

with the presence of specific subunits (*Eaf5* in the NuA4 complex; *Sas4* and *Sas5* in the SAS

complex). Both SAS and NuA4 complexes can catalyze H4K16ac with opposite effects during

cell growth and morphogenesis in *C. albicans* (105). It would be interesting to assess whether

these two complexes exhibit similar activities in Aspergilli. For the NuA3 complex, two

accessory subunits are well conserved among Aspergilli while two others, *Pdp3*, and *Nto1*,

show different PAPs, and *Yng1* is specific to *S. cerevisiae*. *Pdp3* and *Nto1* seem to be lost in

a complementary manner, suggesting a possible functional redundancy (Figure 5). Finally,

two histone deacetylase complexes, Sin3-Rpd3 and HCHC, are fully conserved (Figure 5).

The Sin3-Rpd3 complex is slightly different in *S. cerevisiae* with the specific of the *Ume1*

subunit.

Overall, our analysis showed that the core catalytic modules of complexes involved in

histone PTMs are mostly conserved in all fungi, including Aspergilli, but variability in accessory

subunits could contribute to differences in the localization of histone PTMs along the genome.

Unbiased mass spectrometry captures the diverse histone modifications in *A. nidulans*

and verifies the absence of H3K27 methylation

Our genomic and phylogenetic analyses have identified the presence of multiple chromatin

modifiers in Aspergilli, suggesting that these species have the ability to establish a wide variety

of histone modifications. To assess the presence of histone PTMs predicted in our

evolutionary analyses (Fig 2-5), we extracted histone proteins from *A. nidulans* and performed

unbiased mass spectrometry analyses (Figure 6A and 6B). H3 and H4 peptides with PTMs

corresponding to H3K4me, H3K9me, H3K36me, and H3K79me were detected (Figure 6C),

which we anticipated being catalyzed by SET1, Dim-5, Ash1 and SET2, and Dot1, respectively.

H4K20me, which is catalyzed by SET9 in *S. pombe* (89), was not detected (Figure 6C), which

is likely due to technical reasons because the unmodified form of histone H4 peptide 20-24

containing the K20 residue was not detected. For the acetylation on H3K9, H3K14, H3K18,

H3K23, H4K5, H4K8, H4K12, and H4K16, they were confidently detected (Figure 6C). We did

not observe any peptide that would indicate the presence of H3K27me in *A nidulans*, while we

did detect H3K27ac (Figure 6C and Supplementary Table 6). These results corroborate our

computational analysis that the PRC2 complex is absent in *A. nidulans*, as well as previous

reports on the absence of H3K27me3 in several Aspergilli (17,153,154). Thus, the mass

spectrometry results verify the occurrence of histone modifications predicted to be present

based on our comparative analyses in the fungal genus*Aspergillus*. Moreover, the differences

between the relative abundance of distinct histone modifications provide interesting

information to investigate these abundances and identify specific conditions under which these

modifications increase or decrease in abundance.

Figure 6. Mass spectrometry unlocks the histone code in Aspergilli. A. Schematic
presentation of the in-gel digestion method used to process histones isolated in *A. nidulans*. After
SDS-PAGE separation, gel bands corresponding to histone H3 and H4 are excised and chemically
acylated with propionic anhydride (which derivatizes unmodified and mono-methylated lysines).
Thanks to the derivatization, trypsin cutting at lysines is impaired, and histones are cleaved only
at the C-terminus of arginine. Finally, digested peptides are N-terminally derivatized with phenyl
isocyanate (PIC). **B.** *A. nidulans* histone H3 and H4 sequences. Cleavage sites after PRO-PIC
digestion are shown. Peptides identified in this study are highlighted in light red and labeled using
their starting-ending amino acids. **C.** Relative abundance of post-translational modifications
detected on peptides of histone H3 and H4 from *A. nidulans* using unbiased mass spectrometry
analysis. unmod: unmodified peptide; me1: monomethylation; me2: dimethylation; me3:
trimethylation; ac: acetylation. The term "combinations" refers to multiple modifications occurring
on the same peptide (Supplementary Table 6). **D.** Histone modifications are predicted by genomic

739   and phylogenetic analyses, and/or proved by mass spectrometry in *A. nidulans*. Histone
740   modifications verified by mass spectrometry are labeled as solid dots, while the ones that are only
741   predicted by phylogenic analysis are labeled as solid dots with a red outline.
742

Conclusion

Changes in chromatin organization play an important role in regulating the expression of genes (130). The fungal genus *Aspergillus* is well known for its ability to thrive in a broad range of different environmental niches and the huge potential of secondary metabolite biosynthesis (11,155). However, a detailed and complete overview of occurrence and evolution of chromatin modifiers in Aspergilli is a prerequisite to be able to better understand how these fungi regulate gene expression. In the future, more research is needed to uncover how the absence of *de novo* DNA methylation impacts genome evolution in Aspergilli. We demonstrate that PRC2 is lost throughout Aspergilli, a finding consistent with the lack of H3K27me detection by mass spectrometry. Although histone acetyltransferases and deacetylases are overall conserved, our study also revealed specific duplication and loss events, which require further investigation to understand their impact on chromatin and gene regulation in certain *Aspergillus* species. Moreover, other less common histone modifications like histone lysine succinylation, crotonylation, butyrylation, propionylation, and malonylation, are all likely catalyzed by HATs and HDACs as they all use Coenzyme A (CoA) to complete the reactions (119–121), but may require specific yet to be discovered subunits. In conclusion, our study provides a detailed overview of the evolutionary routes of chromatin modifier complex in the fungal genus *Aspergillus*, and therefore generates the necessary framework to identify targets for functional studies to understand how chromatin regulates gene expression in Aspergilli.

Acknowledgment

The sequence data were produced by the US Department of Energy Joint Genome Institute

(http://www.jgi.doe.gov/) in collaboration with the user community. Especially, we thank the

Scott Baker, Mikael Andersen, and Joseph Spatafora for their permission to use the sequence

data. We thank Jos Houbraken for validating the *Aspergillus* species tree and Joseph Strauss

for providing *A. nidulans* strain. We thank Alessandro Vai for support with analysis of mass

spectrometry data for histone PTMs' identification on *Aspergillus* core histone sequences.

Funding information

Xin Zhang is funded by the Chinese Scholarship Council (CSC) (201907720028). This work

has been supported by EPIC-XS, project number 823839, funded by the Horizon 2020

programme of the European Union.

Authors and contribution

X.Z. formal analysis, investigation, methodology, visualization, writing original draft;

R.N. formal analysis, investigation, methodology, visualization; T.B. funding

acquisition, supervision, resources; J.C. conceptualization, funding acquisition,

methodology, project administrations, supervision, visualization, writing original draft;

M.F.S. conceptualization, funding acquisition, methodology, project administrations,

supervision, visualization, writing original draft.

References

1. Grigoriev I V, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, et al. MycoCosm portal: gearing up for 1000 fungal genomes. Nucleic acids research. 2014;42(D1):D699–704.

2. Labrador M, Corces VG. Setting the boundaries of chromatin domains and nuclear organization. Cell. 2002;111(2):151–4.

3. Luger K, Mader AW, Richmond RK, Sargent DF, Richmond TJ. Crystal structure of the nucleosome resolution core particle at 2.8 Å. Nature. 1997;389:251–60.

4. Luger K. Structure and dynamic behavior of nucleosomes. Current opinion in genetics & development. 2003;13(2):127–35.

5. Jenuwein T, Allis CD. Translating the histone code. Science. 2001;293(5532):1074–80.

6. Peterson CL, Laniel M-A. Histones and histone modifications. Current Biology. 2004;14(14):R546–51.

7. Santos-Rosa H, Schneider R, Bannister AJ, Sherriff J, Bernstein BE, Emre NCT, et al. Active genes are tri-methylated at K4 of histone H3. Nature. 2002;419(6905):407–11.

8. Grewal SIS, Jia S. Heterochromatin revisited. Nature Reviews Genetics. 2007;8(1):35–46.

9. Berger SL. The complex language of chromatin regulation during transcription. Nature. 2007;447(7143):407–12.

10. Xiao B, Wilson JR, Gamblin SJ. SET domains and histone methylation. Current Opinion in Structural Biology. 2003;13(6):699–705.

11. Collemare J, Seidl MF. Chromatin-dependent regulation of secondary metabolite biosynthesis in fungi: is the picture complete? FEMS Microbiology Reviews. 2019;43(6):591–607.

12. Helmlinger D, Papai G, Devys D, Tora L. What do the structures of GCN5-containing complexes teach us about their function? Biochimica et Biophysica Acta - Gene Regulatory Mechanisms. 2021;1864(2).

13. Hajheidari M, Koncz C, Bucher M. Chromatin evolution-key innovations underpinning

morphological complexity. Frontiers in Plant Science. 2019;10:1–12.

14.   Gates LA, Foulds CE, O'Malley BW. Histone marks in the 'drivers seat': functional

roles in steering the transcription cycle. 2018;42(12):977–89.

15.   Jamieson K, Wiles ET, McNaught KJ, Sidoli S, Leggett N, Shao Y, et al. Loss of HP1

causes depletion of H3K27me3 from facultative heterochromatin and gain of

H3K27me2 at constitutive heterochromatin. Genome Research. 2016;26(1):97–107.

16.   Jamieson K, Rountree MR, Lewis ZA, Stajich JE, Selker EU. Regional control of

histone H3 lysine 27 methylation in *Neurospora*. Proceedings of the National

Academy of Sciences of the United States of America. 2013;110(15):6027–32.

17.   Freitag M. Histone Methylation by SET Domain Proteins in Fungi. Annual Review of

Microbiology. 2017;71:413–39.

18.   Kamei M, Ameri AJ, Ferraro AR, Bar-peled Y, Zhao F. IMITATION SWITCH is

required for normal chromatin structure and gene repression in PRC2 target domains.

2021;118(4).

19.   Weiner AKM, Cerón-Romero MA, Yan Y, Katz LA. Phylogenomics of the epigenetic

toolkit reveals punctate retention of genes across eukaryotes. Genome Biology and

Evolution. 2020;1–39.

20.   Choi J, Kim S-H. A genome tree of life for the fungi kingdom. Proceedings of the

National Academy of Sciences. 2017;114(35):9391–6.

21.   Bewick AJ, Hofmeister BT, Powers RA, Mondo SJ, Grigoriev I V., James TY, et al.

Diversity of cytosine methylation across the fungal tree of life. Nature Ecology and

Evolution. 2019;3(3):479–90.

22.   Kouzminova E, Selker EU. Dim-2 encodes a DNA methyltransferase responsible for

all known cytosine methylation in *Neurospora*. The EMBO journal. 2001;20(15):4309–

23.

23.   Nai YS, Huang YC, Yen MR, Chen PY. Diversity of fungal DNA methyltransferases

and their association with DNA methylation patterns. Frontiers in Microbiology.

2021;11.

24.   Samson RA, Visagie CM, Houbraken J, Hong SB, Hubka V, Klaassen CHW, et al.

Phylogeny, identification and nomenclature of the genus *Aspergillus*. Studies in

Mycology. 2014;78(1):141–73.

25.   Cánovas D, Marcos AT, Gacek A, Ramos MS, Gutiérrez G, Reyes-Domínguez Y, et

al. The histone acetyltransferase GcnE (GCN5) plays a central role in the regulation

of *Aspergillus* asexual development. Genetics. 2014;197(4):1175–89.

26.   Palmer JM, Perrin RM, Dagenais TRT, Keller NP. H3K9 methylation regulates growth

and development in *Aspergillus fumigatus*. Eukaryotic Cell. 2008;7(12):2052–60.

27.   Yang K, Liang L, Ran F, Liu Y, Li Z, Lan H, et al. The DmtA methyltransferase

contributes to *Aspergillus flavus* conidiation, sclerotial production, aflatoxin

biosynthesis and virulence. Scientific Reports. 2016;6:1–13.

28.   Palmer JM, Bok JW, Lee S, Dagenais TRT, Andes DR, Kontoyiannis DP, et al. Loss

of CclA, required for histone 3 lysine 4 methylation, decreases growth but increases

secondary metabolite production in *Aspergillus fumigatus*. PeerJ. 2013;2013(1):1–18.

29.   Georgakopoulos P, Lockington RA, Kelly JM. The Spt-Ada-Gcn5 Acetyltransferase

(SAGA) Complex in *Aspergillus nidulans*. PLoS ONE. 2013;8(6).

30.   Eddy SR. Accelerated profile HMM searches. PLoS computational biology.

2011;7(10):e1002195.

31.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Molecular biology and evolution.

2013;30(4):772–80.

32.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics.

2009;25(15):1972–3.

33.   Chernomor O, Von Haeseler A, Minh BQ. Terrace aware data structure for

phylogenomic inference from supermatrices. Systematic biology. 2016;65(6):997–

1008.

34.   Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular

biology and evolution. 2015;32(1):268–74.

35.   Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O. Survey of branch support

methods demonstrates accuracy, power, and robustness of fast likelihood-based

approximation schemes. Systematic biology. 2011;60(5):685–99.

36.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. Nucleic acids research. 2019;47(W1):W256–9.

37.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V, Zdobnov EM. BUSCO:

assessing genome assembly and annotation completeness with single-copy

orthologs. Bioinformatics. 2015;31(19):3210–2.

38.   Boratyn GM, Camacho C, Cooper PS, Coulouris G, Fong A, Ma N, et al. BLAST: a

more efficient report with usability improvements. Nucleic acids research.

2013;41(W1):W29–33.

39.   Slater GSC, Birney E. Automated generation of heuristics for biological sequence

comparison. BMC bioinformatics. 2005;6(1):31.

40.   Rice P, Longden I, Bleasby A. EMBOSS: the European molecular biology open

software suite. Elsevier current trends; 2000.

41.   Hunter JD. Matplotlib: A 2D graphics environment. Computing in science &

engineering. 2007;9(03):90–5.

42.   Waskom ML. Seaborn: statistical data visualization. Journal of Open Source

Software. 2021;6(60):3021.

43.   Noberini R, Restellini C, Savoia EO, Bonaldi T. Enrichment of histones from patient

samples for mass spectrometry-based analysis of post-translational modifications.

Methods. 2020;184:19–28.

44.   He F. Bradford protein assay. Bio-protocol. 2011;45.

45.   Noberini R, Savoia EO, Brandini S, Greco F, Marra F, Bertalot G, et al. Spatial epi-

proteomics enabled by histone post-translational modification analysis from low-

abundance clinical samples. Clinical epigenetics. 2021;13(1):1–16.

46.   Prudhomme N, Gianetto-Hill C, Pastora R, Cheung W-F, Allen-Vercoe E, McLean

895   MD, et al. Quantitative proteomic profiling of shake flask versus bioreactor growth

896   reveals distinct responses of *Agrobacterium tumefaciens* for preparation in molecular

897   pharming. Canadian Journal of Microbiology. 2021;67(1):75–84.

898   47.   Bremang M, Cuomo A, Agresta AM, Stugiewicz M, Spadotto V, Bonaldi T. Mass

899   spectrometry-based identification and characterisation of lysine and arginine

900   methylation in the human proteome. Molecular bioSystems. 2013;9(9):2231–47.

901   48.   Yuan Z-F, Sidoli S, Marchione DM, Simithy J, Janssen KA, Szurgot MR, et al.

902   EpiProfile 2.0: a computational platform for processing epi-proteomics mass

903   spectrometry data. Journal of proteome research. 2018;17(7):2533–41.

904   49.   Pesavento JJ, Mizzen CA, Kelleher NL. Quantitative analysis of modified proteins and

905   their positional isomers by tandem mass spectrometry: human histone H4. Analytical

906   chemistry. 2006;78(13):4271–80.

907   50.   Wade PA, Pruss D, Wolffe AP. Histone acetylation: Chromatin in action. Trends in

908   Biochemical Sciences. 1997;22(4):128–32.

909   51.   Binda O. On your histone mark, SET, methylate! Epigenetics. 2013;8(5):457–63.

910   52.   Wolffe AP, Matzke MA. Epigenetics: Regulation through repression. Science.

911   1999;286(5439):481–6.

912   53.   James TY, Stajich JE, Hittinger CT, Rokas A. Toward a Fully Resolved Fungal Tree of

913   Life. Annual Review of Microbiology. 2020;74(1):291–313.

914   54.   Sun B Da, Houbraken J, Frisvad JC, Jiang XZ, Chen AJ, Samson RA. New species in

915   *Aspergillus* section *Usti* and an overview of *Aspergillus* section *Cavernicolarum*.

916   International Journal of Systematic and Evolutionary Microbiology. 2020;

917   55.   Chen AJ, Frisvad JC, Sun BD, Varga J, Kocsubé S, Dijksterhuis J, et al. *Aspergillus*

918   section *Nidulantes* (formerly *Emericella*): Polyphasic taxonomy, chemistry and

919   biology. Studies in Mycology. 2016;84:1–118.

920   56.   Steenwyk JL, Shen XX, Lind AL, Goldman GH, Rokas A. A robust phylogenomic time

921   tree for biotechnologically and medically important fungi in the genera *Aspergillus* and

Penicillium. mBio. 2019;10(4):1–25.

57.   Vesth TC, Nybo JL, Theobald S, Frisvad JC, Larsen TO, Nielsen KF, et al.

Investigation of inter- and intraspecies variation through genome sequencing of

Aspergillus section *Nigri*. Nature Genetics. 2018;50(12):1688–95.

58.   Kjærbølling I, Vesth T, Frisvad JC, Nybo JL, Theobald S, Kildgaard S, et al. A

comparative genomics study of 23 *Aspergillus* species from section *Flavi*. Nature

Communications. 2020;11(1).

59.   Souza SC, Pereira VM, Moreira SI, Costa SS, Moreira GM, Mendes WO, et al.

Aspergillus trisporus: A new *Jani* section species from Brazilian soil. Current

Research in Environmental and Applied Mycology. 2019;9(1):175–86.

60.   Grunstein M, Gasser SM. Epigenetics in *Saccharomyces cerevisiae*. Cold Spring

Harbor Perspectives in Biology. 2013;5(7).

61.   Nielsen JC, Grijseels S, Prigent S, Ji B, Dainat J, Nielsen KF, et al. Global analysis of

biosynthetic gene clusters reveals vast potential of secondary metabolite production in

Penicillium species. Nature Microbiology. 2017;2(April).

62.   Ko YH, So KK, Chun J, Kim DH. Distinct roles of two dna methyltransferases from

cryphonectria parasitica in fungal virulence, responses to hypovirus infection, and viral

clearance. mBio. 2021;12(1):1–16.

63.   Jurkowski TP, Jeltsch A. On the evolutionary origin of eukaryotic DNA

methyltransferases and Dnmt2. PLoS ONE. 2011;6(11):1–9.

64.   Bestor T. Structure of mammalian DNA methyltransferase as deduced from the

inferred amino acid sequence and direct studies of the protein. Portland Press Ltd.;

1988.

65.   Okano M, Xie S, Li E. Cloning and characterization of a family of novel mammalian

DNA (cytosine-5) methyltransferases. Nature genetics. 1998;19(3):219–20.

66.   Zhang Z, He C, Chen Y, Li B, Tian S. DNA methyltransferases regulate pathogenicity

of *Botrytis cinerea* to horticultural crops. Journal of Fungi. 2021;7(8).

67.   Moller M, Habig M, Lorrain C, Feurtey A, Haueisen J, Fagundes WC, et al. Recent

loss of the Dim2 DNA methyltransferase decreases mutation rate in repeats and

changes evolutionary trajectory in a fungal pathogen. PLoS Genetics. 2021;17(3):1–

27.

68.   Delmas V, Stokes DG, Perry RP. A mammalian DNA-binding protein that contains a

chromodomain and an SNF2/SWI2-like helicase domain. Proceedings of the National

Academy of Sciences. 1993;90(6):2414–8.

69.   Capuano F, Mülleder M, Kok R, Blom HJ, Ralser M. Cytosine DNA methylation is

found in *Drosophila melanogaster* but absent in *Saccharomyces cerevisiae*,

Schizosaccharomyces pombe, and other yeast species. Analytical chemistry.

2014;86(8):3697–702.

70.   Huff JT, Zilberman D. Dnmt1-independent CG methylation contributes to nucleosome

positioning in diverse eukaryotes. Cell. 2014;156(6):1286–97.

71.   Proffitt JH, Davie JR, Swinton D, Hattman S. 5-Methylcytosine is not detectable in

Saccharomyces cerevisiae DNA. Molecular and cellular biology. 1984;4(5):985–8.

72.   Catania S, Dumesic PA, Pimentel H, Nasif A, Stoddard CI, Burke JE, et al.

Evolutionary persistence of DNA methylation for millions of years after ancient loss of

a *de novo* methyltransferase. Cell. 2020;180(2):263-277.e20.

73.   Rountree MR, Selker EU. DNA methylation and the formation of heterochromatin in

Neurospora crassa. Heredity. 2010;105(1):38–44.

74.   Grognet P, Timpano H, Carlier F, Aït-Benkhali J, Berteaux-Lecellier V, Debuchy R, et

al. A RID-like putative cytosine methyltransferase homologue controls sexual

development in the fungus *Podospora anserina*. PLoS genetics.

2019;15(8):e1008086.

75.   Madhani HD. Unbelievable but True: Epigenetics and Chromatin in Fungi. Trends in

Genetics. 2020;

76.   Liu S-Y, Lin J-Q, Wu H-L, Wang C-C, Huang S-J, Luo Y-F, et al. Bisulfite sequencing

reveals that *Aspergillus flavus* holds a hollow in DNA methylation. PloS one.

2012;7(1):e30349.

77.   Spellmon N, Holcomb J, Trescott L, Sirinupong N, Yang Z. Structure and function of

979   SET and MYND domain-containing proteins. International Journal of Molecular

980   Sciences. 2015;16(1):1406–8.

981   78.   Raunser S, Magnani R, Huang Z, Houtz RL, Trievel RC, Penczek PA, et al. Rubisco

982   in complex with Rubisco large subunit methyltransferase. Proceedings of the National

983   Academy of Sciences of the United States of America. 2009;106(9):3160–5.

984   79.   Trievel RC, Flynn EM, Houtz RL, Hurley JH. Mechanism of multiple lysine methylation

985   by the SET domain enzyme Rubisco LSMT. Nature Structural Biology.

986   2003;10(7):545–52.

987   80.   Porras-Yakushi TR, Whitelegge JP, Clarke S. Yeast ribosomal/cytochrome c SET

988   domain methyltransferase subfamily: Identification of Rpl23ab methylation sites and

989   recognition motifs. Journal of Biological Chemistry. 2007;282(17):12368–76.

990   81.   Roguev A, Schaft D, Shevchenko A, Pijnappel WWMP, Wilm M, Aasland R, et al. The

991   *Saccharomyces cerevisiae* Set1 complex includes an Ash2 homologue and

992   methylates histone 3 lysine 4. The EMBO journal. 2001;20(24):7137–48.

993   82.   Zhang X, Tamaru H, Khan SI, Horton JR, Keefe LJ, Selker EU, et al. Structure of the

994   *Neurospora* SET Domain Protein DIM-5, a Histone H3 Lysine Methyltransferase. Cell.

995   2002;111:9.

996   83.   Marina DB, Shankar S, Natarajan P, Finn KJ, Madhani HD. A conserved ncRNA-

997   binding protein recruits silencing factors to heterochromatin through an RNAi-

998   independent mechanism. Genes & development. 2013;27(17):1851–6.

84.   Strahl BD, Grant PA, Briggs SD, Sun Z-W, Bone JR, Caldwell JA, et al. Set2 is a

nucleosomal histone H3-selective methyltransferase that mediates transcriptional

repression. Molecular and cellular biology. 2002;22(5):1298–306.

85.   Xiao T, Hall H, Kizer KO, Shibata Y, Hall MC, Borchers CH, et al. Phosphorylation of

RNA polymerase II CTD regulates H3 methylation in yeast. Genes & development.

2003;17(5):654–63.

86.   Li M, Phatnani HP, Guan Z, Sage H, Greenleaf AL, Zhou P. Solution structure of the

Set2–Rpb1 interacting domain of human Set2 and its interaction with the

hyperphosphorylated C-terminal domain of Rpb1. Proceedings of the National

Academy of Sciences. 2005;102(49):17636–41.

87.   Bicocca VT, Ormsby T, Adhvaryu KK, Honda S, Selker EU. ASH1-catalyzed H3K36

methylation drives gene repression and marks H3K27me2/3-competent chromatin.

eLife. 2018;7:1–19.

88.   Janevska S, Baumann L, Sieber CMK, Münsterkötter M, Ulrich J, Kämper J, et al.

Elucidation of the two H3K36me3 histone methyltransferases Set2 and Ash1 in

Fusarium fujikuroi unravels their different chromosomal targets and a major impact of

Ash1 on genome stability. Genetics. 2018;208:153–71.

89.   Wang Y, Reddy B, Thompson J, Wang H, Noma K, Yates III JR, et al. Regulation of

Set9-mediated H4K20 methylation by a PWWP domain protein. Molecular cell.

2009;33(4):428–37.

90.   Veerappan CS, Avramova Z, Moriyama EN. Evolution of SET-domain protein families

in the unicellular and multicellular Ascomycota fungi. BMC Evolutionary Biology.

2008;8(1):1–20.

91.   Gacek-Matthews A, Noble LM, Gruber C, Berger H, Sulyok M, Marcos AT, et al.

KdmA, a histone H3 demethylase with bipartite function, differentially regulates

primary and secondary metabolism in *Aspergillus nidulans*. Molecular Microbiology.

2015;96(4):839–60.

92.   Bachleitner S, Sørensen JL, Gacek-Matthews A, Sulyok M, Studt L, Strauss J.

Evidence of a demethylase-independent role for the H3K4-specific histone

demethylases in *Aspergillus nidulans* and *Fusarium graminearum* secondary

metabolism. Frontiers in Microbiology. 2019;10.

93.   Kramer HM, Seidl MF, Thomma BPHJ, Cook DE. Local rather than global H3K27me3

dynamics associates with differential gene expression in *Verticillium dahliae*. mBio.

2022 Feb;13(1):e0356621.

94.   Sneeringer CJ, Scott MP, Kuntz KW, Knutson SK, Pollock RM, Richon VM, et al.

Coordinated activities of wild-type plus mutant EZH2 drive tumor-associated

hypertrimethylation of lysine 27 on histone H3 (H3K27) in human B-cell lymphomas.

Proceedings of the National Academy of Sciences. 2010;107(49):20980–5.

95. Tao Y, Neppl RL, Huang Z-P, Chen J, Tang R-H, Cao R, et al. The histone

methyltransferase Set7/9 promotes myoblast differentiation and myofibril assembly.

Journal of Cell Biology. 2011;194(4):551–65.

96. Carrozza MJ, Utley RT, Workman JL, Côté J. The diverse functions of histone

acetyltransferase complexes. Trends in Genetics. 2003;19(6):321–9.

97. Torchia J, Glass C, Rosenfeld MG. Co-activators and co-repressors in the integration

of transcriptional responses. Current Opinion in Cell Biology. 1998;10(3):373–83.

98. Sterner DE, Berger SL. Acetylation of histones and transcription-related factors.

Microbiology and Molecular Biology Reviews. 2000;64(2):435–59.

99. Roth SY, Denu JM, Allis CD. Histone acetyltransferases. Annual review of

biochemistry. 2001;70(1):81–120.

100. Kawahara T, Siegel TN, Ingram AK, Alsford S, Cross GAM, Horn D. Two essential

MYST-family proteins display distinct roles in histone H4K10 acetylation and telomeric

silencing in trypanosomes. Molecular microbiology. 2008;69(4):1054–68.

101. Takechi S, Nakayama T. Sas3 is a histone acetyltransferase and requires a zinc

finger motif. Biochemical and biophysical research communications.

1999;266(2):405–10.

102. Taverna SD, Ilin S, Rogers RS, Tanny JC, Lavender H, Li H, et al. Yng1 PHD Finger

Binding to H3 Trimethylated at K4 Promotes NuA3 HAT Activity at K14 of H3 and

Transcription at a Subset of Targeted ORFs. Molecular Cell. 2006;24(5):785–96.

103. Gómez EB, Espinosa JM, Forsburg SL. *Schizosaccharomyces pombe* mst2+

encodes a MYST family histone acetyltransferase that negatively regulates telomere

silencing. Molecular and Cellular Biology. 2005;25(20):8887–903.

104. Nugent RL, Johnsson A, Fleharty B, Gogol M, Xue-Franzén Y, Seidel C, et al.

Expression profiling of *S. pombe* acetyltransferase mutants identifies redundant

pathways of gene regulation. Bmc Genomics. 2010;11(1):59.

105. Kim J, Park S, Lee JS. Epigenetic control of oxidative stresses by histone

acetyltransferases in *Candida albicans*. Journal of Microbiology and Biotechnology.

2018;28(2):181–9.

106. Millar CB, Xu F, Zhang K, Grunstein M. Acetylation of H2AZ Lys 14 is associated with

genome-wide gene activity in yeast. Genes and Development. 2006;20(6):711–22.

107. Searle NE, Torres-Machorro AL, Pillus L. Chromatin regulation by the NuA4

acetyltransferase complex is mediated by essential interactions between enhancer of

polycomb (Epl1) and Esa1. Genetics. 2017;205(3):1125–37.

108. Allard S, Utley RT, Savard J, Clarke A, Grant P, Brandl CJ, et al. NuA4, an essential

transcription adaptor/histone H4 acetyltransferase complex containing Esa1p and the

ATM-related cofactor Tra1p. EMBO Journal. 1999;18(18):5108–19.

109. Wang X, Chang P, Ding J, Chen J. Distinct and redundant roles of the two MYST

histone acetyltransferases Esa1 and Sas2 in cell growth and morphogenesis of

Candida albicans. Eukaryotic Cell. 2013;12(3):438–49.

110. Meijsing SH, Ehrenhofer-Murray AE. The silencing complex SAS-I links histone

acetylation to the assembly of repressed chromatin by CAF-I and Asf1 in

Saccharomyces cerevisiae. Genes and Development. 2001;15(23):3169–82.

111. Balasubramanian R, Pray-Grant MG, Selleck W, Grant PA, Tan S. Role of the Ada2

and Ada3 transcriptional coactivators in histone acetylation. Journal of Biological

Chemistry. 2002;277(10):7989–95.

112. Winkler GS, Kristjuhan A, Erdjument-Bromage H, Tempst P, Svejstrup JQ. Elongator

is a histone H3 and H4 acetyltransferase important for normal histone acetylation

levels *in vivo*. Proceedings of the National Academy of Sciences of the United States

of America. 2002;99(6):3517–22.

113. Krogan NJ, Greenblatt JF. Characterization of a Six-Subunit Holo-Elongator Complex

Required for the Regulated Expression of a Group of Genes in *Saccharomyces*

cerevisiae. Molecular and Cellular Biology. 2001;21(23):8203–12.

114. Wittschieben B, Otero G, De Bizemont T, Fellows J, Erdjument-Bromage H, Ohba R,

et al. A novel histone acetyltransferase is an integral subunit of elongating RNA

polymerase II holoenzyme. Molecular Cell. 1999;4(1):123–8.

115. Okada Y, Yamagata K, Hong K, Wakayama T, Zhang Y. A role for the elongator

complex in zygotic paternal genome demethylation. Nature. 2010;463(7280):554–8.

116. Booker SJ, Grove TL. Mechanistic and functional versatility of radical SAM enzymes.

F1000 Biology Reports. 2010;2(1).

117. Yuan Y, Yuan H, Yang G, Yun H, Zhao M, Liu Z, et al. IFN-α confers epigenetic

regulation of HBV cccDNA minichromosome by modulating GCN5-mediated

succinylation of histone H3K79 to clear HBV cccDNA. Clinical Epigenetics.

2020;12(1):1–16.

118. Kollenstart L, De Groot AJL, Janssen GMC, Cheng X, Vreeken K, Martino F, et al.

Gcn5 and esa1 function as histone crotonyltransferases to regulate crotonylation-

dependent transcription. Journal of Biological Chemistry. 2019;294(52):20122–34.

119. Jones R. Identification and verification of lysine propionylation and butyrylation in

yeast core histones using PTMap software. Bone. 2014;23(1):1–7.

120. Liu B, Lin Y, Darwanto A, Song X, Xu G, Zhang K. Identification and characterization

of propionylation at histone H3 lysine 23 in mammalian cells. Journal of Biological

Chemistry. 2009;284(47):32288–95.

121. Papanicolaou K, O'Rourke B, Foster DB. Metabolism leaves its mark on the

powerhouse: recent progress in post-translational modifications of lysine in

mitochondria. Frontiers in Physiology. 2014;5 JUL.

122. Ekwall K. Genome-wide analysis of HDAC function. Trends in Genetics.

2005;21(11):608–15.

123. Bernstein BE, Tong JK, Schreiber SL. Genomewide studies of histone deacetylase

function in yeast. Proceedings of the National Academy of Sciences of the United

States of America. 2000;97(25):13708–13.

124. Honda S, Bicocca VT, Gessaman JD, Rountree MR, Yokoyama A, Yu EY, et al. Dual

chromatin recognition by the histone deacetylase complex HCHC is required for

proper DNA methylation in *Neurospora crassa*. Proceedings of the National Academy

of Sciences of the United States of America. 2016;113(41):E6135–44.

125.   Job G, Brugger C, Xu T, Lowe BR, Pfister Y, Qu C, et al. SHREC silences

heterochromatin via distinct remodeling and deacetylation modules. Molecular cell.

2016;62(2):207–21.

126.   Wu J, Suka N, Carlson M, Grunstein M. TUP1 utilizes histone H3/H2B-specific HDA1

deacetylase to repress gene activity in yeast. Molecular Cell. 2001;7(1):117–26.

127.   Kurdistani SK, Grunstein M. Histone acetylation and deacetylation in yeast. Nature

reviews Molecular cell biology. 2003;4(4):276–84.

128.   Li X, Pan L, Wang B, Pan L. The histone deacetylases HosA and HdaA affect the

phenotype and transcriptomic and metabolic profiles of *Aspergillus niger*. Toxins.

2019;11(9).

129.   Lee I, Oh JH, Keats Shwab E, Dagenais TRT, Andes D, Keller NP. HdaA, a class 2

histone deacetylase of *Aspergillus fumigatus*, affects germination and secondary

metabolite production. Fungal Genetics and Biology. 2009;46(10):782–90.

130.   Brosch G, Loidl P, Graessle S. Histone modifications and chromatin dynamics: A

focus on filamentous fungi. FEMS Microbiology Reviews. 2008;32(3):409–39.

131.   Carmen AA, Griffin PR, Calaycay JR, Rundlett SE, Suka Y, Grunstein M. Yeast HOS3

forms a novel trichostatin A-insensitive homodimer with intrinsic histone deacetylase

activity. Proceedings of the National Academy of Sciences of the United States of

America. 1999;96(22):12356–61.

132.   Pérez-Martínez ME, Benet M, Alepuz P, Tordera V. Nut1/Hos1 and Sas2/Rpd3

control the H3 acetylation of two different sets of osmotic stress-induced genes.

Epigenetics. 2020;15(3):251–71.

133.   Li S, Yue Z, Tanaka TU. Smc3 deacetylation by Hos1 facilitates efficient dissolution of

sister chromatid cohesion during early anaphase. Molecular Cell. 2017;68(3):605-

614.e4.

134.   Wang A, Kurdistani SK, Grunstein M. Requirement of Hos2 histone deacetylase for

gene activity in yeast. Science. 2002;298(5597):1412–4.

135.   Suka N, Suka Y, Carmen AA, Wu J, Grunstein M. Highly specific antibodies

determine histone acetylation site usage in yeast heterochromatin and euchromatin.

Molecular cell. 2001;8(2):473–9.

136.   De Nadal E, Zapater M, Alepuz PM, Sumoy L, Mas G, Posas F. The MAPK Hog1

recruits Rpd3 histone deacetylase to activate osmoresponsive genes. Nature.

2004;427(6972):370–4.

137.   Tribus M, Bauer I, Galehr J, Rieser G, Trojer P, Brosch G, et al. A novel motif in

fungal class 1 histone deacetylases is essential for growth and development of

Aspergillus. Molecular biology of the cell. 2010;21(2):345–53.

138.   Kawauchi M, Nishiura M, Iwashita K. Fungus-specific sirtuin HstD coordinates

secondary metabolism and development through control of LaeA. Eukaryotic Cell.

2013;12(8):1087–96.

139.   Klose RJ, Yamane K, Bae Y, Zhang D, Erdjument-Bromage H, Tempst P, et al. The

transcriptional repressor JHDM3A demethylates trimethyl histone H3 lysine 9 and

lysine 36. Nature. 2006;442(7100):312–6.

140.   Tamaru H. Confining euchromatin/heterochromatin territory: Jumonji crosses the line.

Genes and Development. 2010;24(14):1465–78.

141.   Isaac S, Walfridsson J, Zohar T, Lazar D, Kahan T, Ekwall K, et al. Interaction of epe1

with the heterochromatin assembly pathway in *Schizosaccharomyces pombe*.

Genetics. 2007;175(4):1549–60.

142.   Turberfield AH, Kondo T, Nakayama M, Koseki Y, King HW, Koseki H, et al. KDM2

proteins constrain transcription from CpG island gene promoters independently of

their histone demethylase activity. Nucleic acids research. 2019;47(17):9005–23.

143.   Qian S, Wang Y, Ma H, Zhang L. Expansion and functional divergence of Jumonji C-

containing histone demethylases: significance of duplications in ancestral

angiosperms and vertebrates. Plant Physiology. 2015;168(4):1321–37.

144. Hong SH, Cho YW, Yu LR, Yu H, Veenstra TD, Ge K. Identification of JmjC domain-

containing UTX and JMJD3 as histone H3 lysine 27 demethylases. Proceedings of

the National Academy of Sciences of the United States of America.

2007;104(47):18439–44.

145. Agger K, Cloos PAC, Christensen J, Pasini D, Rose S, Rappsilber J, et al. UTX and

JMJD3 are histone H3K27 demethylases involved in HOX gene regulation and

development. Nature. 2007;449(7163):731–4.

146. Dehé PM, Dichtl B, Schaft D, Roguev A, Pamblanco M, Lebrun R, et al. Protein

interactions within the Set1 complex and their roles in the regulation of histone 3

lysine 4 methylation. Journal of Biological Chemistry. 2006;281(46):35404–12.

147. Lewis ZA, Adhvaryu KK, Honda S, Shiver AL, Knip M, Sack R, et al. DNA methylation

and normal chromosome behavior in *Neurospora* depend on five components of a

histone methyltransferase complex, DCDC. PLoS Genetics. 2010;6(11):1–11.

148. Eberharter A, Sterner DE, Schieltz D, Hassan A, Yates JR, Berger SL, et al. The ADA

complex is a distinct histone acetyltransferase complex in *Saccharomyces cerevisiae*.

Molecular and Cellular Biology. 1999;19(10):6621–31.

149. Doyon Y, Côté J. The highly conserved and multifunctional NuA4 HAT complex.

Current Opinion in Genetics and Development. 2004;14(2):147–54.

150. Verzijlbergen KF, van Welsem T, Sie D, Lenstra TL, Turner DJ, Holstege FCP, et al.

A barcode screen for epigenetic regulators reveals a role for the NuB4/HAT-B histone

acetyltransferase complex in histone turnover. PLoS Genetics. 2011;7(10):18–24.

151. Sendra R, Tse C, Hansen JC. The yeast histone acetyltransferase A2 complex, but

not free Gcn5p, binds stably to nucleosomal arrays. Journal of Biological Chemistry.

2000;275(32):24928–34.

152. Grant PA, Duggan L, Côté J, Roberts SM, Brownell JE, Candau R, et al. Yeast Gcn5

functions in two multisubunit complexes to acetylate nucleosomal histones:

Characterization of an ada complex and the saga (spt/ada) complex. Genes and

Development. 1997;11(13):1640–50.

153.   Erlendson AA, Friedman S, Freitag M. A matter of scale and dimensions: chromatin of

chromosome landmarks in the fungi. Microbiology spectrum. 2017;5(4):4–5.

154.   Buscaino A. Chromatin-mediated regulation of genome plasticity in human fungal

pathogens. Genes. 2019;10(11).

155.   Steenwyk JL, Mead ME, Knowles SL, Raja HA, Roberts CD, Bader O, et al. Variation

among biosynthetic gene clusters, secondary metabolite profiles, and cards of

virulence across *Aspergillus* species. Genetics. 2020;216(2):481–97.

1209

Figure captions

Figure 1. Conservation of DNA methyltransferase families in the *Aspergillus* genus and outgroup species. The maximum-likelihood phylogeny of DNA methyltransferase domain (PF00145) found in 94 Aspergilli and 15 outgroup species was determined using IQ-TREE (37,38). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT values over 80. Leaves from Aspergilli with trustworthy support are collapsed and shown as red triangles. The numbers of hits and species are separated by the pipe in the '#Homologs | #Species' column. The number in parentheses after * indicates the additional hits found by the TBLASTN search on NCBI. Additional conserved domains in DNA methyltransferase domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

Figure 2. Conservation of histone methyltransferase families in the *Aspergillus* genus and outgroup species. The maximum-likelihood phylogeny of SET domains (PF00856) found in 94 Aspergilli and 15 outgroup species was determined using IQ-TREE (37,38). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT values over 80. Leaves from Aspergilli with trustworthy support are collapsed and shown as red triangles. Leaves with similar protein architecture but unknown functions covering both Aspergilli and outgroup species are collapsed and shown as grey triangles. The numbers of hits and species are separated by the pipe in the '#Homologs | #Species' column. The number in parentheses after * indicates the additional hits found by the TBLASTN search on NCBI. Additional conserved domains in SET domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

Figure 3. Conservation of MOZ/SAS FAMILY domain-containing histone acetyltransferases in the *Aspergillus* genus and outgroup species. The maximum-likelihood phylogeny of MOZ/SAS FAMILY domains (PF01853) found in 94 Aspergilli and 15 outgroup species was determined using IQ-TREE (37,38). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT values over 80. Leaves from Aspergilli with trustworthy support are collapsed and shown as red triangles. The numbers of hits and species are separated by the pipe in the '#Homologs | #Species' column. '*In vitro*' indicates the acetylation activity of this enzyme was only observed on the substrate *in vitro*. Additional conserved domains in MOZ/SAS FAMILY domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

Figure 4. Conservation of histone deacetylase domain-containing proteins in the *Aspergillus* genus and outgroup species. The maximum-likelihood phylogeny of Histone deacetylase domains (PF00850) found in 94 Aspergilli and 15 outgroup species was determined using IQ-TREE (37,38). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT values over 80. Leaves from Aspergilli with trustworthy support are collapsed and shown as red triangles. The numbers of hits and species are separated by the pipe in the '#Homologs | #Species' column. '*In vitro*' indicates the deacetylation activity of this enzyme was only observed on the substrate *in vitro*. Additional conserved domains in histone deacetylase domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

Figure 5. Presence-absence patterns of subunits that constitute characterized histone modification complexes. The top row with distinct color boxes indicates outgroup species and Aspergilli sections, the order of the species from right to left is the same as in the species tree from top to bottom (Supplementary Figure 1) without *Fusarium oxysporum* and *Aspergillus kawachii* since they were excluded in the later analysis. Each row represents a subunit in the histone modification complexes, the order agrees with the summary list (Supplementary Table 4). Each row shows the presence/absence of a specific complex subunit. For each complex, the first row

1259 represents the catalytic subunit. Each box in the matrix indicates the presence (dark color) or
1260 absence (light color) of a subunit (Supplementary Table 4). The brown boxes indicate additional
1261 hits found with TBLASTN search on NCBI (Supplementary Table 5).
1262
Figure 6. Mass spectrometry unlocks the histone code in Aspergilli. A. Schematic
presentation of the in-gel digestion method used to process histones isolated in *A. nidulans*. After
SDS-PAGE separation, gel bands corresponding to histone H3 and H4 are excised and chemically
acylated with propionic anhydride (which derivatizes unmodified and mono-methylated lysines).
Thanks to the derivatization, trypsin cutting at lysines is impaired, and histones are cleaved only
at the C-terminus of arginine. Finally, digested peptides are N-terminally derivatized with phenyl
isocyanate (PIC). **B.** *A. nidulans* histone H3 and H4 sequences. Cleavage sites after PRO-PIC
digestion are shown. Peptides identified in this study are highlighted in light red and labeled using
their starting-ending amino acids. **C.** Relative abundance of post-translational modifications
detected on peptides of histone H3 and H4 from *A. nidulans* using unbiased mass spectrometry
analysis. unmod: unmodified peptide; me1: monomethylation; me2: dimethylation; me3:
trimethylation; ac: acetylation. The term "combinations" refers to multiple modifications occurring
on the same peptide (Supplementary Table 6). **D.** Histone modifications are predicted by genomic
and phylogenetic analyses, and/or proved by mass spectrometry in *A. nidulans*. Histone
modifications verified by mass spectrometry are labeled as solid dots, while the ones that are only
predicted by phylogenic analysis are labeled as solid dots with a red outline.

Supplementary materials

Supplementary Figure 1. A robust species phylogeny for the fungal genus *Aspergillus*. The species phylogeny was based on a maximum-likelihood phylogeny reconstructed by IQ-TREE with partitioned analysis using 758 BUSCO (Benchmarking Universal Single-Copy Orthologs) genes. The dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT bootstraps values over 80. Species from the same section are labeled by background color with a section name. Two species, *Fusarium oxysporum* and *Aspergillus kawachii*, are included in the species tree, but we don't use these two in the following analyses.

Supplementary Figure 2. Two known histone acyltransferase-domain containing protein groups are conserved in the *Aspergillus* genus and outgroup species together with many unknown proteins. Maximum likelihood phylogeny of Histone acyltransferase domains (PF00583) found in 94 Aspergilli and 15 outgroup species were determined using IQ-TREE (Chernomor et al. 2016; Nguyen et al. 2015). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT bootstraps values over 80. Additional conserved domains in histone acyltransferase domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

Supplementary Figure 3. Five histone demethylase-domain containing protein groups are conserved in *Aspergillus* genus and outgroup species while KDM6B/JMJD3 is lost. Maximum likelihood phylogeny of Jumonji C (JmjC) domain (PF02373) found in 94 Aspergilli and 15 outgroup species were determined using IQ-TREE (Chernomor et al. 2016; Nguyen et al. 2015). The black dot on the branch indicates ultrafast bootstrap values over 95 and SH-aLRT bootstraps values over 80. Additional conserved domains in JmjC domain-containing proteins were identified using the PFAM database (https://pfam.xfam.org).

Supplementary Table 1. List of species included in this study.

Supplementary Table 2. List of BUSCO genes used to build the species tree.

Supplementary Table 3. List of sequence names in the domain trees.

Supplementary Table 4. List of histone modifier complexes reported in the literature.

Supplementary Table 5. Identification of missing chromatin modifiers in the outgroup species using TBLASTN search

Supplementary Table 6. Mass spectrometry result for *A. nidulans*.

Supplementary Data 1. Fasta, trimmed alignment, and maximum-likelihood phylogenetic tree files associated with Figure 1-4 and Supplementary Figure 2-3.

Supplementary Data 2. Fasta, trimmed alignment, and maximum-likelihood phylogenetic tree files associated with Figure 5.