Structural basis for regulation of a tripartite toxin-antitoxin system by dual phosphorylation

René L. Bærentsen[1]*, Stine Vang Nielsen[2]*, Jeppe Lyngsø[3], Francesco Bisiak[1], Jan Skov Pedersen[3],

Kenn Gerdes[4], Michael A. Sørensen[2]†, Ditlev. E. Brodersen[1]†

[1]Department of Molecular Biology and Genetics, Aarhus University, Gustav Wieds Vej 10c, DK-8000 Aarhus C, Denmark, [2]Department of Biology, University of Copenhagen, Ole Maaløes Vej 5, DK-2200 København N, [3]Department of Chemistry and Interdisciplinary Nanoscience Centre (iNANO), Gustav Wieds Vej 14, DK-8000 Aarhus C, [4]Voldmestergade 8, DK-2100 København Ø, Denmark.

*These authors contributed equally.

†To whom correspondence should be addressed (mas@bio.ku.dk, phone +45 35323711 or deb@mbg.au.dk, phone +45 21669001).

Keywords: HipBA; STK; Ser/Thr kinase; TrpS; tRNA synthetase

Running title: Regulation of a tripartite TA system

ABSTRACT

Many bacteria encode multiple toxin-antitoxin (TA) systems targeting separate, but closely related,

cellular functions. The toxin of the *E. coli hipBA* system, HipA, is a kinase that inhibits inhibits

translation via phosphorylation of glutamyl-tRNA synthetase. Enteropathogenic *E. coli* (EPEC)

O127:H6 encodes an additional, tripartite TA module, *hipBST*, for which the HipT toxin was shown

to specifically target tryptophanyl-tRNA synthetase, TrpS. Surprisingly, the function as antitoxin has

been taken over by the third protein, HipS, but the molecular details of how activity of HipT is

controlled remain poorly understood. Here, we show that HipBST is markedly different from HipBA

and that the unique HipS protein, which is homologous to the N-terminal subdomain of HipA, has

evolved to function as antitoxin by breaking the kinase active site. We also show how auto-

phosphorylation at two conserved sites in the kinase toxin serve to dually regulate binding of HipS

and kinase activity. Finally, we demonstrate that the HipBST complex is dynamic and present a

cohesive model for the regulation and activation of this type of three-component system.

INTRODUCTION

Bacteria employ a wide range of mechanisms adapt to changing environments and imminent threats such as antibiotics and phage attack, including activation of toxin-antitoxin (TA) systems that can release intracellular toxins to rapidly alter metabolism or reduce growth (Harms et al., 2018). Canonical (type II) TA systems are small, bicistronic loci that encode a protein toxin and its cognate antidote (antitoxin) that tightly interact to form a higher-order complex in the inactive state, usually capable of controlling transcription via a DNA-binding domain on the antitoxin (Harms et al., 2018). There are also examples of tricistronic TA loci, in which the third component is either a chaperone required for folding of the antitoxin and thus, toxin inhibition (Bordes et al., 2016) or an additional, transcriptional regulator (Hallez et al., 2010; Zielenkiewicz & Ceglowski, 2005). A striking observation is that some microorganisms contain a very large number of similar, paralogous TA systems, such as the human pathogen, *Mycobacterium tuberculosis*, which contains at least 47 paralogous TA systems of the *vapBC*-type, for which the VapC toxin appears to target unique non-coding RNA species (Sharrock et al., 2018; Winther et al., 2016). Likewise, *Salmonella enterica* encodes several paralogous *tacTA* systems in which the TacT toxins function by acetylating the primary amine of amino acids of charged transfer RNA molecules (Grabe et al., 2021). These cases raise important questions about the evolutionary benefit of harbouring multiple, highly similar toxin systems as well as how cross-reactivity is prevented.

In the widespread and diverse *hipBA* (high persister) system, the HipA toxin is a serine-threonine kinase (STKs) (Hanks et al., 1988; Stancik et al., 2018), while the cognate antitoxin, HipB, contains a helix-turn-helix (HTH) motif and inhibits the toxin by a mechanism which is not fully understood (Gerdes et al., 2021; Schumacher et al., 2009). In the most well-studied and canonical *hipBA* system in *Escherichia coli* K-12, the toxin (HipA$_{Ec}$) specifically targets glutamyl-tRNA synthetase (GltX)

by phosphorylation of an ATP-binding motif conserved in type I aminoacyl-tRNA synthetases (Eriani

et al., 1990; Sekine et al., 2003), thereby inhibiting its activity and blocking translation (Germain et

al., 2013). Accumulation of uncharged tRNA^{Glt} subsequently induces the stringent response via RelA-

mediated (p)ppGpp synthesis on starved ribosomes (Haseltine & Block, 1973; Pacios et al., 2020;

Winther et al., 2018). In contrast to most type II TA systems, the cognate antitoxin, HipB does not

directly interact with the HipA active site and inhibition has therefore been proposed to occur by

several mechanisms, including blocking conformational changes in the kinase required for catalysis,

sequestration of HipA on DNA via the HipB DNA binding domain (Schumacher et al., 2009), or

allosterically via placement of a C-terminal Trp residue of HipB into a pocket on HipA (Evdokimov

et al., 2009). In an additional layer of complexity in regulation, HipA$_{Ec}$ is regulated by *trans*

autophosphorylation at a conserved serine (Ser150) situated in a loop (the "Gly-rich loop") near the

active site, which is required for ATP binding and functionally similar to the P loop in eukaryotic

kinases (Huse & Kuriyan, 2002; Schumacher et al., 2012). Phosphorylation at this site causes a

conformational change in which the loop is ejected from its burrowed position inside the active site,

rendering the kinase unable to bind ATP (Schumacher et al., 2012).

Enteropathogenic *E. coli* (EPEC) O127:H6, an important diarrheal pathogen of young children,

contains, in addition to the canonical *hipBA* locus, a homologous tricistronic TA system, *hipBST*,

encoding three separate proteins. HipB, by analogy to HipBA, contains a putative DNA-binding HTH

domain while HipT is a kinase like HipA and functions as toxin by phosphorylating and inactivating

tryptophanyl-tRNA, TrpS (Gerdes et al., 2021; Vang Nielsen et al., 2019). Surprisingly, the third

protein, HipS, which at the sequence level corresponds to the N-terminal subdomain 1 of the larger

HipA kinase, was shown to function as antitoxin by an unknown mechanism (Vang Nielsen et al.,

2019). Moreover, HipT was shown to be subject to autophosphorylation at two conserved serine

positions in the Gly-rich loop, but the functional consequences of this are also not known. Here, we

show that the HipBST complex fom pathogenic *E. coli* is markedly different from HipBA and that

inhibition of HipT is achieved through direct interaction with a conserved residue in HipS. Two

structures of HipT in separate, phosphorylated states further demonstrate that HipT, is found in an

inactive conformation regardless of phosphorylation state inside the HipBST complex. Finally, we

show, using phosphomimetic kinase variants *in vivo* and analysis of phosphorylation patterns and

flexibility *in vitro*, that the two phosphorylation positions control HipT activity and binding of

antitoxin (HipS), and that the HipBST complex is dynamic. Together, our data provide a mechanistic

understanding of how the HipS protein can function as antitoxin and suggests that the corresponding

domain in HipA can have important, regulatory roles as well.

RESULTS

Dual autophosphorylation sites are conserved in HipT kinases

One of the most surprising features of the HipBST system is that HipS shows sequence similarity to

the N-terminal subdomain 1 of HipA, the toxin of the *hipBA* system, while simultaneously

functioning as antitoxin (**Figure S1**) (Vang Nielsen et al., 2019). Moreover, phylogenetic analysis

has demonstrated that HipT and HipA toxins are relatively closely related within a very broad

superfamily of HipA-homologous kinases (Gerdes et al., 2021). To gain further insight into the basis

of this important difference, we initially performed a focused phylogenetic analysis of the 48 HipT

orthologues identified so far among bacterial genomes (Gerdes et al., 2021). Alignment and clustering

reveals a largely bifurcated, phylogenetic tree for which one large branch (the "SIS group") contains

HipT kinases with two serine residues in close proximity in the Gly-rich loop corresponding to the

known autophosphorylation sites in HipT of *E. coli* O127:H6 (**Figure 1a**). In the remainder of the

HipT orthologues, this region contains variable Ser/Thr motifs, including TxT, Sxx, xxT, SxT, and

Figure 1. Phylogenetic analysis of HipT. a. Phylogenetic guide tree of 48 HipT orthologues with sequences motifs (potential phosphorylation sites) indicated on the side. The "SIS group" (red) is by far the largest group followed by the T[IV][TP] group (green), and SIQ group (blue). Sequences used in the alignment in **b** are shown with red letters. Numbers on the branches indicate bootstrap confidence levels. **b.** Sequences of the Gly-rich loop (orange background), including the potential phosphorylation sites for selected HipT orthologues compared to HipA from *E. coli* K-12. Known phosphorylation sites in *E. coli* O127:H6 HipT (top) and *E. coli* K-12 HipA (bottom) are indicated with arrows and conserved sequence motifs with bold white text. **c.** Sequence logo for the Gly-rich loop derived from all 48 HipT sequences. **d.** Consensus motif for the HipT Gly-rich loop with known phosphorylation sites in red. Φ indicates a hydrophobic residue, while Ψ are aliphatic residues.

Txx. We note that the single position between the two Ser/Thr residues invariably is occupied by a

small hydrophobic residue (I, L, or V) and that a small subbranch close to the SIS group represents

an interesting variation with a Gln in the second autophosphorylation position (the "SIQ" group). The

remainder of the Gly-rich loop across a range of HipT kinases is highly similar to HipA$_{Ec}$, which has

the sequence I$\underline{S}$V$\underline{A}$GAQEK (**Figure 1b-c**). We can thus derive a consensus motif for the HipT Gly-

rich loop, which includes several conserved hydrophobic positions in addition to the two Ser/Thr

residues (**Figure 1d**). In summary, phylogenetic analysis reveals that a large fraction of HipT kinases

contain a conserved SIS motif with two potential phosphorylation positions in their Gly-rich loops

but also that this region displays some sequence diversity, which may point to subtle differences in

regulation.

HipBST forms a hexameric higher-order complex

Comparison of the HipBST and HipBA sequences reveals that HipS shows similarity to the N-

terminal subdomain 1 of HipA (**Figure S1 and S2a**) and that both HipB and HipT of *hipBST* contain

N-terminal extensions of around 35 residues not found in HipB and HipA of *hipBA*, respectively

(**Figure 2a and S1**). To understand the molecular basis for the functional and structural differences,

we determined the crystal structure of HipBST from enteropathogenic *E. coli* O127:H6 expressed

with inactive HipT (D233Q) to 2.9 Å by molecular replacement using HipA$_{Ec}$ as search model

(**Figure 2b**). The refined structure (R=19.5% / R$_{free}$=22.8%) has two copies of each of the three

proteins in the asymmetric unit and is generally well-defined for HipB and HipT, while HipS appears

to be more flexible. Analysis by the protein interaction server, PISA (Krissinel & Henrick, 2007),

suggests that the biological assembly is comprised of a dimer of HipBST heterotrimers, which are

held together through dimerisation of HipB (**Figure 2b**). This is consistent with the observation that

the HTH domain is usually found as a dimer in DNA-binding proteins (Brennan & Matthews, 1989).

Figure 2. Crystal structure of the *E. coli* O127:H6 HipBST complex. a. Schematic representation of *E. coli*
O127:H6 HipBST and canonical *E. coli* HipBA showing corresponding proteins and domains. Dashed lines indicate
similar domains and grey areas represent regions missing domains in the crystal structures. HipS (beige) is
structurally similar to the N-subdomain 1 of HipA (dark purple), HipT has an additional N-terminal mini-domain
not found in HipA (light blue) while HipA has a short linker between the two N-subdomains (light purple). **b.** *E.
coli* O127:H6 HipBST forms a hetero-hexameric complex with HipS (beige) on top of HipT (blue) through
dimerisation of HipB (green) also generating a helix-turn-helix DNA-binding motif (HTH, dashed box). **c.** Crystal
structure of the *E. coli* K-12 HipBA complex (PDB: 2WIU) shown as cartoon with HipA in blue purple (N-terminal
subdomain 1) and purple and the HipB homodimer in two shades of green (Evdokimov et al., 2009). **d.** The linker
between N-subdomain 1 and 2 (dark purple) physically linking the two domains in HipA$_{Ec}$. **e.** The N-terminal mini-
domain of HipT (blue), which is absent from HipA.

In this complex, the two HipB monomers interact strongly (interface area = 1,405 Å^2) while the

interface between HipB and HipT is relatively small (513 Å^2). The biological relevance of the hetero-

hexamer is supported by analytical gel filtration (**Figure S2b**) and small-angle X-ray scattering

(SAXS, **Figure S6**), the latter of which showed a major species with a total mass of 129 ± 8 kDa, in

good agreement with the theoretical mass of 125.6 kDa. HipB consists of an α-helical bundle of four

helices including the predicted DNA-binding HTH motif (residues 62-83), which forms a canonical

HTH DNA-binding domain exposed at the bottom of the hetero-hexamer (**Figure 2b**). The helical

bundle is followed by a small β strand that forms an antiparallel sheet through pairing with the

corresponding region in the neighbouring molecule of HipB. The N-terminal extension of HipB (36

residues) is not visible in the electron density, but secondary structure prediction indicates that it

forms a long α-helix (**Figure S1a**). HipS is closely associated with HipT and consists of a solvent-

facing, five-stranded, anti-parallel β sheet of which four strands are located in the N terminus

(residues 1-52). These are followed by a small domain of three helices (residues 53-94) that forms

the interface to HipT followed by the fifth β strand (residues 95-102, **Figure 2b and S3a**).

The overall architecture of the HipBST complex consists of two HipST toxin:antitoxin complexes

separated in space by the dimer of the HipB and in this way differs markedly different from HipBA$_{Ec}$

in which two HipA toxins are closely packed head-to-tail around two HipB antitoxins (**Figure 2c**).

This configuration is key to the proposed way in which HipB can function as antitoxin, namely by

restricting the movement of the N-terminal subdomain of HipA (**Figure 2c**, blue) (Schumacher et al.,

2009). As suggested from sequence analysis, HipS bears a striking structural resemblance to the N-

terminal subdomain 1 of HipA (Schumacher et al., 2012), both with respect to its overall fold and

placement in relation to HipT. As a result of this, the HipS C-terminus overlaps with a extended linker

that bridges N-subdomain 1 to N-subdomain 2 in HipA (**Figure 2d**). HipT corresponds structurally

to the core kinase fold of HipA, except for the presence of an additional N-terminal mini-domain

(residues 1-41), the role of which is possibly to anchor N-subdomain 2 (residues 42-140) to the core

of the kinase in the absence of the physical coupling to HipS **(Figure 2a and 2e)**. Interestingly, the

overall structure of the HipBST complex is reminiscent of the structure of a HipBA complex from

Shewanella oneidensis, which was crystallised in complex with DNA **(Figure S2c)** (Wen et al.,

2014). This suggests that the open conformation is not a specific result the tripartite configuration

and superpositioning of the DNA-binding domains (**Figure S3b**) and analysis of potential interactions

with DNA (**Figure S3c**) suggests HipBST could bind DNA in a similar way.

HipT has an inactive conformation inside the HipBST complex

For HipA$_{Ec}$, two discrete active site conformations are observed depending on phosphorylation state

and whether substrate (ATP) is bound or not (Schumacher et al., 2015; Schumacher et al., 2012;

Schumacher et al., 2009). In its non-phosphorylated form and in the presence of ATP, the flexible

Gly-rich loop is found in an inward conformation in which main chain amino groups on Ala154 and

Gln155 coordinate the ATP phosphate groups (**Figure 3a**) (Schumacher et al., 2009).

Phosphorylation of Ser150, however, causes ejection of the loop through strong interactions of the

phosphate group with active sites residues, which prevents binding of ATP (**Figure 3b**) (Schumacher

et al., 2012). In the HipBST crystal structures, we find the Gly-rich loop of HipT forms a short α-

helix and is in a conformation most similar to the ejected form of the HipA$_{Ec}$ loop despite HipT not

being phosphorylated (**Figure 3c**). In this conformation, Ser59, which is known to undergo

autophosphorylation in HipT, forms interactions to several active site residues, including Asp210 and

the Ser57 auto-phosphorylation site (Vang Nielsen et al., 2019). We conclude that inside the HipBST

complex, HipT in its non-phosphorylated state is maintained in an inactive conformation that does

not support ATP binding.

Figure 3. Trp65 is essential for HipS to function as antitoxin. a. Overview of the HipA active site, bound to ATP

(green, PDB: 3DNT) (Schumacher et al., 2009). Residues interacting with the nucleotide are highlighted as well as

the Gly-rich loop (residues 150-156, green) including Ser150. D309Q is a mutated active site residue. **b.** The active

site of HipA phosphorylated on Ser 150 (green/orange, PDB: 3TPE) with the Gly-rich loop (green) in the outward-

facing conformation (Schumacher et al., 2012). **c.** The active site of HipBS-bound HipT^{D233Q} with the Gly-rich loop

(residues 58-63, ruby) in the outward-facing conformation. D233Q is the mutated active site residue. **d.** Overview

and detailed interactions between HipS (beige) and HipT (blue) with its Gly-rich loop (ruby) and interacting residues

indicated. **e.** Top, HipS Trp65 (beige) is placed in a pocket on the surface of HipT (blue); Bottom, the corresponding

residue (Pro69) in HipA (purple, PDB: 3TPD). **f.** *E. coli* MG1655 harbouring empty pBAD33 vector ("pBAD33")

or pSVN1 (pBAD33::*hipT*) in combination with empty pNDM220 vector ("pNDM220"), pSVN109

(pNDM220::*hipS*), or pSVN178 (pNDM220::*hipS*^{W65A}) as indicated. Plates contained 0.2% glucose (to repress

hipT), 0.2% arabinose (to induce *hipT*), or 0.2% arabinose plus 200 µM IPTG (to induce *hipS*, or *hipS*^{W65A}). The

plates are representative of at least three independent experiments.

HipS inhibits HipT through insertion of a hydrophobic residue near the active site

To understand how HipS, which is homologous to the N-terminal subdomain 1 of HipA, can function as the antitoxin in the HipBST system, we then focused on the specific interactions found between HipS and HipT. There are three major areas of contact between the two proteins with several strong interactions on both sides of the kinase domain, involving both hydrogen bonds and hydrophobic and charged interactions (**Figure 3d**). Near the Gly-rich loop, HipS Glu63 points directly into the HipT active site while Trp65 is wedged into a deep cavity formed by the HipT Gly-rich loop in its outward conformation. While Glu63 is conserved in the N-terminal subdomain of HipA$_{Ec}$ (Asp67) and therefore less likely contribute to functional differences between the two systems, the Trp65 motif (**Figure 3e, top**) is structurally different from the corresponding interaction in HipA$_{Ec}$, which has the helix (HipS α3, **Figure S2a**) in a different orientation and a proline residue at this position (**Figure 3e, bottom**). Moreover, Trp65 is conserved in many HipS orthologues (**Figure S2a**). We hypothesised that placement of the bulky Trp residue inside HipT might block transition of the Gly-rich loop from the outward to the inward conformation, likely required for ATP binding. To test this, we mutated the Trp65 in HipS to alanine and tested the ability of the antitoxin to antagonise the kinase in *E. coli* K-12 on both solid media and in liquid culture (**Figure 3f and S3d**). In both experiments, absence of the bulky Trp65 side chain strongly reduced the antitoxin activity of HipS, confirming that this interaction is critical to inhibition of HipT. We conclude that Trp65 is essential for the ability of HipS to inhibit HipT, presumably either by increasing the affinity of HipS for HipT, or by preventing the Gly-rich loop from adopting the conformation required for ATP binding.

The conformation of HipT in the HipBST complex is independent of phosphorylation

In contrast to HipA$_{Ec}$, which undergoes autophosphorylation at a single site (Ser150), HipT was found to harbour two sites in its Gly-rich loop (Ser57 and Ser59), which are subject to autophosphorylation

individually but not concomitantly (Vang Nielsen et al., 2019). To understand the individual roles of

the two autophosphorylation positions, we next determined crystal structures of two HipBST

complexes with HipT serine residues mutated to alanine individually: HipBST^{S57A} (2.4 Å, **Table S1**)

and HipBST^{S59A} (3.4 Å). Inspection of the electron densities near the active site revealed that in both

cases, the remaining serine residue (i.e. Ser57 in HipBST^{S59A} and Ser59 in HipBST^{S57A}) had become

phosphorylated prior to crystallisation (**Figure 4a and S4a-b).** For HipBST^{S59A}, phosphorylation of

Ser57 was incomplete and could only be modelled in one of the two HipT molecules in the

asymmetric unit (**Figure S4b).** The phosphate group of Ser59 (P-Ser59) forms strong interactions to

Lys161, His212, and the catalytic Asp210 inside the active site (**Figure 4a, top**), while P-Ser57 is

located further away from the active site and forms hydrogen bonds to Tyr162 and Asp210 (**Figure**

4a, bottom). In both structures, however, the Gly-rich loop maintains its outward conformation,

incompatible with ATP binding, suggesting that HipT is inactive. The phosphate group on Ser59

occupies a position very similar to phosphorylated Ser150 in HipA$_{Ec}$, despite the fact that this residue

aligns to Ser57 at the sequence level (**Figure S4c, top and 1b**). This position structurally overlaps

with the γ phosphate of ATP in the structure of ATP-bound HipA$_{Ec}$, suggesting that phosphorylation

prevents nucleotide binding in the kinase active site (**Figure 4b**). Intriguingly, the phosphoryl group

of P-Ser57 overlaps structurally with that of phosphorylated Ser147 in HipA$_{So}$, which corresponds to

HipA$_{Ec}$ Ser150 at the sequence level (**Figure S4c, bottom**). In order to confirm the phosphorylation

state of the HipT variants *in vitro*, we analysed purified HipBST complexes on Phos-tag SDS-PAGE

gels, which contain functional molecules that cause retention of phosphorylated protein species in the

gel (**Figure 4c, top**). Here, we observed a marked difference in migration between HipT from

wildtype HipBST, HipBST^{S59A}, and HipBST^{S57A} confirming that HipT^{S57A} is predominantly

phosphorylated while HipT^{S59A} is only partially modified, consistent with the crystal structures.

Surprisingly, wildtype HipBST complex is completely non-phosphorylated when purified in this way,

Figure 4. The phosphoserine positions in HipT have distinct functional roles. a. Overview of the HipT kinase active site in the D233Q mutant as well as S57A (top) and S59A (bottom) structures. The phosphate groups on Ser57 (in S59A) and Ser59 (in S57A) are shown in orange and relevant nearby residues are highlighted. **b.** Close-up of the HipT active site as observed for the S57A mutant with relevant active site residues shown as sticks. The Gly-rich loop is shown in red with pSer59 indicated. Overlaid is ATP and 2 Mg^{2+} ions derived from the structure of HipA (PDB: 3DNT, salmon/green) (Schumacher et al., 2009). **c.** HipT bands on Phos-tag gels (stained by Coomassie Blue) of purified HipBST complex before (-) and after (+) a Heparin column step to separate various complex species, as well as before (-) and after (+) treatment with 10 mM ATP, as indicated. Phosphorylated ("P-

HipT") and non-phosphorylated ("HipT") protein species are indicated. The gels are representative of two independent experiments. **d.** Top, growth curves measured by OD_{600} of *E. coli* MG1655 harbouring single autophosphorylation mutants pSVN194 (pBAD33::*hipT*^{S57D}, "*hipT* (DIS)"), pSVN195 (pBAD33::*hipT*^{S59D}, "*hipT* (SID)"), pSVN199 (pBAD33::*hipT*^{S57A}, "*hipT* (AIS)"), pSVN201 (pBAD33::*hipT*^{S59A}, "*hipT* (SIA)"), wildtype HipT pSVN1 (pBAD33::*hipT*, "*hipT* (SIS, wt)"), or the empty pBAD33 vector, as indicated, in combination with pSVN109 (pNDM220::*hipS*). *hipT* (ara) and *hipS* (IPTG) expression were induced at the indicated time points. The curves show the mean values of results from at least two independent experiments with error bars indicating standard deviations (hidden when small); Bottom, growth curves of *E. coli* MG1655 harbouring double autophosphorylation mutants pSVN203 (pBAD33::*hipT*$^{S57A+S59A}$, "*hipT* (AIA)"), pSVN205 (pBAD33::*hipT*$^{S57D+S59A}$, "*hipT* (DIA)"), pSVN207 (pBAD33::*hipT*$^{S57A+S59D}$, "*hipT* (AID)"), or pSVN209 (pBAD33::*hipT*$^{S57D+S59D}$, "*hipT* (DID)"), or the empty pBAD33 vector, as indicated, in combination with pSVN109 (pNDM220::*hipS*). *hipT* (ara) and *hipS* (IPTG) expression were induced at the indicated time points. The curves show the mean values of results from at least two independent experiments with error bars indicating standard deviations (hidden when small). **e.** Northern blot analysis probing against tRNA^{trpT} based on RNA extracted from *E. coli* MG1655 harbouring pSVN110 (pNDM220::*hipB-S*) in combination with pSVN1 (pBAD33::*hipT*, "*hipT* (S^{57}IS^{59}, *wt*)"), pSVN194 (pBAD33::*hipT*^{S57D}, "*hipT* (D^{57}IS^{59})"), pSVN195 (pBAD33::*hipT*^{S59D}, "*hipT* (S^{57}ID^{59})"), pSVN209 (pBAD33::*hipT*$^{S57D+S59D}$, "*hipT* (D^{57}ID^{59})") or pSVN205 (pBAD33::*hipT*$^{S57D+S59A}$, "*hipT* (D^{57}IA^{59})") at indicated time points in minutes before (-) and after (+) addition of arabinose (0.2%) to induce wildtype *hipT* or autophosphorylation mutants. Quantification of the bands (represented as % deacylated tRNA) is shown below the gel. The control ("ctrl") is chemically deacylated tRNA. The percentage of deacylated tRNA (bottom) was based on quantification of total tRNA^{trpT}. Results from strains harbouring pSVN1 (pBAD33::*hipT*), pSVN194 (pBAD33::*hipT*^{S57D}), pSVN209 and (pBAD33::*hipT*$^{S57D+S59D}$) are representative of two independent experiments, whereas results from strains harbouring pSVN195 (pBAD33::*hipT*^{S59D}) and pSVN205 (pBAD33::*hipT*$^{S57D+S59A}$) are from single experiments.

but incubation with ATP for 30 minutes leads to a visible increase in phosphorylation of wildtype

HipT while no further phosphorylation seemed to take place for the mutants upon incubation with

ATP (**Figure 4c, bottom**). In summary, we conclude that the conformation of the HipT active site

loop (Gly-rich loop) inside the HipBST complex is independent of the autophosphorylation state.

The two phosphoserine positions in HipT have separate functional roles

We next sought to understand the functional implications of autophosphorylation of HipT at the two

positions. To this end, we designed a set of HipT mutants with either Ala (to prevent

autophosphorylation) or Asp (to mimic the phosphorylated state) in one or both of the two Ser

positions. Liquid culture assays showed that both wildtype HipT $S^{57}IS^{59}$ and HipT $S^{57}ID^{59}$ caused

growth inhibition upon overexpression in *E. coli* K-12 and could be repressed by concomitant

expression of HipS (**Figure 4d, top**) (Vang Nielsen et al., 2019). In contrast to this, growth inhibition

by the HipT $D^{57}IS^{59}$ variant could not be repressed upon induction of HipS. Moreover, mutation to

alanine at either position caused very little ($S^{57}IA^{59}$) or no ($A^{57}IS^{59}$) growth inhibition, suggesting that

the serine residues or their phosphorylated counterparts are important for HipT activity on the target

TrpS (**Figure 4d, top**). To rule out any effects of autophosphorylation of the non-mutated Ser

position, we created double mutants for which both serine residues were mutated into either alanine

or aspartate (**Figure 4d, bottom**). Of these, HipT $D^{57}ID^{59}$ causes immediate growth inhibition while

HipT $D^{57}IA^{59}$ shows a similar but delayed phenotype, and consistent with the observation above,

neither could be repressed by HipS induction. And as before, placing alanine at either position either

completely abolished ($A^{57}IA^{59}$ and $A^{57}ID^{59}$) or strongly delayed ($D^{57}IA^{59}$) growth inhibition. To rule

out any effects due to the fact that HipB is missing in this setup, we finally repeated the experiments

with concomitant induction of both HipB and HipS and obtained very similar results (**Figure S4d**

and S4e).

To confirm that the growth inhibition observed was in fact due to phosphorylation and inactivation of TrpS, we took samples from *E. coli* cultures both before and at several time points after induction of wildtype HipT as well as all variants that caused markedly growth inhibition, extracted total RNA, and performed Northern blotting using probes against tRNA^{trpT} (**Figure 4e**) as well as tRNA^{gltTUVW} and tRNA^{argVYZQ} (**Figure S4f**) as controls. Here, charging of the tRNA is visible as a shift of the probed RNA band and thus allows direct assessment of the activity of TrpS *in vivo* in the presence of the various HipT variants. Induction of all variants caused accumulation of deacylated tRNA^{trpT} (**Figure 4e**) but no accumulation of the two control tRNAs (**Figure S4f**), which strongly suggests that the observed growth effects are due to reduction in the cellular pool of acylated tRNA. Taken together, we conclude that HipT autophosphorylation at Ser57 prevents HipS inhibition and does not prevent phosphorylation of the target, TrpS. On the other hand, phosphorylation at Ser59 does not affect HipS inhibition, indicating that phosphorylation at the two positions serve separate, functional roles.

hipB is both required and sufficient for transcriptional repression

To understand the functional role of HipB, specifically how DNA binding has been separated from the antitoxin function in the tripartite *hipBST* system, we initially constructed a vector-based transcriptional fusion reporter, in which the native *hipBST* promoter region and 5' region of the *hipB* gene was transcriptionally fused to *lacZ* (**Figure 5a, right**). We then expressed various combinations of *hipB*, *hipS*, and *hipT* in *trans* in *E. coli* MG1655 Δ*lacIZYA* cells that do not harbour the *hipBST* operon, utilising two inactivating kinase mutations, *hipT*^{D210Q} (**Figure 5a, left**) and *hipT*^{D233Q} (**Figure S5a**) to avoid toxicity. In both cases, in the absence of expressed *hipBST* components, active transcription could be observed by the blue colour of the colonies (**Figure 5a, left**, top row, pBAD33),

Figure 5. The HipBST complex is dynamic *in vivo*. a. Right, transcriptional reporter assay using a plasmid with the *hipBST* promoter region and a 5' fragment of the *hipB* gene (including 224 bp upstream of *hipB* plus the first 73 bp of the *hipB* gene, shown with a lightening symbol) transcriptionally fused to *lacZ*. Left, *E. coli* TB28 harbouring the reporter plasmid (pSVN141, pGH254::P$_{hipBST}$-*hipB*'-*lacZ*) and empty pBAD33 vector or combinations of *hipB*, *hipS*, and *hipT* in the context of the HipT^{D210Q} inactive mutant; pSVN180 (pBAD33::*hipB-S-T*^{D210Q}, "*hipBST*^{D210Q}"), pSVN182 (pBAD33::*hipB-S*, "*hipBS*"), pSVN184 (pBAD33::*hipB-T*^{D210Q}, "*hipBT*^{D210Q}"), pSVN187 (pBAD33::*hipS-T*^{D210Q}, "*hipST*^{D210Q}"), pSVN189 (pBAD33::*hipB*, "*hipB*"), pSVN190 (pBAD33::*hipS*, "*hipS*"), or pSVN192 (pBAD33::*hipT*^{D210Q}, "*hipT*^{D210Q}") were grown, diluted, and spotted onto YT, X-gal agar plates

containing 0.2% glucose (to repress $hipB/S/T^{D210Q}$), or 0.2% arabinose (to induce $hipB/S/T^{D210Q}$). Results are

representative of two independent experiments. **b.** Left, HipT exchange assay where overexpression of a

catalytically inactive kinase (HipT^{D210Q}) is used to exchange wild type HipT from the HipBST complex *in vivo*.

Right, spot assays of *E. coli* O127 harbouring the empty pBAD33 vector, pSVN220 (pBAD33::$hipT^{D210Q}$, "*hipT*

(S^{57}IS^{59}, *wt*)"), pSVN221 (pBAD33::$hipT^{S57D+D210Q}$, "*hipT* (D^{57}IS^{59})"), pSVN222 (pBAD33::$hipT^{S59D+D210Q}$, "*hipT*

(S^{57}ID^{59})"), pSVN223 (pBAD33::$hipT^{S57A+D210Q}$, "*hipT* (A^{57}IS^{59})"), pSVN224 (pBAD33::$hipT^{S59A+D210Q}$, "*hipT*

(S^{57}IA^{59})"), pSVN225 (pBAD33::$hipT^{S57A+S59A+D210Q}$, "*hipT* (A^{57}IA^{59})"), pSVN226 (pBAD33::$hipT^{S57D+S59A+D210Q}$,

"*hipT* (D^{57}IA^{59})"), pSVN227 (pBAD33::$hipT^{S57A+S59D+D210Q}$, "*hipT* (A^{57}ID^{59})") or pSVN228

(pBAD33::$hipT^{S57D+S59D+D210Q}$, "*hipT* (D^{57}ID^{59})") on YT agar plates containing 0.2% glucose (to repress $hipT^{D210Q}$

autophosphorylation mutants) or 0.2% glucose plus 0.2% arabinose to induce a limited amount of HipT. The column

to the right indicates whether HipT exchange has taken place (+) or not (-). Results are representative of two

independent experiments. **c.** As **b** but with additional overexpression of *hipB* to avoid renewed transcription.

Overnight cultures of *E. coli* O127 harbouring pSVN111 (pNDM220::*hipB*) in combination with the empty

pBAD33 vector, pSVN211 (pBAD33::$hipT^{D210Q}$, "*hipT* (S^{57}IS^{59}, *wt*)"), pSVN212 (pBAD33::$hipT^{S57D+D210Q}$, "*hipT*

(D^{57}IS^{59})"), pSVN213 (pBAD33::$hipT^{S59D+D210Q}$, "*hipT* (S^{57}ID^{59})"), pSVN214 (pBAD33::$hipT^{S57A+D210Q}$, "*hipT*

(A^{57}IS^{59})"), pSVN215 (pBAD33::$hipT^{S59A+D210Q}$, "*hipT* (S^{57}IA^{59})"), pSVN216 (pBAD33::$hipT^{S57A+S59A+D210Q}$, "*hipT*

(A^{57}IA^{59})"), pSVN217 (pBAD33::$hipT^{S57D+S59A+D210Q}$, "*hipT* (D^{57}IA^{59})"), pSVN218 (pBAD33::$hipT^{S57A+S59D+D210Q}$,

"*hipT* (A^{57}ID^{59})") or pSVN219 (pBAD33::$hipT^{S57D+S59D+D210Q}$, "*hipT* (D^{57}ID^{59})") were grown, diluted, and spotted

onto YT agar plates containing 0.2% glucose (to repress *hipT* variants), 0.2% arabinose (to induce *hipT* variants) or

0.2% arabinose and 500 µM IPTG (to induce *hipB*). Results are representative of two independent experiments.

while induction of any construct containing *hipB* efficiently repressed transcription (**Figure 5a, left**, right plate, *hipBST*[D210Q], *hipBS*, *hipBT*[D210Q], and *hipB*). For *hipBST* and *hipBT*, leaky expression on repressing glucose plates was sufficient to reduce transcription. We conclude that *hipB* is both sufficient and required for transcriptional repression of the *hipBST* locus. Given that the *hipBST* and *hipBT* constructs repress transcription even in the absence of inducer, our data are also consistent with HipT (and possibly HipS) stabilising DNA interaction by HipB, as observed for many toxin-antitoxin systems (Harms et al., 2018).

Phosphorylation of HipT Ser57 prevents HipS binding *in vivo*

In principle, the observed lack of HipS antitoxin inhibition of toxicity in the S57D phosphomimetic HipT mutants could be caused by either lack of binding of HipS or hyperactivity of the kinase. To distinguish between these two scenarios, we next tested if inactive HipT can exchange with wildtype HipT inside the HipBST complex *in vivo* in *E. coli* O127:H6. To achieve this, we constructed a series of HipT autophosphorylation variants on a background of inactive HipT[D210Q] and asked if expression would release wildtype HipT from endogenous HipBST complexes encoded by *hipBST* of *E. coli* O127:H6 and thus cause toxicity (**Figure 5b, right**). Both in the absence and presence of the inducer arabinose, all inactive HipT autophosphorylation variants, except those harbouring the phosphomimetic Asp57 ($D^{57}IS^{59}$, $D^{57}IA^{59}$, and $D^{57}ID^{59}$) were able to induce toxicity in this setup, thus suggesting that exchange can take place when Ser57 is not phosphorylated (**Figure 5b, left**). Of these, the $S^{57}IS^{59}$, $A^{57}IS^{59}$, and $S^{57}IA^{59}$ variants had the strongest ability to release wildtype HipT activity while the $S^{57}ID^{59}$, $A^{57}IA^{59}$ and $A^{57}ID^{59}$ variants activated HipT at a lower level. To exclude the possibility that expression of inactive HipT would affect the natural transcriptional repression of the endogenous HipBST complex, we repeated the experiment in a setup where excess HipB is expressed from a separate plasmid using IPTG, knowing that this represses transcription (**Figure 5c,**

right). Due to an uncharacterised growth inhibiting effect observed upon *hipBT* co-expression for the

inactivating kinase mutations, *hipT*^{D210Q} (**Figure 5a, left**) and *hipT*^{D233Q} (**Figure S5a**), we used

genetic constructs with a lower expression level of the HipT mutants for expression in combination

with HipB to obtain the data in figure 5c. Here we found similar results even with full transcriptional

repression, supporting that the observed toxicity is indeed due to wildtype HipT being exchanged

away from preformed endogenous HipBST complexes upon expression of inactive toxin. To confirm

that the growth defects observed were in fact dependent on the *hipBST* system and not a secondary

effect in the host, we performed a similar assay in *E. coli* K-12 with the intact *hipBST* operon fused

to *lacZ* encoding wildtype HipT (**Figure S5b**). Growth inhibition was only observed when inactive

HipT$^{D210Q/D233Q}$ was expressed *in trans* confirming that the phenotype is indeed dependent on the

hipBST operon. To ensure that the low expression levels of the HipT^{D210Q} autophosphorylation

mutants in combination with HipB was not in themselves causing the observed growth inhibitions as

seen for high expression levels (**Figure 5a, left and S5a**), we finally transformed the same vectors

used in *E. coli* O127:H6 into *E. coli* K-12. None of the vectors caused growth inhibition upon

induction (**Figure S5f**), confirming that the phenotype observed in **Figure 5b** must be due to the

presence of the native *hipBST* operon. Finally, to rule out any differences in transcription of the

hipBST promoter between *E. coli* K-12 and O127:H6, we confirmed that mRNA expression can be

repressed in *E. coli* K-12 when *hipBST*^{D210Q} is present *in trans* (**Figure S5c**) and that the

transcriptional start sites (TSS) in the two strains are identical (**Figure S5d-e**). In summary, we

conclude that free HipT molecules are able to exchange with those inside the inhibited HipBST

complex, except in the S57D mutants, suggesting that phosphorylation at Ser57 controls exchange,

likely via control of binding of HipS.

The core HipT kinase domain is flexible in solution

Since the protein exchange experiments suggested that the HipBST complex is dynamic *in vivo*, we

finally performed a careful analysis of the obtained small-angle X-ray scattering (SAXS) data to

understand the structure of the complex in solution (**Figure S6a-b**). We found, that the pair distance

distribution function indicates a maximum complex diameter of 170 Å, significantly larger than the

value calculated from the crystal structure (140 Å) suggesting some degree of structural flexibility

(**Figure S6c-d**). Moreover, fitting of the crystal structure to the solution data yielded a relatively poor

fit, so to accommodate internal flexibility, the HipT structure was split into three parts, roughly

corresponding to the N-terminal mini domain (1-59), the N-terminal subdomain 2 (60-169), and the

core kinase domain (170-335) (**Figure 2a**). Following rigid body refinement against the SAXS data

with soft restraints between the three parts of HipT, including HipS and HipB, and imposing two-fold

symmetry, we obtained a better fit which could be further improved by allowing for partial

dimerisation of the hetero-hexamer (see Material and Methods for details). In this model, the core

kinase domain of HipT has a looser structure than observed in the crystal structure (root mean square

deviation for Cα atoms HipT ~6.5 Å, HipBST ~7 Å) suggesting that the complex in the solution state

allows for internal flexibility in the kinase. In summary, in solution structural analysis confirms that

the HipBST complex has a flexible structure, likely caused by structural dynamics in the HipT core

kinase domain, thus consistent with a model in which HipS (and perhaps HipB) can release and rebind

HipT dynamically.

DISCUSSION

In this paper, we present a detailed structural and functional analysis of the tripartite HipBST toxin-

antitoxin system from enteropathogenic *E. coli* O127:H6. We show that most HipT kinases contain a

conserved SIS motif in their Gly-rich loops, allowing for fine-tuned regulation via dual auto-

phosphorylation. Other members of the family contain various combinations of Ser and Thr at these

positions, which may serve similar roles since Hip toxins are Ser/Thr kinases (Pereira et al., 2011).

Crystal structures of the HipBST complex in several, naturally occuring phosphorylation states

demonstrate that HipT is found in its inactive state inside the HipBST complex regardless of its

autophosphorylation state, consistent with a model whereby HipS, possibly aided by the DNA-

binding protein HipB, keeps the kinase in an inactive conformation and thus overrules any effects of

autophosphorylation. We show that this inhibition is dependent on a conserved Trp65 in HipS, which

couples HipS binding directly to the active site of HipT and the conformation of its Gly-rich loop.

Trp65 is located in a motif that differs both structurally and at the sequence level from the

corresponding region in the N-terminal subdomain 1 of HipA, which is otherwise very similar to

HipS (Gerdes et al., 2021). Together, this suggests that the molecular basis for the role of HipS as

antitoxin in the HipBST complex involves prevention of the Gly-rich loop from transitioning from

the outward, inactive state, to another conformation compatible with ATP binding (**Figure 6**).

Figure 6. Model for the action of HipS as antitoxin in the HipBST system. Structural comparison of the positions

of the Gly-rich loops in HipT D233Q (blue) and *E. coli* K-12 HipA in its ATP-bound conformation (PDB: 3DNT,

green/purple) (Schumacher et al., 2009). Shown is also HipS with Trp65, which is predicted to prevent transition of

the Gly-rich loop from an outward-facing (HipT) to an inward-facing (HipA:ATP) conformation (shown with an

arrow). The active site residues of HipA as well as ATP:Mg^{2+} are shown with sticks on top of the HipT cartoon.

Ser150, which undergoes phosphorylation in HipA, is indicated as well.

The size and topology of the HipT kinase with its lack of the N-terminal subdomain 1 found in HipA is reminiscent of the Cyclin-dependent kinases (Cdks), which are minimal kinases that regulate the eukaryotic cell cycle (**Figure S7a**) (Lim & Kaldis, 2013). Intriguingly, one of these, Cdk5, also regulates translation through direct phosphorylation of glutamyl-prolyl tRNA synthetase (EPRS) (Arif et al., 2011). Despite differences in tertiary structure (**Figure S7b**), the residues involved in the active site align well, suggesting common reaction mechanisms. Cyclin-dependent kinases are stabilised by two hydrophobic regions, the R (regulatory) and C (catalytic) spine (Taylor & Kornev, 2011). Of these, the R spine changes conformation upon an activating phosphorylation by flipping the final active site residue into place and stabilising the kinase by forming an interaction to an aspartate residue in the so-called F helix (Baugh et al., 2015). While several of these structural elements are not present in HipT, the bacterial kinase does feature a conserved hydrophobic core similar to the C spine (**Figure S7c-e**). These regions are more extended than seen in Cdk, in particular due to the presence of two additional, conserved aromatic residues (Y138 and Y232 in HipT, **Figure S7d-e**). In the structure of ATP-bound HipA, the adenine base intercalates right between these and thereby connects the N and C lobes of the kinase. Interestingly, the invariant small hydrophobic residue found between the phosphor-serine positions in HipT (S<u>I</u>S) would connect the adenine base to the N lobe, suggesting that it is conserved to preserve the ATP binding. In general, it thus appears that the similarities between bacterial and eukaryotic kinases may be larger than hitherto appreciated and consequently, that the vast set of knowledge available for the eukaryotic enzymes might help us understand the Hip kinases as well.

A surprising discovery of this study is that HipT was found with phosphorylation at Ser57 or Ser59 when expressed in the context of the inactive HipBST complex. This suggests that some active (HipS-free) HipT exists in the cell and hence compatible with a model whereby the HipBST complex is

dynamic and allows HipT to dissociate *in vitro*. In this context, it is likely that the exposed

conformation of the Gly-rich loop renders it particularly susceptible to phosphorylation by active

HipT kinase. Taken one step further, this could indicate that a secondary effect of HipS binding (apart

from inactivating the kinase) is to expose the kinase to regulation via autophosphorylation in *trans*.

We still don't understand what controls whether HipT becomes modified at one or the other position

but the structures of HipT^{S57A} and HipT^{S59A} could suggest that subtle changes in the loop can make a

big difference (**Figure 4a**). Moreover, since Ser57/Ser59 phosphorylation was observed for both

alanine single mutations *in vitro*, the presence of both serine residues does not appear to be important

for the autophosphorylation activity of HipT. Finally, the observation that the S^{57}IS^{59}, A^{57}IS^{59}, and

S^{57}IA^{59} HipT^{D210Q} variants exhibit a stronger toxicity due to exchange with wildtype HipT than the

A^{57}IA^{59} variant *in vivo* (**Figure 5b-c**) indicates that the hydroxyl groups of the serine residues play a

role in the binding of HipS.

Intriguingly, although the two autophosphorylation positions in HipT occupy a similar space near the

γ-phosphate of ATP, they serve different roles. HipT is active in both phosphomimetic isoforms (S^{57}D

and S^{59}D) while removal of the hydroxyl group on either serine by mutation to alanine (S^{57}IA^{59},

A^{57}IS^{59} and A^{57}IA^{59}) reduces its ability to cause growth inhibition. In contrast to this, we found that

both S^{57}IA^{59} and A^{57}IS^{59} remained capable of autophosphorylation, suggesting that the presense of

both of the serine residues or their phosphorylated counterparts could be important for regulating

whether HipT is active on itself or its target (TrpS). Taken together, these results are consistent with

a model in which autophosphorylation at either position is required for target phosphorylation.

Surprisingly, in contrast to the A^{57}ID^{59} mutant, the D^{57}IA^{59} mutant was still active in inhibiting growth

(but with a delayed effect compared to D^{57}IS^{59}, S^{57}ID^{59}, and D^{57}ID^{59}), suggesting that

phosphorylation of Ser57 results in a kinase, which is more potent against its target compared to

phosphorylation of Ser59. Finally, although the $D^{57}ID^{59}$ mutant is fully active in growth inhibition,

this may be artificial, as dual phosphorylation has not been observed *in vivo* (Vang Nielsen et al.,

2019). Nevertheless, the $D^{57}ID^{59}$ mutant gives us valuable information, as it shows that growth

inhibition does not require the presence of a serine residue and that phosphorylation is likely what

determines the active state. The differences in the ability of the $S^{57}IA^{59}$ and $D^{57}IA^{59}$ mutants to cause

growth inhibition are also striking. The $S^{57}IA^{59}$ mutant has a very small growth inhibitory effect,

whereas $D^{57}IA^{59}$ has a clear effect. One explanation for this observation could be that the $S^{57}IA^{59}$

mutant has a slower rate of autophosphorylation compared to wildtype ($S^{57}IS^{59}$) and that this would

be required for HipT activity. When examining the other autophosphorylation site, Ser59, it is

noteworthy, that the $S^{57}ID^{59}$ mutant is fully active, while the $A^{57}IS^{59}$ and $A^{57}ID^{59}$ mutants are inactive.

Another question raised by the structures is how HipS can be bound in the complex with Ser57 in its

phosphorylated state if this serves to block HipS binding? In this context, it is important to note,

however, that only a minor fraction of the purified complex is phosphorylated (**Figure 4c**),

confirming that if HipT (Ser57-P) binds to HipS at all, it probably binds very poorly. It is also possible

that phosphorylation occurred during purification. In either case, we believe that our data is

compatible with the model for DNA binding and toxin activation for the HipT kinase toxin shown in

Figure 7. Under non-activating conditions, HipBST will be repressing the *hipBST* operon via binding

to promoter DNA. Liberation of HipBST from DNA by an unknown mechanism will result in relaxed

promoter repression and increased transcription from the operon. Free HipT could then form three

populations, non-phosphorylated HipT, HipT P-Ser57, and HipT P-Ser59. Our results indicate that

both phosphorylated forms are active on TrpS and as a result of this, reduce cell growth. Furthermore,

HipT P-Ser59 would readily be able to (re-)bind HipS, and therefore form a stable and inactive

HipBST complex as well as tightly bind the *hipBST* promoter, effectively shutting down the kinase

Figure 7. Proposed model of regulation and activation of the HipBST system. Transcription from the *hipBST*
operon (middle row, left) is repressed by binding of the HipBST hetero-hexamer in which HipT is inactive and the
Gly-rich loop (red) is outward-facing when HipS is bound. The dynamic HipBST complex can disassociate to
generate free HipT, which can phosphorylate another HipT kinase and perhaps TrpS, using ATP.
Autophosphorylation at Ser 57 (top box) severely reduces HipS binding, thereby prolonging the growth inhibitory
effect of HipT by keeping the kinase active and preventing full transcriptional repression. On the contrary,
autophosphorylation at Ser59 (bottom box) stabilises HipS binding and the HipBST complex, reducing the activity
of HipT on TrpS compared to when autophosphorylated at Ser57 and repressing transcription from the *hipBST*
operon.

activity, halting the expression of new protein. On the contrary, HipT P-Ser57 would have reduced

HipS binding and therefore stay in its active conformation. The overall effect on growth could then

be a balance between these two counter-acting responses.

ACKNOWLEDGEMENTS

The authors are indebted to the beamline staff at P14 in EMBL Hamburg, and BioMAX in MaxIV

Lund for help during data collection. This project was funded by grants from the Novo Nordisk

Foundation (NNF18OC0030646 to D.E.B.) and Danish Natural Research Foundation's Centre of

Excellence for Bacterial Stress Response and Persistence (grant number DNRF120).

COMPETING INTERESTS

The authors confirm that there are no competing interests.

AUTHOR CONTRIBUTIONS

Conceptualization, K.G., M.A.S, and D.E.B.; Investigation, R.L.B., S.V.N., F. B., and J.L.; Formal

Analysis, R.L.B, S.V.N., F.B., and J.S.P; Writing – Original Draft, R.L.B., S.V.N., and D.E.B.;

Writing – Review & Editing, R.L.B., S.V.N., J.L., J.S.P., K.G., M.A.S., and D.E.B; Visualization,

R.L.B., S.V.N., M.A.S., and D.E.B.; Funding Acquisition, D.E.B. and K.G.; Resources, J.S.P., K.G.,

M.A.S. and D.E.B; Supervision, K.G., J.S.P., M.A.S., and D.E.B.

MATERIALS AND METHODS

Phylogenetic analysis. The previously identified set of 48 HipT orthologues (Gerdes et al., 2021)

was used for sequence alignment by Clustal Omega (Sievers & Higgins, 2018) at www.ebi.ac.uk and

imported into Jalview (Waterhouse et al., 2009). The phylogenetic tree was visualized using iTOL

(Letunic & Bork, 2019). Reconstruction of the phylogenetic tree was accomplished using IQ-TREE

that uses the Maximum Likelihood approach and Ultrafast bootstrapping via the CIPRES module in

Genious Prime (Minh et al., 2020).

Strains and plasmids. Strains and plasmids are listed in **Table S2**, and DNA oligonucleotides in

Table S3.

Site-directed mutagenesis. Amino acid change HipT$_{S57A}$ was constructed by PCR mutagenesis

(**Table S3**) as previously performed (Vang Nielsen et al., 2019). PCR product was digested with

DpnI, and resulting plasmid transformed into *E. coli* strain DH5α.

Protein purification and structure determination. Expression of the *E. coli* O127:H6 HipBST

complex was done using constructs pSVN78 (HipBST$_{S57A}$), pSVN96 (HipBST$_{D233Q}$), and pMME3

(HipBST$_{S59A}$) derived from pSVN78. The constructs contained the genes coding for HipB, HipS, and

C-terminal hexa-histidine tagged HipT, all separated, and with optimised Shine-Dalgarno sequences

for large scale expression. All constructs contained an IPTG-inducible promoter and ampicillin

resistance gene for selection. For each construct, 2 L cultures with a cell density of OD$_{600}$ = 0.6 were

induced with a final concentration of 1 mM IPTG, and left to express ON at 20°C. Cell pellets were

resuspended in lysis buffer (50 mM Tris-HCl, pH 7.5, 300 mM NaCl, 20 mM imidazole, 5% glycerol,

5 mM BME) together with 1 mM PMSF. Cells were lysed by sonication and the cleared lysate was

applied to a 5 mL HisTrap HP column (Cytiva) equilibrated in lysis buffer, and washed in wash buffer

(50 mM Tris-HCl, pH 7.5, 300 mM NaCl, 40 mM imidazole, 5 mM BME), before eluting with elution

buffer (50 mM Tris-HCl, pH 7.5, 300 mM NaCl, 400 mM imidazole, 5 mM BME). The eluate was

applied directly to a 5 mL Heparin HF column equilibrated in 70% Buffer A (50 mM Tris-HCl, pH

7.5, 5 mM BME) and 30% Buffer B (50 mM Tris-HCl, pH 7.5, 1M NaCl, 5 mM BME) running on

an ÄKTA Pure system (Cytiva). Final separation was achieved after concentrating the Heparin elution

to approximately 8 mg/mL using a Vivaspin column with a 30 kDa cutoff (Sartorius), before applying

the sample to a Superdex 200 10/300 GL (Cytiva) column equilibrated in gel filtration buffer (20

mM Tris-HCl, pH 7.5, 300 mM NaCl, 5 mM BME). Crystals of HipBST[S57A] grew as clusters of thin

plates in a 2 μL drop with a 1:1 ratio of protein to buffer, which consisted of 0.1 M Bicine, pH 9, 8%

MPD and with a protein concentration of 5 mg/mL, and a reservoir of 200 μL. Crystals of all three

mutant forms were obtained by a very similar protocol. All crystals were cryoprotected in the mother

liquor supplemented with 25% MPD before freezing in liquid N_2.

Data collection for HipBST[S57A] was performed at P14 in Hamburg and for HipBST[S59A] and

HipBST[D233Q] at BioMAX in Lund. For HipBST[S59A] and HipBST[D233Q], 7,200 images were collected

with an oscillation of 0.1° and a transmission of 100%. The HipBST[S57A] data set was collected with

an oscillation of 0.1°, 3,600 images and a transmission of 70%. All data was processed in XDS

(Kabsch, 2010), using the CC½ value after scaling as indicator of the data resolution limit (Karplus

& Diederichs, 2012). The space group was confirmed using Pointless (Evans, 2006), and all structures

were determined using molecular replacement with Phaser (McCoy et al., 2007) from the CCP4 suite

(Winn et al., 2011). The data set for HipBST[S57A] used a heavily truncated HipA to obtain initial phases

for HipT, while the data set for HipBST[S59A] and HipBST[D233Q] used the solved structure of HipBST

as a search model. Refinement was performed using Buster (Smart et al., 2012) using 1 big cycle of

20 small cycles of refinement when building, and 5 big cycles of 100 small cycles of refinement

afterwards. Non-crystallographic symmetry (NCS) restraints were used during building since the

asymmetric unit (ASU) contained a dimer but some refinement rounds without NCS were also

included to allow for differences between the molecules in the ASU. Automatic water placement was

used and water molecules with no density were removed. Model building was performed in Coot

(Emsley et al., 2010). All structures were validated by the MolProbity server and Rama-Z scores.

Final R-work/R-free for HipBST$_{D233Q}$, HipBST$_{S57A}$, and HipBST$_{S59A}$, is 0.19/0.23, 0.21/0.24, and

0.20/0.24, respectively.

Phos-tag gel of pre- and post-incubation with ATP. 15% Phos-tag acrylamide gels (Wako) were

cast according to the manufacturer's guidelines, except that 100 µM Phos-tag acrylamide was used to

ensure proper separation between phosphorylated and non-phosphorylated HipT. The non-

phosphorylated HipT control was expressed from pSNN2. The gel was run at 4 °C until the loading

dye reached the bottom of the gel and visualized using standard Coomassie Blue staining as for

normal SDS-PAGE gels. HipBST samples were run directly on the gel orincubated with ATP for 30

minutes at room temperature in a final mixture of: 2 mM KCl, 3.2 µM ZnSO$_4$, 4 mM MgCl$_2$, 66 µM

ATP, 14 µL gel filtration buffer, and 1 mg/mL protein sample.

Spot assays and growth curves. Cultures were grown in liquid YT medium or MOPS minimal

medium + 0.2% glucose at 37°C shaking at 160 rpm and solid medium used was YT agar plates,

which were incubated at 37°C for approximately 16 h. When applicable, media were supplemented

with 25 µg/mL chloramphenicol, 30 µg/mL ampicillin, 25 µg/mL kanamycin and/or 40 µg/mL 5-

bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-gal). For spot assays, cells were grown as

overnight cultures, diluted to obtain the same values of OD$_{600}$, centrifuged at 5,000 rpm for 5 min,

washed in phosphate-buffered saline (PBS), and serially diluted before being spotted onto YT agar

plates containing the indicated amount of inducer/repressor. Gene expression from plasmids carrying

the pBAD promoter was induced by a final concentration of 0.2% arabinose and repressed by 0.2%

glucose. Gene expression from plasmids carrying the P$_{A1/O4/O3}$ promoter was induced by a final

concentration of 200 or 500 μM isopropyl-D-1 thio-galactopyranoside (IPTG), as indicated. Growth

experiments were done in YT medium with addition of relevant antibiotics, diluted from overnight

cultures and grown exponentially for at least 4 h until a seemingly constant doubling time. At OD_{600}

≈ 0.2, arabinose (0.2%) was added to induce wildtype *hipT* or autophosphorylation mutants. After

another 1.5 h, IPTG (200 μM) was added to induce *hipS* or *hipBS*. For each repetition, an independent

colony from the strain was used to start separate cultures. No outliers were rejected.

Northern blot analysis. Total RNA samples were prepared using hot phenol extractions. Cultures

were grown in YT medium (or MOPS minimal medium + 0.2% glucose in the case of the *E. coli*

TB28 strain used for primer extension as described below). At indicated time points 10 mL were

harvested into 1.25 mL ice-cold stop solution (5% phenol, pH 4.3 in absolute ethanol) and pelleted

by centrifugation at 4,000 rpm at 4 ºC for 10 min. Cell samples were resuspended in 200 μL ice-cold

solution I (0.3 M sucrose; 0.01 M Na-Acetate, pH 4.5) and mixed with 200 μL solution II (2% SDS;

0.01 M Na-Acetate, pH 4.5) and 400 μL phenol (pH 4.3). Samples were vortexed for 10 s. before

incubation at 65 ºC for 3 min. followed by freezing in liquid nitrogen and centrifugation at 13,000

rpm at room temperature for 5 min. The upper phase was transferred to a new tube with 400 μL

phenol (pH 4.3) followed by another round of the vortexing, heating, freezing and centrifugation

steps. The RNA was precipitated in 1 mL ice-cold absolute ethanol and 50 μL RNA storage buffer (3

M Na-Acetate, pH 4.7; 10 mM EDTA) at -80 ºC for a minimum of 30 min. RNA was then pelleted

by centrifugation at 13,000 rpm at 4 ºC for 30 min., washed in 70% ice-cold ethanol and air-dried at

room temperature. Finally, samples used for northern blotting analysis were resuspended in 20 μL

RNA storage buffer, whereas samples used for primer extension were resuspended in 20 μL nuclease-

free water. Deacylated control sample used for northern blotting analysis was prepared by mixing 4

μL of any of the total RNA samples with 6 μL 1 M Tris-HCl (pH 9) and 50 μL nuclease-free water

and incubating for 2 hours at 37 ºC. This was followed by the addition of 150 µL ice-cold absolute

ethanol and 4 µL 5 M NaCl and the tube was incubated at -20 ºC for 15 min. The sample was then

centrifuged at 13,000 rpm at 4 ºC for 30 min. The supernatant was discarded, and RNA pellet was

air-dried at room temperature and resuspended in 4 µL RNA storage buffer.

Northern analysis was used for detection and comparison of charged versus uncharged tRNA. 4 µL

RNA samples were mixed with 6 µL tRNA loading buffer (0.1 M Na-succinate, pH 5; 8 M urea;

0.05% bromophenol blue and 0.05% xylene cyanol) and separated on 6% polyacrylamide gels with

8 M urea buffered in 0.1 M Na-succinate (pH 5) and run at 35 V. RNA was transferred to a Hybond-

N+ membrane (Amersham) at 550 mA constant for 1 hour in blotting buffer (0.8 M Tris; 40 mM

EDTA; acetic acid to pH 8.1). Membranes were pre-hybridized at 42 ºC in 6 mL hybridization buffer

(5 x SSPE; 5 x Denhardt's solution; 0.5% SDS; ~0.55 mg/mL herring sperm DNA (Sigma)) for one

hour before addition of 10 µM DNA oligonucleotide probe, which had been radioactively labelled

with [γ-^{32}P]-ATP, for overnight hybridization. Oligonucleotides complementary to the tRNA

anticodons of interest were 5'-end labelled using T4 polynucleotide kinase (Thermo Scientific) and

40 µCi of [γ-^{32}P]-ATP (6000 Ci/mmol) by incubating at 37 ºC for 2 hours. For detection of tRNA^{trpT},

tRNA^{gltTUVW} and   tRNA^{argVYZQ},   oligonucleotides   CCCAACACCCGGTTTTGG,

CCTGTTACCGCCGTGAAAGGG and TCCGACCGCTCGGTTCGTAGC were used, respectively.

Excess probe was removed by washing in 2 x SSC buffer with 0.1% SDS at room temperature and

radiation was visualized by phosphorimaging. Membranes were stripped for re-probing at 95 ºC in

stripping buffer (0.1 x SSPE; 0.5% SDS) until no remaining radiation was detected.

Primer extension analysis. Primer extension was used to map mRNA 5' ends using radioactively

labelled primers following extension by reverse transcription. Specific oligonucleotide primers (4

pmol) were radioactively 5'-end labelled with 130 µCi of $[\gamma\text{-}^{32}\text{P}]$-ATP (6000 Ci/mmol) using T4

polynucleotide kinase (Thermo Scientific) and incubated for 30 min at 37 ºC followed by inactivation

for 15 min at 75 ºC and cooling on ice. To remove excess unincorporated $[\gamma\text{-}^{32}\text{P}]$-ATP, the reaction

was centrifuged through an illustra MicroSpin G-25 Column (Cytiva) at 750 g for 1 min.

Sequencing reactions used as markers of molecular weight were generated using a purified PCR

fragment amplified with primers FP43 and RP38 and 0.1 pmol from pKG127 as template DNA. The

reactions were incubated with radioactively labelled primer (0.4 pmol) and extensions were

performed with DreamTaq DNA polymerase (Thermo Scientific) in the presence of either ddGTP,

ddATP, ddTTP or ddCTP termination mix (Roche). PCRs were carried out and reactions were

stopped by the addition of an equal volume of 2 x formamide loading buffer (95% formamide; 0.025%

bromophenol blue and 0.025% xylene cyanol; 5 mM EDTA; 0.025% SDS).

For extension reactions, radioactively labelled oligonucleotides (0.2 pmol) were hybridized to 10 µg

of total RNA by incubating at 80 ºC for 5 min followed by quick cooling on ice for 5 min. Reverse

transcriptions were carried out using Superscript III reverse transcriptase (Invitrogen) in 1 x First-

Strand buffer plus 10 mM DTT and 1 mM dNTP for 1 hour at 52 ºC and 47 ºC for RP35 and RP37,

respectively. Reactions were stopped by the addition of an equal volume of 2 x formamide loading

buffer. Sequencing and extension reactions were fractionated on 6% polyacrylamide gels with 8 M

urea, buffered in 1 x TBE and run at 45 W. Resulting gels were fixed in fixing solution (50% ethanol;

10% acetic acid), dried and visualized by phosphor imaging. The promoter of *hipBST* was analysed

by primer extension using RP35 and RP37 for mapping of the transcriptional start site from pSVN141

(pGH254::P*hipBST*-*hipB'-lacZ*) in *E. coli* TB28 and the native *hipBST* operon in *E. coli* O127,

respectively.

Small-angle X-ray scattering measurements and analysis. HipBST$_{S57A}$ purified as specified in "Protein purification and structure determination" was used for the SAXS analysis. The SAXS measurements were performed on the optimized NanoSTAR instrument (Bruker AXS) at the iNANO center at Aarhus University (Lyngsø & Pedersen, 2021). It uses a high brilliance Ga metal-jet X-ray source (Excillum), special long optics, a two-pinhole collimation with a home-built 'scatterless' slit pinhole, and a VÅNTEC-2000 (Bruker AXS) microgap 2D gas proportional detector. The sample and buffer were measured in the same flow-through quartz capillary, the scattering from the buffer was subtracted from that of the sample, and the data were converted to absolute scale using the scattering from water. The intensity data, $I(q)$, are displayed as a function of the modulus of the scattering vector $q = 4\pi \sin(\theta)/\lambda$, where 2θ is the scattering angle and λ is the wavelength of the X-rays. The fit to the data in a Guinier plot gave the radius of gyration $R_g = 52.2 \pm 0.2$ Å and the forward scattering $I(0) = 0.0868 \pm 0.005$ cm^{-1}. The molar mass was calculated as: $M = I(0)N_A/(c\Delta\rho_m^2)$, where N_A is Avogadro's number, $c = 1.01$ mg mL^{-1} is the mass concentration, and $\Delta\rho_m = 2.00 \times 10^{10}$ cm g^{-1} is the typical excess scattering length per unit mass for proteins. The radius of gyration and forward scattering can also be obtained from the pair distance distribution function $p(r)$ determined by indirect Fourier transformation (Glatter, 1977; Pedersen et al., 1994) and the values are quite close to those from the Guinier fit: $R_g = 53.8 \pm 0.3$ Å and $I(0) = 0.08651 \pm 0.0005$ cm^{-1}, where the latter corresponds to a mass of $M = 129$ kDa. The maximum in a normalized Kratky plot using the values of R_g and $I(0)$ from the Guinier fit for normalisation agrees with a relatively compact structure, however, the plateau behaviour for $qR_g > 3$ suggest some openness and potential flexibility of the structure of the complexes. The scattering of the complex from the crystal structure was directly compared to the SAXS data and a poor agreement was observed. To improve the agreement, rigid body refinement (RBR) of the structure was performed using previously described methods (Steiner et al., 2018). For the RBR of HipBST in three connected domains, the

first runs showed that there was some degree of oligomerisation present in the sample, since the model

curve is below the measured data at low q. Therefore, a dimer factor $S_{agg}(q) = 1 + \frac{A \sin(qD)}{qD}$ (Larsen

et al., 2020) was multiplied with the intensity calculated for the structure. In the expression, A is the

mass fraction of dimers of the HipBST complex and D is the distance between the two copies in the

dimer. This approach gave good fits to the data.

DATA AND SOFTWARE AVAILABILITY

The accession numbers for the structures reported in this paper are PDB: 7AB3 (HipBST HipT

S57A), 7AB4 (HipBST HipT S59A), and 7AB5 (HipBST HipT D233Q).

REFERENCES

Arif, A., Jia, J., Moodt, R. A., DiCorleto, P. E., & Fox, P. L. (2011, Jan 25). Phosphorylation of glutamyl-prolyl tRNA synthetase by cyclin-dependent kinase 5 dictates transcript-selective translational control. *Proceedings of the National Academy of Sciences of the U S A, 108*(4), 1415-1420. https://doi.org/10.1073/pnas.1011275108

Baugh, L., Phan, I., Begley, D. W., Clifton, M. C., Armour, B., Dranow, D. M., Taylor, B. M., Muruthi, M. M., Abendroth, J., Fairman, J. W., Fox, D., 3rd, Dieterich, S. H., Staker, B. L., Gardberg, A. S., Choi, R., Hewitt, S. N., Napuli, A. J., Myers, J., Barrett, L. K., Zhang, Y., Ferrell, M., Mundt, E., Thompkins, K., Tran, N., Lyons-Abbott, S., Abramov, A., Sekar, A., Serbzhinskiy, D., Lorimer, D., Buchko, G. W., Stacy, R., Stewart, L. J., Edwards, T. E., Van Voorhis, W. C., & Myler, P. J. (2015, Mar). Increasing the structural coverage of tuberculosis drug targets. *Tuberculosis (Edinb), 95*(2), 142-148. https://doi.org/10.1016/j.tube.2014.12.003

Bordes, P., Sala, A. J., Ayala, S., Texier, P., Slama, N., Cirinesi, A. M., Guillet, V., Mourey, L., & Genevaux, P. (2016, Nov 9). Chaperone addiction of toxin-antitoxin systems. *Nat Commun, 7*, 13339. https://doi.org/10.1038/ncomms13339

Brennan, R. G., & Matthews, B. W. (1989, Feb 5). The helix-turn-helix DNA binding motif. *Journal of Biological Chemistry, 264*(4), 1903-1906. https://www.ncbi.nlm.nih.gov/pubmed/2644244

Emsley, P., Lohkamp, B., Scott, W. G., & Cowtan, K. (2010, Apr). Features and development of Coot. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 4), 486-501. https://doi.org/10.1107/S0907444910007493

Eriani, G., Delarue, M., Poch, O., Gangloff, J., & Moras, D. (1990, Sep 13). Partition of tRNA synthetases into two classes based on mutually exclusive sets of sequence motifs. *Nature, 347*(6289), 203-206. https://doi.org/10.1038/347203a0

Evans, P. (2006, Jan). Scaling and assessment of data quality. *Acta Crystallogr D Biol Crystallogr, 62*(Pt 1), 72-82. https://doi.org/10.1107/S0907444905036693

Evdokimov, A., Voznesensky, I., Fennell, K., Anderson, M., Smith, J. F., & Fisher, D. A. (2009,

Aug). New kinase regulation mechanism found in HipBA: a bacterial persistence switch.

Acta Crystallogr D Biol Crystallogr, 65(Pt 8), 875-879.

https://doi.org/10.1107/S0907444909018800

Gerdes, K., Baerentsen, R., & Brodersen, D. E. (2021, Jun 29). Phylogeny Reveals Novel HipA-

Homologous Kinase Families and Toxin-Antitoxin Gene Organizations. *MBio, 12*(3),

e0105821. https://doi.org/10.1128/mBio.01058-21

Germain, E., Castro-Roa, D., Zenkin, N., & Gerdes, K. (2013, Oct 24). Molecular mechanism of

bacterial persistence by HipA. *Molecular Cell, 52*(2), 248-254.

https://doi.org/10.1016/j.molcel.2013.08.045

Glatter, O. (1977). New Method for Evaluation of Small-Angle Scattering Data. *Journal of Applied*

Crystallography, 10(Oct1), 415-421. https://doi.org/Doi 10.1107/S0021889877013879

Grabe, G. J., Giorgio, R. T., Hall, A. M. J., Morgan, R. M. L., Dubois, L., Sisley, T. A., Rycroft, J.

A., Hare, S. A., & Helaine, S. (2021, Dec). Auxiliary interfaces support the evolution of

specific toxin-antitoxin pairing. *Nature Chemical Biology, 17*(12), 1296-1304.

https://doi.org/10.1038/s41589-021-00862-y

Hallez, R., Geeraerts, D., Sterckx, Y., Mine, N., Loris, R., & Van Melderen, L. (2010, May). New

toxins homologous to ParE belonging to three-component toxin-antitoxin systems in

Escherichia coli O157:H7. *Molecular Microbiology, 76*(3), 719-732.

https://doi.org/10.1111/j.1365-2958.2010.07129.x

Hanks, S. K., Quinn, A. M., & Hunter, T. (1988, Jul 1). The protein kinase family: conserved

features and deduced phylogeny of the catalytic domains. *Science, 241*(4861), 42-52.

https://doi.org/10.1126/science.3291115

Harms, A., Brodersen, D. E., Mitarai, N., & Gerdes, K. (2018, Jun 7). Toxins, Targets, and

Triggers: An Overview of Toxin-Antitoxin Biology. *Molecular Cell, 70*(5), 768-784.

https://doi.org/10.1016/j.molcel.2018.01.003

Haseltine, W. A., & Block, R. (1973, May). Synthesis of guanosine tetra- and pentaphosphate

requires the presence of a codon-specific, uncharged transfer ribonucleic acid in the acceptor

site of ribosomes. *Proceedings of the National Academy of Sciences of the U S A, 70*(5),

1564-1568. https://doi.org/10.1073/pnas.70.5.1564

Huse, M., & Kuriyan, J. (2002, May 3). The conformational plasticity of protein kinases. *Cell,*

109(3), 275-282. https://doi.org/10.1016/s0092-8674(02)00741-9

Kabsch, W. (2010, Feb). Xds. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 2), 125-132.

https://doi.org/10.1107/S0907444909047337

Karplus, P. A., & Diederichs, K. (2012, May 25). Linking crystallographic model and data quality.

Science, 336(6084), 1030-1033. https://doi.org/10.1126/science.1218231

Krissinel, E., & Henrick, K. (2007, Sep 21). Inference of macromolecular assemblies from

crystalline state. *Journal of Molecular Biology, 372*(3), 774-797.

https://doi.org/10.1016/j.jmb.2007.05.022

Larsen, A. H., Pedersen, J. S., & Arleth, L. (2020, Aug). Assessment of structure factors for

analysis of small-angle scattering data from desired or undesired aggregates. *Journal of*

Applied Crystallography, 53(4), 991-1005. https://doi.org/10.1107/S1600576720006500

Letunic, I., & Bork, P. (2019, Jul 2). Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. *Nucleic Acids Research, 47*(W1), W256-W259.

https://doi.org/10.1093/nar/gkz239

Lim, S., & Kaldis, P. (2013, Aug). Cdks, cyclins and CKIs: roles beyond cell cycle regulation.

Development, 140(15), 3079-3093. https://doi.org/10.1242/dev.091744

Lyngsø, J., & Pedersen, J. S. (2021). A high-flux automated laboratory small-angle X-ray scattering

instrument optimized for solution scattering. *Journal of Applied Crystallography, 54*(1),

295-305. https://doi.org/10.1107/s1600576720016209

McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., & Read, R. J.

(2007, Aug 1). Phaser crystallographic software. *J Appl Crystallogr, 40*(Pt 4), 658-674.

https://doi.org/10.1107/S0021889807021206

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., &
Lanfear, R. (2020, May 1). IQ-TREE 2: New Models and Efficient Methods for
Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution, 37*(5), 1530-
1534. https://doi.org/10.1093/molbev/msaa015

Pacios, O., Blasco, L., Bleriot, I., Fernandez-Garcia, L., Ambroa, A., Lopez, M., Bou, G., Canton,
R., Garcia-Contreras, R., Wood, T. K., & Tomas, M. (2020, Sep 21). (p)ppGpp and Its Role
in Bacterial Persistence: New Challenges. *Antimicrobial Agents & Chemotherapy, 64*(10).
https://doi.org/10.1128/AAC.01283-20

Pedersen, J. S., Hansen, S., & Bauer, R. (1994). The aggregation behavior of zinc-free insulin
studied by small-angle neutron scattering. *European Biophysics Journal, 22*(6), 379-389.
https://doi.org/10.1007/BF00180159

Pereira, S. F., Goss, L., & Dworkin, J. (2011, Mar). Eukaryote-like serine/threonine kinases and
phosphatases in bacteria. *Microbiology and Molecular Biology Reviews, 75*(1), 192-212.
https://doi.org/10.1128/MMBR.00042-10

Schumacher, M. A., Balani, P., Min, J., Chinnam, N. B., Hansen, S., Vulic, M., Lewis, K., &
Brennan, R. G. (2015, Aug 6). HipBA-promoter structures reveal the basis of heritable
multidrug tolerance. *Nature, 524*(7563), 59-64. https://doi.org/10.1038/nature14662

Schumacher, M. A., Min, J., Link, T. M., Guan, Z., Xu, W., Ahn, Y. H., Soderblom, E. J., Kurie, J.
M., Evdokimov, A., Moseley, M. A., Lewis, K., & Brennan, R. G. (2012, Sep 27). Role of
unusual P loop ejection and autophosphorylation in HipA-mediated persistence and
multidrug tolerance. *Cell Rep, 2*(3), 518-525. https://doi.org/10.1016/j.celrep.2012.08.013

Schumacher, M. A., Piro, K. M., Xu, W., Hansen, S., Lewis, K., & Brennan, R. G. (2009, Jan 16).
Molecular mechanisms of HipA-mediated multidrug tolerance and its neutralization by
HipB. *Science, 323*(5912), 396-401. https://doi.org/10.1126/science.1163806

Sekine, S., Nureki, O., Dubois, D. Y., Bernier, S., Chenevert, R., Lapointe, J., Vassylyev, D. G., &
Yokoyama, S. (2003, Feb 3). ATP binding by glutamyl-tRNA synthetase is switched to the
productive mode by tRNA binding. *EMBO Journal, 22*(3), 676-688.
https://doi.org/10.1093/emboj/cdg053

Sharrock, A., Ruthe, A., Andrews, E. S. V., Arcus, V. A., & Hicks, J. L. (2018). VapC proteins
from Mycobacterium tuberculosis share ribonuclease sequence specificity but differ in
regulation and toxicity. *PLoS One, 13*(8), e0203412.
https://doi.org/10.1371/journal.pone.0203412

Sievers, F., & Higgins, D. G. (2018, Jan). Clustal Omega for making accurate alignments of many
protein sequences. *Protein Science, 27*(1), 135-145. https://doi.org/10.1002/pro.3290

Smart, O. S., Womack, T. O., Flensburg, C., Keller, P., Paciorek, W., Sharff, A., Vonrhein, C., &
Bricogne, G. (2012, Apr). Exploiting structure similarity in refinement: automated NCS and
target-structure restraints in BUSTER. *Acta Crystallogr D Biol Crystallogr, 68*(Pt 4), 368-
380. https://doi.org/10.1107/S0907444911056058

Stancik, I. A., Sestak, M. S., Ji, B., Axelson-Fisk, M., Franjevic, D., Jers, C., Domazet-Loso, T., &
Mijakovic, I. (2018, Jan 5). Serine/Threonine Protein Kinases from Bacteria, Archaea and
Eukarya Share a Common Evolutionary Origin Deeply Rooted in the Tree of Life. *Journal*
of Molecular Biology, 430(1), 27-32. https://doi.org/10.1016/j.jmb.2017.11.004

Steiner, E. M., Lyngso, J., Guy, J. E., Bourenkov, G., Lindqvist, Y., Schneider, T. R., Pedersen, J.
S., Schneider, G., & Schnell, R. (2018, Sep). The structure of the N-terminal module of the
cell wall hydrolase RipA and its role in regulating catalytic activity. *Proteins, 86*(9), 912-
923. https://doi.org/10.1002/prot.25523

Taylor, S. S., & Kornev, A. P. (2011, Feb). Protein kinases: evolution of dynamic regulatory
proteins. *Trends in Biochemical Sciences, 36*(2), 65-77.
https://doi.org/10.1016/j.tibs.2010.09.006

Vang Nielsen, S., Turnbull, K. J., Roghanian, M., Baerentsen, R., Semanjski, M., Brodersen, D. E.,
Macek, B., & Gerdes, K. (2019, Jun 18). Serine-Threonine Kinases Encoded by Split hipA
Homologs Inhibit Tryptophanyl-tRNA Synthetase. *MBio, 10*(3).
https://doi.org/10.1128/mBio.01138-19

Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M., & Barton, G. J. (2009, May 1).
Jalview Version 2--a multiple sequence alignment editor and analysis workbench.
Bioinformatics, 25(9), 1189-1191. https://doi.org/10.1093/bioinformatics/btp033

Wen, Y., Behiels, E., Felix, J., Elegheert, J., Vergauwen, B., Devreese, B., & Savvides, S. N. (2014,

Sep). The bacterial antitoxin HipB establishes a ternary complex with operator DNA and

phosphorylated toxin HipA to regulate bacterial persistence. *Nucleic Acids Research,*

42(15), 10134-10147. https://doi.org/10.1093/nar/gku665

Winn, M. D., Ballard, C. C., Cowtan, K. D., Dodson, E. J., Emsley, P., Evans, P. R., Keegan, R. M.,

Krissinel, E. B., Leslie, A. G., McCoy, A., McNicholas, S. J., Murshudov, G. N., Pannu, N.

S., Potterton, E. A., Powell, H. R., Read, R. J., Vagin, A., & Wilson, K. S. (2011, Apr).

Overview of the CCP4 suite and current developments. *Acta Crystallogr D Biol Crystallogr,*

67(Pt 4), 235-242. https://doi.org/10.1107/S0907444910045749

Winther, K., Tree, J. J., Tollervey, D., & Gerdes, K. (2016, Nov 16). VapCs of Mycobacterium

tuberculosis cleave RNAs essential for translation. *Nucleic Acids Research, 44*(20), 9860-

9871. https://doi.org/10.1093/nar/gkw781

Winther, K. S., Roghanian, M., & Gerdes, K. (2018, Apr 5). Activation of the Stringent Response

by Loading of RelA-tRNA Complexes at the Ribosomal A-Site. *Molecular Cell, 70*(1), 95-

105 e104. https://doi.org/10.1016/j.molcel.2018.02.033

Zielenkiewicz, U., & Ceglowski, P. (2005, Sep). The toxin-antitoxin system of the streptococcal

plasmid pSM19035. *Journal of Bacteriology, 187*(17), 6094-6105.

https://doi.org/10.1128/JB.187.17.6094-6105.2005

TABLE S1. Crystallographic data statistics

	HipBST[D233Q]	HipBST[S57A]	HipBST[S59A]
Data Collection			
Wavelength (Å)	1.54180	0.97625	1.54180
Resolution range (Å)	49.9 – 2.9 (3.0 – 2.9)*	46.9 – 2.4 (2.49 – 2.40)*	50.4 – 3.34 (3.5 – 3.4)*
Space group	C121	C121	C121
Unit cell dimensions			
a, b, c, (Å)	281.66, 106.47, 57.75	281.68, 106.07, 57.56	285.26, 107.15, 58.45
α, β, γ (º)	90, 90.75, 90	90, 90.65, 90	90, 90.71, 90
Total reflections	74,587 (7,335)	625,868 (25,597)	41,777 (4,068)
Unique reflections	37,716 (3,719)	66,117 (6,503)	22,580 (2,193)
Multiplicity	2.0 (2.0)	9.5 (3.9)	1.9 (1.9)
Completeness (%)	99.8 (99.3)	99.9 (99.4)	87.71 (84.84)
R_{meas} (%)	0.13 (1.105)	0.27 (0.08)	0.17 (1.01)
I/σ(I)	6.62 (0.86)	10.3 (0.72)	7.10 (0.93)
$CC_{1/2}$	0.99 (0.37)	0.99 (0.37)	0.97 (0.43)
Refinement			
Average B-factor (Å^2)	98.5	86.1	119.4
protein	98.9	86.5	119.9
ligands	-	166.7	118.6
solvent	80.0	76.7	78.3
No. of reflections	37,706 (3,719)	66,112 (6,504)	22,570 (2,193)
No. of reflections (free)	1,914 (175)	3,364 (326)	1086 (86)
R-work (%)	19.5 (31.8)	21.0 (42.8)	20.2 (29.2)
R-free (%)	22.8 (34.5)	23.8 (46.2)	24.3 (35.4)
Number of			
protein (residues)	1,018	1,017	1,023
solvent (atoms)	203	402	110
ligand (atoms)	-	20	10
rmsd (bonds, Å)	0.011	0.013	0.013
rmsd (angles, degrees)	1.49	1.75	1.66
Rotamer outliers (%)	9.8	6.0	10.07
Clashscore	4.4	2.7	5.8
Ramachandran statistics			
favoured (%)	95.1	96.0	95.6
allowed (%)	4.7	3.6	3.9
outliers (%)	0.2	0.4	0.5
Rama-Z score			
whole	-1.63	-0.44	-0.90
helix	-1.27	-0.40	-0.48
sheet	-1.36	0.21	0.07
loop	-0.75	-0.16	-0.73

*Numbers in parentheses refer to the outermost resolution shell.

TABLE S2. Bacterial strains and plasmids.

E. coli strains	Description	Reference or source
MG1655	Wild-type K12	(Blattner et al., 1997)
O127	O127:H6 strain E2348/69	(Iguchi et al., 2009)
TB28	MG1655Δ*lacIZYA*	Laboratory collection
Plasmid	**Description**[a]	**Reference or source**
pBAD33	p15 *araC* P_{BAD}, Cm^r	(Guzman et al., 1995)
pBAD322T	pMB1 *rop araC* P_{BAD}, Tet^r	(Cronan, 2006)
pNDM220	Mini-R1 *lacI*^{q} $P_{A1/04/03}$, Amp^r	(Gotfredsen & Gerdes, 1998)
pGH254	Mini-R1, *lacZYA* transcriptional fusion vector, Kan^r	Laboratory collection
pET-15b	pBR322 *lacI* P_{T7}, Amp^r	Novagen
pKG127	pUC57::*hipBST*$_{O127}$	(Vang Nielsen et al., 2019)
pSVN1	pBAD33::*hipT*, start codon GTG	(Vang Nielsen et al., 2019)
pSVN68	pUC57::*hipB-S-T*$^{S57A}_{His6}$, optimized SDs for all genes	This work
pSVN78	pET-15b::*hipB-S-T*$^{S57A}_{His6}$, optimized SDs for all genes	This work
pSVN87	pUC57::*hipB-S-T*$^{D210Q}_{His6}$, optimized SDs for all genes	This work

pSVN88	pUC57::$hipB$-S-T^{D233Q}_{His6}, optimized SDs for all genes	This work
pSVN96	pET-15b::$hipB$-S-T^{D233Q}_{His6}, optimized SDs for all genes	This work
pSVN109	pNDM220::$hipS$, optimized SD	(Vang Nielsen et al., 2019)
pSVN110	pNDM220::$hipB$-S, optimized SDs for both genes	(Vang Nielsen et al., 2019)
pSVN111	pNDM220::$hipB$, optimized SD	(Vang Nielsen et al., 2019)
pSVN141	pGH254::P$_{hipBST}$-$hipB$'-laZ, transcriptional P$_{hipBST}$-$hipB$' $lacZ$ fusion	This work
pSVN154	pBAD322T:: $hipB$-S-T^{D210Q}, optimized SDs for all genes	This work
pSVN178	pNDM220::$hipS^{W65A}$, optimized SD	This work
pSVN180	pBAD33::$hipB$-S-T^{D210Q}, optimized SDs for all genes	This work
pSVN181	pBAD33::$hipB$-S-T^{D233Q}, optimized SDs for all genes	This work
pSVN182	pBAD33::$hipB$-S, optimized SDs for both genes	This work
pSVN184	pBAD33::$hipB$-T^{D210Q}, optimized SDs for both genes	This work
pSVN185	pBAD33::$hipB$-T^{D233Q}, optimized SDs for both genes	This work
pSVN187	pBAD33::$hipS$-T^{D210Q}, optimized SDs for both genes	This work
pSVN188	pBAD33::$hipS$-T^{D233Q}, optimized SDs for both genes	This work
pSVN189	pBAD33::$hipB$, optimized SD	This work
pSVN190	pBAD33::$hipS$, optimized SD	This work

pSVN192	pBAD33::*hipT*^{D210Q}, optimized SD	This work
pSVN193	pBAD33::*hipT*^{D233Q}, optimized SD	This work
pSVN194	pBAD33::*hipT*^{S57D}, start codon GTG	This work
pSVN195	pBAD33::*hipT*^{S59D}, start codon GTG	This work
pSVN198	pGH254:: P$_{hipBST}$-*hipBST-lacZ*, transcriptional P$_{hipBST}$-*hipBST lacZ* fusion	This work
pSVN199	pBAD33::*hipT*^{S57A}, start codon GTG	This work
pSVN201	pBAD33::*hipT*^{S59A}, start codon GTG	This work
pSVN203	pBAD33::*hipT*$^{S57A+S59A}$, start codon GTG	This work
pSVN205	pBAD33::*hipT*$^{S57D+S59A}$, start codon GTG	This work
pSVN207	pBAD33::*hipT*$^{S57A+S59D}$, start codon GTG	This work
pSVN209	pBAD33::*hipT*$^{S57D+S59D}$, start codon GTG	This work
pSVN211	pBAD33::*hipT*^{D210Q}, start codon GTG	This work
pSVN212	pBAD33::*hipT*$^{S57D+D210Q}$, start codon GTG	This work
pSVN213	pBAD33::*hipT*$^{S59D+D210Q}$, start codon GTG	This work
pSVN214	pBAD33::*hipT*$^{S57A+D210Q}$, start codon GTG	This work
pSVN215	pBAD33::*hipT*$^{S59A+D210Q}$, start codon GTG	This work
pSVN216	pBAD33::*hipT*$^{S57A+S59A+D210Q}$, start codon GTG	This work

pSVN217	pBAD33::*hipT*$^{S57D+S59A+D210Q}$, start codon GTG	This work
pSVN218	pBAD33::*hipT*$^{S57A+S59D+D210Q}$, start codon GTG	This work
pSVN219	pBAD33::*hipT*$^{S57D+S59D+D210Q}$, start codon GTG	This work
pSVN220	pBAD33::*hipT*^{D210Q}, optimized SD	This work
pSVN221	pBAD33::*hipT*$^{S57D+D210Q}$, optimized SD	This work
pSVN222	pBAD33::*hipT*$^{S59D+D210Q}$, optimized SD	This work
pSVN223	pBAD33::*hipT*$^{S57A+D210Q}$, optimized SD	This work
pSVN224	pBAD33::*hipT*$^{S59A+D210Q}$, optimized SD	This work
pSVN225	pBAD33::*hipT*$^{S57A+S59A+D210Q}$, optimized SD	This work
pSVN226	pBAD33::*hipT*$^{S57D+S59A+D210Q}$, optimized SD	This work
pSVN227	pBAD33::*hipT*$^{S57A+S59D+D210Q}$, optimized SD	This work
pSVN228	pBAD33::*hipT*$^{S57D+S59D+D210Q}$, optimized SD	This work
pSNN1	pET-15b::*hipT*$^{S57A}_{His6}$, optimized SD	This work
pSNN2	pET-15b::*hipT*$^{S57A+D210A}_{His6}$, optimized SD	This work
pMME	pET-15b::*hipB-S-T*$^{S59A}_{His6}$, optimized SDs for all genes	This work

^{a}SD, Shine-Dalgarno sequence.

TABLE S3. Oligonucleotides and primers.

Oligonucleotide	Sequence
FP1(GTG)	CCCCGTCGACGGATCCAAGGAGTTTTATAAGTGGCGAATTGTCGTATTCTG
FP21	GGGGGGTACCGGATCCAAAATAAGGAGGAAAAAAAAATGATCTGCTCAGGACCAC
FP22	CCCCCTCGAGGGATCCAAAATAAGGAGGAAAAAAAAATGCATCGGCGAGTGAAAG
FP43	CCCCGAATTCCTCTCCCGATGAGATCAGC
FP46	GGGGGGTACCGGATCCAAAATAAGGAGGAAAAAAAAATGCATCGGCGAGTGAAAG
FP47	GGGGGGTACCGGATCCAAAATAAGGAGGAAAAAAAAATGGCGAATTGTCGTATTCTG
FP48	GGGGGTCGACCTGCAGAAAATAAGGAGGAAAAAAAAATGGCGAATTGTCGTATTCTG
FP49	CCCCGAATTCCTCTCCCG ATGAGATCAGCTTTC
RP1	CCCCCGCATGCGAATTCGCTCACAGCAGCCCCAGACG
RP11	CCCCCTCGAGAAGCTTTCACAGCAGCCCCAGACG
RP14	GGGGGAATTCAAGCTTTTATTCCTCCCAAGGTAAAATC
RP15	GGGGGAATTCAAGCTTTCACTCGCCGATGCATAG
RP32	CCCCGGATCCTCTGCAACTCCTGGAGTTG
RP35	GTCCTGAGCAGATCATTAAAGAACC

RP37	CGATTGATAAAAGACAAAAACAAGATC
RP38	GTTGGTCGCCGATCGCTTTC
RP42	GGGGGTCGACCTGCAGTCACTCGCCGATGCATAG
RP43	GGGGGGTACCTCACAGCA GCCCCAGACG
hipS W65A Fw	CAGAAGGAGCTCTGCGTCAACGCTA
hipS W65A Rv	TGACGCAGAGCTCCTTCTGGCGC
HipT S57D Fw	GCGTCAACAAAAAGGGATGGATATTTCCGGTT
HipT S57D Rv	GGGCTGGTAACCGGAAATATCCATCCCTTTTT
HipT S59D Fw	GTCAACAAAAAGGGATGAGTATTGACGGTTAC
HipT S59D Rv	TTGGGCTGGTAACCGTCAATACTCATCCCTTT
HipT S59A Fw	GTCAACAAAAAGGGATGAGTATTGCCGGTTAC
HipT S59A Rv	TTGGGCTGGTAACCGGCAATACTCATCCCTTT
HipT S57A + S59A Fw	CGTCAACAAAAAGGGATGGCTATTGCCGGTTAC
HipT S57A + S59A Rv	TTGGGCTGGTAACCGGCAATAGCCATCCCTTTT
HipT S57D + S59A Fw	CGTCAACAAAAAGGGATGGATATTGCCGGTTAC
HipT S57D + S59A Rv	TTGGGCTGGTAACCGGCAATATCCATCCCTTTT

HipT S57A + S59D Fw	CGTCAACAAAAAGGGATGGCTATTGACGGTTAC
HipT S57A + S59D Rv	TTGGGCTGGTAACCGTCAATAGCCATCCCTTTT
HipT S57D + S59D Fw	CGTCAACAAAAAGGGATGGATATTGACGGTTAC
HipT S57D + S59D Rv	TTGGGCTGGTAACCGTCAATATCCATCCCTTTT
hipX S57A Fw	AAGGGATGGCTATTTCCGGTTACCAGCC
hipX S57A Rv	CGGAAATAGCCATCCTTTTTGTTGACG
hipX D210Q Fw	GTAATAACCAGATGCATTTACGAAACTTTGG
hipX D210Q Rv	AAATGCATCTGGTTATTACCCAGCAACCAGG
hipX D233Q Fw	CGGTGTATCAGTTTGTTTCTGTCGCTCCC
hipX D233Q Rv	GAAACAAACTGATACACCGGCGCTAACG
hipBS del Fw	ACGACAATTCGCCATTTTTTTTCCTCCTTATTTTTCTAGAGGG
hipBS del Rv	TTCCCCTCTAGAAAAATAAGGAGGAAAAAAAAATGGCGAAT
hipT D210A Fw	TAAATGCATCGCGTTATTACCCAGCAACAA
hipT D210A Rv	CTGGGTAATAACGCGATGCATTTACGAAACTTT
hipT S57S59A Fw	GGGATGAGTATTGCCGGTTACCAGCCCAAATTGCAA
hipT S57S59A Rv	GTAACCGGCAATACTCATCCCTTTTTGTTGACGCGG

SUPPLEMENTARY METHODS

Construction of plasmids. Construction of plasmids is summarised below.

pSVN68. The mutation in *hipB-S-T*^{S57A}$_{His6}$ with optimized SDs for all three genes was created

using pSVN61 (Vang Nielsen et al., 2019) and primers hipX S57A Fw and hipX S57A Rv in

a site-directed plasmid mutagenesis PCR. Eight reactions were carried out at different

temperatures with a gradient PCR. The samples were pooled and digested with DpnI to digest

the parental plasmid before being transformed into *E. coli* DH5α.

pSVN78. *hipB-S-T*^{S57A}$_{His6}$ with optimized SDs for all three genes was sub-cloned from

pSVN68 by digesting with XbaI and XhoI, purifying the DNA fragment and ligating into pET-

15b.

pSVN87. The mutation in *hipB-S-T*^{D210Q}$_{His6}$ with optimized SDs for all three genes was created

using pSVN61 (Vang Nielsen et al., 2019) and primers hipX D210Q Fw and hipX D210Q Rv

in a site-directed plasmid mutagenesis PCR. Eight reactions were carried out at different

temperatures with a gradient PCR. The samples were pooled and digested with DpnI to digest

the parental plasmid before being transformed into *E. coli* DH5α.

pSVN88. The mutation in *hipB-S-T*^{D233Q}$_{His6}$ with optimized SDs for all three genes was created

using pSVN61 (Vang Nielsen et al., 2019) and primers hipX D233Q Fw and hipX D233Q Rv

in a site-directed plasmid mutagenesis PCR. Eight reactions were carried out at different

temperatures with a gradient PCR. The samples were pooled and digested with DpnI to digest

the parental plasmid before being transformed into *E. coli* DH5α.

pSVN96. *hipB-S-T*^{D233Q}$_{His6}$ with optimized SDs for all three genes was sub-cloned from

pSVN88 by digesting with XbaI and XhoI, purifying the DNA fragment and ligating into pET-

15b.

pSVN141. P_{hipBST}-*hipB*', a fragment containing 224 bp upstream of the *hipB* gene plus the first

73 bp of the *hipB* gene was amplified from pKG127 using primers FP43 and RP32. The

resulting PCR product was digested with EcoRI and BamHI and ligated into pGH254.

pSVN154. *hipB-S-T*^{D210Q} with optimized SDs for all three genes was amplified from pSVN87

using primers FP21 and RP11. The resulting PCR product was digested with KpnI and HindIII

and ligated into pBAD322T.

pSVN178. *hipS*^{W65A} was created by a two-step PCR reaction. Two fragments were amplified

from pSVN109 using primers FP22 and hipS W65A Rv in one reaction and hipS W65A Fw

and RP14 in the other. The resulting two PCR products were joined by a second round of PCR

using both fragments as template DNA and primers FP22 and RP14. The final PCR product

was digested with XhoI and EcoRI and ligated into pNDM220.

pSVN180. *hipB-S-T*^{D210Q} with optimized SDs for all three genes was amplified from pSVN87

using primers FP21 and RP11. The resulting PCR product was digested with KpnI and HindIII

and ligated into pBAD33.

pSVN181. *hipB-S-T*^{D233Q} with optimized SDs for all three genes was amplified from pSVN88

using primers FP21 and RP11. The resulting PCR product was digested with KpnI and HindIII

and ligated into pBAD33.

pSVN182. *hipB-S* with optimized SDs for both genes was amplified from pSVN87 using

primers FP21 and RP14. The resulting PCR product was digested with KpnI and HindIII and

ligated into pBAD33.

pSVN184. *hipB-T*^{D210Q} with optimized SDs for both genes was created by amplifying two PCR

products from pSVN87: *hipB* using primers FP21 and RP42 and *hipT*^{D210Q} using primers FP48

and RP11. The *hipB* fragment was digested with KpnI and PstI, while the *hipT*^{D233Q} fragment

was digested with PstI and HindIII. The two digested fragments were then ligated into pBAD33

using the KpnI and HindIII restriction sites.

pSVN185. *hipB-T*^{D233Q} with optimized SDs for both genes was created by amplifying two PCR

products from pSVN88: *hipB* using primers FP21 and RP42 and *hipT*^{D233Q} using primers FP48

and RP11. The *hipB* fragment was digested with KpnI and PstI, while the *hipT*^{D233Q} fragment

was digested with PstI and HindIII. The two digested fragments were then ligated into pBAD33

using the KpnI and HindIII restriction sites.

pSVN187. *hipS-T*^{D210Q} with optimized SDs for both genes was amplified from pSVN88 using

primers FP46 and RP11. The resulting PCR product was digested with KpnI and HindIII and

ligated into pBAD33.

pSVN188. *hipS-T*^{D233Q} with optimized SDs for both genes was amplified from pSVN87 using

primers FP46 and RP11. The resulting PCR product was digested with KpnI and HindIII and

ligated into pBAD33.

pSVN189. *hipB* with optimized SD was amplified from pSVN87 using primers FP21 and

RP15. The resulting PCR product was digested with KpnI and HindIII and ligated into

pBAD33.

pSVN190. *hipS* with optimized SD was amplified from pSVN87 using primers FP46 and

RP14. The resulting PCR product was digested with KpnI and HindIII and ligated into

pBAD33.

pSVN192. *hipT*^{D210Q} with optimized SD was amplified from pSVN87 using primers FP47 and

RP11. The resulting PCR product was digested with KpnI and HindIII and ligated into

pBAD33.

pSVN193. *hipT*^{D233Q} with optimized SD was amplified from pSVN88 using primers FP47 and

RP11. The resulting PCR product was digested with KpnI and HindIII and ligated into

pBAD33.

pSVN194. *hipT*^{S57D} with start codon GTG was created by a two-step PCR reaction. Two

fragments were amplified from pKG127 using primers FP1(GTG) and HipT S57D Rv in one

reaction and HipT S57D Fw and RP1 in the other. The resulting two PCR products were joined

by a second round of PCR using both fragments as template DNA and primers FP1(GTG) and

RP1. The final PCR product was digested with SalI and SphI and ligated into pBAD33.

pSVN195. *hipT*^{S59D} with start codon GTG was created by a two-step PCR reaction. Two

fragments were amplified from pKG127 using primers FP1(GTG) and HipT S59D Rv in one

reaction and HipT S59D Fw and RP1 in the other. The resulting two PCR products were joined

by a second round of PCR using both fragments as template DNA and primers FP1(GTG) and

RP1. The final PCR product was digested with SalI and SphI and ligated into pBAD33.

pSVN198. P$_{hipBST}$-*hipBST*, a fragment containing 224 bp upstream of the *hipB* gene plus the

native *hipBST* operon was amplified from pKG127 using primers FP49 and RP43. The

resulting PCR product was digested with EcoRI and KpnI and ligated into pGH254.

pSVN199. *hipT*^{S57A} with start codon GTG was amplified from pSVN78 using primers

FP1(GTG) and RP1. The resulting PCR product was digested with SalI and SphI and ligated

into pBAD33.

pSVN201. *hipT*^{S59A} with start codon GTG was created by a two-step PCR reaction. Two

fragments were amplified from pKG127 using primers FP1(GTG) and HipT S59A Rv in one

reaction and HipT S59A Fw and RP1 in the other. The resulting two PCR products were joined

by a second round of PCR using both fragments as template DNA and primers FP1(GTG) and

RP1. The final PCR product was digested with SalI and SphI and ligated into pBAD33.

pSVN203. *hipT*$^{S57A+S59A}$ with start codon GTG was created by a two-step PCR reaction. Two

fragments were amplified from pKG127 using primers FP1(GTG) and HipT S57A + S59A Rv

in one reaction and HipT S57A + S59A Fw and RP1 in the other. The resulting two PCR

products were joined by a second round of PCR using both fragments as template DNA and

primers FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and

ligated into pBAD33.

99 **pSVN205.** *hipT*$^{S57D+S59A}$ with start codon GTG was created by a two-step PCR reaction. Two

100 fragments were amplified from pKG127 using primers FP1(GTG) and HipT S57D + S59A Rv

101 in one reaction and HipT S57D + S59A Fw and RP1 in the other. The resulting two PCR

102 products were joined by a second round of PCR using both fragments as template DNA and

103 primers FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and

104 ligated into pBAD33.

105 **pSVN207.** *hipT*$^{S57A+S59D}$ with start codon GTG was created by a two-step PCR reaction. Two

106 fragments were amplified from pKG127 using primers FP1(GTG) and HipT S57A + S59D Rv

107 in one reaction and HipT S57A + S59D Fw and RP1 in the other. The resulting two PCR

108 products were joined by a second round of PCR using both fragments as template DNA and

109 primers FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and

110 ligated into pBAD33.

111 **pSVN209.** *hipT*$^{S57D+S59D}$ with start codon GTG was created by a two-step PCR reaction. Two

112 fragments were amplified from pKG127 using primers FP1(GTG) and HipT S57D + S59D Rv

113 in one reaction and HipT S57D + S59D Fw and RP1 in the other. The resulting two PCR

114 products were joined by a second round of PCR using both fragments as template DNA and

115 primers FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and

116 ligated into pBAD33.

117 **pSVN211.** *hipT*^{D210Q} with start codon GTG was created by a two-step PCR reaction. Two

118 fragments were amplified from pSVN1 using primers FP1(GTG) and hipX D210Q Rv in one

119 reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products were

120 joined by a second round of PCR using both fragments as template DNA and primers

121 FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

122 pBAD33.

pSVN212. *hipT*$^{S57D+D210Q}$ with start codon GTG was created by a two-step PCR reaction. Two

fragments were amplified from pSVN194 using primers FP1(GTG) and hipX D210Q Rv in

one reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products were

joined by a second round of PCR using both fragments as template DNA and primers

FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

pBAD33.

pSVN213. *hipT*$^{S59D+D210Q}$ with start codon GTG was created by a two-step PCR reaction. Two

fragments were amplified from pSVN195 using primers FP1(GTG) and hipX D210Q Rv in

one reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products were

joined by a second round of PCR using both fragments as template DNA and primers

FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

pBAD33.

pSVN214. *hipT*$^{S57A+D210Q}$ with start codon GTG was created by a two-step PCR reaction. Two

fragments were amplified from pSVN199 using primers FP1(GTG) and hipX D210Q Rv in

one reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products were

joined by a second round of PCR using both fragments as template DNA and primers

FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

pBAD33.

pSVN215. *hipT*$^{S59A+D210Q}$ with start codon GTG was created by a two-step PCR reaction. Two

fragments were amplified from pSVN201 using primers FP1(GTG) and hipX D210Q Rv in

one reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products were

joined by a second round of PCR using both fragments as template DNA and primers

FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

pBAD33.

pSVN216. *hipT*$^{S57A+S59A+D210Q}$ with start codon GTG was created by a two-step PCR reaction.

Two fragments were amplified from pSVN203 using primers FP1(GTG) and hipX D210Q Rv

in one reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products

were joined by a second round of PCR using both fragments as template DNA and primers

FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

pBAD33.

pSVN217. *hipT*$^{S57D+S59A+D210Q}$ with start codon GTG was created by a two-step PCR reaction.

Two fragments were amplified from pSVN205 using primers FP1(GTG) and hipX D210Q Rv

in one reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products

were joined by a second round of PCR using both fragments as template DNA and primers

FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

pBAD33.

pSVN218. *hipT*$^{S57A+S59D+D210Q}$ with start codon GTG was created by a two-step PCR reaction.

Two fragments were amplified from pSVN207 using primers FP1(GTG) and hipX D210Q Rv

in one reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products

were joined by a second round of PCR using both fragments as template DNA and primers

FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

pBAD33.

pSVN219. *hipT*$^{S57D+S59D+D210Q}$ with start codon GTG was created by a two-step PCR reaction.

Two fragments were amplified from pSVN209 using primers FP1(GTG) and hipX D210Q Rv

in one reaction and hipX D210Q Fw and RP1 in the other. The resulting two PCR products

were joined by a second round of PCR using both fragments as template DNA and primers

FP1(GTG) and RP1. The final PCR product was digested with SalI and SphI and ligated into

pBAD33.

pSVN220. *hipT*^{D210Q} with optimized SD was amplified from pSVN211 using primers FP48

and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSVN221. *hipT*^{S57D+D210Q} with optimized SD was amplified from pSVN212 using primers

FP48 and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSVN222. *hipT*^{S59D+D210Q} with optimized SD was amplified from pSVN213 using primers

FP48 and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSVN223. *hipT*^{S57A+D210Q} with optimized SD was amplified from pSVN214 using primers

FP48 and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSVN224. *hipT*^{S59A+D210Q} with optimized SD was amplified from pSVN215 using primers

FP48 and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSVN225. *hipT*^{S57A+S59A+D210Q} with optimized SD was amplified from pSVN216 using primers

FP48 and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSVN226. *hipT*^{S57D+S59A+D210Q} with optimized SD was amplified from pSVN217 using primers

FP48 and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSVN227. *hipT*^{S57A+S59D+D210Q} with optimized SD was amplified from pSVN218 using primers

FP48 and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSVN228. *hipT*$^{S57D+S59D+D210Q}$ with optimized SD was amplified from pSVN219 using primers

FP48 and RP11. The resulting PCR product was digested with PstI and HindIII and ligated into

pBAD33.

pSNN1. *hipT*^{S57A} with optimized SD was amplified from pSVN78 using primers hipBS del Fw

and hipBS del Rv. This resulted in the deletion of HipB and HipS.

pSNN2. *hipT*$^{S57A+D210A}$ with optimized SD was amplified from pSNN1 using primers hipT

D210A Fw and hipT D210A Rv.

pMME3. *hipBST*^{S59A} with optimized SD was amplified from pSVN78 using primers hipT

S57S59A Fw and hipT S57S59A Rv to introduce the mutations A57S and S59A.

 SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Sequence alignment with consensus structural elements of HipBST, HipBA$_{Ec}$

and HipBA$_{So}$. a. Alignment of *E. coli* O127:H6 HipB with HipB from *E. coli* K-12 HipBA

and HipB from *S. oneidensis* HipBA. Secondary structure observed in *E. coli* O127:H6 HipB

is shown above the sequences, except for α1*, which is missing from the crystal structures but

predicted by JPred4 (Drozdetskiy et al., 2015). **b.** Alignment of *E. coli* O127:H6 HipS with the

N-terminal subdomain 1 from *E. coli* K-12 HipA and *S. oneidensis* HipA. The position of the

the Gly-rich interacting helix (α3) of HipS is indicated. Secondary structure observed in *E. coli*

O127:H6 HipS is shown above the sequences **c.** Alignment of *E. coli* O127:H6 HipT with the

main kinase domain of *E. coli* K-12 HipA and *S. oneidensis* HipA. The position of the Gly-

rich loop (residue 58-63) and phosphorylation sites (Ser57 and Ser59), catalytic motif

(Asp210), and Mg^{2+}-binding motif (Asp233), are indicated. Secondary structure observed in

E. coli O127:H6 HipT is shown above the sequences.

Figure S2. Biophysical analysis of the HipBST complex. a. Sequence alignment of *E. coli*

O127:H6 and selected HipS orthologues with the N-terminal subdomain 1 of *E. coli* K-12 HipA

(bottom). Secondary structure as observed for *E. coli* O127:H6 HipS is indicated above the

alignment. Fully conserved residues are shown on a dark orange background, partially

conserved residues in orange text, and regions that interact with HipT are shown on a light grey

background. Trp65 that intercalates in HipT is shown on a blue background. **b.** Size exclusion

chromatography profile as observed for HipBST^{D233Q}. A similar elution profile was observed

for the other HipBST variants used for structural studies. Absorption at 280 (A$_{280}$, blue) and

260 (A$_{260}$, red) nm are shown along with elution volumes of a range of standard proteins (6.5-

440 kDa, as indicated). **c.** The structure of *S. oneidensis* HipBA (PDB: 4PU3, HipA purple,

HipB green) bound to DNA (orange backbone, blue bases) (Wen et al., 2014).

Figure S3. Interactions of HipT with HipS and HipB. a. Overall fold of HipS with secondary structure elements labelled (α helices in red and β sheets in yellow). **b.** Structural comparison between *E. coli* O127:H6 HipB (cartoon, two shades of green) and the corresponding homodimer of *S. oneidensis* HipB (PDB: 4PU3, ribbons, cyan) showing their conserved structure with similar HTH folds (Wen et al., 2014). **c.** Top, structural alignment of the *E. coli* O127:H6 HipBST complex (HipT, blue; HipS, beige; HipB, green) with *S. oneidensis* HipBA bound to DNA (PDB: 4PU3; HipA, purple; HipB, green) (Wen et al., 2014); Below, close-up views of conserved, basic residues on HipA that contact the DNA backbone (left) and the corresponding region in HipT (right). **d.** Growth curves of *E. coli* MG1655 grown in YT medium and harbouring empty pBAD33 vector ("pBAD33") or pSVN1 (pBAD33::*hipT*, "*hipT*"), in combination with empty pNDM220 vector ("pNDM220"), pSVN109 (pNDM220::*hipS*, "*hipS*"), or pSVN178 (pNDM220::*hipS*[W65A], "*hipS*[W65A]") as indicated. At the indicated times, 0.2% arabinose was added to induce *hipT* (long arrow) and 200 μM IPTG was added to induce *hipS* or *hipS*[W65A] (short arrow). The data points represent mean values of results from at least three independent experiments, and error bars show standard deviations (hidden when small).

Figure S4. HipT autophosphorylation at Ser57 and Ser59 affects function differentially.
a. Active site region and Gly-rich loop (ruby) of *E. coli* O127:H6 HipT from the HipBST[S57A] structure showing the 2mFo-DFc difference electron density covering Ser59, contoured at 1.2 σ. The phosphate group was refined to an occupancy of 0.89. **b.** Similar view from the HipBST[S59A] structure showing the 2mFo-DFc difference electron density covering Ser57, contoured at 1.0 σ. The phosphate group was refined to an occupancy of 0.72. **c.** Top, structural alignment of the active sites of *E. coli* O127:H6 HipT S57A (blue) with *E. coli* K-12 HipA

(HipA$_{Ec}$, purple) highlighting the positions of their respective phosphoserine (P-Ser) residues,

as indicated; Bottom, equivalent structural alignment of the active sites of *E. coli* O127:H6

HipT S59A (blue) with *S. oneidensis* HipA (HipA$_{So}$, pale blue). **d.** Growth curves of *E. coli*

MG1655 strains grown in YT medium and harbouring empty pBAD33 vector, pSVN1

(pBAD33::*hipT*, "*hipT* (SIS, *wt*)"), or single autophosphorylation mutants pSVN194

(pBAD33::*hipT*^{S57D}, "*hipT* (DIS)"), pSVN195 (pBAD33::*hipT*^{S59D}, "*hipT* (SID)"), pSVN199

(pBAD33::*hipT*^{S57A}, "*hipT* (AIS)"), or pSVN201 (pBAD33::*hipT*^{S59A}, "*hipT* (SIA)") in

combination with pSVN110 (pNDM220::*hipB-S*, "*hipBS*"). At the indicated times, 0.2%

arabinose was added to induce *hipT* (long arrow) and 200 μM was added to induce *hipBS* (short

arrow). The data points represent mean values of results from two independent experiments,

and error bars show standard deviations (hidden when small). **e.** As in **d** but for double

autophosphorylation mutants pSVN203 (pBAD33::*hipT*$^{S57A+S59A}$, "*hipT* (AIA)"), pSVN205

(pBAD33::*hipT*$^{S57D+S59A}$, "*hipT* (DIA)"), pSVN207 (pBAD33::*hipT*$^{S57A+S59D}$, "*hipT* (AID)"),

and pSVN209 (pBAD33::*hipT*$^{S57D+S59D}$, "*hipT* (DID)"). Data points represent mean values of

results from two independent experiments, and error bars designate standard deviations. **f.**

Northern blot analysis using probe directed against tRNA^{gltTUVW} (top) and tRNA^{argVYZQ}

(bottom) based on RNA extracted from *E. coli* MG1655 harbouring pSVN110

(pNDM220::*hipB-S*) in combination with pSVN1 (pBAD33::*hipT*, "*hipT* (S^{57}IS^{59}, *wt*)"),

pSVN194 (pBAD33::*hipT*^{S57D}, "*hipT* (D^{57}IS^{59})"), pSVN195 (pBAD33::*hipT*^{S59D}, "*hipT*

(S^{57}ID^{59})"), pSVN209 (pBAD33::*hipT*$^{S57D+S59D}$, "*hipT* (D^{57}ID^{59})") or pSVN205

(pBAD33::*hipT*$^{S57D+S59A}$, "*hipT* (D^{57}IA^{59})") at indicated time points in minutes before (-) and

after (+) addition of 0.2% arabinose to induce wildtype *hipT* or autophosphorylation mutants.

Results from strains harbouring pSVN1 (pBAD33::*hipT*), pSVN194 (pBAD33::*hipT*^{S57D}),

pSVN209 and (pBAD33::*hipT*$^{S57D+S59D}$) are representative of two independent experiments,

whereas results from strains harbouring pSVN195 (pBAD33::*hipT*[S59D]) and pSVN205

(pBAD33::*hipT*[S57D+S59A]) are from single experiments.

Figure S5. Transcriptional regulation of the HipBST system of *E. coli* O127:H6. a.

Transcriptional reporter assay using a plasmid with the *hipBST* promoter region and a 5'

fragment of the *hipB* gene (including 224 bp upstream of *hipB* plus the first 73 bp of the *hipB*

gene, shown with a lightening symbol) transcriptionally fused to *lacZ*. *E. coli* TB28 harbouring

the reporter plasmid (pSVN141, pGH254::P$_{hipBST}$-*hipB*'-*lacZ*) and empty pBAD33 vector or

combinations of *hipB*, *hipS*, and *hipT* in the context of the HipT[D233Q] inactive mutant; pSVN180

(pBAD33::*hipB-S-T*[D233Q], "*hipBST*[D233Q]"), pSVN182 (pBAD33::*hipB-S*, "*hipBS*"), pSVN184

(pBAD33::*hipB-T*[D233Q], "*hipBT*[D233Q]"), pSVN187 (pBAD33::*hipS-T*[D233Q], "*hipST*[D233Q]"),

pSVN189 (pBAD33::*hipB*, "*hipB*"), pSVN190 (pBAD33::*hipS*, "*hipS*"), or pSVN192

(pBAD33::*hipT*[D233Q], "*hipT*[D233Q]") were grown, diluted, and spotted onto YT agar plates

containing 40 µg/ml X-gal and 0.2% glucose (to repress *hipB/S/T*[D233Q]) or 0.2% arabinose (to

induce *hipB/S/T*[D233Q]). Results are representative of two independent experiments. **b.**

Overexpression of inactive HipT[D210Q] and HipT[D233Q] increases the level of active wildtype

HipT from the *hipBST* operon transcriptionally fused to *lacZ* in *E. coli* TB28. The strain

harbouring empty pGH254 vector or pSVN198 (pGH254::P$_{hipBST}$-*hipBST*-*lacZ*, "*hipBST*") in

combination with the empty pBAD33 vector, pSVN192 (pBAD33::*hipT*[D210Q], "*hipT*[D210Q]"), or

pSVN193 (pBAD33::*hipT*[D233Q], "*hipT*[D233Q]"), as indicated, were grown, diluted, and spotted

onto YT agar plates containing 40 µg/ml X-gal without 0.2% arabinose (to obtain only a leaky

expression of *hipT*[D210Q/D233Q]), or with 0.2% arabinose (to induce *hipT*[D210Q/D233Q]). Results are

from a single experiment. **c.** Primer extension analysis mapping the transcriptional start site in

E. coli TB28 harbouring pSVN141 (pGH254::P$_{hipBST}$-*hipB'*-*lacZ*), in combination with the

empty pBAD322T vector, or pSVN154 (pBAD322T::*hipB-S-T*[D210Q]), as indicated were grown

and harvested at $OD_{600}\approx0.4$. **d.** Primer extension analysis mapping the transcriptional start site in *E. coli* O127:H6 harbouring pSVN192 (pBAD33::*hipT*[D210Q]). At $OD_{600}\approx0.4$, 0.2% arabinose was added (to induce *hipT*[D210Q]), samples were taken at indicated time points and primer extension performed. **e.** The *hipBST* promoter region showing the transcriptional start site (arrow) and corresponding -10 and -35 sequences (red). The start codon of the *hipB* gene ("Met") and putative Shine-Dalgarno sequence ("SD") are indicated as well. **f.** Spot assays using *E. coli* MG1655 harbouring pSVN111 (pNDM220::*hipB*) in combination with the empty pBAD33 vector, pSVN211 (pBAD33::*hipT*[D210Q], "*hipT* ($S^{57}IS^{59}$, *wt*)"), pSVN212 (pBAD33::*hipT*[S57D+D210Q], "*hipT* ($D^{57}IS^{59}$)"), pSVN213 (pBAD33::*hipT*[S59D+D210Q], "*hipT* ($S^{57}ID^{59}$)"), pSVN214 (pBAD33::*hipT*[S57A+D210Q], "*hipT* ($A^{57}IS^{59}$)"), pSVN215 (pBAD33::*hipT*[S59A+D210Q], "*hipT* ($S^{57}IA^{59}$)"), pSVN216 (pBAD33::*hipT*[S57A+S59A+D210Q], "*hipT* ($A^{57}IA^{59}$)"), pSVN217 (pBAD33::*hipT*[S57D+S59A+D210Q], "*hipT* ($D^{57}IA^{59}$)"), pSVN218 (pBAD33::*hipT*[S57A+S59D+D210Q], "*hipT* ($A^{57}ID^{59}$)"), or pSVN219 (pBAD33::*hipT*[S57D+S59D+D210Q], "*hipT* ($D^{57}ID^{59}$)") spotted onto YT agar plates containing 0.2% glucose (to repress *hipT*[D210Q] autophosphorylation mutants), 0.2% arabinose (to induce *hipT*[D210Q] autophosphorylation mutants) or 0.2% arabinose plus 500 µM IPTG (to induce *hipB*). Results are from a single experiment.

Figure S6. Analysis of HipBST using small-angle x-ray scattering. a. Background-subtracted and normalized SAXS data measured for HipBST[S57A] with indirect Fourier transform fit. **b.** Guinier plot. **c.** $p(r)$ from indirect Fourier transform. **d.** Normalized Kratky plot using $I(0)$ and R_g from the Guinier fit.

Figure S7. HipT is structurally similar to eukaryotic cyclin-dependent kinases (Cdks). a. Domain overview (top) and secondary structure comparison (bottom) between *E. coli* O127:H6

HipT and human CDK5 showing the unique N-terminal mini-domain found in HipT (dashed)

and that the N-terminal subdomain 2 is similar to the N-terminal lobe of CDK5 while the core

kinase domain is similar to the C-terminal lobe. Conserved features and active site residues are

marked with numbers and listed below. **b.** Structural comparison of *E. coli* O127:H6 HipT

(top) and human CDK5 (PDB: 4AU8, bottom) (Malmstrom et al., 2012) shown in rainbow

colours from N (blue) to C (red) terminus. α-helices are shown as cylinders and for CDK5

named in accordance with classical Cdk nomenclature. The two proteins differ in tertiary fold

despite similar topology. **c.** Top, overview structure of Human CDK9 (salmon, PDB: 3BLT)

with the C-spine region shown as surface (cyan) (Baumli et al., 2008); below, details of the

active site bound to ATP with relevant residues indicated. **d.** Top, overview of the *E. coli*

O127:H6 HipT structure with the C/R spine region shown as surface (green); below, details of

the active site. **e.** Top, overview of the *E. coli* K-12 HipA structure (PDB: 3DNT) with the C/R

spine region shown as surface (orange) (Schumacher et al., 2009); below, details of the active

site bound to ATP.

REFERENCES

Baumli, S., Lolli, G., Lowe, E. D., Troiani, S., Rusconi, L., Bullock, A. N., Debreczeni, J. E.,
Knapp, S., & Johnson, L. N. (2008, Jul 9). The structure of P-TEFb (CDK9/cyclin T1),
its complex with flavopiridol and regulation by phosphorylation. *EMBO Journal,*
27(13), 1907-1918. https://doi.org/10.1038/emboj.2008.121

Blattner, F. R., Plunkett, G., 3rd, Bloch, C. A., Perna, N. T., Burland, V., Riley, M., Collado-
Vides, J., Glasner, J. D., Rode, C. K., Mayhew, G. F., Gregor, J., Davis, N. W.,
Kirkpatrick, H. A., Goeden, M. A., Rose, D. J., Mau, B., & Shao, Y. (1997, Sep 5). The
complete genome sequence of Escherichia coli K-12. *Science, 277*(5331), 1453-1462.
https://doi.org/10.1126/science.277.5331.1453

Cronan, J. E. (2006, Mar). A family of arabinose-inducible Escherichia coli expression vectors
having pBR322 copy control. *Plasmid, 55*(2), 152-157.
https://doi.org/10.1016/j.plasmid.2005.07.001

Drozdetskiy, A., Cole, C., Procter, J., & Barton, G. J. (2015, Jul 1). JPred4: a protein secondary
structure prediction server. *Nucleic Acids Research, 43*(W1), W389-394.
https://doi.org/10.1093/nar/gkv332

Gotfredsen, M., & Gerdes, K. (1998, Aug). The Escherichia coli relBE genes belong to a new
toxin-antitoxin gene family. *Molecular Microbiology, 29*(4), 1065-1076.
https://doi.org/10.1046/j.1365-2958.1998.00993.x

Guzman, L. M., Belin, D., Carson, M. J., & Beckwith, J. (1995, Jul). Tight regulation,
modulation, and high-level expression by vectors containing the arabinose PBAD
promoter. *Journal of Bacteriology, 177*(14), 4121-4130.
https://doi.org/10.1128/jb.177.14.4121-4130.1995

Iguchi, A., Thomson, N. R., Ogura, Y., Saunders, D., Ooka, T., Henderson, I. R., Harris, D.,
Asadulghani, M., Kurokawa, K., Dean, P., Kenny, B., Quail, M. A., Thurston, S.,
Dougan, G., Hayashi, T., Parkhill, J., & Frankel, G. (2009, Jan). Complete genome
sequence and comparative genome analysis of enteropathogenic *Escherichia coli*
O127:H6 strain E2348/69. *Journal of Bacteriology, 191*(1), 347-354.
https://doi.org/10.1128/JB.01238-08

Malmstrom, J., Viklund, J., Slivo, C., Costa, A., Maudet, M., Sandelin, C., Hiller, G., Olsson,
L. L., Aagaard, A., Geschwindner, S., Xue, Y., & Vasange, M. (2012, Sep 15).
Synthesis and structure-activity relationship of 4-(1,3-benzothiazol-2-yl)-thiophene-2-
sulfonamides as cyclin-dependent kinase 5 (cdk5)/p25 inhibitors. *Bioorganic and*
Medicinal Chemistry Letters, 22(18), 5919-5923.
https://doi.org/10.1016/j.bmcl.2012.07.068

Schumacher, M. A., Piro, K. M., Xu, W., Hansen, S., Lewis, K., & Brennan, R. G. (2009, Jan
16). Molecular mechanisms of HipA-mediated multidrug tolerance and its
neutralization by HipB. *Science, 323*(5912), 396-401.
https://doi.org/10.1126/science.1163806

Vang Nielsen, S., Turnbull, K. J., Roghanian, M., Baerentsen, R., Semanjski, M., Brodersen,
D. E., Macek, B., & Gerdes, K. (2019, Jun 18). Serine-Threonine Kinases Encoded by
Split hipA Homologs Inhibit Tryptophanyl-tRNA Synthetase. *MBio, 10*(3).
https://doi.org/10.1128/mBio.01138-19

Wen, Y., Behiels, E., Felix, J., Elegheert, J., Vergauwen, B., Devreese, B., & Savvides, S. N.
(2014, Sep). The bacterial antitoxin HipB establishes a ternary complex with operator
DNA and phosphorylated toxin HipA to regulate bacterial persistence. *Nucleic Acids*
Research, 42(15), 10134-10147. https://doi.org/10.1093/nar/gku665
391

Figure S1

Figure S2

a

b

c

S. oneidensis HipBA

Figure S3

a

α1
β4
β3
β2
β1
β5
HipS
α2
α4
α3

b

E. coli O127:H6 HipB
S. oneidensis HipB

c

HipS
HipA$_{So}$
HipT
HipA$_{So}$

HipBA$_{So}$

Arg380
Arg384
Arg383
Arg429

HipBST

Arg198
Arg302
Lys295
Lys302
Arg291

d

ara   IPTG

OD$_{600}$

time (min)

● pBAD33 / pNDM220
■ *hipT* / pNDM220
▲ *hipT* / *hipS*
▼ *hipT* / *hipS*^{W65A}

Figure S4

Figure S5

a

hipB lacZ

	glucose (represses *hipBST*)	arabinose (induces *hipBST*)

pBAD33
hipBST D233Q
hipBS
hipBT D233Q
hipST D233Q
hipB
hipS
hipT D233Q

b

hipB hipS hipT lacZ

	none (leaky *hipT* expression)	arabinose (induces *hipT*)

pGH254 / pBAD33
hipBST / pBAD33
pGH254 / *hipT* D210Q
hipBST / *hipT* D210Q
pGH254 / *hipT* D233Q
hipBST / *hipT* D233Q

c

C T A G - + *hipBST* D210Q

A
T
A
A
C
C
A
+1 → T
T
A
A
T
G
A
A

d

time post induction
min -2 30 60 C T A G *hipT* D210Q

e

-35   -10

CCTTTTTCT**TTGCCA**ATGAGTTATATGAGC**TAAATT**ATAACCATTAATGAAAAATGGTTCATTTGTGAT

ACTTAACGAATTTATCTGACCATAAATGATCTTGTTTTTGTCTTTTATCAATCG**AAAGGT**TCTTTA**ATG**

SD   **Met**

f

hipT D210Q + *hipB*

	glucose (represses *hipT*)	arabinose (induces *hipT*)	arabinose + IPTG (induces *hipB*)

pBAD33 / *hipB*
hipT (S57IS59, *wt*) / *hipB*
hipT (D57IS59) / *hipB*
hipT (S57ID59) / *hipB*
hipT (A57IS59) / *hipB*
hipT (S57IA59) / *hipB*
hipT (A57IA59) / *hipB*
hipT (D57IA59) / *hipB*
hipT (A57ID59) / *hipB*
hipT (D57ID59) / *hipB*

Figure S6

Figure S7

1: Gly-rich motif   3: Glu99/Glu51   5: Asn215/Asn131
2: Lys86/Lys33   4: Asp210/Asp126 6: Asp233/Asp144