A novel nematode species from the Siberian permafrost shares adaptive mechanisms for cryptobiotic survival with *C. elegans* dauer larva

Anastasia Shatilovich[1#], Vamshidhar R. Gade[2,9#], Martin Pippel[3], Tarja T. Hoffmeyer[4], Alexei V. Tchesunov[5], Lewis Stevens[6], Sylke Winkler[2,7], Graham M. Hughes[8], Sofia Traikov[2], Michael Hiller[3,10], Elizaveta Rivkina[1], Philipp H. Schiffer[4*], Eugene W Myers[3], Teymuras V. Kurzchalia[2*]

1. Institute of Physicochemical and Biological Problems in Soil Science RAS, Pushchino, Russia
2. Max Planck Institute for Molecular Cell Biology and Genetics, Dresden, Germany
3. Center for Systems Biology, Dresden, Germany
4. Institute for Zoology, University of Cologne
5. Department of Invertebrate Zoology, Faculty of Biology, M.V. Lomonosov Moscow State University, Moscow, Russia
6. Tree of Life, Wellcome Sanger Institute, Cambridge, UK
7. DRESDEN concept Genome Center, Dresden, Germany
8. School of Biology and Environmental Science, University College Dublin, Belfield, Dublin, Ireland
9. Institute of Biochemistry, ETH Zürich, 8093 Zürich, Switzerland
10. LOEWE Centre for Translational Biodiversity Genomics, Senckenberg Society for Nature Research & Goethe University, Frankfurt am Main, Germany

[#] A.S and V.G contributed equally to this work

*To whom correspondence may be addressed

Email: kurzchalia@mpi-cbg.de

p.schiffer@uni-koeln.de

Abstract

Some organisms in nature have developed the ability to enter a state of suspended metabolism called cryptobiosis[1] when environmental conditions are unfavorable. This state-transition requires execution of a combination of genetic and biochemical pathways[1,2,3] that enable the organism to survive for prolonged periods. Recently, nematode individuals have been reanimated from Siberian permafrost after remaining in cryptobiosis. Preliminary analysis indicates that these nematodes belong to the genera *Panagrolaimus* and *Plectus*[4]. Here, we present precise radiocarbon dating indicating that the *Panagrolaimus* individuals have remained in cryptobiosis since the late Pleistocene (~46,000 years). Phylogenetic inference based on our genome assembly and a detailed morphological analysis demonstrate that they belong to an undescribed species, which we named *Panagrolaimus n. sp*. Comparative genome analysis revealed that the molecular toolkit for cryptobiosis in *Panagrolaimus n. sp.* and in *C. elegans* is partly orthologous. We show that biochemical mechanisms employed by these two species to survive desiccation and freezing under laboratory conditions are similar. Our experimental evidence also reveals that *C. elegans* dauer larvae can remain viable for longer periods in suspended animation than previously reported. Altogether, our findings demonstrate that nematodes evolved mechanisms potentially allowing them to suspend life over geological time scales.

Introduction

Organisms from diverse taxonomic groups can survive extreme environmental conditions, such as the complete absence of water or oxygen, high temperature, freezing, or extreme salinity. The survival strategies of such organisms include a state known as suspended animation or cryptobiosis, in which they reduce metabolism to an undetectable level[6]. Spectacular examples of long-term cryptobiosis include a *Bacillus* spore that was preserved in the abdomen of bees buried in amber for 25 to 40 million years[7], and a 1000 to 1500 years-old *Lotus* seed, found in an ancient lake, that was subsequently able to germinate[8]. Metazoans such as tardigrades, rotifers and nematodes are also known for remaining in cryptobiosis for prolonged periods[9,10]. The longest records of cryptobiosis in nematodes are reported for the Antarctic species *Plectus murrayi*[11] (25.5 years in moss frozen at -20°C), and *Tylenchus polyhypnus*[12] (39 years desiccated in an herbarium specimen).

Intensive research during the last decade has demonstrated that permafrosts (perennially frozen sediments) are unique ecosystems preserving life forms at sub-zero temperatures over thousands of years[13,14,15,16]. Permafrost remains are an exceptional source for discovering a wide variety of unicellular and multicellular living organisms surviving in cryptobiosis for prolonged periods[6,17,18]. The Siberian permafrost is a unique repository for preserving organisms in sub-zero temperatures for millions of years. Expeditions in the past decade have resulted in the revival of several organisms across various taxa from the Siberian permafrost[5,36,37,38]. The possibility to exploit permafrost as a source for reanimating multicellular animals was recognized as early as 1936. A viable Cladocera crustacean *Chydorus sphaericus*, preserved in the Transbaikalian permafrost for several thousand years[39,40], was discovered by P.N. Kapterev, who worked at the scientific station Skovorodino as a GULAG prisoner. Unfortunately, this observation remained unnoticed for many decades. We recently reanimated soil nematodes that were preserved in Siberian permafrost for potentially thousands of years, and initial morphological observations provisionally described them as belonging to the genera *Panagrolaimus* and *Plectus*. Previous studies demonstrated several species of *Panagrolaimus* can undergo cryptobiosis in the form of anhydrobiosis (through desiccation) and cryobiosis (through freezing)[19,20,21,22,23]. In various nematodes, entry into anhydrobiosis is often accompanied by a preparatory phase of exposure to mild desiccation, known as preconditioning[21,24]. This induces a specific re-modelling of the transcriptome, the proteome, and metabolic pathways that enhances survival ability[2,3,25]. Some panagrolaimids possess adaptive mechanisms for rapid desiccation where most of the cellular

water is lost, while others possess freezing tolerance without loss of water at sub-zero
temperatures by inhibiting the growth and recrystallisation of ice crystals[21].

Here we present a high-quality genome assembly, detailed morphological phylogenetic
analysis, and define a novel species, *Panagrolaimus n. sp.* Precise radiocarbon dating indicates
that *Panagrolaimus n. sp* remained in cryptobiosis for about 46,000 years, since the late
Pleistocene. Furthermore, making use of the model organism *C. elegans*, we demonstrate that
C. elegans dauer larvae and *Panagrolaimus* utilize comparable molecular mechanisms to
survive extreme desiccation and freezing, i.e. upregulation of trehalose biosynthesis and
gluconeogenesis.
106

107   Results

108   Discovery site and radiocarbon dating

Previously, we had shown that nematodes from the Siberian permafrost with morphologies
consistent with the genera *Panagrolaimus* and *Plectus* could be reanimated thousands of years
after they had been frozen. Several viable nematode individuals were found in two of the more
than 300 studied samples of permafrost deposits spanning different ages and genesis. Samples
were collected by researchers of the Soil Cryology Lab, Pushchino, Russia, during perennial
paleo-ecological expeditions carried out in the coastal sector of the northeastern Arctic[17]. The
detailed description of the study site (outcrop Duvanny Yar, Kolyma River, Fig.1A), sampling
and revitalizing procedures are provided in Supplementary Information (SI). Like other late
Pleistocene permafrost formations in the northeastern Arctic, Duvanny Yar is composed of
permanently frozen ice-rich silt deposits riddled with large polygonal ice wedges that divide
them into mineral blocks[26,27] (Fig.1B). Sediments include sandy alluvial layers, peat lenses,
buried paleosols and Pleistocene rodent burrows (Fig.1C). The burrow (P-1320), in which
Panagrolaimus nematodes were found (Fig.1D), has been taken from the frozen outcrop wall
at a depth of about 40 m below the surface and about 11 m above river water level in
undisturbed and never thawed late Pleistocene permafrost deposits. The fossil burrow left by
arctic gophers of the genus *Citellus* consists of an entrance tunnel and large nesting chamber
up to 25 cm in diameter [26].

The sterility of permafrost sampling and age of cultivated biota have been discussed in
detail in several reviews[14,28, 29]. Based on previous reports, the age of the organisms found in a
burrow is equal to the freezing time and corresponds to the age of organic matter conserved in
the syncryogenic sediments. This makes it possible to use radiocarbon dating of organic matter

to establish the age of organisms. We performed Accelerator Mass Spectrometry (AMS)

radiocarbon analysis of plant material obtained from studied borrow P-1320 and determined a

direct ^{14}C age of 44,315±405 BP (Institute of Geography, RAS; sample IGAN$_{AMS}$ 9137).

Calibrated age range is 45,839 – 47,769 cal BP (95.4% probability) (Fig.S1).

Like other parthenogenetic *Panagrolaimus*, the newly discovered species is triploid

The revived animal was cultivated in the laboratory for over 100 generations and initially

described as *Panagrolaimus* aff. *detritophagus*[4] based on morphology. We conducted a

detailed morphological analysis of the revived animal (Fig.2, S2, Table S1; BOX1), which

confirmed unambiguously that the animal belongs to the genus of *Panagrolaimus*, in

agreement with a previous phylogenetic analysis of the 18S ribosomal RNA sequence[4].

However, due to the morphological uniformity of *Panagrolaimus*, unusual even for nematodes,

morphology and molecular analysis of a single ribosomal RNA sequence is insufficient to

describe a species. We found the species to be parthenogenetic, which further complicates

description under most species concept. Due to these limitations, we decided to refer to the

phylogenetic species concept, using phylogenetic trees based on multiple genes as markers.

To obtain comprehensive molecular data for phylogenomic species determination, we

generated a genome assembly usinging PacBio HiFi sequencing with long reads (84X

coverage, mean length 14,425 bp). Our analysis of repeat and gene content is described in

Supplementary Table 3. K-mer analysis of the reads clearly indicated that this animal has a

triploid genome (Fig.3A), like other parthenogenetic *Panagrolaimus* species[30]. Despite the

challenges that a triploid genome poses for assembly, we obtained a highly contiguous contig

assembly of the three pseudohaplotypes that comprise almost 266 Mb and thus have a similar

genome size as other parthenogenetic *Panagrolaimus* species[30]. The contig N50 value of all

three pseudohaplotypes is 3.8 Mb. Since these pseudohaplotypes exhibited a noticeable degree

of divergence, we further investigated their relationship by using the apparent homeologs in

our gene predictions to align the longest continuous contigs based on micro synteny (Fig. 3B).

Links between the contigs clearly show the triploid state of the genome.

    BOX 1 Description of *Panagrolaimus n. sp* sp. nov.

Description	Body spindle-shaped and usually almost straight after fixation (Fig.2a, b). Cuticle thin and faintly annulated, 10–11 annules per 10 μm in cervical region, 13–14 in midbody, preanally again 10 annules within 10 μm. Conspicuous convex lateral fields with three incisures 1.5-2 μm wide extended along the body from about ¼–1/3 procorpus length to 2/3 of tail length. In SEM, the lateral field looks like a bolster with a narrow median split. Labial region set off. Mouth opening surrounded with six lips (Fig.2c, d). Anterior sensilla as papillae arranged in two close but separate subsequent circles. Somatic sensilla (i.e., deirids and phasmids) not evident. Buccal cavity cylindro-conoid, and unarmed; its total length 9–13 μm, maximum stoma width 1.7–2.8 μm (Fig.2e). Dorsal stoma wall (dorsal rhabdion) more clearly sclerotized. Anterior part of the buccal cavity comprising cheilostom and gymnostom nearly cylindroids while stegostom conically narrowed and ended with a distinct tight flexion. Pharynx consists of three distinct parts: straight anterior procorpus, narrow medial isthmus and rounded terminal bulb. Procorpus gradually widening to it posterior end, always straight in all specimens, with transversal muscular striation, more prominent in posterior three fourths. Isthmus narrow, cylindroid, bent starkly in nearly all studied specimens. Terminal bulb strongly muscular, with a valvular apparatus at about 40% of the bulb length. Cardia in shape of truncate cone. Intestine (midgut) tissue filled with vacuoles and granules; in the anteriormost region, the granules smaller and look more pallid. Cell borders not visible in the intestine, but internal lumen distinct, sinusoid. No internal content visible in the internal lumen. Ventral excretory-secretory pore and its cuticularized duct situated at the level of anterior part of the bulb. No other details of the excretory-secretory system visible. Vulval lips protruding. Genital branch monodelphic prodelphic and situated dorsally and to the right of the midgut (Fig.2f). Vagina distinctly cuticularized and opens to the elongate uterus. A long oviduct extended anteriad from the anterior uterus; the oviduct folded up and then posteriorward and transforms into an elongate ovary. There are

	one or two ripe eggs in the uterus in most specimens. Tail short conical, with short acute spike-like mucro (Fig.2g).
Etymology	Species name *kolymaensis* (Latin) is derived from the Kolyma River area.
Holotype	Senckenberg Natural History Museum, Frankfurt am Main, Germany (collection number SMF 17067).
Paratypes	Senckenberg Natural History Museum, Frankfurt am Main, Germany (collection numbers SMF 17068, SMF 17069) (eighteen paratypes).
Type locality	Frozen fossil rodent burrow buried in permafrost 45,839 – 47,769 cal BP, 40 meters from the surface, outcrop Duvanny yar, Kolyma River, North-East of Siberia, Russia (68.633410, 159.078800). Frozen material from burrow was collected by Dr.S.Gubin (Soil Cryology Lab, Pushchino, Russia) in august 2002.

Applying a phylogenomic concept to define *Panagrolaimus n. sp* sp. nov.

To place the species in the genus *Panagrolaimus*, we conducted a broader multi-gene

phylogenomic analysis using Maximum likelihood methods. Our analysis of a concatenated,

partitioned alignment of 60 genes, and a coalescence-based approach using a broader set of

12,295 gene trees, retrieves the revived animal as sister to all other sequenced *Panagrolaimus*

species, but as an ingroup to *Propanagrolaimus*[31] (Fig.3C; Fig. S3B-C). Thus, the phylogenetic

placement provides strong evidence that this animal represents a novel species. Furthermore,

there is substantial sequence divergence between this novel species, and *Panagrolaimus* sp.

PS1159 and *Panagrolaimus* sp. ES5, estimated to be on average 2.06 and 2.11 amino acid

substitutions per site in our concatenated alignment, respectively. The substantial divergence

is in line with previous data on ages of *Panagrolaimus* nematodes[30], and more broadly seen in

nematodes, which can be hyper-diverse[32,33]. Our data also contradicts the assumption that

parthenogenesis is a monophyletic trait[30] in the *Panagrolaimus* genus (Fig.3C). Using the gene-

tree reconciliation approach implemented in GRAMPA (Gene-tree Reconciliation Algorithm

with MUL (<u>Mu</u>lti <u>l</u>abelled)-trees for Polyploid Analysis), we explored whether the additional

set of proteins we found in the parthenogenetic species is a result of auto- or allopolyploidy.

Finding these the additional proteins basally branching, outside of the lineage containing both

parthenogenetic and sexual species, suggests that an allopolyploid origin of these extra gene

copies (Fig.S6). Based on the Kolyma River location where the animal was unearthed, we

propose the following taxonomic classification and species name:

Phylum Nematoda Potts, 1932

Class Chromadorea Inglis 1983

Suborder Tylenchina Thorne, 1949

Family Panagrolaimidae Thorne, 1937

Panagrolaimus sp. nov.

C. elegans dauer larvae and Panagrolaimus n. sp. might utilize partially similar

mechanisms to enter and remain in cryptobiotic state for prolonged periods of time

In the absence of established genetic methods in *Panagrolaimus n. sp.*, we referred to

the model *C. elegans* as a comparator system to gain insights into possible pathways for long

term survival[3,4,24,25]. The high-quality genome of *Panagrolaimus n. sp.* allowed us to compare

its molecular toolkit for cryptobiosis with that of *C. elegans*. We used orthology clustering and

phylogenetics to investigate whether the genome of *Panagrolaimus n. sp.* contains genes

previously implicated in cryptobiosis in the *C. elegans* dauer larva. Our analysis showed that,

like other *Panagrolaimus* species[34,35], *Panagrolaimus n. sp.* also encodes orthologs to a *C.*

elegans trehalose phosphate synthase gene (*tps-2*) and to a trehalose phosphatase gene (*gob-1*)

(Fig. 3D, supplementary file Orthology analysis). Furthermore, we found orthologs to all *C.*

elegans enzymes required for polyamine biosynthesis, the TCA cycle, glycolysis,

gluconeogenesis, and glyoxylate shunt (Fig.3D, supplementary file Orthology analysis)

suggesting that *Panagrolaimus n. sp.* might partially utilize similar molecular mechanisms as

C. elegans to facilitate survival of unfavorable conditions.

Our earlier findings established that amongst several developmental stages of *C.*

elegans, only the dauer larva, formed during unfavorable conditions (such as low nutrients and

high population density), could survive anhydrobiosis and exposure to freezing[4,24]. The dauer

larvae is in a hypometabolic state with distinct metabolic properties such as reduced oxygen

consumption and heat dissipation in comparison to other larval stages of *C. elegans*. To survive

extreme desiccation, *C. elegans* dauer larvae (in its hypometabolic state) need to be first

preconditioned at high relative humidity (98% RH) for 4 days[24]. During preconditioning, dauer

larvae upregulate trehalose biosynthesis that ensures their survival to harsh desiccation[24,25]. We

tested whether survival of *Panagrolaimus n. sp.* is also facilitated by preconditioning. As there

is no dauer stage in the *Panagrolaimus* life cycle, we performed our experiments with a
population of all the larval stages and adults. Although a small proportion of *Panagrolaimus*
n. sp. individuals survive harsh desiccation and freezing without preconditioning (Fig.4A), the
mixture of all the larval stages and adults of *Panagrolaimus n. sp.* survive significantly higher
(p value< 0.0001) in proportion to harsh desiccation upon preconditioning (Fig.4A). Similarly,
preconditioning and desiccation further enhanced survival rate of *Panagrolaimus n. sp.* to
freezing (-80°C) (Fig.4A). Like *C. elegans* dauer larva*, Panagrolaimus n. sp.* upregulates
trehalose levels up to 20-fold upon preconditioning (Fig.4B). We previously reported that, to
upregulate trehalose levels upon preconditioning, *C. elegans* dauer larva dissipate their fat
reserves (Triacylglycerols) by activating the glyoxylate shunt and gluconeogenic pathway[25].
Upon preconditioning, we found that triacylglyceride (TAG) levels are significantly decreased
in *Panagrolaimus n. sp.* (Fig.S5A&B). To further investigate whether acetyl-CoA derived from
the degradation of TAGs culminates in trehalose, we applied the previously developed method
of metabolic labelling with ^{14}C-acetate in combination with 2D-TLC [4,25]. The ^{14}C-acetate
metabolized by the worms is incorporated into TAGs. Upon degradation of TAGs, ^{14}C-acetyl
CoA is released which acts as a precursor for trehalose biosynthesis. As shown in Fig. 4C,
preconditioning led to a huge increase of radioactivity in trehalose and to a small increase in
some amino acids (glycine/serine, phenylalanine; panels c and d). Interestingly,
Panagrolaimus n. sp. displayed an additional spot (Fig. 4D, enumerated as 7), that was not
found in *C. elegans*. We identified this spot as trehalose-6-phosphate (Fig.S5C-H), a precursor
of trehalose, based on the fragmentation pattern of the molecule, using mass spectrometry.
Thus, to resist harsh desiccation, like *C. elegans* dauer larvae, *Panagrolaimus n. sp.* might
utilize the glyoxylate shunt and consequently acetate derived from TAGs to synthesize
trehalose. Detection of the immediate precursor (trehalose-6-phosphate) suggests that the flux
of metabolites is intense in *Panagrolaimus n. sp*. Finally, we investigated whether *C. elegans*
dauer larvae can also survive in prolonged cryptobiotic state. Despite preconditioning, the
survival ability of desiccated dauer larvae at room temperature declines very rapidly, with most
larvae dead after almost 10 days (Fig. 4E) (Fig. S4A&B). Direct freezing without any
cryoprotectants at -80℃ leads to instant death of the animals. To test whether combining these
conditions could extend the viability of dauer larvae (Fig. S4A&B), we transferred the
desiccated larvae to -80℃. Remarkably, under these conditions, there was no significant
decline in viability even after 480 days (Fig. 4E). Moreover, after thawing the animals resumed
reproductive growth and produced progeny in numbers like those of animals kept under control

conditions (Fig. 4F). Since we did not observe any reduction in the survival at any time points,

this suggests that the combination of anhydrobiosis and freezing can prolong the survival

ability of dauer larvae. Thus, *C. elegans* dauer larvae, when exposed to combination of

cryptobiotic states can survive for extremely long periods of time.

Discussion

The new nematode species from permafrost can now be placed into the genus *Panagrolaimus*[41],

which contains several described parthenogenetic and gonochoristic species[30,42]. Many

Panagrolaimus display adaptation to survival in harsh environments[21] and the genus includes

the Antarctic species *P. davidi*[22]. The genus *Panagrolaimus* is exceptional in its morphological

uniformity even among nematode species that are hard to classify based on morphology in

general. Thus, species designation via microscopic (including SEM) analysis is unreliable,

which is further complicated by the absence of males in parthenogenetic species. Males have

an important diagnostic feature such as spicules and pericloacal papillae, females differ from

one species to another mainly by morphometrics, where interspecies differences (absolute

measures and ratios) might be subtle. Our specimens are similar based on absolute sizes and

ratios to females of the bisexual species *Panagrolaimus detritophagus*[43]. The only non-

overlapping morphometric character is index "b" (body length: pharynx length): 5.6–6.8 in

Panagrolaimus n. sp. versus 4.4-5.1 in *P. detritophagus*.

Consequently, we turned to phylogenomic methods under the phylogenetic species

concept to place the species on the tree. This showed that this species is an outgroup to other

known *Panagrolaimus* species, raising the possibility of a second independent evolution of

parthenogenesis in the genus, in contrast to previous findings[42,31,30]. Alternatively, the hybrid

origin of parthenogenetic *Panagrolaimus* could influence the phylogenetic positioning of

strains, raising the possibility that the new species is a true sister to the other parthenogenetic

strains. To fully resolve the phylogenetic positioning further, extensive sampling, and genome

sequencing of *Panagrolaimus* species is needed. We found *Panagrolaimus n. sp.* to be triploid

and thus a hybrid origin is possible, as seen in other parthenogenetic *Pangrolaimus*[30]. The

highly contiguous genome of *Panagrolaimus n. sp.* will allow for analyses of this trait in

comparison to other *Panagrolaimus* species currently being genome sequenced.

Our results provide a deeper insight into the homology of molecular and biochemical

mechanisms between *C. elegans* and *P. kolymaensis*, which are not only taxonomically but

also ecologically distinct. *C. elegans* can mostly be found in rotting fruits and plants in

temperate regions[44,45], while *Panagrolaimus* species are globally distributed and prevalent in

leaf litter and soil[42], including in harsh environments[21]. We show through orthology analysis

that the well-studied molecular pathways used by *C. elegans* larvae to enter the dauer state,

such as insulin[46,47] (DAF-11, DAF-2 & DAF-16), TGF-β[48] (DAF-7), steroid[49] (DAF-9, DAF-

12) are present in the genome of the *Panagrolaimus n. sp.* (Fig.S4C). The presence of

homologous genes in two species does not necessarily demonstrate their functionality in both.

Therefore, further functional analyses are needed to study molecular pathways in detail.

Trehalose accumulation (Fig.4B) and depletion of triacylglycerols (Fig.S5A&B) ensures the

functionality of trehalose biosynthesis pathway and utilization of glyoxylate shunt during

desiccation in *Panagrolaimus n. sp.* Without the activity of the enzyme TPS-2 and glyoxylate

shunt, it is unfeasible to synthesize trehalose in nematodes. We do not eliminate the possibility

of other biochemical features that might contribute to desiccation survival ability of

Panagrolaimus n. sp, but with regards to trehalose biosynthesis and the glyoxylate shunt, our

data suggest that molecular tool kit is partially orthologous. In our future studies, we intend to

perform RNAi based experiments to infer the concrete mechanisms. Our results hint at

convergence or parallelism in the molecular mechanisms organizing dauer formation and

cryptobiosis.

As mentioned above, preconditioning enhances the survival of *Panagrolaimus n. sp.*

by rendering them desiccation tolerant. We previously reported that preconditioning elevates

trehalose biosynthesis in *C. elegans* dauer larvae and the elevated trehalose renders desiccation

tolerance by protecting the cellular membranes[24]. It is not surprising that *Panagrolaimus n. sp.*

upregulates trehalose, however the magnitude of trehalose elevation is higher than *C. elegans*

dauer larvae. This indicates that central regulators (DAF-16, DAF-12) of trehalose

upregulation may differentially regulate *tps-2* in *Panagrolaimus n. sp*[50,51,34]. Although

Panagrolaimus n. sp. utilizes the glyoxylate shunt and gluconeogenesis to upregulate trehalose

levels, it is intriguing to observe that they accumulate substantial levels of trehalose-6-

phosphate. Further investigation of this observation using RNAi or inhibitor-based experiments

will provide insights into molecular mechanisms of metabolic regulation in *Panagrolaimus n.*

sp. upon preconditioning. Our findings for the first time demonstrate that *C. elegans* dauer

larvae possess an inherent ability to survive freezing for prolonged periods if they undergo

anhydrobiosis. It is tempting to speculate that undergoing anhydrobiosis might be a survival

strategy of *C. elegans* to survive the seasonal changes in nature.

In summary our findings indicate that by adapting to survive cryptobiotic state for
short time frames in environments like permafrost, some nematode species gained the potential
for individual worms to remain in the state for geological timeframes. This raises the question
whether there is an upper limit to the length of time an individual can remain in the cryptobiotic
state. Long timespans may be limited only by drastic changes to the environment such as strong
fluctuations in ambient temperature, natural radioactivity, or other abiotic factors. These
findings have implications for our understanding of evolutionary processes, as generation times
may be stretched from days to millennia, and long-term survival of individuals of species can
lead to the refoundation of otherwise extinct lineages. This is particularly interesting in the case
of parthenogenetic species, as each individual can found a new population without the need for
mate finding, i.e. evading the cost of sex. Finally, understanding the precise mechanisms of
long-term cryptobiosis and cues that lead to successful revivals can inform new methods for
long term storage of cells and tissues.

Methods

Materials and *C. elegans* strains

[1-14C] -acetate (sodium salt) was purchased from Hartmann Analytic (Braunschweig, Germany). All other chemicals were purchased from Sigma-Aldrich (Taufkirchen, Germany). The *Caenorhabditis* Genetic Centre (CGC) which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440) provided *daf-2(e1370)* and *E. coli NA22* strains.

Genomic DNA isolation from *Panagrolaimus n. sp* nematodes

After isolation (Supplementary Information), to ensure our strain *Panagrolaimus n. sp.* (Pn2-1) can adapt to different laboratories, we grew them for multiple generations. The strain was in culture for several generations during genomic DNA isolation and was frozen after genomic DNA isolation was performed. *Panagrolaimus n. sp.* nematodes (isofemale strain Pn2-1) were grown on several plates of NGM agar plated with *E. coli NA22* bacteria at 20°C. Worms were collected from the plates, washed with water at least three to five times by centrifugation at 1000 g to remove any residual bacteria and any debris. The worm pellet was dissolved in 5 volumes of worm lysis buffer (0.1M Tris-HCl pH=8.5, 0.1M NaCl, 50mM EDTA pH=8.0) and distributed in 1.5 ml of microcentrifuge tubes. These tubes are incubated at -80°C for 20 minutes. 100 µl of Proteinase 'K' (20 mg/ml) was added to each tube and they are incubated at 60°C overnight. 625 µl of cold GTC buffer (4M Guanidinium Thiocynate, 25mM Sodium citrate, 0.5% (v\v) N-lauroylsarcosine, 7%(v/v) Beta Mercaptoethanol) was added to the tube, incubated on ice 30 min, and mixed by inverting every 10 min. 1 volume of phenol–chloroform-isoamyl alcohol (pH=8) was added to the lysate and mixed by inverting the tube 10-15 times. Tubes were centrifuged for 5 min at 10,000 g at 4°C to separate the phases. The upper aqueous phase was carefully collected into a fresh tube. One volume of fresh chloroform was added and mixed by inverting the tubes for 10-15 times and centrifuged for 5 min at 10,000 g at 4°C to separate the phases. One volume of cold 5 M NaCl was added, mixed by inverting the tubes and incubated on ice for 15 min. After incubation these tubes were centrifuged for 15 min at 12,000–16,000 g at 4°C. The supernatant containing the nucleic acids were slowly transferred into a fresh tube. One volume of isopropanol was added to the tube, inverted few times, and incubated on ice for 30 minutes. After incubation, the tubes were centrifuged at 3000 g for 30–45 min at 25°C and the supernatant was discarded without disturbing the pellet. The pellet was washed twice with 1 ml of 70% ethanol, tubes were centrifuged at 3000 g for 5 min and supernatant was discarded and incubated at 37°C for 10-15 min to dry the pellet. The

pellet was resuspended carefully in TE buffer. The quality of the genomic DNA was analyzed

with pulse field gel electrophoresis.

Genome sequencing and assembly

The long insert library was prepared as recommended by Pacific Biosciences according to the

'Procedure & Checklist-Preparing gDNA Libraries Using the SMRTbell® Express Template

Preparation Kit 2.0' protocol. In summary, RNAse treated HMW gDNA was sheared to 20 kb

fragments on the MegaRuptor^{TM} device (Diagenode) and 10 µg sheared gDNA was used for

library preparation. The PacBio SMRTbell^{TM} library was size selected in two fractions (9-

13kb, > 13kb) using the BluePippin^{TM} device with cassette definition of 0.75% DF MarkerS1

3-10 kb Improved Recovery. The second fraction of the size-selected library was loaded with

95 pM on plate on a Sequel SMRT cell (8M). Sequel polymerase 2.0 was used in combination

with the v2 PacBio sequencing primer and the Sequel sequencing kit 2.0EA, with a runtime of

30 hours. We created PacBio CCS reads from the subreads .bam file using PacBio's ccs

command linetool (version4.2.0), outputting 8.5Gb of high-quality CCS reads (HiFi reads N50

of 14.4 kb). HiCanu (version 2.2)[39] was used to create the contig assembly. Blobtools[53] (version

1.1.1) was used to identify and remove bacterial contigs. The final triploid contig assembly

consists of 856 contigs has a N50 of 3.82 Mb and a size of 266Mb. The mitochondrial genome

was created with the mitoHifi pipeline (version 2, https://github.com/marcelauliano/MitoHiFi)

based on the assembled contigs and the closely related reference mitochondrial genome of

Panagrellus redivivus (strain: PS2298/MT8872, ENAaccession: AP017464). The mitoHifi

pipeline identified 49 mitochondrial contigs ranging from 13-32Kb. The final annotated

circular mitochondrial genome has a length of 17467 bp.

To identify pseudohaplotypes in the *Panagrolaimus n. sp.* genome assembly, we

selected the longest isoform of each predicted protein-coding gene in our assembly and in the

C. elegans genome (downloaded from WormBase Parasite, release WBPS15) using AGAT

(version 0.4.0) and clustered them into orthologous groups (OGs) using OrthoFinder (version

2.5.2). We identified OGs that contained three *Panagrolaimus* sequences (i.e. groups that were

present as single-copy in all three pseudohaplotypes) and used these to identify trios of multi-

megabase size contigs derived from the three pseudohaplotypes. Synteny between the three

pseudohaplotypes was visualized using Circos to plot the positions of each homeolog (version

0.69-8).

Genome annotation

RepeatModeler 1.0.8 (http://www.repeatmasker.org/) was used with parameter '-engine ncbi' to create a library of repeat families which was used with RepeatMasker 4.0.9 to soft-mask the *Panagrolaimus* genome. To annotate genes, we cross mapped protein models from an existing *Panagrolaimus* as external evidence in the Augustus based pipeline. The completeness of our predictions was evaluated using BUSCO on the gVolante web interface.

Orthology analysis

We conducted a gene orthology analysis using genomic data from *Panagrolaimus n. sp*, the Plectid nematode species from the permafrost, as well as genomic data from WormBase Parasite (https://parasite.wormbase.org; accessed 17/12/2020): *Caenorhabditis elegans*, *Diploscapter coronatus*, *Diploscapter pachys*, *Halicephalobus mephisto*, *Panagrellus redivivus*, *Panagrolaimus davidi*, *Panagrolaimus* sp. ES5, *Panagrolaimus* sp. PS1159, *Panagrolaimus superbus*, *Plectus sambesii*, and *Propanagrolaimus* sp. JU765. For plectids, genomic resources are scarce. We therefore added transcriptome data of *Plectus murrayi*, *Anaplectus granulosus*, *Neocamacolaimus parasiticus*, and *Stephanolaimus elegans*, with the latter three transcriptomes kindly provided by Dr. Oleksandr Holovachov (Swedish museum of natural history). Transcriptomes for *Anaplectus granulosus*, and *Neocamacolaimus parasiticus* have been published and are readily available [55,56]. All three transcriptomes were assembled *de novo* with Trinity[57]. The exact procedures are described in the respective publications[55,56]. The *Stephanolaimus elegans* transcriptome was assembled using the same methodologies as *Neocamacolaimus parasiticus*.

The *Plectus murrayi* transcriptome was built from raw reads deposited at NCBI (https://sra-downloadb.be-md.ncbi.nlm.nih.gov/sos2/sra-pub-run-13/SRR6827978/SRR6827978.1; accessed 22.12.2020) and assembled using Galaxy Trinity version 2.9.1[58], [57]. All default options were used including *in silico* normalization of reads before assembly. Transdecoder (conda version 5.5.0)[59] was used to translate to amino acid sequence. Identical reads were removed with cd-hit version 4.8.1[60,61], with shorter isoforms removed using the Trinity get_longest_isoform_seq_per_trinity_gene.pl command[59] (Trinity conda version 2.8.5; Anaconda Software Distribution, Conda, Version 4.9.2, Anaconda, Nov. 2020). Amino acid translations of the longest isoforms were extracted with AGAT (Dainat, https://www.doi.org/10.5281/zenodo.3552717) from genome assembly FASTA files and genome annotation GFF3 files using the 'agat_convert_sp_gxf2gxf.pl',

'agat_sp_keep_longest_isoform.pl' and 'agat_sp_extract_sequences.pl' scripts, respectively.

All FASTA headers were modified to allow for simple species assignment of each sequence in

subsequent analysis. Orthology analysis was conducted with OrthoFinder v. 2.5.1[62,63] using

default settings. For genes of interest, we constructed alignments with MAFFT v. 7.475[64] using

the localpair and maxiterate (1000) functions. Spurious sequences and areas that were not well

aligned were removed with Trimal v. 1.4.rev22[65] (procedure stated in supplementary file

Orthology analysis below each phylogeny). We then ran phylogenetic analysis with Iqtree2 v.

2.0.6[66], with -bb 1000 option, testing the model for each analysis (models eventually used

stated in supplementary file Orthology analysis). PFAM domains were explored using

Interproscan v. 5.50-84.0[67]. The phylogenies were visualized with Dendroscope 3.7.6[68] and

figures were created with Inkscape (https://inkscape.org). The majority of our analysis was

performed on the HPC RRZK CHEOPS of the Regional Computing Centre (RRZK) of the

University of Cologne.

Phylogenomics

Sequences of 18S and 28S genes from 44 taxa across the *Propanagrolaimus*, *Panagrolaimus*,

Panagrellus and *Halicephalobus* genera (all listed in Supplementary information) were aligned

(MAFFT L-INS-I v7.475)[64], concatenated [69]and used to infer a species tree using maximum

likelihood via (IQTREE)[70] and partitioned by best-fit models of sequence evolution for both[71].

Nodal support was determined using 1000 bootstrap pseudoreplicates. A further 60 genes from

101 taxa (all listed in Supplementary information) were used to confirm the taxonomic position

using the supermatrix concatenation methods outlined above. Given the limitations of

differential gene sampling, we expanded our phylogenomic analyses to include a coalescence

approach using 12,295 ML gene trees inferred for orthogroups containing the target animal.

Instances of multiple genes per species per group were treated as paralogs/orthologs and

analysed using ASTRAL-Pro[72]. Given the number of copies of genes per orthogroup, we

explored whether auto or alloploidy was the source of extra genes observed using the gene-tree

reconciliation approach implemented in GRAMPA (Gene-tree Reconciliation Algorithm with

MUL (Multi labelled)-trees for Polyploid Analysis)[75]. All gene trees rooted at the midpoint and

the final ASTRAL-pro species tree were used as inputs, with the most parsimonious result

analyzed further.

Desiccation survival assay

C. elegans dauer larvae desiccation assays were performed as described in[24]. *Panagrolaimus n. sp* desiccation assays were performed similarly as described in[24] with mixed population (Mixture of all larval stages and adults) of the nematodes.

Exposure of nematodes to extreme environments

C. elegans dauer larvae or mixed population (Mixture of all larval stages and adults) of *Panagrolaimus n. sp* nematodes were preconditioned and desiccated as described in[24], then transferred to elevated temperature of 34°C, freezing (-80°C) and anoxia. Anoxic environment was generated in a desiccation chamber at 60%RH by flushing the Nitrogen gas into the chamber. The concentration of oxygen inside the chamber was monitored. After each timepoint they were rehydrated with 500 µl of water for 2-3 hours. Rehydrated worms were transferred to NGM agar plates with *E. Coli NA22* as food. Survivors were counted after overnight incubation at 15°C. Each experiment was performed on two different days with at least two technical replicates.

Trehalose quantification from nematode lysates

Trehalose measurements were performed as described in previous reports[25].

Radiolabeling, metabolite extraction and 2D-TLC

The above-mentioned procedures were performed according to previous reports[3,25].

Identification of trehalose-6-phosphate from TLC plates

Normalized aqueous fractions from the non-preconditioned and preconditioned samples were separated by high performance thin layer chromatography (HPTLC), using 1-propanol-methanol-ammonia (32%)-water (28:8:7:7 v/v/v/v) as first, dried for 15 min and 1-butanol-acetone-glacial acetic acid–water (35:35:7:23 v/v/v/v) second dimension respectively. Using the trehalose as a standard on both dimensions of the TLC, the regions of interest were scrapped out from the TLCs. The scraped-out silica was extracted with 10 ml of 50% methanol twice. The fractions were combined, dried under vacuum and dissolved in 100 µl of MS mix solution containing 4:2:1 (Isopropanol:Methanol:Chloroform) with 7.5 mM ammonium formate. Mass spectrometric analysis was performed on a Q Exactive instrument (Thermo Fischer Scientific, Bremen, DE) equipped with a robotic nanoflow ion source TriVersa NanoMate (Advion BioSciences, Ithaca, USA) using nanoelectrospray chips with a diameter of 4.1 µm. The ion

source was controlled by the Chipsoft 8.3.1 software (Advion BioSciences). Ionization voltage

was + 0.96 kV in negative mode; backpressure was set at 1.25 psi. The temperature of the ion

transfer capillary was 200°C; S-lens RF level was set to 50%. FT MS spectra were acquired

within the range of m/z 50–750 at the mass resolution of R m/z 200 = 140000; automated gain

control (AGC) of 3×10^6 and with the maximal injection time of 3000 ms. FT MS/MS spectra

were acquired within the range of m/z 50–750 at the mass resolution of R m/z 200 = 140000;

automated gain control (AGC) of 3×10^4 and with a maximal injection time of 30 s.

Triacylglycerols measurement from *Panagrolaimus n. sp* lysates

Non-preconditioned and preconditioned pellets were lysed in 200 µl of isopropanol with 0.5

mm Zircornium beads twice for 15 min. The lysates were centrifuged at 1300 g for 5 min. The

supernatant was carefully collected without any debris, 20 µl of the lysate was used for protein

estimation. Normalization was performed according to soluble protein levels, supernatant

volumes corresponding to 50-100 µg of proteins were dried in the desiccator. 700 µl of IS

((10:3 (Methyl tert-butyl ether: ethanol)) mix (warmed to room temperature) was added to

dried samples and left on the shaker for 1 hour. The samples were centrifuged at 1400 rpm and

4°C. 140 µl of water was added and left on the shaker for 15 min. These samples were

centrifuged at 13400 rpm for 15 min. The upper organic fraction was collected and transferred

to 1.5 ml glass vial and left for drying in the desiccator. The dried samples were reconstituted

in a volume of 300 µl of 4:2:1 (Isopropanol:Methanol:Chloroform). Volume corresponding to

1 µg was used for injection.

LC-MS/MS analysis was performed on a high-performance liquid chromatography

system (Agilent 1200 HPLC) coupled to a Xevo G2-S QTof (Waters). The samples were

resolved on a reverse phase C18 column (Cortecs C18 2.7um from Waters) with

50:50:0.1:1% (Water:Methanol:Formicacid:1MAmmoniumformate) and 25:85:0.1:1%

(Acetonitrile:Isopropanol:Formic acid:1M Ammonium formate) as mobile phase. The

following gradient program was used: Eluent B from 0 % to 100 % within 12 min; 100 % from

12 min to 17min; 0 % from 17 min to 25 min. The flow rate was set at 0.3 ml/min. The samples

were normalised according to the total protein concentration and the worm numbers. TAG

50:00:00 was used as internal standard. Skylinesoftware

(https://skyline.ms/project/home/software/Skyline/begin.view) was used to analyse the raw

data. TAGs were extracted from Lipidmaps (https://www.lipidmaps.org/) database.

Author contributions

AS, VG, PS, TK conceived and designed the study. AS, VG, TH, MP, AT, GH, MH, ER, PS and TK contributed to the original draft. AS performed cultivation of nematodes, provided samples for scanning electron microscopy and radiocarbon dating. VG performed desiccation survival assays, trehalose measurement, 2D-TLC of metabolites, trehalose-6-phosphate detection, combination of cryptobiosis experiments, genomic DNA isolation from *Panagrolaimus n. sp.*, assembled the data from all the authors, prepared the figures, revised, and submitted the manuscript. MP performed genome assembly. TH conducted orthology analyses and single gene phylogenies and contributed to figures. AT provided morphological description, light, and scanning electron microscopy. LS analyzed the genome assembly and created figures. GMH performed the phylogenetic, MUL analyses and proofread the manuscript. ST performed triacylglycerols measurements, trehalose-6-phosphate detection with suggestions from Andrej Shevchenko. MH annotated supplementary table 3. PS performed genome annotation and supervised TH and LS. EWM supervised MP and had overall responsibility for sequencing and assembly including funding it.

Acknowledgements

VG is thankful to Andrej Shevchenko, members of Kurzchalia lab for helpful discussions and the core facilities of MPI-CBG for assistance. We are grateful to Dr. S. Gubin for field study and sampling, our colleagues in Soil Cryology Lab, Pushchino and North-East Scientific Station in Chersky, Republic of Sakha (Yakutia) for their help and cooperation. The authors thank Long Read Team of the DRESDEN-concept Genome Center, DFG NGS Competence Center, part of the Center for Molecular and Cellular Bioengineering (CMCB), Technische Universität Dresden and MPI-CBG. GMH is funded by a UCD Ad Astra Fellowship. This work was supported by the Russian Foundation for Basic Research (19-29-05003-mk) to AS and ER. VG and TK acknowledge the financial support from the Volkswagen Foundation (Life? research grant 92847). PS and TH are supported by a DFG ENP grant to PS (DFG project 434028868). The authors are thankful to Richard Roy and Jens Bast for critical reading the manuscript. We thank Iain Pattern for suggestions on writing the manuscript. The funders have no role in the design of the study.

584 **Conflict of interest:**

585 The authors declare they have no conflict of interest relating to the content of this article.

References

1. Hibshman, J. D., Clegg, J. S. & Goldstein, B. Mechanisms of Desiccation Tolerance: Themes and Variations in Brine Shrimp, Roundworms, and Tardigrades. *Front. Physiol.* **11**, 592016 (2020).

2. Erkut, C. *et al.* Molecular Strategies of the Caenorhabditis elegans Dauer Larva to Survive Extreme Desiccation. *PLoS ONE* **8**, e82473 (2013).

3. Gade, V. R., Traikov, S., Oertel, J., Fahmy, K. & Kurzchalia, T. V. C. elegans possess a general program to enter cryptobiosis that allows dauer larvae to survive different kinds of abiotic stress. *Sci. Rep.* **10**, 13466 (2020).

4. Shatilovich, A. V. *et al.* Viable Nematodes from Late Pleistocene Permafrost of the Kolyma River Lowland. *Dokl. Biol. Sci.* **480**, 100–102 (2018).

5. Shmakova, L. *et al.* A living bdelloid rotifer from 24,000-year-old Arctic permafrost. *Curr. Biol.* **31**, R712–R713 (2021).

6. Keilin, D. The Leeuwenhoek Lecture - The problem of anabiosis or latent life: history and current concept. *Proc. R. Soc. Lond. Ser. B - Biol. Sci.* **150**, 149–191 (1959).

7. Cano, R. & Borucki, M. Revival and identification of bacterial spores in 25- to 40-million-year-old Dominican amber. *Science* **268**, 1060–1064 (1995).

8. Shen-Miller, J. Sacred lotus, the long-living fruits of China Antique. *Seed Sci. Res.* **12**, 131–143 (2002).

9. Guidetti, R. & Jönsson, K. I. Long-term anhydrobiotic survival in semi-terrestrial micrometazoans. *J. Zool.* **257**, 181–187 (2002).

10. Tsujimoto, M., Imura, S. & Kanda, H. Recovery and reproduction of an Antarctic tardigrade retrieved from a moss sample frozen for over 30 years. *Cryobiology* **72**, 78–81 (2016).

11. Kagoshima, H. *et al.* Multi-decadal survival of an Antarctic nematode, Plectus murrayi, in a -20°C stored moss sample. *CryoLetters* **33**, 280–288 (2012).

12. Steiner, G. & Florence E, A. Resuscitation of the nematode Tylenchus polyhypnus, n. sp., after almost 39 years' dormancy. *J. Wash. Acad. Sci.* **36**, 97–99.

13. Gilichinsky, D. A. *et al.* Microbial Populations in Antarctic Permafrost: Biodiversity, State, Age, and Implication for Astrobiology. *Astrobiology* **7**, 275–311 (2007).

14. Rivkina, E. *et al.* Earth's perennially frozen environments as a model of cryogenic planet ecosystems. *Permafr. Periglac. Process.* **29**, 246–256 (2018).

15. Sipes, K. *et al.* Eight Metagenome-Assembled Genomes Provide Evidence for Microbial
Adaptation in 20,000- to 1,000,000-Year-Old Siberian Permafrost. *Appl. Environ.*
Microbiol. **87**, (2021).

16. Liang, R. *et al.* Genomic reconstruction of fossil and living microorganisms in ancient
Siberian permafrost. *Microbiome* **9**, 110 (2021).

17. Gilichinsky, D. A. & Rivkina, E. M. Permafrost Microbiology. 7 (2011).

18. Yashina, S. *et al.* Regeneration of whole fertile plants from 30,000-y-old fruit tissue buried
in Siberian permafrost. *Proc. Natl. Acad. Sci.* **109**, 4008–4013 (2012).

19. Adhikari, B. N., Wall, D. H. & Adams, B. J. Desiccation survival in an Antarctic nematode:
molecular analysis using expressed sequenced tags. *BMC Genomics* **10**, 69 (2009).

20. Tyson, T. *et al.* A molecular analysis of desiccation tolerance mechanisms in the
anhydrobiotic nematode Panagrolaimus superbus using expressed sequenced tags. *BMC*
Res. Notes **5**, 68 (2012).

21. McGill, L. M. *et al.* Anhydrobiosis and Freezing-Tolerance: Adaptations That Facilitate the
Establishment of Panagrolaimus Nematodes in Polar Habitats. *PLOS ONE* **10**, e0116084
(2015).

22. Thorne, M. A. S., Kagoshima, H., Clark, M. S., Marshall, C. J. & Wharton, D. A. Molecular
Analysis of the Cold Tolerant Antarctic Nematode, Panagrolaimus davidi. *PLoS ONE* **9**,
e104526 (2014).

23. Sandhove, J., Spann, N. & Ristau, K. The Anhydrobiotic Potential of the Terrestrial
Nematodes *Plectus parietinus* and *Plectus velox*: Anhydrobiosis OF *Plectus parietinus* and
P. velox. *J. Exp. Zool. Part Ecol. Genet. Physiol.* **325**, 434–440 (2016).

24. Erkut, C. *et al.* Trehalose Renders the Dauer Larva of Caenorhabditis elegans Resistant to
Extreme Desiccation. *Curr. Biol.* **21**, 1331–1336 (2011).

25. Erkut, C., Gade, V. R., Laxman, S. & Kurzchalia, T. V. The glyoxylate shunt is essential
for desiccation tolerance in C. elegans and budding yeast. *elife* **5**, (2016).

26. Zanina, O. G., Gubin, S. V., Kuzmina, S. A., Maximovich, S. V. & Lopatina, D. A. Late-
Pleistocene (MIS 3-2) palaeoenvironments as recorded by sediments, palaeosols, and
ground-squirrel nests at Duvanny Yar, Kolyma lowland, northeast Siberia. *Quat. Sci. Rev.*
30, 2107–2123 (2011).

27. Murton, J. B. *et al.* Palaeoenvironmental Interpretation of Yedoma Silt (Ice Complex)
Deposition as Cold-Climate Loess, Duvanny Yar, Northeast Siberia: Palaeoenvironmental

692 Interpretation of Yedoma Silt, Duvanny Yar. *Permafr. Periglac. Process.* **26**, 208–288

693 (2015).

694 28.Abramov, A., Vishnivetskaya, T. & Rivkina, E. Are permafrost microorganisms as old as

695 permafrost? *FEMS Microbiol. Ecol.* **97**, fiaa260 (2021).

696 29.Gilichinsky, D. A., Wagener, S. & Vishnevetskaya, T. A. Permafrost microbiology.

697 *Permafr. Periglac. Process.* **6**, 281–291 (1995).

698 30.Schiffer, P. H. *et al.* Signatures of the Evolution of Parthenogenesis and Cryptobiosis in the

699 Genomes of Panagrolaimid Nematodes. *iScience* **21**, 587–602 (2019).

700 31.Schiffer, P. H. *et al.* Developmental variations among Panagrolaimid nematodes indicate

701 developmental system drift within a small taxonomic unit. *Dev. Genes Evol.* **224**, 183–188

702 (2014).

703 32.Dey, A., Chan, C. K. W., Thomas, C. G. & Cutter, A. D. Molecular hyperdiversity defines

704 populations of the nematode Caenorhabditis brenneri. *Proc. Natl. Acad. Sci.* **110**, 11056–

705 11060 (2013).

706 33.Cutter, A. D. Divergence Times in Caenorhabditis and Drosophila Inferred from Direct

707 Estimates of the Neutral Mutation Rate. *Mol. Biol. Evol.* **25**, 778–786 (2008).

708 34.Seybold, A. C., Wharton, D. A., Thorne, M. A. S. & Marshall, C. J. Investigating trehalose

709 synthesis genes after cold acclimation in the Antarctic nematode *Panagrolaimus* sp. DAW1.

710 *Biol. Open* bio.023341 (2017) doi:10.1242/bio.023341.

711 35.Evangelista, C. C. S. *et al.* Multiple genes contribute to anhydrobiosis (tolerance to extreme

712 desiccation) in the nematode Panagrolaimus superbus. *Genet. Mol. Biol.* **40**, 790–802

713 (2017).

714 36.Malavin, S., Shmakova, L., Claverie, J.-M. & Rivkina, E. Frozen Zoo: a collection of

715 permafrost samples containing viable protists and their viruses. *Biodivers. Data J.* **8**, e51586

716 (2020).

717 37.Krivushin, K. *et al.* Two Metagenomes from Late Pleistocene Northeast Siberian

718 Permafrost. *Genome Announc.* **3**, e01380-14 (2015).

719 38.Zhou, J. *et al.* Phylogenetic diversity of a bacterial community determined from Siberian

720 tundra soil DNA. *Microbiology* **143**, 3913–3919 (1997).

721 39.Kapterev, P. Anabiosis in the conditions of permanent congelation. *Izv Akad Nauk SSSR Ser*

722 *Biol* **6**, 1073–1088 (1936).

723 40.Kapterev, P. New data on revitalization of organisms from perpetually frozen grounds. *CR*

724 *Acad Sci URSS* **20**, 315–317 (1938).

41. Andrássy, I. *Klasse Nematoda*. (Gustav Fischer Verlag, 1984).

42. Lewis, S. C. *et al.* Molecular evolution in Panagrolaimus nematodes: origins of

parthenogenesis, hermaphroditism and the Antarctic species P. davidi. *BMC Evol. Biol.* **9**,

15 (2009).

43. Andrássy, I. *Free-living Nematodes of Hungary*. (Hungarian Natural History Museum,

2005).

44. Schulenburg, H. & Félix, M.-A. The Natural Biotic Environment of *Caenorhabditis*

elegans. *Genetics* **206**, 55–86 (2017).

45. Frézal, L. & Félix, M.-A. C. elegans outside the Petri dish. *eLife* **4**, e05849 (2015).

46. Gems, D. *et al.* Two pleiotropic classes of daf-2 mutation affect larval arrest, adult behavior,

reproduction, and longevity in Caenorhabditis elegans. *Genetics* **150**, 129–155 (1998).

47. Vowels, J. J. & Thomas, J. H. Multiple chemosensory defects in daf-11 and daf-21 mutants

of Caenorhabditis elegans. *Genetics* **138**, 303–316 (1994).

48. Ren Peifeng *et al.* Control of C. elegans Larval Development by Neuronal Expression of a

TGF-β Homolog. *Science* **274**, 1389–1391 (1996).

49. Gerisch, B., Weitzel, C., Kober-Eisermann, C., Rottiers, V. & Antebi, A. A Hormonal

Signaling Pathway Influencing C. elegans Metabolism, Reproductive Development, and

Life Span. *Dev. Cell* **1**, 841–851 (2001).

50. Penkov, S. *et al.* Integration of carbohydrate metabolism and redox state controls dauer

larva formation in Caenorhabditis elegans. *Nat. Commun.* **6**, 8060 (2015).

51. Penkov, S. *et al.* A metabolic switch regulates the transition between growth and diapause

in C. elegans. *BMC Biol.* **18**, 31 (2020).

52. Nurk, S. *et al.* HiCanu: accurate assembly of segmental duplications, satellites, and allelic

variants from high-fidelity long reads. *Genome Res.* **30**, 1291–1305 (2020).

53. Laetsch, D. R. & Blaxter, M. L. BlobTools: Interrogation of genome assemblies.

F1000Research **6**, 1287 (2017).

54. Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M. & Borodovsky, M. BRAKER2: automatic

eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a

protein database. *NAR Genomics Bioinforma.* **3**, lqaa108 (2021).

55. Smythe, A. B., Holovachov, O. & Kocot, K. M. Improved phylogenomic sampling of free-

living nematodes enhances resolution of higher-level nematode phylogeny. *BMC Evol. Biol.*

19, 121 (2019).

56. Ahmed, M., Adedidran, F. & Holovachov, O. A draft transcriptome of a parasite
Neocamacolaimus parasiticus (Camacolaimidae, Plectida). *J. Nematol.* **53**, 1–4 (2021).

57. Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

58. Goecks, J., Nekrutenko, A., Taylor, J. & Galaxy Team, T. Galaxy: a comprehensive
approach for supporting accessible, reproducible, and transparent computational research in
the life sciences. *Genome Biol.* **11**, R86 (2010).

59. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the
Trinity platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494–1512 (2013).

60. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

61. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

62. Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157
(2015).

63. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238 (2019).

64. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

65. Capella-Gutierrez, S., Silla-Martinez, J. M. & Gabaldon, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973
(2009).

66. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

67. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*
30, 1236–1240 (2014).

68. Huson, D. H. & Scornavacca, C. Dendroscope 3: An Interactive Tool for Rooted
Phylogenetic Trees and Networks. *Syst. Biol.* **61**, 1061–1067 (2012).

69. Kück, P. & Meusemann, K. FASconCAT: Convenient handling of data matrices. *Mol.*
Phylogenet. Evol. **56**, 1115–1118 (2010).

70. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol.*

Biol. Evol. **32**, 268–274 (2015).

71. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**,

587–589 (2017).

72. Zhang, C., Scornavacca, C., Molloy, E. K. & Mirarab, S. ASTRAL-Pro: Quartet-Based

Species-Tree Inference despite Paralogy. *Mol. Biol. Evol.* **37**, 3292–3307 (2020).

73. Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and Smudgeplot

for reference-free profiling of polyploid genomes. *Nat. Commun.* **11**, 1432 (2020).

74. Kokot, M., Długosz, M. & Deorowicz, S. KMC 3: counting and manipulating k-mer

statistics. *Bioinformatics* **33**, 2759–2761 (2017).

75. Thomas, GWC., Ather, H. S., Hahn, W. Matthew. Gene-Tree Reconciliation with MUL-

Trees to Resolve Polyploidy Events. Systematic Biology 66, 1007-1018 (2017).

Figure legends

Figure 1. Study site.

a) location of the Duvanny Yar outcrope on the Kolyma River, northeastern Siberia, Russia. b) view of the upper part of outcrop composed of ice wedges and permafrost silty deposits. c) lithostratigraphic scheme of deposits, showing location of studied rodent borrow (red circle). d) fossil rodent burrow with herbaceous litter and seeds buried in permafrost deposits; m a.r.l. = meters above river level.

Figure 2. General morphology of *Panagrolaimus n. sp*., female.

Scanning electron pictures (a, c), light microscopy photographs (e, f) and graphic presentations (b, d, g) of holotype: a, b) entire body, c, d) anterior ends, e) anterior body, f) perivulvar body region, g) tail. Abbreviations: l.f. – lateral field, ov – ovary, pro – procorpus of the pharynx, t.b. – terminal bulb of the pharynx, u – uterus with eggs, v – vulva, v.p. – ventral pore. Scale bars: a, d, e, f, g – 20 μm, b – 100 μm, c – 2 μm.

Figure 3. Genome assembly and phylogenomics reveals that the newly discovered _Panagrolaimus n. sp_ species is triploid.

a) Kmer spectra of the _Panagrolaimus n. sp_ PacBio HiFi data. Kmers of length 19 were counted using Jellyfish. b) Circos plot showing the triploid structure of the _Panagrolaimus n. sp_ genome. Lines represent the position of 6,715 homeologs in eight contigs that comprise 39.9 Mb (15%) of the assembly. Homeologs were identified by clustering protein-coding genes into orthogroups using OrthoFinder and selecting groups containing three sequences. Contig IDs and scale is shown. c) Inferred species tree for all taxa. The maximum likelihood tree inferred using a concatenated supermatrix (18S and 28S genes) with bootstrap support values is displayed. All genera are represented as monophyletic clades. _Panagrolaimus n. sp_ is highlighted in red and basal to all other _Panagrolaimus_ taxa. Internal nodes, where all subsequent branches represent identical sequences, are displayed with a black star. d) _Panagrolaimus n. sp._ possesses _C. elegans_ gene orthologs to enzymes required for TCA cycle, glyoxylate shunt, glycolysis, gluconeogenesis, trehalose synthesis, and polyamine synthesis. Black filled circles: Ortholog presence suggested by orthogroup clustering, phylogenetic analysis, and domain architecture. White filled circles: No ortholog found via current analysis. Coloured filled circles: Presence of _Panagrolaimus n. sp._ gene(s), related to several _C. elegans_ genes (all genes of same colour) that are all co-orthologous to that gene (those genes). Label: _C. elegans_ enzyme names and orthogroup that contains that gene according to our orthogroup clustering.

Figure 4. *C. elegans* dauer larvae and *Panagrolaimus n. sp.* might utilize similar mechanisms to survive cryptobiosis.

a) Survival rate of *Panagrolaimus n. sp.* nematodes to desiccation and freezing (-80°C). Error bars indicate standard error of mean of two independent experiments with two technical replicates performed on two different days. Statistical comparison was performed using unpaired t test with Welch correction. n.s. $p > 0.05$, ****$p < 0.0001$. For desiccation (non-preconditioned) $n = 289$, freezing (non-preconditioned) $n = 675$, desiccation (preconditioned at 98%RH) $n = 953$ and freezing (preconditioned at 98%RH) $n = 1295$. b) *Panagrolaimus n. sp.* nematodes and *daf-2(e1370)* dauer larvae upregulate trehalose levels upon preconditioning at 98%RH. Error bars indicate standard error of mean of two independent experiments with three replicates performed on two different days. Statistical comparison was performed using two-way ANOVA with Holm-Sidak's multiple comparison test, ****$p < 0.0001$. c-d) 2D-thin layer chromatography of ^{14}C-acetate labelled metabolites from *Panagrolaimus n. sp.* that were non-preconditioned and preconditioned at 98%RH. Enumerated spots indicate trehalose (1), glucose (2), glutamate (3), glutamine (4), serine/glycine (5) and phenylalanine (6). Representative images from at least two independent experiments performed on two different days. e) Desiccated *daf-2 (e1370)* dauer larvae survive to freezing (-80°C) for an extremely long period. Error bars indicate standard error of mean of two independent experiments with two technical replicates performed on two different days. f) Brood size of desiccated dauer larvae exposed to freezing remain like that non- desiccated dauer larvae. Average brood size is the mean of seven dauer larvae per each condition. Statistical comparison was performed by using non-parametric Kolmogorov-smirnov test n.s $p>0.05$.

Figure S1. Calibration of a radiocarbon (^{14}C) date.

Radiocarbon date (44,315±405 BP) and calibrated age (45,839 -- 47,769 cal BP) of plant material collected from buried borrow P-1320. Radiocarbon ages were converted to calendar age equivalents with the OxCal V.4.4 program using the IntCal20 calibration curve. Pink-shaded area — radiocarbon date with standard deviation; grey-shaded area — radiocarbon date projection on the calibration curve with 95.4% probability.

Figure S2. Morphology of *Panagrolaimus n. sp* sp. n., female.

Graphic presentations of holotype (a,b) and SEM pictures (c-i): a) anterior body, b) female reproductive branch, c–e) anterior end of three different female specimens, f) anterior part of the lateral ridge, g) vulva, h) ventral excretory/secretory pore, i) posterior body with anus and lateral ridge. Scale bars: a,b - 50 μm, c,i - 3 μm, d - 2 μm, e –1 μm, f,h - 5 μm, g - 10 μm, .

Figure S3. Phylogenies inferred for genes sets using both concatenation and coalescence approaches.

a) The ploidy level was analysed with *Smudgeplot* v0.2.1[73]. KMC version 3.1.0[74] was used to count the 21-mers in the PacBio CCS reads. Then we ran *smudgeplot.py* to determine the lower and upper coverage cut-offs. These were determined to be 14 and 380. 21-mers with a coverage between 14 and 380 were filtered with kmc_tools. Then, we computed the k-mer pairs from the filtered 21-mers by running *smudgeplot.py hetkmers*. Finally, the produced a smudgeplot shows an estimated ploidy of 3. b) Species tree inferred for 12,295 gene trees using coalescence approach. The species tree implemented using the coalescence approach with orthogroup gene tree is displayed. Novel species in this study are displayed in red. All nodes have a posterior probability of 1. c) Species tree inferred for 102 taxa. The maximum likelihood tree inferred using a concatenated supermatrix of 60 genes is displayed. Bootstrap values are only displayed for nodes with less than 100% support. The platyhelminth *Macrostum lignano* serves as an outgroup for rooting. The ancestral *Panagrolaimus*, sister to all others within the *Panagrolaimus* genus, is highlighted in red.

Figure S4. Combination of cryptobiotic states enhances survival of *C. elegans* dauer larvae.

a) Desiccated dauer larvae manifest enhanced survival rate to heat stress (34°C). Error bars indicate standard deviation of two independent experiments with two technical replicates. b) Desiccated dauer larvae display enhances survival rate to anoxia. Error bars indicate standard deviation of two independent experiments with two technical replicates. Statistical comparison was performed by paired two tailed t-test. *p<0.05. c) *Panagrolaimus n. sp.* possesses gene orthologs to most genes implicated in dauer formation and metabolism in *C. elegans*. Black filled circles: Ortholog presence suggested by orthogroup clustering, phylogenetic analysis, and domain architecture. White filled circles: No ortholog found via current analysis (in all cases these *C. elegans* genes did not cluster with any *Panagrolaimus* genes in the orthogroup clustering). Label: *C. elegans* enzyme names and orthogroup that contains that gene according to our orthogroup clustering.

Figure S5. *Panagrolaimus n. sp.* reduces triacylglycerols (TAGs) levels and accumulates trehalose-6-phosphate upon preconditioning at 98%RH.

a) 1D-Thin layer chromatography of acetate labelled organic fractions of non-preconditioned (1) and preconditioned (2) *P. kolymaensis*. b) Mass spectrometric quantification of TAG levels of non-preconditioned (1) and preconditioned (2) *P. kolymaensis*. Error bars indicate standard deviation of two biological replicates with two technical replicates. Statistical analysis was performed using unpaired t-test with Welch correction **$p < 0.001$. c-d) non-preconditioned and preconditioned mass spectrum of an empty region, e-f) spot 1 (trehalose), g-h) spot 7 (trehalose-6-phosphate) scraped out and extracted from the 2D-TLC.

1117 **Figure S6. Exploring a possible allopolyploid origin for extra proteins.**

1118 Gene-tree reconciliation was used to determine whether extra sets of proteins across

1119 orthogroups originate through auto- or allopolyploidy. Different copies of proteins (designated

1120 by '+' and '*') suggest an allopolyploid origin. Parthenogenetic species are highlighted in bold.

Figure 1
Shatilovich *et al*

Figure 3
Shatilovich *et al*

A

1   2

TAGs

1-Non-Preconditioned

2-Preconditioned at 98%RH

B

1- Non-Preconditioned

2- Preconditioned at 98%RH

C   Non-preconditioned

Empty Spot

D   Preconditioned at 98%RH

E   Spot 1(Trehalose)(M-1)

89.02   179.05

F   341.11

89.02   179.05

101.02

113.02

G   Spot 7 (Trehalose-6-phosphate)(M-1)

241.01

421.08

H   421.08

78.96

96.97

241.01

Figure S5
Shatilovich *et al*

Figure S6
Shatilovich *et al*