A novel nematode species from the Siberian permafrost shares adaptive mechanisms for cryptobiotic survival with *C. elegans* dauer larva

Anastasia Shatilovich[1#], Vamshidhar R. Gade[2,9#], Martin Pippel[3], Tarja T. Hoffmeyer[4], Alexei V. Tchesunov[5], Lewis Stevens[6], Sylke Winkler[2,7], Graham M. Hughes[8], Sofia Traikov[2], Michael Hiller[3,10], Elizaveta Rivkina[1], Philipp H. Schiffer[4*], Eugene W Myers[3], Teymuras V. Kurzchalia[2*]

1. Institute of Physicochemical and Biological Problems in Soil Science RAS, Pushchino, Russia
2. Max Planck Institute for Molecular Cell Biology and Genetics, Dresden, Germany
3. Center for Systems Biology, Dresden, Germany
4. Institute for Zoology, University of Cologne
5. Department of Invertebrate Zoology, Faculty of Biology, M.V. Lomonosov Moscow State University, Moscow, Russia
6. Tree of Life, Wellcome Sanger Institute, Cambridge, UK
7. DRESDEN concept Genome Center, Dresden, Germany
8. School of Biology and Environmental Science, University College Dublin, Belfield, Dublin, Ireland
9. Institute of Biochemistry, ETH Zürich, 8093 Zürich, Switzerland
10. LOEWE Centre for Translational Biodiversity Genomics, Senckenberg Society for Nature Research & Goethe University, Frankfurt am Main, Germany

[#] A.S and V.G contributed equally to this work

*To whom correspondence may be addressed

Email: kurzchalia@mpi-cbg.de

p.schiffer@uni-koeln.de

Abstract

Some organisms in nature have developed the ability to enter a state of suspended metabolism called cryptobiosis[1] when environmental conditions are unfavorable. This state-transition requires the execution of complex genetic and biochemical programs[1,2,3], that enables the organism to survive for prolonged periods. Recently, nematode individuals have been reanimated from Siberian permafrost after remaining in cryptobiosis. Preliminary analysis indicates that these nematodes belong to the genera *Panagrolaimus* and *Plectus*[4]. Here, we present precise radiocarbon dating indicating that the *Panagrolaimus* individuals have remained in cryptobiosis since the late Pleistocene (~46,000 years). Phylogenetic inference based on our genome assembly and a detailed morphological analysis demonstrate that they belong to an undescribed species, which we named *Panagrolaimus n. sp*. Comparative genome analysis revealed that the molecular toolkit for cryptobiosis in *Panagrolaimus n. sp.* and in *C. elegans* is partly orthologous. We show that biochemical mechanisms employed by these two species to survive desiccation and freezing under laboratory conditions are similar. Our experimental evidence also reveals that *C. elegans* dauer larvae can remain viable for longer periods in suspended animation than previously reported. Altogether, our findings demonstrate that nematodes evolved mechanisms potentially allowing them to suspend life over geological time scales.

Introduction

Organisms from diverse taxonomic groups can survive extreme environmental conditions, such as the complete absence of water or oxygen, high temperature, freezing, or extreme salinity. The survival strategies of such organisms include a state known as suspended animation or cryptobiosis, in which they reduce metabolism to an undetectable level[6]. Spectacular examples of long-term cryptobiosis include a *Bacillus* spore that was preserved in the abdomen of bees buried in amber for 25 to 40 million years[7], and a 1000 to 1500 years-old *Lotus* seed, found in an ancient lake, that was subsequently able to germinate[8]. Metazoans such as tardigrades, rotifers and nematodes are also known for remaining in cryptobiosis for prolonged periods[9,10]. The longest records of cryptobiosis in nematodes are reported for the Antarctic species *Plectus murrayi*[11] (25.5 years in moss frozen at -20°C), and *Tylenchus polyhypnus*[12] (39 years desiccated in a herbarium specimen).

Intensive research during the last decades has demonstrated that permafrosts (perennially frozen sediments) are unique ecosystems preserving life forms at sub-zero temperatures over thousands of years[13,14,15,16]. Permafrost remains are an exceptional source for discovering a wide variety of unicellular and multicellular living organisms surviving in cryptobiosis for prolonged periods[6,17,18]. We recently reanimated soil nematodes that were preserved in Siberian permafrost for potentially thousands of years, and initial morphological observations provisionally described them as belonging to the genera *Panagrolaimus* and *Plectus*. Previous studies demonstrated several species of *Panagrolaimus* can undergo cryptobiosis in the form of anhydrobiosis (through desiccation) and cryobiosis (through freezing)[19,20,21,22,23]. In various nematodes, entry into anhydrobiosis is often accompanied by a preparatory phase of exposure to mild desiccation, known as preconditioning[21,24]. This induces a specific remodelling of the transcriptome, the proteome, and metabolic pathways that enhances survival ability[2,3,25]. Some panagrolaimids evolved adaptive mechanisms for rapid desiccation where most of the cellular water is lost, while others evolved freezing tolerance, without loss of water, at sub-zero temperatures by inhibiting the growth and recrystallisation of ice crystals[21]. However, the genetic and biochemical mechanisms of long term cryptobiosis in these organisms have not yet been investigated in detail.

Here we present a high-quality genome assembly and a detailed morphological, phylogenetic analysis and define a novel species, *Panagrolaimus n. sp.* Precise radiocarbon dating indicates that *Panagrolaimus n. sp* remained in cryptobiosis for about 46,000 years, since late Pleistocene. Furthermore, making use of the powerful model organism *C. elegans*,

we demonstrate that *Panagrolaimus* and *C. elegans* dauer larvae utilize similar adaptive

mechanisms to survive extreme desiccation and freezing, i.e. upregulation of trehalose

biosynthesis and gluconeogenesis.

Results

Discovery site and radiocarbon dating

Previously, we showed that nematodes from the Siberian permafrost with morphologies

consistent with the genera *Panagrolaimus* and *Plectus* could be reanimated thousands of years

after they had been frozen in the permafrost. Several viable nematode individuals were found

in two out of more than 300 studied samples of permafrost deposits of different ages and

genesis, collected by researchers of the Soil Cryology Lab, Pushchino, Russia, in the course of

perennial paleo-ecological expeditions carried out in the coastal sector of the northeastern

Arctic[17]. The detailed description of the study site (outcrop Duvanny Yar, Kolyma River,

Fig.1A), sampling and revitalizing procedures are provided in Supplementary Information (SI).

Like other late Pleistocene permafrost formations in the northeastern Arctic, Duvanny Yar

comprises permanently frozen ice-rich silt deposits riddled with large polygonal ice wedges

and divided by them into mineral blocks[26,27] (Fig.1B). The sediments include sandy alluvial

layers, peat lenses, buried paleosols and Pleistocene rodent burrows (Fig.1C). The burrow (P-

1320), in which *Panagrolaimus* nematodes were found (Fig.1D), has been taken from the

frozen outcrop wall at a depth of about 40 m below the surface and about 11 m above river

water level in undisturbed and never thawed late Pleistocene permafrost deposits. The fossil

burrow left by arctic gophers of the genus *Citellus* consists of entrance tunnel and a large

nesting chamber up to 25 cm in diameter [26].

The sterility of permafrost sampling and age of cultivated biota were discussed in detail

in several reviews[14,28, 29]. Based on previous reports, the age of the organisms found in a burrow

is equal to the freezing time and corresponds to the age of organic matter conserved in the

syncryogenic sediments. This makes it possible to use radiocarbon dating of organic matter to

establish the age of organisms. We performed Accelerator Mass Spectrometry (AMS)

radiocarbon analysis of plant material obtained from studied borrow P-1320 and determined a

direct ^{14}C age of 44,315±405 BP (Institute of Geography, RAS; sample IGAN$_{AMS}$ 9137).

Calibrated age range is 45,839 – 47,769 cal BP (95.4% probability) (Fig.S1).

Like other parthenogenetic *Panagrolaimus* the newly discovered species is triploid

The revived animal was cultivated in the laboratory for over 100 generations and initially described as *Panagrolaimus* aff. d*etritophagus*[4] based on morphology. We conducted a detailed morphological analysis of the revived animal (Fig.2, S2, Table S1;BOX1), which confirmed unambiguously that the animal belongs to the genus of *Panagrolaimus*, in agreement with a previous phylogenetic analysis of the 18S ribosomal RNA sequence[4]. However, due to the morphological uniformity of *Panagrolaimus*, unusual even for nematodes, morphology and molecular analysis of a single ribosomal RNA sequence is insufficient to describe a species. We found the species to be parthenogenetic, which further complicates description under most species concepts.

To obtain comprehensive molecular data for species determination using phylogenomics, we generated a genome assembly. Using PacBio HiFi sequencing, we generated 84X coverage in long reads (average length 14,425 bp). Our analysis of repeat and gene content is described in Supplementary Table 3. K-mer analysis of the reads clearly indicated that this animal has a triploid genome (Fig.3A), similar to other parthenogenetic *Panagrolaimus* species[30]. Despite the challenges that a triploid genome poses for assembly, we obtained a highly-contiguous contig assembly of the three pseudohaplotypes that comprise about 266 Mb and thus have a similar genome size as other parthenogenetic *Panagrolaimus* species[30]. The contig N50 value of all three pseudohaplotypes is 3.8 Mb. Since the three pseudohaplotypes exhibited a noticeable degree of divergence, we further investigated their relationship by using the apparent homeologs in our gene predictions to align long continuous contigs based on micro synteny (Fig. 3B). Links between the contigs clearly shows the triploid state of the genome.

 BOX 1 Description of *Panagrolaimus n. sp* sp. nov.

Description	Body spindle-shaped and usually almost straight after fixation (Fig.2a, b). Cuticle thin and faintly annulated, 10–11 annules per 10 μm in cervical region, 13–14 in midbody, preanally again 10 annules withi 10 μm. Conspicuous convex lateral fields with three incisures 1.5-2 μm wide extended along the body from about ¼–1/3 procorpus length to 2/3 of tail length. In SEM, the lateral field looks as a bolster with a narrow median split. Labial region set off. Mouth opening surrounded with six lips (Fig.2c, d). Anterior sensilla as papillae arranged in two close but separate subsequent circles. Somatic sensilla (i.e., deirids and phasmids) not evident.Buccal cavity cylindro-conoid, and unarmed; its total length 9–13 μm, maximum stoma width 1.7–2.8 μm (Fig.2e). Dorsal stoma wall (dorsal rhabdion) more clearly sclerotized. Anterior part of the buccal cavity comprising cheilostom and gymnostom nearly cylindroids while stegostom conically narrowed and ended with a distinct tight flexion. Pharynx consists of three distinct parts: straight anterior procorpus, narrow medial isthmus and rounded terminal bulb. Procorpus gradually widening to it posterior end, always straight in all specimens, with transversal muscular striation, more prominent in posterior three fourths. Isthmus narrow, cylindroid, bent starkly in nearly all studied specimens. Terminal bulb strongly muscular, with a valvular apparatus at about 40% of the bulb length. Cardia in shape of truncate cone. Intestine (midgut) tissue filled with vacuoles and granules; in the anteriormost region, the granules smaller and look more pallid. Cell borders not visible in the intestine, but internal lumen distinct, sinusoid. No internal content visible in the internal lumen. Ventral excretory-secretory pore and its cuticularized duct situated at the level of anterior part of the bulb. No other details of the excretory-secretory system visible. Vulval lips protruding. Genital branch monodelphic prodelphic and situated dorsally and to the right of the midgut (Fig.2f). Vagina distinctly cuticularized and opens to the elongate uterus. A long oviduct extended anteriad from the anterior uterus; the oviduct folded up and then posteriorward and transforms into an elongate ovary. There are

	one or two ripe eggs in the uterus in most specimens. Tail short conical, with short acute spike-like mucro (Fig.2g).
Etymology	Species name *kolymaensis* (Latin) is derived from the Kolyma River area.
Holotype	Senckenberg Natural History Museum, Frankfurt am Main, Germany (collection number SMF 17067).
Paratypes	Senckenberg Natural History Museum, Frankfurt am Main, Germany (collection numbers SMF 17068, SMF 17069) (eighteen paratypes).
Type locality	Frozen fossil rodent burrow buried in permafrost 45,839 – 47,769 cal BP , 40 meters from the surface, outcrop Duvanny yar, Kolyma River, North-East of Siberia, Russia (68° 37′ 739″ N, 159° 11′ 678″ E). Frozen material from burrow was collected by Dr.S.Gubin (Soil Cryology Lab, Pushchino, Russia) in august 2002.

Phylogenomics to define *Panagrolaimus n. sp* sp. nov.

To place the species in the genus *Panagrolaimus*, we conducted a broader multi-gene
phylogenomic analysis using Maximum likelihood methods. Our analysis of a concatenated,
partitioned alignment of 60 genes, and a coalescence-based approach using a broader set of
12,295 gene trees, retrieves the novel species as sister to all other sequenced *Panagrolaimus*
species, but as an ingroup to *Propanagrolaimus*[31] (Fig.3C; Fig. S3B-C). Thus, the phylogenetic
placement provides strong evidence that this animal belongs to a novel species. Further
supporting this, there is substantial sequence divergence between this novel species, and
Panagrolaimus sp. PS1159 and *Panagrolaimus* sp. ES5, estimated to be on average 2.06 and
2.11 amino acid substitutions per site in our concatenated alignment, respectively. The
substantial divergence is in line with previous data on ages of *Panagrolaimus* nematodes[30], and
more broadly seen in nematodes, which can be hyper-diverse[32,33]. Our data also contradicts the
assumption that parthenogenesis is monophyletic trait[30] in the *Panagrolaimus* genus (Fig.3C).
Based on the Kolyma River location where the animal was unearthed, we propose the following
taxonomic classification and species name:
Phylum Nematoda Potts, 1932
Class Chromadorea Inglis 1983

Suborder Tylenchina Thorne, 1949

Family Panagrolaimidae Thorne, 1937

Panagrolaimus n. sp sp. nov.

C. elegans dauer larvae and *Panagrolaimus n. sp.* utilize similar mechanisms to enter and

remain in cryptobiotic state for prolonged periods of time

In the absence of established genetic methods in *Panagrolaimus n. sp.* we referred to *C. elegans*

as a powerful model system to gain insights into possible pathways for long term survival[3,4,24,25].

The high-quality genome of *Panagrolaimus n. sp.* allowed us to compare its molecular toolkit

for cryptobiosis with that of *C. elegans*. We used orthology clustering and phylogenetics to

investigate whether the genome of *Panagrolaimus n. sp.* contains genes previously implicated

in cryptobiosis in the *C. elegans* dauer larva. Our analysis showed that the *Panagrolaimus n.*

sp. genome encodes orthologs to a *C. elegans* trehalose phosphate synthase gene (*tps-2*) and to

a trehalose phosphatase gene (*gob-1*) (Fig. 3D, supplementary file Orthology analysis).

Furthermore, we found orthologs to all *C. elegans* enzymes required for polyamine

biosynthesis, the TCA cycle, glycolysis, gluconeogenesis, and glyoxylate shunt (Fig.3D,

supplementary file Orthology analysis) suggesting that *Panagrolaimus n. sp.* utilizes a similar

molecular tool kit as *C. elegans* to facilitate survival of unfavorable conditions.

202   Our earlier findings established that among several developmental stages of *C. elegans*,

only the dauer larva, formed during unfavorable conditions (such as low nutrients and high

population density) could survive anhydrobiosis and freezing[4,24]. To survive extreme

desiccation, *C. elegans* dauer larvae need to be first preconditioned at high relative humidity

(98% RH) for 4 days[24]. During preconditioning, dauer larvae upregulate trehalose biosynthesis

that ensures their survival to harsh desiccation[24,25]. We tested whether preconditioning also

facilitates survival of *P. kolymaensis*. Although a small proportion of *Panagrolaimus n. sp.*

individuals survive harsh desiccation and freezing without preconditioning (Fig.4A), mixed

populations of *Panagrolaimus n. sp.* survive significantly (p value< 0.0001) higher to harsh

desiccation upon preconditioning (Fig.4A). Similarly, preconditioning and desiccation further

enhanced survival rate of *Panagrolaimus n. sp.* to freezing (-80°C). Like *C. elegans,*

Panagrolaimus n. sp. upregulate trehalose levels up to 20-fold upon preconditioning (Fig.4B).

We previously reported that, to upregulate trehalose levels upon preconditioning, *C. elegans*

dauer larva dissipate their fat reserves (Triacylglycerols) by activating the glyoxylate shunt and

gluconeogenic pathway[25]. Upon preconditioning, we found that triacylglyceride (TAG) levels

are significantly decreased in *Panagrolaimus n. sp.* (Fig.S5A&B). To further investigate

whether the acetyl-CoA derived from degradation of TAGs culminate in trehalose, we applied

the previously developed method of metabolic labelling with [14]C-acetate in combination with
2D-TLC [4,25]. As shown in Fig. 4C, preconditioning led to huge increase of radioactivity in
trehalose and to smaller extent in some amino acids (glycine/serine, phenylalanine; panels c
and d). Interestingly, *Panagrolaimus n. sp.* displayed an additional spot (Fig. 4D, enumerated
as 7), that was not found in *C. elegans*, which we identified as trehalose-6-phosphate (Fig.S5C-
H), an immediate precursor of trehalose, using mass spectrometry. Thus, to resist harsh
desiccation, like *C. elegans*, *Panagrolaimus n. sp.* utilizes glyoxylate shunt and consequently
acetate derived from TAGs to synthesize trehalose. Detection of the immediate precursor
(trehalose-6-phosphate) suggests the flux of metabolites is intense in the latter.

Finally, we investigated whether *C. elegans* can also survive in prolonged cryptobiotic
state. Despite of preconditioning, the survival ability of desiccated dauer larvae at room
temperature declines very rapidly, with most larvae dead after around 10 days (Fig. 4E) (Fig.
S4A&B). Direct freezing at -80℃, leads to almost instant death of the animals. To test whether
combining these conditions could extend the viability of dauer larvae (Fig. S4A&B), we
transferred the desiccated larvae to -80℃. Remarkably, under these conditions, there was no
significant decline in viability even after 480 days (Fig. 4E). Moreover, after thawing, the
animals resumed reproductive growth and produced progeny in numbers like those of animals
kept under control conditions (Fig. 4F). Since no reduction in survival was observed at any
time point in extended experiments, these results suggest that the combination of anhydrobiosis
and freezing can prolong the survival ability of dauer larvae. Thus, *C. elegans*, when exposed
to combination of cryptobiotic states can survive for extremely long periods of time.

Discussion

Our findings here demonstrate the ability of the nematodes to survive in suspended animation
for geological time scales. Along with the adaptive mechanisms of an organism, a naturally
preserving ecosystem would synergistically aid the survival for prolonged periods. The
Siberian permafrost is a unique repository for preserving organisms in sub-zero temperatures
for millions of years. Expeditions in the past decade revived several organisms across various
taxa from the Siberian permafrost[5,34,35,36]. The possibility to exploit permafrost as a source for
reanimating multicellular animals was recognized already in 1936. A viable Cladocera
crustacean *Chydorus sphaericus* preserved in the Transbaikalian permafrost for several
thousand years[37,38] was discovered by P.N. Kapterev, who worked at the scientific station
Skovorodino as a GULAG prisoner. Unfortunately, this extraordinary observation remained
unnoticed for many decades.

The new species can now be placed into the genus *Panagrolaimus*[39], which contains
several described and undescribed parthenogenetic and gonochoristic species and strains[30,40].
Many *Panagrolaimus* display adaptation to survival in harsh environments[21] and the genus
includes the Antarctic species *P. davidii*[22]. The genus *Panagrolaimus* is exceptional in its
morphological uniformity even among nematode species that are hard to classify based on
morphology in general. Thus, species designation via microscopic (including SEM) analysis is
unreliable, which is further complicated by the absence of males in parthenogenetic species
makes. Males have important diagnostic features as spicules and pericloacal papillae, females
differ from one species to another mainly by morphometrics, whereby the interspecies
differences (absolute measures and ratios) may be very fine. Our specimens are close by
absolute sizes and ratios to females of the bisexual species *Panagrolaimus detritophagus*[41].
The only non- overlapping morphometric character is index "b" (body length: pharynx length):
5.6–6.8 in *Panagrolaimus n. sp.* versus 4.4-5.1 in *P. detritophagus*.

Consequently, we turned to phylogenomics methods to place the species on the tree.
This showed that species is an outgroup to other known *Panagrolaimus* species, raising for the
possibility of a second independent evolution of parthenogenesis in the genus, in contrast to
previous findings[40,31,30]. This will need to be confirmed through further sampling and genome
sequencing of *Panagrolaimus* species. We found *Panagrolaimus n. sp.* to be triploid and thus
a hybrid origin is possible, as seen in other parthenogenetic *Pangrolaimus*[30]. The highly
contiguous genome of *Panagrolaimus n. sp.* will allow for analyses of this trait in comparison
to other *Panagrolaimus* species currently being genome sequenced. The successful
phylogenomic species identification we conducted here demonstrates the utility of this method
for morphologically hard to identify species, in particular when the additional data yielded by
a fully sequenced genome are considered.

Our results provide a deeper insight into the homology of molecular and biochemical
mechanisms between *C. elegans* and *P. kolymaensis*, which are not only taxonomically but
also ecologically distinct. *C. elegans* can mostly be found in rotting fruits and plants in
temperate regions[42,43], while *Panagrolaimus* species are globally distributed and prevalent in
leaf litter and soil[40], including in harsh environments[21]. We show through orthology analysis
that the well-studied molecular pathways used by *C. elegans* larvae to enter the dauer state,
such as insulin[44,45] (DAF-11, DAF-2 & DAF-16), TGF-β[46] (DAF-7), steroid[47] (DAF-9, DAF-
12) are present in the genome of the *Panagrolaimus n. sp.* (Fig.S4C). While further functional

analyses are needed to study molecular pathways detail, our results hint at convergence or
parallelism in the molecular mechanisms organizing dauer formation and cryptobiosis.

As mentioned earlier preconditioning enhances the survival of *Panagrolaimus n. sp.*
to desiccation and freezing. We previously reported that preconditioning induces trehalose
upregulation in *C. elegans* dauer larvae and the elevated trehalose renders desiccation tolerance
by protecting the cellular membranes[24]. It is not surprising that *Panagrolaimus n. sp.*
upregulates trehalose, however the magnitude of trehalose elevation is higher than *C. elegans*
dauer larvae. This indicates that central regulators (DAF-16, DAF-12) of trehalose
upregulation may differentially regulate *tps-2* in *Panagrolaimus n. sp* [48,49]. Although
Panagrolaimus n. sp. utilizes glyoxylate shunt and gluconeogenesis to upregulate trehalose
levels, it is intriguing to observe that they accumulate substantial levels of trehalose-6-
phosphate. Further investigation of this observation using perturbation experiments will
provide insights into metabolic regulation in *Panagrolaimus n. sp.* upon preconditioning. Our
findings for the first time demonstrate that *C. elegans* dauer larvae possess an inherent ability
to survive freezing for prolonged periods if they undergo anhydrobiosis. It is tempting to
speculate that undergoing anhydrobiosis might be a survival strategy of *C. elegans* to survives
seasonal changes in nature.

In summary our findings indicate that by adapting to survive cryptobiotic state for
short time frames in environments like permafrost, some nematode species gained the potential
for individual worms to remain in the state for geological timeframes. This raises the question
whether there is an upper limit to the length of time an individual can remain in the cryptobiotic
state, potentially only limited by drastic changes to the environment such as strong fluctuations
in ambient temperature, natural radioactivity, or other abiotic factors. These findings have
implications for our understanding of evolutionary processes, since generation times could be
stretched from days to millennia, species ages might be much longer than anticipated, and long-
term survival of individuals of species can lead to the refoundation of otherwise extinct lineages.
Finally, understanding the precise mechanisms of long-term cryptobiosis and cues that lead to
successful revivals can inform new methods for long term storage of cells and tissues.
312

Methods

Materials and *C. elegans* strains

[1-14C] -acetate (sodium salt) from Hartmann Analytic (Braunschweig, Germany). All other chemicals were purchased from Sigma-Aldrich (Taufkirchen, Germany). The *Caenorhabditis* Genetic Centre (CGC) provided *daf-2(e1370)* and *E. coli NA22* strains.

Genomic DNA isolation from *Panagrolaimus n. sp* nematodes

Panagrolaimus n. sp. nematodes were grown on several plates of NGM agar plated with *E. Coli NA22* bacteria at 20°C. Worms were collected from the plates, washed with water at least three to five times by centrifugation at 1000 g to remove any residual bacteria and any debris. The worm pellet was dissolved in 5 volumes of worm lysis buffer (0.1M Tris-HCl pH=8.5, 0.1M NaCl, 50mM EDTA pH=8.0) and distributed in 1.5 ml of microcentrifuge tubes. These tubes are incubated at -80°C for 20 minutes. 100 µl of Proteinase 'K' (20 mg/ml) was added to each tube and they are incubated at 60°C overnight. 625 µl of cold GTC buffer (4M Guanidinium Thiocynate, 25mM Sodium citrate, 0.5% (v\v) N-lauroylsarcosine, 7%(v/v) Beta Mercaptoethanol) was added to the tube, incubated on ice 30 min, and mixed by inverting every 10 min. 1 volume of phenol–chloroform-isoamyl alcohol (pH=8) was added to the lysate and mixed by inverting the tube 10-15 times. Tubes were centrifuged for 5 min at 10,000 g at 4 °C to separate the phases. The upper aqueous phase was carefully collected into a fresh tube. One volume of fresh chloroform was added and mixed by inverting the tubes for 10-15 times and centrifuged for 5 min at 10,000 g at 4°C to separate the phases. One volume of cold 5 M NaCl was added, mixed by inverting the tubes and incubated on ice for 15 min. After incubation these tubes were centrifuged for 15 min at 12,000–16,000 g at 4 °C. The supernatant containing the nucleic acids were slowly transferred into a fresh tube. One volume of isopropanol was added to the tube, inverted few times, and incubated on ice for 30 minutes. After incubation, the tubes were centrifuged at 3000 g for 30–45 min at 25°C and the supernatant was discarded without disturbing the pellet. The pellet was washed twice with 1 ml of 70% ethanol, tubes were centrifuged at 3000 g for 5 min and supernatant was discarded and incubated at 37°C for 10-15 min to dry the pellet. The pellet was resuspended carefully in TE buffer. The quality of the genomic DNA was analyzed with pulse field gel electrophoresis.

Genome sequencing and assembly

Long insert library has been prepared as recommended by Pacific Bioscienes according to theprotocol 'Procedure & Checklist-Preparing gDNA Libraries Using the SMRTbell® Express Template Preparation Kit 2.0'. In summary, RNAse treated HMW gDNA has been sheared to 20Kb fragments on the MegaRuptor^{TM} device (Diagenode) and 10 µg sheared gDNA have been used for library preparation. The PacBio SMRTbell^{TM} library was size selected in two fractions (9-13kb, > 13kb) using the BluePippin^{TM} device with cassette definition of 0.75% DF MarkerS1 3-10 kb Improved Recovery. The second fraction of the size selected library has been loaded with 95 pM on plate on a Sequel SMRT cell (8M). Sequel polymerase 2.0 has been used in combination with the v2 PacBio sequencing primer and the Sequel sequencing kit 2.0EA, runtime was 30 hours. We created PacBio CCS reads from the subreads.bam file using PacBio's ccs command linetool (version4.2.0). We obtained 8.5Gb hiqh quality CCS reads (HiFi reads) with a N50 of 14.4 Kb. We ran HiCanu (version 2.2)[39] to create the contig assembly. Blobtools (version 1.1.1,CITE:) was used to identify and remove bacterial contigs. The final triploid contig assembly consists of 856 contigs has a N50 of 3.82 Mb and a size of 266Mb. The mitochondrial genome was created with the mitoHifi pipeline (version 2, https://github.com/marcelauliano/MitoHiFi) based on the assembled contigs and the closely related reference mitochondrial genome of Panagrellus redivivus (strain: PS2298/MT8872, ENAaccession: AP017464). The mitoHifi pipeline identified 49 mitochondrial contigs ranging from 13-32Kb. The final annotated circular mitochondrial genome has length of 17467 bp.

To identify pseudohaplotypes in the *P. kolymanensis* sp. nov genome assembly, we selected the longest isoform of each predicted protein-coding gene in our assembly and in the *C. elegans* genome (downloaded from WormBase Parasite, release WBPS15) using AGAT (version 0.4.0) and clustered them into orthologous groups (OGs) using OrthoFinder (version 2.5.2). We identified OGs that contained three *Panagrolaimus* sequences (i.e. groups that were present as single-copy in all three pseudohaplotypes) and used these to identify trios of multi-megabase size contigs derived from the three pseudohaplotypes. We visualized synteny between the three pseudohaplotypes using Circos to plot the positions of each homeolog (version 0.69-8).

Genome annotation

We used RepeatModeler 1.0.8 (http://www.repeatmasker.org/) with parameter '-engine ncbi' to create a library of repeat families. This library was used with RepeatMasker 4.0.9 to soft-mask the *Panagrolaimus* genome. To annotate genes, we cross mapped protein models from

an existing *Panagrolaimus* as external evidence in the Augustus[51] based pipeline. We evaluated

the completeness of our predictions using BUSCO on the gVolante web interface.

Orthology analysis

We conducted a gene orthology analysis using genomic data from *Panagrolaimus n. sp*, the

plectid nematode species from the permafrost, as well as genomic data from WormBase

Parasite (https://parasite.wormbase.org; accessed 17/12/2020): *Caenorhabditis elegans*,

Diploscapter coronatus, *Diploscapter pachys*, *Halicephalobus mephisto*, *Panagrellus*

redivivus, *Panagrolaimus davidi*, *Panagrolaimus* sp. ES5, *Panagrolaimus* sp. PS1159,

Panagrolaimus superbus, *Plectus sambesii*, and *Propanagrolaimus* sp. JU765. For plectids,

genomic resources are scarce. We therefore added transcriptome data of *Plectus murrayi*,

Anaplectus granulosus, *Neocamacolaimus parasiticus*, and *Stephanolaimus elegans*. The latter

three transcriptomes were kindly provided by Dr. Oleksandr Holovachov (Swedish museum of

natural history). The *Anaplectus granulosus*, and *Neocamacolaimus parasiticus* transcriptomes

have been published [52,53]. All three transcriptomes were assembled de novo with Trinity[54]. The

exact procedures are described in the respective publications[52,53] . The *Stephanolaimus* elegans

transcriptome was assembled in the same way as described for *Neocamacolaimus parasiticus*.

The *Plectus murrayi* transcriptome was built from raw reads deposited at NCBI (https://sra-

[downloadb.be-md.ncbi.nlm.nih.gov/sos2/sra-pub-run-13/SRR6827978/SRR6827978.1](https://sra-downloadb.be-md.ncbi.nlm.nih.gov/sos2/sra-pub-run-13/SRR6827978/SRR6827978.1);

accessed 22.12.2020). The transcriptome was assembled using Galaxy Trinity version 2.9.1[55],

[54] using all default options and including in silico normalization of reads before assembly.

Transdecoder (conda version 5.5.0)[56] was used to translate to amino acid sequence. Identical

reads were removed with cd-hit version 4.8.1[57,58], and the Trinity

get_longest_isoform_seq_per_trinity_gene.pl command[56] (Trinity conda version 2.8.5)

{Anaconda Software Distribution, Conda, Version 4.9.2, Anaconda, Nov. 2020; was used, to

remove shorter isoforms. Amino acid translated longest isoforms from genomic data were

extracted with AGAT {Dainat, https://www.doi.org/10.5281/zenodo.3552717} from genome

assembly FASTA files and genome annotation GFF3 files, using the

agat_convert_sp_gxf2gxf.pl command to convert all GFF3 files to the required GFF3 format,

and the agat_sp_keep_longest_isoform.pl as well as the agat_sp_extract_sequences.pl

commands to extract amino acid translated FASTA files. The headers and names of all FASTA

files were modified to allow for simple species assignment of each sequence in subsequent

analysis. The orthology analysis was conducted with OrthoFinder v. 2.5.1[59,60] using default

settings. For genes of interest, we constructed alignments with MAFFT v. 7.475[61] using the

localpair and maxiterate (1000) functions. We removed spurious sequences and areas that were

not well aligned with Trimal v. 1.4.rev22[62] (procedure stated in supplementary file Orthology

analysis below each phylogeny). We then ran phylogenetic analysis with Iqtree2 v. 2.0.6[63],

with -bb 1000 option, testing the model for each analysis (models eventually used stated in

supplementary file Orthology analysis). We also checked for PFAM domains using

Interproscan v. 5.50-84.0[64]. Part of the analysis was performed on the HPC RRZK CHEOPS

of the Regional Computing Centre (RRZK) of the University of Cologne. The phylogenies

were visualized with Dendroscope 3.7.6[65] and figures were created with Inkscape

{https://inkscape.org}.

Phylogenomics

Sequences of 18S and 28S genes from 44 taxa across the *Propanagrolaimus*, *Panagrolaimus*,

Panagrellus and *Halicephalobus* genera were aligned (MAFFT L-INS-I v7.475)[61],

concatenated [66]and used to infer a species tree using maximum likelihood via (IQTREE)[67] and

partitioned by best-fit models of sequence evolution for both[68]. Nodal support was determined

using 1000 bootstrap pseudoreplicates. A further 60 genes from 101 taxa were used to confirm

the taxonomic position using the supermatrix concatenation methods outlined above. Given the

limitations of differential gene sampling, we expanded our phylogenomic analyses to include

a coalescence approach using 12,295 ML gene trees inferred for orthogroups containing the

target animal. Instances of multiple genes per species per group were treated as

paralogs/orthologs and analysed using ASTRAL-Pro[69].

Desiccation survival assay

C. elegans dauer larvae desiccation assays were performed as described in. *Panagrolaimus n.*

sp desiccation assays were performed with mixed populations of the worms.

Exposure of nematodes to extreme environments

Dauer larvae or *Panagrolaimus n. sp* nematodes were preconditioned and desiccated as

described[24] and then transferred to elevated temperature of 34°C, freezing (-80°C) and anoxia.

Anoxic environment was generated in a desiccation chamber at 60%RH by flushing the

Nitrogen gas into the chamber. The concentration of oxygen inside the chamber was monitored.

After each timepoint they were rehydrated with 500 µl of water for 2-3 hours. Rehydrated

worms were transferred to NGM agar plates with *E. Coli NA22* as food. Survivors were counted

after overnight recovery. Each experiment was performed on two different days with at least

two technical replicates.

Trehalose quantification from nematode lysates

Trehalose measurements were performed as described in previous reports[25].

Radiolabeling, metabolite extraction and 2D-TLC

The above-mentioned procedures were performed according to previous reports[3,25].

Identification of trehalose-6-phosphate from TLC plates

Normalised aqueous fractions from the non-preconditioned and preconditioned samples were

separated by high performance thin layer chromatography (HPTLC), using 1-propanol-

methanol-ammonia (32%)-water (28:8:7:7 v/v/v/v) as first, dried for 15 min and 1-butanol-

acetone-glacial acetic acid–water (35:35:7:23 v/v/v/v) second dimension respectively. Using

the trehalose as a standard on both dimensions of the TLC, the regions of interest were scrapped

out from the TLCs. The scraped-out silica was extracted with 10 ml of 50% methanol twice.

The fractions were combined, dried under vacuum and dissolved in 100 µl of Ms mix solution

containing 4:2:1 (Isopropanol:Methanol:Chloroform) with 7.5 mM ammonium formate. Mass

spectrometric analysis was performed on a Q Exactive instrument (Thermo Fischer Scientific,

Bremen, DE) equipped with a robotic nanoflow ion source TriVersa NanoMate (Advion

BioSciences, Ithaca, USA) using nanoelectrospray chips with a diameter of 4.1 µm. The ion

source was controlled by the Chipsoft 8.3.1 software (Advion BioSciences). Ionization voltage

was + 0.96 kV in negative mode; backpressure was set at 1.25 psi. The temperature of the ion

transfer capillary was 200°C; S-lens RF level was set to 50%. FT MS spectra were acquired

within the range of m/z 50–750 at the mass resolution of R m/z 200 = 140000; automated gain

control (AGC) of 3×10^6 and with the maximal injection time of 3000 ms. FT MS/MS spectra

were acquired within the range of m/z 50–750 at the mass resolution of R m/z 200 = 140000;

automated gain control (AGC) of 3×10^4 and with the maximal injection time of 30 s.

Triacylglycerols measurement from *Panagrolaimus n. sp* lysates

Non-preconditioned and preconditioned pellets were lysed in 200 µl of isopropanol with 0.5

mm Zircornium beads twice for 15 min. The lysates were centrifuged at 1300 g for 5 min. The

supernatant was carefully collected without any debris, 20 µl of the lysate was used for protein

estimation. Normalization was performed according to soluble protein levels, supernatant

volumes corresponding to 50-100 µg of proteins were dried in the desiccator. 700 µl of IS ((10:3 (Methyl tert-butyl ether: ethanol)) mix (warmed to room temperature) was added to dried samples and left on a shaker for 1 hour. The samples were centrifuged at 1400 rpm and 4°C. 140 µl of water was added and left on shaker for 15 min. These samples were centrifuged at 13400 rpm for 15 min. The upper organic fraction was collected and transferred to 1.5 ml glass vial and left for drying in the desiccator. The dried samples were reconstituted in appropriate volume of 300 µl of 4:2:1 (Isopropanol:Methanol:Chloroform) and volume corresponding 1 µg was used for injection.

LC-MS/MS analysis was performed on a high-performance liquid chromatography system (Agilent 1200 HPLC) coupled to a Xevo G2-S QTof (Waters). The samples were resolved on a reverse phase C18 column (Cortecs C18 2.7um from Waters) with 50:50:0.1:1% (Water:methanol:formic acid: 1M Ammonium formate) and 25:85:0.1:1% (Acetonitrile:Isopropanol:Formic acid:1M Ammonium formate) as mobile phase. The following gradient program was used: Eluent B from 0 % to 100 % within 12 min; 100 % from 12 min to 17min; 0 % from 17 min to 25 min. The flow rate was set at 0.3 ml/min. The samples were normalised according to the total protein concentration and the worm numbers. TGD 50:00:00 was used as internal standard. Skyline software was used to analyse the raw data. The TG were extracted from Lipidmaps database.

Author contributions

AS, VG, PS, TK conceived and designed the study. AS, VG, TH, MP, AT, GH, MH, ER, PS and TK contributed to the original draft. AS performed isolation and cultivation of nematodes. VG performed desiccation survival assays, trehalose measurement, 2D-TLC of metabolites, trehalose-6-phosphate detection, combination of cryptobiosis experiments, genomic DNA isolation from *Panagrolaimus n. sp*, assembled the data from all the authors and prepared the figures. MP performed genome assembly. TH conducted orthology analyses and single gene phylogenies and worked on figures. AT provided morphological description, light, and scanning electron microscopy. LS analyzed the genome assembly and created figures. GMH performed the phylogenetic analyses. ST performed triacylglycerols measurements, trehalose-6-phosphate detection with suggestions from Andrej Shevchenko. MH annotated supplementary table 3. PS performed genome annotation and supervised TH and LS. EWM supervised MP and had overall responsibility for sequencing and assembly including funding it.

Acknowledgements

VG is thankful to Andrej Shevchenko, members of Kurzchalia lab for helpful discussions and core facilities of MPI-CBG for assistance. We are grateful to Dr. S. Gubin for field study and sampling, our colleagues in Soil Cryology Lab, Pushchino and North-East Scientific Station in Chersky, Republic of Sakha (Yakutia) for their help and cooperation. The authors thank Long Read Team of the DRESDEN-concept Genome Center, DFG NGS Competence Center, part of the Center for Molecular and Cellular Bioengineering (CMCB), Technische Universität Dresden and MPI-CBG. GMH is funded by a UCD Ad Astra Fellowship. This work was supported by the Russian Foundation for Basic Research (19-29-05003-mk) to AS and ER. VG and TK acknowledge the financial support from the Volkswagen Foundation (Life? research grant 92847). PS and TH are supported by a DFG ENP grant to PS (DFG project 434028868). The authors are thankful to Richard Roy and Jens Bast for critical reading the manuscript. We thank Iain Pattern for suggestions in writing the manuscript. The funders have no role in the design of the study.

Conflict of interest:

The authors declare they have no conflict of interest relating to the content of this article.

References

1. Hibshman, J. D., Clegg, J. S. & Goldstein, B. Mechanisms of Desiccation Tolerance: Themes and Variations in Brine Shrimp, Roundworms, and Tardigrades. *Front. Physiol.* **11**, 592016 (2020).

2. Erkut, C. *et al.* Molecular Strategies of the Caenorhabditis elegans Dauer Larva to Survive Extreme Desiccation. *PLoS ONE* **8**, e82473 (2013).

3. Gade, V. R., Traikov, S., Oertel, J., Fahmy, K. & Kurzchalia, T. V. C. elegans possess a general program to enter cryptobiosis that allows dauer larvae to survive different kinds of abiotic stress. *Sci. Rep.* **10**, 13466 (2020).

4. Shatilovich, A. V. *et al.* Viable Nematodes from Late Pleistocene Permafrost of the Kolyma River Lowland. *Dokl. Biol. Sci.* **480**, 100–102 (2018).

5. Shmakova, L. *et al.* A living bdelloid rotifer from 24,000-year-old Arctic permafrost. *Curr. Biol.* **31**, R712–R713 (2021).

6. Keilin, D. The Leeuwenhoek Lecture - The problem of anabiosis or latent life: history and current concept. *Proc. R. Soc. Lond. Ser. B - Biol. Sci.* **150**, 149–191 (1959).

7. Cano, R. & Borucki, M. Revival and identification of bacterial spores in 25- to 40-million-year-old Dominican amber. *Science* **268**, 1060–1064 (1995).

8. Shen-Miller, J. Sacred lotus, the long-living fruits of China Antique. *Seed Sci. Res.* **12**, 131–143 (2002).

9. Guidetti, R. & Jönsson, K. I. Long-term anhydrobiotic survival in semi-terrestrial micrometazoans. *J. Zool.* **257**, 181–187 (2002).

10. Tsujimoto, M., Imura, S. & Kanda, H. Recovery and reproduction of an Antarctic tardigrade retrieved from a moss sample frozen for over 30 years. *Cryobiology* **72**, 78–81 (2016).

11. Kagoshima, H. *et al.* Multi-decadal survival of an Antarctic nematode, Plectus murrayi, in a -20°C stored moss sample. 9.

12. Aroian, R. V., Carta, L., Kaloshian, I. & Sternberg, P. W. A Free - living Panagrolaimus sp. from Armenia Can Survive in Anhydrobiosis for 8.7 Years. 3.

13. Gilichinsky, D. A. *et al.* Microbial Populations in Antarctic Permafrost: Biodiversity, State, Age, and Implication for Astrobiology. *Astrobiology* **7**, 275–311 (2007).

14. Rivkina, E. *et al.* Earth's perennially frozen environments as a model of cryogenic planet ecosystems. *Permafr. Periglac. Process.* **29**, 246–256 (2018).

15. Sipes, K. *et al.* Eight Metagenome-Assembled Genomes Provide Evidence for Microbial Adaptation in 20,000- to 1,000,000-Year-Old Siberian Permafrost. *Appl. Environ. Microbiol.* **87**, (2021).

16. Liang, R. *et al.* Genomic reconstruction of fossil and living microorganisms in ancient
Siberian permafrost. *Microbiome* **9**, 110 (2021).

17. Gilichinsky, D. A. & Rivkina, E. M. Permafrost microbiology. 7 (2011).

18. Yashina, S. *et al.* Regeneration of whole fertile plants from 30,000-y-old fruit tissue buried
in Siberian permafrost. *Proc. Natl. Acad. Sci.* **109**, 4008–4013 (2012).

19. Adhikari, B. N., Wall, D. H. & Adams, B. J. Desiccation survival in an Antarctic nematode:
molecular analysis using expressed sequenced tags. *BMC Genomics* **10**, 69 (2009).

20. Tyson, T. *et al.* A molecular analysis of desiccation tolerance mechanisms in the
anhydrobiotic nematode Panagrolaimus superbus using expressed sequenced tags. *BMC Res.*
Notes **5**, 68 (2012).

21. McGill, L. M. *et al.* Anhydrobiosis and Freezing-Tolerance: Adaptations That Facilitate the
Establishment of Panagrolaimus Nematodes in Polar Habitats. *PLOS ONE* **10**, e0116084
(2015).

22. Thorne, M. A. S., Kagoshima, H., Clark, M. S., Marshall, C. J. & Wharton, D. A. Molecular
Analysis of the Cold Tolerant Antarctic Nematode, Panagrolaimus davidi. *PLoS ONE* **9**,
e104526 (2014).

23. Sandhove, J., Spann, N. & Ristau, K. The Anhydrobiotic Potential of the Terrestrial
Nematodes *Plectus parietinus* and *Plectus velox*: Anhydrobiosis of *plectus parietinus* and *p.*
velox. *J. Exp. Zool. Part Ecol. Genet. Physiol.* **325**, 434–440 (2016).

24. Erkut, C. *et al.* Trehalose Renders the Dauer Larva of Caenorhabditis elegans Resistant to
Extreme Desiccation. *Curr. Biol.* **21**, 1331–1336 (2011).

25. Erkut, C., Gade, V. R., Laxman, S. & Kurzchalia, T. V. The glyoxylate shunt is essential
for desiccation tolerance in C. elegans and budding yeast. *elife* **5**, (2016).

26. Zanina, O. G., Gubin, S. V., Kuzmina, S. A., Maximovich, S. V. & Lopatina, D. A. Late-
Pleistocene (MIS 3-2) palaeoenvironments as recorded by sediments, palaeosols, and
ground-squirrel nests at Duvanny Yar, Kolyma lowland, northeast Siberia. *Quat. Sci. Rev.*
30, 2107–2123 (2011).

27. Murton, J. B. *et al.* Palaeoenvironmental Interpretation of Yedoma Silt (Ice Complex)
Deposition as Cold-Climate Loess, Duvanny Yar, Northeast Siberia: Palaeoenvironmental
Interpretation of Yedoma Silt, Duvanny Yar. *Permafr. Periglac. Process.* **26**, 208–288
(2015).

28. Abramov, A., Vishnivetskaya, T. & Rivkina, E. Are permafrost microorganisms as old as
permafrost? *FEMS Microbiol. Ecol.* **97**, fiaa260 (2021).

29. Gilichinsky, D. A., Wagener, S. & Vishnevetskaya, T. A. Permafrost microbiology.
Permafr. Periglac. Process. **6**, 281–291 (1995).

30. Schiffer, P. H. *et al.* Signatures of the Evolution of Parthenogenesis and Cryptobiosis in the
Genomes of Panagrolaimid Nematodes. *iScience* **21**, 587–602 (2019).

31. Schiffer, P. H. *et al.* Developmental variations among Panagrolaimid nematodes indicate
developmental system drift within a small taxonomic unit. *Dev. Genes Evol.* **224**, 183–188
(2014).

32. Dey, A., Chan, C. K. W., Thomas, C. G. & Cutter, A. D. Molecular hyperdiversity defines
populations of the nematode Caenorhabditis brenneri. *Proc. Natl. Acad. Sci.* **110**, 11056–
11060 (2013).

33. Cutter, A. D. Divergence Times in Caenorhabditis and Drosophila Inferred from Direct
Estimates of the Neutral Mutation Rate. *Mol. Biol. Evol.* **25**, 778–786 (2008).

34. Malavin, S., Shmakova, L., Claverie, J.-M. & Rivkina, E. Frozen Zoo: a collection of
permafrost samples containing viable protists and their viruses. *Biodivers. Data J.* **8**, e51586
(2020).

35. Krivushin, K. *et al.* Two Metagenomes from Late Pleistocene Northeast Siberian
Permafrost. *Genome Announc.* **3**, e01380-14 (2015).

36. Zhou, J. *et al.* Phylogenetic diversity of a bacterial community determined from Siberian
tundra soil DNA. *Microbiology* **143**, 3913–3919 (1997).

37. Kapterev, P. Anabiosis in the conditions of permanent congelation. *Izv Akad Nauk SSSR Ser*
Biol **6**, 1073–1088 (1936).

38. Kapterev, P. New data on revitalization of organisms from perpetually frozen grounds. *CR*
Acad Sci URSS **20**, 315–317 (1938).

39. Andrássy, I. *Klasse Nematoda*. (Gustav Fischer Verlag, 1984).

40. Lewis, S. C. *et al.* Molecular evolution in Panagrolaimus nematodes: origins of
parthenogenesis, hermaphroditism and the Antarctic species P. davidi. *BMC Evol. Biol.* **9**,
15 (2009).

41. Andrássy, I. *Free-living Nematodes of Hungary*. (Hungarian Natural History Museum,
2005).

42. Schulenburg, H. & Félix, M.-A. The Natural Biotic Environment of *Caenorhabditis elegans*.
Genetics **206**, 55–86 (2017).

43. Frézal, L. & Félix, M.-A. C. elegans outside the Petri dish. *eLife* **4**, e05849 (2015).

44. Gems, D. *et al.* Two pleiotropic classes of daf-2 mutation affect larval arrest, adult behavior,
reproduction and longevity in Caenorhabditis elegans. *Genetics* **150**, 129–155 (1998).

45.Vowels, J. J. & Thomas, J. H. Multiple chemosensory defects in daf-11 and daf-21 mutants
of Caenorhabditis elegans. *Genetics* **138**, 303–316 (1994).

46.Ren Peifeng *et al.* Control of C. elegans Larval Development by Neuronal Expression of a
TGF-β Homolog. *Science* **274**, 1389–1391 (1996).

47.Gerisch, B., Weitzel, C., Kober-Eisermann, C., Rottiers, V. & Antebi, A. A Hormonal
Signaling Pathway Influencing C. elegans Metabolism, Reproductive Development, and
Life Span. *Dev. Cell* **1**, 841–851 (2001).

48.Penkov, S. *et al.* Integration of carbohydrate metabolism and redox state controls dauer
larva formation in Caenorhabditis elegans. *Nat. Commun.* **6**, 8060 (2015).

49.Penkov, S. *et al.* A metabolic switch regulates the transition between growth and diapause
in C. elegans. *BMC Biol.* **18**, 31 (2020).

50.Nurk, S. *et al.* HiCanu: accurate assembly of segmental duplications, satellites, and allelic
variants from high-fidelity long reads. *Genome Res.* **30**, 1291–1305 (2020).

51.Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M. & Borodovsky, M. BRAKER2: automatic
eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a
protein database. *NAR Genomics Bioinforma.* **3**, lqaa108 (2021).

52.Smythe, A. B., Holovachov, O. & Kocot, K. M. Improved phylogenomic sampling of free-
living nematodes enhances resolution of higher-level nematode phylogeny. *BMC Evol. Biol.*
19, 121 (2019).

53.Ahmed, M., Adedidran, F. & Holovachov, O. A draft transcriptome of a parasite
Neocamacolaimus parasiticus (Camacolaimidae, Plectida). *J. Nematol.* **53**, 1–4 (2021).

54.Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

55.Goecks, J., Nekrutenko, A., Taylor, J. & Galaxy Team, T. Galaxy: a comprehensive
approach for supporting accessible, reproducible, and transparent computational research in
the life sciences. *Genome Biol.* **11**, R86 (2010).

56.Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the
Trinity platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494–1512 (2013).

57.Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

58.Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

59.Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157

(2015).

60.Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol.* **20**, 238 (2019).

61.Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

62.Capella-Gutierrez, S., Silla-Martinez, J. M. & Gabaldon, T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973

(2009).

63.Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

64.Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*

30, 1236–1240 (2014).

65.Huson, D. H. & Scornavacca, C. Dendroscope 3: An Interactive Tool for Rooted

Phylogenetic Trees and Networks. *Syst. Biol.* **61**, 1061–1067 (2012).

66.Kück, P. & Meusemann, K. FASconCAT: Convenient handling of data matrices. *Mol.*

Phylogenet. Evol. **56**, 1115–1118 (2010).

67.Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol.*

Biol. Evol. **32**, 268–274 (2015).

68.Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**,

587–589 (2017).

69.Zhang, C., Scornavacca, C., Molloy, E. K. & Mirarab, S. ASTRAL-Pro: Quartet-Based

Species-Tree Inference despite Paralogy. *Mol. Biol. Evol.* **37**, 3292–3307 (2020).

70.Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and Smudgeplot

for reference-free profiling of polyploid genomes. *Nat. Commun.* **11**, 1432 (2020).

71.Kokot, M., Długosz, M. & Deorowicz, S. KMC 3: counting and manipulating k-mer

statistics. *Bioinformatics* **33**, 2759–2761 (2017).

Figure legends

Figure 1. Study site.

a) location of the Duvanny Yar outcrope on the Kolyma River, northeastern Siberia, Russia. b) view of the upper part of outcrop composed of ice wedges and permafrost silty deposits. c) lithostratigraphic scheme of deposits, showing location of studied rodent borrow (red circle). d) fossil rodent burrow with herbaceous litter and seeds buried in permafrost deposits; m a.r.l. = meters above river level.

Figure 2. General morphology of *Panagrolaimus n. sp* sp. n., female.

Scanning electron pictures (a, c), light microscopy photographs (e, f) and graphic presentations (b, d, g) of holotype: a, b) entire body, c, d) anterior ends, e) anterior body, f) perivulvar body region, g) tail. Abbreviations: l.f. – lateral field, ov – ovary, pro – procorpus of the pharynx, t.b. – terminal bulb of the pharynx, u – uterus with eggs, v – vulva, v.p. – ventral pore. Scale bars: a, d, e, f, g – 20 μm, b – 100 μm, c – 2 μm.

 Figure 3. Genome assembly and phylogenomics reveals that the newly discovered

***Panagrolaimus n. sp* species is triploid.**

a) Kmer spectra of the *Panagrolaimus n. sp* PacBio HiFi data. Kmers of length 19 were counted

using Jellyfish. b) Circos plot showing the triploid structure of the *Panagrolaimus kolymensis*

genome. The relationship between eight contigs, representing 39.9 Mb (15%) of the genome,

is shown. Numbers represent the contig IDs. c) Inferred species tree for all taxa. The maximum

likelihood tree inferred using a concatenated supermatrix (18S and 28S genes) with bootstrap

support values is displayed. All genera are represented as monophyletic clades. *Panagrolaimus*

n. sp is highlighted in red and basal to all other *Panagrolaimus* taxa. Internal nodes, where all

subsequent branches represent identical sequences, are displayed with a black star. d)

Panagrolaimus n. sp. possesses *C. elegans* gene orthologs to enzymes required for TCA cycle,

glyoxylate shunt, glycolysis, gluconeogenesis, trehalose synthesis, and polyamine synthesis.

Black filled circles: Ortholog presence suggested by orthogroup clustering, phylogenetic

analysis, and domain architecture. White filled circles: No ortholog found via current analysis.

Coloured filled circles: Presence of *Panagrolaimus n. sp.* gene(s), related to several *C. elegans*

genes (all genes of same colour) that are all co-orthologous to that gene (those genes). Label:

C. elegans enzyme names and orthogroup that contains that gene according to our orthogroup

clustering.

Figure 4. *C. elegans* **dauer larvae and** *Panagrolaimus n. sp.* **utilize similar mechanisms to survive cryptobiosis.**

a) Survival rate of *Panagrolaimus n. sp.* nematodes to desiccation and freezing (-80°C). Error bars indicate standard error of mean of two independent experiments with two technical replicates performed on two different days. Statistical comparison was performed using unpaired t test with Welch correction. n.s. $p > 0.05$, ****$p < 0.0001$. For desiccation (non-preconditioned) n = 289, freezing (non-preconditioned) n = 675, desiccation (preconditioned at 98%RH) n = 953 and freezing (preconditioned at 98%RH) n = 1295. b) *Panagrolaimus n. sp.* nematodes and *daf-2(e1370)* dauer larvae upregulate trehalose levels upon preconditioning at 98%RH. Error bars indicate standard error of mean of two independent experiments with three replicates performed on two different days. Statistical comparison was performed using two-way ANOVA with Holm-Sidak's multiple comparison test, ****$p < 0.0001$. c-d) 2D-thin layer chromatography of ^{14}C-acetate labelled metabolites from *Panagrolaimus n. sp.* that were non-preconditioned and preconditioned at 98%RH. Enumerated spots indicate trehalose (1), glucose (2), glutamate (3), glutamine (4), serine/glycine (5) and phenylalanine (6). Representative images from at least two independent experiments performed on two different days. e) Desiccated *daf-2 (e1370)* dauer larvae survive to freezing (-80°C) for an extremely long period. Error bars indicate standard error of mean of two independent experiments with two technical replicates performed on two different days. f) Brood size of desiccated dauer larvae exposed to freezing remain like that non- desiccated dauer larvae. Average brood size is the mean of seven dauer larvae per each condition. Statistical comparison was performed by using non-parametric Kolmogorov-smirnov test n.s $p>0.05$.

Figure S1. Calibration of a radiocarbon (^{14}C) date.

Radiocarbon date (44,315±405 BP) and calibrated age (45,839 -- 47,769 cal BP) of plant

material collected from buried borrow P-1320. Radiocarbon ages were converted to calendar

age equivalents with the OxCal V.4.4 program using the IntCal20 calibration curve. Pink-

shaded area — radiocarbon date with standard deviation; grey-shaded area — radiocarbon date

projection on the calibration curve with 95.4% probability.

956 **Figure S2. Morphology of *Panagrolaimus n. sp* sp. n., female.**

957 Graphic presentations of holotype (a,b) and SEM pictures (c-i): a) anterior body, b) female

958 reproductive branch, c–e) anterior end of three different female specimens, f) anterior part of the

959 lateral ridge, g) vulva, h) ventral excretory/secretory pore, i) posterior body with anus and

960 lateral ridge. Scale bars: a,b - 50 μm, c,i - 3 μm, d - 2 μm, e –1 μm, f,h - 5 μm, g - 10 μm, .

Figure S3. Phylogenies inferred for genes sets using both concatenation and coalescence approaches.

a) The ploidy level was analysed with *Smudgeplot* v0.2.1[70]. KMC version 3.1.0[71] was used to count the 21-mers in the PacBio CCS reads. Then we ran *smudgeplot.py* to determine the lower and upper coverage cut-offs. These were determined to be 14 and 380. 21-mers with a coverage between 14 and 380 were filtered with kmc_tools. Then, we computed the k-mer pairs from the filtered 21-mers by running *smudgeplot.py hetkmers*. Finally, the produced a smudgeplot shows an estimated ploidy of 3. b) Species tree inferred for 12,295 gene trees using a coalescence approach. The species tree implemented using the coalescence approach with orthogroup gene trees is displayed. Novel species in this study are displayed in red. All nodes have a posterior probability of 1. c) Species tree inferred for 102 taxa. The maximum likelihood tree inferred using a concatenated supermatrix of 60 genes is displayed. Bootstrap values are only displayed for nodes with less than 100% support. The platyhelminth *Macrostum lignano* serves as an outgroup for rooting. The ancestral *Panagrolaimus*, sister to all others within the *Panagrolaimus* genus, is highlighted in red.

Figure S4. Combination of cryptobiotic states enhances survival of *C. elegans* dauer
larvae.

a) Desiccated dauer larvae manifest enhanced survival rate to heat stress (34°C). Error bars
indicate standard deviation of two independent experiments with two technical replicates. b)
Desiccated dauer larvae display enhances survival rate to anoxia. Error bars indicate standard
deviation of two independent experiments with two technical replicates. Statistical
comparison was performed by paired two tailed t-test. *$p<0.05$. c) *Panagrolaimus n. sp.*
possesses gene orthologs to most genes implicated in dauer formation and metabolism in *C.*
elegans. Black filled circles: Ortholog presence suggested by orthogroup clustering,
phylogenetic analysis, and domain architecture. White filled circles: No ortholog found via
current analysis (in all cases these *C. elegans* genes did not cluster with any *Panagrolaimus*
genes in the orthogroup clustering). Label: *C. elegans* enzyme names and orthogroup that
contains that gene according to our orthogroup clustering.

Figure S5. *Panagrolaimus n. sp.* **reduces triacylglycerols (TAGs) levels and accumulates trehalose-6-phosphate upon preconditioning at 98%RH**.

a) 1D-Thin layer chromatography of acetate labelled organic fractions of non-preconditioned (1) and preconditioned (2) *P. kolymaensis*. b) Mass spectrometric quantification of TAG levels of non-preconditioned (1) and preconditioned (2) *P. kolymaensis*. Error bars indicate standard deviation of two independent experiments with two technical replicates. Statistical analysis was performed using unpaired t-test with Welch correction **$p < 0.001$. c-d) non-preconditioned and preconditioned mass spectrum of an empty region, e-f) spot 1 (trehalose), g-f) spot 7 (trehalose-6-phosphate) scraped out and extracted from the 2D-TLC.

Figure 1
Shatilovich *et al*

Figure 3
Shatilovich *et al*

Figure 4
Shatilovich *et al*

Figure S4
Shatilovich *et al*

A

TAGs

1-Non-Preconditioned

2-Preconditioned at 98%RH

B

1- Non-Preconditioned

2- Preconditioned at 98%RH

Non-preconditioned

Preconditioned at 98%RH

C Empty Spot

D

E Spot 1(Trehalose)(M-1)

F

G Spot 7 (Trehalose-6-phosphate)(M-1)

H

Figure S5
Shatilovich *et al*