Sensory plasticity in a socially plastic bee

Rebecca A Boulton*[1,2] and Jeremy Field[3]

[1]Laboratory of Genetics, Plant Sciences Group, Wageningen University & Research,

6708 PB, Netherlands

[2]Biological and Environmental Sciences, University of Stirling, FK9 4NF, United

Kingdom

[3] College of Life and Environmental Sciences, University of Exeter, Penryn campus,

TR10 9FE, United Kingdom

*author for correspondence: rebeccaboulton87@googlemail.com

Abstract

The social Hymenoptera have contributed much to our understanding of the

evolution of sensory systems. Attention has focussed chiefly on how sociality and

sensory systems have evolved together. In the Hymenoptera, the antennal sensilla

are important for optimising the perception of olfactory social information. Social

species have denser antennal sensilla than solitary species, which is thought to

enhance social cohesion through nest-mate recognition. In the current study, we test

whether sensilla numbers vary between populations of the socially plastic sweat bee

Halictus rubicundus from regions that vary in climate and the degree to which

sociality is expressed. We found region level differences in both olfactory and

hygro/thermoreceptive sensilla numbers. We also found evidence that olfactory

sensilla density is developmentally plastic: when we transplanted bees from Scotland

to the south-east of England, their offspring (which developed in the south) had more

olfactory hairs than the transplanted individuals themselves (which developed in

Scotland). The transplanted bees displayed a mix of social (a queen plus workers)

and solitary nesting, but neither individual nor nest phenotype was related to sensilla

density. We suggest that this general, rather than caste-specific sensory plasticity

provides a flexible means to optimise sensory perception according to the most

pressing demands of the environment. Sensory plasticity may support social

plasticity in *H. rubicundus* but does not appear to be causally related to it.

Introduction

Natural selection on sensory abilities has resulted in enormous diversity in the

mechanisms that organisms use to perceive their environment. For example, species

of Lake Tanganyika cichlids that have colonised more complex, rocky habitats have

greater visual acuity than sand-dwelling species (Dobberfuhl et al. 2005). These

mechanisms can also feedback into selective regimes, facilitating micro- and

macroevolutionary change through sensory drive. Sensory drive occurs when local

adaptation results in populations with diverging modes of communication and

perception. This has been proposed to lead to assortative mating between partners

with complementary signalling and perception systems, reducing geneflow and

promoting speciation (Boughman 2002).

The Hymenoptera have emerged as a model system to understand both the

proximate and ultimate factors that underlie sensory evolution. These insects have

antennal receptors with diverse functions and distinct forms that can easily be

characterised within and between species. This work has revealed that both the

density and form of antennal sense organs (sensilla) correlate with diet (Polidori et

al. 2012; 2020), sex (Babu 2019; Do Carmo Queiroz Fialho et al. 2014) and a

cleptoparasitic lifestyle (Wcislo 1995; Galvani et al. 2017). Attention has focussed

particularly on how eusociality has imposed selection to optimise antennal

perception. Eusociality involves reproductive division of labour with a queen/s

producing eggs and workers provisioning the brood. However, sociality takes diverse

forms across the bees and wasps and group sizes vary enormously, providing rich

opportunities for comparative study. Many obligately social species show worker

polyphenism, where some workers specialise on brood care and others on foraging

or guarding the nest (Andersson 1984). Compared with solitary species, social

species require more finely tuned communication systems to distinguish intruders

from kin and gain information from nestmates (Wcislo 1997; Renner and Nieh 2008;

Freeberg et al., 2012; Peckre et al., 2019). Across the Hymenoptera, social species

tend to have greater densities of antennal sensilla that are involved in the perception

of olfactory cues than solitary species (Elgar et al. 2018). This is thought to enhance

social cohesion and performance by facilitating nest-mate recognition (Ozaki et al.

2005; Gill et al. 2013; Couto et al. 2017) and supporting caste polyphenisms

(Spaethe et al. 2007; Riveros and Gronenberg 2010; Gill et al. 2013; Grüter et al.

2017; Elgar et al. 2018) without requiring the production of costly and complex

chemical signals (Kather & Martin 2015).

The olfactory antennal sensilla have been proposed as a pre-adaptation that acts

alongside haplodiploidy to predispose the Hymenoptera to the evolution of sociality

(Couto et al. 2017). A conserved, ancestral olfactory subsystem, involving basiconic

antennal sensilla that detect chemical information from nestmates and neural circuits

that process it, has been suggested to facilitate the evolution of sociality in the

Hymenoptera. This works by allowing nestmate recognition from cuticular

hydrocarbon (CHCs) profiles (Ozaki et al. 2005; McKenzie et al. 2016; Couto et al.

2017; Pask et al. 2017 see also Kather et al., 2016). While empirical evidence for

this subsystem exists for only two species (an ant and a hornet), the halictid bees

have provided support for the possibility that sensory pre-adaptations might

contribute to the evolution of sociality. The Halictidae or sweat bees exhibit a range

of social structures, from solitary nesting through to obligate sociality, including

within-species social polymorphism and social plasticity (Gibbs et al. 2012). A

comparative study by Wittwer et al (2017) has shown that ancestrally solitary species

and their social sister species have equivalent densities of hair-like sensilla

(basiconic and trichoid sensilla) that detect olfactory cues. Species that have

reverted to a solitary existence from sociality, on the other hand, have reduced

densities of these sensilla. This suggests that in the Halictidae, sensitivity to olfactory

cues precedes the evolution of sociality rather than evolving as a consequence of it.

In addition to caste polyphenisms, where individuals always form social groups but

vary in the tasks that they perform, a number of species are facultatively social. In

these species, individuals vary in their propensity to form social groups or nest as

solitary individuals (Gibbs et al. 2012). This social plasticity is thought to be key to

the evolutionary diversification of the Hymenoptera as it provides a genetic base

from which obligate eusociality and caste polyphenism can evolve repeatedly (West-

Eberhard 2003; Rehan & Toth, 2015; Jones et al. 2017; Shell & Rehan, 2018). It also

provides an opportunity to test for correlates of social evolution without the

complication of interspecific differences (see also Schradin 2013; Schradin et al.,

2018). In this study we characterise the antennal sensilla of a facultatively social

species, *Halictus rubicundus,* for the first time. This species shows regional

differences in social behaviour which are not fixed and depend on the climate (see

methods below; Field et al., 2010; 2012). Phylogenetic work suggests that social

plasticity in *H. rubicundus* has evolved from an ancestral state of obligate sociality in

the genus *Halictus* (Danforth 2002).

We measure the density of different antennal sensilla types between populations

which vary in their expression of sociality due to the climate and test whether

sensory plasticity mirrors social plasticity in a transplant experiment with *H.*

rubicundus. Our results represent an important step in disentangling the sequence of

events that leads to the evolution of eusociality and reproductive division of labour,

as well as the evolutionary loss of these traits in the Hymenoptera.

Materials and methods

Study system

Halictus rubicundus is a ground-nesting bee that can be found throughout Europe

and North America in areas with sandy-loamy soil that receive sufficient sunlight. In

Great Britain and Ireland, *H. rubicundus* demonstrates social variation according to

climate (latitude and altitude) but retains social plasticity throughout its range (Field

et al. 2010; 2012). In the north and at high altitudes bees are typically solitary as the

short growing season and cooler climate precludes the bivoltinism that is required for

social living. In a typical year in colder climates, foundresses produce a single brood

of offspring per year, the females of which enter hibernation and emerge to become

foundresses the following spring (figure 1a). In areas further south and at low

altitudes growing seasons tend to be longer and the weather is warmer, so

bivoltinism and sociality can occur more frequently. After overwintering underground,

females produce a first brood (the B1) which become workers and provision a

second brood (B2) which are offspring of either the original foundress or a B1

replacement queen. B1 females are also observed to provision their nests alone and

lay eggs (without workers, i.e. become solitary foundresses) in the same year that

they emerge. The B2 hibernate and emerge the following year to found their own

nests (figure 1b; Field et al. 2010; 2012).

Sociality is extremely plastic in *H. rubicundus* in Great Britain, and when solitary

northern bees are transplanted to the south, the B1 brood will often provision as

workers or solitary foundresses in the same year (Field et al. 2010; 2012; this study).

Here we test for fixed and plastic differences in antennal sensilla density in *H.*

rubicundus across and within populations. We first test whether olfactory sensilla

densities differ between bees from southern populations, where the climate allows

some individuals to form social nests most years, and northern populations with

limited opportunity to form social nests. Second, we test whether antennal sensilla

numbers are developmentally plastic by transplanting bees from the north to the

south. Do the offspring of northern bees that develop in the warmer south have

different numbers of antennal sensilla than their parental generation which

developed in the north? We chose to conduct a north-to-south transplant because

we wanted to test for differences in sensilla number between individual and nest-

level phenotypes, to determine whether social individuals/nests exhibit greater

densities of olfactory sensilla to aid in nestmate recognition. This comparison is

possible only when bees were able to express their full range of social behaviour (i.e.

in the warmer south).

Collection

Halictus rubicundus females were collected from 4 sites in the UK in 2018, 2019 and

2020 (see fig 1c). Two sites in the south-west (SW): Boscastle (Cornwall: N 50° 41'

24" W 4° 41' 24"; N = 12; 2018) and Bodmin (Cornwall, N 50° 30' 36" W 4° 33' 36"; N

= 4; 2019); a site at a mid-latitude (MID): Belfast (Northern Ireland, N 54° 32' 24" W

5° 58' 48"; N = 19; 2020); and a site in the far north of Scotland (SCO): Migdale

(Scotland, N 57° 53' 24" W 4° 15' 0"; N = 15; 2018 and 2020). Previous studies and

our own observations indicate that while a proportion of nests are social in SW (this

study), nests are always non-social in Belfast (MID; Field et al. 2010) and Scotland

(SCO; this study). However, bees from Belfast (MID; Field et al. 2010) and Scotland

(SCO; this study) are socially plastic and may have social nests when moved to

more southerly sites. Bees sampled were expected to be foundresses based on

collection date. We collected females by hand-netting in late June in SCO, and in

May and early June in the SW, when all bees had worn wings indicative of several

weeks of provisioning. In Belfast (MID) we excavated overwintering foundresses in

February.

Transplant

In addition to sampling from native populations, we collected bees that had been

transplanted from Scotland (SCO) to a site in the south-east (SE), the Knepp estate

in West Sussex (N 50° 53' 60" W 0° 21' 36"; N = 47; see fig 1c). We refer to these

transplanted bees as 'T' bees hereafter. Buckets of soil were embedded within a

nest aggregation at the Scottish site during winter 2018-19, and native foundresses

subsequently nested in them during 2019. Their offspring (T) then emerged in late

summer 2019 and hibernated in the buckets. The buckets containing hibernating T

bees were taken from Scotland to the south-east in the spring of 2020, where they

were re-embedded in the ground. The T foundresses were marked with a dot of

enamel paint when they provisioned a B1 brood in the south-east. When the B1

emerged, they were marked with a different colour of enamel paint. Each nest was

marked with a numbered nail at the entrance. This allowed us to distinguish fresh B1

females from the original foundresses and to determine whether nests were solitary

(with a single B1 female or original foundress provisioning alone) or social (with

multiple B1 females provisioning the nest). At the end of the B1 provisioning phase

we collected T foundresses (which had emerged in Scotland and been transplanted

to the SE) and their B1 offspring (which had developed and emerged in the SE; we

refer to these as 'B1' individuals).

Specimen storage and preparation

All specimens were stored in 95% Ethanol until they were prepared for imaging. One

antenna was removed from each bee (whether it was the left or right was recorded)

and mounted on a JEOL aluminium stub (10mm dia x 10mm high) using a PELCO

carbon conductive sticky tab (10mm dia). Specimens were mounted dorsal side up.

Specimens were carbon coated using spectrograpically pure carbon.

Imaging

Mounted and carbon coated specimens were imaged using a TESCAN Vega SEM in

High Vacuum mode at 20.0kV. The scan mode was set to resolution and the

magnification was 485x. Only the two distal segments (11 and 12), which have the

highest density of sensilla, were imaged. The length and area of each antennal

segment was measured during imaging and images of each antennal segment from

the SEM were saved in TIF format.

Figure 1 (A) univoltine lifecycle of *Halictus rubicundus* which prohibits social nesting (B) bivoltine lifecycle of *Halictus rubicundus*

where sociality is possible as the 1st brood (B1) can help to provision a second brood (B2). Note that in (B) the B1 females can

either (i) overwinter and provision the following year as a solitary foundress (ii) provision a nest alone as a solitary foundress in the

year that they emerge or (iii) provision the nest socially with their B1 sisters with a queen (their mother or a sister) in the year that

they emerge. (C) Map depicting the sites where *H. rubicundus* females were studied. Dashed arrow represents the direction of the

transplant experiment (SCO to SE).

Image scoring

Eight distinct types of sensilla with overlapping functions have been characterised in the Hymenoptera (Do Carmo Queiroz Fialho et al. 2014). Placoid sensilla are plate-like structures that are involved in olfaction and chemoreception. Tricoid and basiconic sensilla are hair-like projections with multiple sub-types that are used in olfaction involving contact. Coeloconic, campaniform and ampulliform sensilla are pore-like and are involved in sensing temperature, humidity and CO_2 concentration (Do Carmo Queiroz Fialho et al. 2014). For this study we grouped all sensilla types into three structural/functional groups that allowed for repeatability in scoring: (i) olfactory plate sensilla (sensilla placodea, fig 2), (ii) olfactory hair-type sensilla (sensilla trichodea and basoniconica; fig 2) and (iii) temperature/humidity/CO_2 pore sensilla (coeloconic/campaniform/ampulliform sensilla; fig 2).

Images of antennal segments 11 and 12 were scored using ImageJ software. For each segment, three 50μM x 50μM 'quadrats' were selected, the X and Y coordinates of which were derived by generating a random number using the excel function RANDBETWEEN. Each quadrat was saved as a TIF image and the number of sensilla types i and ii were counted. We counted sensilla type iii across the entire segment as they are distributed unevenly on the surface of the antenna. In some cases, bare patches with no sensilla were present on the surface of the antenna. Any quadrats which fell on the bare areas were discarded and another area was selected. We note that the scorer was not blind to the population of origin as the same person was required to mount, image and score the antennae.

Figure 2 legend labels:
(i) Olfactory plates (placoid sensillae)
(ii) Olfactory hairs (basiconic/trichoid sensillae)
(iii) Temperature/humidity and CO_2 sensitive pores (coeloconic, campaniform and ampulliform sensillae)

Figure 2 SEM image showing a *Halictus rubicundus* antenna tip. (A) shows the distal 2 antennal segments (11 and 12) that were imaged, (B) and (C) show the types (i, ii, and iii) of antennal sensilla that were counted. Scale is given in μM.

Analysis

Repeatability

Thirty-three of the 130 sampled antennae were scored a second time (by the same person) for all sensilla types. The repeatability of image scoring was assessed using the intra-class correlation coefficient, function *ICC* in the R package *irr* (Gamer et al, 2019).

Sensilla numbers

We used linear mixed models in the R package lme4 (Bates et al. 2019) to test for differences in the numbers of sensilla types (i-iii). For all models outlined below, the response variable was total sensilla counted, either i) olfactory plates, total across 3 quadrats for each segment per individual; ii) olfactory hairs, total across 3 quadrats for each segment per individual; or iii) temperature/humidity pores, total across each

segment per individual. We consider the unit of replication (N) to be a single bee. We

did not measure more than one antenna from each bee, but we did measure 2

segments per antenna. As such, Individual was fitted as a random effect in all models.

The package DHARMa was used to determine the most appropriate error structure for

all models (Gaussian) and to interpret residual plots for lack of fit (Hartig, 2020).

Region, development site and social phenotype

To test whether bees from different regions have different numbers of sensilla we

included region as a fixed factor with 3 levels in LMMs: SW (Boscastle and Bodmin

combined; N = 16 individuals), MID (Belfast, Northern Ireland; N = 19 individuals)

and SCO (Migdale, Scotland, far north; N = 15 individuals).

We ran additional models using only data from bees in the transplant experiment to

test whether sensilla density (type i, ii or iii) varies depending on where bees

completed their development (which may suggest developmental plasticity). To do

this we compared bees that developed in Scotland ('T' foundresses) with their

offspring that developed in the south-east (B1 females) after transplantation. LMMs

included a fixed effect of development site, with 2 levels: SCO (bees that emerged in

Scotland and were either collected there (N = 15 individuals, same individuals that

were used in the region models; or 'T' females that were moved to the south-east as

adults, N = 4) and SE (bees that emerged as adults after immature development in

the south-east, the B1 offspring of the bees that were transplanted from Scotland to

the south-east; N = 27). Note that *H. rubicundus* females mate before overwintering

and so both the mother and father of the B1 bees collected in the south-east

originated in Scotland.

Additional LMMs were used to test whether B1 offspring collected in the SE (using

the same individuals used in the *development site* model for which nest phenotypes

were known, N = 14) had higher counts of sensilla if they were workers (hereafter

B1w) from social nests of transplanted females compared to those which left and

started their own solitary nests (hereafter B1sol). Social nests had a queen and one

or more workers (B1w), while solitary nests had a single B1sol foundress that

produced offspring in the same year. For these models we included nest phenotype

as a fixed effect with 2 levels (social, N = 6 or solitary, N = 8). We did not sample

multiple individuals from the same nest, so did not include nest as a random effect.

We did not recover any queens from social nests and so could not test for an effect

of bee phenotype within social nests. We also ran a model to test for effects of

individual, rather than nest-level, phenotype across social and solitary nests, using

the same bees. For this analysis bees were characterised as (1) a future

reproductive, B1hib (a B1 individual that had emerged in 2020 and did not provision

that year, will hibernate and provision in 2021; N = 13); (2) a worker, B1w (a B1

individual that provisioned a nest containing a queen; N = 7); (3) a solitary foundress

from 2019, T (a foundress that emerged in SCO in 2019, was transplanted to the

south-east and laid eggs/provisioned a nest alone without workers); (4) a solitary

foundress from 2020, B1sol (a foundress that emerged in the B1 generation in 2020

in the SE and provisioned a nest alone; N = 4).

Finally, we ran a model using the bees collected in the SE to test whether age-

related wear and tear reduces density of all 3 sensilla types. For these analyses age

was binary: bees were scored as fresh (newly emerged and had not provisioned a

nest; N = 12) or old (had provisioned a nest for several weeks N = 16).

All LMMs included antenna segment number (11 or 12) as a fixed factor. The

interaction effect between region/development type/social phenotype and segment

number was also fitted. We included a random effect of individual as two segments

were imaged for each bee. Models were run using the function *lmer* in the R

package lme4 (Bates et al, 2015).

Results

Repeatability, differences between antennal segments and effects of age

Counts of all sensilla types were highly repeatable (Koo and Li 2016; ICC greater

than 0.85 for all; for a detailed summary see supplementary material table S1).

Olfactory plates (type i) and olfactory hairs (type ii) were found in greater numbers on

the most distal antennal segment, segment 12, while hygro/thermoreceptors (type iii)

were found in higher numbers on segment 11.

We also found no evidence that age-related wear and tear influences the variation in

sensilla density; there was no effect of age on density of any of the three types of

sensilla (type i: X^2 = 0.25, df = 1, p = 0.62; type ii: X^2 = 0.01, df = 1, p = 0.92; type iii:

X^2 = 0.09, df = 1, p = 0.76).

Region

There was a significant effect of region on the counts of type ii (olfactory hairs) and iii

(thermo/hygroreceptors) sensilla, but not type i (olfactory plate receptors; see table

1, figure 3a-c). Pairwise tests show that bees from mid-latitude (MID; Belfast; N = 19)

had significantly more olfactory hairs (type ii) than bees collected from the far north

(Scotland; N=30; p = 0.03; figure 3b), whereas bees from the south-west (N = 16) did

not differ from MID (N = 32; p = 0.06) or Scottish (N = 15; p = 0.93) bees. Bees from

SCO and MID had more thermo/hygro (type iii) receptors than SW bees (p < 0.05;

figure 3c). Bees from the south-west had the most olfactory plate sensilla (type i) but

this was not statistically significant (figure 3a). There were no interaction effects

between segment number and geographic region (table 1).

Table 1 LMM results showing effect of region on antennal sensilla counts across

segments using type II sums of squares (p-values for main effects are calculated

independent from interaction effects which in all cases are non-significant).

Sensilla type	Effect	X^2	df	P
Type i) olfactory plates	Region	2.31	2	0.31
	Segment	19.48	1	1.01×10^{-5}
	Region*Segment	0.45	2	0.80
Type ii) olfactory hairs	Region	7.81	2	0.02
	Segment	41.74	1	1.04×10^{-10}
	Region*Segment	2.70	2	0.26
Type iii) thermo/hygro-receptive	Region	9.14	2	0.01
	Segment	19.17	1	1.20×10^{-5}

Region*Segment	0.35	2	0.84

(a)

(b)

(c)

(d)

(e)

(f)

Figure 3 Numbers of (a) and (d) olfactory plates, i (top row); (b) and (e) olfactory

hairs, ii (middle row); and (c) and (f) hygro/thermoreceptive pores, iii (bottom row),

for *Halictus rubicundus* across: a-c three sampled regions: SW (South-west; mostly

social), MID (Belfast; solitary) and SCO (Scotland; solitary); d-e for *Halictus*

rubicundus originating from Scotland and collected after either developing as larvae

in Scotland (SCO) or after developing as larvae at the transplanted site in the south-

east (SE). Different uppercase letters (A or B) over bars indicate significantly

different counts of sensilla. Black points represent sensilla counts on segment 11

and grey points on segment 12. Box plots show median and quartiles across

segments 11 and 12.

Development site

The density of olfactory hairs (type ii) varied according to where bees originating

from Scotland spent their immature development (table 2; figure 3d-e). Offspring

(B1) of transplanted bees (T) which developed in the south-east (N=27) had more

olfactory hairs (type ii, $p = 0.01$) than conspecifics that developed in their natal

Scottish site (N=19; fig 3e). A similar trend was seen in olfactory plates (I; fig 3d) but

this was not statistically significant ($p = 0.06$). There was no effect of developmental

site on thermo/hygroreceptor numbers (iii; fig 3f).

Table 2 LMM results showing effect of developmental site on antennal sensilla counts across segments (p-values for main effects

are calculated independent from interaction effects which in all cases are non-significant).

Sensilla type	Effect	X^2	df	P
Type i) olfactory plates	Development site	3.44	1	0.06
	Segment	34.82	1	3.6×10^{-9}
	Development site *Segment	0.61	1	0.43
Type ii) olfactory hairs	Development site	6.41	1	0.01
	Segment	61.35	1	4.79×10^{-15}
	Development site *Segment	1.57	1	0.21
Type iii) thermo/hygroreceptive	Development site	0.74	1	0.39
	Segment	33.25	1	8.12×10^{-9}
	Development site *Segment	0.15	1	0.69

Sociality

B1 workers (B1w) from social nests (N= 6) and solitary B1 (B1sol) females that founded new nests alone (N= 8) had equivalent numbers of all 3 sensilla types (i-iii, see table 3; fig 4a-c) and there were no interactions between nest phenotype and antenna segment number. There was some suggestion that solitary bees had higher numbers of olfactory plates (type i), but this was not statistically significant (p = 0.07). Similarly there were no differences in numbers of any sensilla types across different individual phenotypes (B1hib future reproductive emerged in 2020 in SE and did not provision that year, will hibernate and provision in 2021, N = 13; B1w worker emerged in 2020 in the SE, N = 7; T solitary foundress that emerged 2019 in SCO and provisioned in the SE in 2020, N = 4; and B1sol solitary foundress emerged and provisioned in 2020 in SE, N = 4; see table 4; figure 4d-e). These results were the same if B1 solitary foundresses from 2019 (that developed and emerged in SCO) were excluded (see archived code and data; https://github.com/DrBecky-B/Bee.Antennae).

Table 3 LMM results showing effect of nest phenotype (social/solitary) on antennal sensilla counts across segments (p-values for main effects are calculated independent from interaction effects which in all cases are non-significant).

Sensilla type	Effect	X²	df	P
Type i) olfactory plates	Nest phenotype	2.91	1	0.08
	Segment	11.80	1	0.0005
	Nest phenotype*Segment	1.08	1	0.30

Type ii) olfactory hairs	Nest phenotype	0.08	1	0.78
	Segment	42.03	1	8.98×10^{-11}
	Nest phenotype*Segment	1.29	1	0.26
Type iii) thermo/hygroreceptive	Nest phenotype	0.29	1	0.59
	Segment	33.08	1	8.86×10^{-9}
	Nest phenotype*Segment	0.10	1	0.75

394 **Table 4** LMM results showing effect of bee phenotype (future

395 reproductive/worker/solitary foundress 2019 and solitary foundress 2020) on

396 antennal sensilla counts across segments (p-values for main effects are calculated

397 independent from interaction effects which in all cases are non-significant).

Sensilla type	Effect	X^2	df	P
Type i) olfactory plates	Bee phenotype	1.51	3	0.67
	Segment	22.34	1	2.29×10^{-6}
	Bee phenotype*Segment	1.81	3	0.61
Type ii) olfactory hairs	Bee phenotype	1.24	3	0.74
	Segment	42.03	1	8.33×10^{-11}
	Bee phenotype*Segment	0.85	3	0.84
Type iii) thermo/hygroreceptive	Bee phenotype	3.39	3	0.34
	Segment	30.18	1	3.95×10^{-8}
	Bee phenotype*Segment	2.26	3	0.52

Figure 4 Numbers of (a) and (d) olfactory plates (top row); (b) and (e) olfactory hairs

(middle row); and (c) and (f) hygro/thermoreceptive pores, (bottom row); for *Halictus*

rubicundus from Scotland collected from (a-c) social and solitary nests and

according to individual phenotype (d-f) after being transplanted to the south-east in

2020. For figures 4d-4f: B1hib female that emerged in the SE in 2020 but did not

provision that year (will hibernate and provision in 2021); B1sol female that emerged

in the SE in 2020 and provisioned a nest as a solitary foundress; T female that

emerged in SCO in 2019, was transplanted to the south east and provisioned a nest

as a solitary foundress in 2020; B1w female that emerged in the SE in 2020 and

provisioned a social nest in the same year as a worker. Black points represent

sensilla counts on segment 11 and grey points on segment 12. Box plots show

median and quartiles across segments 11 and 12. There were no significant

differences in counts of any sensilla type across the groups.

Discussion

The social Hymenoptera have contributed much to our understanding of the

proximate and ultimate factors that underlie the evolution of sensory systems.

Attention has focussed chiefly on how sociality has imposed selection to optimise

olfactory communication, enhancing social cohesion through nest-mate recognition

and supporting caste polyphenisms (Wcislo 1997; Ozaki et al. 2005; Spaethe et al.

2007; Renner and Nieh 2008; Riveros and Gronenberg 2010; Gill et al. 2013; Couto

et al. 2017; Grüter et al. 2017; Elgar et al. 2018). In the current study we measured

sensilla density in the socially plastic sweat bee *Halictus rubicundus*. We found that

when bees were transplanted from Scotland, where they are solitary, to the south

where they can be social, their offspring had higher densities of olfactory hairs.

These results suggest that the density of olfactory hairs may, like sociality, be

phenotypically plastic in *H. rubicundus*.

We also found evidence for between-region differences in the density of

basiconic/trichoid olfactory hairs (and hygro/thermoreceptive sensilla) for

untransplanted bees, with mid-latitude bees having higher densities than more

northerly Scottish bees. Differences were not perfectly correlated with the expected

degree of sociality, however, because bees from the warm south-west, which is

where we expect sociality to be the dominant strategy did not differ from mid-latitude

or Scottish bees. This contrasts with the results of Wittwer et al (2017) who showed

that in another halictid, *Lasioglossum albipes,* females from social populations have

higher densities of olfactory hairs than conspecifics from solitary populations. Unlike

H. rubicundus, this species is not socially plastic, and fixed differences in social

structure between populations may have resulted in canalising selection on antennal

sensilla density in *L. albipes* but not *H. rubicundus.*

The results from our transplant experiment also suggest that in *H. rubicundus*

olfactory hair density is not canalised and may be phenotypically plastic. Our results

suggest that olfactory hair density may vary depending on the conditions that bees

experience during development. When Scottish bees were transplanted to the south-

east, their offspring that developed there (where air and soil temperatures in June

and July are 2-5°C warmer than their native site; table S2) had greater densities of

trichoid/basiconic hairs. It may be that in *H. rubicunudus* warmer temperatures

during development*,* rather than social phenotype, lead to greater densities of

olfactory hairs. A similar pattern is seen in the parasitoid wasp *Trichogramma*; when

males develop under colder temperatures they have reduced numbers of basiconic

hairs (Pinto et al. 1989). A direct response to temperature might explain why

untransplanted foundresses from the south-west, where bees are often social, but

temperatures are not as high as in the south-east, did not have higher basiconic hair

densities than non-social populations (Fig. 3).

In addition to plasticity within a population, we found evidence that olfactory hair

density (type ii, basiconic/trichoid sensilla) varies across regional populations in *H.*

rubicundus. In contrast we saw no differences in the densities of placoid (plate-like,

type i) olfactory sensilla across regions. This may relate to the function of these

receptor types. Basiconic/trichoid hairs respond to contact with the CHCs of other

individuals, and so are likely to be required for nest-mate recognition and

communication, while placoid plates are thought to be involved in longer range

olfaction to detect food and hosts (Ozaki et al. 2005; McKenzie et al. 2016; Couto et

al. 2017; Pask et al. 2017). While food availability almost certainly varies across

regions we know of no consistent regional variation in diet breadth in this species

(although this remains to be rigorously studied), which fits with the pattern we see

here.

Bees collected from Belfast at a mid-latitude had the highest density of olfactory hair-

like sensilla (basiconic/trichoid; type ii), higher than bees from Scotland (450km north

of Belfast) but statistically equivalent to bees collected in the south-west. We

predicted that *H. rubicundus* collected further south would have the greatest

densities of olfactory sensilla to support social communication between nestmates

(Wittwer et al. 2017; Elgar et al. 2018). In the south, summers are typically warmer

and longer and a higher proportion of the population is expected to exhibit social

behaviour more regularly. While we did see regional differences in olfactory sensilla

density, bees from the south-west did not have the highest densities as we would

have predicted based on clines in sociality. The regional differences we saw may be

related to developmental, potentially temperature-dependent plasticity, which could

limit the scope for fixed population level differences to evolve in the predicted

direction. Indeed, data from nearby weather stations lends support to the possibility

that warmer temperatures during development might lead to higher densities of

olfactory hair-like (basiconic) sensillae (see above; Pinto et al. 1989). The maximum

and average air temperatures when the bees collected in Belfast were developing (in

June and July) was up to 1°C higher than in the south-west and 1-2 °C higher than in

Scotland (although the minimum temperatures in Belfast tend to be lower than in the

south-west; see supplementary table S2 for complete climate data). Longitudinal

studies of these same populations, reciprocal transplant experiments (from south to

north) and more accurate measurements of soil temperature would help to elucidate

the extent to which genetically fixed differences in olfactory hair density and plastic,

temperature-dependent expression contribute to the population level differences we

saw in this trait.

We also found that the density of sensilla (type iii) involved in the perception of

humidity, temperature and CO_2 varies across regions, but unlike olfactory hairs this

does not appear to be plastic. Bees from a mid-latitude and the north of Scotland

(Belfast and Scotland) have more thermo/hygro receptive sensilla than bees from the

south-west (400 km south of Belfast and 850 km south of the Scottish site). The

density of these sensilla does not appear to be plastic, however, as it was not

influenced by where Scottish bees developed or their social phenotype. This pattern

may be a result of a more extreme, variable climate in the north, which leads to

consistent selection for higher densities of thermo/hygro receptors. Sweat bees

including *H. rubicundus* are highly sensitive to temperature and rain. Flight activity is

constrained by low ambient temperatures, lack of sunlight and rainfall (Potts and

Wilmer 1997). In the north, bees experience colder ambient temperatures and

greater rainfall. Bees with more of these receptors may be more sensitive to current

and oncoming weather conditions, so that they have reduced mortality and improved

foraging based on the climate. The potentially severe fitness consequences of

incorrect perception of climatic cues may explain the lack of plasticity in the number

of thermo/hygro receptors if selection has a strong and canalising effect on this trait.

Our results add a new dimension to the growing body of evidence that

developmental temperature may contribute to adaptations which support sociality in

the Hymenoptera. While previous studies suggest that temperatures experienced by

developing larvae contribute to individual differences in social phenotype (i.e. caste

polyphenism: Czekońska and Tofilski 2020; Becher et al. 2009; the development of

status badges: Green et al. 2012; memory formation: Jones et al. 2005 and olfactory

learning: Anton and Rossler 2021), sensilla density in the socially plastic *H.*

rubicundus may be a more general response to the environment that is not directly

related to the social phenotype of the nest or individual (i.e. a worker, solitary

foundress or queen). Scottish bees that developed in the south-east had more

olfactory hairs than their conspecifics that emerged in Scotland irrespective of their

caste or nest phenotype.

In the Halictidae, Wittwer et al (2017) found that halictid bee species that had

reverted back to solitary nesting from a state of sociality had reduced olfactory hair

density compared to social species and ancestrally solitary species. They suggest

that this is because dense olfactory sensilla are a pre-adaptation that facilitates the

evolution of sociality and may contribute to the repeated evolutionary transitions to

sociality seen in the halictid bees. Our results expand on this, suggesting that

olfactory hair density may also be phenotypically plastic. More broadly, plasticity in

traits such as this might contribute to the evolutionary lability of sociality in the

Halictidae and in the Hymenoptera, acting alongside social plasticity in the

hymenopteran ancestor to provide the 'flexible stem' which allowed for repeated

evolutionary transitions to obligate sociality across the order (West-Eberhard 2003;

Wittwer et al. 2017).

Acknowledgements

Many thanks to Gavyn Rollinson and Sharon Uren for SEM training and technical

support. This work is part of a project that received funding from the European

Research Council (ERC) under the European Horizon's 202 research and innovation

programme (grant agreement No. 695744).

Conflict of interest

The authors have no conflict of interests to report.

Ethics

This research adheres to ethical codes of practice at the University of Exeter. No

vertebrates were used in this research. Bee antenna samples were taken from living

and dead stored specimens. Handling during antenna removal of live specimens was

done as quickly as possible to minimize stress to individuals. Bees frequently lose

parts of and whole antennae in nature and we did not see any negative effects on

the survival or provisioning behaviour of live sampled individuals compared to intact

bees. The sample sizes reported were considered appropriate to maximize statistical

power while reducing the number of individuals involved in the experiments.

Individuals were sampled from large populations > 500 individuals and numbers

taken did not pose a threat to the persistence of these populations. The transplant

experiment did risk causing gene flow (due to escape/mating) between Scottish and

native bees in the south east (SE). We chose a transplant site with no records of *H.*

rubicundus to minimize this risk. We have never found any *H. rubicundus* nesting

nearby and the soil in the SE, which is dense clay, is unsuitable for nesting or

hibernation of this species. At the end of the transplant experiment any surviving

bees were taken back to their natal site in Scotland whilst they were hibernating in

the buckets. We did not see any *H. rubicundus* at the site when we returned in the

year following the transplant (2021). As such we believe that the risk of us causing

gene flow between *H. rubicundus* populations is negligible.

Data accessibility

All code and data are archived on Github and can be accessed here:

https://github.com/DrBecky-B/Bee.Antennae

References

Andersson, M. 1984. The evolution of eusociality. Annual review of Ecology,

Evolution and Systematics 15:165–189.

Anton, S., and W. Rössler. 2021. Plasticity and modulation of olfactory circuits in

insects. Cell and Tissue Research 383:149–164.

Babu, M. 2019. Antennal sensilla of the weaver ant *Oecophylla smaragdina* (F.) –

males and females sense differently. Indian Journal of Entomology 81:418-

422.

Bates, D., Maechler, M., Bolker, B and W. Walker. 2015. Fitting Linear Fixed-Effects

Models using lme4. Journal of statistical software 67:1-48

Becher, M. A., H. Scharpenberg, and R. F. A. Moritz. 2009. Pupal developmental

temperature and behavioral specialization of honeybee workers (Apis

mellifera L.). Journal of Comparative Physiology A: Neuroethology, Sensory,

Neural, and Behavioral Physiology 195:673–679.

Boughman, J. W. 2002. How sensory drive can promote speciation. Trends in

Ecology and Evolution 17:571–577.

Charnov, E. L. 1978. Evolution of eusocial behavior: Offspring choice or parental

parasitism? Journal of Theoretical Biology 75:451–465.

Couto, A., A. Mitra, D. Thiéry, F. Marion-Poll, and J. C. Sandoz. 2017. Hornets have

it: A conserved olfactory subsystem for social recognition in hymenoptera?

Frontiers in Neuroanatomy 11:1–12.

Czekońska, K., and A. Tofilski. 2020. Body mass of honey bee drones developing in

constant and in changing temperatures. Apidologie 51:510–518.

Dobberfuhl, A. P., J. F. P. Ullmann, and C. A. Shumway. 2005. Visual acuity,

environmental complexity, and social organization in African cichlid fishes.

Behavioral Neuroscience 119:1648–1655.

Danforth, B. N. 2002. Evolution of sociality in a primitively eusocial lineage of bees.

Proceedings of the National Academy of Sciences 99: 286-290.

Do Carmo Queiroz Fialho, M., C. P. Guss-Matiello, J. C. Zanuncio, L. A. Oliveira

Campos, and J. E. Serrão. 2014. A comparative study of the antennal sensilla

in corbiculate bees. Journal of Apicultural Research 53:392–403.

Elgar, M. A., D. Zhang, Q. Wang, B. Wittwer, H. T. Pham, T. L. Johnson, C. B.

Freelance, et al. 2018. Insect antennal morphology: The evolution of diverse

609   solutions to odorant perception. Yale Journal of Biology and Medicine 91:457–

610   469.

611 Field, J., R. J. Paxton, A. Soro, and C. Bridge. 2010. Cryptic plasticity underlies a

612   major evolutionary transition. Current Biology 20:2028–2031.

613 Field, J., R. J. Paxton, A. Soro, P. Craze, and C. Bridge. 2012. Body size,

614   demography and foraging in a socially plastic sweat bee: A common garden

615   experiment. Behavioral Ecology and Sociobiology 66:743–756.

616 Freeberg, T. M., R. I. Dunbar, and T. J. Ord. 2012. Social complexity as a proximate

617   and ultimate factor in communicative complexity. Philosophical Transactions

618   of the Royal Society B: Biological Sciences 367: 1785-1801.

619 Galvani, G. L., R. A. González-Vaquero, C. Guerra-Navarro, and B. P. Settembrini.

620   2017. Antennal sensilla of cleptoparasitic and non-parasitic bees in two

621   subfamilies of Apidae. Apidologie 48:437–449.

622 Gamer, M., Lemon, J., Fellows, L, and P. Singh. 2019. r: Various Coefficients of

623   Interrater Reliability and Agreement. R package version 0.84.1.

624   https://CRAN.R-project.org/package=irr

625 Gibbs, J., S. G. Brady, K. Kanda, and B. N. Danforth. 2012. Phylogeny of halictine

626   bees supports a shared origin of eusociality for *Halictus* and *Lasioglossum*

627   (Apoidea: Anthophila: Halictidae). Molecular Phylogenetics and Evolution

628   65:926–939.

629 Gill, K. P., E. van Wilgenburg, D. L. Macmillan, and M. A. Elgar. 2013. Density of

630   antennal sensilla influences efficacy of communication in a social insect.

631   American Naturalist 182:834–840.

632 Green, J. P., C. Rose, and J. Field. 2012. The role of climatic factors in the

633   expression of an intrasexual signal in the paper wasp *Polistes dominulus.*

634   Ethology 118:766–774.

635 Groom, S. V. C., and S. M. Rehan. 2018. Climate-mediated behavioural variability in

636   facultatively social bees. Biological Journal of the Linnean Society 125:165–

637   170.

638 Grüter, C., F. H. I. D. Segers, L. L. G. Santos, B. Hammel, U. Zimmermann, and F.

639   S. Nascimento. 2017. Enemy recognition is linked to soldier size in a

640   polymorphic stingless bee. Biology Letters 13: 20170511.

641 Harrison, X. A. 2014. Using observation-level random effects to model

642   overdispersion in count data in ecology and evolution. PeerJ e616.

643 Hartig, F. 2020. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)

644   Regression Models (2017). R package version 0.1, 5.

645 Jones, B. M., C. J. Kingwell, W. T. Wcislo, and G. E. Robinson. 2017. Caste-biased

646   gene expression in a facultatively eusocial bee suggests a role for genetic

647   accommodation in the evolution of eusociality. Proceedings of the Royal

648   Society B: Biological Sciences 284:20162228.

649 Jones, J. C., P. Helliwell, M. Beekman, R. Maleszka, and B. P. Oldroyd. 2005. The

650   effects of rearing temperature on developmental stability and learning and

651   memory in the honey bee, *Apis mellifera*. Journal of comparative physiology.

652   A, Neuroethology, sensory, neural, and behavioral physiology 191:1121–

653   1129.

654 Kather, R and S. J. Martin. 2015. Evolution of Cuticular Hydrocarbons in the

655   Hymenoptera: a Meta-Analysis. Journal of Chemical Ecology 41: 871–883.

Koo, T. K., and M. Y. Li. 2016. A Guideline of Selecting and Reporting Intraclass

Correlation Coefficients for Reliability Research. Journal of Chiropractic

Medicine 15:155–163.

Korner-Nievergelt, F., Roth, T., von Felten, S., Guelat, J., Almasi, B., and P. Korner-

Nievergelt. 2015. Bayesian Data Analysis in Ecology using Linear Models with

R, BUGS and Stan. Elsevier.

McKenzie, S. K., Fetter-Pruneda, I., Ruta, V., and Kronauer, D. J. C. 2016.

Transcriptomics and neuroanatomy of the clonal raider ant implicate an

expanded clade of odorant receptors in chemical communication.

Proceedings of the National Academy of Sciences 113: 14091–14096.

Ozaki, M., A. Wada-Katsumata, K. Fujikawa, M. Iwasaki, F. Yokohari, Y. Satoji, T.

Nisimura, et al. 2005. Behavior: Ant nestmate and non-nestmate

discrimination by a chemosensory sensillum. Science 309:311–314.

Pask, G. M., Slone, J. D., Millar, J. G., Das, P., Moreira, J. A., Zhou, X., Bello, J.,

Berger, S. L., Bonasio, R., Desplan, C., Reinberg, D., Liebig, J., Zwiebel, L. J.,

and Ray, A. 2017. Specialized odorant receptors in social insects that detect

cuticular hydrocarbon cues and candidate pheromones. Nature

Communications 8: 297.

Peckre, L., P. M. Kappeler, P. M., and C. Fichtel. 2019. Clarifying and expanding the

social complexity hypothesis for communicative complexity. Behavioral

Ecology and Sociobiology 73: Article 11.

Pinto, J. D., R. K. Velten, G. R. Platner, and E. R. Oatman. 1989. Phenotypic

plasticity and taxonomic tharacters in *Trichogramma* (Hymenoptera:

Trichogrammatidae). Annals of the Entomological Society of America 82:414–

425.

Potts, S. G., and P. Willmer. 1997. Abiotic and biotic factors influencing nest-site

selection by Halictus rubicundus, a ground-nesting halictine bee. Ecological

Entomology 22:319–328.

Purcell, J. 2011. Geographic patterns in the distribution of social systems in

terrestrial arthropods. Biological Reviews 86:475–491.

Rehan, S. M., and A. L. Toth. 2015. Climbing the social ladder: the molecular

evolution of sociality. Trends in Ecology and Evolution 30: 426–433.

Renner, M. A., and J.C. Nieh. 2008. Bumble bee olfactory information flow and

contact-based foraging activation. Insects Socieaux 55: 417-424.

Riveros, A. J., and W. Gronenberg. 2010. Sensory allometry, foraging task

specialization and resource exploitation in honeybees. Behavioral Ecology

and Sociobiology 64:955–966.

Schradin, C. 2013. Intraspecific variation in social organization by genetic variation,

developmental plasticity, social flexibility or entirely extrinsic factors.

Philosophical Transactions of the Royal Society B: Biological Sciences 368:

20120346.

Schradin, C., L. D. Hayes, N. Pillay, C. Bertelsmeier. 2018. The evolution of

intraspecific variation in social organization. Ethology 124: 527–536.

Shell, W. A., and S. M. Rehan. 2018. Behavioral and genetic mechanisms of social

evolution: insights from incipiently and facultatively social bees. Apidologie 49:

13–30.

Spaethe, J., A. Brockmann, C. Halbig, and J. Tautz. 2007. Size determines antennal

sensitivity and behavioral threshold to odors in bumblebee workers.

Naturwissenschaften 94:733–739.

Wcislo, W. T. 1995. Sensilla numbers and antennal morphology of parasitic and non-

parasitic bees (Hymenoptera : Apoidea). International Journal of Insect

Morphology and Embryology 24:63–81.

Wcislo, W. T. 1997. Social interactions and behavioral context in a solitary bee

Lasioglossum (Dialictus) figueresi (Hymenoptera, Halictidae). Insects Sociaux

44: 199-208.

West-Eberhard, M.J. 2003. Developmental plasticity and evolution. Oxford University

Press, Oxford

Wittwer, B., A. Hefetz, T. Simon, L. E. K. Murphy, M. A. Elgar, N. E. Pierce, and S.

D. Kocher. 2017. Solitary bees reduce investment in communication

compared with their social relatives. Proceedings of the National Academy of

Sciences of the United States of America 114:6569–6574.