Sensory plasticity in a socially plastic bee

Rebecca A Boulton*[1,2] and Jeremy Field[3]

[1]Laboratory of Genetics, Plant Sciences Group, Wageningen University & Research,

6708 PB, Netherlands

[2]Biological and Environmental Sciences, University of Stirling, FK9 4NF, United

Kingdom

[3] College of Life and Environmental Sciences, University of Exeter, Penryn campus,

TR10 9FE, United Kingdom

*author for correspondence: rebeccaboulton87@googlemail.com

Abstract

The social Hymenoptera have contributed much to our understanding of the

evolution of sensory systems. Attention has focussed chiefly on how sociality and

sensory systems have evolved together. In the Hymenoptera, the antennal sensilla

are important for optimising the perception of olfactory social information. Social

species have denser antennal sensilla than solitary bees, which is thought to

enhance social cohesion through nest-mate recognition. In the current study, we test

whether sensilla numbers vary between regional populations of the socially plastic

sweat bee *Halictus rubicundus*. We found region level differences in both olfactory

and hygro/thermoreceptive sensilla numbers. We also found evidence that olfactory

sensilla density is developmentally plastic: when we transplanted bees from Scotland

to the south-east of England, their offspring (which developed in the south) had more

olfactory hairs than the transplanted individuals themselves (which developed in

Scotland). The transplanted bees displayed a mix of social (a queen plus workers)

and solitary nesting, but neither individual nor nest phenotype was related to sensilla

density. We suggest that this general, rather than caste-specific sensory plasticity

provides a flexible means to optimise sensory perception according to the most

pressing demands of the environment, and may support social plasticity in *H.*

rubicundus.

Introduction

Natural selection on sensory abilities has resulted in enormous diversity in the

mechanisms that organisms use to perceive their environment. For example, species

of Lake Tanganyika cichlids that have colonised more complex, rocky habitats have

greater visual acuity than sand-dwelling species (Dobberfuhl et al. 2005). These

mechanisms can also feedback into selective regimes, facilitating micro- and

macroevolutionary change through sensory drive. Sensory drive occurs when local

adaptation results in populations with diverging modes of communication and

perception. This has been proposed to lead to assortative mating between partners

with complementary signalling and perception systems, reducing geneflow and

promoting speciation (Boughman 2002).

The Hymenoptera have emerged as a model system to understand both the

proximate and ultimate factors that underlie sensory evolution. These insects have

antennal receptors with diverse functions and distinct forms that can easily be

characterised within and between species. This work has revealed that both the

density and form of antennal sense organs (sensilla) correlate with diet (Polidori et

al. 2012; 2020), sex (Babu 2019; Do Carmo Queiroz Fialho et al. 2014) and a

cleptoparasitic lifestyle (Wcislo 1995; Galvani et al. 2017). Attention has focussed

particularly on how eusociality has imposed selection to optimise antennal

perception. Eusociality involves reproductive division of labour with a queen/s

producing eggs and workers provisioning the brood. However, sociality takes diverse

forms across the bees and wasps and group sizes vary enormously, providing rich

opportunities for comparative study. Many obligately social species show worker

polyphenism, where some workers specialise on brood care and others on foraging

or guarding the nest (Andersson 1984). Compared with solitary species, social

species require more finely tuned communication systems to distinguish intruders

from kin and gain information from nestmates (Wcislo 1997; Renner and Nieh 2008).

Across the Hymenoptera, social species show tend to have greater densities of

antennal sensilla that are involved in the perception of olfactory cues than solitary

species (Elgar et al. 2018). This is thought to enhance social cohesion and

performance by facilitating nest-mate recognition (Ozaki et al. 2005; Gill et al. 2013;

Couto et al. 2017) and supporting caste polyphenisms (Spaethe et al. 2007; Riveros

and Gronenberg 2010; Gill et al. 2013; Gru□ter et al. 2017; Elgar et al. 2018).

The olfactory antennal sensilla have been proposed as a pre-adaptation that acts

alongside haplodiploidy to predispose the Hymenoptera to the evolution of sociality

(Couto et al. 2017). A conserved, ancestral olfactory subsystem involving the

basiconic antennal sensilla has been suggested to facilitate kin selection by allowing

nestmate recognition from cuticular hydrocarbon (CHCs) profiles (Ozaki et al. 2005;

McKenzie et al. 2016; Couto et al. 2017; Pask et al. 2017). While empirical evidence

for this subsystem exists for only two species (an ant and a hornet), the halictid bees

have also provided support for the possibility that sensory pre-adaptations might

contribute to the evolution of sociality. The Halictidae or sweat bees exhibit a range

of social structures, from solitary nesting through to obligate sociality, including

within-species social polymorphism and social plasticity (Gibbs et al. 2012). A

comparative study by Wittwer et al (2017) has shown that ancestrally solitary species

and their social sister species have equivalent densities of hair-like sensilla

(basiconic and trichoid sensilla) that detect olfactory cues. Species that have

reverted to a solitary existence from sociality on the other hand, have reduced

densities of these sensilla. This suggests that in the Halictidae, sensitivity to olfactory

cues precedes the evolution of sociality rather than evolving as a consequence of it.

In addition to caste polyphenisms, where individuals always form social groups but

vary in the tasks that they perform, a number of species are facultatively social. In

these species, individuals vary in their propensity to form social groups or nest as

solitary individuals (Gibbs et al. 2012). This social plasticity is thought to be key to

the evolutionary diversification of the Hymenoptera as it provides a genetic base

from which obligate eusociality and caste polyphenism can evolve repeatedly (West-

Eberhard 2003; Jones et al. 2017). It also provides an opportunity to test for

correlates of social evolution without the complication of interspecific differences. In

this study we characterise the antennal sensilla of a facultatively social species,

Halictus rubicundus, for the first time. We measure the density of different antennal

sensilla types between regional populations of this socially plastic sweat bee and test

whether sensory plasticity mirrors social plasticity in a transplant experiment. Our

results represent an important step in disentangling the sequence of events that led

to the evolution of eusociality and reproductive division of labour in the

Hymenoptera.

Materials and methods

Study system

Halictus rubicundus is a ground-nesting bee that can be found throughout Europe and North America in areas with sandy-loamy soil that receive sufficient sunlight. In Great Britain and Ireland, *H. rubicundus* demonstrates social variation according to latitude but retains social plasticity throughout its range (Field et al. 2010; 2012). In the north bees are typically solitary as the short growing season and cooler climate precludes the bivoltinism that is required for social living. In a typical year in the north, foundresses produce a single brood of offspring per year, the females of which enter hibernation and emerge to become foundresses the following spring (figure 1a). In the south, growing seasons tend to be longer and the weather is warmer, so bivoltinism and sociality can occur more frequently. After overwintering underground, females produce a first brood (the B1) which become workers and provision a second brood (B2) which are offspring of either the original foundress or a B1 replacement queen. B1 females are also observed to provision their nests alone (without workers) in the same year that they emerge. The B2 hibernate and emerge the following year to found their own nests (figure 1b; Field et al. 2010; 2012).

Sociality is extremely plastic in *H. rubicundus* in Great Britain, and when solitary northern bees are transplanted to the south, the B1 brood will often provision as workers or solitary foundresses in the same year (Field et al. 2010; 2012; this study). Here we test for fixed and plastic differences in antennal sensilla density in *H. rubicundus* across and within populations. We predict that bees from northern populations (with limited opportunity to form social nests) will have fewer olfactory

sensilla than bees from southern populations, where the climate allows some

individuals to form social nests most years. We also test whether antennal sensilla

numbers are developmentally plastic by transplanting bees from the north to the

south. We predict that the offspring of northern bees that develop in the warmer

south will have different numbers of antennal sensilla than their parental generation

which developed in the north. We would also expect to see differences in sensilla

numbers according to nest and individual phenotype, with social individuals

exhibiting greater densities of olfactory sensilla to aid in nestmate recognition.

Collection

Halictus rubicundus females were collected from 4 sites in the UK in 2018, 2019 and

2020 (see fig 1c). Two sites in the south-west (SW): Boscastle (Cornwall: N 50° 41'

24" W 4° 41' 24"; N = 24; 2018) and Bodmin (Cornwall, N 50° 30' 36" W 4° 33' 36"; N

= 8; 2019); a site at a mid-latitude (MID): Belfast (Northern Ireland, N 54° 32' 24" W

5° 58' 48"; N = 38; 2020); and a site in the far north of Scotland (SCO): Migdale

(Scotland, N 57° 53' 24" W 4° 15' 0"; N = 30; 2018 and 2020). Previous studies and

our own observations indicate that while a proportion of nests are social in SW (this

study), nests are always non-social in Belfast (MID; Field et al. 2010) and Scotland

(SCO; this study). However, bees from Belfast (MID; Field et al. 2010) and Scotland

(SCO; this study) are socially plastic and may have social nests when moved to

more southerly sites. Bees sampled were thought to be foundresses based on

collection date. We collected females by hand-netting in May and early June in the

SW, and in late June in SCO. In Belfast (MID) we excavated overwintering

foundresses in February. We note that some bees from the SW (Boscastle) may

have been early emerging B1 but all had fairly worn wings indicative of several

weeks of provisioning.

Transplant

In addition to sampling from native populations, we collected bees that had been

transplanted from Scotland (SCO) to a site in the south-east (SE), the Knepp estate

in West Sussex (N 50° 53' 60" W 0° 21' 36"; N = 95; see fig 1c). Buckets of soil were

embedded within a nest aggregation at the Scottish site during winter 2018-19, and

native foundresses subsequently nested in them during 2019.Their B1 offspring then

emerged in late summer 2019 and hibernated in the buckets. The buckets containing

hibernating Scottish bees were then taken from Scotland to the south-east in the

spring of 2020, where they were re-embedded in the ground. SCO foundresses that

emerged were marked with a dot of enamel paint when they provisioned the B1

brood in the south-east. When the B1 emerged, they were marked with a different

colour of enamel paint. Each nest was marked with a numbered nail at the entrance.

This allowed us to distinguish fresh B1 females from the original foundresses and to

determine whether nests were solitary (with a single B1 female or original foundress

provisioning alone) or social (with multiple B1 females provisioning the nest). At the

end of the B1 provisioning phase we collected SCO foundresses (which had

emerged in Scotland and been transplanted to the SE) and their B1 offspring (which

had developed and emerged in the SE).

Specimen storage and preparation

All specimens were stored in 95% Ethanol until they were prepared for imaging. One

antenna was removed from each bee (whether it was the left or right was recorded)

and mounted on a JEOL aluminium stub (10mm dia x 10mm high) using a PELCO

carbon conductive sticky tab (10mm dia). Specimens were mounted dorsal side up.

Specimens were carbon coated using spectograpically pure carbon.

Imaging

Mounted and carbon coated specimens were imaged using a TESCAN Vega SEM in

High Vacuum mode at 20.0kV. The scan mode was set to resolution and the

magnification was 485x. Only the two distal segments (11 and 12), which have the

highest density of sensilla, were imaged. The length and area of each antennal

segment was measured during imaging and images of each antennal segment from

the SEM were saved in TIF format.

Figure 1 (A) univoltine lifecycle of *Halictus rubicundus* which prohibits social nesting (B) bivoltine lifecycle of *Halictus rubicundus* where sociality is possible as the 1st brood (B1) can provision a second brood (B2). Note that in (B) the B1 females can either (i) overwinter and provision the following year as a solitary foundress (ii) provision a nest alone as a solitary foundress in the year that they emerge or (iii) provision the nest socially with their B1 sisters with a queen (their mother or a sister) in the year that they emerge. (C) Map depicting the sites where *H. rubicundus* females were studied. Dashed arrow represents the direction of the transplant experiment (SCO to SE).

Image scoring

Eight distinct types of sensilla with overlapping functions have been characterised in

the Hymenoptera (Do Carmo Queiroz Fialho et al. 2014). Placoid sensilla are plate-

like structures that are involved in olfaction and chemoreception. Tricoid and

basiconic sensilla are hair-like projections with multiple sub-types that are used in

olfaction involving contact. Coeloconic, campaniform and ampulliform sensilla are

pore-like and are involved in sensing temperature, humidity and CO_2 concentration

(Do Carmo Queiroz Fialho et al. 2014). For this study we grouped all sensilla types

into three structural/functional groups that allowed for repeatability in scoring: (i)

olfactory plate sensilla (sensilla placodea, fig 2), (ii) olfactory hair-type sensilla

(sensilla trichodea and basonica; fig 2) and (iii) temperature/humidity/CO_2 pore

sensilla (coeloconic/campaniform/ampulliform sensilla; fig 2).

Images of antennal segments 11 and 12 were scored using ImageJ software. For

each segment, three 50µM x 50µM 'quadrats' were selected, the X and Y

coordinates of which were derived by generating a random number using the excel

function RANDBETWEEN. Each quadrat was saved as a TIF image and the number

of sensilla types i and ii were counted. We counted sensilla type iii across the entire

segment as they are distributed unevenly on the surface of the antenna. In some

cases, bare patches with no sensilla were present on the surface of the antenna.

Any quadrats which fell on the bare areas were discarded and another area was

selected. We note that the scorer was not blind to the population of origin as the

same person was required to mount, image and score the antennae.

219

Figure 2 SEM image showing a *Halictus rubicundus* antenna tip. (A) shows the

distal 2 antennal segments (11 and 12) that were imaged, (B) and (C) show the

types (i, ii, and iii) of antennal sensilla that were counted. Scale is given in

μM.

Analysis

Repeatability

Thirty-three of the 130 sampled antennae were scored a second time (by the same

person) for all sensilla types. The repeatability of image scoring was assessed using

the intra-class correlation coefficient, function *ICC* in the R package *irr* (Gamer et al,

2019).

Sensilla numbers

We used linear mixed models in the R package lme4 (Bates et al. 2019) to test for

differences in the numbers of sensilla types (i-iii). For all models outlined below, the

response variable was total sensilla counted, either i) olfactory plates, total across 3

quadrats for each segment per individual; ii) olfactory hairs, total across 3 quadrats

for each segment per individual; or iii) temperature/humidity pores, total across each

segment per individual.

Region, development site and social phenotype

To test whether bees from different regions have different numbers of sensilla we

included region as a fixed factor with 3 levels in LMMs: SW (Boscastle and Bodmin

were aggregated as only 8 individuals were collected from Bodmin), MID (Belfast,

Northern Ireland) and SCO (Migdale, Scotland, far north).

We ran additional models to test for within population differences in sensilla density

and determine whether this trait is plastic. To do this we included only bees that

developed in Scotland (SCO). LMMs included a fixed effect of development site, with

2 levels: SCO (bees that emerged in Scotland and were either collected there or

were moved to the south-east as adults) and SE (bees that emerged as adults after

immature development in the south-east, the B1 offspring of the bees that were

transplanted from Scotland to the south-east). Note that *H. rubicundus* females mate

before overwintering and so both the mother and father of the B1 bees collected in

the south-east originated in Scotland.

Additional LMMs were used to test whether B1 offspring collected in the SE had

higher counts of sensilla if they were from social nests of transplanted females

compared to those which started their own solitary nests. Social nests had a queen

and one or more workers, while solitary nests had a single B1 foundress that

produced offspring in the same year. For these models we included nest phenotype

(social or solitary) as a fixed effect with 2 levels. We did not sample multiple

individuals from the same nest, so did not include nest as a random effect. We did

not recover any queens from social nests and so could not test for an effect of bee

phenotype within social nests. We did run a model to test for effects of individual,

rather than nest-level, phenotype across social and solitary nests. For this analysis

bees were characterised as (1) a future reproductive (a B1 individual that had

emerged in 2020 and did not provision that year at all); (2) a worker (a B1 individual

that provisioned a nest containing a queen); (3) a solitary foundress from 2019 (a

foundress that emerged in SCO in 2019 and provisioned a nest alone and laid eggs);

(4) a solitary foundress from 2020 (a foundress that emerged in the B1 generation in

2020 in the SE and provisioned a nest alone).

Finally, we ran a model using the bees collected in the SE to test whether age-

related wear and tear reduces density of all 3 sensilla types. For these analyses age

was binary: bees were scored as fresh (newly emerged and had not provisioned a

nest; N = 26) or old (had provisioned a nest for several weeks N = 40).

All LMMs included antenna segment number (11 or 12) as a fixed factor. The

interaction effect between region/development type/social phenotype and segment

number was also fitted. We included a random effect of individual as two segments

were imaged for each bee. Models were run using the function *lmer* in the R

package lme4 (Bates et al, 2015). The package DHARMa was used to determine the

most appropriate error structure for all models (Gaussian) and to interpret residual

plots for lack of fit (Hartig, 2020).

Results

Repeatability and effects of age

Counts of all sensilla types were highly repeatable (Koo and Li 2016; ICC greater

than 0.85 for all, for a detailed summary see supplementary material table S1).

We also found no evidence that age-related wear and tear influences the variation in

sensilla density; there was no effect of age on density of any of the three types of

sensilla (type i: X^2 = 0.25, df = 1, p = 0.62; type ii: X^2 = 0.01, df = 1, p = 0.92; type iii:

X^2 = 0.09, df = 1, p = 0.76).

Differences between antennal segments

299 Olfactory plates (type i) and olfactory hairs (type ii) were found in greater numbers on

300 the most distal antennal segment, segment 12, while hygro/thermoreceptors (type iii)

301 were found in higher numbers on segment 11. There were no interaction effects

302 between segment number and region (table 1).

Region

There was a significant effect of region on the counts of type ii (olfactory hairs) and iii

(thermo/hygroreceptors) sensilla, but not type i (olfactory plate receptors; see table

1, figure 3a-c). Pairwise tests show that bees from a mid-latitude (MID; Belfast; N =

38) had significantly more olfactory hairs (type ii) than bees collected from the far

north (Scotland; N=30; p = 0.03; figure 3b) but bees from the south-west (N = 32)

had numbers of olfactory hairs that were statistically equivalent to MID (N = 32; p =

0.06) and Scottish (p = 0.93) bees. Bees from SCO and MID had more thermo/hygro

(type iii) receptors than SW bees (p < 0.05; figure 3c). Bees from the south-west had

the most olfactory plate sensilla (type i) but this was not statistically significant (figure

3a).

Table 1 LMM results showing effect of region on antennal sensilla counts across

segments using type II sums of squares (p-values for main effects are calculated

independent from interaction effects which in all cases are non-significant).

Sensilla type	Effect	X^2	df	P
Type i) olfactory plates	Region	2.31	2	0.31
	Segment	19.48	1	1.01×10^{-5}
	Region*Segment	0.45	2	0.80
Type ii) olfactory hairs	Region	7.81	2	0.02
	Segment	41.74	1	1.04×10^{-10}
	Region*Segment	2.70	2	0.26
Type iii) thermo/hygro-receptive	Region	9.14	2	0.01
	Segment	19.17	1	1.20×10^{-5}
	Region*Segment	0.35	2	0.84

Figure 3 Numbers of (a) and (d) olfactory plates, i; (b) and (e) olfactory hairs, ii; and

(c) and (f) hygro/thermoreceptive pores, iii, for *Halictus rubicundus* across: a-c three

sampled regions: SW (South-west; mostly social), MID (Belfast; solitary) and SCO

(Scotland; solitary); d-e for *Halictus rubicundus* originating from Scotland and

collected after developing as larvae in Scotland (SCO) or after developing as larvae

at the transplanted site in the south-east (SE). Different uppercase letters over bars

indicate significantly different counts of sensilla. Black points represent sensilla

counts on segment 11 and grey points on segment 12. Box plots show median and

quartiles across segments 11 and 12.

Development site

The density of olfactory hairs (type ii) varied according to where bees originating

from Scotland spent their immature development (table 2; figure 3d-e). Offspring of

transplanted bees which developed in the south-east (N=59) had more olfactory

hairs (type ii, p = 0.01) than conspecifics that developed in their natal Scottish site

(N=36; fig 3e). A similar trend was seen in olfactory plates (I; fig 3d) but this was not

statistically significant (p = 0.06). There was no effect of origin on

thermo/hygroreceptor numbers (iii; fig 3f).

Table 2 LMM results showing effect of origin of emergence on antennal sensilla

counts across segments (p-values for main effects are calculated independent from

interaction effects which in all cases are non-significant).

Sensilla type	Effect	X^2	df	P
Type i) olfactory plates	Development site	3.44	1	0.0
	Segment	34.82	1	3.6
	Development site *Segment	0.61	1	0.4
Type ii) olfactory hairs	Development site	6.41	1	0.0
	Segment	61.35	1	4.7
	Development site *Segment	1.57	1	0.2
Type iii) thermo/hygroreceptive	Development site	0.74	1	0.3
	Segment	33.25	1	8.1
	Development site *Segment	0.15	1	0.6

Sociality

B1 workers from social nests (N= 12) and solitary B1 females that founded new

nests alone (N=18) had equivalent sensilla numbers across all 3 categories (i-iii, see

table 3; fig 4a-c) and there were no interactions between nest phenotype and

antenna segment number. There was some suggestion that solitary bees had higher

numbers of olfactory plates (type i), but this was not statistically significant (p = 0.07).

Similarly there were no differences in numbers of any sensilla types across different

individual phenotypes (B1 future reproductive emerged in 2020 in SE and did not

provision that year, will overwinter, N = 11; B1 worker emerged in 2020 in the SE, N

= 7; B1 solitary foundress that emerged 2019 in SCO and provisioned in the SE in

2020, N = 4; and B1 solitary foundress emerged and provisioned in 2020 in SE, N =

4; see table 4; figure 4d-e).

Table 3 LMM results showing effect of nest phenotype (social/solitary) on antennal

sensilla counts across segments (p-values for main effects are calculated

independent from interaction effects which in all cases are non-significant).

Sensilla type	Effect	X^2	df	P
Type i) olfactory plates	Nest phenotype	2.91	1	0.08
	Segment	11.80	1	0.0005
	Nest phenotype*Segment	1.08	1	0.30
Type ii) olfactory hairs	Nest phenotype	0.08	1	0.78
	Segment	42.03	1	8.98×10^{-11}
	Nest phenotype*Segment	1.29	1	0.26
Type iii) thermo/hygroreceptive	Nest phenotype	0.29	1	0.59

		33.08	1	8.86×10^{-9}
Segment		33.08	1	8.86×10^{-9}
Nest phenotype*Segment		0.10	1	0.75

Table 4 LMM results showing effect of bee phenotype (future

reproductive/worker/solitary foundress 2019 and solitary foundress 2020) on

antennal sensilla counts across segments (p-values for main effects are calculated

independent from interaction effects which in all cases are non-significant).

Sensilla type	Effect	X^2	df	P
Type i) olfactory plates	Bee phenotype	1.51	3	0.67
	Segment	22.34	1	2.29×10^{-6}
	Bee phenotype*Segment	1.81	3	0.61
Type ii) olfactory hairs	Bee phenotype	1.24	3	0.74
	Segment	42.03	1	8.33×10^{-11}
	Bee phenotype*Segment	0.85	3	0.84
Type iii) thermo/hygroreceptive	Bee phenotype	3.39	3	0.34
	Segment	30.18	1	3.95×10^{-8}
	Bee phenotype*Segment	2.26	3	0.52

Figure 4 Numbers of (a) and (d) olfactory plates, i; (b) and (e) olfactory hairs, ii; and

(c) and (f) hygro/thermoreceptive pores, iii; for *Halictus rubicundus* from Scotland

collected from (a-c) social and solitary nests and according to individual phenotype

(d-f) after being transplanted to the south-east in 2020. For figures 4d-4f: F-2021

female that emerged in the SE in 2020 but did not provision that year; F-2020 female

that emerged in the SE in 2020 and provisioned a nest as a solitary foundress; F-

2019 female that emerged in SCO in 2019 and provisioned a nest as a solitary

foundress in 2020; W-2020 female that emerged in the SE in 2020 and provisioned a

social nest in the same year as a worker. Black points represent sensilla counts on

segment 11 and grey points on segment 12. Box plots show median and quartiles

across segments 11 and 12. There were no significant differences in counts of any

sensilla type across the groups.

Discussion

The social Hymenoptera have contributed much to our understanding of the

proximate and ultimate factors that underlie the evolution of sensory systems.

Attention has focussed chiefly on how sociality has imposed selection to optimise

olfactory communication, enhancing social cohesion through nest-mate recognition

and supporting caste polyphenisms (Wcislo 1997; Ozaki et al. 2005; Spaethe et al.

2007; Renner and Nieh 2008; Riveros and Gronenberg 2010; Gill et al. 2013; Couto

et al. 2017; Gru□ter et al. 2017; Elgar et al. 2018).

In the current study we measured sensilla density in the socially plastic sweat bee

Halictus rubicundus. We found that when Scottish bees were transplanted from

Scotland, where they are solitary, to the south where they can be social, they

produced offspring with higher densities of olfactory hairs. These results suggest that

the density of olfactory hairs may, like sociality, be phenotypically plastic in *H.*

rubicundus. We also found evidence for between-region differences in the density of

basiconic/trichoid olfactory hairs (and hygro/thermoreceptive sensilla) for

untransplanted bees, but these differences did not correlate with the expected

degree of sociality.

The results from our transplant experiment suggest that olfactory hair density is

phenotypically plastic and depends on the conditions that bees experience during

development. When Scottish bees were transplanted to the south-east, their

offspring that developed there (where air and soil temperatures in June and July are

2-5°C warmer than their native site; table S2) had greater densities of

trichoid/basiconic hairs. It may be that this trait is plastic and warmer temperatures

during development lead to greater densities of olfactory hairs. A similar pattern is

seen in the parasitoid wasp *Trichogramma*; when males develop under colder

temperatures they have reduced numbers of basiconic hairs (Pinto et al. 1989).

Social plasticity may be involved in the maintenance of this developmental plasticity.

Northern and mid-latitude bees can have social nests when the climate allows (Field

et al. 2010; 2012; this study), and the plasticity in olfactory hair density that we saw

in this study may support social plasticity, facilitating social olfactory communication

only in favourable climates, when it is likely to be needed.

In addition to plasticity within a population, we also saw evidence that olfactory hair

density (type ii, basiconic/trichoid sensilla) varies across regional populations in *H.*

rubicundus. In contrast we saw no differences in the densities of placoid (plate-like,

type i) olfactory sensilla across regions. This may relate to the function of these

receptor types. Basiconic/trichoid hairs respond to contact with the CHCs of other

individuals, and so are likely to be required for nest-mate recognition and

communication, while placoid plates are thought to be involved in longer range

olfaction to detect food and hosts (Ozaki et al. 2005; McKenzie et al. 2016; Couto et

al. 2017; Pask et al. 2017). We know of no clinal variation in diet breadth in this

species (although this remains to be rigorously studied), which fits with the pattern

we see here.

Bees collected from Belfast at a mid-latitude had the highest density of olfactory hair-

like sensilla (type ii), higher than bees from Scotland (450km north of Belfast) but

statistically equivalent to bees collected in the south-west. We predicted that *H.*

rubicundus collected further south would have the greatest densities of olfactory

sensilla to support social communication between nestmates (Wittwer et al. 2017;

Elgar et al. 2018). In the south, summers are typically warmer and longer and a

higher proportion of the population is expected to exhibit social behaviour more

regularly. While we did see regional differences in olfactory sensilla density, bees

from the south-west did not have the highest densities as we would have predicted

based on clines in sociality. The regional differences we saw may be related to

developmental plasticity, which could limit the scope for fixed population level

differences to evolve in the predicted direction. Longitudinal studies of these same

populations and additional, reciprocal transplant experiments (from south to north)

would help to elucidate the extent to which genetically fixed differences in olfactory

hair density and plastic expression contribute to the population level differences we

saw in this trait.

We also found that the density of sensilla (type iii) involved in the perception of

humidity, temperature and CO_2 varies across regions, but unlike olfactory hairs this

does not appear to be plastic. Bees from a mid-latitude and the north of scotland

(Belfast and Scotland) have more thermo/hygro receptive sensilla than bees from the

south-west (400 km south of Belfast and 850 km south of the Scottish site). The

density of these sensilla does not appear to be plastic as it was not influenced by

where the bees developed or their social phenotype. This pattern may be a result of

a more extreme, variable climate in the north, which leads to consistent selection on

higher densities of thermo/hygro receptors. Sweat bees including *H. rubicundus* are

highly sensitive to temperature and rain. Flight activity is constrained by low ambient

temperatures, lack of sunlight and rainfall (Potts and Wilmer 1997). In the north,

bees experience colder ambient temperatures and greater rainfall. Bees with more of

these receptors may be more sensitive to current and oncoming weather conditions,

so that they have reduced mortality and improved foraging based on the climate. The

potentially severe fitness consequences of incorrect perception of climatic cues may

explain the lack of plasticity in the number of thermo/hygro receptors if selection has

a strong and canalising effect on this trait.

Our results add a new dimension to the growing body of evidence that

developmental temperature may contribute to sociality in the Hymenoptera. While

previous studies suggest that temperatures experienced by developing larvae

contribute to individual differences in social phenotype (i.e. caste polyphenism:

Czekońska and Tofilski 2020; Becher et al. 2009; the development of status badges:

Green et al. 2012; memory formation: Jones et al. 2005 and olfactory learning: Anton

and Rossler 2021), our results in the socially plastic *H. rubicundus* point to a more

general response to the environment that is not directly related to the social

phenotype of the nest or individual (i.e. a worker, solitary foundress or queen).

Scottish bees that developed in the south-east had more olfactory hairs than their

conspecifics that emerged in Scotland irrespective of their caste or nest phenotype.

In the Halictidae, Wittwer et al (2017) found that halictid bee species that had

reverted back to solitary nesting from a state of sociality had reduced olfactory hair

density compared to social species and ancestrally solitary species. They suggest

that this is because dense olfactory sensilla are a pre-adaptation that facilitates the

evolution of sociality and may contribute to the repeated evolutionary transitions to

sociality seen in the halictid bees. Our results expand on this, suggesting that

olfactory hair density may also be phenotypically plastic. More broadly, plasticity in

traits such as this might contribute to the evolutionary lability of sociality in the

Halictidae and in the Hymenoptera, acting alongside social plasticity in the

hymenopteran ancestor to provide the 'flexible stem' which allowed for repeated

evolutionary transitions to obligate sociality across the order (West-Eberhard 2003;

Wittwer et al. 2017).

Acknowledgements

Many thanks to Gavyn Rollinson and Sharon Uren for SEM training and technical

support. This work is part of a project that received funding from the European

Research Council (ERC) under the European Horizon's 202 research and innovation

programme (grant agreement No. 695744).

Conflict of interest

The authors have no conflict of interests to report.

Ethics

This research adheres to ethical codes of practice at the University of Exeter. No

vertebrates were used in this research. Bee antenna samples were taken from living

and dead stored specimens. Handling during antenna removal of live specimens was

done as quickly as possible to minimize stress to individuals. Bees frequently lose

parts of and whole antennae in nature and we did not see any negative effects on

the survival or provisioning behaviour of live sampled individuals compared to intact

bees. The sample sizes reported were considered appropriate to maximize statistical

power while reducing the number of individuals involved in the experiments.

Individuals were sampled from large populations > 500 individuals and numbers

taken did not pose a threat to the persistence of these populations.

Data accessibility

All code and data are archived on Github and can be accessed here:

https://github.com/DrBecky-B/Bee.Antennae

References

Andersson, M. 1984. The evolution of eusociality. Annual review of Ecology,

Evolution and Systematics 15:165–189.

Anton, S., and W. Rössler. 2021. Plasticity and modulation of olfactory circuits in

insects. Cell and Tissue Research 383:149–164.

Babu, M. 2019. Antennal sensilla of the weaver ant *Oecophylla smaragdina* (F.) –

males and females sense differently. Indian Journal of Entomology 81:418-

422.

Bates, D., Maechler, M., Bolker, B and W. Walker. 2015. Fitting Linear Fixed-Effects

Models using lme4. Journal of statistical software 67:1-48

Becher, M. A., H. Scharpenberg, and R. F. A. Moritz. 2009. Pupal developmental

temperature and behavioral specialization of honeybee workers (Apis

mellifera L.). Journal of Comparative Physiology A: Neuroethology, Sensory,

Neural, and Behavioral Physiology 195:673–679.

Boughman, J. W. 2002. How sensory drive can promote speciation. Trends in

Ecology and Evolution 17:571–577.

Charnov, E. L. 1978. Evolution of eusocial behavior: Offspring choice or parental

parasitism? Journal of Theoretical Biology 75:451–465.

Couto, A., A. Mitra, D. Thiéry, F. Marion-Poll, and J. C. Sandoz. 2017. Hornets have

it: A conserved olfactory subsystem for social recognition in hymenoptera?

Frontiers in Neuroanatomy 11:1–12.

Czekońska, K., and A. Tofilski. 2020. Body mass of honey bee drones developing in

constant and in changing temperatures. Apidologie 51:510–518.

Dobberfuhl, A. P., J. F. P. Ullmann, and C. A. Shumway. 2005. Visual acuity,

environmental complexity, and social organization in African cichlid fishes.

Behavioral Neuroscience 119:1648–1655.

Do Carmo Queiroz Fialho, M., C. P. Guss-Matiello, J. C. Zanuncio, L. A. Oliveira

Campos, and J. E. Serrão. 2014. A comparative study of the antennal sensilla

in corbiculate bees. Journal of Apicultural Research 53:392–403.

Elgar, M. A., D. Zhang, Q. Wang, B. Wittwer, H. T. Pham, T. L. Johnson, C. B.

Freelance, et al. 2018. Insect antennal morphology: The evolution of diverse

solutions to odorant perception. Yale Journal of Biology and Medicine 91:457–

469.

Field, J., R. J. Paxton, A. Soro, and C. Bridge. 2010. Cryptic plasticity underlies a

major evolutionary transition. Current Biology 20:2028–2031.

Field, J., R. Paxton, A. Soro, P. Craze, and C. Bridge. 2012. Body size, demography

and foraging in a socially plastic sweat bee: A common garden experiment.

Behavioral Ecology and Sociobiology 66:743–756.

Galvani, G. L., R. A. González-Vaquero, C. Guerra-Navarro, and B. P. Settembrini.

2017. Antennal sensilla of cleptoparasitic and non-parasitic bees in two

subfamilies of Apidae. Apidologie 48:437–449.

Gamer, M., Lemon, J., Fellows, L, and P. Singh. 2019. r: Various Coefficients of

Interrater Reliability and Agreement. R package version 0.84.1.

https://CRAN.R-project.org/package=irr

Gibbs, J., S. G. Brady, K. Kanda, and B. N. Danforth. 2012. Phylogeny of halictine

bees supports a shared origin of eusociality for *Halictus* and *Lasioglossum*

(Apoidea: Anthophila: Halictidae). Molecular Phylogenetics and Evolution

65:926–939.

Gill, K. P., E. van Wilgenburg, D. L. Macmillan, and M. A. Elgar. 2013. Density of

antennal sensilla influences efficacy of communication in a social insect.

American Naturalist 182:834–840.

Green, J. P., C. Rose, and J. Field. 2012. The role of climatic factors in the

expression of an intrasexual signal in the paper wasp *Polistes dominulus.*

Ethology 118:766–774.

Groom, S. V. C., and S. M. Rehan. 2018. Climate-mediated behavioural variability in

facultatively social bees. Biological Journal of the Linnean Society 125:165–

170.

Grüter, C., F. H. I. D. Segers, L. L. G. Santos, B. Hammel, U. Zimmermann, and F.

S. Nascimento. 2017. Enemy recognition is linked to soldier size in a

polymorphic stingless bee. Biology Letters 13: 20170511.

Harrison, X. A. 2014. Using observation-level random effects to model

overdispersion in count data in ecology and evolution. PeerJ e616.

Hartig, F. 2020. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)

Regression Models (2017). R package version 0.1, 5.

Jones, B. M., C. J. Kingwell, W. T. Wcislo, and G. E. Robinson. 2017. Caste-biased

gene expression in a facultatively eusocial bee suggests a role for genetic

accommodation in the evolution of eusociality. Proceedings of the Royal

Society B: Biological Sciences 284:20162228.

Jones, J. C., P. Helliwell, M. Beekman, R. Maleszka, and B. P. Oldroyd. 2005. The

effects of rearing temperature on developmental stability and learning and

memory in the honey bee, *Apis mellifera*. Journal of comparative physiology.

A, Neuroethology, sensory, neural, and behavioral physiology 191:1121–

1129.

Koo, T. K., and M. Y. Li. 2016. A Guideline of Selecting and Reporting Intraclass

Correlation Coefficients for Reliability Research. Journal of Chiropractic

Medicine 15:155–163.

Korner-Nievergelt, F., Roth, T., von Felten, S., Guelat, J., Almasi, B., and P. Korner-

Nievergelt. 2015. Bayesian Data Analysis in Ecology using Linear Models with

R, BUGS and Stan. Elsevier.

McKenzie, S. K., Fetter-Pruneda, I., Ruta, V., and Kronauer, D. J. C. 2016.

Transcriptomics and neuroanatomy of the clonal raider ant implicate an

expanded clade of odorant receptors in chemical communication.

Proceedings of the National Academy of Sciences 113: 14091–14096.

Ozaki, M., A. Wada-Katsumata, K. Fujikawa, M. Iwasaki, F. Yokohari, Y. Satoji, T.

Nisimura, et al. 2005. Behavior: Ant nestmate and non-nestmate

discrimination by a chemosensory sensillum. Science 309:311–314.

Pask, G. M., Slone, J. D., Millar, J. G., Das, P., Moreira, J. A., Zhou, X., Bello, J.,

Berger, S. L., Bonasio, R., Desplan, C., Reinberg, D., Liebig, J., Zwiebel, L. J.,

and Ray, A. 2017. Specialized odorant receptors in social insects that detect

cuticular hydrocarbon cues and candidate pheromones. Nature

Communications 8: 297.

Pinto, J. D., R. K. Velten, G. R. Platner, and E. R. Oatman. 1989. Phenotypic

plasticity and taxonomic tharacters in *Trichogramma* (Hymenoptera:

Trichogrammatidae). Annals of the Entomological Society of America 82:414–

425.

Potts, S. G., and P. Willmer. 1997. Abiotic and biotic factors influencing nest-site

selection by Halictus rubicundus, a ground-nesting halictine bee. Ecological

Entomology 22:319–328.

Purcell, J. 2011. Geographic patterns in the distribution of social systems in

terrestrial arthropods. Biological Reviews 86:475–491.

Renner, M. A., and J.C. Nieh. 2008. Bumble bee olfactory information flow and

contact-based foraging activation. Insects Socieaux 55: 417-424.

Riveros, A. J., and W. Gronenberg. 2010. Sensory allometry, foraging task

specialization and resource exploitation in honeybees. Behavioral Ecology

and Sociobiology 64:955–966.

Spaethe, J., A. Brockmann, C. Halbig, and J. Tautz. 2007. Size determines antennal

sensitivity and behavioral threshold to odors in bumblebee workers.

Naturwissenschaften 94:733–739.

Wcislo, W. T. 1995. Sensilla numbers and antennal morphology of parasitic and non-

parasitic bees (Hymenoptera□: Apoidea). International Journal of Insect

Morphology and Embryology 24:63–81.

Wcislo, W. T. 1997. Social interactions and behavioral context in a solitary bee

Lasioglossum (Dialictus) figueresi (Hymenoptera, Halictidae). Insects Sociaux

44: 199-208.

West-Eberhard, M.J. 2003. Developmental plasticity and evolution. Oxford University

Press, Oxford

Wittwer, B., A. Hefetz, T. Simon, L. E. K. Murphy, M. A. Elgar, N. E. Pierce, and S.

D. Kocher. 2017. Solitary bees reduce investment in communication

compared with their social relatives. Proceedings of the National Academy of

Sciences of the United States of America 114:6569–6574.