LKB1 acts as a critical brake for the glucagon-mediated fasting response

Suehelay Acevedo-Acevedo*, Megan L. Stefkovich*, Sun Woo Sophie Kang, Rory P. Cunningham, Constance M. Cultraro, and Natalie Porat-Shliom‡.

Thoracic and GI Malignancies Branch, Center for Cancer Research, National Cancer Institute, NIH, Bethesda, MD.

* These authors contributed equally to this work.

‡ To whom correspondence should be addressed: Natalie Porat-Shliom, Thoracic and GI Malignancies Branch, Center for Cancer Research, National Cancer Institute, Bethesda, MD 20892; poratshliomn@mail.nih.gov; Tel. (240) 760-6847.

Key words: LKB1/STK11, fasting, glucagon, liver zonation, intravital microscopy

Abstract

As important as the fasting response is for survival, an inability to shut it down once nutrients become available can lead to exacerbated disease and severe wasting. The liver is central to transitions between feeding and fasting states, with glucagon being a key initiator of the fasting response in the liver. However, the precise regulatory mechanisms behind glucagon-induced fasting are unclear. One potential mediator of these transitions is Liver Kinase B1 (LKB1) given its role in nutrient sensing. Here, we show LKB1 knockout mice have a severe wasting and prolonged fasting phenotype despite increased food intake. By applying RNA sequencing and intravital microscopy we show that loss of LKB1 leads to a dramatic reprogramming of the hepatic lobule through robust upregulation of periportal genes and functions. This is mediated through the opposing effect LKB1 has on glucagon signaling and gene expression. Conclusion: our findings show that LKB1 acts as a brake to the glucagon-mediated fasting response resulting in "periportalization" of the hepatic lobule and whole-body metabolic inefficiency. These findings reveal a new mechanism by which hepatic metabolic compartmentalization is regulated by nutrient-sensing.

Introduction

Fasting is a conserved, adaptive response to nutrient deprivation that involves metabolic and transcriptional changes. The liver is critical in this response by sequentially activating catabolic and inhibiting anabolic processes to meet energy demands. Glucagon is at the crux of regulating the hepatic metabolic adaptation to fasting. When circulating glucose levels drop, glucagon stimulates hepatic glucose output via its release from glycogen storages or *de novo* synthesis through gluconeogenesis, as well as fatty acid oxidation and ketogenesis which provide crucial fuels to peripheral tissues during extended fasting [1]. Equally important is the ability to shut down the fasting response once nutrients become available. This action prevents catabolic processes from continuing needlessly and preventing gluconeogenesis, lipolysis, and ketogenesis to go unchecked. Indeed, impairment of fasting inhibition is observed in Non-Alcoholic Fatty Liver Disease (NAFLD) patients where gluconeogenesis is elevated [2]. Indeed, glucagon-like peptide 1 (GLP-1) agonists have been successful in treatment of obesity, type 2 diabetes, and NAFLD via attenuation of the glucagon response [3-5]. Fine tuning of this complex response is temporally regulated by glucagon through activation of transcription factor cascades leading to sequential waves of gene expression that optimize energy usage and prevent tissue damage [6].

Recently, *Cheng at al*. demonstrated that glucagon is also involved in the spatial regulation of hepatic metabolism [7, 8]. The spatial segregation of hepatic metabolism, also known as liver zonation, permits simultaneous operation of conflicting pathways [9-11]. This is achieved through the organization of hepatocytes in polarized, hexagonal lobules in which six portal vessels deliver oxygen, nutrients, and hormones (Sup Fig 1E) that flows directionally towards a single central vein. Several factors regulate zonation by affecting differential gene expression along the periportal (PP) to pericentral (PC) axis. For example, hypoxia and Wnt ligands activate β-catenin signaling resulting in the expression of target genes in PC cells [12, 13]. On the other hand, glucagon controls gene expression in the PP zone [8]. While the 'push and pull' relationship between Wnt ligand and glucagon have been established as a central determinant exerting zone-dependent expression patterns, it is not clear how hepatocytes integrate those signals during periods of fasting to maintain homeostasis.

One potential mediator is Liver kinase B1 (LKB1), given its role as a nutrient sensor that activates AMP-activated Protein Kinase (AMPK) and members of the AMPK-related (ARKs) family. LKB1 is also an important regulator of hepatic polarity [14-17], a feature that may be important for the spatial organization of the lobule. Loss of LKB1 in whole body [18] and liver-specific knockout [16, 17, 19] results in stark metabolic changes resembling cachexia, a condition that causes extreme weight loss and muscle wasting. Hepatic LKB1 was shown to suppress gluconeogenesis through the activation of Salt-induced Kinase, [20, 21], and conversely, mice lacking LKB1 display elevated hepatic gluconeogenesis (Shan 2016), driven by amino acid catabolism [19]. Despite its established involvement in whole-body homeostasis and gluconeogenesis, it is not clear how LKB1 loss of function results in such a severe metabolic phenotype.

Here, we show that hepatocyte-specific deletion of LKB1 (KO) leads to a prolonged fasting response resulting in a severe wasting phenotype. At the transcriptional level, liver from KO mice displayed upregulated glucagon-induced genes throughout the hepatic lobule, many of which are restricted to PP regions in wild type (WT) mice. Our results show that LKB1 fine-tunes glucagon signals across the hepatic lobule to balance the spatial extent of the fasting response which is critical for whole-body energy homeostasis. These findings further understanding of hepatic physiology and factors regulating liver zonation.

Material and methods

Reagents

Primary antibodies for western blots: LKB1 (3047S, 1:1000; Cell Signaling Technologies); β-tubulin (2146S, 1:1000; Cell Signaling Technologies); phosphoenolpyruvate carboxykinase 1 (PCK1 [ab28455], 1:1000, Abcam); Glyceraldehyde 3-phosphate dehydrogenase (GAPDH, 2118S, 1:1000; Cell Signaling Technologies). Secondary antibodies used: goat anti-mouse IgG, HRP conjugate (12-349, 1:1000, Millipore Sigma), goat anti-rabbit IgG, HRP conjugate (12-348, 1:1000, Millipore Sigma).

Primary antibodies used for immunofluorescence: Alexa Fluor 421, 647- conjugated CD324 (E-cadherin, 1:50, added during secondary antibody incubation, BioLegend); and Cytochrome P450 2E1 (Cyp2e1, ab28146, 1:200, Abcam). Secondary antibodies

were purchased from Jackson ImmunoResearch. Alexa Fluor 647- phalloidin (1:200, Invitrogen) to label actin.

Mice

The study was conducted according to the animal protocols approved by the National Cancer Institute (NCI)-Bethesda Animal Care and Use Committee (ACUC) protocol number LCMB037. C57BL/6J and LKB1 fl/fl mice (Jackson Laboratories) with free access to water and a standard diet (NIH-31 Open Formula, Envigo). Experiments were done during the light cycle.

LKB1 KO

LKB1 fl/fl were injected adeno-associated virus 8 (AAV8; 1.25 x 10^{11} GC) expressing Cre recombinase under the hepatocyte thyroid hormone-binding globulin (TBG) promoter (Vector Biolabs). AAV8 was delivered by tail vein injection to LKB1 fl/fl and C57BL/6J at 6 weeks of age and analyzed 4 weeks after. As expected, male mice had higher body weights and lower fat mass than female mice (data not shown). However, since no sex by KO interaction was observed for any measurement, male and female data was combined.

Liver lysate and western blots

Livers were snap-frozen in liquid nitrogen. Fifty miligram of tissue was homogenized in 3 mL RIPA buffer (Invitrogen) with 1X Halt protease and phosphatase inhibitor (Invitrogen). Homogenate was centrifuged at 13,000 rpm for 20 minutes at 4°C, and the supernatant was collected. Protein concentration was determined using a Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific). Lysates were boiled and 20-60μg of protein was loaded onto Mini-PROTEAN® TGX™ Precast Gels (Bio-Rad). SDS-PAGE gels ran for 50 minutes at 125 V. Trans-Blot Turbo Mini 0.2 μm Nitrocellulose or polyvinylidene difluoride (PVDF) Transfer Packs and protein transfer onto membranes was achieved using the low or mixed molecular weight (MW) preprogrammed protocol in a Trans-Blot Turbo transfer system (Bio-Rad). Membranes were blocked for 1 hour at room temperature in 5% milk or 5% BSA (Millipore Sigma) in 1X Tris-buffered saline + 0.1% Tween 20 (TBST). Membranes were rinsed once with 1X TBST and primary antibodies diluted in 5% BSA in 1X TBST were added at desired concentrations. Blots were incubated in primary antibodies overnight at 4°C.

The primary antibody was removed, membranes were washed 3 times in 1X TBST, and secondary antibodies diluted in 5% milk or BSA in 1X TBST were added. Blots were incubated in the secondary antibodies for 1 hour at room temperature. After incubation, membranes were rinsed 3 times with 1X TBST and the blots were incubated in Clarity ECL western blot substrate solution (Bio-Rad) for 5 minutes. Membranes were imaged using the ChemiDoc Imaging System (Bio-Rad).

Liver tissue fixation and immunofluorescence staining

Mice were anesthetized with intraperitoneally (i.p.) injection of 250 mg/kg xylazine and 50 mg/kg ketamine. The liver was fixed by transcardial perfusion of ice-cold PBS for 2 minutes followed by ice-cold 4% paraformaldehyde (PFA) in PBS at a rate of 5 mL/min. Livers were harvested and stored in 4% PFA in PBS overnight followed by sectioning using a Leica Vibratome (50-75 µm thick). Sections were blocked with 0.5% Triton X-100 and 20% FBS in PBS for 1 hour at room temperature. Sections were incubated with the primary antibody for 1-3 days at 4°C. Tissues were washed in PBS for 1 hour and incubated with secondary antibodies overnight at 4°C. Sections were mounted with 80% glycerol in PBS.

Histology

Paraffin-embedding, sectioning and histological staining were done by Histoserv, Inc. Histology slides were scanned and evaluated by the Molecular Pathology Unit National Cancer Institute (NCI).

Glucose and insulin tolerance tests

Fasted mice (6h) were i.p injected with 0.75 units/kg insulin (Humulin R, Lilly Pharmaceuticals) or 1 g/kg 50% Dextrose (Hospira), for insulin and glucose tolerance tests, respectively. Blood was collected from the tail vein and glucose was measured using testing strips (Ascencia Diabetes Care).

Metabolic evaluation

Measurements were carried on *ad libitum* fed and individually housed mice by the Mouse Metabolism Core at NIDDK.

Weight and body composition analysis. To calculate the change in body weight (BW), the final BW was subtracted from the initial BW and divided by the initial BW.

Body composition measurements using an Echo MRI (EchoMRI LLC, Houston, TX). Specifically, the fat and lean mass of the mice was measured during their natural fasting period (1:00 pm).

Food intake and energy balance analysis. Food was weighed weekly. Food intake was converted to kcal consumed/mouse/day using the energy density of the standard diet (3.1 kcal/g). Energy balance was calculated using the method described in (Ravussin et al., 2013). Briefly, the fat mass, lean mass, and body weight measurements at 4 weeks following Cre injection were subtracted from measurements taken the previous week. The difference in fat mass and lean mass values from week 3 to 4 converted to kcal (fat= 9 kcal/g, lean= 1 kcal/g). The energy gain in the body (kcal) was obtained by adding fat and lean mass values. Finally, the energy expenditure was calculated by subtracting the accumulated energy in the body from the food intake of week 4.

Serum analysis. Elisa kits were used according to manufacturer's instructions to measure β-hydroxybutyrate (K632, Biovision), FGF21 (EZRMFGF21-26K, EMD Millipore), and glucagon (10-1281-01, Mercodia).

Confocal microscopy and image processing

Images were acquired using a Leica SP8 inverted confocal laser scanning microscope using 63X oil objective 1.4 NA. E-cadherin staining was used to identify PP regions of the hepatic lobule. Images were cropped contrast enhanced, background subtracted, a median filter was applied using ImageJ. Mean gray values were measured for PP and PC regions, and ratios of PC over PP were determined.

Intravital microscopy

Mice were anesthetized with 2% isoflurane administered through a nose cone. Prior to surgery, mice were injected with heparin diluted 1:10 in 0.9% (w/v) NaCl to prevent liver ischemia. Abdominal fur was shaved, and left lobe was exposed using surgical scissors. The mouse was positioned on the stage of a Leica SP8 inverted confocal laser scanning microscope fitted with a custom stage insert. Transparent ultrasound transmission gel (Parker Aquasonic 100) was used around the incision to maintain the tissue from drying.

Glucose uptake assay. The microscope stage was heated to 34°C and a 20X objective, 0.75 NA was used for imaging. Eighteen mg/kg of 2,000,000 MW

tetramethylrhodamine-dextran (Invitrogen) was injected retro-orbitally to label the sinusoids. IVM time-lapse were acquired every 2.58 seconds for 10 minutes (512 x 512 pixels and a pixel size of 1.14 µm). After image acquisition had begun, the fluorescent glucose analog 2-(N-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)Amino)-2-Deoxyglucose (2-NBDG; 6.83 µg/kg BW, Invitrogen) dissolved in 50% dextrose (1 g/kg BW) was injected retro-orbitally. Quantification of glucose uptake and retention was performed as previously described [22].

RNA isolation

Three mice per group were sacrificed and the left lobe was snap-frozen in liquid nitrogen. ~20 mg per sample was homogenized using a TissueLyser II (Qiagen) and total RNA using the Qiagen RNeasy Mini Kit (#74104). The RNA yield was determined using a High-sensitivity RNA ScreenTape Analysis (Agilent) on an Agilent TapeStation Instrument. Samples with a RIN > 8.0 were sent for bulk mRNA sequencing.

Bulk mRNA sequencing

Sequencing was done by the CCR Sequencing Facility at the Frederick National Laboratory for Cancer Research. TruSeq mRNA library kit (Illumina) was used to prepare the mRNA library following the manufacturer's instructions. 200ng of total RNA was used as the input to an mRNA capture with oligo-dT coated magnetic beads. The mRNA was fragmented, and then a random-primed cDNA synthesis was performed. To prepare the library, the resulting double-strand cDNA was used as the input to a standard Illumina library prep with end-repair, adapter ligation and PCR amplification being performed. The final purified product was then quantitated by qPCR before cluster generation and sequencing. Six mRNA-Seq samples were pooled and sequenced on NextSeq using paired-end sequencing and a read length of 76 (2 x 76 cycles).

Sequencing data analysis

Raw data files were processed using the HiSeq Real Time Analysis software (RTA 2.11.3). Files were demultiplexed and conversion of binary base calls and qualities into fastq format was performed using the Illumina bcl2fastq2.17. Cutadapt (version 1.18) was utilized to trim adapters and low-quality bases from the sequencing reads and the trimmed reads were mapped to mouse reference genome NCBI m37/mm9

and Gencode annotation M9 using STAR (version 2.7.0f) with two-pass alignment option. RSEM (version 1.3.1) was used for gene and transcript quantification.

Raw counts files were imported into the NIH Data Analysis Portal (NIDAP) for analysis in collaboration with the CCR Collaborative Bioinformatics Resource (CCBR). Genes with low counts were removed and the data was normalized by quantile normalization. Batch correction was performed on the data to account for sex differences in the samples. Differential expression of genes (DEG) and gene set enrichment analysis (GSEA) were performed. A threshold of FDR < 0.05, FC > 1.5, FC < -1.5 was utilized to determine significance. Heatmaps were generated using hierarchical clustering on the NIDAP platform. The Venn diagrams were generated using Venny 2.1 (Oliveros, 2007). The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus [23] and are accessible through GEO Series accession number GSE192404 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE192404).

Statistical Analysis

A minimum of 3 mice were averaged per condition. As no sex by LKB1 KO interaction were observed for any of the data, male and female values were pooled. Unpaired t-tests were performed for data with equal variance and a Welch's t-test was performed for data with unequal variance. For comparing WT and KO mice within PP or PC regions, a two-way ANOVA with Bonferroni post hoc was used, or comparing across time, a two-way repeated measures ANOVA was used. All data are shown as mean ± SEM. Statistical significance was determined by a p-value < 0.05 using GraphPad Prism (version 8.4.2). For RNA seq data, genes that had an FDR < 0.05, FC < -1.5, and FC > 1.5 were considered to be regulated by LKB1 depletion.

Results

LKB1 KO mice are in a prolonged fasting state resulting in a wasting phenotype.
Liver-specific LKB1 KO mice were generated via tail vein injection of AAV8 Cre recombinase with a TGB promoter in LKB1 floxed mice at 6 weeks of age and analyzed 4 weeks after. Over 90% reduction in LKB1 protein was achieved (Sup Fig 1A) with some residual LKB1 detected, likely originating from non-parenchymal cells. As previously reported in a model where LKB1 was depleted in neonates [17], pAMPK levels were significantly reduced in KO livers (Sup Fig 1B). Overall, KO mice were smaller than WT controls which was also reflected by significant weight loss over the

course of four weeks (Fig 1A and B). Weight loss occurred despite increased food intake and increase in cumulative energy balance, indicating poorer energy efficiency in KO mice (Fig 1C and D). Weight loss in KO mice included both fat and lean mass (Fig 1E and F). Additionally, substantial reduction in glycogen storage was observed in liver sections from KO mice, primarily around PC and mid-lobular regions (Fig 1G and Sup Fig 1E). The wasting phenotype observed is consistent with previous whole body [18] and liver-specific LKB1 KO models [16, 17, 19]. Previously, we have shown that deletion of LKB1 at birth via Albumin Cre excision displayed a similar wasting phenotype to the one we describe here using AAV8 Cre injection in mature mice [17]. This result demonstrates a vital role for liver-specific LKB1 in maintaining metabolic homeostasis in both neonate and adult mice.

We next examined whether KO mice have elevated markers of prolonged fasting/starvation. Ketone bodies are produced in the liver during extended fasting, starvation or exercise (Rui, 2014; White and Venkatesh, 2011) through fatty acid oxidation (Puchalska and Crawford, 2017). Indeed, serum level of the ketone body β-hydroxybutyrate was increased in KO mice (Fig 1H). In addition, we measured serum levels of starvation-associated hormone fibroblast growth factor 21 (FGF21) [24] and found it was higher in KO mice (Fig 1I). FGF21 levels have been shown to promote lipolysis in diet-induced obese mice, leading to weight loss [25]. Consistently, serum free fatty acid levels were also elevated in KO mice (Fig 1J). Despite these elevated markers of prolonged fasting serum glucagon levels in KO mice were significantly reduced (Sup Fig 1C), likely driven by the hyperglycemia in KOs and shown here (Fig 2A), as previously reported [20, 21]. On the other hand, insulin levels were unchanged (Sup Fig 1D). Collectively, these results demonstrate elevated markers indicative of prolonged fasting, consistent with the observed wasting phenotype.

Glucose retention is reduced in LKB1 KO hepatocytes.

LKB1 has a critical role in glycemic control [20, 21]. Indeed, KO mice were severely hyperglycemic (Fig 2A) and displayed glucose intolerance (Fig 2A and B), but insulin-sensitive, indicating peripheral tissues were not affected by depletion of LKB1. This disruption in glycemic state and the non-uniform effect on glycogen storage (Fig 1G), suggests increased hepatic glucose efflux. To test this, we developed an assay to measure hepatic glucose mobilization in live anesthetized mice using time-lapse IVM

[22] To this aim, we traced the fluorescent glucose analog 2-(*N*-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)Amino)-2-Deoxyglucose (2-NBDG) (Fig 2D-F). Using this approach, the rate of glucose uptake and retention in PP and PC hepatocytes was measured. In WT mice, 2-NBDG signal appeared in portal vein areas within seconds of injection and was taken up by hepatocytes after 6 minutes (Fig 2D, Movie S1). Although serum glucose concentrations are reportedly higher in the portal vein [26], we found no difference in the rate of glucose uptake between PP and PC hepatocytes in WT mice (Fig 2E). However, in KO mice, the rate of uptake in both PP and PC hepatocytes was significantly lower compared to WT and accumulation of 2-NBDG signal was not observed in PC regions (Fig 2E). Additionally, we compared the ability of the hepatocytes to retain glucose by calculating the ratio of 2-NBDG fluorescent signal in hepatocytes over sinusoids. Hepatic glucose retention was significantly decreased in KO mice in PP and PC regions consistent with robust glucose efflux (Fig 2F).

Upregulation of PP genes and pathways in LKB1 KO mice

To better understand the metabolic changes in LKB1-deficient livers, we performed RNA sequencing and examined its impact on gene expression. Out of 11,669 identified genes, 1,635 were differentially expressed in the KO liver compared to WT, with roughly a 1:1 ratio of upregulated to downregulated genes (Fig 3A). Gene set enrichment analysis (GSEA) and Ingenuity Pathway Analysis (IPA) were used to analyze the differentially expressed genes (DEG). We found 699 significantly enriched pathways, a subset of which is shown in Fig 3B. Interestingly, many pathways that are associated with PP functions including mitochondria oxidative phosphorylation (Sup Fig 2A-C), TCA cycle, the previously reported gluconeogenesis [21] and amino acids catabolism [19] were enriched in KO hepatocytes (Fig 3B). Pathways associated with PC functions, on the other hand, such as lipogenesis and bile acid metabolism, were downregulated (Fig 3B) suggesting a bias towards PP genes in the KO liver. To investigate this idea further, we used a list of PP and PC genes compiled by Cheng *et al.* [8] and compared them with the DEG identified in the LKB1 KO mice. Indeed, out of 270 PP and 461 PC genes, 89 and 157 were regulated in the LKB1 KO livers, respectively. Strikingly, within these regulated genes, there was substantial upregulation of PP and downregulation of PC genes (Fig 3C), suggesting the involvement of LKB1 in the spatial division of metabolic functions.

Given these changes in zone-restricted gene expression in KO livers, immunofluorescence localization studies were performed to examine the spatial distribution of selected enzymes in liver sections. Based on gene expression, gluconeogenesis is restricted to PP hepatocytes [10]. Consistently, protein expression of the rate-limiting enzyme Phosphoenolpyruvate Carboxykinase 1 (PCK1) was confined to PP hepatocytes in WT (Fig 3D and E). However, in the KO lobule, its distribution was uniform across both PP and PC hepatocytes (Fig 3D and E). Similar results were obtained with the PP urea cycle enzyme Arginase 1, (Sup Fig 2D and E). The spatial localization of PC enzymes was also investigated, including Cyp2e1 which is involved in xenobiotic metabolism. Cyp2e1 is restricted to PC hepatocytes in the WT, and this pattern was lost in KO due to reduced expression levels and limited distribution around the central vein (Fig 3F and G). Notably, the distribution of E-cadherin, which was used to identify PP regions, expanded into PC regions in KO sections reinforcing that idea of "periportalization" of the hepatic lobule (Fig 3F, compare WT and KO). Similar results were obtained with the PC lipogenic enzyme acetyl-CoA carboxylase 1 (ACC1), (Sup Fig 1F and G). Overall, in the absence of LKB1, the liver defaults to a PP metabolic program that overrides PC-related gene expression and functions. This loss in zonation indicates that LKB1 signaling is important in maintaining hepatocyte heterogeneity.

LKB1 regulates glucagon target genes, but not Wnt/b-catenin

To understand how LKB1 affects liver zonation IPA analysis was performed to identify potential regulators based on the DEG. Interestingly, glucagon was identified as the top upstream activator of LKB1 (Fig 4A). Since glucagon controls liver zonation by promoting the expression of PP genes [8] we next investigated the relationship between LKB1- and glucagon-regulated genes. To this aim, DEG identified in LKB1 KO was compared with those identified in glucagon KO mice [8]. Of the 297 DEGs identified in glucagon KO livers, 40% overlapped with DEGs in LKB1 KO (Fig 4B). Interestingly, the overlapping genes inversely correlated (Fig 4C) suggesting that LKB1 has an overall inhibitory impact on glucagon induced gene expression. In glucagon KO mice, PC genes were upregulated [8], whereas, in LKB1 KO, despite reduced glucagon levels (Sup 1C), PP genes were enhanced, thus reinforcing the notion that LKB1 negatively regulates glucagon-induced transcriptional changes. Similar analysis of Wnt target genes revealed that out of the 1003 DEG, 20% of the LKB1 KO overlapped (Fig 4D).

However, there was no correlation in expression between the overlapping genes (Fig 4E). These results suggest that upregulation in PP and decreased PC gene expression in LKB1 deficient hepatocytes is driven by uninhibited glucagon signaling rather than changes to Wnt/b-catenin. Taken together, our data suggests that glucagon is an activator of LKB1 which subsequently suppresses glucagon-induced gene expression. This critical negative feedback loop of glucagon signaling prevents continuation of the fasting response and whole body wasting.

Glucagon is involved in the fasting response through activation of transcription factors. Initially, glucagon stimulates cAMP/CREB pathway which leads to expression of additional transcription factors and a second wave of gene expression facilitating the temporal regulation of the fasting response [6]. Therefore, expression levels of transcription factors involved in the fasting response in the LKB1 KO were examined. Significant increases in Nr4a1 (6-fold) and Nr4a2 (16-fold) were found in KO mice. These factors are involved in upregulating gluconeogenesis and the expression of FGF21 [27, 28]. Additionally, increased expression of Ppargc1a (7-fold) was observed which was previously described in LKB1 KO mice [20], and is involved in FAO and ketogenesis [29]. These transcription factors are part of the delayed response to glucagon and may explain the exacerbated and prolonged fasting phenotype (Fig 1). Lastly, Fig 4G summarizes our findings in a model. We identified a negative feedback loop in which LKB1 keeps the glucagon-mediated fasting response under control. Depletion of LKB1 results in unrestrained activation of glucagon mediated pathways including gluconeogenesis, glucose efflux, ketogenesis and lipolysis. These zone-restricted functions extend across the hepatic lobule in mutant mice leading to loss of liver zonation. These findings explain the severe wasting phenotypes resulting in overall weight loss, inability to store fuels and metabolic inefficiency. This study highlights the possibility that glucagon-mediated fasting is not only transcriptionally controlled in time [6], but also spatially regulated by LKB1 across the hepatic lobule.

Discussion

Fasting is an essential adaptive response to simultaneously mobilize nutrients as a fuel source while conserving energy expenditure. During diseases such as NAFLD and type 2 diabetes, an inability to shut down this fasting response results in persistent gluconeogenesis and lipolysis, despite adequate nutrient consumption. This excessive

mobilization of fuels worsens the inability to arrest the fasting response, and further exacerbates disease progression. Here, we show that LKB1 is vital for the inhibition of the fasting response. Our findings that LKB1 KO mice have a wasting phenotype (Fig 1A-F) despite increased food intake (Fig 1C) and circulating substrates in the blood (Fig 1 and 2A) are indicative of disruption in the homeostatic control of energy balance and suggestive of overall metabolic inefficiency. Energy expenditure allows the evaluation of overall metabolic efficiency by measuring the ability to convert ingested energy into fat and protein [30]. In an attempt to maintain bodyweight during fasting, energy expenditure is decreased by reducing physical activity [31] and diet-induced thermogenesis [32]. Supporting this, Just and colleagues previously showed that liver specific LKB1 KO had decreased locomotor activity (Just et al 2020). As expected, we measured significantly higher energy expenditure in KO mice despite being in a phenotypical fasting state (Fig 5I), reflecting metabolic inefficiency in LKB1 KO mice. Although we do not know the cause for this inefficiency, it is possible that due to the severe hyperglycemia in KO mice, the diet-induced thermogenesis is increased.

LKB1 is a key regulator of glucose homeostasis [21]. Hepatocytes shift between glucose uptake for storage and efflux during fed and fasted states, respectively. To measure the changes in glucose uptake and release, we developed an IVM-based assay to measure glucose mobilization in space and time [22]. We measured markedly reduced glucose uptake and retention in KO hepatocytes indicating they are in constitutive efflux state (Fig 2D-F). These results are consistent with the severe hyperglycemia and glucose intolerance (Fig 2A and B), reduced glycogen storage (Fig 1G), and upregulation of gluconeogenesis genes (Fig 3B and D-E). Interestingly, Berndt and colleagues used modeling to propose that PP hepatocytes release and PC hepatocytes take up glucose [33]. However, *in vivo*, we did not detect any difference in the rate of glucose uptake and/or retention between PP and PC hepatocytes in WT or KO (Fig 3A-C). Similar kinetics in PP and PC are likely achieved through differential expression of high- and low-affinity glucose transporters across the lobule [10] and redistribution of these transporters when glucose availability shifts. Our results underscore the importance of measurements of physiological parameters in the intact tissue.

This unique spatial organization of hepatocytes (Sup Fig 1E) results in the formation of nutrient, oxygen, hormone, and morphogen gradients, which in turn, shape hepatocytes' gene expression. Indeed, a recent study employing single-cell RNA sequencing demonstrated that 50% of liver genes are zonated [34]. Key regulators of this zonation were identified and include Wnt/b-catenin, glucagon, hypoxia, H-ras, and growth and thyroid hormones [10]. However, these do not account for all the zonated genes, highlighting the existence of additional regulators of zonation. Our findings show that LKB1 plays a significant role in liver zonation. In the absence of LKB1, the spatial division of hepatic metabolism is abolished. Instead, we observed a striking upregulation of PP-associated genes, expansion of their spatial distribution and downregulation of PC-related genes (Fig 3C-G and Sup Fig 2D-G). In a previous study, Cheng *et al*. showed that glucagon oppose Wnt/b -catenin activity by inducing PP gene expression [8]. Our analysis identified glucagon as a probable upstream activator of LKB1. Furthermore, we show a strong inverse correlation between glucagon and LKB1 regulated genes. This suggests a negative feedback loop between glucagon and LKB1, leading to fine tuning of hepatocyte's response to glucagon. Indeed, Yang et al. showed that glucagon receptor ablation led to downregulation of PP associated pathways and upregulation of PC ones, reinforcing the notion that LKB1 influences glucagon signaling [35]. This type of regulation is reminiscent of the balancing act between Wnt/b-catenin and its negative regulator Adenomatous Polyposis Coli Protein (APC) in the regulation of PC genes [36, 37]. Previous reports also indicated a cross talk between LKB1 and Wnt signaling in different models [38-40]. However, our data shows no correlation between the expression of Wnt and LKB1 regulated genes (Fig 2D and E) suggesting that the effects of LKB1 on zonation are largely independent of Wnt. While mechanistic studies to elucidate how LKB1 counterbalances glucagon are warranted, we propose that hepatocyte nutrient sensing through LKB1, together with extrinsic factors such as glucagon, orchestrate the patterned gene expression observed in the hepatic lobule.

Our study demonstrates that loss of LKB1 in the liver leads to a prolonged fasting response, which contribute to weight loss, decreased glycogen, lean mass, and adipose tissue reserve and an overall decrease in metabolic efficiency (Fig 1). For the first time, we identify LKB1 as a vital inhibitor of the glucagon-mediated fasting response to prevent a wasting phenotype. Attenuation of the glucagon response via

GLP-1 agonists has become one of the most effective drug targets for obesity, type 2 diabetes, and NAFLD, therefore, increasing our understanding the regulatory mechanisms behind glucagon signaling is vital to improve new pharmacological therapeutics [3, 5]. Here, the wasting phenotype caused by uncontrolled glucagon signaling was observed alongside complete ablation of metabolic zonation in the hepatic lobule, raising the provocative possibility that the fasting response is not only regulated transcriptionally in time [6] but also in space within the hepatic lobule. This could be critical for the organism's survival in sustaining the fasting response if nutrients are unavailable. The lobule's anatomy allows for increased capacity of PP-like hepatocytes and glucagon-LKB1 signaling axis may be at the nexus of the spatio-temporal regulation of the fasting response. Given the conserved role of LKB1 in cell polarity, it is possible it is also involved in the organization of the hepatic lobule which represents polarity in the tissue scale. Mechanistically, it is unclear how these zone-specific changes influence metabolic efficiency seen in LKB1 KO mice. Future studies that attempt to restore liver zonation and rescue the wasting phenotype in LKB1 KO mice would identify its disruption as the primary cause of metabolic inefficiency in these mice. Nonetheless, this study reveals a new mechanism by which hepatic metabolic compartmentalization is regulated by nutrient sensing.

Contact information

Natalie Porat-Shliom, Center for Cancer Research, National Cancer Institute. Building 10, Room 12C207 MSC 1919, Bethesda, MD 20892. Phone: 240-760-6847. Email: natalie.porat-shliom@nih.gov

Abbreviations

LKB1	Liver Kinase B1
NAFLD	Non-Alcoholic Fatty Liver Disease
GLP-1	Glucagon-like peptide 1
PP	Periportal
PC	Pericentral
AMPK	AMP-activated Protein Kinase
ARK	AMPK-related
KO	Knock out
WT	Wild Type
PCK1	phosphoenolpyruvate carboxykinase 1
GAPDH	Glyceraldehyde 3-phosphate dehydrogenase
AAV8	Adeno-Associated Virus 8
TBS	Tris-buffered saline

TBST	Tris-buffered saline triton
PBS	Phosphate-buffered saline
FBS	Fetal Bovine serum
PFA	Paraformaldehyde
I.p.	intraperitoneally
BW	Body weight
FGF21	Fibroblast growth factor 21
ACC1	Acetyl-CoA carboxylase
NaCl	Sodium Chloride
2-NBDG	2-(N-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)Amino)-2-Deoxyglucose
PCR	Polymerase chain reaction
qPCR	Quantitative polymerase chain reaction
DEG	Differential expression of genes
GSEA	Gene set enrichment analysis
NIDAP	NIH Data Analysis Portal
IPA	Ingenuity pathway analysis
FC	Fold change
FDR	False discovery rate

Financial Support

This work was supported by the Intramural Research Program at the NIH, National Cancer Institute (1ZIABC011828). The authors have no conflicts to report.

Acknowledgment

We thank members of the Porat-Shliom lab, Drs. Will Printz, Win Arias and Julie Donaldson for critical reading of the manuscript.

Figure legend

Figure 1. **Characterization of changes in whole-body energy homeostasis in WT and KO mice.** A) Representative image of WT and KO mice. B) Percent change in body weight. C) Daily food intake. D) Cumulative energy balance per mouse per day. E) Final fat mass and F) final lean mass. G) PAS staining (glycogen) in liver sections from WT and KO. *PP; **PC. H) Serum levels of b-hydroxybutyrate. I) Serum levels of FGF21. J) Serum levels of free fatty acids. In all the above experiments, N = 16-18 mice per measurement. Data are shown as mean ± SEM. P-values were calculated using an unpaired Student's t-test. *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$, ****$p < 0.0001$, and ns= not significant. PP = periportal, PC = pericentral.

Figure 2. **IVM of glucose uptake and retention in WT and KO mice.** A) Fasting blood glucose from WT and KO mice. B) Glucose tolerance test (GTT) and C) insulin tolerance test (ITT) of WT and KO mice. D) Representative time-lapse frames of hepatic glucose uptake *in vivo* using the fluorescent glucose probe 2-NBDG (white). Mice were injected with 2-NBDG approximately 10 seconds into image acquisition. Dextran (red) was used to label the vasculature. Time = minutes: seconds. Dashed white circles indicate portal veins. CV = central vein. E) Quantification of glucose uptake rate (positive pixels/second). F) Quantification of glucose retention shown as the ratio of mean fluorescence intensity of 2-NBDG in hepatocytes to sinusoids in PP hepatocytes and PC hepatocytes. Scale bar= 100 μm. Data are shown as mean $\pm$ SEM. P-values were calculated using an unpaired Student's t-test for fasting glucose, a two-way ANOVA for WT vs KO within PP or PC, or repeated measures two-way ANOVA for WT vs KO across time. N= 3 mice. $*p < 0.05$, $**p < 0.005$, $***p < 0.0005$, and $****p < 0.00001$.

Figure 3. **LKB1 regulates PP genes and function.** A) Volcano plot showing global gene expression from WT and KO livers. N=3 mice per group. B) Enrichment of select upregulated (red) and downregulated (black) pathways in KO mice. C) Gene set enrichment plots showing expression of PP- and PC-associated genes, upper and lower graphs, respectively, in LKB1 KO livers. Gene list from [8] was used for the comparison. D) Immunofluorescence showing the distribution of PCK1 (grey) in WT and KO liver sections. PP regions are labeled with E-cadherin (magenta). E) Quantification of PCK1 distribution is shown as the ratio of PCK1 intensity in PC and PP hepatocytes. F) Immunofluorescence showing the distribution of Cyp2e1 (grey) in WT and KO liver sections. PP regions are labeled with E-cadherin (magenta). G) Quantification of Cyp2e1distribution is shown as the ratio of Cyp2e1 mean grey value in PC and PP. Four-8 PP or PC regions were analyzed per mouse and averaged, n=3 mice per group. Scale bar= 50 μm. Data are shown as mean $\pm$ SEM; p-values were calculated using an unpaired Student's t-test or a Welch's t-test.

Figure 4. **LKB1 negatively regulates glucagon-induced genes, but not Wnt-regulated genes.** A) Upstream regulators of LKB1 identified using Ingenuity Pathway Analysis (IPA). B) Venn diagram of differentially expressed genes in LKB1 KO (current study) and glucagon KO [8] showing 119 overlapping genes. C) The correlation between

LKB1 KO and glucagon KO genes. D) Venn diagram of differentially expressed genes in LKB1 KO (current study), and Wnt-regulated [34] showing 191 overlapping genes. E) The correlation between LKB1 KO and Wnt-regulated genes. F) Fold change of transcription factors in WT versus KO mice. G) Graphical summary of results, showing how loss of LKB1 allows glucagon signaling to cause excessive periportalization of the lobule.

Supplementary Figures

Figure S1. **Confirmation of LKB1 KO.** A) Western blot showing protein levels of LKB1 in WT and KO and quantification. B) Total AMPK and pAMPK (Thr172) and their representative Western blots in WT and KO mice. C) Serum glucagon and D) insulin levels. E) Schematic visualization of the hepatic lobule. Data are shown as mean $\pm$ SEM; p-values were calculated using an unpaired Student's t-test or a Welch's t-test. *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$, and ****$p < 0.00001$.

Figure S2. **Upregulation of PP genes and function in LKB1 KO.** A) Fold change of genes involved in oxidative phosphorylation and mitochondrial transport, and B) their representative heat map in WT and KO mice. C) IPA representation of changes in oxidative phosphorylation genes in KO mice. D) Immunofluorescence showing the distribution of Arginase 1 (grey) in WT and KO liver sections. PP regions are labeled with E-cadherin (magenta). E) Quantification of Arginase 1 distribution is shown as the ratio of Arginase 1 intensity in PC and PP. F) Immunofluorescence showing the distribution of ACC 1 (grey) in WT and KO liver sections. PP regions are labeled with E-cadherin (magenta). G) Quantification of ACC 1 distribution is shown as the ratio of ACC 1 intensity in PC and PP. 4-8 veins were analyzed per mouse and averaged, n=3 mice per group. Scale bar= 50 μm. All data are shown as mean ± SEM, n= 3-5 mice. P-values were calculated using an unpaired Student's t-test (Prism 8.4.2), *p-value<0.05, **p-value=0.005, ***p-value=0.0005.

Supplementary Movies

Movie S1. **IVM of 2-NBDG uptake in WT liver.** Vasculature was labeled by dextran (red) which was injected prior to imaging and 2-NBDG is shown in green, which was injected after the first image was acquired. Images were acquired every 2.5 seconds

for 5 minutes.

Movie S2. **IVM of 2-NBDG uptake in KO liver.** Vasculature was labeled by dextran (red) which was injected prior to imaging and 2-NBDG is shown in green, which was injected after the first image was acquired. Images were acquired every 2.5 seconds for 5 minutes.

References

1. Jiang, G.; Zhang, B. B., Glucagon and regulation of glucose metabolism. *Am J Physiol Endocrinol Metab* **2003,** *284* (4), E671-8.
2. Samuel, V. T.; Shulman, G. I., Nonalcoholic Fatty Liver Disease as a Nexus of Metabolic and Hepatic Diseases. *Cell Metab* **2018,** *27* (1), 22-41.
3. Rosenstock, J.; Wysham, C.; Frias, J. P.; Kaneko, S.; Lee, C. J.; Fernandez Lando, L.; Mao, H.; Cui, X.; Karanikas, C. A.; Thieu, V. T., Efficacy and safety of a novel dual GIP and GLP-1 receptor agonist tirzepatide in patients with type 2 diabetes (SURPASS-1): a double-blind, randomised, phase 3 trial. *Lancet* **2021,** *398* (10295), 143-155.
4. Drucker, D. J., Mechanisms of Action and Therapeutic Application of Glucagon-like Peptide-1. *Cell Metab* **2018,** *27* (4), 740-756.
5. Wilding, J. P. H.; Batterham, R. L.; Calanna, S.; Davies, M.; Van Gaal, L. F.; Lingvay, I.; McGowan, B. M.; Rosenstock, J.; Tran, M. T. D.; Wadden, T. A.; Wharton, S.; Yokote, K.; Zeuthen, N.; Kushner, R. F.; Group, S. S., Once-Weekly Semaglutide in Adults with Overweight or Obesity. *N Engl J Med* **2021,** *384* (11), 989.
6. Goldstein, I.; Hager, G. L., The Three Ds of Transcription Activation by Glucagon: Direct, Delayed, and Dynamic. *Endocrinology* **2018,** *159* (1), 206-216.
7. Ravnskjaer, K.; Madiraju, A.; Montminy, M., Role of the cAMP Pathway in Glucose and Lipid Metabolism. *Handb Exp Pharmacol* **2016,** *233*, 29-49.
8. Cheng, X.; Kim, S. Y.; Okamoto, H.; Xin, Y.; Yancopoulos, G. D.; Murphy, A. J.; Gromada, J., Glucagon contributes to liver zonation. *Proc Natl Acad Sci U S A* **2018,** *115* (17), E4111-E4119.
9. Jungermann, K.; Sasse, D., Heterogeneity of Liver Parenchymal-Cells. *Trends Biochem Sci* **1978,** *3* (9), 198-202.
10. Ben-Moshe, S.; Itzkovitz, S., Spatial heterogeneity in the mammalian liver. *Nat Rev Gastroenterol Hepatol* **2019,** *16* (7), 395-410.
11. Cunningham, R. P.; Porat-Shliom, N., Liver Zonation - Revisiting Old Questions With New Technologies. *Front Physiol* **2021,** *12*, 732929.
12. Kietzmann, T., Metabolic zonation of the liver: The oxygen gradient revisited. *Redox Biology* **2017,** *11*, 622-630.
13. Preziosi, M.; Okabe, H.; Poddar, M.; Singh, S.; Monga, S. P., Endothelial Wnts regulate beta-catenin signaling in murine liver zonation and regeneration: A sequel to the Wnt-Wnt situation. *Hepatol Commun* **2018,** *2* (7), 845-860.
14. Fu, D.; Wakabayashi, Y.; Ido, Y.; Lippincott-Schwartz, J.; Arias, I. M., Regulation of bile canalicular network formation and maintenance by AMP-activated protein kinase and LKB1. *Journal of Cell Science* **2010,** *123* (19), 3294-3302.

15. Fu, D.; Wakabayashi, Y.; Lippincott-Schwartz, J.; Arias, I. M., Bile acid stimulates hepatocyte polarization through a cAMP-Epac-MEK-LKB1-AMPK pathway. *P Natl Acad Sci USA* **2011,** *108* (4), 1403-1408.
16. Homolya, L.; Fu, D.; Sengupta, P.; Jarnik, M.; Gillet, J. P.; Vitale-Cross, L.; Gutkind, J. S.; Lippincott-Schwartz, J.; Arias, I. M., LKB1/AMPK and PKA control ABCB11 trafficking and polarization in hepatocytes. *PLoS One* **2014,** *9* (3), e91921.
17. Porat-Shliom, N.; Tietgens, A. J.; Van Itallie, C. M.; Vitale-Cross, L.; Jarnik, M.; Harding, O. J.; Anderson, J. M.; Gutkind, J. S.; Weigert, R.; Arias, I. M., Liver Kinase B1 Regulates Hepatocellular Tight Junction Distribution and Function In Vivo. *Hepatology* **2016,** *64* (4), 1317-1329.
18. Shan, T.; Xiong, Y.; Kuang, S., Deletion of Lkb1 in adult mice results in body weight reduction and lethality. *Sci Rep* **2016,** *6*, 36561.
19. Just, P. A.; Charawi, S.; Denis, R. G. P.; Savall, M.; Traore, M.; Foretz, M.; Bastu, S.; Magassa, S.; Senni, N.; Sohier, P.; Wursmer, M.; Vasseur-Cognet, M.; Schmitt, A.; Le Gall, M.; Leduc, M.; Guillonneau, F.; De Bandt, J. P.; Mayeux, P.; Romagnolo, B.; Luquet, S.; Bossard, P.; Perret, C., Lkb1 suppresses amino acid-driven gluconeogenesis in the liver. *Nat Commun* **2020,** *11* (1), 6127.
20. Shaw, R. J.; Lamia, K. A.; Vasquez, D.; Koo, S. H.; Bardeesy, N.; Depinho, R. A.; Montminy, M.; Cantley, L. C., The kinase LKB1 mediates glucose homeostasis in liver and therapeutic effects of metformin. *Science* **2005,** *310* (5754), 1642-6.
21. Patel, K.; Foretz, M.; Marion, A.; Campbell, D. G.; Gourlay, R.; Boudaba, N.; Tournier, E.; Titchenell, P.; Peggie, M.; Deak, M.; Wan, M.; Kaestner, K. H.; Goransson, O.; Viollet, B.; Gray, N. S.; Birnbaum, M. J.; Sutherland, C.; Sakamoto, K., The LKB1-salt-inducible kinase pathway functions as a key gluconeogenic suppressor in the liver. *Nat Commun* **2014,** *5*, 4535.
22. Stefkovich, M. L.; Kang, S. W. S.; Porat-Shliom, N., Intravital Microscopy for the Study of Hepatic Glucose Uptake. *Curr Protoc* **2021,** *1* (5), e139.
23. Edgar, R.; Domrachev, M.; Lash, A. E., Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* **2002,** *30* (1), 207-10.
24. Lin, X. L.; Liu, Y. B.; Hu, H. J., Metabolic role of fibroblast growth factor 21 in liver, adipose and nervous system tissues. *Biomed Rep* **2017,** *6* (5), 495-502.
25. Inagaki, T.; Dutchak, P.; Zhao, G.; Ding, X.; Gautron, L.; Parameswara, V.; Li, Y.; Goetz, R.; Mohammadi, M.; Esser, V.; Elmquist, J. K.; Gerard, R. D.; Burgess, S. C.; Hammer, R. E.; Mangelsdorf, D. J.; Kliewer, S. A., Endocrine regulation of the fasting response by PPARalpha-mediated induction of fibroblast growth factor 21. *Cell Metab* **2007,** *5* (6), 415-25.
26. Pagliassotti, M. J.; Myers, S. R.; Moore, M. C.; Neal, D. W.; Cherrington, A. D., Magnitude of Negative Arterial-Portal Glucose Gradient Alters Net Hepatic Glucose Balance in Conscious Dogs. *Diabetes* **1991,** *40* (12), 1659-1668.
27. Pei, L.; Waki, H.; Vaitheesvaran, B.; Wilpitz, D. C.; Kurland, I. J.; Tontonoz, P., NR4A orphan nuclear receptors are transcriptional regulators of hepatic glucose metabolism. *Nat Med* **2006,** *12* (9), 1048-55.
28. Min, A. K.; Bae, K. H.; Jung, Y. A.; Choi, Y. K.; Kim, M. J.; Kim, J. H.; Jeon, J. H.; Kim, J. G.; Lee, I. K.; Park, K. G., Orphan nuclear receptor Nur77 mediates fasting-induced hepatic fibroblast growth factor 21 expression. *Endocrinology* **2014,** *155* (8), 2924-31.

29. Settembre, C.; De Cegli, R.; Mansueto, G.; Saha, P. K.; Vetrini, F.; Visvikis, O.; Huynh, T.; Carissimo, A.; Palmer, D.; Klisch, T. J.; Wollenberg, A. C.; Di Bernardo, D.; Chan, L.; Irazoqui, J. E.; Ballabio, A., TFEB controls cellular lipid metabolism through a starvation-induced autoregulatory loop. *Nature Cell Biology* **2013,** *15* (6), 647-+.

30. Brooks, S. P. J., Chapter 9 Fasting and refeeding: Models of changes in metabolic efficiency. In *Cell and Molecular Response to Stress*, Storey, K. B.; Storey, J. M., Eds. Elsevier: 2001; Vol. 2, pp 111-127.

31. Farooq, A.; Chamari, K.; Sayegh, S.; El Akoum, M.; Al-Mohannadi, A. S., Ramadan daily intermittent fasting reduces objectively assessed habitual physical activity among adults. *Bmc Public Health* **2021,** *21* (1).

32. Minghelli, G.; Schutz, Y.; Whitehead, R.; Jequier, E., Seasonal changes in 24-h and basal energy expenditures in rural Gambian men as measured in a respiration chamber. *Am J Clin Nutr* **1991,** *53* (1), 14-20.

33. Berndt, N.; Horger, M. S.; Bulik, S.; Holzhutter, H. G., A multiscale modelling approach to assess the impact of metabolic zonation and microperfusion on the hepatic carbohydrate metabolism. *PLoS Comput Biol* **2018,** *14* (2), e1006005.

34. Halpern, K. B.; Shenhav, R.; Matcovitch-Natan, O.; Toth, B.; Lemze, D.; Golan, M.; Massasa, E. E.; Baydatch, S.; Landen, S.; Moor, A. E.; Brandis, A.; Giladi, A.; Avihail, A. S.; David, E.; Amit, I.; Itzkovitz, S., Single-cell spatial reconstruction reveals global division of labour in the mammalian liver. *Nature* **2017,** *542* (7641), 352-356.

35. Yang, J.; MacDougall, M. L.; McDowell, M. T.; Xi, L.; Wei, R.; Zavadoski, W. J.; Molloy, M. P.; Baker, J. D.; Kuhn, M.; Cabrera, O.; Treadway, J. L., Polyomic profiling reveals significant hepatic metabolic alterations in glucagon-receptor (GCGR) knockout mice: implications on anti-glucagon therapies for diabetes. *BMC Genomics* **2011,** *12*, 281.

36. Benhamouche, S.; Decaens, T.; Godard, C.; Chambrey, R.; Rickman, D. S.; Moinard, C.; Vasseur-Cognet, M.; Kuo, C. J.; Kahn, A.; Perret, C.; Colnot, S., Apc tumor suppressor gene is the "zonation-keeper" of mouse liver. *Dev Cell* **2006,** *10* (6), 759-70.

37. Sekine, S.; Lan, B. Y.; Bedolli, M.; Feng, S.; Hebrok, M., Liver-specific loss of beta-catenin blocks glutamine synthesis pathway activity and cytochrome p450 expression in mice. *Hepatology* **2006,** *43* (4), 817-25.

38. Charawi, S.; Just, P. A.; Savall, M.; Abitbol, S.; Traore, M.; Metzger, N.; Ravinger, R.; Cavard, C.; Terris, B.; Perret, C., LKB1 signaling is activated in CTNNB1-mutated HCC and positively regulates beta-catenin-dependent CTNNB1-mutated HCC. *J Pathol* **2019,** *247* (4), 435-443.

39. Spicer, J.; Rayter, S.; Young, N.; Elliott, R.; Ashworth, A.; Smith, D., Regulation of the Wnt signalling component PAR1A by the Peutz-Jeghers syndrome kinase LKB1. *Oncogene* **2003,** *22* (30), 4752-4756.

40. Sun, T. Q.; Lu, B. W.; Feng, J. J.; Reinhard, C.; Jan, Y. N.; Fantl, W. J.; Williams, L. T., PAR-1 is a Dishevelled-associated kinase and a positive regulator of Wnt signalling. *Nature Cell Biology* **2001,** *3* (7), 628-636.

Figure 1 Acevedo-Acevedo and Stefkovich et al.

Figure 2 Acevedo-Acevedo and Stefkovich et al.

Figure 3 Acevedo-Acevedo and Stefkovich et al.

Figure 4 Acevedo-Acevedo and Stefkovich et al.

Figure S1 Acevedo-Acevedo and Stefkovich et al.

Figure S2 Acevedo-Acevedo and Stefkovich et al.