**MUCOSECRETORY LUNG DISEASE: DIFFERENT ASSEMBLIES OF JAG1 AND JAG2**

**DETERMINE TRACHEOBRONCHIAL CELL FATE**

Susan D. Reynolds[1,5]

Cynthia L. Hill[1]

Alfahdah Alsudayri[1]

Scott W. Lallier[1]

Saranga Wijeratne[2]

ZhengHong Tan[3]

Tendy Chiang[3]

Estelle Cormet-Boyaka[4]

1.  Center for Perinatal Research, Nationwide Children's Hospital, Columbus, OH, USA,

43215

2.  Institute for Genomic Medicine, Nationwide Children's Hospital, Columbus, OH, USA,

43215

3.  Center for Regenerative Medicine, Nationwide Children's Hospital, Columbus, OH, USA

43215.

4. School of Veterinary Medicine, The Ohio State University, Columbus, OH, USA, 43210

5. Contact Information: Susan D. Reynolds, PhD, Nationwide Children's Hospital, 700

Children's Drive, Columbus, OH 43205, Phone: (614) 355-6716, Fax: (614) 355-6675,

Email: susan.reynolds@nationwidechildrens.org

**Word count:** 6414

**Figures:** 7

**Tables:** 0

**References:** 67

**Supplemental Information:** yes

**Running Head:** JAG1 and JAG2 determine cell fate

**Key words:** airway epithelial stem cell, Notch signaling, Jagged, cell fate, goblet cell, ciliated

cell, biotinylation, ubiquitination, protein trafficking

**Author Contributions**

SDR: Conception and design, acquisition of laboratory data, interpretation of data, manuscript

preparation and review.

CLH:  Acquisition of laboratory data, manuscript review

AA: Acquisition of laboratory data, manuscript review

SWL: Acquisition of laboratory data, manuscript review

SW: RNAseq and scRNAseq analysis, manuscript review

ZHT: scRNAseq analysis, manuscript review

TC: Data review and interpretation, manuscript review

ECB: Data review and interpretation, manuscript review

**Funding:** Cystic Fibrosis Foundation Research Grant (REYNOL17XX0-20XX0*),* Cystic Fibrosis

Foundation Therapeutics, Inc. Research Development Program Grant (MCCOY19R0),

Nationwide Children's Hospital Cell Based Therapy Program (47306-0002-1217, SDR), and

Nationwide Children's Hospital Technology Development Award (Technology 2019-015, SDR).

**ABSTRACT (200 WORDS):**

Mucosecretory lung disease compromises airway epithelial function and is characterized by

goblet cell hyperplasia and ciliated cell hypoplasia. These cell types are derived from

tracheobronchial stem/progenitor cells via a Notch dependent mechanism. Although specific

arrays of Notch receptors regulate cell fate determination, the function of the ligands Jagged1

(JAG1) and Jagged2 (JAG2) is unclear. This study used primary human bronchial air-liquid-

interface cultures, gamma secretase inhibition, and neutralizing antibodies to show: 1) JAG1

and JAG2 were necessary for secretory progenitor cell fate determination; 2) JAG2 suppressed

squamous differentiation; and 3) pausing of the ciliated cell differentiation process after Notch

inhibition. Histological, cell fractionation, cell surface biotinylation, and ubiquitination analyses

demonstrated that all cells were JAG1 positive but that little JAG1 was present on the cell

surface. In contrast, JAG2 was expressed in a positive-negative pattern and was abundant on

the cell surface. Glycogen synthase kinase 3 (GSK3) and tankyrase inhibition studies showed

that GSK3 regulated JAG2 trafficking, and that this mechanism was WNT-independent.

Collectively, these data indicate that variation in JAG2 trafficking creates regions of high,

medium, and low ligand expression. Thus, distinct assemblies of JAG1 and JAG2 may regulate

Notch signal strength and determine the fate of tracheobronchial stem/progenitor cells.

**INTRODUCTION**

The conducting airway epithelium protects the lung from inspired environmental agents via

mucociliary clearance. Effective removal of pathogens, irritants, and toxins (including cigarette

smoke and air pollutants) requires the coordinated activity of two specialized airway epithelial

cell types, the multiciliated cell (termed a ciliated cell) and secretory cells (including goblet cells

and club cells). These cell types are descendants of the tracheobronchial epithelial tissue

specific stem cell (TSC) which regenerates the pseudostratified airway epithelium and resides in

the trachea and bronchi of the mouse and upper respiratory tract of humans. Lineage-tracing in

mice (1-4) and clonal analysis in human (5) indicated that the TSC was a basal cell subtype.

Previous studies reported that tracheobronchial ciliated and secretory cells were present in

approximately equal numbers and that their frequency was regulated by Notch signaling

(reviewed in (6)). Specifically, the Notch Lateral Induction Pathway (Notch pathway) regulates

basal and secretory cell differentiation (reviewed in (7)). Notch signaling occurs between

adjacent cells which express Notch ligands (Jagged (JAG) 1 or 2 or Delta-like (DLL) 1, 3, or 4)

or Notch receptors (NOTCH1, 2, 3, or 4) (6, 8). Following receptor ligation, ADAM10 or

ADAM17 cleaves NOTCH at an extracellular site and the $\gamma$-secretase complex cleaves NOTCH

at an intracellular site. This process releases the NOTCH intracellular domain which forms a

transcriptional complex with several proteins including RBPJK and activates expression of the

HES and HEY family of transcriptional repressors (9-11).

Chronic lung disease is frequently associated a disruption of mucociliary clearance. In asthma,

chronic obstructive pulmonary disease (COPD), chronic bronchitis (CB), and idiopathic

pulmonary fibrosis (IPF) goblet cell hyperplasia and excess mucus secretion drive the

mucosecretory phenotype. Single cell RNAseq (scRNAseq) analysis of asthma, IPF, and COPD

indicated that Notch signaling was aberrant (12-14) and functional studies indicated that

cigarette smoke, an agent which increases risk of developing COPD, CB, or IPF, activated

Notch signaling (15). Mucociliary clearance can also be disrupted by ozone and cigarette smoke

exposure or infection with SARS-CoV2 and/or respiratory syncytial virus. These agents damage

the ciliated cell, often resulting in cell death and ciliated cell hypoplasia. Notch pathway

components were identified as susceptibility genes for lung damage (16), viral entry (17), and

hyperinflammation (18). Consequently, treatments which specifically target Notch pathway

components have the potential to normalize ciliated and goblet cell frequency, restore

mucociliary clearance, and improve the health of people with chronic lung disease.

Several studies defined the Notch receptors responsible for generation of ciliated and secretory

cells in mouse and human (13, 19-23). However, the Notch ligand(s) that is responsible for

pathway activity is unclear. Studies in mice indicate that DLL ligands regulate generation of

neuroepithelial cells but are unlikely to be involved in ciliated and secretory cell differentiation

(24). In contrast, over-expression of *Jag2* mRNA relative to *Jag1* in basal cells (21), treatment-

dependent increases in *Jag1* and *Jag2* mRNA (25), and co-localization of the two mRNAs (24,

26) suggested overlapping roles for JAG1 and JAG2 in basal cell fate determination. Conditional

knockout of *Jag1* or *Jag2* early in mouse lung development supported this overlapping

mechanism but other studies identified JAG2 as the dominant ligand (24, 26). Finally, a specific

role for JAG1 in maintenance of secretory cell fate and suppression of ciliated cell fate was

indicated by experiments involving *Jag1* overexpression in immortalized human basal cells (27),

analysis of *Jag1* knockout in mice (28), and treatment of mice with a JAG1 neutralizing antibody

(29) or a *Jag1* antisense oligonucleotide (30).

A potential explanation for the disparate Jag results is the focus on the fully differentiated

ciliated or goblet cell. Ciliated cell subtypes have been identified by scRNAseq (31, 32), while

histological analysis of the mouse tracheobronchial epithelium (33) and the Xenopus epidermis

(reviewed in (34)) defined ciliated cell differentiation intermediates. These intermediates can be

detected by immunostaining for acetylated tubulin (ACT, (35)). Similarly, multiple secretory cell

subtypes have been identified, with the MUC5B-expressing goblet cell being the major subtype

(36). Goblet cell differentiation intermediates can be identified by immunostaining for MUC5B.

This study examined Notch signaling during differentiation of tracheobronchial TSC and

progenitor cells and focused on roles for JAG1 and JAG2. Cell fate determination was evaluated

using human bronchial air-liquid-interface cultures, a well-described and widely-utilized model of

the tracheobronchial epithelium (37). Treatment with γ-secretase inhibitors and analysis of

differentiation intermediates was used to examine Notch regulation of cell fate. Roles for JAG1

and JAG2 in differentiation of TSC and the secretory progenitor were evaluated using

histological and biochemical approaches. JAG1 and JAG2 necessity was examined using a

JAG1 peptide and anti-JAG1 and anti-JAG2 neutralizing antibodies.

**METHODS**

*Human TSC recovery, expansion, and ALI culture:* Human bronchial basal cells were recovered

as previously described (38, 39). The modified conditional reprogramming culture (mCRC)

method (39) was adapted from Suprynowicz et al (40). The major change from this protocol was

the use of irradiated NIH3T3 fibroblast feeder layers. Human basal cells were differentiated

using previously published methods (39). Initial studies used three culture media: Wu (41), Half

& Half (H&H, (35), or complete Pneumacult (Stem Cell Technology, Vancouver, BC, Canada).

Culture medium is defined in Results and in Figure Legends.

*ALI culture treatment protocol*: The specific treatment and analysis days and concentration of

agonists/antagonists are presented in Results and Figure Legends. Small molecule agonists

and antagonists were purchased from Tocris (Minneapolis, MN) and dissolved in DMSO at

1000x the treatment concentration. Vehicle controls were treated with an equal volume of

DMSO. Jag1 peptide was purchased from Anaspec (Freemont, CA) and dissolved in 1x

Phosphate Buffered Saline (PBS) at 500x the treatment concentration. Vehicle controls were

treated with an equal volume of PBS. Neutralizing antibodies specific for Jag1 (anti-JAG1.b70)

and Jag2 (anti-JAG2.b33) were provided by Genentech (29) and were dissolved in PBS at 100x

the treatment concentration. Vehicle controls were treated with an equivalent volume of PBS.

*Immunofluorescence analysis and quantification*: Differentiation was evaluated in ALI cultures.

Cultures were fix with 4% paraformaldehyde/1% sucrose/1x PBS. For analysis of intracellular

markers cultures were permeabilized and blocked with 0.1% NP40/5% bovine serum albumin/1x

PBS. For analysis of cell surface markers, cultures were fixed with 10% neutral buffered

formalin and blocked with 5% bovine serum albumin/1x PBS. Goblet cells and differentiation

intermediates were identified by MUC5B staining using rabbit-anti-MUC5B (Sigma, 1/500).

Ciliated cells and differentiation intermediates were identified by ACT staining using mouse-anti-

ACT (Sigma, 1/8000). JAG1 was evaluated using a mouse-anti-JAG1 C-terminal specific

antibody (BD Biosciences #612346, 1/50). JAG2 was evaluated using a rabbit-anti-JAG2 N-

terminal specific antibody (Cell Signaling C23D2, 1/50) and a rabbit-anti-JAG2 C-terminal

specific antibody (Cell Signaling C83A8, 1/50). HES1 was evaluated using a goat-anti-HES1

(Santa Cruz #sc-13842, 1/40). Rbpjk was evaluated using a rabbit-anti-RBPJK (Sigma

#AB5790, 1/100). Primary antibodies were detected with ALEXA-488 or ALEXA-594 labeled

secondary antibodies (Jackson Immunological, 1/500). Nuclei were detected with DAPI. Cell

type frequency was quantified as previously described (39).

*Cell fractionation:* Cells were separated into cytoplasmic and nuclear fractions using a NE-PER

kit (Thermo Scientific, P178833). 4x Laemmli's Buffer/10% β-mercaptoethanol (BME) was

diluted 1/4 into the protein fraction and the mixture was incubated at 96 ˚C for 10 minutes.

*Cell lysis:* RIPA buffer (42) was used to lyse cells for immunoprecipitation, poly-ubiquitination,

and western blot analysis. The apical and basal surfaces of ALI cultures were washed with

Hams/F12 at 4 ˚C for 10 min. Wash solution was aspirated and 270 µl RIPA buffer was added to

the apical surface. The cell layer was disrupted with a p200 pipette tip and then incubated for 20

min at 4 ˚C. The lysate was collected in a 1 ml microcentrifuge tube, sonicated 5 x 30 sec in a

Bioruptor (Diagenode) at 4 ˚C, and centrifuged at 11,176g for 10 min at 4 ˚C. The supernatant

was stored at -80 ˚C.

*Immunoprecipitation:* Immunoprecipitation was performed using a modified version of the Pierce

Protein Magnetic Bead protocol (Thermo 88848). For each sample, 125 µl of beads were pre-

washed three times with RIPA Buffer and resuspended in RIPA Buffer with proteinase (Roche;

11836153001) and phosphatase inhibitors (Roche; 04906837001). Washed beads (25 µl) were

added to 400-600 ug protein per sample and the lysate was cleared by rocking for 1 hr at 4 ˚C.

The cleared lysate was incubated with a titration-determined amount of antibody (usually 1:100).

This mixture was rocked for 1 hr at 4 ˚C. The remaining 100 µl of washed beads were added to

the sample and rocked for 2 hr at 4 ˚C. The beads were collected using a neodymium magnet,

washed three times with RIPA Buffer, resuspended in 1x Laemmli's Buffer plus 2.5% BME, and

incubated at 96 ˚C for 10 minutes. The supernatant was kept as Unbound Fraction.

*Cell surface biotinylation:* A modified version of the Pierce Cell Surface Protein Biotinylation and

Isolation Kit (Thermo A44390) protocol was used to isolate surface proteins. The recommended

biotinylation incubation time was increased to 45 minutes and incubations were done on ice.

*Poly-ubiquitination:* Total Poly-ubiquitinated proteins were isolated using a LifeSensors Tandem

Ubiquitin Binding Entities (TUBEs) Kit (LifeSensors, UM411M) and a modified protocol. The cell

lysate was incubated with 100 µl washed, unconjugated beads at 4 ˚C on a rotating platform for

30 minutes. A bead titration study demonstrated an optimal ratio of 100 µl TUBE-conjugated

beads per 1-2 mg protein in a total of 1 ml lysis buffer. The cleared lysate and washed TUBE

beads were incubated at 4 ˚C on a rotating platform for 30 minutes. The TUBE beads were

collected using a magnet, washed three times with PBST, and divided into two aliquots. One

aliquot was treated with buffer and the second aliquot was treated with 2 µl of 10 µM broad-

spectrum deubiquitinase for 2hr at 30˚C with rocking. The beads were resuspended in 1x

Laemmli's Buffer/2.5% BME and incubated for 10 minutes.

*Western blot analysis and quantification*: Protein concentration, gel electrophoresis and transfer,

and blotting were done as previously reported (42). In some instances, blots were stripped by

incubation with 1X Western Reprobe (G Biosciences) for 45 minutes at 55 ˚C followed by three

washes with Tris buffered saline with TWEEN-20 (TBS-T, Cell Signaling Technology) at room

temperature. Antibodies used for western blot studies: JAG2-N-terminal (Cell Signaling C23D2,

1:1000); JAG2-C-terminal (Cell Signaling C83A8, 1:1000); GSK3 (Millipore 05-412, 1:2000);

pGSK3 (Millipore 0-413, 1:1000); 14-3-3 (Millipore AB9748-I, 1:2000); TOPO II (Abcam 109524,

1:10000); Sodium Potassium ATPase (NAK, Cell Signaling 3010S, 1:2500); JAG1-C-terminal

(BD Biosciences #612346, 1:1000); α-Tubulin (Sigma T6793, 1:2000); N-terminal β-catenin Cell

Signaling #9581X, 1:1000); C-terminal β-catenin (BD Transduction #610154, 1:2000); and

Keratin 5 (BioLegend #905501, 1:2000).

*RNA preparation and RNAseq gene expression analysis:* RNA was purified using a RNeasy Kit

(Qiagen,74106) according to the manufacturer's directions. RNA-sequencing (RNAseq) was

conducted by the Institute for Genomic Medicine at Nationwide Children's Hospital. On average,

32 million paired-end 151 bp RNA-Seq reads were generated for each sample (the range was

25 to 37 million). Each sample was aligned to the GRCh38.p9 assembly of the Human

reference from NCBI (http://www.ncbi.nlm.nih.gov/assembly/GCF_000001405.35/) using

version 2.6.0c of the RNA-Seq aligner STAR

(http://bioinformatics.oxfordjournals.org/content/29/1/15). Transcript features were identified

from the GFF file provided with the assembly from Gencode (v28) and raw coverage counts

were calculated using featureCounts (http://bioinf.wehi.edu.au/featureCounts/). The raw RNA-

Seq gene expression data was normalized and post-alignment statistical analyses were

performed using DESeq2 (http://genomebiology.com/2014/15/12/550) and custom analysis

scripts written in R. Comparisons of gene expression and associated statistical analysis were

made between different conditions of interest using the normalized read counts. All fold change

values are expressed as test condition / control condition, where values less than one are

denoted as the negative of its inverse (note that there will be no fold change values between –1

and 1, and that the fold changes of "1" and "-1" represent the same value). Transcripts were

considered significantly differentially expressed using a 10% false discovery rate (DESeq2

adjusted p value <= 0.1). GSEA was used to identify treatment dependent changes in signaling.

*scRNAseq data analysis*: Single cell RNAseq methods and analytical techniques are presented

in detail in (13) and (43). The raw reads were processed with Cell Ranger software 3.0.2 (10X

Genomics) using hg38 transcriptome reference from GENCODE 25 annotation, and a digital

expression matrix was generated with cell barcodes as rows and gene identities as columns.

Each dataset was run with SoupX analysis package to remove contaminant 'ambient' RNA

derived from lysed cells during isolation and capture (44). To minimize doublet contamination,

for each dataset the Scrublet package was used to detect neotypic doublets (45) and a quantile

thresholding was performed to identify high UMI using a fit model generated using multiplet's

rate to recovered cells proportion, as previously described (43). For all data, quality control and

filtering were performed to remove cells with low number of expressed genes (threshold

n>=200) and elevated expression of apoptotic transcripts (threshold mitochondrial genes <

15%). Only genes detected in at least 3 cells were included. Data were normalized using the

Scran package (46), using a low resolution clustering to use as input for the size factor

estimation and then log transformed. Data were processed with Principal Component Analysis

(PCA) using the 5000 most variable genes as input, and clustering was performed with the

Leiden algorithm. Batch correction for the two samples was performed with the package BBKNN

(Polanski et al., 2019). Partition-based graph abstraction (PAGA) (47) was added to Uniform

Manifold Approximation and Projection (UMAP) with the option init_pos='paga' (48) to obtain a

two-dimensional visualization. To identify differentially expressed genes between clusters and

representative of both datasets, Model-based Analysis of Single-cell Transcriptomics (MAST)

(49) was used. To characterize cell subsets in our integrated dataset, previously published lung

cell subtype specific gene lists (43) were used to create cell type-specific gene signatures using

a strategy previously described (50). The steps described above were generated using Seurat

3.0 in R 4.0 or Scanpy 1.7.0 in Python 3.8.5. Plots were generated within Seurat and Scanpy or

using the ggplot2 package.

*Statistics*: All statistical analyses were performed using Graph Pad Prism. Data normality was

evaluated by the Shapiro-Wilk test, the Anderson-Darling test, and/or the D'Agostino & Pearson

test. For normally distributed data sets, differences were evaluated using Student's t-test. For

non-normally distributed data sets, differences were evaluated by the Mann-Whitney test and

data are presented as the median and the interquartile range. Trends were analyzed by

regression analysis. The definition of center and dispersion precision measures are indicated in

the Figure Legends. Data sets containing multiple variables were analyzed by analysis of

variance (ANOVA) and a post hoc Tukey test (normally distributed data sets) or Kruskal Willis

test (non-normally distributed data sets). Sample size is indicated in the Figure Legends. P-

values $< 0.05$ were considered significant.

*Study approval:* The Institutional Review Board at Nationwide Children's Hospital approved the

human studies. Written informed consent and assent was obtained from every participant.

**RESULTS**

*Ciliated and goblet cell differentiation*: Since kinetics and cell type frequency vary across model

systems, differentiation was evaluated in three differentiation media. Quantification

demonstrated that cell density did not vary as a function of medium or time (SFig 1A) and that

the frequency of ciliated cells increased significantly between differentiation days 7 and 14 in all

three media (SFig 1B). The frequency of mature goblet cells did not vary between differentiation

days 7 and 14 in Wu or complete Pneumacult medium but increased significantly in Half&Half

(H&H) medium (SFig 1C). Based on the finding that H&H medium supported an increase in

ciliated and goblet cell frequency over days 7-14 and generated similar numbers of goblet and

ciliated cells on day 14, this medium was used to evaluate the signaling mechanisms that

regulate production of ciliated and goblet cells.

*Intermediate ciliated and goblet cell phenotypes*: Previous work demonstrated that ciliated cell

differentiation involved a set of intermediate phenotypes (reviewed in (34)). In Stage I, cells exit

the cell cycle and express a primary cilium (SFig 2A-B). This phenotype was detected at early

time points and the frequency of cells with a primary cilium peaked on differentiation day 6 (Fig

1A). During Stages II and III, new centrioles are generated and dock at the apical cell surface.

Cells that contained many basal bodies and occasional short cilia (bristle cells, SFig 2C-D) were

identified on days 8-12 (Fig 1B). Finally, during Stage IV, each docked centriole (now termed a

basal body) nucleates a motile 9+2 ciliary axoneme and the ciliary appendage is formed. Cells

defined many long motile cilia (ciliated cells, SFig 2E-F) were detected on days 8-12 (Fig 1C).

Goblet cells also exhibited three morphological phenotypes that were defined by low, medium,

and high expression of MUC5B (SFig 2G-H). All three intermediates were detected at all time

points (Fig 1D-F). The frequency of MUC5B-low and MUC5B-medium cells peaked on day 8

while the frequency of MUC5B-high cells increased through day 12 (Fig 2F). These data

indicated that the first wave of ciliated and goblet cell differentiation was completed the process

by day 12.

*Notch regulation of ciliated and goblet cell differentiation*: Previous studies demonstrated that

treatment with DAPT, a γ-secretase inhibitor (GSI), inhibited Notch signaling, increased the

frequency of ciliated cells, and decreased the frequency of mature goblet cells (35). To

determine if this paradigm applied to the first wave of differentiation, H&H cultures were treated

with DAPT on days 0-4, 2-6, 4-8, or 8-12. DAPT treatment did not alter cell density at any time

point (SFig 3).

DAPT treatment significantly increased the frequency of cells with a primary cilium on day 4 (Fig

1A), significantly increased the frequency of bristle cells on days 4, 6, and 8 (Fig 1B), and

significantly increased the frequency of ciliated cells on days 8 and 12 (Fig 1C). DAPT treatment

did not alter the frequency of MUC5B-low or MUC5B-medium cells at any time point (Fig 1D-E).

However, DAPT treatment significantly decreased the frequency of MUC5B-high cells on days 8

and 12 (Fig 1F). Similar results were gathered in experiments using a second GSI, LY411575

(LY, data not shown). Collectively, these data indicated that Notch signaling was active during

the first wave of differentiation.

*Pausing of the Notch signal:* Accumulation of ciliated cell differentiation intermediates suggested

that GSI treatment paused the ciliation process. To investigate this mechanism, cultures were

treated with LY for 4 days and recovered (recovery) or treated with LY for 8 days (continuous,

Fig 2A). On differentiation day 12, ciliated and goblet cell intermediates were quantified. Cells

with a primary cilium were not detected in these cultures (Fig 2B). Both bristle and ciliated cells

were detected, and their frequency did not vary between the recovery and continuous treatment

protocols (Fig 2B). Goblet cells defined by low, medium, and high expression of MUC5B were

identified but their frequency did not vary with treatment. Since continuous Ly treatment did not

cause an increase in bristle or ciliated cell frequency, these data indicated that GSI treatment

paused ciliated cell differentiation.

Since Notch signaling in the tracheobronchial epithelium has been attributed to JAG1 and

JAG2, the impact of GSI treatment on JAG1 or JAG2 protein abundance was determined.

Cultures were treated with LY or DAPT on days 4 and 6 and protein expression was analyzed

by western blot on day 8. GSI treatment did not alter JAG1 abundance (Fig 2C). In contrast, GSI

treatment significantly increased the abundance of full-length JAG2 (Fig 2D). These data

suggested that GSI treatment paused the differentiation process by increasing JAG2.

*Cellular localization of JAG1 and JAG2*: A JAG2-dependent pausing mechanism was

unanticipated and supported an in-depth analysis of JAG1 and JAG2 distribution on

differentiation day 7. Immunofluorescence analysis detected JAG1 in cultures that were

differentiating in each test medium (SFig 4A-C, 4J-K) and showed that JAG1 was highly

localized to the perinuclear/nuclear compartments. All cells were positive for JAG1. JAG2 was

also detected in all three culture conditions but was most abundant in cultures that were

differentiating in H&H medium (SFig 4D-F, 4J-K). In contrast with JAG1, JAG2 was highly

enriched in the cortical domain. All cells were positive for JAG2; however, expression varied

from low to high.

*JAG1 subcellular location and molecular weight*: The subcellular location of JAG1 on

differentiation day 7 was unexpected and led to analysis at additional time points. On

proliferation day 3 (Fig 3A, D, E) and differentiation day 2 (Fig 3B), JAG1 was expressed in a

subset of cells and was in the perinuclear/nuclear domain. JAG1 did not colocalize with β-

catenin, a component of the adherens junction (Fig 3F-G). On differentiation day 4 (Fig 3C),

JAG1 was detected in most cells and some variation in protein abundance was noted.

Previous studies reported that Notch ligands can be cleaved by γ-secretase resulting in creation

of a transcriptionally active 25 kDa C-terminal peptide (51-53). Consequently, JAG1 molecular

weight was determined using a JAG1 antibody which detected the C-terminal domain. On days

4 and 8, all samples contained the full-length (~130 KDa) JAG1 protein (Fig 3H). To further

examine JAG1 subcellular localization, cytoplasmic and nuclear fractions were isolated on day

8. Detection of Topoisomerase II in the nuclear fraction and alpha tubulin in the cytoplasmic

fraction (Fig 3I) demonstrated successful separation. JAG1 was limited to the cytoplasmic

fraction and indicated that the protein was perinuclear rather than nuclear.

Since the previous studies indicated that most if not all JAG1 was in a perinuclear compartment

and that JAG1 was not acting as transcription factor, the relationship between transcription and

translation was evaluated. *Jag1* mRNA was quantified by RNAseq and JAG1 protein levels

were quantified by Western blot. *Jag1* abundance decreased as a function of time (Fig 3J). In

contrast, JAG1 abundance was constant over time (Fig 3K). These data suggested that a post-

translational mechanism regulated JAG1 abundance.

*JAG2 subcellular location and molecular weight*: JAG2 localization was further evaluated on

proliferation day 5 and differentiation days 4 and 8 (Fig 4A-C). The cortical pattern was

observed on proliferation day 5 and JAG2-high cells were apparent on differentiation day 4. By

differentiation day 8, JAG2-positive and -negative cells were noted. At each time point, JAG2

colocalized with β-catenin. On differentiation day 8, groups of cells expressing nuclear HES1

and RBPJK were detected (Fig 4D) and indicated focal Notch signaling.

The molecular weight of JAG2 was determined using two antibodies which were specific to the

N-terminal or C-terminal domains. This study detected full-length (~145 KDa) JAG2 on days 4

and 8 (Fig 4E). To further examine JAG2 subcellular localization, co-immunoprecipitation was

used to determine if JAG2 was associated with the adherens junction. An N-terminal specific β-

catenin antibody was used for immunoprecipitation and the precipitate was analyzed with N-

terminal and C-terminal specific β-catenin antibodies. β-catenin was successfully isolated (Fig

4F) and JAG2 was identified in the β-catenin complex.

Since the previous studies suggested that much of JAG2 was trafficking to or from the plasma

membrane, the relationship between transcription and translation was evaluated using RNAseq

and western blots. *Jag2* abundance decreased as a function of time (Fig 4G). In contrast, JAG2

abundance was constant over time (Fig 4H). Like JAG1, these data suggested a post-

translational mechanism regulated JAG2 abundance.

*Regulation of JAG2 abundance*: Identification of JAG2-positive and JAG2-negative cells on

differentiation day 8 suggested that JAG2 was degraded in some cells. A candidate regulator is

the WNT/β-catenin pathway (35). To examine this mechanism, ALI cultures were treated with

various concentrations of CHIR99021 (CHIR, a WNT/β-catenin agonist, (54)) or XAV939 (XAV,

a WNT/β-catenin antagonist, (55)) on differentiation days 4 and 6 and cell lysates were collected

on day 8. Western blot analysis demonstrated that neither CHIR nor XAV treatment altered the

abundance of JAG1 (Fig 5A). In contrast, both treatments caused a significant decrease in

JAG2 abundance (Fig 5B). Interestingly, the CHIR effect was limited to lower doses (5 and 10

μM); whereas the XAV effect was observed at all three doses.

*Impact of CHIR and XAV on the transcriptome:* CHIR and XAV are typically thought to regulate

WNT/β-catenin pathway by altering gene expression. To evaluate this mechanism, ALI cultures

were treated with vehicle, 10 µM CHIR or 10 µM XAV on differentiation days 4 and 6 and RNA

was purified on day 8. RNA-sequencing and Gene Set Expression Analysis (GSEA)

demonstrated that CHIR upregulated expression of WNT/β-catenin target genes including

*Axin2*, *Lef1*, *Tcf7* (SFig 5A-B, SFig 6). These genes contributed to the GSEA Leading-Edge and

had a high impact on the GSEA Enrichment Score (an indicator of changes in pathway activity).

Other GSEA Leading-Edge genes were *Wnt5B* and known pathway antagonists (*Axin1, Dkk1*,

*Dkk4,* and *Nkd1*). Since these genes and *Axin2* inhibit the WNT/β-catenin pathway and *Lef1*

and *Tcf7* repress gene expression, these data suggested that CHIR antagonized the WNT/β-

catenin pathway on day 8.

In keeping with its perceived function as a WNT/β-catenin pathway antagonist, XAV treatment

downregulated expression of *Axin2*, *Lef1*, and *Tcf7*, and *Wnt5B* (SFig 5C-D, SFig 6). However,

most GSEA Leading-Edge WNT/β-catenin pathway genes were typical of the Notch pathway

(*Dll1*, *Hey1*, *Jag1*, *Jag2*, *Notch1*, and *Notch4*). Analysis of the Notch gene set indicated that

XAV treatment caused a significant downregulation of Notch signaling (SFig 5G-H). A similar

analysis of the Notch gene set in CHIR treated cultures also indicated downregulation (SFig 5E-

F). Although the high FDR q-value suggested cautious interpretation of the CHIR-Notch

pathway interaction, many of the GSEA Leading-Edge genes were shared between the CHIR

and XAV groups (*Dll1*, *Jag1*, *Notch1*, *Lfng*, *Prkca*, *Psenen*, *Sap30,* and St3gal6, SFig 5E-H,

SFig 6).

Although *Jag1* and *Jag2* contributed to the Enrichment Scores, neither gene product was

consistently included in the GSEA Leading-Edge gene sets. Consequently, the impact of CHIR

and XAV treatment on gene expression was evaluated. *Jag1* and *Jag2* abundance did not vary

by treatment (SFig 5I-J). Collectively, the gene expression data indicated that the main effect of

CHIR or XAV treatment was to antagonize WNT/β-catenin and Notch signaling and that

decreased signaling could not be attributed to a change in *Jag1* or *Jag2* gene expression.

*WNT independent regulation of JAG2 abundance*: To examine the impact of WNT antagonism

on JAG1 and JAG2 abundance, ALI cultures were treated with vehicle, LGK974 (LGK) a

porcupine inhibitor which prevents WNT secretion, or DKK1 which inhibits WNT signaling by

binding the LRP5/6 coreceptor. Neither LGK treatment (Fig 5A-B) nor DKK1 treatment (Fig 5C-

D) altered JAG1 or JAG2 abundance. These data suggested that a WNT-independent

mechanism regulated the abundance of JAG2 in CHIR and XAV treated cultures.

*CHIR regulation of cell fate*: While the decrease in JAG2 abundance in response to CHIR or

XAV treatment was unexpected, a previous study reported that both drugs altered secretory and

ciliated cell differentiation (35). To further examine changes in cell fate, ALI cultures were

treated with CHIR on days 4 and 6 and cells were isolated for single cell RNAseq in day 8.

Fourteen clusters (0-13) were identified (Fig 5E, F) and 10 clusters differed significantly

between vehicle and CHIR treated cultures (Fig 5G). In vehicle treated controls, clusters 6, 8,

and 10 exhibited a mitotic signature (STable 1). The frequency of cells in these clusters

increased in CHIR treated cultures. Cluster 1 was prominent in vehicle treated cultures and

contained basal cells which exhibited a matrix production signature. This cluster over-expressed

*Jag2* and *Wnt4* and was nearly absent in CHIR treated cultures. Clusters 4, 0, 3 and 12

contained cells within the secretory lineage and clusters 4, 0 and 3 expressed Hes4, Hes1,

and/or Hes2. Cluster 3 contained secretory primed basal cells that were defined by expression

of *Ceacam6*, *Scgb1A1*, and *Scgb3A2*. Cells within cluster 3 were less mature than those in

cluster 12. Clusters 4, 0 and 3 were significantly decreased in CHIR treated cultures. Clusters

13 and 5 contained ciliated cell lineage cells. Cluster 13 was enriched in early-stage ciliated

cells defined by expression of *Mcidas* and cluster 5 contained more mature ciliated cells. Cluster

2 was increased in CHIR treated cultures and clusters 7 and 9 were unique to CHIR treated

cultures. Cluster 7 was distinguished by down-regulation of WNT pathway antagonists (*Axin2*,

*Dkk2*, *Dkk4*, *Nkd1*), and the presence of cell motility and keratinization signatures. The

keratinization signature was upregulated in clusters 2 and 9. Since keratinization was previously

associated with squamous differentiation in the tracheobronchial epithelium (56), these data

indicated that CHIR treatment promoted this fate and suggested that JAG2 suppressed

squamous differentiation.

*JAG1 and JAG2 trafficking to the cell surface*: To further investigate roles for JAG1 and JAG2 in

Notch signaling, cell surface protein was evaluated by immunofluorescence analysis of non-

permeabilized cultures. JAG1 was not detected on the cell surface on differentiation day 8. In

contrast, JAG2 was detected on the surface of many cells (Fig 6A-B). Subsequent

permeabilization and staining detected both JAG1 and JAG2 and illustrated their typical

intracellular distributions (Fig 6C-F).

Cell surface biotinylation was used to evaluate JAG1 and JAG2 trafficking to the plasma

membrane. These studies detected sodium/potassium ATPase (NKA) a known cell surface

protein as well as JAG1 and JAG2 (Fig 6G). However, cell surface JAG2 was 5-10 times more

abundant than JAG1. Since interaction with a Notch receptor results in ligand ubiquitination and

internalization, polyubiquitination of JAG1 and JAG2 was analyzed. Little or no JAG1 was

detected in the bound (ubiquitin-positive) fraction (Fig 6H, upper panel). In contrast, JAG2 was

detected in the bound fraction and the molecular weight of the captured JAG2 decreased after

treatment with a deubiquitinase (Fig 6H). These biochemical studies indicated that JAG2

participated in Notch-signaling and raised the possibility that JAG1 was not involved.

*Phosphorylation of JAG2*: Immunofluorescence analysis demonstrated that CHIR treatment

caused a striking redistribution of JAG2 to a perinuclear location (Fig 6I-J) and was suggestive

of JAG2 degradation. JAG2 was not detected by immunofluorescence staining in XAV treated

cells (data not shown). CHIR is a potent and highly selective inhibitor of GSK3 (54). In contrast,

XAV inhibits tankyrase and regulates GSK3 by altering the kinase's localization within the cell

(55). Since both drugs decrease GSK3 activity and decrease JAG2 abundance, it was possible

that the shared mechanism was a GSK3-dependent decrease in JAG2 phosphorylation. To

evaluate this mechanism, differentiation day 8 cell lysates were immunoprecipitated with an

anti-pSerine/pThreonine (pS/pT) antibody and western blots were probed for JAG2 and positive

controls (KRT5 and YWHA). This study detected JAG2, KRT5 and YWHA in the bound fraction

(Fig 6K) and indicated that JAG2 was phosphorylated on Serine and/or Threonine. Collectively,

these data suggested that normal JAG2 trafficking was dependent on phosphorylation by GSK3.

*Roles for JAG1 and JAG2 in Notch Signaling*: To determine if JAG1 was involved in Notch

signaling, cultures were treated with a JAG1 DSL-domain peptide on differentiation days 8 and

10 and cultures were fixed on differentiation day 12. Immunofluorescence analysis of ACT and

MUC5B was used to detect differentiation intermediates. Cells expressing a primary cilium were

not detected (Fig 7A). However, vehicle treated cultures contained a small number of bristle

cells and many ciliated cells. Treatment with the JAG1 peptide caused a significant increase in

the frequency of bristle cells but did not alter the frequency of ciliated cells (Fig 7A). The same

cultures did not contain MUC5B-low cells (Fig 7B). However, MUC5B-medium, and MUC5B-

high cells were detected. Treatment with JAG1 peptide did not alter the frequency of MUC5B-

medium cells but significantly decreased the frequency of MUC5B-high cells. These data

suggested two JAG1-dependent processes. First, that the JAG1 peptide inhibited Notch

signaling resulting in increased basal-to-bristle cell differentiation. Second, the decreased

maturation of MUC5B-medium cells combined with the increase in bristle cell frequency raised

the possibility that JAG1 peptide treatment promoted MUC5B-medium to ciliated cell

differentiation.

Since the JAG1 peptide is 71% identical to the JAG2-DSL domain, it was unclear if the JAG1

peptide results could be attributed to JAG1. Further, the JAG1 peptide results were consistent

with reports that used JAG1 peptide to antagonize Notch signaling (57) but disagreed with those

reporting that this reagent agonized Notch signaling (58, 59). Consequently, additional studies

used neutralizing antibodies to inhibit JAG1 or JAG2 activation of Notch receptors (29).

To address the limitations of the JAG1 peptide studies, ALI cultures were treated with

neutralizing antibodies on days 8 and 10 and ciliated and secretory cell differentiation was

assayed on day 12. Treatment with anti-JAG1 or anti-JAG2 significantly increased bristle cell

frequency and decreased ciliated cell frequency (Fig 7C-D). Neutralizing antibody treatment did

not alter the frequency of MUC5B-medium cells but significantly decreased the frequency

MUC5B-high cells (Fig 7E-F). These data suggested that loss of either JAG1 or JAG2 caused

MUC5B-medium cells to generate bristle cells. To address the finding that anti-JAG1 or anti-

JAG2 treatment decreased ciliated cell frequency, ALI cultures were treated with a mixture of

anti-JAG1 and anti-JAG2. This treatment significantly decreased bristle cell frequency but did

not alter ciliated cell frequency (Fig 7G-H). The combined neutralizing antibody treatment did

not alter the frequency of MUC5B medium cells and significantly decreased the frequency of

MUC5B-high cells (Fig 7I-J). These data in combination with the single antibody study indicated

that either JAG1 or JAG2 inhibited bristle cell to ciliated cell differentiation.

**DISCUSSION**

This study demonstrated that JAG1 and JAG2 cooperate to determine the fate of the MUC5B-

medium (secretory progenitor) cell (SFig 7A). The neutralizing antibody studies found that both

JAG1 and JAG2 were required to generate a mature MUC5B-high goblet cell. In contrast,

neutralization of either JAG1 or JAG2 allowed the MUC5B-medium cell to generate a bristle cell.

Finally, the CHIR and XAV studies indicated that degradation of JAG2 resulted in a squamous

fate (SFig 7B). While these data support the paradigm that high Notch signal strength selected

the goblet cell fate, the possibility of moderate and low Notch intensity within the secretory

lineage was novel.

Previous studies indicated that cell fate was determined by different assemblies of Notch

receptors (13, 20). The present study suggests that different assemblies of JAG ligands

determine Notch signal intensity. First, JAG1 was detected in all differentiating cells. Second, a

striking change in cell morphology coincided with the emergence of JAG2 positive and negative

cells and raised the possibility that increased polarization concentrated JAG2 at cell-cell

junctions (60). When combined with the histological and biochemical data, these studies

indicated that JAG1 might establish a minimum ligand concentration while changes JAG2

trafficking result in cells that express high or low levels of cell surface ligand. Thus, it is possible

that the MUC5B-medium cell fate is determined by its location in a ligand-low (JAG1 only),

ligand-medium (mixture of JAG1+/JAG2+ and JAG1+/JAG2- cells), or a ligand-high

(JAG1+/JAG2+) environment (Graphical Abstract).

This study also indicated that JAG2 paused the ciliated cell differentiation process (SFig 7C).

Initial support for pausing came from the finding that GSI treatment increased the frequency of

ciliated cell differentiation intermediates. These data suggested that Notch signaling fluctuated

from "off" to "on" as the cell progressed from one morphological intermediate to the next. These

data challenged the idea that ciliated differentiation occurred in the absence of a Notch signal

but also raised the possibility that the GSI-sensitive target was the ligand rather than the

receptor. Support for the latter idea comes from previous demonstration that JAG1 and JAG2

were cleaved by γ-secretase resulting in production of an intracellular domain which functions

as a transcription factor (51-53). Failure to detect the ~25 kDa intracellular domain may reflect

low sensitivity in the ALI model relative to ligand over-expression systems.

Evidence of pausing also challenged the notion that ciliated cell differentiation was continuous

and suggested that differentiation occurred in incremental steps. Demonstration that expression

of a ciliated transcription network gene product, *cMyb,* fluctuated over time and regulated

progression through the ciliated cell differentiation program (61) supported this interpretation.

Although the neutralizing antibody studies indicated that bristle-to-ciliated cell differentiation

occurred only when JAG1 and JAG2 were blocked, western blot analysis correlated the GSI-

dependent signal with an increase in full-length JAG2 abundance. Since gene expression

studies indicated that GSI treatment did not alter *Jag2* (or *Jag1*) mRNA abundance, it is

possible that pausing involved a change in JAG2 protein trafficking.

Previous studies showed that transmembrane proteins, including JAG1 and JAG2, transit the

endocytic recycling compartment as they move to the plasma membrane. This compartment is

positioned under the nucleus in adherent cells, and sorts endocytosed proteins that are recycled

to the plasma membrane, enter the lysosome for degradation, or are sequestered in a

perinuclear site termed the pericentrion (62). A protein's trafficking pattern is determined in part

by phosphorylation and ubiquitination and changes in the rate of movement through the various

endosomal compartments regulate Notch signaling (63).

This study identified a striking difference in JAG1 and JAG2 localization within the cytoplasm of

differentiating cells. Cell fractionation studies indicated that JAG1 was highly enriched in the

perinuclear domain. Demonstration that JAG1 was not poly-ubiquitinated indicated that most

JAG1 was in the endocytic recycling compartment and not sequestered in the pericentrion. In

contrast, the subcellular distribution and polyubiquitination of JAG2 were consistent with

endocytosis, recycling through the endocytic pathway, and post-signaling degradation via the

lysosome. Increased JAG2 protein levels in GSI-treated cultures and decreased JAG2 levels in

CHIR and XAV treated cultures suggested that JAG2 routing could be altered and that JAG2

and GSK3 might be co-localized in multivesicular bodies (64, 65).

A previous study reported that NOTCH1 trafficking to the cell surface was inhibited by GSK3-

mediated phosphorylation (66) while the present study showed that GSK3 inhibition resulted in

JAG2 degradation. Consequently, GSK3 activity might determine if a cell traffics JAG2 or

NOTCH1 to the cell surface. Although the WNT/β-catenin pathway was a likely regulator of

GSK3 activity, our previous report found that CHIR and XAV (35) inhibited ciliated cell and

goblet cell commitment and suggested a complex mechanism. The present study indicates that

both CHIR and XAV antagonized WNT/β-catenin signaling and that the main effect of these

drugs was downregulation of Notch signaling. Although these data suggest that Notch signaling

is dependent on β-catenin, a role for WNT signaling was discounted by studies which inhibited

WNT secretion or WNT interaction with LRP6/5. A similar WNT ligand-independent role for

GSK3 in cell fate specification was previously reported for iPSC that were undergoing directed

differentiation (67) and suggested that establishment of tracheobronchial ligand- and receptor-

expressing cells was regulated by a GSK3-dependent and WNT-independent mechanism.

**CONCLUSION**

This study demonstrates that JAG1 and JAG2 are necessary for cell fate specification within the

tracheobronchial epithelium. Differences in JAG1 and JAG2 trafficking raise the possibility that

distinct assemblies of these ligands generate high, medium, and low Notch signaling

environments within the differentiating epithelium (Graphical Abstract). Further analysis of JAG

trafficking has the potential to identify treatments which could normalize goblet and ciliated cell

frequency in people who have chronic mucosecretory lung diseases and those whose ciliated

cell population is depleted by exposure to environmental agents.

**ACKNOWLEDGEMENTS**

This project was funded by a Research Grant from Cystic Fibrosis Foundation (REYNOL17XX0-

20XX0, SDR), the Nationwide Children's Hospital Cell Based Therapy Program (47306-0002-

1217, SDR), and a Nationwide Children's Hospital Technology Development Award

(Technology 2019-015, SDR). We acknowledge the Abigail Wexner Research Institute Center

for Genomics Research which conducted the RNA sequencing and analysis, and the C3 RDP

which provided human bronchial cells (MCCOY19R0). JAG1 and JAG2 neutralizing antibodies

were provided by Genentech, Inc. under a Materials Transfer Agreement.

**FIGURE LEGENDS**

*Graphical Abstract:* Different assemblies of JAG1 and JAG2 may determine Notch signal

strength and cell fate within the tracheobronchial epithelium. A cell which interacts with JAG1+

cells (blue squares) receives a low Notch signal (light yellow square). A cell which interacts with

a mixture of JAG1+ and JAG1+/JAG2+ cells (purple squares) receives a medium (med) Notch

signal (medium yellow square). A cell which interacts with JAG1+/JAG2+ cells receives a high

Notch signal (bright yellow square).

*Figure 1: Notch regulation of ciliated and goblet cell differentiation.* Human bronchial basal cells

were differentiated in air-liquid-interface cultures using H&H medium. Cells were treated with

vehicle (DMSO) or 25 µM DAPT as follows: treatment on differentiation days 0 and 2 and fix on

day 4, treatment on differentiation days 2 and 4 and fix on day 6, treatment on differentiation

days 4 and 6 and fix on day 8, or treatment on differentiation days 8 and 10 and fix on day 12.

Dual immunofluorescence was used to detect acetylated tubulin and MUC5B. Nuclei were

stained with DAPI. **A.** Frequency of cells with a primary cilium. **B.** Frequency of bristle cells. **C.**

Frequency of ciliated cells. **D.** Frequency of MUC5B-low cells. **E.** Frequency of MUC5B-medium

cells. **F.** Frequency of MUC5B-high cells. Mean ± SD, N=6.

*Figure 2: Impact of intermittent or continuous Notch inhibition of ciliated and goblet cell*

*differentiation.* **A-C**. Human bronchial basal cells were differentiated in air-liquid-interface

cultures using H&H medium. Dual immunofluorescence was used to detect acetylated tubulin

and MUC5B. Nuclei were stained with DAPI. (**A**) Experimental design. Recovery study: Cells

were treated with 10 µM Ly on differentiation days 4 and 6 and with vehicle (DMSO) on

differentiation days 8 and 10. Cultures were fixed on day 12. Continuous study: Cells were

treated with 10 µM Ly on differentiation days 4, 6, 8, and 10. Cultures were fixed on day 12. (**B**)

Frequency of ciliated cell differentiation intermediates. P-cells with a primary cilium, B-bristle

cells, M-ciliated cells. (**C**) Frequency of goblet cell differentiation intermediates. L-MUC5B-low

cells, M-MUC5B-medium cells, H-MUC5B-high cells. Mean ± SD, N=3. **D-E**. Western blot

analysis of JAG1 (**D**) and JAG2 (**E**) abundance in cultures that were treated with DMSO, 10 µM

LY, or 25 µM DAPT on differentiation days 8 and 10 and harvested on differentiation day 12.

Mean ± SD, N=3.

*Figure 3: JAG1 subcellular location and length.* Human bronchial basal cells were differentiated

in air-liquid-interface cultures using H&H medium. **A-C.** Immunofluorescence was used to detect

JAG1 (red) on proliferation day 3 (**A**), differentiation day 2 (**B**), or differentiation day 4 (**C**).

Nuclei were detected with DAPI (blue). **D-E**. High magnification of the region indicated in panel

A. Arrows indicate JAG1 positive cells. **F-G.** Dual immunofluorescence analysis of JAG1 (green)
and β-catenin (β-Cat, red) on differentiation day 4. Nuclei were detected with DAPI (blue).
Three-color (**F**) and two color (**G**) images of the same region. **H.** Western blot analysis of JAG1
on differentiation days 4 and 8. The full-length band is labeled JAG1. Three samples per time
point. **I.** Cells were cultured to differentiation day 8, separated into cytoplasmic (C), nuclear (N),
insoluble (I) fractions, or were unfractionated (U), and proteins were analyzed by western blot.
Top panel: TOPOII, a nuclear protein; middle panel: alpha tubulin (αTUB), a cytoplasmic
protein; Bottom panel: JAG1. Two different samples were analyzed. **J.** Analysis of *Jag1* mRNA
abundance on differentiation days 2, 4, and 8. Mean ± SD (n=3). **K.** Western blot analysis of
JAG1 protein abundance. Mean ± SD (n=3).

*Figure 4: JAG2 subcellular location and length.* Human bronchial basal cells were differentiated
in air-liquid-interface cultures using H&H medium. **A-C.** Immunofluorescence was used to detect
JAG2 (red) and β-catenin (green) on proliferation day 5 (**A**), differentiation day 4 (**B**), or
differentiation day 8 (**C**). Nuclei were detected with DAPI (blue). Each panel presents single
color and merged images. Arrows: white, cortical JAG2; yellow, JAG2-high cells; blue, JAG2-
positive cells; pink, JAG2-negative cells. **D.** Immunofluorescence analysis of HES1 (red) and
RBPJK (red) on differentiation day 8. Nuclei were detected with DAPI (blue). **E.** Western blot
analysis of JAG2 on differentiation days 4 (lane 1) and 8 (lane 2). The full-length band is labeled
JAG2. **F**. Co-immunoprecipitation analysis of JAG2 and β-catenin on differentiation day 8.
Proteins were immunoprecipitated with an N-terminus (N-term) specific β-catenin antibody.
Precipitates were analyzed for β-catenin using a C-terminus (C-term) specific β-catenin
antibody, N-term specific β-catenin antibody, or a JAG2 N-term specific antibody. Three
samples were analyzed. **G.** Analysis of *Jag2* mRNA abundance on differentiation days 2, 4, and
8. Mean ± SD (n=3). **H.** Western blot analysis of JAG2 protein abundance. Mean ± SD, N=3. **G.**

*Figure 5: Regulation of JAG1 and JAG2 abundance.* Human bronchial basal cells were
differentiated in air-liquid-interface cultures using H&H medium. **A-B.** On differentiation days 4
and 6, the cultures were treated with vehicle, (Veh, DMSO), CHIR99021 (CHIR), XAV939
(XAV), or LGK974 (LGK) and protein lysates were collected on day 8. Drug concentrations are
in μM. Western blots were used to quantify JAG1 (**A**) and JAG2 (**B**) abundance. Mean ± SD,
N=3. **C-D.** Cultures were treated with Veh or DKK1 on differentiation days 0 and 2 and lysed on
day 4 or treated on differentiation days 4 and 6 and lysed on day 8. Drug concentrations are in
ng/ml. Western blots were used to quantify JAG1 (**C**) and JAG2 (**D**) abundance. Mean ± SD,
N=3. **E-F**. Uniform manifold approximation and projection (UMAP) analysis of cells that were

treated with vehicle (**E**) or CHIR (**F**) on days 4 and 6 and harvested for scRNAseq analysis on

day 8. **G.** Number of normalized cells per Surat Cluster.

*Figure 6: JAG1 and JAG2 trafficking to the cell surface.* Human bronchial basal cells were

differentiated in air-liquid-interface cultures using H&H medium. **A-B**. On differentiation day 8,

cultures were fixed under non-permeabilization conditions and JAG1 (green) and JAG2 (red)

were detected by dual immunofluorescence staining. Nuclei were stained with DAPI (blue).

Panel B is a higher magnification of the region indicated in panel A. **C-F.** After imaging, the

cultures were permeabilized and re-stained for JAG1 (green), JAG2 (red), and DAPI (blue).

Arrows: antigen-positive cells. **G.** On differentiation day 8, cell surface proteins were labeled

with biotin, biotinylated proteins were recovered, and the bound (BIO) and unbound (UnB)

fractions were evaluated by western blot. Two samples were evaluated. **H.** On differentiation

day 8, cell lysates were reacted with TUBE beads which bind to proteins decorated with 4 or

more ubiquitin moieties. The bound fraction was treated with vehicle (-DUB) or a broad

spectrum debubiquitinase (+DUB). Western blots were generated from the unfractionated (In),

unbound (Un), -DUB, and +DUB samples and were evaluated for JAG1 and JAG2.

*Ubiquitinated JAG2, **deubiquitinated JAG2. Immunofluorescence was used to detect JAG2 in

Veh (**I**) and CHIR (5 µM, **J**) treated cultures on differentiation day 8. Arrows: red, JAG2-high cell;

white, JAG2-negative cell; yellow, redistributed JAG2. Asterisks: JAG2-low cells. **K.** Cells were

cultured to differentiation day 8, lysed, and immunoprecipitated with a phospho-serine/phospho-

threonine (pSer/pThr) antibody. Precipitates were analyzed for JAG2 using the C-terminus

specific antibody, KRT5 antibody, or a pan-YWHA antibody. Two samples were analyzed.

*Figure 7: Roles for JAG1 and JAG2 in ciliated and goblet cell differentiation.* Human bronchial

basal cells were differentiated in air-liquid-interface cultures using H&H medium. Treatments

were on differentiation days 8 and 10 and cultures were fixed on day 12. Differentiation was

quantified by immunofluorescence analysis of acetylated tubulin, MUC5B, and DAPI. **A-B.** Cells

were treated with vehicle (DMSO) or 40 µM JAG1 peptide. Frequency of ciliated cell

differentiation intermediates. P-cells with a primary cilium, B-bristle cells, M-ciliated cells (**A**).

Frequency of goblet cell differentiation intermediates (**B**). L-MUC5B-low cells, M-MUC5B-

medium cells, H-MUC5B-high cells. **C-F.** Cells were treated with vehicle (PBS) or 25 µg/ml

neutralizing antibody to JAG1 (αJ1) or JAG2 (αJ2). Frequency of bristle cells (**C**). Frequency of

ciliated cells (**D**). Frequency of MUC5B-medium cells (**E**). Frequency of MUC5B-high cells (**F**).

**G-J.** Cells were treated with PBS or 25 µg/ml αJ1 and 25 µg/ml αJ2. Frequency of bristle cells

(**G**). Frequency of ciliated cells (**H**). Frequency of MUC5B-medium cells (**I**). Frequency of

MUC5B-high cells (**J**). Mean ± SD, N=6.

*Supplemental Figure 1: Ciliated and goblet cell differentiation as a function of medium type.*

Human bronchial basal cells were differentiated in air-liquid-interface cultures using three

media: Wu, H&H (H&H), and complete Pneumacult (cPC). Cultures were fixed on days 7 and

14. **A.** Cell density was determined by quantifying the number of DAPI-stained nuclei per unit

area. **B.** Ciliated cells were identified by acetylated tubulin (ACT) staining and their frequency

was reported as the number ACT+ cells/number nuclei. **C.** Goblet cells were identified by

MUC5B staining and their frequency was reported as the number MUC5B+ cells/number nuclei.

Mean ± SD, N=6.

*Supplemental Figure 2: Intermediate ciliated and goblet cell phenotypes.* Human bronchial basal

cells were differentiated in air-liquid-interface cultures using H&H medium. **A-B.** Cells defined by

a primary cilium were identified by ACT staining on differentiation day 4. Panel B is a high

magnification image of the region identified in panel A. Arrows: white, primary cilium; yellow,

mitotic figure.  **C-D.** Cells with bristle morphology were identified by ACT staining on

differentiation day 8. Panel D is a high magnification image of the region identified in panel C.

Arrows: bristle cells.  **E-F.** Ciliated cells defined were identified by ACT staining on differentiation

day 12. Panel F is a high magnification image of the region identified in panel E. Arrows:

Ciliated cells. G-H. Goblet cells were identified by MUC5B staining on differentiation day 8.

Panel H is a high magnification image of the region identified in panel G. Arrows identify

MUC5B-low, -medium, and -high goblet cell intermediates.

*Supplemental Figure 3: Cell density in vehicle and DAPT treated cultures.* Human bronchial

basal cells were differentiated in air-liquid-interface cultures using H&H medium. Cells were

treated with vehicle (DMSO) or 25 µM DAPT as follows: treatment on differentiation days 0 and

2 and fix on day 4, treatment on differentiation days 2 and 4 and fix on day 6, treatment on

differentiation days 4 and 6 and fix on day 8, or treatment on differentiation days 8 and 10 and

fix on day 12. Nuclei were stained with DAPI and quantified. Mean ± SD, N=6.

*Supplemental Figure 4: JAG1 and JAG2 localization as a function of medium type.* Human

bronchial basal cells were differentiated in air-liquid-interface cultures using three media: Wu (**A,**

**D, G, J**), H&H (H&H, **B, E, H, K**), and complete Pneumacult (cPC, **C, F, I, L**). Cultures were

fixed on day 7. JAG1 (green, **A-C**) and JAG2 (red, **D-F**) were detected by dual

immunofluorescence. Nuclei were detected with DAPI (blue, **G-I**). Merged images (**J-L**). Arrows:

yellow, JAG2 expression; white, low JAG2 expression. Scale bar in panel A, 100 µm.

*Supplemental Figure 5: Gene expression and gene set enrichment analysis.* ALI cultures were
treated with vehicle, 10 µM CHIR or 10 µM XAV on differentiation days 4 and 6 and RNA was
purified on day 8. Transcriptional changes were interrogated by RNA-sequencing and Gene Set
Expression Analysis (GSEA). **A.** GSEA for WNT/β-catenin genes in vehicle and CHIR treated
cultures. **B.** Heat map representation of WNT/β-catenin genes in vehicle and CHIR treated
cultures. Each column is a sample: samples 1-3 vehicle treated (VEH), samples 4-6 CHIR
treated. Each row is a gene. Red indicates upregulation. Blue indicates downregulation.
Leading-edge genes are indicated by the green line. Jag1 and Jag2 are indicated by the
asterisks. **C.** GSEA for WNT/β-catenin genes in vehicle and XAV treated cultures. **D.** Heat map
representation of WNT/β-catenin genes in vehicle and XAV treated cultures. Each column is a
sample: samples 1-3 vehicle (VEH treated), samples 7-9 XAV treated. Each row is a gene. Red
indicates upregulation. Blue indicates downregulation. Leading-edge genes are indicated by the
green line. Jag1 and Jag2 are indicated by the asterisks. **E.** GSEA for Notch genes in vehicle
and CHIR treated cultures. **F.** Heat map representation of Notch genes in vehicle and CHIR
treated cultures. Each column is a sample: samples 1-3 vehicle (VEH treated), samples 4-6
CHIR treated. Each row is a gene. Red indicates upregulation. Blue indicates downregulation.
Leading-edge genes are indicated by the green line. Jag1 and Jag2 are indicated by the
asterisks. **G.** GSEA for Notch genes in vehicle and XAV treated cultures. **H.** Heat map
representation of Notch genes in vehicle and XAV treated cultures. Each column is a sample:
samples 1-3 vehicle (VEH treated), samples 7-9 XAV treated. Each row is a gene. Red
indicates upregulation. Blue indicates downregulation. Leading-edge genes are indicated by the
green line. Jag1 and Jag2 are indicated by the asterisks. **I.** Analysis of *Jag1* gene expression in
vehicle, CHIR, and XAV treated cultures. Data are presented as the Log2(fold change). Mean ±
SD (n=3). No significant differences were detected. **J.** Analysis of *Jag2* gene expression in
vehicle, CHIR, and XAV treated cultures. Data are presented as the Log2(fold change). Mean ±
SD (n=3). No significant differences were detected.

*Supplemental Figure 6: Analysis of leading-edge genes.* ALI cultures were treated with vehicle,
10 µM CHIR or 10 µM XAV on differentiation days 4 and 6 and RNA was purified on day 8.
Transcriptional changes were interrogated by RNA-sequencing and Gene Set Expression
Analysis (GSEA). Leading-edge genes, which determine the gene set Enrichment Score, were
extracted and organized to indicate similarities/differences across the treatments and gene sets.
Red indicates downregulation. Blue indicates upregulation. Transc: Transcriptional targets of
WNT/β-catenin were based on literature reports. (transc): WNT/β-catenin targets identified by

similarity. Protein: Targets of glycogen synthase kinase (GSK3) and tankyrase (TNKS) were

based on literature reports. (protein): GSK3 and TNKS targets identified by similarity.

*Supplemental Figure 7:* Roles for JAG1 and JAG2 in tracheobronchial cell fate decisions. **A**.

JAG1 or JAG2 permit Basal Progenitor Cell (BPC) differentiation to a Secretory Progenitor Cell

(SPC) and SPC differentiation to a bristle cell (green light). Both JAG1 and JAG2 are required

for SPC differentiation to a goblet cell (green light). **B**. JAG2 inhibits the Squamous Cell fate

(red light). **C.** JAG2 pauses progression from one ciliated cell differentiation intermediate to the

next (yellow light).

**REFERENCES**

1.      Hong KU, Reynolds SD, Watkins S, Fuchs E, Stripp BR. Basal cells are a multipotent

progenitor capable of renewing the bronchial epithelium. The American journal of pathology.

2004;164(2):577-88.

2.      Hong KU, Reynolds SD, Watkins S, Fuchs E, Stripp BR. In vivo differentiation potential

of tracheal basal cells: Evidence for multipotent and unipotent subpopulations. American journal

of physiology Lung cellular and molecular physiology. 2003.

3.      Ghosh M, Brechbuhl HM, Smith RW, Li B, Hicks DA, Titchner T, et al. Context-

dependent differentiation of multipotential keratin 14-expressing tracheal basal cells. American

journal of respiratory cell and molecular biology. 2011;45(2):403-10.

4.      Rock JR, Onaitis MW, Rawlins EL, Lu Y, Clark CP, Xue Y, et al. Basal cells as stem

cells of the mouse trachea and human airway epithelium. Proceedings of the National Academy

of Sciences of the United States of America. 2009;106(31):12771-5.

5.      Engelhardt JF, Schlossberg H, Yankaskas JR, Dudus L. Progenitor cells of the adult

human airway involved in submucosal gland development. Development. 1995;121(7):2031-46.

6.      Carraro G, Stripp BR. A new notch for lung stem cells. Cell stem cell. 2015;16(2):107-9.

7.      Kiyokawa H, Morimoto M. Notch signaling in the mammalian respiratory system,

specifically the trachea and lungs, in development, homeostasis, regeneration, and disease.

Dev Growth Differ. 2020;62(1):67-79.

8.      Radtke F, Raj K. The role of Notch in tumorigenesis: oncogene or tumour suppressor?

Nature reviews Cancer. 2003;3(10):756-67.

9.      Kopan R, Ilagan MX. The canonical Notch signaling pathway: unfolding the activation

mechanism. Cell. 2009;137(2):216-33.

10.     Ilagan MX, Kopan R. SnapShot: notch signaling pathway. Cell. 2007;128(6):1246.

11.     Castel D, Mourikis P, Bartels SJ, Brinkman AB, Tajbakhsh S, Stunnenberg HG. Dynamic

binding of RBPJ is determined by Notch signaling status. Genes & development.

2013;27(9):1059-71.

12.     Vieira Braga FA, Kar G, Berg M, Carpaij OA, Polanski K, Simon LM, et al. A cellular

census of human lungs identifies novel cell states in health and in asthma. Nature medicine.

2019;25(7):1153-63.

13.     Carraro G, Mulay A, Yao C, Mizuno T, Konda B, Petrov M, et al. Single Cell

Reconstruction of Human Basal Cell Diversity in Normal and IPF Lung. American journal of

respiratory and critical care medicine. 2020.

14.     Goldfarbmuren KC, Jackson ND, Sajuthi SP, Dyjack N, Li KS, Rios CL, et al. Dissecting

the cellular specificity of smoking effects and reconstructing lineages in the human airway

epithelium. Nature communications. 2020;11(1):2485.

15.     Bodas M, Moore AR, Subramaniyan B, Georgescu C, Wren JD, Freeman WM, et al.

Cigarette Smoke Activates NOTCH3 to Promote Goblet Cell Differentiation in Human Airway

Epithelial Cells. American journal of respiratory cell and molecular biology. 2021;64(4):426-40.

16.     Verhein KC, McCaw Z, Gladwell W, Trivedi S, Bushel PR, Kleeberger SR. Novel Roles

for Notch3 and Notch4 Receptors in Gene Expression and Susceptibility to Ozone-Induced

Lung Inflammation in Mice. Environmental health perspectives. 2015;123(8):799-805.

17.     Xu F, Gao J, Bergmann S, Sims AC, Ashbrook DG, Baric RS, et al. Genetic Dissection

of the Regulatory Mechanisms of Ace2 in the Infected Mouse Lung. Front Immunol.

2020;11:607314.

18.    Qin L, Qiu KZ, Hu CP, Wu GJ, Wang LL, Tan YR. Bronchial Epithelial Cells Promote the

Differentiation of Th2 Lymphocytes in Airway Microenvironment through Jagged/Notch-1

Signaling after RSV Infection. Int Arch Allergy Immunol. 2019;179(1):43-52.

19.    Tsao PN, Wei SC, Wu MF, Huang MT, Lin HY, Lee MC, et al. Notch signaling prevents

mucous metaplasia in mouse conducting airways during postnatal development. Development.

2011;138(16):3533-43.

20.    Morimoto M, Nishinakamura R, Saga Y, Kopan R. Different assemblies of Notch

receptors coordinate the distribution of the major bronchial Clara, ciliated and neuroendocrine

cells. Development. 2012;139(23):4365-73.

21.    Rock JR, Gao X, Xue Y, Randell SH, Kong YY, Hogan BL. Notch-dependent

differentiation of adult airway basal stem cells. Cell stem cell. 2011;8(6):639-48.

22.    Gomi K, Arbelaez V, Crystal RG, Walters MS. Activation of NOTCH1 or NOTCH3

signaling skews human airway basal cell differentiation toward a secretory pathway. PloS one.

2015;10(2):e0116507.

23.    Reid AT, Nichol KS, Chander Veerati P, Moheimani F, Kicic A, Stick SM, et al. Blocking

Notch3 Signaling Abolishes MUC5AC Production in Airway Epithelial Cells from Individuals with

Asthma. American journal of respiratory cell and molecular biology. 2020;62(4):513-23.

24.    Stupnikov MR, Yang Y, Mori M, Lu J, Cardoso WV. Jagged and Delta-like ligands

control distinct events during airway progenitor cell differentiation. eLife. 2019;8.

25.    Paul MK, Bisht B, Darmawan DO, Chiou R, Ha VL, Wallace WD, et al. Dynamic changes

in intracellular ROS levels regulate airway basal stem cell homeostasis through Nrf2-dependent

Notch signaling. Cell stem cell. 2014;15(2):199-214.

26.    Mori M, Mahoney JE, Stupnikov MR, Paez-Cortez JR, Szymaniak AD, Varelas X, et al.

Notch3-Jagged signaling controls the pool of undifferentiated airway progenitors. Development.

2015;142(2):258-67.

27.    Gomi K, Staudt MR, Salit J, Kaner RJ, Heldrich J, Rogalski AM, et al. JAG1-Mediated

Notch Signaling Regulates Secretory Cell Differentiation of the Human Airway Epithelium. Stem

Cell Rev Rep. 2016;12(4):454-63.

28.    Zhang S, Loch AJ, Radtke F, Egan SE, Xu K. Jagged1 is the major regulator of Notch-

dependent cell fate in proximal airways. Developmental dynamics : an official publication of the

American Association of Anatomists. 2013;242(6):678-86.

29.    Lafkas D, Shelton A, Chiu C, de Leon Boenig G, Chen Y, Stawicki SS, et al. Therapeutic

antibodies reveal Notch control of transdifferentiation in the adult lung. Nature.

2015;528(7580):127-31.

30.    Carrer M, Crosby JR, Sun G, Zhao C, Damle SS, Kuntz SG, et al. Antisense

Oligonucleotides Targeting Jagged 1 Reduce House Dust Mite-induced Goblet Cell Metaplasia

in the Adult Murine Lung. American journal of respiratory cell and molecular biology.

2020;63(1):46-56.

31.    Plasschaert LW, Zilionis R, Choo-Wing R, Savova V, Knehr J, Roma G, et al. A single-

cell atlas of the airway epithelium reveals the CFTR-rich pulmonary ionocyte. Nature.

2018;560(7718):377-81.

32.    Montoro DT, Haber AL, Biton M, Vinarsky V, Lin B, Birket SE, et al. A revised airway

epithelial hierarchy includes CFTR-expressing ionocytes. Nature. 2018;560(7718):319-24.

33.    Jain R, Pan J, Driscoll JA, Wisner JW, Huang T, Gunsten SP, et al. Temporal

relationship between primary and motile ciliogenesis in airway epithelial cells. American journal

of respiratory cell and molecular biology. 2010;43(6):731-9.

34.    Boutin C, Kodjabachian L. Biology of multiciliated cells. Current opinion in genetics &

development. 2019;56:1-7.

35.    Malleske DT, Hayes D, Jr., Lallier SW, Hill CL, Reynolds SD. Regulation of human

airway epithelial tissue stem cell differentiation by beta-catenin, P300, and CBP. Stem Cells.

2018.

36.    Seibold MA, Smith RW, Urbanek C, Groshong SD, Cosgrove GP, Brown KK, et al. The

idiopathic pulmonary fibrosis honeycomb cyst contains a mucocilary pseudostratified epithelium.

PloS one. 2013;8(3):e58658.

37.    Fulcher ML, Randell SH. Human nasal and tracheo-bronchial respiratory epithelial cell

culture. Methods Mol Biol. 2013;945:109-21.

38.    Hayes D, Jr., Kopp BT, Hill CL, Lallier SW, Schwartz CM, Tadesse M, et al. Cell Therapy

for Cystic Fibrosis Lung Disease: Regenerative Basal Cell Amplification. Stem Cells Transl Med.

2018.

39.    Reynolds SD, Rios C, Wesolowska-Andersen A, Zhuang Y, Pinter M, Happoldt C, et al.

Airway Progenitor Clone Formation is Enhanced by Y-27632-dependent Changes in the

Transcriptome. American journal of respiratory cell and molecular biology. 2016.

40.    Suprynowicz FA, Upadhyay G, Krawczyk E, Kramer SC, Hebert JD, Liu X, et al.

Conditionally reprogrammed cells represent a stem-like state of adult epithelial cells.

Proceedings of the National Academy of Sciences of the United States of America.

2012;109(49):20035-40.

41.    Wu R. Culture of normal human airway epithelial cells and measurement of mucin

synthesis and secretion. Methods in molecular medicine. 2000;44:31-9.

42.    Lallier SW, Hill CL, Nichols DP, Reynolds SD. Protein Abundance Determination: An

Optimized Western Blot Workflow. Ann Clin Lab Sci. 2019;49(4):507-12.

43.    Carraro G, Langerman J, Sabri S, Lorenzana Z, Purkayastha A, Zhang G, et al.

Transcriptional analysis of cystic fibrosis airways at single-cell resolution reveals altered

epithelial cell states and composition. Nature medicine. 2021;27(5):806-14.

44.    Young MD, Behjati S. SoupX removes ambient RNA contamination from droplet-based

single-cell RNA sequencing data. Gigascience. 2020;9(12).

45.    Wolock SL, Lopez R, Klein AM. Scrublet: Computational Identification of Cell Doublets in

Single-Cell Transcriptomic Data. Cell Syst. 2019;8(4):281-91 e9.

46.    Lun AT, McCarthy DJ, Marioni JC. A step-by-step workflow for low-level analysis of

single-cell RNA-seq data with Bioconductor. F1000Res. 2016;5:2122.

47.    Wolf FA, Hamey FK, Plass M, Solana J, Dahlin JS, Gottgens B, et al. PAGA: graph

abstraction reconciles clustering with trajectory inference through a topology preserving map of

single cells. Genome biology. 2019;20(1):59.

48.    Becht E, McInnes L, Healy J, Dutertre CA, Kwok IWH, Ng LG, et al. Dimensionality

reduction for visualizing single-cell data using UMAP. Nature biotechnology. 2018.

49.    Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK, et al. MAST: a flexible

statistical framework for assessing transcriptional changes and characterizing heterogeneity in

single-cell RNA sequencing data. Genome biology. 2015;16:278.

50.    Tirosh I, Izar B, Prakadan SM, Wadsworth MH, 2nd, Treacy D, Trombetta JJ, et al.

Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq.

Science. 2016;352(6282):189-96.

51.    LaVoie MJ, Selkoe DJ. The Notch ligands, Jagged and Delta, are sequentially

processed by alpha-secretase and presenilin/gamma-secretase and release signaling

fragments. The Journal of biological chemistry. 2003;278(36):34427-37.

52.    Ikeuchi T, Sisodia SS. The Notch ligands, Delta1 and Jagged2, are substrates for

presenilin-dependent "gamma-secretase" cleavage. The Journal of biological chemistry.

2003;278(10):7751-4.

53.    Metrich M, Bezdek Pomey A, Berthonneche C, Sarre A, Nemir M, Pedrazzini T. Jagged1

intracellular domain-mediated inhibition of Notch1 signalling regulates cardiac homeostasis in

the postnatal heart. Cardiovasc Res. 2015;108(1):74-86.

54.    An WF, Germain AR, Bishop JA, Nag PP, Metkar S, Ketterman J, et al. Discovery of

Potent and Highly Selective Inhibitors of GSK3b.  Probe Reports from the NIH Molecular

Libraries Program. Bethesda (MD)2010.

55.    Huang SM, Mishina YM, Liu S, Cheung A, Stegmeier F, Michaud GA, et al. Tankyrase

inhibition stabilizes axin and antagonizes Wnt signalling. Nature. 2009;461(7264):614-20.

56.    Araya J, Cambier S, Markovics JA, Wolters P, Jablons D, Hill A, et al. Squamous

metaplasia amplifies pathologic epithelial-mesenchymal interactions in COPD patients. The

Journal of clinical investigation. 2007;117(11):3551-62.

57.    Sun J, Luo Z, Wang G, Wang Y, Wang Y, Olmedo M, et al. Notch ligand Jagged1

promotes mesenchymal stromal cell-based cartilage repair. Exp Mol Med. 2018;50(9):1-10.

58.    Dontu G, Jackson KW, McNicholas E, Kawamura MJ, Abdallah WM, Wicha MS. Role of

Notch signaling in cell-fate determination of human mammary stem/progenitor cells. Breast

Cancer Res. 2004;6(6):R605-15.

59.    Hellstrom M, Phng LK, Hofmann JJ, Wallgard E, Coultas L, Lindblom P, et al. Dll4

signalling through Notch1 regulates formation of tip cells during angiogenesis. Nature.

2007;445(7129):776-80.

60.    Ezratty EJ, Pasolli HA, Fuchs E. A Presenilin-2-ARF4 trafficking axis modulates Notch

signaling during epidermal differentiation. The Journal of cell biology. 2016;214(1):89-101.

61.    Tan FE, Vladar EK, Ma L, Fuentealba LC, Hoh R, Espinoza FH, et al. Myb promotes

centriole amplification and later steps of the multiciliogenesis program. Development.

2013;140(20):4277-86.

62.    El-Osta MA, Idkowiak-Baldys J, Hannun YA. Delayed phosphorylation of classical

protein kinase C (PKC) substrates requires PKC internalization and formation of the pericentrion

in a phospholipase D (PLD)-dependent manner. The Journal of biological chemistry.

2011;286(22):19340-53.

63.    Fortini ME, Bilder D. Endocytic regulation of Notch signaling. Current opinion in genetics

& development. 2009;19(4):323-8.

64.    Taelman VF, Dobrowolski R, Plouhinec JL, Fuentealba LC, Vorwald PP, Gumper I, et al.

Wnt signaling requires sequestration of glycogen synthase kinase 3 inside multivesicular

endosomes. Cell. 2010;143(7):1136-48.

65.    Vinyoles M, Del Valle-Perez B, Curto J, Vinas-Castells R, Alba-Castellon L, Garcia de

Herreros A, et al. Multivesicular GSK3 sequestration upon Wnt signaling is controlled by p120-

catenin/cadherin interaction with LRP5/6. Molecular cell. 2014;53(3):444-57.

66.    Zheng L, Conner SD. Glycogen synthase kinase 3beta inhibition enhances Notch1

recycling. Molecular biology of the cell. 2018;29(4):389-95.

67.    de Carvalho A, Strikoudis A, Liu HY, Chen YW, Dantas TJ, Vallee RB, et al. Glycogen

synthase kinase 3 induces multilineage maturation of human pluripotent stem cell-derived lung

progenitors in 3D culture. Development. 2019;146(2).

## Graphical Abstract

# Figure 1

# Figure 2

## A

|  | Day 4 | Day 8 | Day 12 |
|---|---|---|---|
| Recovery | LY | DMSO | |
| Continuous | LY | | |

## B

## C

## D

## E

Figure 3

# Figure 4

Figure 5

# Figure 6

Not Permeabilized        Permeabilized

A
JAG1
JAG2
DAPI
5μm

B
JAG1
JAG2
DAPI
20 μm

C
JAG1
20 μm

D
JAG2
20 μm

E
DAPI
20 μm

F
Merge
20 μm

G
| | Bio | UnB | Bio | UnB |
|------|-----|-----|-----|-----|
| JAG1 | | | | |
| JAG2 | | | | |
| NKA | | | | |

H
| | In | Un | -DUB | +DUB |
|------|----|----|------|------|
| JAG1 | | | | |
| JAG2 | | | * | ** |

I
JAG2
DAPI
10 μm

J
JAG2
DAPI
10 μm

K
pSer/pThr IP
Western Blot    1    2
JAG2
KRT5
YWHA

Figure 7

# Supplemental Figure 1

A

B

C

# Supplemental Figure 2

# Supplemental Figure 3

# Supplemental Figure 4

## Diferentiation Day 7

Wu Medium      H&H Medium      cPC Medium

A — Jag1    B    C

100 μm

D — Jag2    E    F

G — DAPI    H    I

J — Merge    K    L

# Supplemental Figure 5

**A**

Enrichment Score = -0.427
FDR q-value = 0.0332
Leading Edge Genes = 14

**C**

Enrichment Score = +0.578
FDR q-value = 0.0169
Leading Edge Genes = 13

**E**

Enrichment Score = +0.368
FDR q-value = 0.506
Leading Edge Genes = 9

**G**

Enrichment Score = +0.567
FDR q-value = 0.0379
Leading Edge Genes = 10

# Supplemental Figure 6

| Pathway / Gene | WNT | | NOTCH | | Target | | | Full Gene Name |
|---|---|---|---|---|---|---|---|---|
| Treatment | CHIR | XAV | CHIR | XAV | bCat | GSK3 | TNKS | |
| *Axin2* | Up | Down | | | Transc | (Protein) | Protein | *Axis Inhibition Protein 2* |
| *Lef1* | Up | Down | | | Transc | | | Lymphoid Enhancer Binding Factor 1 |
| *Tcf7* | Up | Down | | | Transc | | | Transcription Factor 7 |
| *Wnt5b* | Up | Down | | | | | | Wnt Family Member 5B |
| *Dll1* | | Down | Down | Down | Transc | | | Delta Like Canonical Notch Ligand 1 |
| *Jag1* | | Down | Down | Down | Transc | | | Jagged Canonical Notch Ligand 1 |
| *Notch1* | | Down | Down | Down | | Protein | Protein | Notch Receptor 1 |
| *Lfng* | | | Down | Down | | | | LFNG O-Fucosylpeptide 3-Beta-N-Acetylglucosaminyltransferase |
| *Prkca* | | | Down | Down | | | | Protein Kinase C Alpha |
| *Psenen* | | | Down | Down | | | | Presenilin Enhancer, Gamma-Secretase Subunit |
| *Sap30* | | | Down | Down | | | | Sin3A Associated Protein 30 |
| *St3gal6* | | | Down | Down | | | | ST3 Beta-Galactoside Alpha-2,3-Sialyltransferase 6 |
| *Axin1* | Up | | | | | Protein | Protein | *Axis Inhibition Protein 1* |
| *Ccnd2* | Up | | | | | | | Cyclin D2 |
| *Ctnnb1* | Up | | | | | | Protein | Catenin Beta 1 |
| *Dkk1* | Up | | | | Transc | | | Dickkopf WNT Signaling Pathway Inhibitor 1 |
| *Dkk4* | Up | | | | (Transc) | | | Dickkopf WNT Signaling Pathway Inhibitor 4 |
| *Dvl2* | Up | | | | | | | Dishevelled Segment Polarity Protein 2 |
| *Fdz1* | Up | | | | | | | Frizzled Class Receptor 1 |
| *Nkd1* | Up | | | | | | (Protein) | NKD Inhibitor Of WNT Signaling Pathway 1 |
| *Psen2* | Up | | | | | | | Presenilin 2 |
| *Ptch1* | Up | | | | | | | Patched 1 |
| *Gnai1* | | Down | | | | | | G Protein Subunit Alpha I1 |
| *Hey1* | | Down | | | | | | Hes Related Family BHLH Transcription Factor With YRPW Motif 1 |
| *Jag2* | | Down | | | | | | Jagged Canonical Notch Ligand 2 |
| *Notch4* | | Down | | | | Protein | (Protein) | Notch Receptor 4 |
| *Skp2* | | Down | | | | | | Hes Related Family BHLH Transcription Factor With YRPW Motif 1 |
| *Ccdn1* | | | Down | | | | | Cyclin D1 |
| *Dtx4* | | | Down | | | | | Deltex E3 Ubiquitin Ligase 4 |
| *Maml2* | | | Down | | | | | Mastermind Like Transcriptional Coactivator 2 |
| *Arrb1* | | | | Down | | | | Arrestin Beta 1 |
| *Heyl* | | | | Down | | | | Hes Related Family BHLH Transcription Factor With YRPW Motif Like |

🟥 Upregulated in treated cells
🟦 Downregulated in treated cells
Transc: transcriptional
(Transc): based on similarity
(Protein): based on similarity

# Supplemental Figure 7

## A

BPC

JAG1 or JAG2

SPC

JAG1 or JAG2 → Bristle

JAG1 and JAG2

Goblet Cell

## B

BPC

Squamous

JAG2

SPC

Basal Progenitor Cell (BPC)
Secretory Progenitor Cell (SPC)

Stop Light Analogy
Green: Permitted fate
Yellow: Paused fate
Red: Inhibited fate

## C

Primary Cilium

JAG2

Bristle

JAG2

Ciliated Cell