A predicted developmental and evolutionary morphospace for grapevine leaves

Daniel H. Chitwood[1,2,*], Joey Mullins[1]
[1]Dept. Horticulture, [2]Dept. Computational Mathematics, Science & Engineering, Michigan State University, East Lansing, MI 48823 USA

*Correspondence: Daniel H. Chitwood, chitwoo9@msu.edu

Word count: 4,268

ABSTRACT

Using conventional statistical approaches there exist powerful methods to classify shapes. Embedded in morphospaces is information that allows us to visualize theoretical leaves. These unmeasured leaves are never considered nor how the negative morphospace can inform us about the forces responsible for shaping leaf morphology. Here, we model leaf shape using an allometric indicator of leaf size, the ratio of vein to blade areas. The borders of the observable morphospace are restricted by constraints and define an orthogonal grid of developmental and evolutionary effects which can predict the shapes of possible grapevine leaves. Leaves in the genus *Vitis* are found to fully occupy morphospace available to them. From this morphospace we predict the developmental and evolutionary shapes of grapevine leaves that are not only possible, but exist, and argue that rather than explaining leaf shape in terms of discrete nodes or species, that a continuous model is more appropriate.

INTRODUCTION

Leaf shape across plants is diverse and spectacular, but it is not random. Development, evolution, and the environment sculpt leaf shape in specific ways (Chitwood and Sinha, 2016). Leaves allometrically expand, first shown by Stephen Hales through pin pricks on developing fig leaves that were displaced differentially along the length versus the width of the leaf (1727). The developmental programing of leaves changes from node-to-node resulting in changing leaf shapes. Goethe described this process as "metamorphosis" and in terms of the mutable, changing internal state of leaves (1817). Environment modulates leaf size and serrations, as observed by Bailey and Sinnott (1915) who used the distribution of entire leaves across latitudes to estimate the temperatures of paleoclimates. If we measure leaf shape across the seed plants, clear demarcations between phylogenetic groups are observed (Li et al., 2018). We have measured enough leaf shapes to know the borders and demarcations of what exists and the processes that shape leaves in specific ways.

The shapes of grapevine leaves have been measured under intense scrutiny and with
purpose. Originally through morphometric techniques developed by Louis Ravaz (1902),
the field of ampelography ("vine" + "process of measuring") sought to discern, using
leaves and other features of the vine, American *Vitis* species that were new to Europeans
and would eventually be used as rootstocks against Phylloxera. Eventually the
techniques would be famously applied to wine grape varieties by Pierre Galet (1979;
1985; 1988; 1990; 2000; Chitwood, 2020). Morphometric techniques have been used to
genetically study the basis of leaf shape in grapevines (Chitwood et al., 2014; Demmings
et al., 2019), how grapevine leaves develop (Chitwood et al., 2016a), the effects of
environment (Chitwood et al., 2016b; Baumgartner et al., 2020), and to show that
increases in vein length compensate for leaf area lost to lobing (Migicovsky et al.,
2022a). Modeling has been used in several ways, including calculating average shapes of
grapevine varieties while preserving features (Martínez et al., 1995; 1997a; 1997b; 1999),
modeling development across grapevine shoots (Bryson et al., 2020), and using leaf
allometry, specifically the ratio of vein to blade areas, as a proxy of leaf size and to
measure the effects of year-to-year variation in leaf shape (Chitwood et al., 2021). For
grapevines, as for many other types of leaves, we have extensively measured and
modeled leaf shape, allowing us to discern genetic, developmental, and environmental
effects with great power.
61
62   But what about leaves that are not available for us to measure? Using what we know
63   about the underlying structure of leaf morphospaces across genotypic, developmental,
64   and environmental effects, and making modeling assumptions about what is and is not
65   possible, could we compare what we have measured and observed against the
66   boundaries of what we know is possible?
67
Here, we measure the shapes of over 8900 grapevine leaves and model them against an
allometric indicator of leaf size, vein-to-blade ratio, across *Vitis* species. The expansion
of blade area at the expense of that for veins is found to be a principal determinant of
the resulting morphospace, as much so as differences in leaf shape between species.
These developmental and evolutionary forces that sculpt leaf shape are independent
and lie orthogonal to each other. Using an inverse transform of the Principal
Component Analysis (PCA) space, theoretical leaves missing from the data are
reconstructed. We find that the borders of the grapevine leaf morphospace are sharply
defined by developmental constraints of lobing and the ratio of vein-to-blade area and
that leaves in the genus *Vitis* fully occupy the space available to them. Rather than
discrete stages of development or species, for leaf shape, the morphospace is better
described continuously as a grid defined by developmental and evolutionary effects
from which any leaf shape in the genus *Vitis* can be predicted.
81

82 **MATERIALS AND METHODS**
83
This work uses two sources of genetic material to sample grapevine leaf shape, referred
to as "New York germplasm" and "California populations". The first is the USDA
germplasm repository in Geneva, NY which samples mostly North American *Vitis* species
leaves (although not exclusively) as a developmental series, keeping track of the node
the leaves arise from. These leaves tend to be more entire (again, not exclusively so). The
second source of materials are segregating populations in California from E. & J. Gallo
Winery (the exact identity of which is proprietary). The parentage of this material arises
from *Vitis vinifera, V. mustangensis,* and *V. piasezkii* species and is more deeply lobed
than the New York germplasm material (again, this is not always the case). Only mature,
fully expanded leaves from the middle of the shoot were sampled from this population.
This population was not sampled as a developmental series and the node the leaves
arise from was not recorded. The New York germplasm allows models of leaf
development to be estimated whereas the California populations sample additional leaf
shapes throughout the genus *Vitis*. More specific information about each of these
materials is given below.
99
New York germplasm material
As described in Bryson et al., 2020 (and copied verbatim here for convenience), leaves
were collected from 209 vines at the USDA germplasm repository vineyard in Geneva,
New York, USA. Samples were taken from the same vines during the second week of
June, annually, in 2013 and 2015–2017. The vines sampled represent 11 species
(*Ampelopsis glandulosa* (Wall.) Momiy. var. *brevipedunculata* (Maxim.) Momiy., *V.*
acerifolia Raf., *V. aestivalis* Michx., *V. amurensis* Rupr., *V. cinerea* (Engelm.) Millardet, *V.*
coignetiae Pulliat ex Planch., *V. labrusca* L., *V. palmata* Vahl, *V. riparia* Michx., *V.*
rupestris Scheele, and *V. vulpina* L.), four hybrids
(*V. ×andersonii* Rehder, *V. ×champinii* Planch., *V. ×doaniana* Munson ex Viala,
and *V. ×novae-angliae* Fernald), and 13 *Vitis* vines, designated as *Vitis* spp., for which
original species assignments from the germplasm collection are lacking. Starting at the
shoot tip (with shoot order noted for each leaf), leaves greater than ~1 cm in length
were collected in stacks and stored in a cooler in labeled plastic bags with ventilation
holes. Within two days of collection, the leaves were arranged on a large-format Epson
Workforce DS-50000 scanner (Tokyo, Japan) in the order they were collected, with a
small number near each leaf indicating which node it came from and a ruler for scale
within the image file. The image files were named with the vine identification number,
followed by a sequential lowercase letter if multiple scans were needed. The original
scans are available on Dryad (Chitwood et al., 2020).
120
California populations material

As described in Migicovsky et al., 2022a (and copied verbatim here for convenience),
leaves were sampled from seedlings of five biparental *Vitis* populations located in
Madera County, California, USA. 500 seedlings were planted in the vineyard. 450
seedlings shared a seed parent, DVIT 2876. The remaining 50 seedlings had DVIT 2876
as a grandparent. DVIT 2876 'Olmo b55-19' is a compound-leafed accession from the
USDA-ARS National Clonal Germplasm repository, suspected to include *V. piasezkii*
Maximowicz, as one of its parents (or grandparents). The populations were created to
examine variation in leaf lobing. The vines were composed of 125 individuals from a
DVIT 2876 x unnamed *V. vinifera* selection cross (Pop1), 100 individuals from a DVIT
2876 x a different unnamed *V. vinifera* selection cross (Pop2), 150 individuals from a
DVIT 2876 x unnamed *Vitis* hybrid cross (Pop3), 75 individuals from a DVIT 2876 x a
different unnamed *Vitis* hybrid cross (Pop4), and 50 individuals from a seedling (DVIT
2876 x unnamed *V. vinifera* selection) x DVIT 3374 (*V. mustangensis* Buckley) cross
(Pop5). The vines sampled were planted in 2017. They were trained to a unilateral
cordon and spur pruned. Leaf samples were collected on June 22 and July 12 2018, then
again in 2019 on June 14, 19, and July 4. Across the sampling dates within a given year,
a total of three mature, representative leaves were sampled from each of the vines and
placed into labeled plastic bags. The plastic bags were stored in a cooler during
collection and scanned, abaxial side down, later the same day using a flatbed scanner.
Files were named using the accession identification number. The original scans are
available on Dryad (Migicovsky et al., 2022b).
143
Data Analysis
Twenty one landmarks (**Figure 1A**) were placed on one half of each leaf outlining the
midvein, distal vein, proximal vein, and the most proximal branching vein of each of
these major veins as well as distal and proximal lobe sinuses using ImageJ (Abràmoff et
al., 2004). Two landmarks are placed at the base of each vein to measure the width.
Landmarks were superimposed through scaling, translation, rotation, and reflection
using Generalized Procrustes Analysis with the shapes (Dryden and Mardia, 2016)
package in R.
152
Data was analyzed using Python and Jupyter notebooks (Kluyver et al., 2016). Code to
reproduce the analysis in this manuscript can be found at the *Github* repository
DanChitwood/grapevine_morphospace: https://
https://github.com/DanChitwood/grapevine_morphospace). The Jupyter notebook
(grapevine_morphospace.ipynb) comments on the code and also contains a narrative to
guide the reader through the analysis. Calculation of distal lobing is according to Galet
(1979), as the ratio of the distance of the distal sinus to the petiolar junction divided by
the distance of the distal lobe tip to the petiolar junction, such that the distal lobing
value of a completely dissected leaf is 0 and the value of a completely entire leaf is 1.

Figure 1: Grapevine leaf morphology. A) Counting from the shoot tip, Vitis cinerea leaves from node positions 1 (left) and 5 (right), each with respective scale bar, are expanded in detail from the same leaves shown in the panel below. The 21 landmarks used in this study are indicated, as well as ampelographic nomenclature naming morphological features. Note that in the younger leaf that vasculature takes up relatively more area than in the mature leaf. B) For seven different grapevine species analyzed in this study, leaves from the shoot tip to the shoot base are shown with scale bar. Leaf area increases from the shoot tip to the middle of the shoot due to leaf expansion, whereas increases in leaf size from the shoot base to the middle of the shoot in mature leaves are due to heteroblasty.

162 Calculation of the natural log of the ratio of vein to blade area, ln($vein\ to\ blade\ ratio$), is

164 as described in Chitwood et al. (2021) using the shoelace algorithm, also known as

165 Gauss' area formula, to calculate polygon areas as originally described by Meister
166 (1769), where n is the number of polygon vertices defined by x and y coordinates:
167

$$\frac{1}{2}|x_1y_2 + x_2y_3 + \ldots + x_{n-1}y_n + x_ny_1 - x_2y_1 - x_3y_2 - \ldots - x_ny_{n-1} - x_1y_n|$$

Principal Component Analysis (PCA) (and calculation of its inverse) was performed using
the scikit learn decomposition PCA module (Pedregosa et al., 2011). Modeling of
$\ln(vein\ to\ blade\ ratio)$, $\ln(leaf\ area)$, and landmarks as polynomial functions of each
other and shoot position was performed using the np.polyfit and np.poly1d functions
from NumPy (Oliphant, 2006). The curve_fit function from SciPy (Virtanen et al., 2020)
was used to fit a reciprocal function of $\ln(leaf\ area)$ across the shoot. Pandas
(McKinney, 2010) and Matplotlib (Hunter, 2007) were used for data analysis and
visualization.
178
179 **RESULTS**
180
Developmental models of leaf expansion
Previously, we modeled leaf shape continuously across grapevine shoots as a
polynomial function of each Procrustes landmark coordinate value as a function of
normalized node position. Normalized node position is the node number counting from
the shoot tip divided the total number of leaves in a shoot, such that node number is
converted to a 0 to 1 scale, from tip to base (Bryson et al., 2020). We also previously
described the natural log of the ratio of vein to blade area, $\ln(vein\ to\ blade\ ratio)$,
which is more sensitive to leaf area than size itself due to the exponential increases in
blade relative to vein area during development (Chitwood et al., 2021). Before we
explore the limits of the grapevine leaf morphospace, we must first model shape across
development to understand how continuous developmental trajectories change
between species during evolution. But it is important to first understand two
developmental processes that affect leaf size and shape across grapevine shoots. At the
shoot tip and base leaves are smaller (and accordingly $\ln(vein\ to\ blade\ ratio)$ is higher)
than the middle of the shoot where leaves are larger (and $\ln(vein\ to\ blade\ ratio)$ lower)
(**Figure 1B**). At the shoot tip leaves are young and at the shoot base they are mature.
The increases in leaf area (and decreases in $\ln(vein\ to\ blade\ ratio)$) from the shoot tip
to the middle of the shoot are mostly due to the expansion of young leaves as they
mature. However, the increases in leaf area (and decreases in $\ln(vein\ to\ blade\ ratio)$)
from the shoot base to the middle of the shoot occur in mature leaves that have already
expanded. The size and shape differences between mature leaves at the shoot base are
due to heteroblasty, node-to-node differences in leaf morphology that result from the
temporal development of the shoot apical meristem, and not from leaf expansion.

Below, we create models of leaf development to focus on allometric changes due to leaf
expansion and its relationship to the grapevine leaf morphospace. To do so requires us
to separate these confounding effects on leaf shape and size across the grapevine shoot
to the best of our ability.

Figure 2: Modeling ln(vein to blade ratio) and ln(leaf area) as a function of normalized node position. A) The natural log of the ratio of vein-to-blade area, ln(vein to blade ratio), and B) the natural log of leaf area, ln(leaf area), are modeled as 2nd degree polynomials of normalized node position (where 0 is the shoot tip and 1 is the shoot base). The normalized node position values corresponding to the minimum ln(vein to blade ratio) and maximum ln(leaf area) values are indicated by a magenta vertical line and the inverse of the golden ratio is indicated by a gold vertical line. C) In order to model developmental changes due to leaf expansion separate from heteroblastic effects, leaves from the shoot tip to the normalized node position value corresponding to the ln(vein to blade ratio) minimum were isolated and modeled as a reciprocal function of normalized node position. Extrapolated values are shown in dashed line and function asymptotes in purple. D) A linear model of ln(leaf area) as a function of ln(vein to blade ratio).

209
We plotted $\ln(vein\ to\ blade\ ratio)$ versus normalized node position (**Figure 2A**), which
can be modeled as a second-degree polynomial. $\ln(vein\ to\ blade\ ratio)$ is highest at the
shoot tip and reaches its minimum in the middle of the shoot. As expected,
$\ln(leaf\ area)$ versus relative node position correspondingly increases in the middle of
the shoot compared to the shoot tip and base (**Figure 2B**). A curiosity that is perhaps
coincidental, we note that the corresponding normalized node position to the minimum
$\ln(vein\ to\ blade\ ratio)$ and maximum $\ln(leaf\ area)$ values are close to the inverse of
the golden ratio (**Figure 2A-B**). Although this may arise as a developmental
phenomenon, it could also be spurious and warrants further investigation.
219
From previous work we know that allometric changes during grapevine leaf expansion
dominate the morphospace (Chitwood et al., 2016a; Chitwood et al., 2016b; Bryson et
al., 2020). We therefore took leaves from the shoot tip to the normalized node position
value corresponding to the minimum $\ln(vein\ to\ blade\ ratio)$ value across the shoot
(**Figure 2A**) to model shape changes associated with leaf expansion. Assuming that
$\ln(vein\ to\ blade\ ratio)$ approaches ∞ as a normalized node position value of 0 is
approached (leaf initiation, where vein area would dominate) and that another
asymptote is approached as leaves mature (where blade area dominates) a reciprocal
function was fit to the data (**Figure 2C**). Using the model, the context of the collected
data compared to extrapolated leaf shapes that remain unsampled (for example, young
leaf primordia or leaves that continue to mature incrementally past the leaves collected
in this study) can be understood. From these expanding leaves a linear model of
$\ln(leaf\ area)$ as a function of $\ln(vein\ to\ blade\ ratio)$ can be fit (**Figure 2D**). From this
model, using a scaleless measure of leaf shape alone, leaf size can be predicted.
Importantly, for the expanding leaves selected for modeling above, their
$\ln(vein\ to\ blade\ ratio)$ values are always decreasing, and their leaf area values are
always increasing moving away from the shoot tip, separating and unconfounding these
effects from those of heteroblasty (**Figure 1B**).
238
By modeling Procrustes-adjusted coordinate values as a polynomial function of
$\ln(vein\ to\ blade\ ratio)$, we can visualize and compare the developmental trajectories of
different grapevine species (**Figure 3**). Theoretical leaves for the six most represented
Vitis species and *Ampelopsis glandulosa* var. *brevipedunculata* across ten equally spaced
$\ln(vein\ to\ blade\ ratio)$ values from the maximum to minimum (inclusive), show the
shape changes associated with leaf expansion and evolutionary differences between
species. Leaf expansion is mostly achieved through increases in blade area relative to
vein, as well as other changes, such as a wider leaf. These developmental changes in
shape are conserved and distinct from species differences, which affect a different set of
shape features, especially the depth of the distal lobe. These shape changes are

allometric and occur concomitantly with exponential decreases in leaf size. The
developmental models of leaf expansion described above will be projected onto the
morphospace described below to anchor and contextualize the space and to quantify
and compare evolutionary versus developmental sources of shape variance across
grapevine leaves.

Figure 3: Developmental models of leaf shape. Fitting each coordinate value of 21 landmarks as a second-degree polynomial of ln(vein to blade ratio), continuous models of expanding leaves for the seven species shown were created. Inclusive of the maximum and minimum ln(vein to blade ratio) values for each species, corresponding to young and mature leaves, respectively, leaves corresponding to ten equally spaced time points were reconstructed. Estimated leaf areas were estimated from ln(vein to blade ratio) values and 1 cm scale bars for each leaf are shown. Leaf areas are indicated by color.

Morphospace

The developmental models of leaf expansion described above are from a dataset, the
"New York germplasm", where leaves were sampled from shoots and their node position

was recorded. These leaves, from the USDA germplasm repository in Geneva, NY sample
mostly (although not exclusively) North American *Vitis* species that tend to have more
entire leaves (although there are highly dissected leaf samples in the dataset). Largely
missing is shape variation from *V. vinifera* and other highly dissected species. To
supplement the New York germplasm leaves, we added leaves from segregating
populations designed to sample highly lobed genetic material, derived from *V. vinifera*,
V. mustangensis, and *V. piasezkii*, called the "California populations". All leaves from the
California populations are mature, creating an opportunity to predict and extrapolate
the development of these leaves from the New York germplasm. Although not
representing the entirety of mature leaf shape variation within *Vitis*, the two datasets
together comprehensively sample it.
270
To visualize the relationship of New York germplasm to California populations datasets,
and how developmental versus evolutionary sources of leaf shape variation compare, we
performed a Principal Component Analysis (PCA). PCA decomposes multivariate data, in
essence rotating and projecting it onto orthogonal axes (principal components) that
more efficiently explain variation in the data than the original measurements (in this
case, Procrustes-adjusted coordinate values). The inverse of this transformation can be
used to reverse calculate original data, which we will later use to visualize theoretical
leaves in the morphospace. PC1 and PC2 explain 39.7% and 17.6% of the variance in the
data, respectively (~57.3% of the total variance). Within this space, the NY germplasm
and CA population data are roughly orthogonal (perpendicular) to each other (**Figure
4**). One interpretation is that the more entire leaves of the NY germplasm data run along
a developmental continuum, whereas the California populations data only represents
mature leaves but falls on a separate axis representing leaves that are more dissected.
The empty space not covered within the ranges of the two datasets would be predicted
to be the missing developmental variation from the deeply lobed leaves in the California
populations data. Two pieces of evidence support the above interpretation. First, if
developmental models of leaf expansion are projected onto the morphospace, they are
collinear with the distribution of the New York germplasm data, consistent with this axis
of the data representing developmental variation. Second, if $\ln(vein\ to\ blade\ ratio)$
values for theoretical leaves calculated from the inverse transform of the morphospace
are projected back onto it (**Figure 4A**) they too are collinear with the NY germplasm
data. Similarly distal lobing, which varies across species (**Figures 1 and 3**), can also be
calculated and projected back onto the morphospace (**Figure 4B**). Distal lobing runs at
roughly right angles to $\ln(vein\ to\ blade\ ratio)$ values and the CA populations data is
collinear with it. The CA populations data intersects with the NY germplasm data in a
location defined by low $\ln(vein\ to\ blade\ ratio)$ values, consistent with these being

mature leaves.

Figure 4: Morphospace. A morphospace calculated from a Principal Component Analysis (PCA) of all leaves from the New York germplasm (black) and California populations (white). A) ln(vein to blade ratio) values and B) distal lobing values were calculated from reconstructed leaves throughout the morphospace using its inverse transform and colored by magma and virdis color schemes, respectively, as indicated. To orient and contextualize the space, developmental models for seven grapevine species were projected into the space, as indicated by colored lines. Isolines for A) ln(vein to blade ratio) values (solid lines) and B) distal lobing values (dashed lines) are shown and their values provided in the respective plots.

298
299

If developmental variation (indicated by $\ln(vein\ to\ blade\ ratio)$ values, **Figure 4A**) and
evolutionary variation between species (indicated by distal lobing values, **Figure 4B**) are
roughly orthogonal to each other, then even though unsampled, the shapes of
developing leaves that are highly dissected that are missing from the CA populations
data could be predicted. The ability to make this prediction rests on the assumption that
highly dissected leaves would follow a developmental trajectory similar to more entire
species. Evidence that this is the case is observed for the developmental model of
Ampelopsis glandulosa var. *brevipedunculata* (**Figure 4**), which is collinear like the other
models with $\ln(vein\ to\ blade\ ratio)$ values and occupies a space with low distal lobing
values, consistent with its deeply lobed morphology.
310
Beyond stages of leaf development and different species, the morphospace of
grapevine leaves can be described more quantitatively and comprehensively using
$\ln(vein\ to\ blade\ ratio)$ and distal lobing values that define it continuously. Isolines that
fall along the same $\ln(vein\ to\ blade\ ratio)$ and distal lobing values can be calculated so
that they extend to the borders of observable morphospace and sample, in a grid-like
fashion, the space inside. These isolines also sample inferred leaf shapes not
represented in the sampled data, including the missing developmental series from the
CA populations data and leaf primordia younger than those sampled. Theoretical,
reconstructed leaves at the intersection of $\ln(vein\ to\ blade\ ratio)$ and distal lobing
isolines, that sample the limits of the observable morphospace, exhibit the distinct
changes in shape associated with development and evolution (**Figure 5**). Across
developmental series regardless of how deeply lobed leaves are, $\ln(vein\ to\ blade\ ratio)$
decreases and leaves become wider as they expand and increase in size. Similarly, as
$\ln(vein\ to\ blade\ ratio)$ isolines traverse orthogonally to distal lobing isolines, the depth
of the distal lobe is preserved regardless of developmental stage and comprises
evolutionary differences in grapevine leaf shape that are independent of development.
327
DISCUSSION
PC1 and PC2 together explain round 57.3% of the variance in the data, but they
represent the first two major, orthogonal sources of variance and as described (**Figure**
4) highlight natural axes in the data that delimit developmental and evolutionary
boundaries that constrain observable grapevine leaf shapes. $\ln(vein\ to\ blade\ ratio)$ and
distal lobing are only indicators of multivariate signatures of leaf development and
evolution, respectively, that lie orthogonal to each other and define a grid in which
grapevine leaves fully occupy to its limits. One set of boundaries is indicated by distal
lobing values (dashed isolines in **Figure 4B**), defined by leaves with values approaching
zero and completely dissected (like *A. glandulosa* var. *brevipedunculata* or *V. piasezkii*) or
nearly equal to one and lacking any significant lobing (like *V. rupestris*). The other set of
boundaries is indicated by $\ln(vein\ to\ blade\ ratio)$ values (solid isolines in **Figure 4A**)

that asymptotically define developmental constraints. Higher $\ln(vein\ to\ blade\ ratio)$
values are associated with young, expanding leaves in which vein area initially
dominates the leaf until the blade exponentially expands. The developmental models
presented in this analysis work from the assumption that young leaf primordia approach
an asymptote consisting entirely of vein area at initiation (**Figure 2C**). In leaves that are
nearly fully expanded the opposite is true, and they are defined by lower
$\ln(vein\ to\ blade\ ratio)$, in which a small amount of vein area remains, but that blade will
always allometrically expand at a faster rate than vein and approach an asymptote in
which vein area is vanishingly small (**Figure 2C**).
349
The morphospace is unexpectedly simple, providing a predictive framework and
empirical insight into theoretical biological concepts. While the New York germplasm
and California populations data sample most shape variation in *Vitis*, the developmental
information for highly dissected species was missing. Because developmental and
evolutionary axes are nearly orthogonal to each other and describe additive signatures
of leaf morphology, where developmental progressions in leaf shape are conserved
across species and variation defining differences between species is maintained
throughout their development, to extrapolate the leaf shapes missing in this space was
straightforward (**Figure 5**). In theory we talk about evolutionary and developmental
forces describing the organismal form, but definition is lacking: to what degree do they
act separately or are confounded together, do they act additively or do interaction
effects predominate? In the case of grapevine leaves, development and evolution are
orthogonal and acting independently of each other to such an extent that rather than
describe leaf shape as arising from discrete nodes or species, a continuous model
defined by indicators like $\ln(vein\ to\ blade\ ratio)$ and distal lobing is more efficient
(**Figure 4**). It is also an open question to what degree developmental constraint and
selection would limit the full manifestation of phenotype across a morphospace. For the
example of grapevine leaf shape, the boundaries of the morphospace are well defined
by developmental constraint and it appears that development and evolution have fully
sampled the space, up to the borders (**Figure 4**).
370
Although reconstructing leaves from a PCA morphospace is routine statistically, this
work focuses on interpretation and how we can use morphometrics to see shape and
natural phenomena through different lenses. Embedded in the morphology of
morphospaces we measure are the constraints by which development and evolution are
modulating natural forms. Measured in sufficient quantities and making reasonable
assumptions about the limits of our models, we can begin to deduce and quantify

constraint, and predict the extent of what is phenotypically possible.

Figure 5: Theoretical leaves. 100 theoretical leaves reconstructed from the intersection of ten, equally spaced ln(vein to blade ratio) and distal lobing isolines, corresponding to orthogonal developmental and evolutionary changes, respectively, across grapevine leaf morphospace. ln(vein to blade ratio) and distal lobing values are shown and leaf areas indicated by color.

378
379

DATA AND CODING AVAILABILITY STATEMENT

The code to reproduce this analysis can be found at the *Github* repository
DanChitwood/grapevine_morphospace:
https://github.com/DanChitwood/grapevine_morphospace. The original leaf scans used
to produce the landmarks are archived on Dryad (Chitwood et al., 2020; Migicovsky et al,
2022b).
387

388　**AUTHOR CONTRIBUTIONS**
389

390　JM developed methods and tools, performed data analysis, and reviewed the
391　manuscript. DHC conceived of the project, performed data analysis, and wrote the
392　manuscript.
393

394　**FINANCIAL SUPPORT**
395

396　This work is supported by the National Science Foundation Plant Genome Research
397　Program award number 1546869. This project was supported by the USDA National
398　Institute of Food and Agriculture, and by Michigan State University AgBioResearch.
399

400　**REFERENCES**
401

402　Abràmoff, M. D., Magalhães, P. J., and Ram, S. J. (2004). Image processing with ImageJ.
403　*Biophotonics International* 11, 36-42.
404

405　Bailey, I. W., and Sinnott, E. W. (1915). A botanical index of Cretaceous and Tertiary
406　climates. *Science* 41 (1066), 831-4.
407

408　Baumgartner, A., Donahoo, M., Chitwood, D. H., and Peppe, D. J. (2020). The influences
409　of environmental change and development on leaf shape in *Vitis*. *American Journal of*
410　*Botany* 107, 1-13.
411

Bryson, A. E., Brown, M. W., Mullins, J., Dong, W., Bahmani, K., Bornowski, N., Chiu, C.,
Engelgau, P., Gettings, B., Gomezcano, F., Gregory, L. M., Haber, A. C., Hoh, D., Jennings,
E. E., Ji, Z., Kaur, P., Kenchanmane Raju, S. K., Long, Y., Lotreck, S., Mathieu, D. T.,
Ranaweera, T., Ritter, E. J., Sadohara, R., Shrote, R. Z., Smith, K. E., Teresi, S. J., Venegas, J.,
Wang, H., Wilson, M. L., Tarrant, A. R., Frank, M. H., Migicovsky, Z., Kumar, J., VanBuren,
R., Londo, J. P., and Chitwood, D. H. (2020). Composite modeling of leaf shape along
shoots discriminates *Vitis* species better than individual leaves. *Applications in Plant*
Sciences 8 (12), e11404.
420

421 Chitwood, D. H., and Sinha, N. R. (2016). Evolutionary and environmental forces
422 sculpting leaf development. *Current Biology* 26 (7), R297-306.
423
424 Chitwood, D. H, Ranjan, A., Martinez, C. C., Headland, L. R., Thiem, T., Kumar, R.,
425 Covington, M. F., Hatcher, T., Naylor, D. T., Zimmerman, S., et al. (2014). A modern
426 ampelography: a genetic basis for leaf shape and venation patterning in grape. *Plant*
427 *Physiology* 164, 259-272.
428
429 Chitwood, D. H., Klein, L. L., O'Hanlon, R., Chacko, S., Greg, M., Kitchen, C., Miller, A. J.,
430 and Londo, J. P. (2016a). Latent developmental and evolutionary shapes embedded
431 within the grapevine leaf. *New Phytologist* 210, 343-355.
432
433 Chitwood, D. H., Rundell, S. M., Li, D. Y., Woodford, Q. L., Tommy, T. Y., Lopez, J. R.,
434 Greenblatt, D., Kang, J., and Londo, J. P. (2016b). Climate and developmental plasticity:
435 interannual variability in grapevine leaf morphology. *Plant Physiology* 170, 1480-1491.
436
437 Chitwood, D. H., VanBuren, R., Migicovsky, Z., Frank, M., and Londo, J. (2020). Data from:
438 Latent developmental and evolutionary shapes embedded within the grapevine
439 leaf. *Dryad*. Dataset, doi.org/10.5061/dryad.zkh189377.
440
441 Chitwood, D. H. (2020). The shapes of wine and table grape leaves: an ampelometric
442 study inspired by the methods of Pierre Galet. *Plants, People, Planet* 00, 1-16.
443
444 Chitwood, D. H., Mullins, J., Migicovsky, Z., Frank, M., VanBuren, R., and Londo, J.P.
445 (2021). Vein-to-blade ratio is an allometric indicator of climate-induced changes in
446 grapevine leaf size and shape. *American Journal of Botany* 108 (4), 571-579.
447
448 Demmings, E. M., Williams, B. R., Lee, C. R., Barba, P., Yang, S., Hwang, C. F., Resich, B. I.,
449 Chitwood, D. H., and Londo, J. P. (2019). Quantitative Trait Locus Analysis of Leaf
450 Morphology Indicates Conserved Shape Loci in Grapevine. *Frontiers in Plant Science* 10,
451 1373.
452
453 Dryden, I. L, and Mardia, K. V. (2016). Statistical shape analysis: with applications in
454 R. Vol. 995. Hoboken, New Jersey, USA: John Wiley & Sons.
455
456 Galet, P. (1979). *A practical ampelography* (L.T. Morton, Trans.). Ithaca, USA: Cornell
457 University Press.
458
459 Galet, P. (1985). *Précis d'ampélographie pratique*, 5 ed., Montpellier, France: Déhan.
460

461   Galet, P. (1988). *Cépages et vignobles de France*, vol. I, *Les vignes américaines*.
462   Montpellier, France: Déhan.
463
464   Galet, P. (1990). *Cépages et vignobles de France*, vol. II. *L'ampélographie française*.
465   Montpellier, France: Déhan.
466
467   Galet, P. (2000). *Dictionnaire encyclopédique des cépages*. Paris, France: Hachette.
468
469   Goethe, J. W. (1817). *Goethe's Werk, Italienische Reise*. Stuttgart, Germany: Dreizehnter
470   Band.
471
472   Hales, S. (1727). *Vegetable Staticks: Or, an Account of Some Statical Experiments on the*
473   *Sap in Vegetables: Being an Essay Towards a Natural History of Vegetation. Also, a*
474   *Specimen of an Attempt to Analyse the Air, by a Great Variety of Chymio-statical*
475   *Experiments; which Were Read at Several Meetings Before the Royal Society* (Vol. 1). p.
476   344, Figs. 43-45. London, England: W. and J. Innys and T. Woodward.
477
478   Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science &*
479   *Engineering* 9, 90-95.
480
481   Kluyver, T., Ragan-Kelley, B., Pérez, F., Granger, B. E., Bussonnier, M., Frederic, J., et al.,
482   and Willing, C. (2016). Jupyter Notebooks-a publishing format for reproducible
483   computational workflows. In F. Loizides & B. Schmidt (Eds.), *Positioning and Power in*
484   *Academic Publishing: Players, Agents and Agendas* pp. 87-90.
485
486   Li, M., An, H., Angelovici, R., Bagaza, C., Batushansky, A., Clark, L., Coneva, V., Donoghue,
487   M. J., Edwards, E., Fajardo, D., et al. (2018). Topological data analysis as a morphometric
488   method: using persistent homology to demarcate a leaf morphospace. *Frontiers in Plant*
489   *Science* 9, 553.
490
491   Martínez, M. C., Loureiro, M. D., and Mantilla, J. L. G. (1995). Importancia y validez de
492   distintos parámetros ampelométricos de hoja adulta para la diferenciación de cepas de
493   Vitis vinifera L., de distintos cultivares. *Invest. Agr. Prod. Prot. Veg.* 9, 377-389.
494
495   Martínez, M. C., Boursiquot, J. M., Grenan, S., and Boidron, R. (1997a). *Étude*
496   *ampelométrique de feuilles adultes de somaclones du cv. Grenache* N (Vitis vinifera L.).
497   *Can. J. Bot.* 75, 333-345.
498

Martínez, M. C., Grenan, S., Boursiquot, J. M. (1997b). Variabilidad de algunos caracteres
ampelográficos y de producción, en somaclones del cultivar Grenache N (Vitis vinifera
L.). *Acta Hortic*. 18, 271-280.

Martínez, M., and Grenan, S. S. (1999). A graphic reconstruction method of an average
vine leaf. *Agronomie, EDP Sciences* 19, 491-507.

McKinney, W. (2010). Data structures for statistical computing in python. In *Proceedings*
of the 9th Python in Science Conference. Vol. 445, pp. 51-56.

Meister, A. L. F. (1769). *Generalia de genesi figurarum planarum et inde pendentibus*
earum affectionibus, Nov. Com. Gött. (in Latin) 1, 144.

Migicovsky, Z., Swift, J. F., Helget, Z., Klein, L. L., Ly, A., Maimaitiyiming, M., Woodhouse,
K., Fennell, A., Kwasniewski, M., Miller, A. J., Cousins, P., Chitwood, D. H. (2022a).
Increases in vein length compensate for leaf area lost to lobing in grapevine. *bioRxiv*,
https://doi.org/10.1101/2022.03.15.484490

Migicovsky, Z., Swift, J. F., Helget, Z., Klein, L. L., Ly, A., Maimaitiyiming, M., Woodhouse,
K., Fennell, A., Kwasniewski, M., Miller, A. J., Cousins, P., Chitwood, D. H. (2022b).
Increases in vein length compensate for leaf area lost to lobing in grapevine. *Dryad*,
https://doi.org/10.5061/dryad.3ffbg79m8

Oliphant, T. E. (2006). A guide to NumPy. USA: Trelgol Publishing.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D.,
Brucher, M., Perrot, M., and Duchesnay, E. (2011). Scikit-learn: Machine learning in
Python. *Journal of Machine Learning Research* 12, 2825-2830.

Ravaz, L. (1902). *Les vignes américaines: Porte-greffes et producteurs directs*. Montpellier
and Paris, France: Goulet. Digitized by Google Books from Cornell University. Public
domain.

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,
Burovski, E., Peterson, P., Weckesser, W., Bright, J. and Van Der Walt, S.J. (2020). SciPy
1.0: fundamental algorithms for scientific computing in Python. *Nature methods*, 17 (3),
261-272

FIGURE LEGENDS

Figure 1: Grapevine leaf morphology. A) Counting from the shoot tip, *Vitis cinerea*
leaves from node positions 1 (left) and 5 (right), each with respective scale bar, are
expanded in detail from the same leaves shown in the panel below. The 21 landmarks
used in this study are indicated, as well as ampelographic nomenclature naming
morphological features. Note that in the younger leaf that vasculature takes up relatively
more area than in the mature leaf. **B)** For seven different grapevine species analyzed in
this study, leaves from the shoot tip to the shoot base are shown with scale bar. Leaf
area increases from the shoot tip to the middle of the shoot due to leaf expansion,
whereas increases in leaf size from the shoot base to the middle of the shoot in mature
leaves are due to heteroblasty.

Figure 2: Modeling $\ln(vein\ to\ blade\ ratio)$ and $\ln(leaf\ area)$ as a function of
normalized node position. A) The natural log of the ratio of vein-to-blade area,
$\ln(vein\ to\ blade\ ratio)$, and **B)** the natural log of leaf area, $\ln(leaf\ area)$, are modeled
as 2nd degree polynomials of normalized node position (where 0 is the shoot tip and 1
is the shoot base). The normalized node position values corresponding to the minimum
$\ln(vein\ to\ blade\ ratio)$ and maximum $\ln(leaf\ area)$ values are indicated by a magenta
vertical line and the inverse of the golden ratio is indicated by a gold vertical line. **C)** In
order to model developmental changes due to leaf expansion separate from
heteroblastic effects, leaves from the shoot tip to the normalized node position value
corresponding to the $\ln(vein\ to\ blade\ ratio)$ minimum were isolated and modeled as a
reciprocal function of normalized node position. Extrapolated values are shown in
dashed line and function asymptotes in purple. **D)** A linear model of $\ln(leaf\ area)$ as a
function of $\ln(vein\ to\ blade\ ratio)$.

Figure 3: Developmental models of leaf shape. Fitting each coordinate value of 21
landmarks as a second-degree polynomial of $\ln(vein\ to\ blade\ ratio)$, continuous models
of expanding leaves for the seven species shown were created. Inclusive of the
maximum and minimum $\ln(vein\ to\ blade\ ratio)$ values for each species, corresponding
to young and mature leaves, respectively, leaves corresponding to ten equally spaced
time points were reconstructed. Estimated leaf areas were estimated from
$\ln(vein\ to\ blade\ ratio)$ values and 1 cm scale bars for each leaf are shown. Leaf areas
are indicated by color.

Figure 4: Morphospace. A morphospace calculated from a Principal Component
Analysis (PCA) of all leaves from the New York germplasm (black) and California
populations (white). **A)** $\ln(vein\ to\ blade\ ratio)$ values and **B)** distal lobing values were
calculated from reconstructed leaves throughout the morphospace using its inverse
transform and colored by magma and virdis color schemes, respectively, as indicated. To
orient and contextualize the space, developmental models for seven grapevine species

were projected into the space, as indicated by colored lines. Isolines for **A)**
$\ln(vein\ to\ blade\ ratio)$ values (solid lines) and **B)** distal lobing values (dashed lines) are
shown and their values provided in the respective plots.
580
581 **Figure 5: Theoretical leaves.** 100 theoretical leaves reconstructed from the intersection
582 of ten, equally spaced $\ln(vein\ to\ blade\ ratio)$ and distal lobing isolines, corresponding
583 to orthogonal developmental and evolutionary changes, respectively, across grapevine
584 leaf morphospace. $\ln(vein\ to\ blade\ ratio)$ and distal lobing values are shown and leaf
585 areas indicated by color.