The grapevine leaves that are (or were) not: constraints on leaf development and choosing to see what seemingly is not

Daniel H. Chitwood[1,2,*], Joey Mullins[1]
[1]Dept. Horticulture, [2]Dept. Computational Mathematics, Science & Engineering, Michigan State University, East Lansing, MI 48823 USA

*Correspondence: Daniel H. Chitwood, <u>chitwoo9@msu.edu</u>

ABSTRACT

The shapes of leaves have been extensively measured and modeled. Using conventional statistical approaches there exist powerful methods to discern between groups and classify shapes. Embedded in calculated morphospaces is information that allows us to visualize theoretical leaves. These unmeasured leaves, that may or may not exist, are never considered in their relation to empirically observable leaves nor how the negative morphospace can inform us about the forces responsible for shaping leaf morphology. Using landmark data for 8465 grapevine leaves, we model leaf shape using an allometric indicator of leaf size, the ratio of vein to blade areas. The borders of the observable morphospace are restricted by the constraint of the subregions of the leaf reducing to nothing. Assuming that self-intersecting polygonal regions in leaves are a developmental constraint, we define a region of morphospace that represents leaf shapes that are possible but not observed. We end with a discussion of how we choose not to observe what seemingly does not exist and the ramifications of this perspective to survivorship bias in the sciences in general.

INTRODUCTION

Leaf shape across plants is diverse and spectacular, but it is not random. Development, evolution, and the environment sculpt leaf shape in specific ways (Chitwood and Sinha, 2016). Leaves allometrically expand, first shown by Stephen Hales through pin pricks on developing fig leaves that were displaced differentially along the length versus the width of the leaf (1727). The developmental programing of leaves changes from node-to-node resulting in changing leaf shapes. Goethe described this process as "metamorphosis" and in terms of the mutable, changing internal state of leaves (1817). Environment modulates leaf size and serrations, as observed by Bailey and Sinnott (1915) who used the distribution of entire leaves across latitudes to estimate the temperatures of paleoclimates. If we measure leaf shape across the seed plants, clear demarcations between phylogenetic groups are observed (Li et al., 2018). We have measured enough leaf shapes to know the borders and demarcations of what exists and the processes that shape leaves in specific ways.

The shapes of grapevine leaves have been measured under intense scrutiny and with
purpose. Originally through morphometric techniques developed by Louis Ravaz (1902),
the field of ampelography ("vine" + "process of measuring") sought to discern, using
leaves and other features of the vine, American *Vitis* species that were new to Europeans
and would eventually be used as rootstocks against Phylloxera. Eventually the
techniques would be famously applied to wine grape varieties by Pierre Galet (1979;
1985; 1988; 1990; 2000; Chitwood, 2020). Morphometric techniques have been used to
genetically study the basis of leaf shape in grapevines (Chitwood et al., 2014; Demmings
et al., 2019), how grapevine leaves develop (Chitwood et al., 2016a), as well as the
effects of environment (Chitwood et al., 2016b; Baumgartner et al., 2020). Modeling has
been used in several ways, including calculating average shapes of grapevine varieties
while preserving features (Martínez et al., 1995; 1997a; 1997b; 1999), modeling
development across grapevine shoots (Bryson et al., 2020), and using leaf allometry,
specifically the ratio of vein to blade areas, as a proxy of leaf size and to measure the
effects of year-to-year variation in leaf shape (Chitwood et al., 2021). For grapevines, as
for many other types of leaves, we have extensively measured and modeled leaf shape,
allowing us to discern genetic, developmental, and environmental effects with great
power.

But what about leaves that are not available for us to measure? Or what about leaves
that never existed? Using what we know about the underlying structure of leaf
morphospaces across genotypic, developmental, end environmental effects, and making
modeling assumptions about what is and is not possible, could we compare what we
have measured and observed against the boundaries of what we know is possible to
reconstruct leaves that *might have been* or *could be*?

Here, we measure the shapes of 8465 grapevine leaves and model them against an
allometric indicator of leaf size, vein-to-blade ratio, across *Vitis* species. The expansion
of blade area at the expense of that for veins is found to be a principal determinant of
the resulting morphospace. Using an inverse transform of the Principal Component
Analysis (PCA) space, theoretical leaves around the border of empirically observable
leaves are visualized. These border leaves are defined by the reduction of blade or vein
area (or both) to the point that the areas of subregions in the leaf are reduced to
nothing. Assuming that self-intersecting polygonal regions represent theoretical leaves
that cannot exist, a region of morphospace containing shapes that are *possible* but not
observed is defined using this developmental constraint. We end with a discussion of
how we choose not to observe what seemingly does not exist and the ramifications of
this perspective to survivorship bias in the sciences.

MATERIALS AND METHODS

As described in Bryson et al., 2020 (and copied verbatim here for convenience), a total of 8465 leaves were collected from 209 vines at the USDA germplasm repository vineyard in Geneva, New York, USA. Samples were taken from the same vines during the second week of June, annually, in 2013 and 2015–2017. The vines sampled represent 11 species (*Ampelopsis glandulosa* (Wall.) Momiy. var. *brevipedunculata* (Maxim.) Momiy., *V. acerifolia* Raf., *V. aestivalis* Michx., *V. amurensis* Rupr., *V. cinerea* (Engelm.) Millardet, *V. coignetiae* Pulliat ex Planch., *V. labrusca* L., *V. palmata* Vahl, *V. riparia* Michx., *V. rupestris* Scheele, and *V. vulpina* L.), four hybrids (*V. ×andersonii* Rehder, *V. ×champinii* Planch., *V. ×doaniana* Munson ex Viala, and *V. ×novae-angliae* Fernald), and 13 *Vitis* vines with an unassigned identity. Starting at the shoot tip (with shoot order noted for each leaf), leaves greater than ~1 cm in length were collected in stacks and stored in a cooler in labeled plastic bags with ventilation holes. Within two days of collection, the leaves were arranged on a large-format Epson Workforce DS-50000 scanner (Tokyo, Japan) in the order they were collected, with a small number near each leaf indicating which node it came from and a ruler for scale within the image file. The image files were named with the vine identification number, followed by a sequential lowercase letter if multiple scans were needed. The original scans are available on Dryad (Chitwood et al., 2020). 21 landmarks were placed on one half of each leaf outlining the midvein, distal vein, proximal vein, and the most proximal branching vein of each of these major veins as well as distal and proximal lobe sinuses using ImageJ (Abràmoff et al., 2004). Two landmarks are placed at the base of each vein to measure the width. The exact placement and correspondence of landmark number to leaf features can be found in Bryson et al., 2020. Landmarks were superimposed through scaling, translation, rotation, and reflection using Generalized Procrustes Analysis with the shapes (Dryden and Mardia, 2016) package in R. It is the Procrustes-adjusted landmark coordinates from Bryson et al., 2020 that are used as the starting point for the analysis presented here.

Data was analyzed using Python and Jupyter notebooks (Kluyver et al., 2016). Code to reproduce the analysis in this manuscript can be found at the *Github* repository DanChitwood/leaves_that_are_or_were_not: https://github.com/DanChitwood/leaves_that_are_or_were_not). The Jupyter notebook (.ipynb) comments on the code and also contains a narrative to guide the reader through the analysis. Calculation of the natural log of the ratio of vein to blade area, $\ln(vein\ to\ blade\ ratio)$, is as described in Chitwood et al. (2021) using the shoelace algorithm, also known as Gauss' are formula, to calculate polygon areas as originally described by Meister (1769), where n is the number of polygon vertices defined by x and y coordinates:

$$\frac{1}{2}|x_1y_2 + x_2y_3 + \ldots + x_{n-1}y_n + x_ny_1 - x_2y_1 - x_3y_2 - \ldots - x_ny_{n-1} - x_1y_n|$$

Principal Component Analysis (PCA) (and calculation of its inverse) was performed using
the scikit learn decomposition PCA module (Pedregosa et al., 2011). Modeling of
$\ln(vein\ to\ blade\ ratio)$, principal components, and landmarks as polynomial functions of
each other and shoot position was performed using the np.polyfit and np.poly1d
functions from NumPy (Oliphant, 2006). An implementation of the Bentley Ottman
sweep-line algorithm (Bentley and Ottman, 1979) was used to detect self-intersecting
polygons in leaf shapes (Kiers and Barton, 2010). Pandas (McKinney, 2010) and
Matplotlib (Hunter, 2007) were used for data analysis and visualization.

RESULTS

Allometry

Previously, we modeled leaf shape
continuously across grapevine shoots
as a polynomial function of each
Procrustes landmark coordinate value
as a function of relative node position.
Relative node position is the node
number counting from the shoot tip
divided the total number of leaves in a
shoot, such that node number is
converted to a 0 to 1 scale, from tip to

base (Bryson et al., 2020). We also
previously described the natural log of
the ratio of vein to blade area,
$\ln(vein\ to\ blade\ ratio)$, which is more
sensitive to leaf area than size itself due
to the exponential increases in blade
relative to vein area during
development (Chitwood et al., 2021).

Figure 1: Modeling *ln(vein to blade ratio)* as a function of normalized shoot position. The natural log of the ratio of vein-to-blade area, *ln(vein to blade ratio)*, is modeled as a 2nd degree polynomial of normalized shoot position (where 0 is the shoot tip and 1 is the shoot base). The normalized shoot position value corresponding to the minimum *ln(vein to blade ratio)* value is 0.6379505834434538 (magenta vertical line). This value has a 3.22% difference from the value of the inverse of the golden ratio, 0.61803398875 (gold vertical line).

Before we explore the limits of the grapevine leaf morphospace (and beyond), we must
first model shape across development to understand how continuous developmental
trajectories change between species during evolution. We plotted
$\ln(vein\ to\ blade\ ratio)$ versus relative shoot position (**Figure 1**), which can be modeled
as a second-degree polynomial. $\ln(vein\ to\ blade\ ratio)$ is highest at the shoot tip and
reaches its minimum in the middle of the shoot. At the shoot tip, veins dominate young,

small leaves before blade area starts to exponentially expand pushing the ratio value
down. $\ln(vein\ to\ blade\ ratio)$ increases again at the shoot base where the first leaves
that emerge are smaller than those that emerge subsequently. A curiosity that is
perhaps coincidental, we note that the corresponding relative node position to the
minimum $\ln(vein\ to\ blade\ ratio)$ value is within 3.22% of the inverse of the golden ratio
(**Figure 1**). Although this may arise as a developmental phenomenon, it could also be
spurious and warrants further investigation.
168
From previous work we know that allometric changes during grapevine leaf expansion
dominate the morphospace (Chitwood et al., 2016a; Chitwood et al., 2016b; Bryson et
al., 2020). We therefore performed a Principal Component Analysis (PCA) on Procrustes-
adjusted landmark coordinates and modeled this space as a function of
$\ln(vein\ to\ blade\ ratio)$ (**Figure 2**). PCA decomposes multivariate data, in essence
rotating and projecting it onto orthogonal axes (principal components) that more
efficiently explain variation in the data than the original measurements (in this case,
Procrustes-adjusted coordinate values). The inverse of this transformation can be used
to reverse calculate original data, which we will later use to visualize theoretical leaves in
the morphospace. PC1 and PC2 explain 90.2% and 9.6% of the variance in the data,
respectively, and can modeled as simply second-degree polynomial functions of
$\ln(vein\ to\ blade\ ratio)$ that for some species can almost be linear (**Figures 2A,B**). These
models can subsequently be combined in a visualization of PC2 vs. PC1, showing the
developmental trajectory of different species in the first two principal components that
explain 99.8% of all variance (**Figure 2C**). We can then visualize all the data, colored by
$\ln(vein\ to\ blade\ ratio)$, and superimpose on this the developmental trajectories of
different *Vitis* and *Ampelopsis* species as modeled by $\ln(vein\ to\ blade\ ratio)$ (**Figure
2D**). Note that $\ln(vein\ to\ blade\ ratio)$ is the major source of variance in the
morphospace, varying strongly and continuously by both PC1 and PC2.
$\ln(vein\ to\ blade\ ratio)$ also makes a strong, straight border in this space along its
lowest values (white in **Figure 2D**) that correspond to large, mature leaves with a small
ratio of vein-to-blade areas. The modeled trajectories of different species each traverse
through higher (young, small leaves) and lower (mature, large leaves)
$\ln(vein\ to\ blade\ ratio)$ values, but the different species trajectories separate
orthogonally to this direction. As such, development as measured by
$\ln(vein\ to\ blade\ ratio)$ and genetic variation as measured through different species
define the morphospace.
196
197 We can visualize the continuous allometric changes in leaf shape across the modeled
198 developmental trajectories (**Figure 3**). We can both visualize modeled leaf shapes
199 themselves and keep track of the proportion of the leaf occupied by each region and

Figure 2: Developmental trajectories of leaf development through Principal Component Analysis (PCA) space by species. A-B) 2nd degree polynomials modeling PC1 (A) and PC2 (B) values by *ln(vein to blade ratio)* for three species (*Vitis riparia*, gold; *V. amurensis*, lavender; *Ampelopsis glandulosa* var. *brevipedunculata*, green). **C)** Data points and models plotted as PC2 vs. PC1. Regions corresponding to mature, old leaves and young, small leaves are indicated. **D)** Models for 11 species (indicated by color) and data points colored by *ln(vein to blade ratio)* (a white-to-black gradient, scale provided). Percent variance explained by each PC are indicated in axis labels.

clearly see how ln($vein\ to\ blade\ ratio$) corresponds to the vein and blade regions it
measures. Young and small leaves have relatively high ratios of vein area (yellow in
Figure 3). As leaves expand in size and mature, blade regions expand at an exponential
rate before plateauing, and reduce the relative proportion of vein area in the leaf. Note
that in *A. glandulosa* var. *brevipedunculata*, which is deeply lobed, the ratios of vein and
blade area are more stable across ln($vein\ to\ blade\ ratio$) values compared to more
entire species. In one of the most entire species, *V. amurensis*, the decreases in vein area
as the leaf expands are more dramatic, perhaps corresponding to the fact that there is
indeed more blade that expands.

Figure 3: Allometric changes in leaf shape during development. Changes in the proportional area different regions of the leaf occupy (middle lobe, lavender blue; vasculature, yellow; distal lobe, light green; distal sinus, dark green; proximal lobe, orange; proximal sinus, magenta) are modeled as developmental trajectories across 100 theoretical time points for three different species: *V. riparia*, *V. amurensis*, and *A. glandulosa* var. *brevipedunculata*. Theoretical grapevine leaves are reconstructed for timepoints 0, 12, 24, 36, 48, 60, 72, 84, and 96 for each species. Note that the trajectories run from mature, large leaves on the left to young, small leaves on the right.

Figure 4: Theoretical grapevine leaves at the limits of what is empirically observed. Principal Component Analysis (PCA) space plotting individual leaves colored by *ln(vein to blade ratio)* (a white-to-black gradient, scale provided) and modeled developmental trajectories of different species (indicated by color). Three lines are drawn that encompass the shape of all empirically measured leaves. Theoretical grapevine leaves are reconstructed for five equidistant points along each line (large gray dots), pointed to by lines outside of the plot. Different regions of the leaf are indicated by color: middle lobe, lavender blue; vasculature, yellow; distal lobe, light green; distal sinus, dark green; proximal lobe, orange; proximal sinus, magenta.

Morphospace

The morphospace is clearly defined (**Figure 2D**): for the leaves we have measured there is a strong, continuous gradient of $\ln(\textit{vein to blade ratio})$ values and for the highest values that correspond to mature, large leaves in which blade has expanded at the expense of veins, there is a clearly delineated border that the developmental trajectories of species terminate at. For the leaves that we have measured, what do leaves at the border of the morphospace look like, the most "extreme" leaves? Drawing three, arbitrary lines around the data and taking five equidistant points in morphospace along each, theoretical leaves were reconstructed using PCA loadings and the inverse transformation (**Figure 4**). The most obvious varying feature of these border leaves is that which most strongly affects the structure of the morphospace itself: $\ln(\textit{vein to blade ratio})$. Along the defined line of border leaves with low $\ln(\textit{vein to blade ratio})$ values (white in **Figure 4**), the vein area is almost non-existent relative to blade in reconstructed leaves. The line opposite it in the border triangle, the leaves with high $\ln(\textit{vein to blade ratio})$ values (black in **Figure 4**), the opposite is true, leaves have thick veins and reduced blade area. These two lines intersect at a point with high PC1 and low PC2 values that combines the reductions in blade and vein into an unsustainable crisis: an imploding leaf with no blade or vein area. Opposite this vertex, the line connecting the spaces of low and high $\ln(\textit{vein to blade ratio})$ values shows what one would expect: a gradual transition between a leaf with almost no vein area to one where it predominates.

If we take Procrustes-adjusted landmarks to be a model of leaf shape, then we can consider instances that violate assumptions of reality on which that model is based. In this specific case, if any of the polygonal areas that comprise a reconstructed leaf self-intersect and cross themselves, this is a leaf that cannot exist because of developmental constraint: it simply is not possible. This does not imply that the condition of non-intersecting polygonal areas accurately represents the actual boundaries of constraint, only that for our model of leaf shape, that it is clearly a violation of what is possible. Expanding beyond the PCA morphospace and reconstructing theoretical leaves, we can use the Bentley Ottman sweep-line algorithm to test if leaves contain any self-intersecting polygonal regions (**Figure 5**). We can also measure $\ln(\textit{vein to blade ratio})$ values of these theoretical leaves to help us understand why such constraints arise as well as visualize the leaves themselves (**Figure 6**).

On at least two sides and a point of intersection of the empirically observed distribution of the grapevine leaves, the border of leaves that are self-intersecting comes close to that of those that we measured. These borders are along the sharply defined line of leaves with low $\ln(\textit{vein to blade ratio})$ values in which blade is greatly expanded at the expense of vein regions and along the opposite border with high $\ln(\textit{vein to blade ratio})$

Figure 5: Self-intersecting leaves. 100 theoretical leaves, equidistantly spaced along PC1 and PC2, were reconstructed. The range was selected to find the limits of theoretical leaves with any self-intersection in any sub-region of the leaf. For each reconstructed leaf, *ln(vein to blade ratio)* was calculated (a white-to-black gradient, scale provided). The boundary that encompasses theoretical leaves with no self-intersection of any region is indicated by a solid, black line. Empirically measured leaves are indicated by points (dodger blue).

values in which vein areas predominate at the expense of blade (**Figures 5-6**). Also, the "point of implosion" where these two axes intersect, where the reduction of both vein and blade area simultaneously reduces the areas of all regions to the point of non-existence, corresponds closely to that observed in measured leaves. If we take this apex and we extend the two lines of low and high $ln(vein\ to\ blade\ ratio)$ values outward, we find that there is a substantial region of morphospace of leaves *without* self-intersecting regions that extends beyond the region of empirically measured leaves in this study. These are leaves, given the caveats of our model, that potentially

could have, *do*, or *can* exist but have not been measured.

DISCUSSION

In exploring the leaves beyond the boundaries that have been empirically observed and
are described in this manuscript, it is important that we distinguish between different
types. First (1), the leaves that exist or existed: there are (1a) the leaves that we have
measured here and (1b) the leaves that we have not measured but *do* or *have* existed.
Although relatively comprehensive across *Vitis* spp. excluding varieties of *V. vinifera*
(discussed in the following paragraphs below), the leaves empirically observed in this
manuscript do not come close to measuring all *Vitis* leaves. There are (1bi) extant
species and genotypes known or unknown that are not represented, (1bii) all the
environments and circumstances those extant genotypes have existed in, (1biii) extinct
species and genotypes, and (1biv) the environments and circumstances that extinct
grapevine leaves existed in. Then there are (2) the leaves that have not existed or maybe
never will. There are (2a) leaves that *could* have existed or *can*, that represent (2ai)
evolutionary, genetic, and environmental possibilities that remain unsampled or never
were sampled or (2aii) leaves that *cannot* exist and never have and never will. This last
category is what is referred to as *developmental constraint* here and, given the
limitations of the model of grapevine leaf shape, is being measured by observing self-
intersecting polygonal areas.

Figure 6: Morphospace. 100 reconstructed grapevine leaves across PC1 and PC2. Rounded coordinates for each theoretical leaf are indicated along the axes. The boundary that encompasses theoretical leaves with no self-intersection of any region is indicated by a solid black line. The boundary that encompasses nearly all empirically measured leaves is indicated by a dashed black line. Different regions of the leaf are indicated by color: middle lobe, lavender blue; vasculature, yellow; distal lobe, light green; distal sinus, dark green; proximal lobe, orange; proximal sinus, magenta. Alpha transparency is used so that overlapping regions in leaves can be seen.

In describing the features of leaves that define the morphospace, $\ln(vein\ to\ blade\ ratio)$
is the strongest, but a detail needs to be clarified. High PC1 and PC2 values correspond
to a region with high $\ln(vein\ to\ blade\ ratio)$ values (black in **Figure 5**). The border of
leaves with self-intersecting polygonal regions opposite of this space has the lowest
observed $\ln(vein\ to\ blade\ ratio)$ values (the band of white in **Figure 5**), but then it
appears that $\ln(vein\ to\ blade\ ratio)$ increases again in the area defined by low PC1 and
PC2 values. The shoelace algorithm can be used to calculate the areas of self-
intersecting polygons, and these are non-negative values. The area defined by low PC1
and PC2 values has vein areas so reduced that they have "imploded" and self-
intersected on themselves (observe reconstructed leaves from this area in **Figure 6**),
increasing $\ln(vein\ to\ blade\ ratio)$ values in a misleading way. Unlike **Figure 5** shows, the
morphospace is divided into regions of strongly high and low $\ln(vein\ to\ blade\ ratio)$
values, with empirically measured leaves in the middle. With respect to developmental
constraint, it likely does limit observable grapevine leaf shapes. The most mature, largest
leaves encounter a constraint in which vein widths cannot be further reduced and blade
area is maximized (low PC1 and PC2 values); the youngest, smallest leaves cannot
eliminate any more blade region at the expense of vasculature (high PC1 and PC2
values); and then there is the "point of implosion" (high PC1, low PC2 values), in which
reductions in both blade and vein area threaten the existence of any leaf area at all
(**Figure 6**).
312
The most interesting region of the morphospace identified in this study is that in which
no self-intersection is detected but there are no empirically measured leaves (**Figure 6**).
A characteristic of these leaves is a "croissant" shape: it appears as if the leaf is bending
over upon itself. A first thought is that no empirically measured leaves occupy this space
because of the fact we measured leaf halves, that there is no way these shapes can make
a whole leaf and it is an artifact of the analysis. However, these half leaves can be put
together into whole leaves, it will just be that the petiolar veins and sinus (that are
comprised of the "proximal lobe" and "proximal sinus" regions in this study) will overlap.
An overlapping petiolar sinus is a historical ampelographic feature, especially in *V.*
vinifera varieties (Ravaz, 1902; Galet, 1979; 1985; 1988; 1990, 2000). We do have scans
and have performed landmark analysis in *V. vinifera* varieties previously (Chitwood et al.,
2014; Chitwood, 2020), but we used a fewer number of landmarks that do not measure
the widths of veins (and therefore do not measure vein area). Additionally, the *V. vinifera*
leaves that have been measured do not include young leaves, which are mostly
contributing to leaves with high $\ln(vein\ to\ blade\ ratio)$ values. It is outside of the scope
of this manuscript to measure these leaves, but it is an important next step.
Landmarking and projecting these leaves onto the morphospace will reveal how much
of the space without developmental constraints but not currently occupied by the leaves
measured in this study is occupied and if it is truly empty. It might be possible that

developmental constraint, defined by self-intersecting regions of the leaf, is sufficient to
explain the boundaries of extant grapevine leaves and likewise that grapevine leaves
fully occupy the space available to them.
335
Although reconstructing leaves from a PCA morphospace is routine statistically, this
work focuses on interpretation and how we can use morphometrics to see shape and
natural phenomena through different lenses. Embedded in the morphology of
morphospaces we measure are the constraints by which development and evolution are
modulating natural forms. Measured in sufficient quantities and making reasonable
assumptions about the limits of our models, we can begin to deduce and quantify
constraint, drawing a line between that which is impossible and that which is possible
but may not exist (or once existed). Both in past and present, morphometrics has been
used for nefarious purposes to distinguish groups of people and justify exclusion,
discrimination, and erasure (Clayton, 2020), and it is from this past that the current
scientific focus on classification and parametric statistics is based. But we can *choose* to
explore our data more fully and focus on more than only what exists (or that we choose
to acknowledge). Failing to do so is a type of survivorship bias: by only considering that
which exists, is present, survived, or that we selectively acknowledge, we are led to false
conclusions and commit logical errors (Wald, 1943; Wallis, 1980). This philosophy can be
extended to how we conduct ourselves and relate to others in science as well, from how
we address and prevent demographic biases in the sciences (Madzima and MacIntosh,
2021; Field and Rajewski, 2021) to the response of the academy to racial injustices and
the pandemic (Montgomery, 2020). By choosing to see what seemingly is not—who we
exclude around us but nonetheless exist, who have made contributions we neglect, and
who are part of the community and its history but have not been properly
acknowledged—we can see a bigger picture and better understand the dynamics that
underlie the spaces that we study and inhabit.
359
360 **DATA AVAILABILITY STATEMENT**
361
The code to reproduce this analysis can be found at the *Github* repository
DanChitwood/leaves_that_are_or_were_not:
https://github.com/DanChitwood/leaves_that_are_or_were_not. The Procrustes landmark
data was originally used in Bryson et al., 2020 and is posted at the *Github* repository
DanChitwood/grapevine_shoots: https://github.com/DanChitwood/grapevine_shoots.
The original leaf scans used to produce the landmarks are archived on Dryad (Chitwood
et al., 2020).
369
370 **AUTHOR CONTRIBUTIONS**
371

372 JM developed methods and tools, performed data analysis, and reviewed the
373 manuscript. DHC conceived of the project, performed data analysis, and wrote the
374 manuscript.
375

376 **FUNDING**
377
378 This work is supported by the National Science Foundation Plant Genome Research
379 Program award number 1546869. This project was supported by the USDA National
380 Institute of Food and Agriculture, and by Michigan State University AgBioResearch.
381

382 **REFERENCES**
383
384 Abràmoff, M. D., Magalhães, P. J., and Ram, S. J. (2004). Image processing with ImageJ.
385 *Biophotonics International* 11, 36-42.
386
387 Bailey, I. W., and Sinnott, E. W. (1915). A botanical index of Cretaceous and Tertiary
388 climates. *Science* 41 (1066), 831-4.
389
390 Baumgartner, A., Donahoo, M., Chitwood, D. H., and Peppe, D. J. (2020). The influences
391 of environmental change and development on leaf shape in *Vitis*. *American Journal of*
392 *Botany* 107, 1-13.
393
394 Bentley, J. L., and Ottmann, T. A. (1979). Algorithms for reporting and counting
395 geometric intersections. *IEEE Transactions on Computers* C-28 (9), 643–647.
396
Bryson, A. E., Brown, M. W., Mullins, J., Dong, W., Bahmani, K., Bornowski, N., Chiu, C.,
Engelgau, P., Gettings, B., Gomezcano, F., Gregory, L. M., Haber, A. C., Hoh, D., Jennings,
E. E., Ji, Z., Kaur, P., Kenchanmane Raju, S. K., Long, Y., Lotreck, S., Mathieu, D. T.,
Ranaweera, T., Ritter, E. J., Sadohara, R., Shrote, R. Z., Smith, K. E., Teresi, S. J., Venegas, J.,
Wang, H., Wilson, M. L., Tarrant, A. R., Frank, M. H., Migicovsky, Z., Kumar, J., VanBuren,
R., Londo, J. P., and Chitwood, D. H. (2020). Composite modeling of leaf shape along
shoots discriminates *Vitis* species better than individual leaves. *Applications in Plant*
Sciences 8 (12), e11404.
405
406 Chitwood, D. H., and Sinha, N. R. (2016). Evolutionary and environmental forces
407 sculpting leaf development. *Current Biology* 26 (7), R297-306.
408
Chitwood, D. H, Ranjan, A., Martinez, C. C., Headland, L. R., Thiem, T., Kumar, R.,
Covington, M. F., Hatcher, T., Naylor, D. T., Zimmerman, S., et al. (2014). A modern

ampelography: a genetic basis for leaf shape and venation patterning in grape. *Plant*
Physiology 164, 259-272.

Chitwood, D. H., Klein, L. L., O'Hanlon, R., Chacko, S., Greg, M., Kitchen, C., Miller, A. J.,
and Londo, J. P. (2016a). Latent developmental and evolutionary shapes embedded
within the grapevine leaf. *New Phytologist* 210, 343-355.

Chitwood, D. H., Rundell, S. M., Li, D. Y., Woodford, Q. L., Tommy, T. Y., Lopez, J. R.,
Greenblatt, D., Kang, J., and Londo, J. P. (2016b). Climate and developmental plasticity:
interannual variability in grapevine leaf morphology. *Plant Physiology* 170, 1480-1491.

Chitwood, D. H., VanBuren, R., Migicovsky, Z., Frank, M., and Londo, J. (2020). Data from:
Latent developmental and evolutionary shapes embedded within the grapevine
leaf. *Dryad*. Dataset, doi.org/10.5061/dryad.zkh189377.

Chitwood, D. H. (2020). The shapes of wine and table grape leaves: an ampelometric
study inspired by the methods of Pierre Galet. *Plants, People, Planet* 00, 1-16.

Chitwood, D. H., Mullins, J., Migicovsky, Z., Frank, M., VanBuren, R., and Londo, J.P.
(2021). Vein-to-blade ratio is an allometric indicator of climate-induced changes in
grapevine leaf size and shape. *American Journal of Botany* 108 (4), 571-579.

Clayton, A. (2020). How eugenics shaped statistics. *Nautilus* 28 October. Retrieved 24
January 2022 https://nautil.us/how-eugenics-shaped-statistics-9365/

Demmings, E. M., Williams, B. R., Lee, C. R., Barba, P., Yang, S., Hwang, C. F., Resich, B. I.,
Chitwood, D. H., and Londo, J. P. (2019). Quantitative Trait Locus Analysis of Leaf
Morphology Indicates Conserved Shape Loci in Grapevine. *Frontiers in Plant Science* 10,
1373.

Dryden, I. L, and Mardia, K. V. (2016). Statistical shape analysis: with applications in
R. Vol. 995. Hoboken, New Jersey, USA: John Wiley & Sons.

Field, S., and Rajewski, A. (2021). Challenges facing LGBTQ+ early-career scientists and
how to engage in changing the status quo. *Plant Cell* 33, 1859-1862.

Galet, P. (1979). *A practical ampelography* (L.T. Morton, Trans.). Ithaca, USA: Cornell
University Press.

Galet, P. (1985). *Précis d'ampélographie pratique*, 5 ed., Montpellier, France: Déhan.

Galet, P. (1988). *Cépages et vignobles de France*, vol. I, *Les vignes américaines*.
Montpellier, France: Déhan.

Galet, P. (1990). *Cépages et vignobles de France*, vol. II. *L'ampélographie française*.
Montpellier, France: Déhan.

Galet, P. (2000). *Dictionnaire encyclopédique des cépages*. Paris, France: Hachette.

Goethe, J. W. (1817). *Goethe's Werk, Italienische Reise*. Stuttgart, Germany: Dreizehnter
Band.

Hales, S. (1727). *Vegetable Staticks: Or, an Account of Some Statical Experiments on the*
Sap in Vegetables: Being an Essay Towards a Natural History of Vegetation. Also, a
Specimen of an Attempt to Analyse the Air, by a Great Variety of Chymio-statical
Experiments; which Were Read at Several Meetings Before the Royal Society (Vol. 1). p.
344, Figs. 43-45. London, England: W. and J. Innys and T. Woodward.

Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science &*
Engineering 9, 90-95.

Kiers, B., and Barton, C. (copyrights 2010 and 2015, respectively). Poly Point
Intersections. *Github*. https://github.com/ideasman42/isect_segments-bentley_ottmann.
Retrieved 27 January 2022.

Kluyver, T., Ragan-Kelley, B., Pérez, F., Granger, B. E., Bussonnier, M., Frederic, J., et al.,
and Willing, C. (2016). Jupyter Notebooks-a publishing format for reproducible
computational workflows. In F. Loizides & B. Schmidt (Eds.), *Positioning and Power in*
Academic Publishing: Players, Agents and Agendas pp. 87-90.

Li, M., An, H., Angelovici, R., Bagaza, C., Batushansky, A., Clark, L., Coneva, V., Donoghue,
M. J., Edwards, E., Fajardo, D., et al. (2018). Topological data analysis as a morphometric
method: using persistent homology to demarcate a leaf morphospace. *Frontiers in Plant*
Science 9, 553.

Madzima, T. F., and MacIntosh, G. C. (2021). Equity, diversity, and inclusion efforts in
professional societies: intention versus reaction. *Plant Cell* 33, 3189-319.

Martínez, M. C., Loureiro, M. D., and Mantilla, J. L. G. (1995). Importancia y validez de
distintos parámetros ampelométricos de hoja adulta para la diferenciación de cepas de
Vitis vinifera L., de distintos cultivares. *Invest. Agr. Prod. Prot. Veg*. 9, 377-389.
492

493 Martínez, M. C., Boursiquot, J. M., Grenan, S., and Boidron, R. (1997a). *Étude*
494 *ampelométrique de feuilles adultes de somaclones du cv. Grenache* N (Vitis vinifera L.).
495 *Can. J. Bot*. 75, 333-345.
496

497 Martínez, M. C., Grenan, S., Boursiquot, J. M. (1997b). Variabilidad de algunos caracteres
498 ampelográficos y de producción, en somaclones del cultivar Grenache N (Vitis vinifera
499 L.). *Acta Hortic*. 18, 271-280.
500

501 Martínez, M., and Grenan, S. S. (1999). A graphic reconstruction method of an average
502 vine leaf. *Agronomie, EDP Sciences* 19, 491-507.
503

504 McKinney, W. (2010). Data structures for statistical computing in python. In *Proceedings*
505 *of the 9th Python in Science Conference*. Vol. 445, pp. 51-56.
506

507 Meister, A. L. F. (1769), *Generalia de genesi figurarum planarum et inde pendentibus*
508 *earum affectionibus*, Nov. Com. Gött. (in Latin) 1, 144.
509

510 Montgomery, B. L. (2020). Planting equity: Using what we know to cultivate growth as a
511 plant biology community. *Plant Cell* 32, 3372-3375.
512

513 Oliphant, T. E. (2006). A guide to NumPy. USA: Trelgol Publishing.
514

515 Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
516 Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D.,
517 Brucher, M., Perrot, M., and Duchesnay, E. (2011). Scikit-learn: Machine learning in
518 Python. *Journal of Machine Learning Research* 12, 2825-2830.
519

520 Ravaz, L. (1902). *Les vignes américaines: Porte-greffes et producteurs directs*. Montpellier
521 and Paris, France: Goulet. Digitized by Google Books from Cornell University. Public
522 domain.
523

524 Wald, A. (1943). A Method of Estimating Plane Vulnerability Based on Damage of
525 Survivors. Statistical Research Group, Columbia University. CRC 432 — reprint from July
526 1980.
527

528 Wallis, W. A. (1980). The Statistical Research Group, 1942-1945: Rejoinder. *Journal of the*
529 *American Statistical Association*. 75 (370), 334–335.

530
531 **FIGURE LEGENDS**
532

Figure 1: Modeling $\ln(vein\ to\ blade\ ratio)$ as a function of normalized shoot
position. The natural log of the ratio of vein-to-blade area, $\ln(vein\ to\ blade\ ratio)$, is
modeled as a 2nd degree polynomial of normalized shoot position (where 0 is the shoot
tip and 1 is the shoot base). The normalized shoot position value corresponding to the
minimum $\ln(vein\ to\ blade\ ratio)$ value is 0.6379505834434538 (magenta vertical line).
This value has a 3.22% difference from the value of the inverse of the golden ratio,
0.61803398875 (gold vertical line).

Figure 2: Developmental trajectories of leaf development through Principal
Component Analysis (PCA) space by species. A-B) 2nd degree polynomials modeling
PC1 (A) and PC2 (B) values by $\ln(vein\ to\ blade\ ratio)$ for three species (*Vitis riparia*,
gold; *V. amurensis*, lavender; *Ampelopsis glandulosa* var. *brevipedunculata*,
green). **C)** Data points and models plotted as PC2 vs. PC1. Regions corresponding to
mature, old leaves and young, small leaves are indicated. **D)** Models for 11 species
(indicated by color) and data points colored by $\ln(vein\ to\ blade\ ratio)$ (a white-to-black
gradient, scale provided). Percent variance explained by each PC are indicated in axis
labels.

Figure 3: Allometric changes in leaf shape during development. Changes in the
proportional area different regions of the leaf occupy (middle lobe, lavender blue;
vasculature, yellow; distal lobe, light green; distal sinus, dark green; proximal lobe,
orange; proximal sinus, magenta) are modeled as developmental trajectories across 100
theoretical time points for three different species: *V. riparia*, *V. amurensis*, and *A.*
glandulosa var. *brevipedunculata*. Theoretical grapevine leaves are reconstructed for
timepoints 0, 12, 24, 36, 48, 60, 72, 84, and 96 for each species. Note that the trajectories
run from mature, large leaves on the left to young, small leaves on the right.

Figure 4: Theoretical grapevine leaves at the limits of what is empirically
observed. Principal Component Analysis (PCA) space plotting individual leaves colored
by $\ln(vein\ to\ blade\ ratio)$ (a white-to-black gradient, scale provided) and modeled
developmental trajectories of different species (indicated by color). Three lines are
drawn that encompass the shape of all empirically measured leaves. Theoretical
grapevine leaves are reconstructed for five equidistant points along each line (large gray
dots), pointed to by lines outside of the plot. Different regions of the leaf are indicated
by color: middle lobe, lavender blue; vasculature, yellow; distal lobe, light green; distal
sinus, dark green; proximal lobe, orange; proximal sinus, magenta.

Figure 5: Self-intersecting leaves. 100 theoretical leaves, equidistantly spaced along PC1 and PC2, were reconstructed. The range was selected to find the limits of theoretical leaves with any self-intersection in any sub-region of the leaf. For each reconstructed leaf, $\ln(vein\ to\ blade\ ratio)$ was calculated (a white-to-black gradient, scale provided). The boundary that encompasses theoretical leaves with no self-intersection of any region is indicated by a solid, black line. Empirically measured leaves are indicated by points (dodger blue).

Figure 6: Morphospace. 100 reconstructed grapevine leaves across PC1 and PC2. Rounded coordinates for each theoretical leaf are indicated along the axes. The boundary that encompasses theoretical leaves with no self-intersection of any region is indicated by a solid black line. The boundary that encompasses nearly all empirically measured leaves is indicated by a dashed black line. Different regions of the leaf are indicated by color: middle lobe, lavender blue; vasculature, yellow; distal lobe, light green; distal sinus, dark green; proximal lobe, orange; proximal sinus, magenta. Alpha transparency is used so that overlapping regions in leaves can be seen.