Condensation of a nuclear mRNA export factor regulates mRNA transport during stress

Stephanie Heinrich[1,#], Maria Hondele[1,2], Désirée Marchand[1], Carina Patrizia Derrer[1], Mostafa Zedan[1], Alexandra Oswald[1], Federico Uliana[1], Roberta Mancini[1], David Grunwald[3], Karsten Weis[1,#]

[1] Institute of Biochemistry, ETH Zurich, Otto-Stern-Weg 3, 8093 Zurich, Switzerland
[2] Biozentrum, University of Basel, Klingelbergstrasse 50/70, 4056 Basel, Switzerland
[3] University of Massachusetts Medical School, RNA Therapeutics Institute, 368 Plantation Street, Worcester MA 01605, USA
[#] Corresponding author, lead contact: karsten.weis@bc.biol.ethz.ch

Highlights

- The nuclear Poly(A)-binding protein Nab2 forms condensates upon acute cellular depletion of the DEAD-box ATPase Dbp5
- Nab2 dimerization but not its intrinsically disordered regions (IDRs) are essential for condensation *in vitro* and *in vivo*
- Glucose stress leads to global mRNA retention in the nucleus and depends on Nab2 condensation
- Nab2 condensation confers selectivity of RNA retention during glucose stress
- Proper Nab2 condensation is essential for survival after prolonged stress

Summary

Unidirectional transport of mRNA from the nucleus to the cytoplasm via nuclear pore complexes is an essential step in the gene expression of all eukaryotes. Although factors involved in mRNA transport have been characterized, a comprehensive mechanistic understanding of this critical process and its regulation is lacking. Here, we use real-time single RNA imaging to demonstrate that acute depletion of the budding yeast DEAD-box ATPase Dbp5 causes rapid nuclear accumulation of mRNAs *in vivo* and dramatic changes in nuclear dynamics of RNA export factors. In particular, the essential export factor Nab2 ceases to shuttle between the nucleus and cytoplasm and forms an RNA-dependent condensate throughout the nucleus. Phase-separation can be recapitulated *in vitro*, with Nab2 forming RNA-dependent liquid droplets, which depend on the presence of Dbp5. Intriguingly, in glucose stress, condensation of Nab2 blocks bulk mRNA export while selectively allowing the passage of stress-induced mRNAs from the nucleus to the cytoplasm to elicit a timely cellular stress response. This is accompanied by a lowered abundance of the DEAD-box ATPase Dbp5 at the cytoplasmic sites of nuclear pore complexes, which leads to the formation of the Nab2 condensates. Our results suggest that cells use selective mRNA retention in nuclear Nab2 condensates to re-wire mRNA export and to regulate gene expression during stress.

Keywords

RNA, mRNA export, phase separation, Nab2, DEAD-box ATPase Dbp5, glucose stress

Introduction

Gene expression in eukaryotic cells is intricately regulated and aided by subcellular compartmentalization. The nuclear envelope serves as a physical barrier and separates nuclear transcription from cytoplasmic translation processes. Embedded in the nuclear membrane are nuclear pore complexes (NPC) that regulate the passage of macromolecules to and from the nucleus, including the export of messenger RNAs (mRNA) that relay the genetic information to the protein translation machinery in the cytoplasm (reviewed in (Ashkenazy-Titelman et al., 2020; De Magistris, 2021)). mRNAs are extensively processed in the nucleus and undergo a coordinated series of modifications before export, which includes the addition of a 5' m7G cap structure, removal of introns through splicing, and 3' end processing for poly(A) tail synthesis (reviewed in (Herzel et al., 2017; Rambout and Maquat, 2020; Shi and Manley, 2015)). To facilitate these processes, a range of RNA-binding proteins (RBPs) form a dynamic mRNA coat on the nascent mRNA, either co-transcriptionally or post-transcriptionally in the nucleoplasm. The orderly assembly of this coat is essential for forming an export-competent mRNA-protein (mRNP) particle (reviewed in (Ashkenazy-Titelman et al., 2020; Singh et al., 2015)). Early binding proteins include the cap-binding complex (CBC; comprising Cbc1 (CBP80 in vertebrates) and Cbc2 (CBP20)), which binds co-transcriptionally to the 5'-cap of nascent mRNAs (Izaurralde et al., 1994). Another early binding complex, the transcription and export (TREX) complex (consisting of the multi-subunit THO subcomplex, the DExD/H-box ATPase Sub2 (UAP56), and the RNA binding protein Yra1 (ALYREF)) connects 3' end processing with mRNA export (Katahira, 2012; Pühringer et al., 2020; Sträßer et al., 2002; Sträßer and Hurt, 2001). Similar to TREX, the abundant class of serine-arginine-rich (SR) proteins (e.g., yeast Npl3) has been implicated in a multitude of mRNA processing steps, such as co-transcriptional splicing, 3' end processing and nuclear export (reviewed in (Wegener and Müller-McNicoll, 2019)). The export factor Nab2 (ZC3H14) is recruited to the poly(A) tail of the pre-mRNA and is involved in nuclear mRNA surveillance protecting mRNAs against degradation by the nuclear exosome (Anderson et al., 1993; Hector et al., 2002; Schmid et al., 2012, 2015). Nab2 has been shown to dimerize on RNA and to play a role in proper 3' end processing of nascent transcripts to terminate polyadenylation and facilitate compaction of the mRNP (Aibara et al., 2017; Alpert et al., 2020; Fasken et al., 2019). In addition, it directly interacts with the NPC component Mlp1 (Green et al., 2003). The translocation of mature transcripts through NPCs is mediated by the export factor Mex67:Mtr2 (NXF1:NXT1), which functions akin to a mobile nuclear pore protein facilitating transport by binding to both mRNPs as well as phenylalanine–glycine (FG)-containing NPC components (nucleoporins)(Ashkenazy-Titelman et al., 2020; Derrer et al., 2019). On the cytoplasmic face of the NPC, the essential DEAD-box ATPase Dbp5 (DDX19B), aided by its interaction partners Nup159 (NUP214), Gle1 and inositol hexakisphosphate (IP$_6$), has been suggested to remodel emerging mRNPs by removing export factors such as Nab2 from the mRNA, thereby facilitating their release into the cytoplasm (Alcázar-Román et al., 2006; Tran et al., 2007; Weirich et al., 2006). However, these models are mainly based on *in vitro* studies (Montpetit et al., 2011; Noble et al., 2011; Tran et al., 2007), and the *in vivo* function of Dbp5 in directional transport remains unclear.

In addition to membrane-based compartmentalization, cells also organize their content in membraneless organelles through liquid-liquid phase separation (LLPS)(reviewed in (Banani et al., 2017; Shin and Brangwynne, 2017). Formation of these biomolecular condensates depends on weak, multivalent interactions and proteins undergoing LLPS often contain low-complexity domains with intrinsically disordered regions (IDRs). IDRs are frequently found in RNA-binding proteins and RNA can enhance their LLPS (reviewed in (Roden and Gladfelter, 2021)). LLPS is an evolutionarily ancient and conserved self-organizing principle (Hondele et al., 2019; Riback et al., 2017). Within a cell, many structures can be

defined as membraneless organelles and they have been implicated in various aspects of gene expression regulation (reviewed in (Banani et al., 2017)). For example, stress conditions such as nutrient depletion or acute temperature shifts have been shown to induce the formation of cytoplasmic condensates such as P-bodies (PB) or stress granules (SG) (Kroschwald et al., 2015; Sheth and Parker, 2003). If LLPS contributes to or controls directional mRNA export is unknown.

Here, we report that nuclear condensation of the mRNA export factor Nab2 selectively regulates mRNA transport in glucose stress, which is modulated by the activity of the DEAD-box ATPase Dbp5. Nab2 condensation leads to the retention of bulk mRNAs in the nucleus but allows selective export of stress-induced transcripts, which is impaired in Nab2 mutants that affect the state of the condensate. Our results establish condensation as a regulator of mRNA transport during cellular stress and provide mechanistic insights into mRNA export regulation.

Results

Acute Dbp5 depletion causes nuclear mRNA accumulation and Nab2 mobility reduction

To characterize the function of Dbp5 in directional mRNA export *in vivo*, we employed the auxin-inducible acute protein depletion system in the budding yeast *Saccharomyces cerevisiae* (Nishimura 2009, Fig. 1A, 'degron'). Addition of auxin led to the efficient and specific depletion of Dbp5 with a half-time of approx. 12 min (Fig. S1A,B), without affecting the overall NPC composition based on the quantification of selected nucleoporins using a previously established method (NuRIM (Rajoo et al., 2018); Fig. S1C). Dbp5 depletion did not affect protein export as an NLS-NES-2xGFP reporter did not enrich in the nucleus after auxin addition. In contrast, blocking Xpo1-dependent export using Leptomycin B (LMB) in an LMB-sensitive Xpo1 mutant (Stade et al., 1997) strongly impaired nuclear export of the GFP reporter (Fig. S1D-G). Furthermore, addition of LMB to cells that were already Dbp5-depleted shifted the GFP reporter distribution towards the nucleus. This shows that nuclear protein import and export remain functional when Dbp5 is depleted (Fig. S1D-G).

Upon Dbp5 depletion, we observed a global nuclear retention of poly(A) RNA using fluorescence *in situ* hybridization (FISH) with an oligo(dT)30 probe, similar to previous reports using temperature-sensitive *dbp5* alleles (Fig. S1H)(Hodge et al., 1999; Snay-Hodge et al., 1998). To investigate whether this oligo(dT)30 staining corresponds to the nuclear retention of mRNA, we examined the export of individual mRNAs. To this end, we marked transcripts of an essential gene, glutamine-fructose-6-phosphate amidotransferase (*GFA1)*, with the PP7-PCP stem loop labeling system ((Larson et al., 2011), Fig. 1A) in the Dbp5 degron background. In the presence of Dbp5, the GFA1-PP7 transcripts were mostly localized in the cytoplasm (Fig. 1B). Upon Dbp5 depletion, transcripts accumulated in the nucleus close to the nuclear periphery. Interestingly, the mobility of nuclear mRNAs was reduced in the absence of Dbp5 based on single particle tracking results obtained from a custom-built microscope ((Smith et al., 2015), Fig. 1C, Video S1, Video S2). To test whether the dynamics of nuclear export factors was also affected, we performed fluorescence recovery after photobleaching (FRAP) measurements after Dbp5 depletion. The nuclear dynamics of the export factors Yra1 and Npl3 as well as an NLS-GFP(3x) control did not change when Dbp5 was depleted (Fig. S2A-F). However, the nuclear mobility of the poly(A)-binding protein Nab2 was dramatically reduced (Fig. 1D-F), with a >5-fold reduction of recovery half-time in the absence of Dbp5. Intriguingly, FRAP of approximately half of the nucleus showed very slow recovery of Nab2 in the entire bleached region (Fig. 1F). The reduction in Nab2 mobility is RNA-dependent as additional depletion of a polymerase II subunit (Rpb2-AID) reverted the phenotype (Fig.

1G,H) by reducing the amount of poly(A) RNA in the nucleus (Fig. S2G). The mRNA-dependent mobility of Nab2 in the nucleus controlled by the DEAD-box ATPase Dbp5 is reminiscent of PBs or SGs which are mRNA-dependent cytoplasmic condensates regulated by the DEAD box ATPases, Dhh1 and Ded1, respectively (Hilliker et al., 2011; Hondele et al., 2019; Mugler et al., 2016; Teixeira and Parker, 2007). Therefore, our results suggest that Nab2 forms an mRNA-dependent condensate in the nucleus *in vivo*, whose dynamics depend on the presence of Dbp5.

Nab2 phase-separates *in vitro*, which is reverted by active Dbp5
The sequence of Nab2 shows many hallmarks of RNA-binding proteins that can undergo LLPS. In addition to an N-terminal domain required for Mlp1 binding and mRNA export, Nab2 contains a Q-rich domain of unknown function (QQQP), an RGG domain with an internal nuclear localization signal (NLS), and 7 zinc fingers (ZnF), of which ZnF5-7 are essential for RNA binding (Fig. S2H)(Anderson et al., 1993; Marfatia et al., 2003; Soniat et al., 2013). Furthermore, according to PONDR, a predictor of naturally disordered regions (Li et al., 1999; Romero et al., 1997), the Q-rich domain and the RGG domain are intrinsically disordered (Fig. S2H). We therefore wanted to examine whether Nab2 can undergo phase separation *in vitro*. Intriguingly, in the presence of poly(A) RNA recombinant Nab2 forms liquid droplets at close to endogenous concentrations (Fig. 2A, Fig. S2I,J). These droplets are dynamic as they had the ability to fuse and grow over time (Video S3), and were mildly sensitive to the aliphatic alcohol 1,6-hexanediol, consistent with them being formed via liquid-liquid phase separation (Fig. S2K,L).
Nab2 condensates form at physiological salt concentrations and are pH-dependent (Fig. 2B). In addition, Nab2 can form condensates with other single-nucleoside polymers (poly(U)/poly(G)/poly(C), Fig. 2C), suggesting that Nab2 does not exclusively bind to the poly(A) tail. Indeed, previous reports have shown that Nab2 can bind also non-poly(A) RNA *in vitro* (Anderson et al., 1993) and associates with mRNA regions upstream of the poly(A) tail *in vivo* (Tuck and Tollervey, 2013).
We next wanted to understand the interplay between Nab2 and Dbp5. *In vitro* work by Tran *et al.* demonstrated that Dbp5 is able to replace Nab2 from poly(A) RNA (Tran et al., 2007). Strikingly, co-incubation of Nab2 with wild type Dbp5 and its ATPase-activators Gle1 and IP$_6$ (Alcázar-Román et al., 2006; Weirich et al., 2006) prevented condensate formation (Fig. S2M). This was dependent on Dbp5 as Gle1 and IP$_6$ alone did not block droplet formation (Fig. S2M). Furthermore, a dominant-active Dbp5 mutant, Dbp5$^{\Delta N90}$, which functions independently of Gle1 and IP$_6$ (Montpetit et al., 2011), also inhibits Nab2 condensation (Fig. 2D). Conversely, the ATP-locked form of Dbp5, Dbp5$^{\Delta N90-DQAD}$, is unable to affect Nab2 condensation (Fig. 2D), suggesting that the ATP hydrolysis cycle is required to inhibit Nab2 condensation.
Together our results indicate that RNA-dependent Nab2 condensation can be regulated by the ATPase activity of Dbp5 and that loss of Dbp5 activity triggers Nab2 condensation.

Glucose stress causes reduction of nuclear Nab2 dynamics and bulk nuclear mRNA retention
Having established that Nab2 forms a condensate *in vitro*, we next asked if Nab2 condensation can also be observed in physiologically relevant conditions in cells. It has been previously observed that Dbp5 re-localizes from the NPC to the nucleoplasm in specific stress conditions, such as acute ethanol stress (Izawa et al., 2005; Takemura et al., 2004). We therefore tested Dbp5 localization upon nutrients stress. Interestingly, NPC localization of Dbp5, but not its cytoplasmic NPC-interaction partners Nup159 and Gle1, is reduced both upon acute (30 min –DEX) and prolonged (24 hrs –DEX) glucose starvation. (Fig. 3A,B and S3A,B). Furthermore, in these starvation conditions, export of bulk mRNA is blocked (Fig. 3C,D). Intriguingly, glucose starvation resulted in the formation of predominantly nuclear Nab2 foci,

while other export factors such as Yra1 and Npl3 were unaffected (Fig. 3C,D and S3C,D). Discrete Nab2 foci form during short-term glucose withdrawal and focus formation is enhanced upon long-term glucose starvation (Fig. 3C,D). Nab2 foci are mostly resistant to 1,6-hexanediol treatment, which dissolves cytoplasmic membraneless organelles such as P-bodies, but not stress granules (Fig. S4A-C) (Kroschwald et al., 2015). Nab2 focus formation is accompanied by nuclear accumulation of poly(A) RNA in a punctate pattern overlapping with Nab2 foci (Fig. 3C, D). The poly(A) RNA that is retained in these foci seem to correspond to intact transcripts as they also show co-localization with the cap-binding complex proteins Cbc1 and Cbc2 ((Izaurralde et al., 1994), Fig. S4G). Nab2-mRNA foci are reversible, as release from starvation conditions into medium containing glucose led to the full dispersal of Nab2 in the nucleus within an hour (Fig. S4D). These results suggest that Nab2 condensation confers nuclear enrichment of bulk mRNA in long-term starvation.

We then tested Nab2 mobility in short-term and long-term starvation. Similar to acute Dbp5 depletion, we observed a reduction in nuclear dynamics after 30 min of acute glucose starvation (Fig. 3E,F). This phenotype was even more prominent upon long-term glucose starvation (24 hrs) (Fig. 3G,H). The reduction in dynamics was specific for Nab2 as the mobility of other export factors such as Npl3 and Yra1 was not affected (Fig. S3E-H). Nab2 mobility reduction was dependent on the presence of mRNA and treatment with the transcription inhibitor thiolutin (Tipper, 1973) restored Nab2 mobility to the levels of non-stressed cells (Fig. S4E,F). We next wanted to examine whether Nab2 focus formation is triggered by the loss of Dbp5 from the NPC during long-term stress. Intriguingly, when Dbp5 is prevented to leave the NPC, by using a strain in which Dbp5 is permanently fused to its interaction partner Nup159 ('Dbp5 tether', Dbp5-$^{\Delta N}$Nup159, (Adams and Wente, 2020)), Nab2 focus formation is strongly reduced and the mobility of Nab2 is only mildly affected in long-term glucose withdrawal (Fig. 3I-K and S3I). This suggests that reduction of Dbp5 abundance at the NPC in response to glucose starvation is required for Nab2 focus formation.

We conclude that upon glucose starvation, RNA-dependent Nab2 focus formation correlates with reduction of Dbp5 abundance at the NPC to facilitate bulk mRNA retention in the nucleus.

Protein-protein dimerization, but not predicted IDRs, drives Nab2 condensation

Nab2 is predicted to contain several IDRs including an RGG and a polyQ domain (Fig. S2H). To identify the domains in Nab2 that are responsible for the alterations in nuclear dynamics in glucose stress, we tested if removal of the IDR domains would affect condensation behavior *in vitro*. Deleting the Q-rich domain did not abolish condensate formation, but slightly shifted the pH-dependency to higher pH values (Fig. S5A). Deleting the RGG domain and replacing it with an exogenous nuclear import signal (ΔRGG^{SV40NLS}, with the SV40NLS replacing the endogenous NLS found in the RGG domain) also affected the pH sensitivity of Nab2, which was even more pronounced than for Nab2$^{\Delta Q}$-GFP (Fig. S5A). Surprisingly, when both domains were deleted (Nab2$^{\Delta Q-\Delta RGGSV40NLS}$-GFP, 'IDR mutant'), condensates could still form, even though the major regions expected to be critical for condensation were removed. In addition, droplets with this Nab2 variant even formed in the absence of RNA (Fig. 4A) and were less dynamic and more solid-like compared to their wild type counterparts, as shown by a reduction in recovery by FRAP (Fig. 4B). We then tested a Nab2 variant that has been previously described to affect mRNA compaction by impairing Nab2 dimerization on RNA (Nab2^{F450A}, (Aibara et al., 2017)). Intriguingly, this Nab2 dimerization mutant displayed a strong reduction in condensate formation (Fig. 4A). Thus, our results suggest that Nab2 condensate formation does not depend on the IDRs but instead requires the protein's ability to dimerize on RNA, Furthermore, the major disordered regions within the protein, in particular the RGG domain, help to keep Nab2 condensates dynamic.

The type of Nab2 condensate determines the degree of nuclear mRNA retention

How does the ability of Nab2 to undergo phase separation affect nuclear focus formation and RNA retention *in vivo*? Whereas the expression of a Nab2 variant that lacks the Q-rich domain (ΔQ) does not result in any growth disadvantages, deletion of the RGG domain (ΔRGG^{SV40NLS}) leads to a severe growth impairment (Fig, S5B). Intriguingly, the combined deletion of both the Q and the RGG domain (ΔQ-ΔRGG^{SV40NLS}, 'IDR mutant') rescues the growth defect of the ΔRGG^{SV40NLS} mutant (Fig. S5B), and we therefore continued our experiments with this mutant (Fig. S5C). Consistent with our *in vitro* results, both Nab2^{ΔQ}-GFP and Nab2$^{ΔQ-ΔRGG-SV40NLS}$-GFP still formed foci in the nucleus upon long-term glucose starvation (Fig. 4C and Fig. S5E) coinciding with a reduction in nuclear mobility compared to WT in FRAP (Fig. 4D,E and Fig. S5F,G). In contrast, the dimerization mutant Nab2^{F450A} formed almost no nuclear foci (Fig. 4C) and also did not show a reduction in nuclear mobility (Fig. 4F). This suggests that the degree to which Nab2 mutants form condensates *in vitro* reflects focus formation and nuclear mobility *in vivo*.

Intriguingly, the capability of Nab2 to form foci in stress is also linked to the ability to retain RNA in the nucleus: while the IDR mutant shows an increased nuclear poly(A) RNA signal, the dimerization mutant displays decreased nuclear poly(A) RNA retention during stress (Fig. 5A,B). This implies that the level of Nab2 condensate formation determines the degree of nuclear RNA retention in long-term glucose starvation. Of note, the Nab2 IDR mutant also shows a slight nuclear mRNA retention phenotype already in log growing cells (Fig. 5A, S5H), suggesting that the RGG domain is important for efficient mRNA export both in unstressed and stressed cells.

Nab2 condensation confers selective RNA retention in stress

In order to identify which mRNAs are retained in nuclear Nab2 foci in stress, and examine if this altered in the Nab2 mutant background (Nab2$^{ΔQ-ΔRGGSV40NLS}$-GFP and Nab2^{F450A}-GFP) we next purified poly(A)-RNA from whole cells cultured in +glucose and 24 hrs –glucose conditions, as well as from nuclear fractions of cells starved for 24 hrs in –glucose (Fig. S5J,K and Supplementary Table 4). As expected, we detected a global rearrangement of the transcriptional landscape between the +glucose and 24 hrs –glucose conditions on a whole cell level, (Fig. S6A), in agreement with RNA FISH quantifications showing a 5-fold reduction of total poly(A) RNA abundance in 24 hrs –glucose compared to +glucose (Fig. S5I). However, for a given condition, the RNA sequencing reads of the wild type and the mutants did not differ (Fig. S6A) and a stress response was initiated to a similar extent in all strains at the total RNA level (Fig. S6B). By contrast, differences in the nuclear RNA fractions could be observed between the wild type and the Nab2 mutants in the 24 hrs –glucose revealing an increase in sequencing reads (CPM, counts per million) for the IDR mutant Nab2$^{ΔQ-ΔRGGSV40NLS}$ compared to wild type, whereas the dimerization mutant Nab2^{F450A} did not significantly differ from the wild type (Fig. 5C and S6D). Intriguingly, the IDR mutant displayed selective nuclear enrichment of certain transcripts whereas others did not show any selective enrichment (Fig. 5C). Gene ontology (GO) term analysis of the most abundant enriched transcripts (CPM >500 (318 genes) representing 42% of the total reads; Fig. 5C (blue dots), Fig. 5D (purple) and Fig. S6E,F) revealed the highest GO scores in stress response (oxidation-reduction processes, cellular response to heat), carbohydrate metabolic processes and glucose import. Transcripts that were not differentially enriched fraction (Fig. 5D, grey) showed highest GO scores in translation, rRNA processing, glycolytic process/gluconeogenesis and proton transport (i.e. 'pro-proliferation' mRNAs). To confirm our RNA sequencing results, we also performed RNA FISH experiments using probes against individual mRNAs that either belong to the class of differentially enriched mRNAs (*BTN2, HSP42, HXT6*, Fig. 6A-F) or to the not differentially enriched mRNAs (*DED1*, Fig. 6G,H).

Indeed, these experiments confirmed nuclear RNA retention and condensate selectivity for the tested transcripts.

Our results reveal that wild-type Nab2 retains pro-proliferation mRNAs in the nucleus that are not induced in stress allowing the selective export of stress-induced mRNAs (Fig. 6). The altered condensate state seen in the IDR mutant Nab2$^{\Delta Q-\Delta RGGSV40NLS}$ prevents the selective transport of stress-induced mRNAs, which results in their nuclear accumulation.

Correct Nab2 condensation is essential for survival after prolonged stress

Finally, we also asked whether there is a physiological consequence of the altered mRNA retention in long-term stress. We therefore washed cells acutely into medium without glucose and incubated them for 5 days before allowing them to recover on rich medium plates containing dextrose. Compared to wild type, the IDR mutant Nab2$^{\Delta Q-\Delta RGGSV40NLS}$ displays a slight growth disadvantage on rich medium plates in log growing cells, which gets exacerbated after prolonged glucose withdrawal (Fig. 7A) and when stress is maintained after dextrose recovery by incubating cells at different temperatures (Fig. 7A and S7A). Furthermore, gradual glucose depletion also affects the recovery of the IDR mutant as cells recovered from early stationary phase (24 hrs after mid-log phase) show a similar growth impairment as after acute glucose withdrawal (Fig. S7B).

In summary, our results suggests that aberrant mRNA localization caused by altered Nab2 condensation affects survival in and recovery from glucose stress conditions.

Discussion

Here we show that the mRNA export factor Nab2 can form biomolecular condensates *in vitro* and *in vivo*, which are counteracted by the activity of the DEAD-box ATPase Dbp5. Our results provide evidence that the nuclear RNA-dependent Nab2 condensates globally retain bulk mRNAs in the nucleus in physiologically relevant stress conditions. Furthermore, our results suggest that a group of stress-induced mRNAs do not accumulate in the nucleus and predominantly localize in the cytoplasm. Therefore, we propose a model in which stress-related transcripts can escape nuclear retention to elicit a timely cellular stress response, suggesting that cells use selective confinement in or release from nuclear condensates to re-wire mRNA export during stress (Fig. 7B). Nuclear retained mRNAs are intact as both the cap-binding complex as well as the poly(A)-binding protein Nab2 enrich in stress-induced nuclear RNA foci, and largely represent pro-proliferation transcripts (Fig. 5C,D and Fig. S4G). Storing them in the nucleus during glucose starvation might be a cellular strategy required for re-initiation of cell growth after stress release, which could allow cells to quickly re-start protein production in a transcription-independent fashion, minimizing energy consumption at early stages of re-growth. Nuclear retained mRNAs tend to be longer (Fig. S6G) and more likely stem from intron-containing genes (Fig. S6H) than the RNAs that evade nuclear retention (e.g., those induced in glucose stress, Fig. 5D), although it is unclear if those mRNAs still contain introns once they are present in nuclear Nab2 condensates. Nab2 has been implicated in splicing regulation and artificial depletion of Nab2 from the nucleus leads to transcriptional read-through which produces chimeric transcripts with retained introns (Alpert et al., 2020; Soucek et al., 2016). RNA length has been reported to correlate with enrichment in cytoplasmic stress granules and poor translation efficiency (Khong et al., 2017), suggesting that short transcripts, such as mRNAs responsible for the cellular stress response, could evade both nuclear and cytoplasmic condensates to allow for their preferential translation in stress conditions. Furthermore, nuclear mRNA retention has previously been reported in mammalian systems, where it serves to rapidly modify the neuronal

transcriptome upon stimulation (Mauger et al., 2016), or to buffer gene expression noise (Bahar Halpern et al., 2015). Buffering mechanisms could also be at play in yeast, which needs to be investigated in the future.

We show that Nab2 condensate formation is regulated by the activity of Dbp5 (Fig. 2D and Fig. S2M), and that forcing Dbp5 to nuclear pore complexes (NPCs) via fusion to $^{\Delta N}$Nup159 prevents stress-induced Nab2 mobility reduction and nuclear focus formation (Fig. 3I-K). Previous reports have demonstrated a transient Nab2 and Dbp5 interaction at the NPC using a splitVenus system (Adams and Wente, 2020). We speculate that Dbp5 could generally act to dissolve RNA-containing condensates. Consistent with this, overexpression of the mammalian ortholog of Dbp5, DDX19, prevents stress granule formation in an RNA-binding dependent manner (Hochberg-Laufer et al., 2019; Tauber et al., 2020). However, the local ATPase activation through its interaction partner Gle1, which exclusively resides at the NPC, could target Dbp5's activity largely towards specific RNPs such as Nab2-bound mRNAs that pass the NPC during export. In line with this, Dbp5-$^{\Delta N}$Nup159 localization has been shown to be sufficient for viability (Adams and Wente, 2020), suggesting that the essential role of Dbp5 is restricted to the NPC.

Nuclear Nab2 condensation depends on several domains that modulate the degree of condensation and the material properties of the condensates controlling the degree of nuclear RNA retention (Fig. 5A,B). Surprisingly, when we deleted the major regions predicted to be causal for condensation (Fig. S2H), the Q-rich domain and the RGG domain, we observed a stronger Nab2 condensation phenotype, indicating that these domains act together to function as 'liquefiers' of condensates or to block condensation altogether. Similar observations have been made for the stress granule component Ded1, where deletion of evolutionarily preserved residues in its N-terminal intrinsically disordered region (IDR) lead to constitutive SG formation (Iserman et al., 2020).

Conversely, Nab2 condensation (and nuclear focus formation/RNA retention) depends on its ability to dimerize on RNA (Nab2^{F450A}, Fig. 4A,C and 5A). The dimerization mutation lies within the zinc fingers (ZnF) essential for RNA binding, highlighting the multivalent binding mode of Nab2 as well as the importance of non-IDR regions for condensate formation. Indeed, Yang *et al.* have previously shown that increasing the number of ZnFs promotes cytoplasmic condensation of the SG component G3BP1, highlighting how altering RNA binding strength and potentially the valency of interaction affects condensate formation (Yang et al., 2020).

Glucose starvation is accompanied by an intracellular pH drop from pH 7.5 to pH 6-6.5 (Dechant et al., 2014; Joyner et al., 2016; Orij et al., 2009). Intriguingly, altered condensation of the IDR mutant Nab2$^{\Delta Q-\Delta RGGSV40NLS}$ resulted in pH insensitivity *in vitro* (Fig. 4A) and a slight nuclear enrichment of poly(A) RNA already in +DEX conditions (Fig. 5A and Fig. S5H), suggesting that Nab2 condensation could also play a role in mRNA export in exponentially growing cultures. Future research needs to determine to which extent mRNP (micro)condensation is also involved in mRNA export in unstressed cells, and if the essential function Dbp5 in mRNA export is to locally prevent or modulate mRNP condensation.

Materials and Methods

Supplementary Table 1 (Strain list)

Strain name	Genotype	Figure	Reference
KWY6116	*MAT A his3∆1 leu2∆0 ura3∆0 NDC1::NCD1-yomKate2-URA3 GFA1::GFA1-24xPP7stemloops::loxP-kanR-loxP pMET25-PP7CP-yEGFP::LEU2*	1B,C	This study
KWY6353	*MAT A his3∆1 leu2∆0 ura3∆0 NDC1::NCD1-yomKate2-URA3 GFA1::GFA1-24xPP7stemloops pMET25-PP7CP-yEGFP::LEU2*	1B,C	This study
KWY8262	*MAT A leu2 his3 NAB2::NAB2-yEGFP-URA3 DBP5::DBP5-3V5-IAA7-KANMX trp1-1:: dsRed-HDEL*	1D-H	This study
KWY8263	*MAT alpha leu2 NAB2::NAB2-yEGFP-URA3 DBP5::DBP5-3V5-IAA7-KANMX trp1-1:: dsRed-HDEL HIS::pGPD1-osTIR-HIS*	1D-H	This study
KWY8258	*MAT A leu2 his3 NAB2::NAB2-yEGFP-URA3 DBP5::DBP5-3V5-IAA7-KANMX RPB2::RPB2-3V5-IAA7-KANMX trp1-1:: dsRed-HDEL*	1G,H	This study
KWY8395	*MAT A leu2 his3 NAB2::NAB2-yEGFP-URA3 DBP5::DBP5-3V5-IAA7-KANMX RPB2::RPB2-3V5-IAA7-KANMX trp1-1:: dsRed-HDEL pNH603-pGPD1-osTIR::HIS*	1G,H	This study
KWY6567	*MAT alpha his3-11,15 trp1-1 ura3-1 ade2-1 DBP5::DBP5-link-yEGFP::CaURA3 TRP1:: dsRed-HDEL*	3A,B S3A,B S3I	Rajoo *et al.* PNAS 2018
KWY9296	*MAT alpha his3∆1 leu2∆0 lys2∆0 ura3∆0 NAB2::NAB2-yEGFP-HISMX*	3C-H, 4C,D,G 5A-C 6A-H 7A S4A-F S5C,D,H -K S6A-H S7A,B	This study
KWY9399	*MAT alpha his3∆1 leu2∆0 lys2∆0 ura3∆0 NAB2::NAB2-∆QQQP-∆RGG(∆104-263)-SV40NLS-yEGFP-HISMX*	4C,E,G 5A-C 6A-H 7A S5C,H-K S6A-H S7A,B	This study
KWY9823	*MAT alpha his3∆1 leu2∆0 lys2∆0 ura3∆0 NAB2::NAB2-F450A-GFP-HISMX*	4C,F,G 5A-C 6A-H 7A S5D,H-K S6A-H S7A,B	This study
KWY10392	*MAT A his3-11,15 ura3-1 leu2-3 trp1-1Nab2-GFP::HIS*	3I-K	This study
KWY10394	*ura3-1 his3-11 leu2-3 dbp5∆::KAN nup159∆::KAN pDBP5-NUP159∆N::LEU NAB2::NAB2-GFP-HIS*	3I-K	This study

KWY7462	*MAT alpha TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX pNH603-pGPD1-osTIR::HIS*	S1A	This study
KWY6037	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP84::NUP84-link-yEGFP::CaURA3 TRP1::dsRed-HDEL*	S1C	Rajoo *et al.* PNAS 2018
KWY6041	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP133::NUP133-link-yEGFP::CaURA3 TRP1::dsRed-HDEL*	S1C	Rajoo *et al.* PNAS 2018
KWY6048	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP82::NUP82-link-yEGFP::CaURA3 TRP1::dsRed-HDEL*	S1C	Rajoo *et al.* PNAS 2018
KWY6053	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP60::NUP60-link-yEGFP::CaURA3 TRP1::dsRed-HDEL*	S1C	Rajoo *et al.* PNAS 2018
KWY6054	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP159::NUP159-link-yEGFP::CaURA3 TRP1::dsRed-HDEL*	S1C S3A,B	Rajoo *et al.* PNAS 2018
KWY6921	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP159::NUP159-link-yEGFP::CaURA3 TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX*	S1C	This study
KWY7056	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP159::NUP159-link-yEGFP::CaURA3 TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX pNH603-pGPD1-osTIR::HIS*	S1C	This study
KWY7407	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP133::NUP133-link-yEGFP::CaURA3 TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX*	S1C	This study
KWY7408	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP82::NUP82-link-yEGFP::CaURA3 TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX*	S1C	This study
KWY7412	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP60::NUP60-link-yEGFP::CaURA3 TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX*	S1C	This study
KWY7444	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP133::NUP133-link-yEGFP::CaURA3 TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX pNH603-pGPD1-osTIR::HIS*	S1C	This study
KWY7446	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP82::NUP82-link-yEGFP::CaURA3 TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX pNH603-pGPD1-osTIR::HIS*	S1C	This study
KWY7458	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 NUP60::NUP60-link-yEGFP::CaURA3 TRP1::dsRed-HDEL DBP5::DBP5-GFP-IAA7-KANMX pNH603-pGPD1-osTIR::HIS*	S1C	This study
KWY8118	*MAT A his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX DBP5::DBP5-3V5-IAA7-KANMX xpo1Δ::HYGMX pRS313-XPO1(LMBsensitive)::HIS3MX pADH1-NLS-NES-2xGFP::URA3*	S1B,D-G	This study
KWY10381	*MAT A his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX DBP5::DBP5-3V5-IAA7-KANMX xpo1Δ::HYGMX pRS313-XPO1(LMBsensitive)::HIS3MX pADH1-NLS-NES-2xGFP::URA3 pNH605-pGPD1-osTIR::LEU2*	S1D-G	This study
KWY9559	*MAT alpha his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 NAB2::NAB2-yEGFP-HISMX DBP5-3V5-IAA7::KANMX*	S1H	This study
KWY9579	*MAT alpha his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 NAB2::NAB2-yEGFP-HISMX DBP5-3V5-IAA7::KANMX LEU2::pNH605-pGPD1-osTIR1*	S1H, S2G	This study
KWY7642	*MAT A his3Δ1 leu2Δ0 met15Δ0 DBP5::DBP5-3V5-IAA7-KANMX YRA1::yEGFP-YRA1-NATMX*	S2A,B	This study
KWY7652	*MAT A his3Δ1 leu2Δ0 met15Δ0 DBP5::DBP5-3V5-IAA7-KANMX YRA1::yEGFP-YRA1-NATMX TRP1::pGPD1-osTIR-hphNT1*	S2A,B	This study
KWY8009	*MAT A his3Δ1 leu2Δ0 met15Δ0 TRP1::dsRed-HDEL-NATMX*	S2C,D	This study

	DBP5::DBP5-3V5-IAA7-KANMX NPL3::NPL3-yEGFP-URA3		
KWY8010	*MAT A his3Δ1 leu2Δ0 met15Δ0 TRP1::dsRed-HDEL-NATMX DBP5::DBP5-3V5-IAA7-KANMX NPL3::NPL3-yEGFP-URA3 pNH603-pGPD1-osTIR::HIS*	S2C,D	This study
KWY7855	*MAT A his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX DBP5::DBP5-3V5-IAA7-KANMX p(PHO4-NLS-GFP3 ::URA3)*	S2E,F	This study
KWY7949	*MAT A his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX DBP5::DBP5-3V5-IAA7-KANMX HIS::pNH603-pGPD1-osTIR p(PHO4-NLS-GFP3 ::URA3)*	S2E,F	This study
KWY8259	*MAT alpha leu2 NAB2::NAB2-yEGFP::URA3 DBP5::DBP5-3V5-IAA7::KANMX RPB2::RPB2-3V5-IAA7-KANMX trp1-1::dsRed-HDEL-TRP1 HIS::pGPD1-osTIR-HIS*	S2G	This study
KWY8050	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 DHH1::DHH1-yEGFP-URA3*	S2I,J	This study
KWY8650	*MAT A his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 TRP1::dsRed-HDEL-URA3 NAB2::NAB2-yEGFP-HIS3*	S2I,J S5B	This study
KWY6051	*MATalpha his3-11,15 trp1-1 ura3-1 ade2-1 GLE1::GLE1-link-yEGFP::CaURA3 TRP1::dsRed-HDEL*	S3A,B	Rajoo *et al.* PNAS 2018
KWY6268	*MAT A his3Δ1 leu2Δ0 met15Δ0 YRA1:: NATMX-yEGFP-YRA1*	S3C,E,F	This study
KWY8008	*MAT A his3Δ1 leu2Δ0 met15Δ0 TRP1::dsRed-HDEL-NATMX DBP5::DBP5-3V5-IAA7-KANMX NPL3::NPL3-yEGFP-URA3*	S3D,G,H	This study
KWY10277 (SWY6285)	*MAT A ura3-1 his3-11 leu2-3 dbp5Δ::KAN nup159Δ::KANN GFP-DBP5-NUP159ΔN::LEU*	S3I	Adams & Wente, PLoS Genetics 2020
KWY7690	*MAT A his3Δ1 leu2Δ0 met15Δ0 CBC2::CBC2-GFP(S65T)-HIS3MX*	S4G	Huh *et al.* Nature 2003
KWY7696	*MAT A his3Δ1 leu2Δ0 met15Δ0 CBC1::CBC2-GFP(S65T)-HIS3MX*	S4G	Huh *et al.* Nature 2003
KWY8215	*MAT alpha his3Δ1 leu2Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX NAB2::NAB2-GFP-Y66L-IAA7-KANMX pNH603-pGPD1-osTIR::HIS*	S5B	This study
KWY9288	*MAT alpha his3Δ1 leu2Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX NAB2::NAB2-GFP-Y66L-IAA7-KANMX pNH603-pGPD1-osTIR::HIS p(Nab2-yEGFP-LEU2 (CEN))*	S5B	This study
KWY9289	*MAT alpha his3Δ1 leu2Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX NAB2::NAB2-GFP-Y66L-IAA7-KANMX pNH603-pGPD1-osTIR::HIS p(Nab2-ΔRGG(Δ201-263)-SV40NLS-yEGFP-LEU2 (CEN))*	S5B	This study
KWY9290	*MAT alpha his3Δ1 leu2Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX NAB2::NAB2-GFP-Y66L-IAA7-KANMX pNH603-pGPD1-osTIR::HIS p(Nab2-ΔQQQP-ΔRGG(Δ104-263)-SV40NLS-yEGFP-LEU2 (CEN))*	S5B	This study
KWY8666	*MAT A his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0 TRP1::dsRed-HDEL-natMX NAB2-ΔQQQP(Δ104-169)-yEGFP-HISMX*	S5C	This study
KWY10234	*MAT? his3Δ1? leu2Δ0? met15Δ0? lys2Δ0? ura3Δ0 NAB2-ΔQQQP(Δ104-169)-yEGFP-HISMX*	S5E-G	This study

Supplementary Table 2 (Plasmid List)

Plasmid name	description	Figure	Published

pKW2499	6xHIS-TEV-ΔN90-Dbp5 (KAN)	2C	Montpetit *et al.* Nature 2011
pKW2501	6xHIS-TEV-ΔN90-Dbp5-DQAD (KAN)	2C	This study
pKW3714	6xHis-V5-Dbp5-mCherry (AMP)	S5B	This study
pKW4404	6xHis-V5-Gle1 s.c. [244-538] (AMP)	S5B	This study
pKW4814	6xHis-TwinStrep-PreScission-Nab2-yEGFP (AMP)	2A-C 4A S5B-D	This study
pKW4817	6xHis-TwinStrep-PreScission-Nab2-ΔRGG(Δ201-263)-SV40NLS-yEGFP (AMP)	S8G	This study
pKW4818	6xHis-TwinStrep-PreScission-Nab2-ΔQQQP-ΔRGG(Δ104-263)-SV40NLS-yEGFP (AMP)	4A	This study
pKW4954	6xHis-TwinStrep-PreScission-Nab2-F450A-yEGFP (AMP)	4A	This study
pKW4957	6xHis-TwinStrep-PreScission-Nab2-ΔQQQP(Δ104-169)-yEGFP (AMP)	S8G	This study

Supplementary Table 3 (Oligo list (smFISH))

Oligo	Sequence	Modification	Company
dT30	TTT TTT TTT TTT TTT TTT TTT TTT TTT TTT	TAMRA (3' end)	Microsynth
BTN2	1 GGCAGCTGTTCAAAAACACA 2 CGAATGTAGGGGCTGACTAA 3 ATAGCTCACTGGAGAACTGC 4 TTTCCTCCTTTTACATTCTG 5 GCTCTTAGGTTAGCTTTTAT 6 ATTTGCATCGCTTTTTTTTG 7 TGCATACCTCAAAGGTTCTG 8 TAACCATTTGGTGTTTCAGC 9 GTTCATATGGAATCCGCTTA 10 ACCTAATTTCTCATTAACGT 11 TGTAATGGTTCTCCTTTAGC 12 TCTTGGACAACATGGTAAGT 13 TCCGCTTCATCTTCAACATA 14 GCAGATCTCAATAGAGCATC 15 GGCTCTAAAATCCAGATCTT 16 TCCTTAGCAATTTTCTTTCC 17 TTCTATTTCGTAGTCTTGGA 18 ATTCATCACCTCTATGATTC 19 TCCTTCTTACTCAATATGCT 20 CTTGGTCTAGAGAGAATTCC 21 CAACTCCACAGCCAATAACA 22 ATATTTTTCTCTCGAGCCAT 23 CCTCATGCTTAACTAAACCA 24 ATGCCACCTTCGGAAATTTC 25 TTATTGGCATCTTTGGTTCG 26 GACTCGTCTATTTTGGATTC 27 AGACATGTTAACATCATCGT 28 CTTCTTCCTCCTTCAAAGAT	Cal Fluor Red 590 (3' end)	BioCat

	29	TTTGGTTAGTGGTTCTTTCG		
	30	TCCATTTTTTTATTTGGTCT		
	31	TGCATCAATCTTTCTTCCTC		
	32	CTGTTCTGATTTTCTGCTTT		
	33	TTTTGCCTTTCTTCATCTTC		
	34	CAATCTGGCTTCCTTCTCTT		
	35	TCAAAGATTCTTTCCTTGCC		
	36	CTCTTGGTTTTTTGCTTATT		
	37	TTGCAATTTTTTTTGCTGGG		
	38	TCTCAGAGATAGGCAATGAT		
	39	TTATTTTTATTGCTGGCCTC		
	40	TATCACTTTCTGCTGAACCA		
	41	GTCGTATCAGAATCACTGTT		
	42	ACCAGACACAGAGAAATCCA		
	43	GAAGCGTGTTTTTTTAGTGT		
	44	CTCAACGTCTTCTAATAGGG		
	45	TGTATCTGTCAACTTCCTCA		
	46	GGAGATCTGCTTAGGGACTC		
	47	CCTCAATAATAGAGTTTCCC		
HSP42	1	AGAGATAGGGATGGTTGATA	Cal Fluor Red 590 (3' end)	BioCat
	2	CCAGTTTGGTTGGATAATGC		
	3	TTTGTGGTTGGCGAGGATAT		
	4	ACGTGCACTTGTCCATAATG		
	5	ATGATGACGAGGATGATGCC		
	6	AGGAACACCATTGTATCTGC		
	7	TCCAGGGAACTGGTAGTAAT		
	8	CAGGACTATAGTAATAGGCT		
	9	ACCATCCTCGTCATCATAAC		
	10	ACCATATCTTCGTCTTGGTC		
	11	CATCTTCCTGTCTTGTAGTG		
	12	TGGATATCTTCTCGAGTTGC		
	13	TCCTGGCAGTATTACAATGG		
	14	CGTTTAAACTGTTTGCCTGT		
	15	GAACACCTATTAACGCGGTT		
	16	TGGTTCAGTGCCTTCATATG		
	17	CTGTTCGGTATTTGCTTCAA		
	18	TTCTTATCCTTTTCTTCTCC		
	19	TTTGGGTGCTTCAGACTTAT		
	20	TTTTTCTTTGTTGGTTTCGC		
	21	ATTCCTCCAGCTGATTCAAA		
	22	TTTTTTGGCTAATGGTGGTC		
	23	TGTAGGTGAGCGAACGATGA		
	24	GTCAGGTATTGGGGAAGGCG		
	25	CAGGCTTGGATACTTGTAAC		
	26	AATGGTAAGTCCATTCTCGT		
	27	CATAGACATTCACTTCTGGT		
	28	AACTACGTAAGTGTCCTCGG		
	29	TAGCACCTGGTAACGCAAGA		

	30	TCAATGTGGAAAGCCCTAGA		
	31	CTCATGAGAAGATGGATGGT		
	32	TCGATCTTACCCTTGATGAG		
	33	TTTCATCAATGCCCACTCTG		
	34	AGTTCCGTGATCTTCAAGAA		
	35	TCTCTCAAACGCACCATATT		
	36	GCAGCACGGGGAATTTAACG		
	37	ATTTCTTCGTCCTTAATGCG		
	38	ACCGTTGTTGTAAGTAGCTT		
	39	TAGGCACCTTAATTTGTAGT		
	40	GCTTTTCAGTGTCATTGACA		
	41	AATGGCGATCCTCTTTTTTG		
	42	CAATTCTTCGTCGGGTATTT		
	43	CAATTTTCTACCGTAGGGTT		
HXT6	1	AGGAGTTTGCTCTGCAATAG	Cal Fluor Red 590 (3' end)	BioCat
	2	TCAACAGCAGAGAGATGCTC		
	3	GATAAAACCGAGTGGGAGGC		
	4	TCTTTCAGCCTTGTTTGATG		
	5	ACCTTCACCATAAGCTTTTA		
	6	CTGGTCTCTTTGGAATTTCA		
	7	GAGACAGTGACATAGGCAGA		
	8	ACGAAACCACCAAAGGCGAT		
	9	AAATGGTACCAGTATCCCAA		
	10	AGTACCATCTTTATGCTTCA		
	11	ACCAGTTCTAACCTTAGACA		
	12	CACCAATGGCACAACCAATG		
	13	CCGTACATATCACCCAATTT		
	14	CGACAATCAAACCCACCTTA		
	15	AACGGCAATACCACCAACAC		
	16	GCTTTGGGGATACTTCAGAA		
	17	CAAGAGACTAAAGTACCCCT		
	18	ACCGGCAGTAATCATCAATT		
	19	GTTGGTACAGTAACCCAAGA		
	20	TGGAGTAGTTCTTAGTACCG		
	21	ACCTAATGGAACTCTCCATT		
	22	TCATAAACAAAGCCCAGGCA		
	23	ACGTGGAGACTCTGGAACAA		
	24	TTCGATCTTACCGACTTCAG		
	25	ACGGCAATAGAACGTTTGGC		
	26	TGGATCATCAACAGCAACCT		
	27	AGCTTCGACTTCAGCCAAAA		
	28	TTACCAGCTAATTTCTCTGC		
	29	ACTAAACAATTCACCCCAGG		
	30	AAACGCTGAAGGACCTTTGT		
	31	TTGAATCATAGCACCCATGA		
	32	CACCTGTCAATTGTTGTAGA		
	33	ACTCAAACCAACAGCCTTGA		
	34	ACCCCATAGCAAACAAGTAC		

	35	TACAAGCAGTCATGGATGCA		
	36	ATTTGGCCATAATCTGGTGA		
15	37	CTTGGAAGATGGTTGGTCTT		
	38	AGCCCATGTAGTAGCAAAAC		
	39	AGACTTGACTCTCAATGGGA		
	40	GCAGCTGTAGCAATAGACAT		
	41	TCAAGAAACCCACAACCAA		
	42	CAGCCCATGAAAACGTAACC		
	43	CCCTTAGTTTCTGGAACAAC		
	44	TGGTGTTGACTTCTTCCAAA		
	45	CATGGTAGAACACCTTCTTC		
	46	TAGATGGTGGAACCCATGAG		
	47	TTATCATCGTGAGCCATTTC		
	48	TTTGGTGCTGAACATTCTCT		
DED1	1	AATTTTGCACTTGTTCGCTC	Cal Fluor Red 590 (3' end)	BioCat
	2	TAACCATTCTCGTTGTTGTC		
	3	GTTTTCCTCTTAAGTGAGGA		
	4	ACTGTTATTTCTGGCACTTC		
	5	CCGTTGTTGTTATTGTAGTT		
	6	TTGTTGCTAAAGAAGCTGCC		
	7	TTGTTTCCACCGAAGAAACC		
	8	CACCAAATATGGCGATCTCG		
	9	TGGAAATTTGGATCCTCGGG		
	10	CGTCCACTGGAATATCATCG		
	11	TGGTTCAGGAACATCCTTAC		
	12	TTCCAATAACAATCCGTCCA		
	13	TGGTGAAACGGGCCAATTTG		
	14	GGAGTATTTTTGCACAGGTG		
	15	TGATTCGGACAACACTGGGA		
	16	GAGACTCTGGTTGAGGAGAT		
	17	GGCCTTTCTTTGGTAAAAGG		
	18	TCTCTAGTTGGAGCCATAAT		
	19	ATCGAAAATTTGGGTGGCCA		
	20	TTGACCCAGGATCTATAAGT		
	21	TTACCAATTGGAGAACCACC		
	22	AGATCGCAACCACGTTCAAT		
	23	TTCAAACGACCTGGAGTAGC		
	24	TTTACCACGTTCCAACAAGT		
	25	TACTTGACGTTGGCCAAAGA		
	26	ATCAGCTTCATCTAGAACCA		
	27	GTTCGAAACCCATATCCAAC		
	28	CAGGAGTCATATCACAGTCT		
	29	ATCAGAGTTTGTCTTTCACC		
	30	TCAGCGGGAAAAGTAGCTGA		
	31	AATCACGGGCCAAATGTTGG		
	32	AACCGACTCTACCAACAGAC		
	33	AGGCTGACTTCTTATCTTGA		
	34	AGTCAAACCGTCAGTGGATG		

	35 TTGATCTGCCATTCTCTTAG		
	36 GCATGATCAGGAAATCGGTC		
	37 ATGAATGGCGGTAGCTCTAA		
	38 TTCTCTCTCAGATTGGGTAC		
	39 ACAGCTGTGGCAACCAATAA		
	40 ATAACGTGGGTGACGTTTGG		
	41 GACCAGTTCTACCAATTCTA		
	42 AAAAGGCAGTGGCAAGACCG		
	43 TCTTGGTTAGCTTCAGTCAA		
	44 CGTCCTTCAAGAATGATGGG		
	45 AATCTCTGTTGTTGTTGCGA		
	46 GAGTTATCTCTGCTTCTTGA		
	47 GAAGACTTGGAATCGCTACC		
	48 CACCACCAAGAAGAGTTGTT		

Yeast strain construction

Saccharomyces cerevisiae strains were constructed using standard yeast genetic techniques either by transformation of a CEN plasmid, a linearized plasmid, or of a PCR amplification product with homology to the target site (Longtine et al., 1998). All yeast strains expressing a GFP-tagged protein were tagged with a yeast-optimized monomeric EGFP version (A206K).

Nab2-ΔQ and Nab2-ΔQ-ΔRGG^{SV40NLS} mutant strains were constructed using a CRISPR/Cas9 strategy (Laughery et al., 2015). In brief, gRNA sites were determined using Benchling (settings: single guide of 20 nt length and an NGG PAM site (SpCas9, 3' side)). For both Nab2-ΔQ and Nab2-ΔQ-ΔRGG^{SV40NLS} the cut site lies within the Q-rich region (gRNA regions amplified off pML104 (Laughery et al., 2015): (A) 'upstream' gRNA: GTCCCTAGTTGTGGCTGAAGgttttagagctagaaatagcaagttaaaataaggctagtccg + cggaaacgctatacctccacc; (B) 'downstream' gRNA: gcgaggaagcggaagagcg + ctagctctaaaacCTTCAGCCACAACTAGGGACgatcatttatctttcactgcggagaag; KWY9296 was transformed with stitched gRNA regions, SwaI-digested pML104 and PCR amplification products with homology to the target site (ΔQ: amplified homology product from genomic DNA excluding DNA region coding for aa104-169; -ΔQ-ΔRGG^{SV40NLS} amplified homology product from plasmid pKW4774 where DNA region coding for aa104-263 was replaced by SV40NLS sequence (AAAGGCCTAAGAAGAAGAGGAAAGTATTA)).

Yeast Cell Culturing

Cells were cultured in synthetic complete media containing 2% glucose (Formedium, CSC0205) (for microscopy, RNA FISH and nuclear extraction assay). Cells were inoculated from saturated overnight cultures into fresh medium and grown at 25°C to OD$_{600}$ 0.6–0.8. For acute glucose withdrawal, cells were then washed 3 times in synthetic complete media without glucose, resuspended in the same medium and cultured for the indicated time (30 min or 24 hours).

Acute degron depletion was performed by addition of auxin (500 µM indole-3-acetic acid (IAA); 500 mM stock in ethanol) for the indicated time points (typically 1 or 2 hours). As a solvent control, cells were treated with ethanol.

Single-molecule fluorescence *in situ* hybridization (smFISH)

Probe design: DNA probes coupled to CAL Fluor Red 590 (Stellaris Biosearch Technologies, synthesized by BioCat) were used for smFISH, targeting gene-specific moieties for single molecule visualization. *dT30*

DNA probes coupled to TAMRA (synthesized by Microsynth AG) targeting poly(A) stretches were used for poly(A) RNA visualization.

Methodology: Samples were processed as described in (Derrer et al., 2019). Briefly, cultures were fixed for 30 min with 4% paraformaldehyde (Electron Microscopy Sciences, 15714). Cells were washed once with buffer B (1.2 M sorbitol and 100 mM KHPO4, pH 7.5, at 4°C), then spheroplasted for 20 min using 1% 20T zymolyase in 1.2 M sorbitol, 100 mM KHPO4, pH 7.5, 20 mM vanadyl ribonuclease complex (New England Biolabs, S1402S), and 20 µM β-mercaptoethanol. To stop the spheroplasting reaction, cells were washed once with buffer B, and then transferred into pre-hybridization buffer (10% formamide (Merck Millipore, S4117), 2x SSC (saline-sodium citrate buffer; Life Technologies, AM9624). Separately, 0.5 µL of probe mix (stock 25 µM) was mixed with 2 µL of salmon-sperm DNA (10 mg/ml; Life Technologies, 15632-011) and 2 µL yeast transfer RNA (10mg/ml; Life Technologies, AM7119). The probe mix was denatured in 50 µL of hybridization buffer F (20% formamide and 10mMNaHPO4, pH7.0) for 3 min at 95 °C, and then mixed with 50 µL hybridization buffer H (4x SSC, 4 mg/ml acetylated BSA (Sigma Fluka Aldrich, B8894-1.5mL), 20 mM vanadyl ribonuclease complex). This final hybridization mixture was added to the pre-hybridized cell pellet (which corresponds to approx. 3 OD_{600}) and incubated overnight at 37 °C. The next day, cells were washed in 1x PBS/DAPI and incubated for 3 min, washed again in 1x PBS and then stored at 4 °C.

Microscopy for live cell imaging and smFISH

For live cell imaging and NuRIM fluorescence intensity quantification experiments, cells were transferred to a 384 well plate (Brooks, Matrical MGB101-1-2-LG-L) treated with Concanavalin A (stock solution of 1 mg/ml, air-dried), and spun down for 1 min at 100 rcf.

For smFISH, approx. 1 OD_{600} of fixed cells were spread onto a glass slide using a 12mm cover slip (Huber, 10.0360.52) to obtain a single layer of cells prior to imaging. Alternatively, fixed cells were transferred to a 384 well plate (Brooks, Matrical MGB101-1-2-LG-L) treated with Concanavalin A (stock solution of 1 mg/ml, air-dried), and spun down for 1 min at 100 rcf, and washed twice with 1x PBS to obtain a single layer of cells.

All imaging experiments were performed at 25 °C. Microscopy was performed using an inverted epi-fluoresence microscope (Nikon Ti) equipped with a Spectra X LED light source and a Hamamatsu Flash 4.0 sCMOS camera using a 100X Plan-Apo objective NA 1.4 and NIS Elements software. For live cell imaging and smFISH image analysis, images were processed using FIJI software and plots were generated in GraphPad Prism 8 (GraphPad). For NuRIM quantification, images were processed using NuRIM code (Rajoo et al., 2018; Vallotton et al., 2019) in MATLAB. Briefly, this method uses the HDEL-DsRed signal to generate a mask that segments the nuclear envelope. This mask is then transferred to the GFP channel to extract average intensity information from this mask.

Fluorescence recovery after photobleaching (FRAP)

A) Yeast cells:

200 µL of a culture with OD 0.6-0.8 was added to a Concanavalin A treated imaging well (IBIDI, µ-Slide 8-well, 80827), and spun down for 1 min at 100 rcf. Microscopy was performed on a Leica TCS SP8 STED/FLIM/FCS or a Leica TCS SP8-AOBS microscope using a 63x 1.4NA Oil HC PL APO CS2 objective. A bidirectional scanner at a speed of 8,000 Hz, NF488/561/633, an Airy unit of 1.5, and a FRAP booster for bleaching were applied for every FRAP experiment using a PMT detector (500–551 nm). An image size of 504 x 50 pixel and a zoom of 4 were used together with line accumulation of three, yielding a frame rate of 74.07 frames/s with a pixel dwell time of 73 ns. 50 prebleach and 300 postbleach frames were acquired. A 488-nm argon laser line was used at 30%

power. An elliptical region covering approximately one half of the cell nucleus was bleached. The bleaching was performed using 80% argon laser power for 195 ms. Imaging was conducted with 1.5% laser intensity (with a gain of 570 for Nab2-GFP and GFP-Yra1, and 560 for Npl3-GFP) to illuminate the GFP during prebleach and postbleach acquisition.

B) In vitro droplets:

Protein droplets were set up according to (Hondele et al., 2019). In brief, reactions were pipetted in 384-well microscopy plates (Brooks, Matrical MGB101-1-2-LG-L). 3 μL recombinant GFP-tagged protein (stock 150 μM to a final concentration of 20 μM) was mixed at room temperature with 20 μL of a master mix (14.5 μL LSB-100 (100 mM KCl, 30 mM HEPES-KOH pH 7.4, 2 mM $MgCl_2$), 0.9 μL CRM reconstitution mix (40 mM ATP, 40 mM $MgCl_2$, 200 mM creatine phosphate, 70 U ml−1 creatine kinase), 1.8 μL ATP/MgCl2 (100 mM), 0.9 μL BSA (10 mg mL−1), 0.9 μL poly(A) RNA (1 mg mL-1), 0.9 μL 1M HEPES pH 6.8, 0.1 μL RNasin Plus (Promega). Microscopy was performed on a Leica TCS SP8 STED/FLIM/FCS or a Leica TCS SP8-AOBS microscope using a 63x 1.4NA Oil HC PL APO CS2 objective. A bidirectional scanner at a speed of 8,000 Hz, NF488/561/633, an Airy unit of 1.5, and a FRAP booster for bleaching were applied for every FRAP experiment using a PMT detector (500–551 nm). An image size of 504 x 50 pixel and a zoom of 4 were used together with line accumulation of three, yielding a frame rate of 74.07 frames/s with a pixel dwell time of 73 ns. 50 prebleach and 800 postbleach frames were acquired. A 488-nm argon laser line was used at 30% power. Imaging was conducted with 1.5% laser intensity (with a gain of 570) to illuminate the GFP during prebleach and postbleach acquisition. The bleach was performed using 80% argon laser power for 195 ms. The bleaching was performed by defining an elliptical region covering approximately one half of the droplet.

C) Image analysis: 1) yeast cells: Fluorescence recovery in the bleached region was evaluated by subtracting the extracellular background from the intensity of the bleached region, and the values were then bleach-corrected by normalizing for total cell intensity (Bancaud et al., 2010). 2) droplets: Fluorescence recovery in the bleached region was evaluated by subtracting the background signal outside the droplet from the intensity of the bleached region, and the values were then bleach-corrected by normalizing for total droplet intensity. All curves were fitted with an exponential function in a custom-written MATLAB script. Plots were generated in GraphPad Prism 8 (GraphPad).

Nucleus extraction

The protocol was adapted from (Aris and Blobel, 1991) and (Niepel et al., 2017). Cells were inoculated in a 10 mL pre-culture of synthetic complete media containing 2% glucose (SCD, Formedium, CSC0205), and cultured for 8-10 hrs at 25 °C. Cells were then back-diluted in 500 mL SCD to OD_{600} 0.006, grown for 16-18 hrs at 25 °C to OD_{600} 0.8 and harvested onto a cellulose ester membrane filter (Whatman 10400921) using a vacuum filtration pump. Cells were then washed from the membrane with 40 mL ddH$_2$O into a 50 mL Falcon tube, spun down for 2 min at 845 rcf, resuspended in 25 mL buffer 1 (0.1M Tris-HCl pH7.4, 10mM DTT) and incubated shaking at 30 °C for 20 min in a horizontal shaker at 150 rpm. Cells were then spun down for 2 min at 845 rcf, washed in 10 mL buffer 2 (20mM KPi pH 7.4, 1.2M Sorbitol), and spun down again for 2 min at 845 rcf. For cell wall digestion (spheroplasting), cells were resuspended in 50 mL buffer 3 (20mM KPi pH 7.4, 1.2M Sorbitol, 10 mM ribonucleoside vanadyl complex (NEB S1402S, pre-warmed at 65 °C), 0.08mg/ml Zymolyase-20T (Amsbio 326921)) and incubated for 60 min at 30 °C in a rotation wheel. Spheroplasts were spun at 3000 rcf for 2 min, washed in 10 mL buffer 4 (20mM KPi pH 6.4, 1.2M Sorbitol), and spun again at 3000 rcf for 2 min. To release nuclei, spheroplasts were carefully resuspended in 20 mL buffer F (20% Ficoll, 20mM KPi pH 6.4, 5 mM

RVC, 0.005mg/ml Pepstatin A, 1x protease inhibitor cocktail (PIC, Sigma P8215-5mL), 250 µg/mL DAPI) and put on ice. To mechanically assist nuclei release, the lysate was then processed with 5 slow strokes in a dounce homogenizer on ice (this step is not essential as most of the nuclei are already released upon resuspension in buffer F). To separate nuclei from remaining cells/spheroplasts, the lysate was spun for 5 min at 3460 rcf at 4 °C. Afterwards, the supernatant was carefully removed with a 25 mL pipet until ca. 2 mL was left (to avoid pipetting the cell/spheroplast pellet) and transferred to a fresh 15 mL Falcon tube on ice. For nucleus enrichment and removal of other organelles, a sucrose gradient was prepared as follows: ice-cold 5 mL 2.5 M sucrose was added to the bottom of an open-top plastic tube (Beckman-Coulter 344058), carefully topped up with ice-cold 5 mL 1.875 M sucrose, which in turn was carefully topped up with ice-cold 5 mL 1.7 M sucrose. Lastly, the lysate supernatant was carefully added on top of the 1.7 M sucrose layer, and the full tube was transferred to a pre-cooled metal holder of a SW32Ti ultracentrifuge rotor (Beckman-Coulter). Tubes were spun for 30 min at 103.000 rcf at 4 °C, which resulted in 'cloudy' layer formation between the 1.7 M and 1.875 M sucrose interface (containing mostly cytoplasmic organelles, fraction 1) and between the 1.875 M and 2.5 M sucrose layer (containing nuclei, fraction 2). The enriched nucleus fraction 2 (typically around 4 mL) was then harvested by carefully pipetting away the top layers. To reduce the sucrose molarity, 4 mL of ice-cold buffer 5 (20mM KHPO4 pH 6.45) was added to the nucleus fraction 2, mixed and spun for 4 min at 3460 rcf at 4 °C. The supernatant was carefully removed with a 10 mL pipet until ca. 2 mL was left to separate any remaining cell debris from the nuclear fraction. 2x500 µL aliquots were taken for DNA content determination as a proxy for the amount of harvested nuclei. Enriched nuclei were then immediately processed for RNA extraction (see below).

RNA extraction from nuclear sucrose gradient fraction
The enriched nucleus fraction was transferred to a fresh open-top plastic tube (Beckman-Coulter 344058) and spun for 30 min at 20.000 rcf for 30 min in SW32Ti rotor. The pellet was resuspended in 500 µL RNase-free TES buffer (10 mM TrisHCl pH 7.5, 10 mM EDTA, 0.5% SDS) and transferred to screw-cap tubes. 500 µL acid-saturated phenol was added, the mixture was vortexed and incubated for 30 min in a thermo block at 65 °C (shaking at 150 rpm). Afterwards, the tube was put on ice for 5 min and then spun for 10 min at 13.000 rcf at 4 °C. The upper aqueous phase was transferred to a fresh RNase-free tube, mixed with 500 µL acid-saturated phenol, vortexed and spun for 5 min at 13.000 rcf at 4 °C. The upper aqueous phase was transferred to a chloroform-resistant screw-cap tube, mixed with 500 µL chloroform, vortexed and spun for 5 min at 13.000 rcf at 4 °C. The upper aqueous phase was transferred to a fresh RNase-free tube, mixed with 60 µL 3M NaOAc pH 5.2 and 700 µL isopropanol, and incubated overnight at -20 °C. The next morning, the mixture was spun for 40 min at 13.000 rcf at 4 °C, the supernatant was decanted and 300 µL 75% EtOH was added, followed by centrifugation for 5 min at 13.000 rcf at 4 °C. The supernatant was removed and the pellet was air-dried for 10 min on ice, resuspended in nuclease-free H2O and stored at -80 °C.

RNA extraction from whole cell lysates
Cells were inoculated in a 10 mL pre-culture of synthetic complete media containing 2% glucose (SCD, Formedium, CSC0205), and cultured for 8-10 hrs at 25 °C. Cells were then back-diluted in 400 mL SCD to OD_{600} 0.006, grown for 16-18 hrs at 25 °C to OD_{600} 0.8 and harvested onto a cellulose ester membrane filter (Whatman 10400921) using a vacuum filtration pump. Cells were scraped off the membrane with a metal rod and snap-frozen in a 50 mL Falcon tube containing liquid nitrogen. To lyse the cells, the cell pellet was resuspended in 400 uL TES buffer (10mM TrisHCl pH7.5, 10mM EDTA, 0.5% SDS), followed by addition of 400 uL acid-saturated phenol. The samples were incubated at 65°C for 1 hour

with constant vortexing. Afterwards, the tube was put on ice for 5 min and then spun for 10 min at 13.000 rcf at 4 °C. The upper aqueous phase was transferred to a fresh RNase-free tube, mixed with 500 µL acid-saturated phenol, vortexed and spun for 5 min at 13.000 rcf at 4 °C. The upper aqueous phase was transferred to a chloroform-resistant screw-cap tube, mixed with 500 µL chloroform, vortexed and spun for 5 min at 13.000 rcf at 4 °C. The upper aqueous phase was transferred to a fresh RNase-free tube, and the RNA was precipitated by adding 50uL 3M NaOAc pH5.2 and 500 uL isopropanol to 500 uL sample, and incubated at -20 °C for at least 2 hours. The next morning, the mixture was spun for 30 min at 13.000 rcf at 4 °C, the supernatant was decanted and washed with 300 µL 75% EtOH, followed by centrifugation for 5 min at 13.000 rcf at 4 °C. The RNA pellet was eventually resuspended in nuclease-free water.

RNA seq library preparation and sequencing
For polyA selection, 150 ug of total RNA in 200 uL binding buffer (10 mM Tris-HCl pH 7.5, 0.5 M LiCl, 6.7 mM EDTA) was first denatured at 80°C for 2 min, and immediately incubated on ice. 150 µL of resuspended oligo-dT beads were first washed twice with binding buffer, then mixed with the total RNA samples, and incubated for 5 min at room temperature. The beads were then washed twice with washing buffer (10 mM Tris-HCl pH 7.5, 0.15 M LiCl, 1 mM EDTA), then resuspended in 40 µL 10 mM Tris pH 7 followed by incubation at 80°C for 2 min to elute the RNA. The polyA+ RNA was used to generate strand-specific sequencing libraries using the NEXTflex Rapid Directional Illumina RNA-Seq Library Prep Kit [BioO] according to the manufacturer's instructions. The libraries were sequenced on an Nextseq2000 sequencer. Reads were adaptor trimmed using cutadapt, reads were mapped using tophat2 after rRNA reads subtraction using botwtie2, and quantified using featurecount.
For nuclear extracts, spike-in reads were summed per sample, and sample normalization within a replicate was performed by calculating the spike-in ratios between Nab2-GFP (set to '1'), Nab2-GFP:Nab2$^{\text{ΔQ-ΔRGG-SV40NLS}}$-GFP and Nab2-GFP:Nab2$^{\text{F450A}}$-GFP. Spike-in correction of the CPM values was then performed by multiplying the spike-in ratio value of each sample with the CPM value of each gene.

Protein expression and purification
All recombinantly expressed Nab2 proteins were amplified from yeast genomic DNA of the yeast strains used in this study and cloned into pETMCN-based expression vector with a cleavable N-terminal 6×His-TwinStrep tag plus a C-terminal mGFPA206K tags. Plasmids are listed in Supplementary Table 3. The plasmids were transformed into chemically competent E. coli BL21* under the selection of 0.1 mg/mL ampicillin. Pre-cultures were grown in LB at 37 °C overnight, and diluted 1:100 into rich medium the next morning. Cells were grown at 37 °C to an OD$_{600}$ of 0.6 and induced with final 200 mM IPTG. Cells were then grown overnight at 18 °C, collected and resuspended in 30 ml lysis buffer (500 mM NaCl, 25 mM Tris-HCl pH 7.5, 10 mM imidazole, 10% glycerol, protease inhibitors) per cell pellet from 2L of culture. After cell lysis by EmulsiFlex (Avestin), the 6×His-tagged proteins were affinity extracted with Ni2+ sepharose in small columns, supplemented with prescission (3C) protease for cleavage of the 6xHis-TwinStrep tag, and dialysed into storage buffer (200 mM NaCl, 25 mM Tris-HCl pH 7.5, 10 mM imidazole, 5% glycerol). The next day, protein solutions were further concentrated and purified by size exclusion with a Superdex 200 column on an AEKTA purifier (GE Life Sciences) in the final storage buffer (200 mM NaCl, 25 mM Tris-HCl pH 7.5, 10 mM imidazole, 5% glycerol). Protein expression levels, His eluates and gel filtration fractions were analysed by SDS–PAGE. Clean Superdex elution fractions were pooled, concentrated to approximately 500 µM using Millipore Amicon Centrifugation units, snap frozen as 20 µL aliquots in siliconized tubes in liquid nitrogen and stored at −80 °C. Dbp5ΔN constructs were purified according to the same protocol.

***In vitro* droplet formation assay**

A) Nab2-GFP wild-type and mutant experimental setup:

Protein droplets were set up according to (Hondele et al., 2019). In brief, reactions were pipetted in 384-well microscopy plates (Brooks, Matrical MGB101-1-2-LG-L). 2 μL recombinant GFP-tagged protein (stock 195 μM to a final concentration of 23 μM) was mixed at room temperature with 15 μL of a master mix (10.875 μL low-salt buffer (LSB; variable: LSB-100 (100 mM KCl, 30 mM HEPES-KOH pH 7.4, 2 mM MgCl$_2$) or LSB-150 (150 mM KCl, 30 mM HEPES-KOH pH 7.4, 2 mM MgCl$_2$)), 0.675 μL CKM reconstitution mix (40 mM ATP, 40 mM MgCl$_2$, 200 mM creatine phosphate, 70 U ml−1 creatine kinase), 1.35 μL ATP/MgCl2 (100 mM), 0.675 μL BSA (10 mg mL−1), 0.675 μL poly(A) RNA (1 mg mL-1), 0.675 μL 1M HEPES (variable: pH 6.6/6.8/7.0/7.2/7.4), 0.075 μL RNasin Plus (Promega). For final concentration of 11.5 μL, 1 μL Nab2-GFP (from 195 μM stock) was mixed with 16 μL of the aforementioned master mix. For final concentration of 34.5 μL, 3 μL Nab2-GFP (from 195 μM stock) was mixed with 14 μL of the aforementioned master mix.

Droplets were imaged 20-30 min after setup at 25 °C. Microscopy was performed using an inverted epi-fluorescence microscope (Nikon Ti) equipped with a Spectra X LED light source and a Hamamatsu Flash 4.0 sCMOS camera using a 100X Plan-Apo objective NA 1.4 and NIS Elements software. Images were processed using FIJI software.

B) Nab2+Dbp5 experimental setup:

2 μl of a 100 μM Nab2 solution was mixed with a total of 3 μl Dbp5ΔN protein (WT or mutant) mixed with Dbp5 storage buffer (200 mM NaCl, 25 mM Tris pH 7.5, 5 % glycerol) to achieve the final concentration as indicated. The final volume of 5 μl protein solution was mixed with 20 μL master mix (16 μL LSB100 buffer (100 mM KCl, 30 mM HEPES-KOH pH 7.4, 2 mM MgCl$_2$), 1 μL 10 mg/ml BSA, 1 μL 100 mM ATP / 100 mM MgCl$_2$, 1 μl HEPES-KOH pH 6.2, 1 μL poly(A) 1 mg/mL). Reactions were incubated for 10 minutes at room temperature and images recorded as described above.

Acknowledgements

We thank S. Wente and A. Corbett for providing yeast strains and plasmids, ScopeM and J. Kusch for microscopy support, L. Malinovska for experimental assistance and discussions, and the Weis lab for critical comments on the manuscript. This work was supported by an EMBO long-term fellowship to S.H. and M.H. (ALTF 290-2014, EMBOCOFUND2012, GA-2012-600394 to S.H.; ALTF 870-2014 to M.H.). M.H. was supported by a Human Frontier Science Program (HFSP) postdoctoral fellowship (LT000914/2015) and an ETH postdoctoral fellowship (FEL-37-14-2). This work was supported by the Swiss National Science Foundation (SNF 31003A_179275 and CRSII5_193740 to K.W.).

Competing Interest Statement

The authors have declared no competing interest.

Author contributions

Conceptualization, S.H., M.H. and K.W.; Methodology, S.H., M.H., D.G. and K.W.; Investigation, S.H., M.H., C.P.D., D.M., M.Z., R.M.; Formal analysis, S.H., M.Z., A.O., F.U.; Writing – Original Draft, S.H.; Writing – Review & Editing, S.H., M.H., D.M., M.Z., R.M., A.O., F.U. and K.W.; Funding Acquisition, S.H., M.H. and K.W.; Resources, K.W.; Supervision, S.H. and K.W.

References

Adams, R.L., and Wente, S.R. (2020). Dbp5 associates with RNA-bound Mex67 and Nab2 and its localization at the nuclear pore complex is sufficient for mRNP export and cell viability. PLOS Genet. *16*, e1009033.

Aibara, S., Gordon, J.M.B., Riesterer, A.S., McLaughlin, S.H., and Stewart, M. (2017). Structural basis for the dimerization of Nab2 generated by RNA binding provides insight into its contribution to both poly(A) tail length determination and transcript compaction in Saccharomyces cerevisiae. Nucleic Acids Res. *45*, 1529–1538.

Alcázar-Román, A.R., Tran, E.J., Guo, S., and Wente, S.R. (2006). Inositol hexakisphosphate and Gle1 activate the DEAD-box protein Dbp5 for nuclear mRNA export. Nat. Cell Biol. *8*, 711–716.

Alpert, T., Straube, K., Oesterreich, F.C., and Neugebauer, K.M. (2020). Widespread Transcriptional Readthrough Caused by Nab2 Depletion Leads to Chimeric Transcripts with Retained Introns. Cell Rep. *33*.

Anderson, J.T., Wilson, S.M., Datar, K.V., and Swanson, M.S. (1993). NAB2: a yeast nuclear polyadenylated RNA-binding protein essential for cell viability. Mol. Cell. Biol. *13*, 2730–2741.

Aris, J.P., and Blobel, G. (1991). Isolation of yeast nuclei. In Methods in Enzymology. *194*, 735–749.

Ashkenazy-Titelman, A., Shav-Tal, Y., and Kehlenbach, R.H. (2020). Into the basket and beyond: the journey of mRNA through the nuclear pore complex. Biochem. J. *477*, 23–44.

Bahar Halpern, K., Caspi, I., Lemze, D., Levy, M., Landen, S., Elinav, E., Ulitsky, I., and Itzkovitz, S. (2015). Nuclear Retention of mRNA in Mammalian Tissues. Cell Rep. *13*, 2653–2662

Banani, S.F., Lee, H.O., Hyman, A.A., and Rosen, M.K. (2017). Biomolecular condensates: organizers of cellular biochemistry. Nat. Rev. Mol. Cell Biol. *18*, 285–298.

Bancaud, A., Huet, S., Rabut, G., and Ellenberg, J. (2010). Fluorescence Perturbation Techniques to Study Mobility and Molecular Dynamics of Proteins in Live Cells: FRAP, Photoactivation, Photoconversion, and FLIP. Cold Spring Harb. Protoc. *2010*, pdb.top90.

De Magistris, P. (2021). The Great Escape: mRNA Export through the Nuclear Pore Complex. Int. J. Mol. Sci. *22*, 11767.

Dechant, R., Saad, S., Ibáñez, A.J., and Peter, M. (2014). Cytosolic pH Regulates Cell Growth through Distinct GTPases, Arf1 and Gtr1, to Promote Ras/PKA and TORC1 Activity. Mol. Cell *55*, 409–421.

Derrer, C.P., Mancini, R., Vallotton, P., Huet, S., Weis, K., and Dultz, E. (2019). The RNA export factor Mex67 functions as a mobile nucleoporin. J. Cell Biol. *218* (*12*), 3967–3976

Fasken, M.B., Corbett, A.H., and Stewart, M. (2019). Structure–function relationships in the Nab2 polyadenosine-RNA binding Zn finger protein family. Protein Sci. *28*, 513–523.

Green, D.M., Johnson, C.P., Hagan, H., and Corbett, A.H. (2003). The C-terminal domain of myosin-like protein 1 (Mlp1p) is a docking site for heterogeneous nuclear ribonucleoproteins that are required for mRNA export. Proc. Natl. Acad. Sci. *100*, 1010–1015.

Hector, R.E., Nykamp, K.R., Dheur, S., Anderson, J.T., Non, P.J., Urbinati, C.R., Wilson, S.M., Minvielle-Sebastia, L., and Swanson, M.S. (2002). Dual requirement for yeast hnRNP Nab2p in mRNA poly(A) tail length control and nuclear export. EMBO J. *21*, 1800–1810.

Herzel, L., Ottoz, D.S.M., Alpert, T., and Neugebauer, K.M. (2017). Splicing and transcription touch base: co-transcriptional spliceosome assembly and function. Nat. Rev. Mol. Cell Biol. *18*, 637–650.

Hilliker, A., Gao, Z., Jankowsky, E., and Parker, R. (2011). The DEAD-Box Protein Ded1 Modulates Translation by the Formation and Resolution of an eIF4F-mRNA Complex. Mol. Cell *43*, 962–972.

Hochberg-Laufer, H., Schwed-Gross, A., Neugebauer, K.M., and Shav-Tal, Y. (2019). Uncoupling of nucleo-cytoplasmic RNA export and localization during stress. Nucleic Acids Res. *47*, 4778–4797.

Hodge, C.A., Colot, H.V., Stafford, P., and Cole, C.N. (1999). Rat8p/Dbp5p is a shuttling transport factor that interacts with Rat7p/Nup159p and Gle1p and suppresses the mRNA export defect of xpo1-1 cells. EMBO J. *18*, 5778–5788.

Hondele, M., Sachdev, R., Heinrich, S., Wang, J., Vallotton, P., Fontoura, B.M.A., and Weis, K. (2019). DEAD-box ATPases are global regulators of phase-separated organelles. Nature. *573 (7772)*, 144-148

Huang, D.W., Sherman, B.T., Tan, Q., Collins, J.R., Alvord, W.G., Roayaei, J., Stephens, R., Baseler, M.W., Lane, H.C., and Lempicki, R.A. (2007). The DAVID Gene Functional Classification Tool: a novel biological module-centric algorithm to functionally analyze large gene lists. Genome Biol. *8*, R183.

Iserman, C., Desroches Altamirano, C., Jegers, C., Friedrich, U., Zarin, T., Fritsch, A.W., Mittasch, M., Domingues, A., Hersemann, L., Jahnel, M., et al. (2020). Condensation of Ded1p Promotes a Translational Switch from Housekeeping to Stress Protein Production. Cell *181*, 818-831.e19.

Izaurralde, E., Lewis, J., McGuigan, C., Jankowska, M., Darzynkiewicz, E., and Mattaj, I.W. (1994). A nuclear cap binding protein complex involved in pre-mRNA splicing. Cell *78*, 657–668.

Izawa, S., Takemura, R., Ikeda, K., Fukuda, K., Wakai, Y., and Inoue, Y. (2005). Characterization of Rat8 localization and mRNA export in Saccharomyces cerevisiae during the brewing of Japanese sake. Appl. Microbiol. Biotechnol. *69*, 86–91.

Joyner, R.P., Tang, J.H., Helenius, J., Dultz, E., Brune, C., Holt, L.J., Huet, S., Müller, D.J., and Weis, K. (2016). A glucose-starvation response regulates the diffusion of macromolecules. ELife *5*, e09376.

Katahira, J. (2012). mRNA export and the TREX complex. Biochim. Biophys. Acta BBA - Gene Regul. Mech. *1819*, 507–513.

Khong, A., Matheny, T., Jain, S., Mitchell, S.F., Wheeler, J.R., and Parker, R. (2017). The Stress Granule Transcriptome Reveals Principles of mRNA Accumulation in Stress Granules. Mol. Cell *68*, 808-820

Kroschwald, S., Maharana, S., Mateju, D., Malinovska, L., Nüske, E., Poser, I., Richter, D., and Alberti, S. (2015). Promiscuous interactions and protein disaggregases determine the material state of stress-inducible RNP granules. ELife *4*, e06807.

Larson, D.R., Zenklusen, D., Wu, B., Chao, J.A., and Singer, R.H. (2011). Real-Time Observation of Transcription Initiation and Elongation on an Endogenous Yeast Gene. Science *332*, 475–478.

Laughery, M.F., Hunter, T., Brown, A., Hoopes, J., Ostbye, T., Shumaker, T., and Wyrick, J.J. (2015). New vectors for simple and streamlined CRISPR-Cas9 genome editing in Saccharomyces cerevisiae. Yeast Chichester Engl. *32*, 711–720.

Li, X., Romero, P., Rani, M., Dunker, A.K., and Obradovic, Z. (1999). Predicting Protein Disorder for N-, C-, and Internal Regions. Genome Inform. Workshop Genome Inform. *10*, 30–40.

Longtine, M.S., Mckenzie III, A., Demarini, D.J., Shah, N.G., Wach, A., Brachat, A., Philippsen, P., and Pringle, J.R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in Saccharomyces cerevisiae. Yeast *14*, 953–961.

Marfatia, K.A., Crafton, E.B., Green, D.M., and Corbett, A.H. (2003). Domain Analysis of the Saccharomyces cerevisiae Heterogeneous Nuclear Ribonucleoprotein, Nab2p dissecting the requirements for Nab2p-facilitated Poly(A) RNA export. J. Biol. Chem. *278*, 6731–6740.

Mauger, O., Lemoine, F., and Scheiffele, P. (2016). Targeted Intron Retention and Excision for Rapid Gene Regulation in Response to Neuronal Activity. Neuron *92*, 1266–1278.

Montpetit, B., Thomsen, N.D., Helmke, K.J., Seeliger, M.A., Berger, J.M., and Weis, K. (2011). A conserved mechanism of DEAD-box ATPase activation by nucleoporins and InsP6 in mRNA export. Nature *472*, 238–242.

Mugler, C.F., Hondele, M., Heinrich, S., Sachdev, R., Vallotton, P., Koek, A.Y., Chan, L.Y., and Weis, K. (2016). ATPase activity of the DEAD-box protein Dhh1 controls processing body formation. ELife *5*, e18746.

Niepel, M., Farr, J.C., Rout, M.P., and Strambio-De-Castillia, C. (2017). Rapid isolation of functionally intact nuclei from the yeast Saccharomyces. BioRxiv 162388.

Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T., and Kanemaki, M. (2009). An auxin-based degron system for the rapid depletion of proteins in nonplant cells. Nat. Methods *6*, 917–922.

Noble, K.N., Tran, E.J., Alcázar-Román, A.R., Hodge, C.A., Cole, C.N., and Wente, S.R. (2011). The Dbp5 cycle at the nuclear pore complex during mRNA export II: nucleotide cycling and mRNP remodeling by Dbp5 are controlled by Nup159 and Gle1. Genes Dev. *25*, 1065–1077.

Orij, R., Postmus, J., Beek, A.T., Brul, S., and Smits, G.J. (2009). In vivo measurement of cytosolic and mitochondrial pH using a pH-sensitive GFP derivative in Saccharomyces cerevisiae reveals a relation between intracellular pH and growth. Microbiology *155*, 268–278.

Pühringer, T., Hohmann, U., Fin, L., Pacheco-Fiallos, B., Schellhaas, U., Brennecke, J., and Plaschka, C. (2020). Structure of the human core transcription-export complex reveals a hub for multivalent interactions. ELife *9*, e61503.

Rajoo, S., Vallotton, P., Onischenko, E., and Weis, K. (2018). Stoichiometry and compositional plasticity of the yeast nuclear pore complex revealed by quantitative fluorescence microscopy. Proc. Natl. Acad. Sci. 201719398.

Rambout, X., and Maquat, L.E. (2020). The nuclear cap-binding complex as choreographer of gene transcription and pre-mRNA processing. Genes Dev. *34*, 1113–1127.

Riback, J.A., Katanski, C.D., Kear-Scott, J.L., Pilipenko, E.V., Rojek, A.E., Sosnick, T.R., and Drummond, D.A. (2017). Stress-Triggered Phase Separation Is an Adaptive, Evolutionarily Tuned Response. Cell *168*, 1028-1040.e19.

Roden, C., and Gladfelter, A.S. (2021). RNA contributions to the form and function of biomolecular condensates. Nat. Rev. Mol. Cell Biol. *22*, 183–195.

Romero, P., Obradovic, Z., and Dunker, A.K. (1997). Sequence Data Analysis for Long Disordered Regions Prediction in the Calcineurin Family. Genome Inform. Workshop Genome Inform. *8*, 110–124.

Schmid, M., Poulsen, M.B., Olszewski, P., Pelechano, V., Saguez, C., Gupta, I., Steinmetz, L.M., Moore, C., and Jensen, T.H. (2012). Rrp6p Controls mRNA Poly(A) Tail Length and Its Decoration with Poly(A) Binding Proteins. Mol. Cell *47*, 267–280.

Schmid, M., Olszewski, P., Pelechano, V., Gupta, I., Steinmetz, L.M., and Jensen, T.H. (2015). The Nuclear PolyA-Binding Protein Nab2p Is Essential for mRNA Production. Cell Rep. *12(1),* 128-139.

Sheth, U., and Parker, R. (2003). Decapping and Decay of Messenger RNA Occur in Cytoplasmic Processing Bodies. Science *300*, 805–808.

Shi, Y., and Manley, J.L. (2015). The end of the message: multiple protein–RNA interactions define the mRNA polyadenylation site. Genes Dev. *29*, 889–897.

Shin, Y., and Brangwynne, C.P. (2017). Liquid phase condensation in cell physiology and disease. Science *357(6357),* eaaf4382.

Singh, G., Pratt, G., Yeo, G.W., and Moore, M.J. (2015). The Clothes Make the mRNA: Past and Present Trends in mRNP Fashion. Annu. Rev. Biochem. *84*, 325–354.

Smith, C., Lari, A., Derrer, C.P., Ouwehand, A., Rossouw, A., Huisman, M., Dange, T., Hopman, M., Joseph, A., Zenklusen, D., et al. (2015). In vivo single-particle imaging of nuclear mRNA export in budding yeast demonstrates an essential role for Mex67p. J. Cell Biol. *211*, 1121–1130.

Snay-Hodge, C.A., Colot, H.V., Goldstein, A.L., and Cole, C.N. (1998). Dbp5p/Rat8p is a yeast nuclear pore-associated DEAD-box protein essential for RNA export. EMBO J. *17*, 2663–2676.

Soniat, M., Sampathkumar, P., Collett, G., Gizzi, A.S., Banu, R.N., Bhosle, R.C., Chamala, S., Chowdhury, S., Fiser, A., Glenn, A.S., et al. (2013). Crystal structure of human Karyopherin β2 bound to the PY-NLS of Saccharomyces cerevisiae Nab2. J. Struct. Funct. Genomics *14*, 31–35.

Soucek, S., Zeng, Y., Bellur, D.L., Bergkessel, M., Morris, K.J., Deng, Q., Duong, D., Seyfried, N.T., Guthrie, C., Staley, J.P., et al. (2016). Evolutionarily Conserved Polyadenosine RNA Binding Protein Nab2 Cooperates with Splicing Machinery To Regulate the Fate of Pre-mRNA. Mol. Cell. Biol. *36*, 2697–2714.

Stade, K., Ford, C.S., Guthrie, C., and Weis, K. (1997). Exportin 1 (Crm1p) Is an Essential Nuclear Export Factor. Cell *90*, 1041–1050.

Sträßer, K., and Hurt, E. (2001). Splicing factor Sub2p is required for nuclear mRNA export through its interaction with Yra1p. Nature *413*, 648–652.

Strässer, K., Masuda, S., Mason, P., Pfannstiel, J., Oppizzi, M., Rodriguez-Navarro, S., Rondón, A.G., Aguilera, A., Struhl, K., Reed, R., et al. (2002). TREX is a conserved complex coupling transcription with messenger RNA export. Nature *417*, 304–308.

Takemura, R., Inoue, Y., and Izawa, S. (2004). Stress response in yeast mRNA export factor: reversible changes in Rat8p localization are caused by ethanol stress but not heat shock. J. Cell Sci. *117*, 4189–4197.

Tauber, D., Tauber, G., Khong, A., Treeck, B.V., Pelletier, J., and Parker, R. (2020). Modulation of RNA Condensation by the DEAD-Box Protein eIF4A. Cell *180(3),* 411-426.

Teixeira, D., and Parker, R. (2007). Analysis of P-Body Assembly in Saccharomyces cerevisiae. Mol. Biol. Cell *18*, 2274–2287.

Tipper, D.J. (1973). Inhibition of Yeast Ribonucleic Acid Polymerases by Thiolutin. J. Bacteriol. *116(1)*, 245-56

Tran, E.J., Zhou, Y., Corbett, A.H., and Wente, S.R. (2007). The DEAD-Box Protein Dbp5 Controls mRNA Export by Triggering Specific RNA:Protein Remodeling Events. Mol. Cell *28*, 850–859.

Tuck, A.C., and Tollervey, D. (2013). A Transcriptome-wide Atlas of RNP Composition Reveals Diverse Classes of mRNAs and lncRNAs. Cell *154*, 996–1009.

Vallotton, P., Rajoo, S., Wojtynek, M., Onischenko, E., Kralt, A., Derrer, C.P., and Weis, K. (2019). Mapping the native organization of the yeast nuclear pore complex using nuclear radial intensity measurements. Proc. Natl. Acad. Sci. *116*, 14606–14613.

Wegener, M., and Müller-McNicoll, M. (2019). View from an mRNP: The Roles of SR Proteins in Assembly, Maturation and Turnover. In: Oeffinger M., Zenklusen D. (eds) The Biology of mRNA: Structure and Function. Advances in Experimental Medicine and Biology, vol 1203. Springer, Cham. https://doi.org/10.1007/978-3-030-31434-7_3.

Weirich, C.S., Erzberger, J.P., Flick, J.S., Berger, J.M., Thorner, J., and Weis, K. (2006). Activation of the DExD/H-box protein Dbp5 by the nuclear-pore protein Gle1 and its coactivator InsP6 is required for mRNA export. Nat. Cell Biol. *8*, 668–676.

Yang, P., Mathieu, C., Kolaitis, R.-M., Zhang, P., Messing, J., Yurtsever, U., Yang, Z., Wu, J., Li, Y., Pan, Q., et al. (2020). G3BP1 Is a Tunable Switch that Triggers Phase Separation to Assemble Stress Granules. Cell *181*, 325-345.e28.

Figure legends

1) **Acute Dbp5 depletion causes nuclear mRNA accumulation and Nab2 condensation**

   A) Schematic representation of the auxin-inducible protein depletion tool and the single RNA visualization in vivo. (upper) Upon addition of Auxin, degron-tagged Dbp5 gets recognized by TIR1 and targeted for proteasomal degradation (Nishimura et al., 2009). (lower) 24 PP7 stem loops (PP7sl) are integrated after the STOP codon of a low-copy number gene of interest. A second locus drives the expression of GFP-tagged PP7 coat protein (PP7CP-GFP) that can bind to PP7sl with high affinity, thereby allowing single RNA visualization (Larson et al., 2011).

   B) Cells expressing GFA1-PP7sl, PP7CP-GFP, the nuclear rim marker Ndc1-mKate and the Dbp5-degron with or without TIR1 are treated for 1 hr with auxin (IAA) before imaging using a custom-built microscope (Smith et al., 2015). Scale bar 5 μm.

   C) Quantification of mean squared displacement (MSD) of PP7-labeled GFA1 mRNA in cells from B). Images were taken every 15 ms. Shown are means ± SD of three biological replicates with ≥265 RNA particles per condition.

   D) FRAP of Nab2-GFP cells expressing the Dbp5-degron with or without TIR1 treated for 2 hr with auxin (IAA). FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥16 cells per replicate are shown.

   E) Half time of recovery retrieved from fitting individual FRAP curves from D) with a single component model. Shown are means ± SEM of three biological replicates. * p=0.028, unpaired t-test.

   F) Representative FRAP images of a single nucleus from D)

   G) FRAP of Nab2-GFP cells expressing either only the Dbp5-degron or both the Dbp5-degron and the Rpb2-degron with or without TIR1 treated for 2 hr with auxin (IAA). FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥15 cells per replicate are shown.

   H) Half time of recovery retrieved from fitting individual FRAP curves from G) with a single component model. Shown are means ± SEM of three biological replicates. * p=0.0271 (–TIR1(+Dbp5) vs. +TIR1(–Dbp5), * p=0.0232 +TIR1(–Dbp5) vs. +TIR1(–Dbp5, –Rbp2), ns p=0.6109 ((–TIR1(+Dbp5) vs. +TIR1(–Dbp5, –Rbp2)), unpaired t-test.

2) **Nab2 phase-separates *in vitro*, which gets reverted by active Dbp5**

   A) Recombinant Nab2-GFP droplets were formed in LSB150 buffer to a final KCl concentration of 94 mM in the presence or absence of 0.04 mg/mL poly(A) RNA and imaged at 25 °C 30 min after setup. Scale bar 10 μm.

   B) Recombinant Nab2-GFP droplets were formed in LSB100 or LSB150 buffers to a final KCl concentration of 63 or 94 mM, respectively, at the indicated pH in the presence of 0.04 mg/mL poly(A) RNA and imaged at 25 °C 30 min after setup. Scale bar 10 μm.

   C) Representative images and phase diagram of recombinant Nab2-GFP droplets, which were formed in LSB150 buffer (to a final KCl concentration of 94 mM) containing the indicated single-nucleoside polymers and imaged at 25 °C 20 min after setup. Scale bar 5 μm.

   D) Recombinant Nab2-GFP droplets were formed without Dbp5 or in the presence of increasing amounts of Dbp5ΔN90 or the ATPase-locked Dbp5-ΔN90-DQAD and imaged at 25 °C 30 min after setup. Nab2 9 μM in LSB100 (final 63 mM KCl), 0.04 mg/mL poly(A) RNA, Dbp5 concentrations 14/27/39 μM. Scale bar 10 μm.

3) **Glucose stress causes reduction of nuclear Nab2 dynamics and bulk nuclear RNA retention**
   A) Cells expressing Dbp5-GFP and dsRed-HDEL were grown in medium + DEX or acutely shifted for 30 min or 24 hrs into medium without DEX. Scale bar 5 μm.
   B) Quantification of Dbp5-GFP intensity at the nuclear envelope by NuRIM in the conditions shown in A). Shown are means ± SEM of three biological replicates. At least 350 cells were analyzed per condition and replicate. Dbp5: ** p=0.0082 (+DEX vs. 30 min –DEX), * p=0.022 (+DEX vs. 24 hrs –DEX), unpaired t-test.
   C) RNA FISH against poly(A) using an oligo(dT)30 probe was performed in fixed cells expressing Nab2-GFP. Samples were taken in +DEX or after acute starvation in medium without DEX for 30 min. Insets show single nucleus (2.16x magnification). White arrows indicate Nab2-GFP focus. Scale bar 5 μm.
   D) RNA FISH against poly(A) RNA using an oligo(dT)30 probe was performed in fixed cells expressing Nab2-GFP. Samples were taken in +DEX or after acute starvation in medium without DEX for 24 hrs. Insets show single nucleus (2.16x magnification). White arrows indicate Nab2-GFP focus. Scale bar 5 μm.
   E) FRAP of Nab2-GFP cells in +DEX or 30 min –DEX. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥15 cells per replicate are shown.
   F) Half time of recovery retrieved from fitting individual FRAP curves from E) with a single component model. Shown are means ± SEM of three biological replicates. * p=0.04, unpaired t-test.
   G) FRAP of Nab2-GFP cells in +DEX or 24 hrs –DEX. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥15 cells per replicate are shown.
   H) Half time of recovery retrieved from fitting individual FRAP curves from G) with a single component model. Shown are means ± SEM of three biological replicates. * p=0.01, unpaired t-test.
   I) Representative images of Nab2-GFP cells in wild type (WT) or Dbp5-$^{\Delta N}$Nup159 ('Dbp5 tether') background in +DEX or 24 hrs –DEX. Scale bar 5 μm.
   J) FRAP of cells in I). FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥15 cells per replicate are shown.
   K) Half time of recovery retrieved from fitting individual FRAP curves from J) with a single component model. Shown are means ± SEM of three biological replicates. ns p=0.4425, * p=0.0425, ** p=0.0067 (wild type 24 hrs –DEX vs. Dbp5-$^{\Delta N}$Nup159 +DEX), ** p=0.0032 (Dbp5-$^{\Delta N}$Nup159 +DEX vs. 24 hrs –DEX), unpaired t-test.

4) **Nab2 dimerization, but not RGG domain function, drives condensation *in vitro***
   A) GFP-tagged recombinant Nab2 wild type or Nab2 mutant droplets were formed in LSB150 buffer to a final KCl concentration of 94 mM at the indicated pH in the presence of 0.04 mg/mL polyA RNA and imaged at 25 °C 30 min after setup. Right side: phase diagram of droplet formation at pH 6.8 with the indicated protein and RNA concentrations. Scale bar 10 μm.
   B) FRAP of 20 μM Nab2-GFP and Nab2$^{\Delta Q\text{-}\Delta RGG\text{-}SV40NLS}$-GFP droplets 30 min after setup with 0.04 mg/mL polyA RNA. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥14 droplets per replicate are shown.
   C) Representative images of GFP-tagged Nab2 wild type and mutant cells in +DEX conditions and after acute shift and incubation for 24 hrs in –DEX. Scale bar 5 μm.

D-F) FRAP of cells shown in C). D) Nab2-GFP, E) Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP, F) Nab2^{F450A}-GFP. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥15 cells per replicate are shown.

G) Half time of recovery retrieved from fitting individual FRAP curves from D-F) with a single component model. Shown are means ± SEM of three biological replicates. * p=0.0155, ** p=0.0082, ns p=0.3694, unpaired t-test.

5) The degree of Nab2 condensation determines the level of nuclear mRNA retention

A) RNA FISH against poly(A) RNA using an oligo(dT)30 probe was performed in fixed cells expressing Nab2-GFP, Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP or Nab2^{F450A}-GFP. Samples were taken in +DEX or after acute starvation in medium without DEX for 24 hrs. Insets show single nucleus (2.16x magnification). Scale bar 5 μm.

B) Quantification of nuclear poly(A) RNA signal intensity in 24 hrs –DEX condition in A). Data is normalized to Nab2-GFP. Shown are means ± SEM of five biological replicates, n>100 cells per replicate. ** p=0.003 (Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP), ** p=0.0023 (Nab2-GFP vs. Nab2^{F450A}-GFP), ** p=0.0011 (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), unpaired t-test.

C) Plot of ratios of nuclear RNA sequencing reads (CPM, counts per million). Single dots represent individual genes. Inset shows most abundant reads (blue dots, >500 CPM in Nab2-GFP). Boxes highlight differentially enriched genes (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP : Nab2-GFP; purple, fold change ratio >1.5) and not differentially enriched genes (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP : Nab2-GFP; grey, fold change ratio <1.2).

D) Gene ontology (GO) term analysis of most abundant reads (blue dots) from C) using DAVID Functional Annotation Clustering tool (Huang et al., 2007).

6) Nab2 condensation confers selective mRNA retention *in vivo* during stress

A) RNA FISH against *BTN2* was performed in fixed cells expressing Nab2-GFP, Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP or Nab2^{F450A}-GFP. Samples were harvested in +DEX or after acute starvation in medium without DEX for 24 hrs. Scale bar 5 μm.

B) Quantification of 24 hrs –DEX condition in A). Shown are means ± SEM of five biological replicates, n>100 cells per replicate. ** p=0.0018 (Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP), ns p=0.2943 (Nab2-GFP vs. Nab2^{F450A}-GFP), ** p=0.0064 (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), unpaired t-test.

C) RNA FISH against *HSP42* was performed in fixed cells expressing Nab2-GFP, Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP or Nab2^{F450A}-GFP. Samples were harvested in +DEX or after acute starvation in medium without DEX for 24 hrs. Scale bar 5 μm.

D) Quantification of 24 hrs –DEX condition in C). Data is normalized to Nab2-GFP. Shown are means ± SEM of three biological replicates, n>100 cells per replicate. * p=0.0249 (Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP), ns p=0.4480 (Nab2-GFP vs. Nab2^{F450A}-GFP), * p=0.0342 (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), unpaired t-test.

E) RNA FISH against *HXT6* was performed in fixed cells expressing Nab2-GFP, Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP or Nab2^{F450A}-GFP. Samples were harvested in +DEX or after acute starvation in medium without DEX for 24 hrs. Scale bar 5 μm.

F) Quantification of 24 hrs –DEX condition in E). Data is normalized to Nab2-GFP. Shown are means ± SEM of five biological replicates, n>100 cells per replicate. ** p=0.0081 (Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP), ns p=0.2777 (Nab2-GFP vs. Nab2^{F450A}-GFP), * p=0.0135 (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), unpaired t-test.

G) RNA FISH against *DED1* was performed in fixed cells expressing Nab2-GFP, Nab2$^{\Delta Q\text{-}\Delta RGG\text{-}SV40NLS}$-GFP or Nab2^{F450A}-GFP. Samples were harvested in +DEX or after acute starvation in medium without DEX for 24 hrs. Scale bar 5 μm.

H) Quantification of 24 hrs –DEX condition in G). Data is normalized to Nab2-GFP. Shown are means ± SEM of three biological replicates, n>100 cells per replicate. ns p=0.8324 (Nab2-GFP vs. Nab2$^{\Delta Q\text{-}\Delta RGG\text{-}SV40NLS}$-GFP), ns p=0.5458 (Nab2-GFP vs. Nab2^{F450A}-GFP), ns p=0.9009 (Nab2$^{\Delta Q\text{-}\Delta RGG\text{-}SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), unpaired t-test.

7) **Proper Nab2 condensation is essential for survival after prolonged stress**

A) Growth recovery assay of cells expressing Nab2-GFP, Nab2$^{\Delta Q\text{-}\Delta RGG\text{-}SV40NLS}$-GFP or Nab2^{F450A}-GFP. Mid-log cultures grown in +DEX or acutely starved in medium without DEX for 24 hrs or 120 hrs were recovered as 1:5 serial dilutions (starting OD$_{600}$ 0.2) on rich medium plates containing glucose at either 25 °C for 2 days or 37 °C for 1 day.

B) Model depicting how nuclear Nab2 condensation blocks export of bulk mRNA and allows selective transport of stress-induced mRNAs.

Supplemental Figures

1) **Acute depletion of Dbp5 does not affect NPC composition or NPC transport function; related to Figure 1**

A) Left: Time lapse imaging of Dbp5-IAA7-GFP strain expressing TIR1 protein after addition of either 500 μM IAA or solvent control (EtOH). Scale bar 2 μm. Right: GFP intensity quantification of >200 cells per condition, green: one-phase decay fit, half-life = 12.3 min.

B) Growth assay on rich medium plates containing 500 μM IAA or solvent control (EtOH). Cells of a Dbp5-IAA7 strain with or without expressing TIR1 protein were spotted as 1:5 serial dilutions (starting OD$_{600}$ 0.2) and incubated at 25 °C for 2 days.

C) Quantification of GFP-tagged nucleoporin (NUP) intensity at the nuclear rim in wild type *DBP5* background (wt), and in the presence (-TIR1) or absence (+TIR1) of the Dbp5 degron (Dbp5-AID) after incubation with 500 μM IAA for 0 min or 120 min. Shown are means ± SEM of three biological replicates, n>400 cells.

D) Mid-log cultures of Dbp5-IAA7 *xpo1-1* cells expressing an NLS-NES-2xGFP reporter and the nuclear rim marker dsRed-HDEL with or without TIR1 were treated for 1 hr with either 500 μM IAA, 200 μM leptomycin B (LMB) or 1 hr 500 μM IAA followed by 1 hr 200 μM LMB before imaging. Scale bar 5 μm.

E) Quantification of cells in B). Plot shows nuclear to cytoplasmic (N:C) GFP intensity ratio of NLS-NES-2xGFP reporter. Shown are means of three biological replicates, n>100 cells per replicate.

F) Time lapse imaging of cells in D). Scale bar 5 μm.

G) Quantification of time lapse imaging in F) shows nuclear to cytoplasmic (N:C) GFP intensity ratio of NLS-NES-2xGFP reporter. Means ± SEM (dotted lines) of three biological replicates, n>100 cells per replicate.

H) RNA FISH against the indicated RNAs was performed in cells expressing Dbp5-IAA7 + Ndc1-GFP as a nuclear rim marker with or without TIR1. Samples were harvested after 2 hrs incubation with 500 μM IAA. Scale bar 5 μm.

2) **Acute depletion of Dbp5 does not alter mobility of Yra1, Npl3 or a nuclear GFP reporter; *in vitro* Nab2-GFP condensates are impaired in the presence of Dbp5, Gle1 and IP6; related to Figure 1 and 2**

   A) FRAP of GFP-Yra1 cells expressing the Dbp5-degron with or without TIR1. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥13 cells per replicate are shown.

   B) Half time of recovery retrieved from fitting individual FRAP curves from A) with a single component model. Shown are means ± SEM of three biological replicates. ns p=0.4818, unpaired t-test.

   C) FRAP of Npl3-GFP cells expressing the Dbp5-degron with or without TIR1. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥12 cells per replicate are shown.

   D) Half time of recovery retrieved from fitting individual FRAP curves from C) with a single component model. Shown are means ± SEM of three biological replicates. ns p=0.7217, unpaired t-test.

   E) FRAP of NLS-GFP(3x) cells expressing the Dbp5-degron with or without TIR1. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥12 cells per replicate are shown.

   F) Half time of recovery retrieved from fitting individual FRAP curves from E) with a single component model. Shown are means ± SEM of three biological replicates. ns p=0.2642, unpaired t-test.

   G) RNA FISH against poly(A) RNA using an oligo(dT)30 probe was performed in cells expressing Dbp5-IAA7, or Dbp5-IAA7 and Rpb2-IAA7, with or without TIR1 protein. Samples were harvested after 2 hrs incubation with 500 μM IAA. Scale bar 5 μm.

   H) Domain architecture of Nab2 (top), together with PONDR prediction of naturally disordered regions (bottom) (VL-XT predictor, trained on *V*ariously characterized *L*ong disordered regions and two trained on *X*-ray characterized *T*erminal disordered regions). Black bars indicate predicted disordered regions.

   I) Concentration determination of nuclear and cytoplasmic Nab2-GFP relative to Dhh1-GFP (approx. cellular concentration of 2 μM) as reference for *in vitro* condensate setup. Shown are means ± SEM of three biological replicates.

   J) Representative images depicting the nuclear and cytoplasmic compartment that was measured, respectively, and the calculated intensity ratios of Nab2-GFP vs. Dhh1-GFP, suggesting that Nab2-GFP has a nuclear concentration of approx. 24 μM. Scale bar 5 μm.

   K) Recombinant Nab2-GFP droplets were formed in LSB100 buffer to a final KCl concentration of 63 mM in the presence or absence of 0.04 mg/mL polyA RNA. Time lapse imaging at 25 °C was started 60 min after setup upon addition of water or 5% 1,6-hexanediol. Scale bar 5 μm.

   L) Quantification of K). Shown are means ± SD of GFP droplet intensity of two biological replicates.

   M) Recombinant Nab2-GFP droplets were formed without or in the presence of the indicated proteins and imaged at 25 °C 20 min after setup. Nab2 8 μM in LSB100 pH 6.2, 0.04 mg/mL polyA RNA, 12 μM Gle1, 40 μM IP$_6$, Dbp5-mCherry concentrations: 24/48 μM. Scale bar 10 μm.

3) **Glucose stress does not affect Gle1 and Nup159 localization, or FRAP recovery kinetics of Yra1 and Npl3; related to Figure 3**

A) Cells expressing Dbp5-GFP, Gle1-GFP or Nup159-GFP, and dsRed-HDEL were grown in medium + DEX or acutely shifted for 30 min or 24 hrs into medium without DEX. Scale bar 5 μm. Dbp5-GFP data also shown in Fig. 3A.

B) Quantification of Dbp5-GFP, Gle1-GFP or Nup159-GFP intensity at the nuclear envelope by NuRIM in the conditions shown in A). Shown are means ± SEM of three biological replicates. At least 350 cells were analyzed per condition and replicate. Dbp5-GFP data also shown in Fig. 3A. Dbp5: ** p=0.0082 (+DEX vs. 30 min –DEX), * p=0.022 (+DEX vs. 24 hrs –DEX); Gle1: n.s. p=0.4342 (+DEX vs. 30 min –DEX), * p=0.0368 (+DEX vs. 24 hrs –DEX); Nup159: n.s. p=0.2453 (+DEX vs. 30 min –DEX), * p=0.0499 (+DEX vs. 24 hrs –DEX); unpaired t-test.

C) Representative images of GFP-Yra1 cells in +DEX or 24 hrs –DEX. Scale bar 5 μm.

D) Representative images of Npl3-GFP cells in +DEX or 24 hrs –DEX. Scale bar 5 μm.

E) FRAP of GFP-Yra1 cells in +DEX or 24 hrs –DEX. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥15 cells per replicate are shown.

F) Half time of recovery retrieved from fitting individual FRAP curves from E) with a single component model. Shown are means ± SEM of three biological replicates. ns p=0.154, unpaired t-test.

G) FRAP of Npl3-GFP cells in +DEX or 24 hrs –DEX. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥14 cells per replicate are shown.

H) Half time of recovery retrieved from fitting individual FRAP curves from G) with a single component model. Shown are means ± SEM of three biological replicates. ns p=0.679, unpaired t-test.

I) Representative images of wild type (Dbp5-GFP) or GFP-Dbp5-$^{\Delta N}$Nup159 ('GFP-Dbp5 tether') cells in +DEX or 24 hrs –DEX. Scale bar 5 μm.

4) Nab2-GFP focus formation in glucose stress is RNA-dependent, reversible, and mildly affected upon addition of 1,6-hexanediol *in vivo*; related to Figure 3

A-C) Time lapse imaging of Nab2-GFP cells that were treated with water or 5% 1,6-hexanediol in A) +DEX, B) 30 min –DEX or C) 24 hrs –DEX. Scale bar 5 μm.

D) Time lapse imaging of Nab2-GFP cells after release from 24 hrs Dextrose starvation into +DEX medium. Scale bar 5 μm.

E) FRAP of Nab2-GFP cells in +DEX, 30 min –DEX or pre-treated for 15 min with 3 μg/mL thiolutin before starving cells for 30 min in –DEX. FRAP curves are normalized to prebleach and postbleach values. Mean curves ± SEM of three biological replicates with ≥14 cells per replicate are shown.

F) Half time of recovery retrieved from fitting individual FRAP curves from E) with a single component model. Shown are means ± SEM of three biological replicates. * p=0.0454, ns p=0.1437, unpaired t-test.

G) RNA FISH against poly(A) RNA using an oligo(dT)30 probe was performed in fixed cells expressing Cbc1-GFP or Cbc2-GFP. Samples were harvested in +DEX or after acute starvation in medium without DEX for 24 hrs. Scale bar 5 μm.

5) Characterization of Nab2 mutants *in vivo* and *in vitro*, and nucleus isolation efficiency; related to Figure 4 and 5

A) GFP-tagged recombinant Nab2$^{\Delta RGG-SV40NLS}$ or Nab2$^{\Delta Q}$ mutant droplets were formed in LSB150 buffer at the indicated pH in the presence of 0.04 mg/mL polyA RNA and imaged at 25 °C 30 min after setup. Scale bar 10 μm.

B) Growth assay on rich medium plates containing 500 μM IAA or solvent control (EtOH). The indicated strains were spotted as 1:5 serial dilutions (starting OD$_{600}$ 0.2) and incubated at 25 °C for 2 days.

C-D) Growth assay on rich medium plates. The indicated strains were spotted as 1:5 serial dilutions (starting OD$_{600}$ 0.2) and incubated at 25 °C for 2 days.

E) Representative images of Nab2$^{\Delta Q}$-GFP cells in +DEX or 24 hrs –DEX. Scale bar 5 μm.

F) FRAP of Nab2$^{\Delta Q}$-GFP cells in +DEX or 24 hrs –DEX. FRAP curves are normalized to prebleach and postbleach values. Mean curves + SEM of three biological replicates with ≥14 cells per replicate are shown.

G) Half time of recovery retrieved from fitting individual FRAP curves from F) with a single component model. Shown are means ± SEM of three biological replicates. * p=0.0132, unpaired t-test.

H) Quantification of the ratio of mean nuclear poly(A) RNA signal intensity versus mean cellular poly(A) RNA signal intensity in +DEX and 24 hrs –DEX from RNA FISH experiment shown in Figure 5A. Shown are means ± SEM of five biological replicates, n>100 cells per replicate. +DEX: * p=0.0168 (Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP), n.s. p=0.3346 (Nab2-GFP vs. Nab2^{F450A}-GFP), n.s. p=0.0557 (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), –DEX: * p=0.0398 (Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP), n.s. p=0.2159 (Nab2-GFP vs. Nab2^{F450A}-GFP), * p=0.0161 (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), unpaired t-test.

I) Quantification of the ratio of cellular (left) and nuclear (right) poly(A) RNA signal intensity comparing +DEX and 24 hrs –DEX from RNA FISH experiment shown in Figure 5A. Shown are means ± SEM of five biological replicates, n>100 cells per replicate. Cellular ratios: n.s. p=0.3076 (Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP), n.s. p=0.3645 (Nab2-GFP vs. Nab2^{F450A}-GFP), n.s. p=0.0733 (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), Nuclear ratios: ** p=0.008 (Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP), n.s. p=0.1616 (Nab2-GFP vs. Nab2^{F450A}-GFP), ** p=0.0023 (Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP vs. Nab2^{F450A}-GFP), unpaired t-test.

J) RNA FISH against poly(A) RNA using an oligo(dT)30 probe was performed on spheroplasts or extracted nuclei from indicated strains harvested after 24 hrs starvation in –DEX. BF, bright field. Scale bar 5 μm.

K) Quantification of extracted nuclei fraction in J). Shown are means ± SEM of 3 biological replicates.

6) **RNA sequencing analysis of whole cell extracts and nuclear extracts; related to Figure 5**

A) Similarity matrix of individual replicates of whole cell RNA sequencing reads, and hierarchical clustering (one minus Pearson correlation) using Morpheus software (Broad Institute, https://software.broadinstitute.org/morpheus).

B) GO term analysis of 204 genes upregulated in 24 hrs –DEX compared to +DEX (pval< 0.05 and log2 Fold Change> 3) in the indicated strains.

C) Similarity matrix of individual replicates of nuclear RNA sequencing reads, and hierarchical clustering (one minus Pearson correlation) using Morpheus software (Broad Institute).

D) Comparison of nuclear RNAseq reads (CPM, counts per million), left: Nab2-GFP vs. Nab2$^{\Delta Q-\Delta RGG-SV40NLS}$-GFP, right: Nab2-GFP vs. Nab2^{F450A}-GFP. Each dot represents one gene. Grey line represents a 1:1 ratio. Shown are average CPMs of three biological replicates.

E) Distribution of average reads of most abundant genes (>500 CPM in Nab2-GFP, 318 genes) from D). Box-and-whiskers plot, min to max. **** p<0.001, ns p=0.3439, unpaired t-test.

F) Comparison of whole cell RNAseq reads (CPM, counts per million) of differentially enriched vs. not differentially enriched genes of most abundant genes (>500 CPM in Nab2-GFP, 318 genes). Box-and-whiskers plot, 10[th] to 90[th] percentile.

G) Comparison of transcript length (log10) of differentially enriched (149 genes) vs. not differentially enriched (100 genes) vs. all transcripts (5506). Box-and-whiskers plot, 10[th] to 90[th] percentile. **** p<0.001, ns p=0.2563, Mann Whitney test.

H) Comparison of intron-containing transcripts of differentially enriched (3 out of 149 genes) vs. not differentially enriched (14 out of 100 genes) vs. all transcripts (263 out of 5506).

7) Proper Nab2 condensation is essential for survival after glucose stress, both acutely and in stationary phase; related to Figure 7

A) Growth recovery assay of cells expressing Nab2-GFP, Nab2$^{\Delta Q\text{-}\Delta RGG\text{-}SV40NLS}$-GFP or Nab2^{F450A}-GFP. Mid-log cultures grown at 25 °C in +DEX or acutely starved in medium without DEX 120 hrs (5 days) were recovered as 1:5 serial dilutions (starting OD$_{600}$ 0.2) on rich medium plates containing glucose at either 18 °C for 4 days, 25 °C for 2 days, 30 °C for 1 day or 37 °C for 1 day.

B) Growth recovery assay of cells expressing Nab2-GFP, Nab2$^{\Delta Q\text{-}\Delta RGG\text{-}SV40NLS}$-GFP or Nab2^{F450A}-GFP. Cultures in +DEX were grown from mid-log into stationary phase for 1 day at 25 °C and subsequently recovered as 1:5 serial dilutions (starting OD$_{600}$ 0.2) on rich medium plates containing glucose at either 18 °C for 4 days, 25 °C for 2 days, 30 °C for 1 day or 37 °C for 1 day.

Video S1 Cells expressing GFA1-PP7sl, PP7CP-GFP, the nuclear rim marker Ndc1-mKate and the Dbp5-degron without TIR1 are treated for 1 hr with Auxin (IAA) before imaging using a custom-built microscope (Smith et al., 2015). Scale bar 5 μm. related to Figure 1.

Video S2 Cells expressing GFA1-PP7sl, PP7CP-GFP, the nuclear rim marker Ndc1-mKate and the Dbp5-degron with TIR1 are treated for 1 hr with Auxin (IAA) before imaging using a custom-built microscope (Smith et al., 2015). Scale bar 5 μm. related to Figure 1.

Video S3 Recombinant Nab2-GFP droplets were formed in LSB100 buffer to a final KCl concentration of 63 mM in the presence or absence of 0.04 mg/mL poly(A) RNA and imaged every 2 min for 2 hrs at 25 °C directly after setup. Scale bar 10 μm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

	+ DEX	24 hrs – DEX	120 hrs –DEX	
Nab2-GFP				
Nab2$^{ΔQ-ΔRGG-SV40NLS}$-GFP				regrowth at 25 °C
Nab2^{F450A}-GFP				
Nab2-GFP				
Nab2$^{ΔQ-ΔRGG-SV40NLS}$-GFP				regrowth at 37 °C
Nab2^{F450A}-GFP				

B