Effects of global change on animal biodiversity in boreal forest landscape: an assemblage dissimilarity analysis

Ilhem Bouderbala[1,2,4,*] Guillemette Labadie[2], Jean-Michel Béland[3], Yan Boulanger[3], Christian Hébert[3], Patrick Desrosiers[1,4,5], Antoine Allard[1,4], Daniel Fortin[2,4]

August 18, 2022

[1] Département de physique, de génie physique et d'optique, Université Laval, Québec (Qc), Canada G1V 0A6

[2] Département de biologie, Université Laval, Québec (Qc), Canada G1V 0A6

[3] Centre de foresterie des Laurentides, Service canadien des forêts, Québec (Qc), Canada G1V 4C7

[4] Centre interdisciplinaire en modélisation mathématique de l'Université Laval, Québec (Qc), Canada G1V 0A6

[5] Centre de recherche CERVO, Québec (Qc), Canada G1J 2G3

Abstract

Despite an increasing number of studies highlighting the impacts of climate change on boreal species, the main factors that will drive changes in species assemblages remain ambiguous. We quantify two climate-induced pathways based on direct and indirect effects on species occupancy and assemblage dissimilarity under different harvest management scenarios. The direct climate effects illustrate the impact of climate variables while the indirect effects are reflected through the changes in land cover composition. To understand the main causes in assemblage dissimilarity, we analyze the regional and the latitudinal species assemblage dissimilarity by decomposing it into *balanced variation in species occupancy and occurrence* and *occupancy and occurrence gradient*. We develop empirical models to predict the distribution of more than 100 bird and beetle species in the Côte-Nord region of Québec over the next century. Our results show that the two pathways are complementary and alter biodiversity, mainly caused by balanced variation in species occupancy and occurrence. At the regional scale, both effects have an impact on decreasing the number of winning species. Yet, responses are much larger in magnitude under mixed climate effects (a mixture of direct and indirect effects). Regional assemblage dissimilarity reached 0.77

*ilhem.bouderbala.1@ulaval.ca

and 0.69 under mixed effects versus 0.09 and 0.10 under indirect effects for beetles and birds, respectively, between RCP8.5 and baseline climate scenarios when considering harvesting. Therefore, inclusion of climatic variables considers aspects other than just those related to forest landscapes, such as life cycles of animal species. Latitudinally, assemblage dissimilarity increased following the climate conditions pattern. Our analysis contributes to the understanding of how climate change alters biodiversity by reshaping community composition and highlights the importance of climate variables in biodiversity prediction.

Keywords: Biodiversity modeling, climate change, anthropogenic disturbance, assemblage dissimilarity, assemblage dissimilarity decomposition, boreal forests, latitudinal gradient.

1 Introduction

Global climate warming affects the functionality of ecosystems by modifying forest composition and biomass, which in turn has repercussions for biodiversity and species assemblages (Kelly and Goulden, 2008; Hillebrand et al., 2010; Thuiller et al., 2011; Pachauri et al., 2014; Zhang and Liang, 2014). In fact, many studies have anticipated that anthropogenic radiative forcing will alter boreal biodiversity and ecosystems (Pachauri et al., 2014; Tremblay et al., 2018; Cadieux et al., 2020). For example, global scale predictions have shown that the potential high emissions of greenhouse gases would lead mainly to negative effects on biodiversity (Pachauri et al., 2014). By contrast, other results have been published anticipating an increase in biodiversity during this century, which is referred *the northern biodiversity paradox* (Matthews et al., 2004; Morin and Thuiller, 2009; Berteaux et al., 2010). This paradox suggests that the ecological niches of some "southerly" species would increase in size due to expansion beyond the northern periphery of their ranges (Berteaux et al., 2010).

In Canada, temperature has increased by 1.7 °C since 1948, twice as fast as the global average (Bush and Lemmen, 2019). This increase in temperature could lead to the northward migration of thermophilous hardwood tree species to the detriment of boreal conifers, particularly mid-to-late-successional species (Duveneck et al., 2014; Boulanger et al., 2017; Boulanger and Puigdevall, 2021). Moreover, climate change is expected to influence wildfire activity directly (Boulanger and Puigdevall, 2021), which would favour pioneer and fire-adapted boreal tree species (Boulanger et al., 2017). Significant changes in species composition are expected within the transition zone between boreal and temperate biomes, where several tree species are currently reaching their thermal limits (Brice et al., 2020; Boulanger and Puigdevall, 2021). Global warming could

⁴⁴ also drive the occurrence of more extreme climatic events, including severe droughts (Kumar, 2013; Masson-
⁴⁵ Delmotte et al., 2018), thereby reducing the productivity of several boreal tree species through increasing
⁴⁶ metabolic respiration (Girardin et al., 2016). Climate-induced changes in insect outbreak regimes, notably
⁴⁷ those in spruce budworm (SBW) (*Choristoneura fumiferana*) can lead to severe defoliation and death of
⁴⁸ firs (*Abies* spp.) and spruces (*Picea* spp.), and alter forest successional pathways (Pureswaran et al., 2015;
⁴⁹ Labadie et al., 2021).

⁵⁰ Global change might influence biodiversity through behavioral, morphological and physiological changes
⁵¹ of organisms or might even influence gene flow (Naeem et al., 2012; Wisz et al., 2013; Sergio et al., 2018;
⁵² Matuoka et al., 2020; Boulanger and Puigdevall, 2021; Micheletti et al., 2021). The implications of global
⁵³ change on biodiversity could be synthesized through direct and indirect effects. On one hand, the indirect
⁵⁴ effects are associated with habitat change (e.g., vegetation) that could occur through various events such
⁵⁵ as natural and anthropogenic disturbances (Boulanger et al., 2017). On the other hand, direct effects are
⁵⁶ mainly characterized by the influence of climate and meteorological conditions on the organisms (Micheletti
⁵⁷ et al., 2021). Both types of effects should be considered when studying the impacts of global warming on
⁵⁸ ecosystem functioning. For example, ecosystem services can depend on the magnitude of both direct and
⁵⁹ indirect pathways on global biodiversity. Understanding both types of effects should thus help identify the
⁶⁰ best conservation actions that are required to maintain valued services. If habitat-based global change is
⁶¹ projected to decrease species occupancy, land management actions could be adapted by targeting vegetation
⁶² restoration. Otherwise, species translocation actions could be adopted if direct changes are anticipated that
⁶³ would cause a decline in species occupancy (Micheletti et al., 2021).

⁶⁴ Climate and land-use changes may interact in complex ways to impact changes in wildlife habitats (Bentz
⁶⁵ et al., 2010; Tremblay et al., 2018). For example, an increase in disturbance rate due to warmer conditions
⁶⁶ could favor the regeneration of warm-adapted, pioneer tree species (Brice et al., 2020; Boulanger and Puigde-
⁶⁷ vall, 2021), which could affect biodiversity through adjustments in species ecological traits. Moreover, the
⁶⁸ synergistic effects of harvest and natural disturbances with direct climate change could alter species as-
⁶⁹ semblages, and, therefore, biodiversity. In this work, we investigated future climate-induced variations in
⁷⁰ bird and beetle assemblages in Québec's boreal forest under different forest harvesting scenarios. This will
⁷¹ enhance our knowledge regarding the interactive effects of climate change and forest use on future biodiver-
⁷² sity. We assessed effects of future climate conditions on these assemblages by comparing forest landscapes
⁷³ that were simulated under two anthropogenic radiative-forcing scenarios, i.e., Representative Concentration
⁷⁴ Pathway (RCP) 4.5 and RCP 8.5 (van Vuuren et al., 2011) with landscapes that are simulated under av-
⁷⁵ erage historical (baseline) climate conditions. Forest landscapes were simulated through a spatially explicit

raster-based forest landscape model to capture key forest ecosystem processes. We further assessed how

forest management may affect future assemblages by simulating species occupancy if current harvest levels

are maintained vs there would be no harvest activities. We analyze the effects of global change in two ways:

(1) indirectly, i.e., habitat-based change (Wisz et al., 2013; Boulanger and Puigdevall, 2021; Micheletti et al.,

2021) and (2) through a mixed effect combining indirect and direct effects, i.e., those stemming from climate

variables *per se* (Thuiller et al., 2018). The choice of the two pathways was made to quantify separately their

implications on biodiversity in a cascading configuration. More precisely, starting first with habitat effects,

through Habitat-Only-Based Models (HOBMs), and adding afterward meteorological conditions, through

Climate-Habitat-Based Models (CHBMs). We also determined the main drivers of assemblage dissimilarity

following the two pathways.

We opted for a species distribution modeling (SDM) approach (Guisan and Zimmermann, 2000; Guisan

and Thuiller, 2005; Peterson et al., 2011) to model the single-species occurrence probability based on extensive

field surveys of beetles and birds. We analyzed the assemblage structure based on continuous occurrence

probabilities (probability-based) (Guisan and Thuiller, 2005; Grenié et al., 2020) to avoid overprediction

risks (Gelfand et al., 2005; Dubuis et al., 2011; Grenié et al., 2020). It has been previously demonstrated

that probability-based richness provided a better fit of the actual richness than would the threshold-based

approach (Grenié et al., 2020). We computed dissimilarity measures (Baselga, 2010; Albouy et al., 2012;

Baselga, 2013) between different climate scenarios under the two harvesting levels. In this context, we

adapted continuous-based decomposition in the context of occurrence probabilities to detail the two com-

ponents of *Beta-diversity*: *balanced variation in species abundances* and *abundance gradient*, which are

generalizations of turnover and nestedness (Baselga, 2013). Quantifying the main causes of change in biodi-

versity could be very helpful in assessing the potential underlying determinants because species replacement

and nestedness, for example, are two different ways of generating assemblage dissimilarity (Baselga, 2013).

2 Material and methods

2.1 Study area and occurrence data

The study area is located in the Côte-Nord region of Québec, Canada ($48°N$ to $53°N$, $65°W$ to $71°W$),

within an area of 114118 km^2 (Fig. 1). The northern part of the study area belongs to the black spruce-

feather moss bioclimatic domain, and is dominated by black spruce (*Picea mariana* [Mill.] BSP) and balsam

fir (*Abies balsamea* [L.] Miller). Wildfires are the major natural disturbances in northern areas that have

yet to be logged (Boucher et al., 2017). The southern part of the study area belongs to the eastern balsam

fir - white birch (*Betula papyrifera* Marsha.) subdomain, mostly dominated by balsam fir and white spruce

(*Picea glauca* [Moench] Voss) mixed with white (paper) birch. Forest harvesting had been the main source of

forest disturbance since the late 1990s in this latter area (Bouchard and Pothier, 2011). In Québec, logging

affected around 0.8% of public forest annually (Bureau du Forestier en chef, 2010). This part of the territory

also has been affected by an outbreak of the SBW that began in 2006 and which is still ongoing. Tree

mortality began around 2015 and has subsequently increased.

Figure 1: Study area with the presence-absence distribution of the two taxa.

We used presence-absence data that were collected between 2004 and 2018 to model species distributions.

Given that we were mainly focusing on the impacts of fire and harvesting, we wanted to remove sites that

were located in stands heavily damaged by the SBW outbreak (Ministère des Forêts de la Faune et des Parcs,

2018) by using a cumulative index of defoliation severity from 2007 to 2018 (Labadie et al., 2021). Annual

defoliation severity was based on aerial surveys characterizing damage that was incurred by SBW since 2006

(Ministère des Forêts de la Faune et des Parcs, 2018) and was classified between 0 and 3, with 3 indicating

the highest level of defoliation. The cumulative severity of the outbreak was obtained by summing the annual

severity in Labadie et al. (2021). Sites with cumulative severity values of 10 and above were discarded from

analyses. This threshold attests for a strong impact of the SBW over a long period, which modifies stand

structure and therefore affects species presence. Indeed, Labadie et al. (2021) showed (see Fig. 2 in this

paper) that, at this level, the response of shrubs is clearly stronger that of herbaceous plants (no overlap in

the 95% CI), which indicates important changes in forest structure that modify habitat of many species.

We used the data from the Atlas of Breeding Birds (Atlas des oiseaux nicheurs du Québec, 2018), which

were based on species occurrences that were detected using unlimited distance 5-minute point counts (Bibby

et al., 2000), which were collected during the breeding season (late May to mid-July) between 2010 and 2018. For beetles, we merged different databases that had been collected in 2004, 2005, 2007 and 2011 (Janssen et al., 2009; Légaré et al., 2011; Bichet et al., 2016). In addition, we used data from 54 sites that had been sampled in 2018 in the northern part of the territory, along the northeast principal road going to Labrador. The sampling protocols were characterized by one multidirectional flight-interception trap per site, which sampled flying beetles, and four meshed pitfall traps per site, which sampled epigeic beetles during their peak period of activity (June-August) (Janssen et al., 2009; Bichet et al., 2016). For beetles, we used species-level identifications wherever possible; otherwise, we standardized the identification to the genus level (around 92% initial identifications were considered at the species level).

2.2   Predictor variables

To predict species occurrence, we used climate and land cover variables that were grouped in two classes of models: Climate-Habitat-Based (CHBMs) and Habitat-Only-Based (HOBMs). These two model classes were designed to study climate-induced effects, as follows: CHBMs for the mixed effects; and HOBMs for the indirect climate effects. Initially, we generated 22 potential climate variables at a 250-m resolution using the BioSim platform (Régnière et al., 2017), including the annual average of temperature, precipitation and water deficit between 2004 and 2018 (see Table. 1 in Supporting Information for the description of all potential predictor variables). BioSIM simulates daily maximum and minimum temperatures, precipitation, water deficit, mean daily relative humidity and wind speed by matching georeferenced sources of daily weather data to spatially georeferenced points. BioSIM uses spatial regression to adjust weather data for differences in latitude, longitude, and elevation between the sources of weather data and each field location (for more details, see (Boulanger et al., 2018a)). In our case, the spatially referenced points were 15 000 points that were randomly located across the entire Province of Québec, whereas weather data were daily data originating from discrete weather stations that were located within the province. We generated climate variables at a 250 m scale by spatially interpolating data from the 15 000 random points using kriging and elevation as a drifting variable. Land cover maps from the Canadian National Forest Inventory (NFI) were used to generate land cover variables based on k-nearest-neighbor interpolation at a 250-m resolution that was referenced to the year 2001 (Beaudoin et al., 2014). To estimate forest composition, we used the relative importance of tree species groups (conifer and deciduous species), treed land and tree canopy-closure maps from these NFI data to generate five natural land cover classes: closed-canopy conifer forest; open-canopy mature conifer forest; mixed forest; open area; and others (Labadie et al., 2021) (for more details, see Table 2 in Supporting Information). We used four of natural land cover categories together with six disturbance

classes. For the disturbed stands (by fire or harvest), we subdivided them into three age-classes: $[0, 10]$,

$]10, 20]$ and $]20, 50]$ years. In addition, we considered stand age and the distance to the nearest burned area

as potential predictor variables. Stand age maps were based on the year 2001 (Beaudoin et al., 2014) with

an update according to the registered fire and harvest disturbances between 2004 and 2018. Furthermore,

it has been mentioned in Bichet et al. (2016) and Zhao et al. (2013) that the influence of the landscape

varied between 400 m for beetles to 1000 m for birds. Consequently, we used a matrix of 20 pixels centred

on the focal pixel (21 pixels in total) to calculate frequencies of the land cover (see Table.1 and Table. 2 in

Supporting Information for descriptions of the 10 land cover variables that were used in the study).

2.3   Climate and forest management scenarios

We obtained future climate projections from the Canadian Earth System Model version 2 (CanESM2) for

RCP 4.5 and RCP 8.5 for the period 2071-2100, which were further downscaled to a 10-km resolution using

ANUSPLIN (McKenney et al., 2013). Future monthly normals at each random point that was previously

used to assess baseline climate were directly assessed from changes that were observed between the 1981-

2010 period and future projections in the 10-km cell in which the random point was located. Daily time

series were stochastically generated for each random point from these future monthly normals using BioSIM.

Future climate variables at each random point were calculated by averaging these daily values from 30

BioSIM simulations (Boulanger et al., 2018a). Climate scenarios varied depending upon the mean annual

temperature that was expected to increase between 3 °C (RCP4.5) and 7.5 °C (RCP8.5) throughout the

southern boreal region by 2100 (compared to 2000, see Fig. 2), while average precipitation was projected to

increase between 7% and 10% regionally, with relatively small differences among scenarios (Boulanger et al.

(2018b); Boulanger and Puigdevall (2021), see Fig. 2).

The forest landscapes were simulated using the spatially explicit raster-based forest landscape model

LANDIS-II (Scheller and Mladenoff, 2004), which simulates stand- and landscape-scale processes, including

forest succession, seed dispersal and natural (wildfires and spruce budworm outbreaks) and anthropogenic

(harvest) disturbances. This model has been extensively used in Québec over the last decade and has been

thoroughly validated under various forest conditions (Boulanger et al., 2017; Taylor et al., 2017; Tremblay

et al., 2018; Boulanger and Puigdevall, 2021). Forest landscapes were initialized for the year 2000 condi-

tions using the NFI attribute maps (Beaudoin et al., 2014) and provincial sample plots. Tree growth and

regeneration as well as wildfires were climate-sensitive in simulations. We simulated two levels of harvesting

scenarios according to a gradient of harvesting pressure, from no harvesting to high intensity harvesting,

¹⁸⁷ similar to current management practices in Québec (harvest applied to 8% of the harvestable upland area

¹⁸⁸ per 10 years). The Biomass Harvest extension was used to simulate forest harvests (Gustafson et al., 2000).

¹⁸⁹ Only stands that contained tree cohorts that were greater than 60-years-old were allowed to be harvested.

¹⁹⁰ Mean harvested patch size varied between 40 km² and 150 km². Harvest rates were held constant throughout

¹⁹¹ the simulations unless sufficient numbers of stands did not qualify for harvest. In the latter case, harvesting

¹⁹² continued proceed until stands were no longer available. Simulations were performed from 2000 to 2100

¹⁹³ using a 10-yr time step. Climate sensitive parameters for simulations that were performed under RCP 4.5

¹⁹⁴ and RCP 8.5 were set to change in 2011-2020, 2041-2050 and 2071-2080. Forest succession, wildfire, SBW

¹⁹⁵ outbreaks and harvesting were simulated using Biomass Succession v5.0, Base Fire v4.0, BDA v4.0 and the

¹⁹⁶ Biomass Harvest v5.0 extensions, respectively. Many more details regarding LANDIS-II simulations that

¹⁹⁷ were performed in this study can be found in Labadie et al. (2022) as well as in Boulanger et al. (2017) and

Tremblay et al. (2018).

Figure 2: The distribution of the land cover and three climate variables under the simulated management scenarios in 2100. **A-B**, **F-G** and **K-L** represent the distribution of land cover over the six study scenarios that were classified into four large cover classes: *Natural*, which included conifer dense, conifer open, mixed wood and open habitat; *Fire* and *Cut* for the land cover disturbed by fire and Harvest; and *Others* for the rest. The barplots represent the frequency of each class in the map. **D-F**, **J-L** and **P-R** represent the distribution of mean temperature, precipitation, and water deficit for the three climate scenarios **Baseline**, **RCP4.5** and **RCP8.5**, respectively.

¹⁹⁸

2.4   The modelling strategy

We used two classes of models, i.e., Habitat-Only-Based models (HOBMs) and Climate-Habitat-Based models (CHBMs), to study indirect climate effects and mixed climate effects, respectively. The mixed effects represented a combination of direct and indirect effects (see filled arrows in Fig. 3). We divided our modeling strategy into five steps (see Fig. 4). We aimed to study the effect of climate change under different forest harvesting scenarios. Therefore, we compared baseline climate scenario with RCP 4.5 and RCP 8.5 under two forest harvesting scenarios (NoHarvest and Harvest), hence the use of two baseline scenarios (see Step 5 in Fig. 4).

The modelling procedure that is described here was repeated for the two model classes (CHBMs and HOBMs) according to potential predictor variables under each class. We standardized the variables to facilitate model convergence (MacKenzie et al., 2017) prior to model calibration that was based on the corresponding database. Steps 2 and 3 correspond to a cross-validation loop, in which we split the date into 10 folds of relatively equal size, so that at each step, 9 folds were used for training and the one remaining fold was used for testing.

Figure 3: Framework used for the study. The indirect effects were generated through the change in forest composition and the direct effects included the effects of temperature, precipitation and water deficit. Abbreviations: Climate-Habitat-Based Models (CHBMs), Habitat-Only-Based Models (HOBMs).

2.4.1   Step 1: Data pre-processing

We used the following procedure for preparing the two databases: (1) we removed all of the sites that were strongly affected by the spruce budworm outbreak; and (2) we included only the more common species, with

Figure 4: The modeling strategy that was used under Habitat-Only-Based models (HOBMs) and Climate-Habitat-Based models (CHBMs). Step 1 was dedicated to database preparation before starting the cross-validation. In Step 2, we performed the cross-validation and computed the performance metrics in Step 3. In Step 4, we calibrated SDMs only for the selected species $AUC \geq 0.7$ and computed the prediction maps for the six specified scenarios. The last step was devoted to the assemblage analysis.

216   a minimum record of 1% and 5% presence among sites for birds and beetles, respectively. These percentages

217   made sure that we only modeled species that occurred at ≥ 24 sites for birds and ≥ 14 sites for beetles.

2.4.2   Step 2: Model estimation

First, we removed the highly correlated variables, based on pairwise Pearson correlation coefficient (r), and

retained the 5 most important predictor variables (Zurell et al., 2020). To do so, we fitted a univariate

Generalized linear model (GLM) with linear and quadratic terms for each variable; we ranked the predictors

according to their importance, using the Akaike information criterion (AIC); we removed the highly correlated

variables ($|r| > 0.60$). To reduce the separation in the regressions, we removed any predictor with a standard

deviation value greater than 50 through a stepwise procedure.

We started with a generalized linear mixed model (GLMM) (package 'lme4', (Bates et al., 2015)) with a

random intercept to account for differences between sampling years. We developed six and three full potential

models differing only in the fixed effects for CHBMs and HOBMs model classes, respectively (see Table 3 in

Supporting Information). We used interaction terms between the best temperature variable (where the AIC

criterion of the corresponding univariate regression is the lowest among all selected temperature variables)

with distance to the nearest burned stand and with stand age variables to include the effect of latitudinal
variation.

The best models were selected based on each full-model under 10-fold cross-validation by minimizing the
AIC criterion (package 'MuMIn', (Barton, 2015)) under the following conditions: (1) limiting the number of
terms in the model between 1 and $\lceil min(N_{Pre}, N_{Abs})/5 \rceil$ without counting the intercept by specifying 1 in
5 rule, where N_{Pre} and N_{Abs} represented the number of respective presence and absence records, to avoid
overparameterized models; and (2) respecting the principle of marginality when the interaction terms were
included in the model. For each species and for each test dataset, we calculated the occurrence probabilities
matrix for the calibrated models. In total, we retained six models for CHBMs and three for the HOBMs.

2.4.3   Step 3: Model selection

Once the occurrence probabilities were computed for the complete dataset for each species, we calculated
the following performance metrics: (1) specificity; (2) sensitivity; (3) the area under the curve AUC; and
(4) the true skill statistic TSS (Araújo et al., 2005; Allouche et al., 2006). The package 'AUC' (Ballings
and Van den Poel, 2013) was used to calculate the Receiver Operating Characteristic (ROC) and the AUC;
we used the package 'PresenceAbsence' (Freeman and Freeman, 2012) for the other performance metrics.
Species were excluded if no model yielded $AUC \geq 0.7$ (Araújo et al., 2005; Hosmer et al., 2013). For the
selected species, we used the model with the highest AUC for projections.

2.4.4   Step 4: Calibration and prediction

248 We estimated the parameters of the final model for the selected species based on the full dataset using
249 the same procedure that was described for cross-validation. We subsequently used the simulated predictor
250 variable maps for the six study scenarios and computed the occurrence probability maps.

2.4.5   Step 5: Assemblage and occupancy analysis

252 We used the following indices to compare species assemblages between scenarios:

253 **Regional occupancy probability** (ROP):   ROP was calculated as the regional average of the occurrence
254 probabilities for the study area (Bichet et al., 2016). We also used the percentage of change in ROP between
255 the reference (Ref) and the simulated (Sim) scenarios (see step 5 in Fig. 4 for the definitions of Ref and

256 Sim scenarios): $\Delta\overline{ROP}_{Sim,Ref} = \left(\dfrac{\overline{ROP}_{Sim} - \overline{ROP}_{Ref}}{\overline{ROP}_{Ref}}\right) \times 100$, where $\overline{ROP}_s$ represented the average of

257 the ROP over all species under the scenario s given by:

$$
\begin{aligned}
\overline{ROP}_s &= \frac{1}{N_{species}} = \sum_{i=1}^{N_{species}} ROP_{i,s}, \\
ROP_{i,s} &= \frac{1}{N_{pixel}} = \sum_{j=1}^{N_{pixel}} P_{s,i,j},
\end{aligned}
\tag{1}
$$

258   where $N_{species}$, N_{pixel} and $P_{s,i,j}$ represented respectively the species number, the number of pixels in the

259 study area and the occurrence probability of the species i for the scenario s at the cell j.

260 **Dissimilarity measures:**   Jaccard Index of Dissimilarity was used to assess climate change effects on

261 species assemblage (Baselga, 2010, 2013; Legendre, 2014; Bichet et al., 2016; Barton et al., 2016; Belliard

262 et al., 2018; Scherrer et al., 2020). We computed two different Jaccard indices, both based on Bray-Curtis

263 dissimilarity measure, regional (RJ) and spatial (SJ) Jaccard indices computed as follows:

$$
\begin{aligned}
RBC_{Sim,Ref} &= \sum_{i=1}^{N_{species}} \frac{|ROP_{Sim,i} - ROP_{Ref,i}|}{\sum_{i'=1}^{N_{species}} (ROP_{Sim,i'} + ROP_{Ref,i'})}, \\
SBC_{Sim,Ref,j} &= \sum_{i=1}^{N_{species}} \frac{|P_{Sim,i,j} - P_{Ref,i,j}|}{\sum_{i'=1}^{N_{species}} (P_{Sim,i',j} + P_{Ref,i',j})}, \\
RJ_{Sim,Ref} &= \frac{2RBC_{Sim,Ref}}{1 + RBC_{Sim,Ref}}, \qquad SJ_{Sim,Ref,j} = \frac{2SBC_{Sim,Ref,j}}{1 + SBC_{Sim,Ref,j}}.
\end{aligned}
\tag{2}
$$

264 $RBC_{Sim,Ref}$ and $SBC_{Sim,Ref,j}$ represented respectively the regional Bray-Curtis dissimilarity between

265 the scenarios Sim and Ref, the spatial Bray-Curtis at the cell j selected randomly.

266   From Fig. 5, we compared the performance of the Jaccard index that was based on our continuous

267 output (SJ_{OP}) and the traditional Jaccard index that was based on the presence-absence transformation

268 ($SJ_{Incidence}$). From the simulation results, the SJ_{OP} yields results that were close to $SJ_{Incidence}$; moreover,

269 performance increased with the number of species that were analyzed (see Fig. 5I). Furthermore, we added

270 two situations with weaker binarization (Bad and Medium cases in Fig. 5) to illustrate that a gap can be

271 generated between the two indices that is mainly due mainly to binarization error.

272 **Beta ratio:**   We separated the Bray-Curtis dissimilarity index into two additive components main drivers

273 in assemblage dissimilarity: (1) the occurrence gradient (BC_grad); and (2) balanced variation in species

274 occurrence (BC_bal). We used these notations instead of the abundance gradient and balanced variation in

Figure 5: A generic Incidence-based vs. Occurrence Probabilities Jaccard dissimilarity. The three levels of binarization concerned the rank choice of the occurrence probabilities that were simulated from a mixture of uniform distributions under constraints. **(a)-(b)-(c)** represented the three simulated occurrence probabilities and the corresponding binarized maps with a 0.5 threshold for one species and under one scenario. **(d)-(e)-(f)** showed the frequency of the occurrence probability at the different pixels. **(g)-(h)-(i)** concerned the absolute difference between Jaccard dissimilarity based on the binarized data and the continuous version by varying the number of species.

species abundance (Baselga, 2013), given that we worked with occurrence and occupancy probabilities. We presented a detailed explanation for each component, as follows:

- If $BC\_grad_{Sim,Ref} = 0$ ($BC_{Sim,Ref} = BC\_bal_{Sim,Ref}$), means a **total** absence of differences in occurrences between the two scenarios. Furthermore, if $BC_{Sim,Ref} \simeq 0$, the assemblage structure remained almost the same. If $BC_{Sim,Ref} \simeq 1$, the occurrence of species in one scenario was almost perfectly balanced by the occurrence of species in the other scenario.

- If $BC\_bal_{Sim,Ref} = 0$ ($BC_{Sim,Ref} = BC\_grad_{Sim,Ref}$), this means that all species occurrence changes from one scenario to the next were in **the same direction**. If $BC_{Sim,Ref} \simeq 1$, this implies that the difference in occurrence between the two scenarios is too large and in the same direction in almost all species.

To measure the fraction of each component of Bray-Curtis dissimilarity, we used the BC_ratio given by

$$BC\_ratio_{Sim,Ref} = \frac{BC\_grad_{Sim,Ref}}{BC_{Sim,Ref}}.$$ If $BC\_ratio > 0.5$, the assisted community change was caused mostly by the occurrence gradient, whereas a value smaller than 0.5 indicated a dominance of balanced change in occurrence in the assemblage dissimilarity (Albouy et al., 2012). We used the package 'betapart' (Baselga and Orme, 2012) for the assemblage analysis.

The BC_ratio and its components were illustrated by using occurrence probabilities. The same formalism was adapted regionally through the regional occupancy probabilities.

3   Results

Of 231 candidate species of birds and beetles, 127 and 108 species were selected for projection (with $AUC \geq$ 0.7) respectively for CHBMs and HOBMs, with an average of AUC between 0.76 (± 0.07) and 0.79 (± 0.08), and TSS between 0.43 (± 0.13) and 0.53 (± 0.16) (mean $\pm$ SD). For bird species, *stand age* was the most frequently selected predictor variable with 43.9% and 71% of selection in CHBMs and HOBMs, respectively (see Fig. 6). For beetle species, mean temperature of the warmest month (*WarmMTmean*) and *stand age* were the most frequently selected variables under CHBMs and HOBMs with 33.7% and 40.3% of selection, respectively (see Fig. 6).   .

3.1   Occurrence and regional occupancy

The analysis of the occurrence and occupancy helped us to evaluate the direction and magnitude of potential community changes following global change. The difference between the indirect and mixed climate effects was visualized through $\Delta\overline{ROP}_{Sim,Ref}$ (Fig. 7A). The magnitude of changes in occupancy compared to the baseline reference scenario was larger when we included the climate variables (CHBMs) when compared to the models including only habitat variables (HOBMs). The change in occupancy was slightly more significant for birds, compared to beetles under HOBMs. Under CHBMs, we observed that occupancy increased for birds but decreased for beetles when comparing the same radiative-forcing scenarios (RCP4.5 and RCP8.5) to the baseline (the projection maps also demonstrated this result; compare Fig. 8E to Fig. 8F and Fig. 8A to Fig. 8B). However, we observed a decrease in the percentage of winner species with climate change for the two taxa, except for birds under HOBMs with no harvest (Fig. 11; see Fig. 1 in Supporting Information responses of five variable classes on eight winner and loser species). Moreover, to observe the change at the species level, Table 5 to Table 9 in the Supporting Information contained the percentage of change in the

Figure 6: Percentage of the predictor variables that were included in the species regressions for the two taxa (See Table. 1 in Supporting Information for the variable descriptions), **A.** for Habitat-Only-Based-Models (HOBMs) and **B.** for Climate-Habitat-Based Models (CHBMs). Land cover abbreviations: Conifer Dense (CD), Conifer Open (CO), Mixed Wood (MW), Open Habitat (OH), Disturbance by Fire (DF), and Disturbance by Cut (DC). For example, under CHBMs, 21% and 44% of beetle and bird species, respectively, selected Stand age variable .

Figure 7: Regional dissimilarity and change in occupancy. **A.** Bray-Curtis ratio (BC_ratio) with the percentage of change in the regional occupancy probability ($\Delta \overline{ROP}$) under the four compared landscapes. **B** Bird and Beetle Jaccard dissimilarity measures. Abbreviations: Climate-Habitat-Based Model (**CHBM**), Habitat-Only-Based Model (**HOBM**).

regional occupancy probability for the accepted species under the two model classes (CHBMs and HOBMs).

3.2   Regional species assemblage change

An increase in assemblage dissimilarity was observed when comparing RCP4.5 and RCP8.5 to the baseline for both taxa. Based on CHBMs, regional dissimilarity from RCP4.5 to RCP8.5 increased respectively by 0.20

Figure 8: The average potential occurrence maps for each taxon according to two model classes (CHBMs and HOBMs) and four scenarios: BaselineNoHarvest (**A, E, I, M**), RCP8.5NoHarvest (**B, F, J, N**), BaselineHarvest (**C, G, K, O**) and RCP8.5Harvest (**D, H, J, P**).

and 0.14 for birds and beetles under no harvest (Fig. 7B-C). Also, based on HOBMs, regional dissimilarity from RCP4.5 to RCP8.5 increased respectively by 0.07 and 0.08 for birds and beetles under no harvest. This regional dissimilarity was mainly incurred through balanced variation in species occupancy for both taxa ($BC\_ratio < 0.5$) (see Fig. 7A).

We also observed an increase in assemblage dissimilarity from HOBMs to CHBMs for both taxa. Inclusion of the climate variables reshaped assemblage structure strongly under the two forest management levels (see differences in the Jaccard dissimilarity index between HOBMs and CHBMs, Fig. 7B-C).

3.3   Latitudinal changes in species assemblage

The inclusion of climate variables produced a latitudinal gradient in projections of assemblage dissimilarity. We used CHBMs to analyze how latitudinal changes in temperature and other climate variables would affect assemblage structure. A clear increasing pattern was observed in assemblage dissimilarity heading north,

especially for beetle species. For birds, a slight increase in dissimilarity was observed, compared to that of beetles (see the difference between Fig. 9A and Fig. 9C).

Figure 9: Spatial analysis under CHBMs. **A, C.** Latitudinal variation of spatial Jaccard dissimilarity of **RCP4.5Harvest** and **RCP8.5Harvest** compared to reference scenario **BaselineHarvest**. **B, D.** Latitudinal variation in the spatial BC_ratio. The shaded areas represented the standard deviation. Latitudes are given in the UTM coordinate system.

Our models predict that beetles would show greater sensitivity to climate variations, given that an increase in dissimilarity was observed even for a medium level of climate change (Figs. 9A), i.e., RCP 4.5 following changes in temperature between the baseline and the two forcing scenarios. In Figs 10A-B, we depicted spatial change in $Tmin$ and $WarmMTmean$ between RCP8.5 and baseline climate scenarios. $Tmin$ and $WarmMTmean$ were the most frequently selected respective predictor variables for birds and beetles (see Fig. 6B for the selection percentage). Furthermore, the observed latitudinal dissimilarity gradient was derived mainly from balanced variation in occurrence ($BC\_ratio < 0.5$ in Figs. 9B, D). Yet, the two taxa behaved in a contrasting manner with regard to their latitudinal variation in the occurrence gradient component of BC dissimilarity.

Figure 10: Temperature difference maps between RCP8.5 and baseline of annual minimum temperature (**A**) and mean temperature of the warmest month (**B**) in °C. Map scale: The dark red and the dark blue represented respectively the highest and the lowest values. The horizontal and vertical marginal plots respectively show the longitudinal and latitudinal distribution of the temperature differences.

Figure 11: Percentage of winner species per taxon under the four landscapes that were compared and the two model classes (CHBMs and HOBMs). Species i was considered a winner if the regional occupancy probability under the simulated scenario was higher than under the reference scenario ($ROP_{Sim,i} > ROP_{Ref,i}$).

4   Discussion

The present study reveals several intricate effects of global change on animal species assemblages. We showed that the two climate-induced pathways that were studied increased dissimilarity in species assemblage mostly by inducing species turnover, under both forest harvesting scenarios. Of the two pathways, the models integrating both climate and habitat effects strongly modified the assemblage composition more than did the models based only on habitat effects. In fact, the difference in magnitude between the two effects was due to the mismatch that was generated by rapid climate variation compared to slower vegetation change (Wu et al., 2015; Stralberg et al., 2015; Micheletti et al., 2021). Both climate effects caused the decrease in number of winner species from RCP 8.5 to RCP 4.5 compared to the baseline scenario, under the two

forest harvesting scenarios. However, some species remained winners under both climate effects such as

Stenichnus turbatus (Casey) [Staphylindae] and *Chipping Sparrow*, while others changed from winner to

loser species, such *Stenichnus perforatus* (Schaum) [Staphylinidae] and *Greater Yellowlegs* (see Table. 6-9

in Supporting Information). Nevertheless, we observed almost an opposite feedback between the two taxa

regarding changes in the regional occupancy. This could result from physiological differences between taxa

in their individual response to climate.

Our analysis predicts pronounced variations in assemblage composition for both birds and beetles by

2100 that will be caused by climate change under any of the harvest levels. Whereas a rise in anthropogenic

radiative forcing should increase dissimilarity in species assemblage following each of the two climate path-

ways, the magnitude of change differed considerably between the two pathways. For example, the minimum

and the maximum assemblage dissimilarities were 0.49 and 0.14 under climate-habitat and habitat-only

models respectively. This result could explain the subsequent mismatch between the climate and the biota

(Stralberg et al., 2015; Wu et al., 2015; Micheletti et al., 2021). For example, Brice et al. (2020) noted that

under climate change the variation in climate conditions would be faster than the capacity of tree species

to migrate, a discrepancy that is expected to create a gap between the climatic niche and the observed

distribution of species.

Regarding the direction of change at the taxon level, we observed that birds and beetles responded in

almost opposite directions. In fact, the increase in bird occupancy that resulted from the mixed combination

between direct and indirect climate effects coincided in overall with the *northern biodiversity paradox* that

was emphasized by Matthews et al. (2004), Morin and Thuiller (2009), and Berteaux et al. (2010), which

anticipated an increase in biodiversity in northern protected areas during this century. More precisely, the

northern biodiversity paradox suggests that climate change could lead to a potential increase in the abundance

of many species for which the low temperatures are currently a limiting factor (Berteaux et al., 2010). For

beetles, we observed a substantial decrease in regional occupancies following an increase in climate change.

This outcome aligned with the global scale biodiversity trajectory predicting mostly negative effects of high

emissions of greenhouse gas on biodiversity and ecosystem services (Pachauri et al., 2014). Given that beetles

are poikilotherms, their internal temperatures vary according to the temperature of the environment, while

birds are homeotherms, the internal temperatures of which are physiologically regulated. Our results suggest

the potential presence of cold-habitat beetle species in the north. With increasing radiative forcing (Fig. 8F),

their occurrence probability decreases as climate conditions extend beyond their tolerance limits. However,

those species may also be affected by the competition with southern species invading northern habitats. This

could explain the different global implications of direct climate change on beetles and birds. Our conclusions

accord with those of other studies based on limitations to phenotypic plasticity and evolvability of critical thermal tolerances in insects (Gaston and Chown, 1999; Terblanche et al., 2007; García-Robledo et al., 2016). For instance, García-Robledo et al. (2016) demonstrated the role of critical thermal temperature on insect tolerance to global warming in the context of elevation, which was used as a proxy for latitude in context of global warming. The authors demonstrated that insects found at middle elevations and on mountain tops were less tolerant to temperature increases than were species located at lower latitudes.

Under mixed climate effects, we observed substantial latitudinal changes in biodiversity with increasing radiative forcing, with an almost complete change in beetle assemblages in the northern portion of the study area. Latitudinal patterns were also noted by Brice et al. (2019), who reported a northward decrease in temporal $Beta - diversity$ of tree species between past (1970-1980) and present (2000-2016) periods. Our results warn about the possibility that direct climatic effects could increase over the next several decades, particularly for insects. This latitudinal gradient could be a result of the increase in temperature after a change in global climate forcing (Holland and Bitz, 2003). However, the climatic envelope that was used for calibration could increase the effects of the climate change on species occurrence compared to a study that is based upon a greater latitudinal range.

In conclusion, based on implicit assumptions regarding individual species responses to climate change, our analyses identified potential repercussions of two climate-induced pathways that were based on direct and indirect effects on the assemblage composition of two taxonomic groups regionally and latitudinally, under two different forest harvesting scenarios. The observed change was derived mainly by species turnover regardless of harvest level. Moreover, based on the studied species pool and despite the possibility that other species could arrive, we could expect large differences in occupancy between the two studied taxa, which are explained probably by their ecological trait differences. This could indicate the potential range of change in boreal species concerning novel environmental conditions.

Acknowledgements

This work was supported by the Sentinel North program of Laval University, funded by the Canada First Research Excellence Fund. A.A., D.F., C.H., and P.D. were also supported by the Natural Sciences and Engineering Research Council of Canada (NSERC). We acknowledge Calcul Québec and Compute Canada for their technical support and computing infrastructures. We thank also the Québec Breeding Bird Atlas for supplying data. We would also like to thank the following partners: Regroupement QuébecOiseaux, Environment and Climate Change Canada and Birds Canada, together with all the volunteer participants

who gathered data for the project. We are grateful to Louis-Paul Rivest for his valuable statistical suggestions and advice. Special thanks are due to Alexandre Terrigeol for his valuable efforts regarding data collection, Georges Pelletier and Nicolas Bédard for beetle identification and W.F.J. Parsons for his English revision.

References

Albouy, C., Guilhaumon, F., Araújo, M. B., Mouillot, D., and Leprieur, F. (2012). Combining projected changes in species richness and composition reveals climate change impacts on coastal Mediterranean fish assemblages. *Global Change Biology*, 18(10):2995–3003.

Allouche, O., Tsoar, A., and Kadmon, R. (2006). Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, 43(6):1223–1232.

Araújo, M. B., Pearson, R. G., Thuiller, W., and Erhard, M. (2005). Validation of species–climate impact models under climate change. *Global Change Biology*, 11(9):1504–1513.

Atlas des oiseaux nicheurs du Québec (2018). Données obtenues en réponse à une demande présentée aux bureaux de l'Atlas (www.atlas-oiseaux.qc.ca). Regroupement QuébecOiseaux, Service canadien de la faune d'Environnement Canada et Études d'Oiseaux Canada. Québec, QC, Canada.

Ballings, M. and Van den Poel, D. (2013). Package 'AUC'. https://cran.r-project.org/web/packages/AUC.

Barton, A. D., Irwin, A. J., Finkel, Z. V., and Stock, C. A. (2016). Anthropogenic climate change drives shift and shuffle in North Atlantic phytoplankton communities. *Proceedings of the National Academy of Sciences of the United States of America*, 113(11):2964–2969.

Barton, K. (2015). Package 'MuMIn'. https://cran.r-project.org/web/packages/MuMIn.

Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, 19(1):134–143.

Baselga, A. (2013). Separating the two components of abundance-based dissimilarity: balanced changes in abundance vs. abundance gradients. *Methods in Ecology and Evolution*, 4(6):552–557.

Baselga, A. and Orme, C. D. L. (2012). betapart: an R package for the study of beta diversity. *Methods in Ecology and Evolution*, 3(5):808–812.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using lme4. https://cran.r-project.org/web/packages/lme4.

Beaudoin, A., Bernier, P., Guindon, L., Villemaire, P., Guo, X., Stinson, G., Bergeron, T., Magnussen, S., and Hall, R. (2014). Mapping attributes of Canada's forests at moderate resolution through kNN and MODIS imagery. *Canadian Journal of Forest Research*, 44(5):521–532.

Belliard, J., Beslagic, S., Delaigue, O., and Tales, E. (2018). Reconstructing long-term trajectories of fish assemblages using historical data: the Seine River Basin (France) during the last two centuries. *Environmental Science and Pollution Research*, 25(24):23430–23450.

Bentz, B. J., Rgnire, J., Fettig, C. J., Hansen, E. M., Hayes, J. L., Hicke, J. A., Kelsey, R. G., Negron, J. F., and Seybold, S. J. (2010). Climate change and bark beetles of the western United States and Canada: Direct and indirect effects. *BioScience*, 60(8):602–613.

Berteaux, D., de Blois, S., Angers, J.-F., Bonin, J., Casajus, N., Darveau, M., Fournier, F., Humphries, M. M., McGill, B., Larivée, J., et al. (2010). The CC-Bio Project: studying the effects of climate change on Quebec biodiversity. *Diversity*, 2(11):1181–1204.

Bibby, C. J., Burgess, N. D., Hill, D. A., Hillis, D. M., and Mustoe, S. (2000). *Bird census techniques*. Elsevier.

Bichet, O., Dupuch, A., Hébert, C., Le Borgne, H., and Fortin, D. (2016). Maintaining animal assemblages through single-species management: the case of threatened caribou in boreal forest. *Ecological Applications*, 26(2):612–623.

Bouchard, M. and Pothier, D. (2011). Long-term influence of fire and harvesting on boreal forest age structure and forest composition in eastern Québec. *Forest Ecology and Management*, 261(4):811–820.

Boucher, Y., Perrault-Hébert, M., Fournier, R., Drapeau, P., and Auger, I. (2017). Cumulative patterns of logging and fire (1940–2009): consequences on the structure of the eastern Canadian boreal forest. *Landscape Ecology*, 32(2):361–375.

Boulanger, Y., Parisien, M.-A., and Wang, X. (2018a). Model-specification uncertainty in future area burned by wildfires in Canada. *International Journal of Wildland Fire*, 27(3):164–175.

Boulanger, Y. and Puigdevall, J. P. (2021). Boreal forests will be more severely affected by projected anthropogenic climate forcing than mixedwood and northern hardwood forests in eastern Canada. *Landscape Ecology*, 36(6):1725–1740.

Boulanger, Y., Taylor, A. R., Price, D. T., Cyr, D., McGarrigle, E., Rammer, W., Sainte-Marie, G., Beaudoin, A., Guindon, L., and Mansuy, N. (2017). Climate change impacts on forest landscapes along the Canadian southern boreal forest transition zone. *Landscape Ecology*, 32(7):1415–1431.

Boulanger, Y., Taylor, A. R., Price, D. T., Cyr, D., and Sainte-Marie, G. (2018b). Stand-level drivers most important in determining boreal forest response to climate change. *Journal of Ecology*, 106(3):977–990.

Brice, M.-H., Cazelles, K., Legendre, P., and Fortin, M.-J. (2019). Disturbances amplify tree community responses to climate change in the temperate–boreal ecotone. *Global Ecology and Biogeography*, 28(11):1668–1681.

Brice, M.-H., Vissault, S., Vieira, W., Gravel, D., Legendre, P., and Fortin, M.-J. (2020). Moderate disturbances accelerate forest transition dynamics under climate change in the temperate–boreal ecotone of eastern North America. *Global Change Biology*, 26(8):4418–4435.

Bureau du Forestier en chef (2010). Bilan d'aménagement forestier durable au Québec 2000-2008. Gouvernement du Québec, Roberval, Québec, 290 p.

Bush, E. and Lemmen, D. S., editors (2019). *Canada's Changing Climate Report. Government of Canada, Ottawa, ON, Canada, 444 p.*

Cadieux, P., Boulanger, Y., Cyr, D., Taylor, A. R., Price, D. T., Sólymos, P., Stralberg, D., Chen, H. Y., Brecka, A., and Tremblay, J. A. (2020). Projected effects of climate change on boreal bird community accentuated by anthropogenic disturbances in western boreal forest, Canada. *Diversity and Distributions*, 26(6):668–682.

Dubuis, A., Pottier, J., Rion, V., Pellissier, L., Theurillat, J.-P., and Guisan, A. (2011). Predicting spatial patterns of plant species richness: a comparison of direct macroecological and species stacking modelling approaches. *Diversity and Distributions*, 17(6):1122–1131.

Duveneck, M. J., Scheller, R. M., White, M. A., Handler, S. D., and Ravenscroft, C. (2014). Climate change effects on northern Great Lake (USA) forests: A case for preserving diversity. *Ecosphere*, 5(2):art23.

Freeman, E. and Freeman, M. E. (2012). Package 'PresenceAbsence'. https://cran.r-project.org/web/packages/PresenceAbsence/.

García-Robledo, C., Kuprewicz, E. K., Staines, C. L., Erwin, T. L., and Kress, W. J. (2016). Limited tolerance by insects to high temperatures across tropical elevational gradients and the implications of

global warming for extinction. *Proceedings of the National Academy of Sciences of the United States of America*, 113(3):680–685.

Gaston, K. J. and Chown, S. L. (1999). Elevation and climatic tolerance: A test using dung beetles. *Oikos*, 86(3):584–590.

Gelfand, A. E., Schmidt, A. M., Wu, S., Silander Jr, J. A., Latimer, A., and Rebelo, A. G. (2005). Modelling species diversity through species level hierarchical modelling. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 54(1):1–20.

Girardin, M. P., Bouriaud, O., Hogg, E. H., Kurz, W., Zimmermann, N. E., Metsaranta, J. M., de Jong, R., Frank, D. C., Esper, J., Büntgen, U., Guo, X. J., and Bhatti, J. (2016). No growth stimulation of Canada's boreal forest under half-century of combined warming and CO_2 fertilization. *Proceedings of the National Academy of Sciences of the United States of America*, 113(52):E8406–E8414.

Grenié, M., Violle, C., and Munoz, F. (2020). Is prediction of species richness from stacked species distribution models biased by habitat saturation? *Ecological Indicators*, 111:105970.

Guisan, A. and Thuiller, W. (2005). Predicting species distribution: offering more than simple habitat models. *Ecology Letters*, 8(9):993–1009.

Guisan, A. and Zimmermann, N. E. (2000). Predictive habitat distribution models in ecology. *Ecological Modelling*, 135(2):147–186.

Gustafson, E. J., Shifley, S. R., Mladenoff, D. J., Nimerfro, K. K., and He, H. S. (2000). Spatial simulation of forest succession and timber harvesting using landis. *Canadian Journal of Forest Research*, 30(1):32–43.

Hillebrand, H., Soininen, J., and Snoijs, P. (2010). Warming leads to higher species turnover in a coastal ecosystem. *Global Change Biology*, 16(4):1181–1193.

Holland, M. M. and Bitz, C. M. (2003). Polar amplification of climate change in coupled models. *Climate Dynamics*, 21(3):221–232.

Hosmer, D. W., Lemeshow, S., and Sturdivant, R. X. (2013). *Applied logistic regression, third edition*. John Wiley & Sons.

Janssen, P., Fortin, D., and Hébert, C. (2009). Beetle diversity in a matrix of old-growth boreal forest: influence of habitat heterogeneity at multiple scales. *Ecography*, 32(3):423–432.

Kelly, A. E. and Goulden, M. L. (2008). Rapid shifts in plant distribution with recent climate change. *Proceedings of the National Academy of Sciences of the United States of America*, 105(33):11823–11826.

Kumar, P. (2013). Hydrology: Seasonal Rain Changes. *Nature Climate Change*, 3(9):783–784.

Labadie, G., Hardy, C., Boulanger, Y., Vanlandeghem, V., Hebblewhite, M., and Fortin, D. (2022). Global changes impact a threatened species through alteration of predator-prey dynamics. *To be submitted to Nature Climate Change.*

Labadie, G., McLoughlin, P. D., Hebblewhite, M., and Fortin, D. (2021). Insect-mediated apparent competition between mammals in a boreal food web. *Proceedings of the National Academy of Sciences of the United States of America*, 118(30).

Légaré, J.-P., Hébert, C., and Ruel, J.-C. (2011). Alternative silvicultural practices in irregular boreal forests: Response of beetle assemblages. *Silva Fennica*, 45(5):937–956.

Legendre, P. (2014). Interpreting the replacement and richness difference components of beta diversity. *Global Ecology and Biogeography*, 23(11):1324–1334.

MacKenzie, D. I., Nichols, J. D., Royle, J. A., Pollock, K. H., Bailey, L., and Hines, J. E. (2017). *Occupancy estimation and modeling: inferring patterns and dynamics of species occurrence*. Elsevier.

Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P. R., et al. (2018). Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty. Technical report, IPCC.

Matthews, S., O'Connor, R., Iverson, L. R., and Prasad, A. M. (2004). Atlas of climate change effects in 150 bird species of the Eastern United States. Gen. Tech. Rep. NE-318. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 340 p.

Matuoka, M. A., Benchimol, M., de Almeida-Rocha, J. M., and Morante-Filho, J. C. (2020). Effects of anthropogenic disturbances on bird functional diversity: A global meta-analysis. *Ecological Indicators*, 116:106471.

McKenney, D., Pedlar, J., Hutchinson, M., Papadopol, P., Lawrence, K., Campbell, K., Milewska, E., Hopkinson, R. F., and Price, D. (2013). Spatial climate models for Canada's forestry community. *Forestry Chronicle*, 89(05):659–663.

Micheletti, T., Stewart, F. E. C., Cumming, S. G., Haché, S., Stralberg, D., Tremblay, J. A., Barros, C., Eddy, I. M. S., Chubaty, A. M., Leblond, M., Pankratz, R. F., Mahon, C. L., Van Wilgenburg, S. L.,

Bayne, E. M., Schmiegelow, F., and McIntire, E. J. B. (2021). Assessing pathways of climate change
effects in SpaDES: An application to bBoreal Landbirds of Northwest Territories Canada. *Frontiers in*
Ecology and Evolution, 9(October).

Ministère des Forêts de la Faune et des Parcs (2018). Aires infestées par la tordeuse des bourgeons de
l'épinette au québec en 2018, 20 p. Technical report, Québec, gouvernement du Québec, Direction de la
protection des forêts.

Morin, X. and Thuiller, W. (2009). Comparing niche-and process-based models to reduce prediction uncer-
tainty in species range shifts under climate change. *Ecology*, 90(5):1301–1313.

Naeem, S., Duffy, J. E., and Zavaleta, E. (2012). The functions of biological diversity in an age of extinction.
Science, 336(6087):1401–1406.

Pachauri, R. K., Allen, M. R., Barros, V. R., Broome, J., Cramer, W., Christ, R., Church, J. A., Clarke,
L., Dahe, Q., Dasgupta, P., et al. (2014). *Climate change 2014: synthesis report. Contribution of Working*
Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change.
IPCC, Gland, Switzerland.

Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martínez-Meyer, E., Nakamura, M., and
Araújo, M. B. (2011). *Ecological niches and geographic distributions (MPB-49)*, volume 49. Princeton
University Press.

Pureswaran, D. S., De Grandpré, L., Paré, D., Taylor, A., Barrette, M., Morin, H., Régnière, J., and
Kneeshaw, D. D. (2015). Climate-induced changes in host tree–insect phenology may drive ecological
state-shift in boreal forests. *Ecology*, 96(6):1480–1491.

Régnière, J., Saint-Amant, R., Béchard, A., and Moutaoufik, A. (2017). BioSIM 11: User's manual. Technical
report, Natural Resources Canada, Canadian Forest Service, Laurentian Forestry Centre, Québec, QC
Canada.

Scheller, R. M. and Mladenoff, D. J. (2004). A forest growth and biomass module for a landscape simulation
model, LANDIS: design, validation, and application. *Ecological Modelling*, 180(1):211–229.

Scherrer, D., Mod, H. K., and Guisan, A. (2020). How to evaluate community predictions without thresh-
olding? *Methods in Ecology and Evolution*, 11(1):51–63.

Sergio, F., Blas, J., and Hiraldo, F. (2018). Animal responses to natural disturbance and climate extremes:
a review. *Global and Planetary Change*, 161:28–40.

Stralberg, D., Bayne, E. M., Cumming, S. G., Sólymos, P., Song, S. J., and Schmiegelow, F. K. (2015). Conservation of future boreal forest bird communities considering lags in vegetation response to climate change: A modified refugia approach. *Diversity and Distributions*, 21(9):1112–1128.

Taylor, A. R., Boulanger, Y., Price, D. T., Cyr, D., McGarrigle, E., Rammer, W., and Kershaw, J. A. (2017). Rapid 21st century climate change projected to shift composition and growth of Canada's Acadian Forest Region. *Forest Ecology and Management*, 405:284–294.

Terblanche, J. S., Deere, J. A., Clusella-Trullas, S., Janion, C., and Chown, S. L. (2007). Critical thermal limits depend on methodological context. *Proceedings of the Royal Society B: Biological Sciences*, 274(1628):2935–2943.

Thuiller, W., Guéguen, M., Bison, M., Duparc, A., Garel, M., Loison, A., Renaud, J., and Poggiato, G. (2018). Combining point-process and landscape vegetation models to predict large herbivore distributions in space and time— A case study of *Rupicapra rupicapra*. *Diversity and Distributions*, 24(3):352–362.

Thuiller, W., Lavergne, S., Roquet, C., Boulangeat, I., Lafourcade, B., and Araujo, M. B. (2011). Consequences of climate change on the tree of life in Europe. *Nature*, 470(7335):531–534.

Tremblay, J. A., Boulanger, Y., Cyr, D., Taylor, A. R., Price, D. T., and St-Laurent, M.-H. (2018). Harvesting interacts with climate change to affect future habitat quality of a focal species in eastern Canada's boreal forest. *PLoS ONE*, 13(2):1–25.

van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K. (2011). The representative concentration pathways: an overview. *Climatic Change*, 109(1):5.

Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., Dormann, C. F., Forchhammer, M. C., Grytnes, J.-A., Guisan, A., Heikkinen, R. K., Høye, T. T., Kühn, I., Luoto, M., Maiorano, L., Nilsson, M.-C., Normand, S., Öckinger, E., Schmidt, N. M., Termansen, M., Timmermann, A., Wardle, D. A., Aastrup, P., and Svenning, J.-C. (2013). The role of biotic interactions in shaping distributions and realised assemblages of species: implications for species distribution modelling. *Biological Reviews*, 88(1):15–30.

Wu, D., Zhao, X., Liang, S., Zhou, T., Huang, K., Tang, B., and Zhao, W. (2015). Time-lag effects of global vegetation responses to climate change. *Global Change Biology*, 21(9):3520–3531.

Zhang, Y. and Liang, S. (2014). Changes in forest biomass and linkage to climate and forest disturbances over northeastern china. *Global Change Biology*, 20(8):2596–2606.

610 Zhao, Q., Azeria, E. T., Le Blanc, M.-L., Lemaître, J., and Fortin, D. (2013). Landscape-scale disturbances

611   modified bird community dynamics in successional forest environment. *PLoS ONE*, 8(11):1–11.

612 Zurell, D., Zimmermann, N. E., Gross, H., Baltensweiler, A., Sattler, T., and Wüest, R. O. (2020). Testing

613   species assemblage predictions from stacked and joint species distribution models. *Journal of Biogeography*,

614   47(1):101–113.

A Regional change in occupancy and BC-ratio

B Regional Jaccard dissimilarity

NoHarvest	Harvest	Tmean	Precipitation	WaterDeficit

Baseline: A, B, C, D, E

RCP4.5: F, G, H, I, J

RCP8.5: K, L, M, N, O

Natural | Other | Fire | Cut

			Temperature	Precipitation	Water deficit	Fire	Harvest
Birds	CHBM	**Winner:** American Crow	⬆ (green)	⬇ (red)			
		Loser: Wilson's Snipe	⬇ (red)	⬆ (green)			
	HOBM	**Winner:** Palm Warbler				⬆ (green)	⬇ (red)
		Loser: Golden-crowned Kinglet				⬇ (red)	
Beetles	CHBM	**Winner:** Stenichnus turbatus	⬆ (green)				
		Loser: Liogluta terminalis	⬇ (red)		⬇ (red)	⬆ (green)	
	HOBM	**Winner:** Gnathacmaeops pratensis				⬆ (green)	
		Loser: Phymaphora pulchella				⬇ (red)	

Global change modeling

A Tmin:RCP8.5-BASE

B WarmMTmean:RCP8.5-BASE