Pan-cancer association of DNA repair deficiencies with whole-genome mutational

patterns

Simon G. Sørensen[1,2], Amruta Shrikhande[3], Gustav A. Poulsgaard[1,2], Mikkel H. Christensen[1,2],

Johanna Bertl[1,2], Eva R. Hoffmann[3]* and Jakob S. Pedersen[1,2,4]*

[1]Department of Molecular Medicine (MOMA), Aarhus University Hospital, Palle Juul-Jensens

Boulevard 99, 8200, Aarhus N, Denmark.

[2]Department of Clinical Medicine, Aarhus University, Palle Juul-Jensens Boulevard 82, 8200,

Aarhus N, Denmark.

[3]DNRF Center for Chromosome Stability, Department of Cellular and Molecular Medicine,

Faculty of Health and Medical Sciences, University of Copenhagen, Denmark

[4]Bioinformatics Research Center (BiRC), Aarhus University, 8000, Aarhus C, Denmark.

*corresponding authors: jakob.skou@clin.au.dk, eva@sund.ku.dk

ABSTRACT

DNA repair deficiencies in cancers may result in characteristic mutational patterns, as

exemplified by deficiency of *BRCA1/2* and efficacy prediction for PARP-inhibitors. We trained

and evaluated predictive models for loss-of-function (LOF) of 145 individual DDR genes based

on genome-wide mutational patterns, including structural variants, indels, and base-substitution

signatures. We identified 24 genes whose deficiency could be predicted with good accuracy,

including expected mutational patterns for *BRCA1/2*, *MSH3/6*, *TP53*, and *CDK12* LOF variants.

CDK12 is associated with tandem-duplications, and we here demonstrate that this association

can accurately predict gene deficiency in prostate cancers (area under the ROC curve=0.97).

Our novel associations include mono- or biallelic LOF variants of *ATRX, IDH1, HERC2,*

CDKN2A, PTEN, and *SMARCA4,* and our systematic approach yielded a catalogue of

predictive models, which may provide targets for further research and development of

treatment, and potentially help guide therapy.

Keywords: DNA damage response, repair deficiency, predictive modelling, personalised

medicine, mutational signatures.

LIST OF ABBREVIATIONS

AUROC | Area under the receiver operator characteristic

CNS | Central nervous system

DDR | DNA damage response

HMF | Hartwig medical foundation

HRD | Homologous recombination deficiency

Indel | Insertion or deletion

LASSO | least absolute shrinkage and selection operator

LASSO regression | Logistic regression with LASSO regularisation

LOF | Loss of function

LOH | Loss of heterozygosity

MMR | Mismatch repair

MSI | Microsatellite instability

PCAWG | Pan-cancer analysis of whole genomes

PR-AUC | Precision-recall area under the curve

PR-AUC-E | Precision-recall area under the curve enrichment from baseline

SBS | Single base substitution

SV | Structural variant

UV | Ultraviolet

VUS | Variants of unknown significance

WGS | Whole genome sequence

WT | Wild type

-d | Deficiency

INTRODUCTION

The DNA damage response (DDR) and repair pathways are central to the genetic integrity of cells, and deficiencies may cause mutational patterns genome-wide[1–3]. Some DNA repair deficiencies are known to modulate the response to therapies: *BRCA1/2* deficiency renders cancers susceptible to treatment with PARP inhibitors[4], mismatch repair-deficient cancers are sensitive to checkpoint inhibitors[5] but resistant to alkylating agents such as temozolamide[6], and *CDK12*-mutated cancers have a suggested sensitivity to CHK1 inhibitors[7]. Because of this, efforts have been made to annotate inactivating mutations in DDR genes[8]. However, the approach is limited by the lack of functional impact annotation of most variants, which are generally denoted as "variants of unknown significance" (VUS). Moreover, loss of gene activity could also occur by other means, such as transcriptional silencing.

A complementary approach is to investigate whether DNA repair deficiencies can be identified by DNA mutational patterns, also referred to as 'mutational scars'. This approach has been pioneered for homologous recombination deficiency caused by *BRCA1/2* deficiencies (-d), which can be successfully predicted by measuring the accumulation of small deletions with neighbouring microhomologous sequences[2,9,10], such as done by the HRDetect algorithm by Davies *et al.*[9]. The association with microhomologous deletions is due to the use of microhomology-mediated endjoining (MMEJ) to repair double-strand breaks in homologous recombination (HR) deficient tumours[11,12]. Likewise, mismatch repair-deficiency causes an elevated rate of mono- and dinucleotide repeat indels genome-wide, a genetic phenotype denoted microsatellite instability[13,14]. Mutations in other DNA repair genes have also been associated with mutational patterns, including the tumour suppressor gene *TP53*, which is associated with increased structural rearrangements and whole genome duplications[15,16] *CDK12* which is associated with a genome-wide phenotype of large tandem duplications[17,18].

The scope of this approach can now be evaluated systematically across DDR genes by

exploiting available whole cancer genomes from thousands of patients[19,20].

To achieve this, mutations observed genome-wide may be condensed into mutational summary

statistics for predictive modelling, including statistics based on single-base substitutions (SBS),

indels, and different types of structural variants (SV). The SBSs are statistically assigned to so-

called SBS signatures that are catalogued and enumerated within the COSMIC database[21].

Some of these are associated with specific DNA repair deficiencies as well as genotoxic

exposures, such as ultraviolet (UV) light and smoking. Each SBS signature captures the relative

frequency of the different mutation types and their flanking nucleotides[22].

Here we performed a systematic screen for DDR gene deficiencies that can be predicted

through their association with genome-wide mutational patterns. We developed a generic

approach to train predictive statistical models that identify associations with individual mutational

summary statistics that capture the mutational patterns, including SBS signatures, indels, and

large SVs. We applied it to 736 DDR gene deficiencies, considering both monoallelic and

biallelic loss of function (LOF), identified across 32 cancer types, in a combined set of whole

cancer genomes from 6,065 patients[19,20]. The underlying aim was to identify novel associations

with potential biological relevance and to evaluate whether DDR deficiencies can be predicted

with sufficiently high certainty to have a potential for clinical application.

Our analysis revealed 24 DDR genes where deficiencies are associated with specific mutational

summary statistics in individual cancer types across 48 predictive models. These results

recapitulated the expected associations between mutational patterns and deficiencies of

BRCA1/2, TP53, MSH3/6, and *CDK12*. We supplemented this knowledge by providing a

predictive model of *CDK12* deficiency that achieved high accuracy (area under the receiver

operator characteristic (AUROC) = 0.97) in prostate cancer. Furthermore, we present

unexpected predictive models of several DDR deficiencies; *ATRX* and *IDH1* deficiency in

cancers of the central nervous systems, *HERC2* and *CDKN2A* deficiency in skin, *PTEN*

deficiency in cancers of the central nervous system and uterus, and *SMARCA4* deficiency in

cancers of unknown primary.

RESULTS

DDR gene deficiencies across 6,065 whole cancer genomes

We compiled and analysed 2,568 whole genome sequences (WGS) from The Pan-Cancer

Analysis of Whole Genomes (PCAWG)[19] and 3,497 WGS from the Hartwig Medical Foundation

(HMF)[20]. In total we investigated 6,065 whole cancer genomes of 32 cancer types (**Fig. 1a;**

Additional file 1, Supplementary Table 1).

For each genome, we evaluated 736 known DNA damage response (DDR) genes for both

germline and somatic loss of function (LOF) events[23–25]. We annotated both mono- and biallelic

loss-of-function (LOF) events, where each event could be either a single nucleotide variant, an

indel, or a loss-of-heterozygosity (LOH) (**Fig. 1b,c; Additional file 1, Supplementary Table 2**).

Pathogenicity of single base substitutions (SBSs) and indels was evaluated using a combination

of CADD scores (>25; 0.3% most pathogenic variants) [26] and ClinVar annotation, when

available (**Methods**).

We inferred a total of 8,408 biallelic DDR gene deficiencies, primarily through a combination of

somatic or germline variants (SBSs and indels) with pathogenic potential (n=1,702), or LOH

events combined with a single pathogenic germline (n=3,562) or somatic (n=3,078) variant

(SBS or indel; **Fig. 1b**). On average we observed a single, biallelic DDR gene loss per patient,

with some tumours showing extreme rates of somatic pathogenic mutations (**Fig. 1c**). As

expected, *TP53* deficiency (*TP53*-d) was the most frequent LOF event (81 biallelic and 1,746

monoallelic events; 29% of tumours affected; **Additional file 1, Supplementary Table 2; Fig.**

1d), while 70% of DDR genes had biallelic deficiency in less than ten tumours across all cancer

types (511/736; **Fig. 1d**). Among monoallelic events, we identified 15,063 pathogenic germline

(59%) and 10,336 somatic (41%) events.

Whole genome mutational patterns

We collected mutational summary statistics for each cancer genome, which were used as

features for the downstream predictive models (**Fig. 1e,f; Additional file 2, Supplementary**

Table 3-4). For SBSs, we evaluated exposure towards predefined sets of cohort-specific SBS

signatures[22,27]. Short indels and SVs were simply categorised and counted: Deletions were sub-

categorised based on surrounding sequence repetitiveness and presence of microhomology.

SVs were sub-categorised by type (tandem duplications, inversions, deletions, and

translocations), five size ranges (not relevant for translocations), and cluster presence

(**Methods**). Several SBS signatures as well as some types of indels have suggested etiologies

(collected in **Additional file 1, Supplementary Table 5**).

Statistical modelling of DDR gene deficiencies

For the downstream statistical analysis, we restricted our focus to DDR genes in cancer types

with more than five biallelic LOF events in either PCAWG or HMF (n=194) or more than ten

monoallelic LOF events (n=341) (**Additional file 1, Supplementary Table 6**). Using *BRCA2*-d

in the set of HMF breast cancer tumours (n=645) as an example, we observed biallelic LOF

events in 17 (2.6%; 14 germline, 3 somatic) tumours and monoallelic LOF events in 7 (1.1%; 4

germline, 3 somatic) tumours (**Additional file 1, Supplementary Table 8**)**.** We further observed

variants of unknown significance (VUS) events in 53 tumours (8.2%; 42 germline, 11 somatic),

which were excluded from the analysis. The remaining *BRCA2* wild-type (WT) tumours (n=568;

88.1%) were used as a background set for training the predictive models (**Fig. 2a)**. The high

fraction of germline pathogenic variants diminishes the probability of a reverse-causal

relationship between the loss of *BRCA2* and the associated mutation patterns.

For each of the 535 groups of tumours we trained a LASSO regression model and evaluated the

ability to discriminate between deficient and WT tumours (**Methods**). For *BRCA2*-d we

observed a strong association with the number of deletions at sites of microhomology (**Fig. 2b**),

with a median of 608 deletions per patient in *BRCA2*-d breast cancers versus 81 in *BRCA2* WT

breast cancers, in agreement with prior findings[2,9,10]. The LASSO regression also included non-

clustered inversions 10-100Kb and clustered tandem duplications 1−10Kb, although both show

high variance among tumours for both deficient and WT (**Fig. 2b**) and have considerably

smaller coefficients, ultimately contributing little influence on overall predictive performance (**Fig.**

2c).

Notably, some models include features with negative coefficients. The biologically interpretation

would be that tumours with a certain gene deficiency have fewer mutations attributed to a

particular mutation pattern. Negative features were excluded in the development of the

HRDetect algorithm[9], but we include them as we cannot rule out the possibility that a DDR

deficiency protects from specific types of mutagenesis. Though not distinguishable in this study,

we suggest that negative coefficient features may derive in three ways: First, they may stem

from enhanced repair; second, they may stem from the decomposition of mutation counts into

signatures; and third, the mutated tumours may represent a subclass of patients in terms of age,

gender, or tumour subtype with specific mutational patterns.

Evaluating model performance

For each model, we evaluated the predictive performance using the area-under-the-receiver-

operating-curve (AUROC) score as well as the precision-recall area-under-the-curve (PR-AUC)

score. The PR-AUC score is a more robust measure for unbalanced data sets [28]; however, the

expected value for non-informative (unskilled) models equals the fraction of true positives and

thus varies between models. Therefore, we used the PR-AUC enrichment over the true-positive

rate (PR-AUC-E) as our selection criteria for predictive models. For the downstream analysis,

we included (shortlisted) models with PR-AUC-E that was substantial (>0.2; more than two

standard deviations above the mean across all 535 models) and significant (Benjamin-Hochberg

false discovery rate, FDR<0.05; Monte Carlo simulations) (**Methods**; **Additional file 1,**

Supplementary Table 7). In addition, we tested each model in the same cohort in the other

data set whenever one or more tumours had mono- or biallelic loss of the gene. Due to the

difference in biology between the two sets, we did not use this to explicitly evaluate the

predictive performance of models but have included the PR-AUC-E values from the tests in

Supplementary Table 6 and 7.

In the example of *BRCA2*-d in breast cancers of the HMF data set, our model achieved an

AUROC of 0.93 and a PR-AUC-E of 0.29 (**Fig. 2d,e; Additional file 1, Supplementary Table**

7). Out of 30,000 permuted LOF-sets, none had a similar or higher PR-AUC score and we

considered the model significant with a p-value < 3×10^{-5} (FDR adjusted q-value < 6×10^{-4}) (**Fig.**

2f,g). The model achieved a PR-AUC-E of 0.37 when tested on the PCAWG data set,

suggesting that the model may generalise across both metastatic and non-metastatic tumours.

This was further supported by the independent discovery of a similar model in the PCAWG data

set, which had an almost similar predictive power in the HMF data (PR-AUC-E = 0.19;

Supplementary Table 7). Notably, the *BRCA2*-d model did not include non-clustered deletions

<100Kb, SBS signature 3, and SBS signature 8, all features which have been associated with

BRCAness[9]. However, SBS signature 3 and non-clustered deletions 1-10Kb are included in the

model when the deletions at sites of microhomology are omitted from the input data set,

suggesting that they are excluded during feature selection due to high positive correlation with

the number of deletions at sites of microhomology among HMF breast cancers (Pearson corr. >

0.7; **Fig. 2h**; **Additional file 1, Supplementary Table 9**).

Our selection criteria resulted in 48 shortlisted predictive models across 24 DDR genes (**Fig. 3a;**

Additional file 1, Supplementary Table 7). As exemplified for *BRCA2*, each model is specified

by a set of predictive features representing mutational patterns associated with DDR gene LOF.

We divided the models into four groups based on aetiology and origin: models of *BRCA1/2*-d

(eight models of *BRCA2*-d and a single model of *BRCA1*-d; **Fig. 3b**); models of monoallelic

TP53-d (11 models; **Fig. 3c**); models of various monoallelic gene deficiencies derived from

colorectal cancer patients (eight models; **Fig. 3d**); and models including other DDR genes and

cancer types, including previously undescribed associations (20 models; **Fig. 3e**).

Recapitulation and predictive modelling of expected associations with *BRCA1/2*

deficiency

Five models predicted biallelic loss of *BRCA2* in cancers of the ovary, prostate, pancreas, and

breast. In addition, three models predicted *BRCA2* monoallelic loss in cancers of the pancreas,

breast, and prostate. Finally, we derived a single model of biallelic *BRCA1* loss in ovarian

cancer (**Fig. 3c; Fig. 4a**)**.** All models significantly outperformed their Monte-Carlo simulations

(q<0.05; Benjamini Hochberg FDR control) and had PR-AUC-E above 0.2 (**Fig. 4b,c**).

All *BRCA2*-d models were predominantly predicted by deletions at sites of microhomology,

consistent with the role of *BRCA2* in homologous recombination and suppression of

microhomology-mediated end joining[29]. Both clustered and non-clustered tandem duplications in

the range of 1 to 100Kb were included as features for various models, though with much smaller

predictive power. This agrees with what was identified for *BRCA2* deficient tumours in prior

studies[9,10]. The biallelic breast cancer model based on PCAWG further included SBS signature

3 as a predictive feature (**Fig. 4d**). Contrasting to the models of *BRCA2*-d, *BRCA1*-d in ovarian

cancer was exclusively associated with clustered and non-clustered tandem duplications (1-

10Kb; **Fig. 4d**). This aligns with prior studies[9,10]**,** which also found *BRCA1*-d to be closely

associated with a tandem-duplicator phenotype. In general, *BRCA1* and *BRCA2* were subject to

predominantly germline pathogenic events, and not a single deletion at a site of

microhomology, suggesting the expected forward causality (**Supplementary Table 8**). As for

the loss of *BRCA2*, the model for loss of *BRCA1* loss in ovary had sufficient predictive power

(PR-AUC-E = 0.3) in the other data set, suggesting that the model works independently of the

metastatic capacity of the tumour (**Supplementary Table 7**).

***TP53* deficiencies associate with increased numbers of structural variants**

We detected 11 predictive models (four based on PCAWG and seven on HMF) of monoallelic

TP53-d across cancers of the breast, skin, ovary, uterus, neuro-endocrine tissues (NET), biliary

gland, head and neck, pancreas, and the central nervous system (CNS) (**Fig. 4e**). These

predictive models performed with PR-AUC-E values ranging from 0.21 to 0.48 in breast and

biliary gland cancers, respectively. Similarly, AUROC values ranged from 0.48 to 0.88, again in

breast and biliary gland cancers (**Fig. 4f,g**). In line with existing literature[30], *TP53-d* is

associated with a significantly increased number of structural variants across the genome

(Wilcoxon rank-sum test; **Fig. 4h,i**), except in skin cancers. The models of *TP53* loss in head

and neck, skin, breast, and the biliary gland performed well (PR-AUC-E above 0.2) in the other

data set, suggesting that the predictive performance generalises independent of metastatic

tumour state (**Supplementary Table 7**).

Colorectal cancer models derived from hypermutated mismatch repair deficient tumours

In the HMF colorectal cancers, we discovered eight predictive gene deficiency models (*MSH3,*

SMC2, SMC6, BMPR2, CLASP2, SRCAP, *UBR5,* and *UVRAG)* of monoallelic LOF with PR-

AUC-E ranging from 0.21 (*CLASP2*) to 0.62 (*MSH3*) (AUROC ranging from 0.68 for *SRCAP*

deficiency to 0.94 for *MSH3* deficiency; **Fig. 4j,k,l**).

We suggest that the high number of models of monoallelic deficiencies may arise from spurious

LOF events in DDR genes in a subset of colorectal cancers that are hypermutated. Some

colorectal cancers are signified by mismatch repair deficiencies, such as LOF of *MSH3* or

MSH6, creating a high number of deletions in repetitive DNA[13,14]. Indeed, we found that this

pattern was most profound among the *MSH3* mutated cancers (**Fig. 4m,n**). Furthermore, we

found co-mutation with *MSH3* across the tumours underlying each model, ranging from 20%

(*SRCAP*) to 33% (*BMPR2*) of the mutated tumours (**Fig. 4p**). This suggests that the models

(except for the model of *MSH3*-d) might be the consequence of the hypermutator phenotype. In

other words, the causality may be reversed in these cases, and the mutational process driven

by *MSH3*-d may have caused the majority of their LOF events. This notion is supported by

investigating the features of the models. All eight models are characterised by a single, primary

predictive feature: Insertions (*SMC2, SMC6, BMPR2, UBR5, and UVRAG*), deletions in

repetitive DNA (*MSH3, SRCAP*), or deletions not flanked by repetitive or microhomologous DNA

(*CLASP2*) (**Fig. 4o**). Each of these features has a high positive correlation (Pearson corr. >

0.93) with the number of deletions in repetitive DNA. This correlation suggests that all eight

models relate to a genome-instability phenotype, which may be driven by the *MSH3* co-mutated

tumours or, potentially, a concurrent deficiency of other genes within the mismatch repair

system (**Additional file 1, Supplementary Table 9**). Notably, the deficiency models of *UBR5,*

BMPR2, CLASP2, and *SMC6* all had PR-AUC-E above 0.2 in the other data set, suggesting

that these genes are associated with the mismatch repair phenotype regardless of metastatic

state (**Supplementary Table 7**).

Biallelic LOF of *MSH6* associated with increased number of deletions in repetitive DNA in

prostate cancer

MSH6, a gene implicated in mismatch repair and microsatellite stability[14]**,** was mutated in both

alleles in 7 out of 342 HMF prostate cancer patients. We observed pathogenic indels in *MSH6* in

all 7 tumours, but only one of these in mono- or dinucleotide repeat DNA. *MSH6* deficiency

could be predicted with high accuracy (PR-AUC-E=0.25; AUROC=0.98) by an enrichment of

deletions in repetitive DNA (**Fig. 5a,b; Additional file 1, Supplementary Table 7**). This is

consistent with existing findings of mismatch repair deficiency and its presence in metastatic

prostate cancers[31].

High predictive power for *CDK12* deficiency

CDK12 encodes a kinase that regulates transcriptional and post-transcriptional processes of the

DDR[32–34]. We found that *CDK12*-d prostate cancers had an increased number of mid- and large-

sized tandem duplications 100Kb - 10Mb, compared to CDK12-WT (**Fig. 5c,d**). Several studies

have observed similar tandem duplication phenotypes in ovarian cancers [17,18,35] and castration

resistant prostate cancers[36,37]. In agreement, nine of the ten patients in our data set were

treated with drugs associated with castration resistance (4 Enzalutamide, 3 Abiraterone, 1

Cabazitaxel,1 Pembrolizumab)[38].

Whereas the tandem duplication phenotype has been previously associated with loss of *CDK12*,

in this study we present the first predictive algorithm utilising and quantifying the high predictive

value of these patterns (PR-AUC-E=0.73 and AUROC=0.97). Indeed, the loss of CDK12 has

been demonstrated to sensitise cancer cells to CHK1-[7] and PARP-inhibitors[39,40].

We went on to test the *CDK12*-d model across other cancer types (**Fig. 5c**). As expected, we

observed predictive power in cancers of the ovary and breast, though at a lower level (PR-AUC-

E=0.19 and AUROC=0.72). No predictive power was observed for the remaining cancer types.

We only observed a single tumour with ballelic LOF of *CDK12* in PCAWG, but the predictive

model was able to correctly identify this tumour, and reached a PR-AUC-E of 0.99 in the

PCAWG data set.

Novel predictive gene deficiency models

The shortlisted predictive LOF models further include biallelic LOF of *ATRX, PTEN, HERC2,*

MEN1, SMARCA4, BAP1, and *RB1,* as well as monoallelic LOF of *ATRX, IDH1, PTEN,*

CDKN2A, ARID1A, TP53BP1, HERC2, and *RB1* (**Fig. 3e; Additional file 1, Supplementary**

Table 7). The number of mutated tumours underlying each model varied from eight (biallelic

SMARCA-d in cancers of unknown primary) to 22 (monoallelic *HERC2*-d in skin). We here

present the models of *ATRX, IDH1, SMARCA4, CDKN2A, PTEN, and HERC2*; remaining

models may be found in the supplementary material (**Additional file 1, 3, 4, Supplementary**

Table 7, Supplementary Fig. 1).

Predictive models of *ATRX*-d and *IDH1*-d in CNS cancers

We found that *ATRX*-d (monoallelic: PR-AUC-E=0.21, AUROC=0.71; biallelic: PR-AUC-E=0.23,

AUROC=0.76; **Fig. 6a**) and *IDH1*-d (monoallelic: PR-AUC-E=0.24, AUROC=0.82; **Fig. 6b**) in

CNS cancers were both predicted by a decreased number of SBS signature 8 mutations. In

addition, *ATRX*-d was further predicted by non-clustered inv. 10-100Kb, although with a small

coefficient and hence contributing limited discriminatory power. We discovered that 7 of11 *IDH1*

mutated tumours were also *ATRX* mutated (28-fold enrichment compared to *IDH1* WT skin

cancers; p=1.2x10[-8], Fisher's exact test) and that both *ATRX* and *IDH1* were predominantly hit

by somatic mutations (14 of 15 for *ATRX*, 11 of 11 for *IDH1*).

Co-mutation between *IDH1* and *ATRX* is well-described in gliomas[41]. Interestingly, LOF of either

gene is associated with lack of SBS signature 8, a signature associated with BRCAness[9,42] and

late-replication errors[43]. This suggests that CNS cancers with *IDH1/ATRX* deficiency are not

subject to the same DNA lesions or repair processes as other CNS cancers and they may

potentially belong to a separate patient subclass, though we could not identify evidence of this.

Predictive model for *SMARCA4*-d in cancer of unknown primary

We discovered eight tumours (HMF) out of 77 with cancers of unknown primary with

SMARCA4-d (biallelic) that could be predicted with relatively high accuracy (PR-AUC-E=0.44;

AUROC=0.85; **Fig. 6c**). These tumours showed an enrichment of SBS signature 27 (a signature

first detected in myeloid cancers [44]), which has been suggested to be a sequencing artefact

though it also displays strong strand bias [21] (**Additional file 1, Supplementary Table 5**).

Among cancers of unknown primary, SBS signature 27 correlates strongly with SBS signature 4

(Pearson corr.=0.96), despite the signatures different composition (cosine similarity of

signatures = 0.17) (**Additional file 1, Supplementary Table 9; Fig. 6h,i**). This suggests that

SBS signature 4 may also be associated with *SMARCA4*-d and we indeed found that its

predictive performance (PR-AUC-E of 0.43; AUROC=0.83) was almost equivalent to SBS

signature 27. Signature 4 is associated with smoking across several cancer types[45,46];

interestingly, *SMARCA4*-d is seen in aggressive thoracic sarcomas[47] and strongly enriched

among patients with a history of smoking[48]. Its gene product, BRG1, has been suggested as a

lung cancer transcriptional regulator of genes that induce tumour proliferation[49] and

metastasis[50]. We evaluated the ability of SBS signature 4 to predict *SMARCA4*-d in other

cancer types and found a significant predictive association in lung cancer, though much lower

than for cancers of unknown primary (**Fig. 6j**). We also found a significant ability to predict

SMARCA4-d by the number of SBS signature 27 mutations in cancers of the NET (Wilcoxon

test, one-tailed p=0.002) and head and neck (p=4.8×10^{-7}), but we could not evaluate this in lung

cancers, as the signature is not among its set of cohort-specific signatures.

Twelve cancers had a high posterior probability of *SMARCA4*-d (**Fig. 6j**) despite being

SMARCA4 WT and lacking pathogenic events. No other single DDR gene was mutated among

all twelve tumours, with *TP53* having the most LOF events (6 of 12 cancers). Given that cancers

of unknown primary have disparate origins, this raises the possibility that the patients with high

posterior probability of *SMARCA4*-d may have metastasized from a cancer type or subtype with

both high levels of SBS signature 4 and 27, as well as high incidence of *SMARCA4*-d. Further

studies are thus needed to clarify if the observed associations can be explained through such

an ascertainment bias rather than causatively by *SMARCA4*-d.

Predictive model for monoallelic *CDKN2A*-d in skin cancer

Both germline and somatic variants in *CDKN2A* are known to predispose for melanoma[51]. In the

PCAWG skin cancer cohort, we found that a monoallelic predictive model of *CDKN2A*-d

achieved relatively high accuracy (PR-AUC-E=0.28; AUROC=0.82). Its predictive features are

enrichment of deletions at sites of microhomology, non-clustered inv. (100Kb-1Mb), and SBS

signature 7, in order of predictive importance (**Fig. 6d**). Apart from the inversions, these features

are also significantly associated with biallelic *CDKN2A*-d in the HMF skin cancer cohort

(**Additional file 4, Supplementary Fig. 2a,b**) and are included in the corresponding predictive

model, though it had lower predictive performance (PR-AUC-E=0.008; AUROC=0.542) and was

not shortlisted (**Additional file 1, Supplementary Table 6**). In both HMF and PCAWG, most of

the observed LOF events are somatic, with the majority being biallelic (16 of 18 in PCAWG; 32

of 33 in HMF; **Additional file 1, Supplementary Table 8**). The presence of deletions at sites of

microhomology suggests a possible reduction in error-free double-stranded break repair in

combination with an increased accumulation of SBS signature 7.

Predictive model for *HERC2-d* in metastatic skin cancer

HERC2-d has been associated with susceptibility to developing melanoma[52]. We found *HERC2*-

d predictable in HMF skin cancer patients (biallelic: PR-AUC-E=0.24, AUROC=0.73;

monoallelic: PR-AUC-E=0.23, AUROC=0.66; **Fig. 6e)**, primarily based on enrichment of

deletions in non-microhomologous and non-repetitive regions.

HERC2 encodes a protein ligase that modulates the activity of P53[53]. We observed that seven

of the nine tumours with biallelic *HERC2*-d also had a monoallelic pathogenic event in *TP53* (8-

fold enrichment; p=4.6x10^{-6}, Fisher's exact test). Tumours that are co-mutated in *TP53*

(monoallelic) and *HERC2* (mono- or biallelic) showed a significantly higher number of deletions

compared to tumours with LOF in either gene alone (Wilcoxon test, One-tailed p<0.031) and

cancers that are wild-type for both genes (p<7.2x10⁻⁷) (**Additional file 4, Supplementary Fig.**

2c).

***PTEN* deficiency associates with fewer structural variants in CNS and uterine cancers**

PTEN is a tumour suppressor gene found in various cancer types[54,55] and its deficiency was

found to be associated with mutational patterns in uterine and CNS cancers (**Additional file 1,**

Supplementary Table 7). In CNS, we acquired two identical models from the HMF data set, as

we observed no monoallelic events without a second hit (mono- and biallelic PR-AUC-E=0.37,

AUROC=0.74; **Fig. 6f**). In uterine cancer, we acquired significant predictive models from both

HMF and PCAWG (HMF, biallelic: PR-AUC-E=0.37, AUROC=0.74; HMF, monoallelic: PR-AUC-

E=0.36, AUROC=0.79; PCAWG, monoallelic: PR-AUC-E=0.22, AUROC=0.74; **Fig. 6g**). In

addition, the models of *PTEN* loss in uterus had predictive power in the other data sets,

suggesting signal robustness and independence of metastatic state (the PCAWG derived model

had PR-AUC-E of 0.55 in the HMF data; the HMF derived model had a PR-AUC-E of 0.27 in

PCAWG; **Supplementary Table 7**). The model of biallelic *PTEN*-d in uterine cancers is

primarily based on depletion of non-clustered inv. 10-100Kb, whereas both the HMF and

PCAWG models of monoallelic deficiency are primarily based on depletion of non-clustered

tandem duplications (10-100Kb) (**Fig. 6g**). In contrast, our models of *PTEN*-d in CNS cancers

from HMF were based on depletion of both non-clustered deletions 1-10Kb and non-clustered

translocations (**Fig. 6f**).

DISCUSSION

The cancer-specific, incomplete repair of endogenous and exogenous DNA lesions leave

specific genome-wide mutational patterns. Their detection provides potentially powerful

information on the fidelity of individual DNA repair pathways and response to chemo- and

immunotherapies. Taking advantage of a large pan-cancer data set, our analysis shows that

mutational patterns are associated with DNA repair defects across a wide range of cancers and

repair mechanisms. In this study, we have contributed concrete initial predictive algorithms for

mutational patterns of several DDR gene deficiencies, with potential use for clinical intervention.

The clinical scope of our approach is exemplified by recent regulatory approval for PARPi

administration, which was supported by statistical predictions of homologous recombination

deficiency (HRD) based on mutational patterns[56] We have similarly predicted *CDK12* deficiency

(*CDK12*-d) which has previously been associated with a tandem duplication phenotype– with

power similar to that of HRD detection. This suggests a clinical benefit of clinical application of

predictive algorithms to test for *CDK12*-d, specifically when considering the application of

CHK1-inhibitors in prostate, ovary, and breast cancer treatment[7].

We also find high predictive power for *MSH6*-d in prostate cancers by counting deletions in

repetitive DNA, also known as the microsatellite instability phenotype (MSI)[57]. It is common

practice to search for signs of MSI in colorectal cancers, endometrial cancers, and aggressive

prostate cancers. This prediction is most commonly made using a panel of repetitive DNA

regions, such as the Bethesda panel[58]. Here we demonstrated high predictive power in prostate

cancers based on mutation summary statistics, which may be routinely extracted from whole

genome sequencing. The correct identification of tumours with high MSI supports the

administration of immune checkpoint blockade treatment, as the mutational phenotype leads to

an increased expression of neo-peptides and thus an increased sensitivity towards the immune

response[59,60].

Our systematic approach identified a similar model of *PTEN* LOF in uterine cancer in the

PCAWG data and the HMF data; either model also having predictive power in the other data

set. The re-discovery in either dataset grants further trust to this model, and further experimental

investigation is warranted to understand the underlying aetiology of the mutation patterns, and

the potential for clinical benefits.

Recent studies have suggested the use of CDK4/6 inhibition in treating *SMARCA4* deficient

tumours[61]. We discovered that a large subset of cancers of unknown primary (>10%) have

SMARCA4 deficiency and that these cancers can be accurately predicted from their whole

genome accumulation of SBS signature 4 or SBS signature 27 mutations. This is a clinically

challenging cancer type due to the lack of a primary cancer to guide treatment and any such

prediction may serve in selecting treatment for these complex cancers, given additional

experimental evaluation.

Of note, the inclusion of monoallelic events allowed for larger sets of tumours with DRR gene

LOF, thereby increasing the power of our study when causative associations with mutational

patterns exist. This is for instance the case for *TP53,* which is known to be functionally impacted

by monoallelic events[62]. We shortlisted 11 models of TP53-d, which were all monoallelic. We

further found that inclusion of monoallelic LOF events matched the predictive power for *ATRX*

biallelic LOF in CNS cancers and *HERC2* biallelic LOF in skin cancers, suggesting that single

genetic hits are sufficient to affect repair and ensuing mutational patterns for these genes.

Incorrectly annotated LOF events in DDR genes may affect both the ability to discover

associations with mutational patterns and the performance of any associated predictive models.

On the other hand, highly stringent LOF criteria may result in true LOF events being overlooked,

and create small deficiency sets with insufficient power to discover true associations. For the

annotation of germline LOF events, we restricted our focus to rare variants (population

frequency ≤0.5%), as more common variants are likely benign. Given the high number of

variants evaluated overall, false positive germline LOF calls are expected. However, they are

unlikely to associate with specific mutational patterns and thus are unlikely to contribute

significant, shortlisted false positive predictive models of DDR deficiency. Likewise, in cancers

with hypermutator phenotypes, somatic LOF events may be caused by and associated with

specific mutational patterns. This leads to predictive models that in effect detect instances of

reverse causality, as discussed for the monoallelic models found in colorectal cancer. We

investigated whether reverse causality was a likely explanation for an association captured by a

predictive model by evaluating whether the annotated LOF events matched the predictive

mutational features (**Additional file 1, Supplementary Table 9**). In general, biallelic LOF

criteria is less sensitive to wrongly annotated LOF events than monoallelic criteria, as double

hits are rare compared to single hits. For both mono- and biallelic predictive models, the data

permutations ensure that the observed association between the DDR gene LOF events and

given mutational patterns are surprising.

Our conservative LOF curation, and a lack of expression or protein level data, is expected to

cause some LOF events to be missed. This may explain a high posterior probability of a

particular gene LOF in some tumours, but no evidence of genetic disruption (as seen in **Figure

5e** and **Figure 6h**). This highlights the scope of mutation-pattern-based predictions, in particular

in tumours without canonical DDR gene LOF events. At a very least, such tumours could be

considered for further scrutiny for LOF of the individual DDR gene that they are predicted to

have lost.

Our systematic approach provides a proof-of-concept that will become increasingly powerful as

the available data sets increase in number and size. For the current data sets, consistent

validation of detected associations was challenging due to small cohorts and differences in

cancer biology. To establish the basis for any future synthetic-lethality uses of our predictive

models, it would be desirable to establish causal relationships. This has been beyond the scope

of this study but could be achieved by whole-genome-sequencing of cell-lines or organoids with

individual DDR gene knockout. Alternatively, the loss of DDR genes can be explored in animal

models as previously done for several DDR genes including *BRCA1/2*[63], MMR genes[64,65], and

TP53[66].

The current study yields a catalogue of predictive models that captures both known and novel

associations between DDR gene deficiencies and mutational patterns. The included DDR genes

may provide targets for research and development of treatment. With further optimization,

predictive models such as these may guide the selection of therapy by adding certainty of

disabled or compromised repair deficiency phenotypes found in cancers of individual patients.

Materials and methods

Data

The analysis was conducted on 6,098 whole cancer genomes and included a set of 6,065 whole

cancer genomes after filtering (see below). The data came from two independent data sets: the

Pan-Cancer analysis of Whole Genomes (PCAWG; tumours = 2,583; ICGC study ID.

EGAS00001001692)[19] and the Hartwig Medical Foundation (HMF; tumours = 3,515; Acc. Nr.

DR-044)[20]. The data sets contain tumours from 32 cancer types and represent diverse patient

groups in terms of age, gender, and disease history. PCAWG consists of tumours both with and

without metastasis, whereas HMF exclusively consists of donors with tumours showing

metastatic capability. To best relate the two data sets, all tumours are catalogued by the site of

primary disease. For 77 of the metastatic HMF tumours, the primary site was unknown, and

these are annotated as such (**Figure 1a; Additional file 1, Supplementary Table 1**).

Curating samples

In the HMF data, we selected the earliest available sample whenever multiple samples existed

for the same patient. For the PCAWG data, we used the official whitelist, obtained at the ICGC

resources (https://dcc.icgc.org/pcawg), and for patients with more than one sample, we always

selected the earliest, whitelisted sample available. We discarded six PCAWG samples of

bone/soft tissue, as these were discarded in prior publications based on the PCAWG data set[27]

leaving 2,568 PCAWG samples and 3,497 HMF samples across 32 sites of the body. A list of all

Donor IDs, Sample IDs, and primary sites of disease may be found in **Additional file 1,**

Supplementary Table 1.

Curating variants for loss-of-function annotation

We annotated variants across a set of 736 genes related to the DNA damage response

(**Additional file 1, Supplementary Table 2**). The set of genes is combined from three sources,

Knijnenburg *et al.*[24]; Pearl *et al.*[23]; and Olivieri *et al.*[25].

We filtered the downloaded variant call files (VCF) of all samples for variants between the

Ensembl (GRCH37/hg19) start- and end position of each DDR gene (see coordinates in

Additional file 1, Supplementary Table 2). We included variants classified as PASS in the

VCF files and variants of the PCAWG data set supported by at least two of our four variant

callers. Furthermore, we discarded all variants which occurred in more than 200 samples across

the two data sets in order to avoid noise arising from single nucleotide polymorphisms (SNPs)

called as single nucleotide variants (SNVs), SNPs with high frequency in particular populations

and possibly technical artefacts. We also discarded somatic variants with variant allele fractions

(VAF) below 0.2 and variants where gnomAD (V2.1.1) showed a germline population frequency

above 0.5%[67].

Annotating pathogenic variants and mutations

We annotated all variants and mutations with CADD phred scores (V1.6) in order to separate

likely pathogenic variants from likely benign variants[26]. All variants with CADD phred scores of

25 or higher were considered pathogenic, whereas non-synonymous mutations with CADD

phred scores below 25 and above 10 were considered variants of unknown significance (VUS).

Variants with CADD phred scores below 10 were considered benign. A CADD phred score of 25

is a conservative threshold[68], and only includes the 0.3% most-likely pathogenic variants. In

addition, we annotated all variants with their status in the ClinVar database (when present).

Combining ClinVar and CADD phred scores ensured that we would be able to discover

associations across all DDR genes, not only genes with ClinVar annotations, which are

expected to be incomplete for any given gene.

Annotating loss of heterozygosity and deep deletions

We also used the copy number profiles of each sample to discover genes with biallelic or

monoallelic losses of parts of genes. Any overlap between a gene and a copy number loss was

indicated as a loss-of-function event, under the assumption that losing any part of the protein

coding DNA is detrimental to the complete protein product. We considered events in which the

minor allele copy number was below 0.2 to be a loss of heterozygosity (LOH), whereas we

considered events with a total tumour copy number below 0.3 (major and minor allele

summarised) as deep deletions. This cutoff is adapted from the work of Nguyen *et al*.[10].

Annotating biallelic and monoallelic gene hits

We considered genes with a single pathogenic variant, either somatic or germline, to be

monoallelic hit (n = 25,399). In cases where the monoallelic hit is accompanied by a LOH, we

considered the event a biallelic loss (n = 6,640). Finally, genes that were completely depleted in

a sample were considered biallelic lost (n = 66). We did not consider a single LOH event as a

monoallelic loss due to the broad impact and high frequency of such events, with ~25 times

higher rates of LOH than deep deletions (**Figure 1b,c**).

Mutational patterns of single base substitutions, indels and structural variants

We summarised the genome-wide set of somatic mutations by using Signature Tools Lib,

developed by Degaspari *et al.*[27] Firstly, we used Signature Tools Lib to count single nucleotide

variants by base change and context, and assigned these counts to a set of organ-specific

signatures, as recommended by Degaspari *et al*. We then have converted the organ-specific

signature exposures to reference signature exposures using the conversion matrix found in the

supplementary material of the paper by Degaspari *et al.*[27] For twelve signatures (N1-N12) this

was not possible, and these signatures are considered to be novel. Likewise, two signatures

associated with mismatch repair (MMR) deficiency, MMR1 and MMR2, are associated with

several of the COSMIC MMR signatures. For these, we preserved the signature label as given

by Signature Tools Lib. For cohorts with no organ-specific signatures (cancers of unknown

primary, neuroendocrine tissue, thymus, urinary tract, penis, testis, small intestine, vulva, double

primary, and the adrenal gland), we have assigned mutations directly to the full set of 30

reference signatures (COSMIC signatures 1-30). Note that we excluded age-related signature 1

from the modelling, as this signature confounded by acting as a proxy for the age of the patient

so that the model would be learning to differentiate old from young patients rather than patients

with activity of different mutational processes or repair deficiencies.

Secondly, Signature Tools Lib was used to count indels: insertions were simply counted,

whereas deletions were separated by the DNA context of the deletion, being microhomologous,

repetitive, or none of the two. Microhomology was defined by whether the deleted sequence

was similar to the region immediately 3' of the breakpoint, as this indicates repair by

microhomology-mediated end joining. Repetitive deletions were defined by whether there is a

repeat of the indel at the 3' end of the breakpoint[27]. We decided to not further compose indels

into signatures, for the ease of interpretation; the type of deletion has a clear relation to the

mechanism of repair, which is what we are investigating in this paper.

Finally, we used Signature Tools Lib to count structural variants. SVs were separated into two

groups, clustered and non-clustered based on the average distance between rearrangements

(the inter-rearrangement distance). Regions with at least 10 breakpoints having at least 10

times smaller inter-rearrangement distance than the average across the genome of that cancer,

were considered clusters[69]. SVs were further divided based on the type of mutation: Deletions,

tandem duplications, inversions, and translocations. Deletions, tandem duplications, and

inversions were further divided by size, with intervals being 1-10Kb, 10-100Kb, 100Kb - 1Mb, 1-

10Mb, and finally mutations with lengths above 10Mb. As with indels, we decided not to

decompose SV counts into rearrangement signatures, to ease interpretability and because SV

signatures are less established in the field. The per-sample exposure to each summary statistic

may be seen in **Additional file 1, Supplementary Table 3** (log transformed and scaled in

Additional file 1, Supplementary Table 4; see method below) and the suggested aetiology

(when existing) may be seen in **Additional file 1, Supplementary Table 5**[21,42].

Preparing subsets of data for modelling

We separated the two data sets and divided the tumours into their respective cancer-type

cohorts. We stratified the data for each DDR gene; designating whether the patient had a

biallelic pathogenic variant, monoallelic pathogenic variant, or no pathogenic variants. We

excluded tumours with inconclusive variants (CADD phred >10, <25) within the gene. For

modelling of cancers with biallelic variants in a gene, we selected all DDR gene-cohort

combinations where more than five tumours had biallelic variants in the same gene (n=194.

Likewise, we selected all cases with more than ten tumours with monoallelic variants (n = 341)

(**Additional file 1, Supplementary Table 6**). This setup means that each sample may be

included in developing several models but occur only once in each model as either mutated or

non-mutated. Biallelic mutated tumours were also included in the models of monoallelic loss, as

we consider biallelic loss a special case of monoallelic loss.

Generating models using LASSO regression

For each model we calculated per-sample weights to counter the imbalance in the data, so that

the weight of each sample was one minus the proportion of tumours with this mutational status

(either pathogenic or non-pahogenic/wildtype). We then used logistic regression model with

least absolute shrinkage and selection operator (LASSO) regularisation (henceforth LASSO

regression), with the alpha parameter at 1, from the R-package "glmnet" (v4.0)[70] to select

features with predictive power. The LASSO regression was selected to achieve a sparse set of

associated features that could be readily interpreted. This is in line with the methodology used in

HRDetect by Davies *et al.*[9].

For each of k cross-fold validation sets (with k being the number of biallelic or monoallelic hit

tumours, respectively), we ran a LASSO regression with a nested 5-fold cross-validation and the

assigned weights (as mentioned above to counter imbalance in the data), to produce a set of

lambda values. From the LASSO regression we selected the lambda value corresponding to the

binomial deviance which is one standard deviation away from the observed minimal binomial

deviance of the regression, in order to avoid overfitting (overfitting may occur if taking the

lambda of the minimal binomial deviance)[70]. In a few cases, the LASSO regression converged

to larger sets of features than what is justifiable by the low number of mutated tumours, and so

we limited the number of features to one feature per ten mutated tumours, rounded up and

added one (e.g. 12 mutated tumours gives basis for maximum three features). For each of the k

cross-folds we used the model to predict the left-out data and used these predictions for the

evaluation of the model performance.

Finally, we generated a model by including the entire data set. This was done to train the best

possible model in terms of selected features and coefficients, and this is the model reported in

the main figures, whereas the performance measures shown in the main figures are derived

from the k-fold cross-validation.

Evaluating model predictive performance

We used k-fold cross validation to get a measure of the predictive performance of each model.

Due to the strong imbalance in the data, we use the precision-recall area-under-the-curve (PR-

AUC) enrichment from the true-positive rate (PR-AUC-E) as our statistic of performance, while

also reporting traditional AUROC scores. The true-positive rate is the baseline that would be

expected for a non-informed model. By subtracting the true-positive rate from the PR-AUC of

each model, we can compare the performance between models despite different sample sizes.

Model selection

We selected models for further analysis (our shortlist) based on both their PR-AUC-E

performance (effect size) and their significance. Across the 535 initial models, we observed a

mean PR-AUC-E of 0.04 with a standard deviation of 0.099 (90%-quantile = 0.18) (**Additional**

file 1, Supplementary Table 6). The PR-AUC-E threshold was set at 0.2 and thus roughly two

standard deviations above the overall mean (**Additional file 1, Supplementary Table 7**).

To evaluate significance, we generated a null-distribution for each of the 535 initial models. This

was done by permuting the mutation state of the underlying tumours and then running the

LASSO regression on the permuted data. We ran a minimum of 10,000 permutations (n) for

each of the 535 models, storing the PR-AUC-E for each permutation. For models where the

number (r) of permutations leading to PR-AUC-E values as or more extreme than the original

model was smaller than five, we ran an additional 10,000 permutations, up to a maximum of

30,000 permutations. A p-value (p=$(r + 1)/n$) was calculated for each model (**Additional file 1,**

Supplementary Table 6). The Benjamini-Hochberg procedure was used to control the false

discovery rate and resulting adjusted p-values (q-values) smaller than 0.05 were considered

significant (**Additional file 1, Supplementary Table 7**).

In conclusion, 48 models of 24 genes were shortlisted for further investigation. Of these, we

describe 43 in the main text and, more briefly, 5 in **Supplementary text** and **Supplementary**

figure 1. The inclusion into the main manuscript was done by collegial evaluation of interest to

the field and does not represent a definitive difference in quality of the models.

DECLARATIONS

Ethics approval and consent to participate

We analysed data generated and made available by the Pan-Cancer Analysis of Whole

Genomes (PCAWG)[19] Consortium of the International Cancer Genome Consortium (ICGC) and

The Cancer Genome Atlas (TCGA) as well as the Hartwig Medical Foundation[20]. The research

conforms to the principles of the Helsinki Declaration.

Availability of Data and Materials

The data sets supporting the conclusions of this article are from the Pan-Cancer Analysis of

Whole Genomes and from the Hartwig Medical Foundation. The public parts of the PCAWG

data set is available at https://dcc.icgc.org/releases/PCAWG, whereas controlled files may be

accessed through gbGaP and daco, as instructed on this site https://docs.icgc.org/pcawg/data/.

The ICGC study ID of the project is EGAS00001001692.

The HMF data used in this project may be found by accession code DR-044 and can be

obtained upon request at the Hartwig Medical Foundation

(https://www.hartwigmedicalfoundation.nl/en).

Code availability

The code needed to reproduce the analysis will be made available at

https://github.com/SimonGrund/DDR_Predict , including data pre-processing, modelling of the

535 predictive models and modelling of ≥10,000 data permutation null models for each of the

535 models.

Competing interests

The authors declare no competing interests.

Funding

ERH and AS were funded by the Novo Nordisk Foundation (NNF15OC0016662), Cancer

Research UK (C23210/A7574), and the Danish National Research Foundation (Center grant,

DNRF115). JSP, SGS, GAP, and MHC were funded by the Independent Research Fund

Denmark | Medical Sciences (8021-00419B), the Danish Cancer Society (R307-A17932),

Aarhus University Research Foundation (AUFF-E-2020-6-14), a PhD stipend from Aarhus

University, and a stipend from Research Foundation of Central Region Denmark (A2972).

Author contributions

The project was conceived and designed by JSP and SGS, with early contributions by JB. The

data analysis was carried out by SGS under supervision of JSP with input from MHC, including

data preparation, exploratory analysis, and statistical modelling. SGS generated all figures and

tables. SGS, AS, GAP, ERH, and JSP interpreted the results. SGS, ERH and JSP wrote the

manuscript. All authors accepted the manuscript.

Acknowledgements

We thank the Pan-Cancer Analysis of Whole Genomes (PCAWG), The International Cancer

Genome Consortium (ICGC), and The Cancer Genome Atlas (TCGA) for access to whole

cancer genomes. We would also like to thank the Hartwig Medical Foundation (HMF) and the

Center for Personalized Cancer Treatment (CPCT) for creating and providing access to

metastatic whole cancer genome data. We thank Jenny Gruhn and Amy V. Kaucher for valuable

comments on the manuscript. We thank Dr. Andrea Degaspari for aid in using Signature Tools

Lib for summarising mutation patterns.

REFERENCE LIST

1. Lindahl, T. Instability and decay of the primary structure of DNA. *Nature* **362**, 709–715

(1993).

2. Nik-Zainal, S. *et al.* Mutational Processes Molding the Genomes of 21 Breast Cancers. *Cell*

vol. 149 979–993 (2012).

3. Volkova, N. V. *et al.* Mutational signatures are jointly shaped by DNA damage and repair.

Nat. Commun. **11**, 2169 (2020).

4. Bryant, H. E. *et al.* Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-

ribose) polymerase. *Nature* vol. 447 346–346 (2007).

5. Le, D. T. *et al.* PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N. Engl. J. Med.*

372, 2509–2520 (2015).

6. von Bueren, A. O. *et al.* Mismatch repair deficiency: a temozolomide resistance factor in

medulloblastoma cell lines that is uncommon in primary medulloblastoma tumours. *Br. J.*

Cancer **107**, 1399–1408 (2012).

7. Paculová, H. *et al.* BRCA1 or CDK12 loss sensitizes cells to CHK1 inhibitors. *Tumour Biol.*

39, 1010428317727479 (2017).

8. Landrum, M. J. *et al.* ClinVar: public archive of relationships among sequence variation and

human phenotype. *Nucleic Acids Res.* **42**, D980–5 (2014).

9. Davies, H. *et al.* HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on

mutational signatures. *Nat. Med.* **23**, 517–525 (2017).

10. Nguyen, L., W M Martens, J., Van Hoeck, A. & Cuppen, E. Pan-cancer landscape of

homologous recombination deficiency. *Nat. Commun.* **11**, 5584 (2020).

11. McVey, M. & Lee, S. E. MMEJ repair of double-strand breaks (director's cut): deleted

sequences and alternative endings. *Trends Genet.* **24**, 529–538 (2008).

12. Nussenzweig, A. & Nussenzweig, M. C. A backup DNA repair pathway moves to the

forefront. *Cell* **131**, 223–225 (2007).

13. Umar, A. *et al.* Defective mismatch repair in extracts of colorectal and endometrial cancer

cell lines exhibiting microsatellite instability. *J. Biol. Chem.* **269**, 14367–14370 (1994).

14. Edelmann, W. *et al.* The DNA mismatch repair genes Msh3 and Msh6 cooperate in

intestinal tumor suppression. *Cancer Res.* **60**, 803–807 (2000).

15. Lanni, J. S. & Jacks, T. Characterization of the p53-dependent postmitotic checkpoint

following spindle disruption. *Mol. Cell. Biol.* **18**, 1055–1064 (1998).

16. Gorgoulis, V. G. *et al.* Activation of the DNA damage checkpoint and genomic instability in

human precancerous lesions. *Nature* **434**, 907–913 (2005).

17. Popova, T. *et al.* Ovarian Cancers Harboring Inactivating Mutations in CDK12 Display a

Distinct Genomic Instability Pattern Characterized by Large Tandem Duplications. *Cancer*

Res. **76**, 1882–1891 (2016).

18. Menghi, F. *et al.* The Tandem Duplicator Phenotype Is a Prevalent Genome-Wide Cancer

Configuration Driven by Distinct Gene Mutations. *Cancer Cell* **34**, 197–210.e5 (2018).

19. ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium. Pan-cancer analysis of

whole genomes. *Nature* **578**, 82–93 (2020).

20. Priestley, P. *et al.* Pan-cancer whole-genome analyses of metastatic solid tumours. *Nature*

575, 210–216 (2019).

21. Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids*

Res. **47**, D941–D947 (2019).

22. Alexandrov, L. B. & Stratton, M. R. Mutational signatures: the patterns of somatic mutations

hidden in cancer genomes. *Curr. Opin. Genet. Dev.* **24**, 52–60 (2014).

23. Pearl, L. H., Schierz, A. C., Ward, S. E., Al-Lazikani, B. & Pearl, F. M. G. Therapeutic

opportunities within the DNA damage response. *Nat. Rev. Cancer* **15**, 166–180 (2015).

24. Knijnenburg, T. A. *et al.* Genomic and Molecular Landscape of DNA Damage Repair

Deficiency across The Cancer Genome Atlas. *Cell Rep.* **23**, 239–254.e6 (2018).

25. Olivieri, M. *et al.* A Genetic Map of the Response to DNA Damage in Human Cells. *Cell*

182, 481–496.e21 (2020).

26. Rentzsch, P., Witten, D., Cooper, G. M., Shendure, J. & Kircher, M. CADD: predicting the

deleteriousness of variants throughout the human genome. *Nucleic Acids Research* vol. 47

D886–D894 (2019).

27. Degasperi, A. *et al.* A practical framework and online tool for mutational signature analyses

show inter-tissue variation and driver dependencies. *Nat Cancer* **1**, 249–263 (2020).

28. Davis, J. & Goadrich, M. The relationship between Precision-Recall and ROC curves. in

Proceedings of the 23rd international conference on Machine learning 233–240

(Association for Computing Machinery, 2006).

29. Ceccaldi, R. *et al.* Homologous-recombination-deficient tumours are dependent on Polθ-

mediated repair. *Nature* **518**, 258–262 (2015).

30. Hanel, W. & Moll, U. M. Links between mutant p53 and genomic instability. *J. Cell.*

Biochem. **113**, 433–439 (2012).

31. Graham, L. S. *et al.* Mismatch repair deficiency in metastatic prostate cancer: Response to

PD-1 blockade and standard therapies. *PLoS One* **15**, e0233260 (2020).

32. Blazek, D. *et al.* The Cyclin K/Cdk12 complex maintains genomic stability via regulation of

expression of DNA damage response genes. *Genes Dev.* **25**, 2158–2172 (2011).

33. Marqués, F. *et al.* A new subfamily of high molecular mass CDC2-related kinases with

PITAI/VRE motifs. *Biochem. Biophys. Res. Commun.* **279**, 832–837 (2000).

34. Li, X., Chatterjee, N., Spirohn, K., Boutros, M. & Bohmann, D. Cdk12 Is A Gene-Selective

RNA Polymerase II Kinase That Regulates a Subset of the Transcriptome, Including Nrf2

Target Genes. *Sci. Rep.* **6**, 21455 (2016).

35. Li, Y. *et al.* Patterns of somatic structural variation in human cancer genomes. *Nature* **578**,

112–121 (2020).

36. Wu, Y.-M. *et al.* Inactivation of CDK12 Delineates a Distinct Immunogenic Class of

Advanced Prostate Cancer. *Cell* vol. 173 1770–1782.e14 (2018).

37. Rescigno, P. *et al.* Characterizing CDK12-Mutated Prostate Cancers. *Clinical cancer*

research: an official journal of the American Association for Cancer Research vol. 27 566–

574 (2021).

38. Sumanasuriya, S. & De Bono, J. Treatment of Advanced Prostate Cancer—A Review of

Current Therapies and Future Promise. *Cold Spring Harb. Perspect. Med.* **8**, (2018).

39. Bajrami, I. *et al.* Genome-wide profiling of genetic synthetic lethality identifies CDK12 as a

novel determinant of PARP1/2 inhibitor sensitivity. *Cancer Res.* **74**, 287–297 (2014).

40. Joshi, P. M., Sutor, S. L., Huntoon, C. J. & Karnitz, L. M. Ovarian cancer-associated

mutations disable catalytic activity of CDK12, a kinase that promotes homologous

recombination repair and resistance to cisplatin and poly(ADP-ribose) polymerase

inhibitors. *J. Biol. Chem.* **289**, 9247–9253 (2014).

41. Mukherjee, J. *et al.* Mutant IDH1 Cooperates with ATRX Loss to Drive the Alternative

Lengthening of Telomere Phenotype in Glioma. *Cancer Res.* **78**, 2966–2977 (2018).

42. Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature*

578, 94–101 (2020).

43. Singh, V. K., Rastogi, A., Hu, X., Wang, Y. & De, S. Mutational signature SBS8

predominantly arises due to late replication errors in cancer. *Commun Biol* **3**, 421 (2020).

44. Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic cells. *Nat. Genet.*

47, 1402–1407 (2015).

45. Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* vol.

500 415–421 (2013).

46. Nik-Zainal, S. *et al.* The genome as a record of environmental exposure. *Mutagenesis*

gev073 (2015) doi:10.1093/mutage/gev073.

47.   Sauter, J. L. *et al.* SMARCA4-deficient thoracic sarcoma: a distinctive clinicopathological

entity with undifferentiated rhabdoid morphology and aggressive behavior. *Mod. Pathol.* **30**,

1422–1432 (2017).

48.   Rekhtman, N. *et al.* SMARCA4-Deficient Thoracic Sarcomatoid Tumors Represent

Primarily Smoking-Related Undifferentiated Carcinomas Rather Than Primary Thoracic

Sarcomas. *J. Thorac. Oncol.* **15**, 231–247 (2020).

49.   Dagogo-Jack, I. *et al.* Clinicopathologic Characteristics of BRG1-Deficient NSCLC. *J.*

Thorac. Oncol. **15**, 766–776 (2020).

50.   Concepcion, C. P. *et al.* SMARCA4 inactivation promotes lineage-specific transformation

and early metastatic features in the lung. *Cancer Discov.* (2021) doi:10.1158/2159-

8290.CD-21-0248.

51.   Liu, L. *et al.* Mutation of the CDKN2A 5'UTR creates an aberrant initiation codon and

predisposes to melanoma. *Nat. Genet.* **21**, 128–132 (1999).

52.   Amos, C. I. *et al.* Genome-wide association study identifies novel loci predisposing to

cutaneous melanoma. *Hum. Mol. Genet.* **20**, 5012–5023 (2011).

53.   Cubillos-Rojas, M. *et al.* The E3 ubiquitin protein ligase HERC2 modulates the activity of

tumor protein p53 by regulating its oligomerization. *J. Biol. Chem.* **289**, 14782–14795

(2014).

54.   Li, J. *et al.* PTEN, a putative protein tyrosine phosphatase gene mutated in human brain,

breast, and prostate cancer. *Science* **275**, 1943–1947 (1997).

55.   Liaw, D. *et al.* Germline mutations of the PTEN gene in Cowden disease, an inherited

breast and thyroid cancer syndrome. *Nat. Genet.* **16**, 64–67 (1997).

56.   Fda. FDA approves niraparib for HRD-positive advanced ovarian cancer. (2019).

57.   Boland, C. R. & Goel, A. Microsatellite instability in colorectal cancer. *Gastroenterology*

138, 2073–2087.e3 (2010).

58. Umar, A. *et al.* Revised Bethesda Guidelines for hereditary nonpolyposis colorectal cancer

(Lynch syndrome) and microsatellite instability. *J. Natl. Cancer Inst.* **96**, 261–268 (2004).

59. Abida, W. *et al.* Analysis of the Prevalence of Microsatellite Instability in Prostate Cancer

and Response to Immune Checkpoint Blockade. *JAMA Oncol* **5**, 471–478 (2019).

60. Antonarakis, E. S. *et al.* Clinical Features and Therapeutic Outcomes in Men with

Advanced Prostate Cancer and DNA Mismatch Repair Gene Mutations. *Eur. Urol.* **75**, 378–

382 (2019).

61. Xue, Y. *et al.* SMARCA4 loss is synthetic lethal with CDK4/6 inhibition in non-small cell lung

cancer. *Nat. Commun.* **10**, 557 (2019).

62. Malcikova, J. *et al.* Monoallelic and biallelic inactivation of TP53 gene in chronic

lymphocytic leukemia: selection, impact on survival, and response to DNA damage. *Blood*

114, 5307–5314 (2009).

63. Evers, B. & Jonkers, J. Mouse models of BRCA1 and BRCA2 deficiency: past lessons,

current understanding and future prospects. *Oncogene* **25**, 5885–5897 (2006).

64. Reitmair, A. H. *et al.* MSH2 deficient mice are viable and susceptible to lymphoid tumours.

Nat. Genet. **11**, 64–70 (1995).

65. Prolla, T. A. *et al.* Tumour susceptibility and spontaneous mutation in mice deficient in

Mlh1, Pms1 and Pms2 DMA mismatch repair. *Nat. Genet.* **18**, 276–279 (1998).

66. Donehower, L. A. The p53-deficient mouse: a model for basic and applied cancer studies.

Semin. Cancer Biol. **7**, 269–278 (1996).

67. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in

141,456 humans. *Nature* **581**, 434–443 (2020).

68. Itan, Y. *et al.* The mutation significance cutoff: gene-level thresholds for variant predictions.

Nat. Methods **13**, 109–110 (2016).

69. Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome

sequences. *Nature* **534**, 47–54 (2016).

70. Friedman, J., Hastie, T. & Tibshirani, R. Regularization paths for generalized linear models

via coordinate descent. *J. Stat. Softw.* (2010).

Figure titles and legends

Figure 1: Cancer-types, DDR gene deficiencies, and mutational patterns

a Cohort sizes for the 32 cancer types comprising the 6,065 whole cancer genomes collected

from the Hartwig Medical Foundation (HMF; n=3,497) and the PanCancer Analysis of Whole

Genomes (PCAWG; n=2,568). **b** mono- and biallelic loss-of-function (LOF) events were

annotated across 736 DDR genes based on both pathogenic variants and copy-number-losses

(loss of heterozygosity; LOH), overall and (**c**) per patient (**d**) with varying no. of LOF events per

DDR gene (x-axis; logarithmic). **e** Whole genome mutational patterns were represented as

summary statistics and used as input features for the predictive models of DDR gene deficiency.

Concretely, each patient was annotated with the number of single-base-substitutions (SBS) that

are accounted to each SBS signature (Alexandrov *et al.*[22]; Degaspari *et al.*[27]), number of indels

divided by context (mh = microhomology; rep = repetitive), and (**f**) number of structural variants

divided by clusterness, size, and type (del = deletion; inv = inversion; tds = tandem duplication;

trans = translocation)

Figure 2: Predictive modelling of BRCA2 deficiencies in the HMF breast cancers

a Mutational status of *BRCA2* across 645 HMF breast cancer patients. **b** Mutational summary

statistics for the HMF breast cancer patients divided by biallelic *BRCA2* loss-of-function (LOF;

red) and *BRCA2* wild type (WT; grey) (selected predictive features in bold). **c** Predictive features

and their coefficients for model of biallelic *BRCA2* loss with predictive performance measured in

(**d**) AUROC and (**e**) PR-AUC (PR-AUC-E=PR-AUC - baseline=0.29; **Methods**). **f** Distributions of

AUROC and (**g**) PR-AUC-E values obtained from 30,000 random data permutations compared

to observed values (punctuated lines). **h** Correlation between selected predictive features

(horizontal) and other highly correlated (Pearson corr. >0.65) mutational features (vertical).

Figure 3: Predictive models of DDR gene deficiencies

a The precision-recall AUC enrichment PR-AUC-E; x-axis) and significance (FDR; logarithmic y-axis) of the 535 predictive models (one model per gene with more than five biallelic or more

than ten tumours either mono- or biallelic mutated in either HMF or PCAWG in any one cancer

type; **Methods**). Significance (q-value representing FDR) evaluated by counting equally or

more-extreme PR-AUC-E values across >10,000 permuted data sets and applying Benjamini-

Hochberg FDR control. Models with FDR below 0.05 and PR-AUC-E above 0.2 are shortlisted

(**Methods**). **b** Shortlisted predictive models of deficiency of *BRCA1* or *BRCA2*; (**c**) *TP53*

monoallelic predictive models; (**d**) monoallelic gene deficiency models across colorectal cancer

patients; (**e**) and remaining gene deficiency models not contained in the other sub-groups.

Numbers indicate the number of mutated out of the total number of tumours included in the

development of each model.

Figure 4: Predictive models with anticipated aetiology or origin

a Overview of predictive models for *BRCA1*-d and *BRCA2*-d, showing data source, type of

model, and LOF-set statistics (**b**) PR-AUC-E, (**c**) AUROC, (**d**) and the predictive features and

their coefficient for individual models. **e,f,g** Overview of predictive models of *TP53*-d (as in a-c).

h For each cohort the number of structural variants (x-axis; logarithmic) for *TP53* LOF tumours

(red) versus *TP53* wild type tumours (grey) and (**i**) the significance of their difference (two-sided

Wilcoxon rank-sum test). **j,k,l** Predictive models of gene deficiencies in colorectal cancers (as in

a-c). **m** Number of deletions in repetitive DNA (as in h) and (**n**) its significance (as in i)**. o** The

predictive features of each model (as in d) and (**p**) the percentage of tumours that are co-

mutated with *MSH3*.

Figure 5

Figure 5: *CDK12* mutated prostate tumours are predicted by tandem duplications

a Biallelic predictive model *MSH6*-d in HMF prostate tumours and its PR-AUC-E and AUROC. **b**

Boxplots of mutation counts between tumours that are *MSH6*-d loss-of-function (red) and *MSH6*

wild-type (grey), (mutation counts are normalised and log-transformed; **Methods). c** Biallelic

predictive model for *CDK12*-d with performance measures (as in a). **b** Boxplots of mutation

counts for *CDK12* deficient and wildtype tumours (as in b). **e** *CDK12*-d predictive performance

for different cancer types. Predicted probability of *CDK12*-d (x-axis) for tumours with *CDK12*

LOF (red) and *CDK12* wild type (grey) are shown for prostate cancer, breast and ovary cancer,

and all other cancer types (y-axis).

Figure 6

Figure 6: Novel predictive models of DDR gene deficiencies

a Predictive model of *ATRX*-d and its PR-AUC-E, AUROC, and selected features and their

coefficients. Same information for predictive models of (**b)** *IDH1*-d*,* (**c**) *SMARCA4*-d, (**d**)

CDKN2A-d, (**e**) *HERC2*-d, and (**f**) *PTEN*-d in CNS cancers and (**g**) uterine cancers*.*

h Number of SBS sig. 27 and SBS sig. 4 (y-axis; logarithmic) among tumours of unknown

primary with *SMARCA4* biallelic LOF (red) or wild-type (grey). **i** Pearson correlation between the

per-tumour number (tumours of unknown primary; HMF) of SBS signature 27 (y-axis) and SBS

signature 4 (x-axis; logarithmic) mutations, with an overlaid linear model (blue) and its 95%

confidence interval (grey). **j** Using a model trained to predict *SMARCA4* biallelic LOF in HMF

cancers of unknown primary, we evaluate the predictive power across cohorts (one-tailed

Wilcoxon rank-sum test) displaying significant cohorts separately (colours as in **a**).

Additional files

Additional file 1: Supplementary tables 1,2 and 4-9; Microsoft Excel (xlsx)

Additional file 1, Supplementary Table 1: All included tumours and their primary tumour

locations

Additional file 1, Supplementary Table 2: 736 DDR genes, hg19 coordinates and the number of

pathogenic events across 6,065 cancer genomes

Additional file 1, Supplementary Table 5: Proposed Etiologies of base substitution signatures

Additional file 1, Supplementary Table 6: All models (n=535)

Additional file 1, Supplementary Table 7: Shortlisted models (n=48)

Additional file 1, Supplementary Table 8: Pathogenic events in each of the 535 LOF-sets

Additional file 1, Supplementary Table 9: Correlation between features in shortlisted models

Additional file 2: Supplementary table 3-4; zip-compressed; tab-separated values (.tsv), may be

opened in Microsoft Excel

Additional file 1, Supplementary Table 3: All SBS signature contributions, indels counts, and SV

counts, per sample

Additional file 1, Supplementary Table 4: All SBS signature contributions, indels counts, and SV

counts, per sample, log-transformed and scaled to z-scores

1002

1003   Additional file 3: Supplementary text; pdf (.pdf)

1004   Supplementary Text: Significant models that are not described in the main text

1005

1006   Additional file 4: Supplementary figures; pdf (.pdf)

1007   Supplementary Fig. 1: Shortlisted predictive models of DDR gene deficiencies not included in

1008   the main manuscript

1009   Supplementary Fig. 2: *CDKN2A*-d in PCAWG and HMF skin cancers, *HERC2*-d in HMF skin

1010   cancers, and *ARID1A*-d in PCAWG prostate cancers