ANTIPSYCHOTIC DRUGS SELECTIVELY DECORRELATE LONG-RANGE INTERACTIONS

IN DEEP CORTICAL LAYERS

Matthias Heindorf[1] & Georg B. Keller[1,2,3]

[1]*Friedrich Miescher Institute for Biomedical Research, Basel, Switzerland*
[2]*Faculty of Science, University of Basel, Basel, Switzerland*
[3]*Lead contact, correspondence to: georg.keller@fmi.ch*

Psychosis is characterized by a diminished ability of the brain to distinguish externally driven

activity patterns from self-generated activity patterns. Antipsychotic drugs are a class of small

molecules with relatively broad binding affinity for a variety of neuromodulator receptors that, in

humans, can prevent or ameliorate psychosis. How these drugs influence the function of cortical

circuits, and in particular their ability to distinguish between externally and self-generated activity

patterns, is still largely unclear. To have experimental control over self-generated sensory

feedback we used a virtual reality environment in which the coupling between movement and

visual feedback can be altered. We then used widefield calcium imaging to determine the cell type

specific functional effects of antipsychotic drugs in mouse dorsal cortex under different conditions

of visuomotor coupling. By comparing cell type specific activation patterns between locomotion

onsets that were experimentally coupled to self-generated visual feedback and locomotion onsets

that were not coupled, we show that deep cortical layers were differentially activated in these

two conditions. We then show that the antipsychotic drug clozapine disrupted visuomotor

integration at locomotion onsets also primarily in deep cortical layers. Given that one of the key

components of visuomotor integration in cortex is long-range cortico-cortical connections, we

tested whether the effect of clozapine was detectable in the correlation structure of activity

patterns across dorsal cortex. We found that clozapine as well as two other antipsychotic drugs,

aripiprazole and haloperidol, resulted in a strong reduction in correlations of layer 5 activity

between cortical areas and impaired the spread of visuomotor prediction errors generated in

visual cortex. Our results are consistent with the interpretation that a major functional effect of

antipsychotic drugs is a selective alteration of long-range layer 5 mediated communication.

INTRODUCTION

Under normal conditions, self-generated movements are coupled to sensory feedback in different

modalities. Locomotion, for example, is coupled to proprioceptive feedback from muscle

movements, somatosensory and auditory feedback from touching the ground, vestibular feedback

from translational acceleration, and visual feedback in the form of expanding visual flow. This

coupling is key to our brain's ability to learn internal models of the world (Jordan and Rumelhart,

1992). Internal models are the transformations between cortical areas and can be used to predict

the sensory consequences of movement. Hallucinations and delusions can be explained as a

misconfiguration of these internal models: inaccurate predictions of the sensory consequences of

movements can result in self-generated activity that is not correctly distinguished from externally

driven activity (Fletcher and Frith, 2009; Frith, 2005).

Self-generated movements result in broad activation of most regions of the brain (Musall et al.,

2019; Stringer et al., 2019), including some primary sensory areas like visual cortex (Keller et al.,

2012; Saleem et al., 2013) where they have been shown to depend on visual context (Pakan et al.,

2016). The framework of predictive processing postulates that in cortex, these movement related

signals are internal model based predictions of the sensory consequences of movement that are

compared to externally generated bottom-up input to compute prediction errors (Jiang and Rao,

2021; Keller and Mrsic-Flogel, 2018). Evidence for this interpretation has mainly come from the

discovery of movement related prediction error responses in a variety of different cortical regions

and species (Attinger et al., 2017; Audette et al., 2021; Ayaz et al., 2019; Eliades and Wang, 2008;

Heindorf et al., 2018; Keller et al., 2012; Keller and Hahnloser, 2009; Stanley and Miall, 2007; Zmarz

and Keller, 2016), and the observation that top-down inputs from motor areas of cortex appear to

convey movement related predictions of sensory input to sensory areas of cortex (Leinweber et al.,

2017; Schneider et al., 2018). The neurons on which the opposing bottom-up input and top-down

prediction converge are referred to as prediction error neurons and are thought to reside primarily

in layer 2/3 (L2/3) of cortex (Jordan and Keller, 2020). To complete the circuit, there also needs to be

a population of neurons that integrate over prediction error responses. These are referred to as

internal representation neurons. It is still unclear which cortical neuron type functions as internal

representation neurons, but based on anatomical arguments it has been speculated that they can be

found in deep cortical layers (Bastos et al., 2012).

In humans, externally driven and self-generated activity patterns can be experimentally separated at

least partially, for example, by showing subjects a visual stimulus and later asking them to visualize

the same stimulus (Le Bihan et al., 1993). This type of experiment is considerably more difficult to

perform in animals. Using movement as a proxy for self-generated activity patterns, however, using

a virtual environment, we can compare a condition in which self-generated and externally generated

activity patterns are correlated to a condition in which they are not. In the visual system, we can use

a virtual reality environment to experimentally couple or decouple locomotion from visual flow

feedback. We will refer to the former as a closed loop condition and to the latter as an open loop

condition. Assuming the predictive processing model is correct, we should find differences in the

activation patterns in dorsal cortex in response to locomotion onsets as a function of whether they

occurred with or without coupled visual flow feedback. Both prediction error neurons as well as

internal representation neurons should respond differentially. Thus, we set out to investigate

whether there are differences between closed and open loop activation patterns of mouse dorsal

cortex, and whether these differences are neuron type specific. If hallucinations are indeed the

result of aberrant activation of internal representations, possibly as a result of misconfigured

internal models, antipsychotic drugs should alter the differential activation patterns in these

neurons.

RESULTS

Tlx3 positive layer 5 (L5) intratelencephalic (IT) neurons distinguished closed and open loop
visuomotor coupling

To investigate neuron type specific differences in activity patterns at locomotion onsets with and

without self-generated visual feedback in mouse dorsal cortex, we quantified locomotion onset

activity using widefield calcium imaging (Wekselblatt et al., 2016). A GCaMP variant was expressed

using either a retro-orbital injection of an AAV-PHP.eB vector (Chan et al., 2017), or a cross between

a Cre line and the Ai148 GCaMP6 reporter line (Daigle et al., 2018) (see Methods; **Figure 1A**, **Table**

S2). Mice expressed a GCaMP variant either brain wide (C57BL/6), or in genetically identified subsets

of excitatory or inhibitory neurons using Cre driver lines. Please note, we use the term 'brain wide'

here to describe an expression pattern that was determined by the intersection of the AAV-PHP.eB

tropism and the expression pattern of EF1α or hSyn1 promoters but that was otherwise brain wide

(see Discussion). Prior to the start of the imaging experiments, mice were implanted with a crystal

skull cranial window (Kim et al., 2016). We used crystal skull preparations as we found that

hemodynamic artifacts were smaller in this preparation compared to those observed in clear skull

preparations (**Figures S1A and S1B**). For all experiments, mice were head fixed and free to locomote

on a spherical treadmill (**Figure 1B**). Activity from dorsal cortex was imaged while mice were in a

closed loop visuomotor condition, in which locomotion on the spherical treadmill controlled forward

movement in a virtual corridor. Following this, mice were exposed to an open loop condition during

which mice were presented with a replay of the visual flow they had self-generated in the preceding

closed loop condition.

To test whether there are differences between closed and open loop locomotion onset responses,

we first compared these in mice that expressed GCaMP brain wide (C57BL/6). Consistent with

previous findings (Musall et al., 2019; Stringer et al., 2019), we found that locomotion onset resulted

in broad activation across most of dorsal cortex in both conditions (**Figure 1C**). However, comparing

locomotion onsets in mice that expressed GCaMP6 only in Tlx3 positive L5 IT neurons, we found that

the activation pattern was strikingly different between closed and open loop conditions (**Figure 1D**).

While locomotion onset in the open loop condition resulted in an increase in activity across dorsal

cortex, similar to that observed in C57BL/6 mice with brain wide GCaMP expression, locomotion

onset in the closed loop condition resulted in a very different activation pattern: In posterior parts of

dorsal cortex, the activity decreased, while in anterior parts of the dorsal cortex, the activity

increased. To quantify these activation patterns across mice, we selected six regions of interest:

primary visual cortex (V1), antero-medial secondary visual cortex (V2am), retrosplenial cortex (RSC),

primary motor cortex (M1), anterior cingulate cortex at the level of bregma (A24b), and secondary

motor cortex (M2) (**Figure 1E**). In C57BL/6 mice that expressed GCaMP brain wide, closed loop

locomotion onset resulted in similar activity patterns across these six regions (**Figure 1F**). This

activity pattern was not markedly different from that observed during locomotion onsets in an open

loop condition (**Figure 1G**). To quantify the contribution of visual flow onset to the closed loop

locomotion onset activity, we measured responses to visual flow onsets that occurred in the absence

of locomotion in the open loop condition (**Figure 1H**). In C57BL/6 mice with brain wide GCaMP

expression, visual flow onsets resulted in responses that were small compared to those observed

during locomotion onset, even in V1. Note that a visual flow onset is a relatively weak visual stimulus

compared to a drifting grating stimulus. A grating stimulus is typically composed of a gray screen or a

standing grating stimulus that suddenly transitions to a drifting grating, and thus the visual flow

undergoes a very high acceleration. A visual flow onset in an open loop condition is the replay of

previously self-generated visual flow and thus exhibits much lower visual flow acceleration. The

absence of a strong visual flow onset response in C57BL/6 mice that expressed GCaMP brain wide is

consistent with a high similarity of closed and open loop locomotion onset responses.

Different from what we found in mice that expressed GCaMP brain wide, in mice that expressed

GCaMP6 in Tlx3 positive L5 IT neurons, the closed loop locomotion onset patterns differed markedly

when compared across the six regions of interest (**Figure 1I**). Moreover, these activation patterns

were different from those observed during open loop locomotion onsets (**Figure 1J**). While these

responses could partially be explained by hemodynamic occlusion, the time course of hemodynamic

occlusion in Tlx3-Cre mice that expressed eGFP in L5 IT neurons was similar to that observed in

C57BL/6 mice that expressed eGFP brain wide and mainly resulted in a decrease of fluorescence

(**Figures S1C and S1D**). In Tlx3 positive L5 IT neurons, open loop visual flow onsets that occurred in

the absence of locomotion resulted in quantifiable responses in V1 and V2am (**Figure 1K**), but the

closed loop locomotion onset responses in V1 could not be explained by the sum of locomotion and

visual flow onset responses (**Figures S2A and S2C**). Most of the widefield signals we record here

likely originate in compartments in superficial layers which are composed of axons and dendrites of

local neurons, and long-range inputs (Allen et al., 2017) (see Discussion). To test how well the

widefield signals recorded in V1 reflected local somatic activity in V1, we repeated the experiments

using two-photon calcium imaging of somatic activity in Tlx3 positive L5 IT neurons. We found that

while there were differences between the responses measured with widefield imaging and those

measured at the soma, both for locomotion onset responses and visual responses, these differences

did not change our conclusions (**Figure S3**).

To quantify the differences between closed and open loop locomotion onset responses of L5 IT

neurons in relation to those observed in other cortical neuron types, we repeated the experiments

for the selection of excitatory and inhibitory cortical neuron types (**Figures S4A-S4I**). For each

neuron type, we quantified the similarity of the activity patterns associated with closed and open

loop locomotion onsets by calculating the correlation coefficient between the activity traces in a

window around locomotion onset (-5 s to +3 s) for each region of interest (see Methods).

Quantifying the differences in V1, we find that the similarity in Tlx3 positive L5 IT neurons was lower

than that observed in all other neuron types tested, except for Ntsr1 positive L6 neurons (**Figure

S4J**). Tlx3 positive L5 IT neurons and Ntsr1 positive L6 neurons also exhibit a lower similarity than

other neuron types when comparing the similarity across all regions of interest (**Figure 1L**). Both in

V1 across mice, as well as when comparing across all areas, we found the highest similarity between

closed and open loop locomotion onsets across regions in C57BL/6 mice that expressed GCaMP

brain wide. Thus, among the neuron types tested, excitatory neurons of deep cortical layers

exhibited the largest differences between closed and open loop locomotion related activation across

all the selected cortical regions. The remarkable aspect of these findings is not that there are

differences between closed and open loop locomotion onsets, particularly in visual regions of cortex

one would perhaps expect to find differences, but that these differences are larger specifically in the

deep cortical layers.

To quantify how the interaction between locomotion and visual flow feedback changes across dorsal

cortex, we computed correlations between calcium activity and locomotion speed for each region of

interest, for closed and open loop conditions separately. In C57BL/6 mice that expressed GCaMP

brain wide, this correlation was comparable between closed and open loop conditions, was highest

in posterior regions of dorsal cortex, and lower in more anterior regions (**Figures 2A and 2B**). In mice

that expressed GCaMP6 in Tlx3 positive L5 IT neurons, the correlation between locomotion speed

and calcium activity in the closed loop condition was negative in posterior regions of dorsal cortex

and positive in anterior regions. Conversely, in open loop conditions, the correlation was positive

throughout dorsal cortex (**Figures 2C and 2D**). To test whether it was the coupling of visual flow that

resulted in a reduction of the correlation in the closed loop condition or whether the presence of an

uncoupled visual stimulus resulted in an increase of correlation in the open loop condition, we also

compared these correlations to those observed when mice were locomoting in relative darkness

(see Methods). In C57BL/6 mice that expressed GCaMP brain wide, the correlation of calcium activity

with locomotion in darkness was indistinguishable from that in closed or open loop (**Figure 2B**). In

mice that expressed GCaMP6 in Tlx3 positive L5 IT neurons, an interesting pattern emerged. In visual

regions of dorsal cortex, the correlation between locomotion speed and calcium activity in relative

darkness resembled that observed in open loop conditions, while in frontal regions, it resembled

that observed in closed loop conditions. In visual regions, the coupling of visual flow resulted in a

reduction of correlation, while in frontal regions the presence of an uncoupled visual stimulus

resulted in an increase of correlations. Thus, the difference between closed and open loop

conditions likely had different origins in visual and frontal regions of dorsal cortex. Overall, we

concluded that the coupling of locomotion and resulting visual flow feedback not only activated Tlx3

positive L5 IT neurons differentially but did so throughout dorsal cortex, well beyond visual regions.

Visuomotor prediction errors originated in visual cortical regions and had opposing effects on Tlx3
positive L5 IT activity

A consequence of the natural coupling of locomotion and visual flow feedback is that movements

are a good predictor of self-generated sensory feedback. In our closed loop experiments, forward

locomotion was a strong predictor of backward visual flow. It has been speculated that such

predictions are used to compute prediction errors in L2/3 of cortex (Jordan and Keller, 2020).

Prediction errors come in two variants: Positive prediction errors that signal more visual flow than

predicted and negative prediction errors that signal less visual flow than predicted (Keller and Mrsic-

Flogel, 2018; Rao and Ballard, 1999). To probe for negative and positive prediction error signals, we

quantified the responses to visuomotor mismatch events and to unexpected visual stimuli

(representing less and more visual flow input than expected, respectively). Visuomotor mismatch

was introduced by briefly (1 s) halting visual flow during self-generated feedback in the closed loop

condition. Visual responses were elicited by presenting full-field (covering the entire toroidal screen)

drifting grating stimuli to a passively observing mouse (see Methods). Please note, both types of

prediction errors can be probed using a variety of stimuli, but the two we are using here, in our

experience, are the most robust at eliciting strong responses in V1. In C57BL/6 mice that expressed

GCaMP brain wide, both visuomotor mismatch and grating stimuli resulted in increases of activity

that were strongest and appeared first in visual regions of dorsal cortex (**Figures 3A-3C and S2B**). In

mice that expressed GCaMP6 in Tlx3 positive L5 IT neurons, visuomotor mismatch resulted in an

increase of activity in V1, while grating onsets resulted in a decrease of activity in V1 (**Figures 3D-3F**

and S2D). Thus, in dorsal cortex, both mismatch and drifting grating stimuli primarily activated visual

cortex but resulted in opposing activation patterns in Tlx3 positive L5 IT neurons. An increase in

activity to negative prediction errors and a decrease to positive prediction errors would be

consistent with either a population of negative prediction error neurons or a type of internal

representation neuron (Rao and Ballard, 1999). Given that L5 neurons in V1 do not exhibit the

opposing influence of top-down and bottom-up inputs necessary to compute prediction errors

(Jordan and Keller, 2020), we speculated that Tlx3 positive L5 IT neurons might function as a type of

internal representation neuron.

The antipsychotic drug clozapine alters visuomotor integration in L5

The activation of an internal representation is the brain's best guess at what the stimuli in the

environment are. We speculated that the activation of such an internal representation would be

particularly susceptible to substances that are known to reduce illusory percepts, like antipsychotic

drugs. We thus quantified the changes in dorsal cortex activity associated with a single

intraperitoneal injection of an antipsychotic drug (clozapine). In C57BL/6 mice that expressed

GCaMP brain wide, we found that closed and open loop locomotion onsets were almost unaffected

by clozapine (**Figures 4A and 4B**), while there was a small increase in open loop visual flow onset

responses (**Figure 4C**). In mice that expressed GCaMP6 in Tlx3 positive L5 IT neurons, however,

clozapine fundamentally changed both closed and open loop locomotion onset responses. In V1,

both types of locomotion onset now resulted in a massive increase in activity (**Figures 4D and 4E**).

Conversely, open loop visual flow onset responses remained largely unchanged (**Figure 4F**). These

effects could not be explained by clozapine induced changes in hemodynamic occlusion (**Figure S5A**)

or locomotion behavior (**Figures S5B and S5C**). Clozapine also had an opposing effect on the average

correlation of activity with locomotion speed in Tlx3 positive L5 IT neurons. While the average

correlation between activity and locomotion speed was increased in C57BL/6 mice that expressed

GCaMP brain wide, the same measure was decreased in mice that expressed GCaMP6 in Tlx3

positive L5 IT neurons (**Figure S5D**). Finally, we confirmed that this effect was also present when

recording somatic activity in V1. Using two-photon imaging, after clozapine administration we

observed an increase of somatic responses during locomotion onsets that was very similar to that

observed with widefield imaging (**Figures S5E and S5F**). Thus, consistent with the speculation that

Tlx3 positive L5 IT neurons might function as internal representation neurons, clozapine exhibited a

substantially stronger influence on locomotion onset responses in Tlx3 positive L5 IT neurons than

would be expected from the effect of clozapine on brain wide responses.

Antipsychotic drugs decouple long-range cortico-cortical activity

L5 IT neurons are likely the primary source of long-range cortical communication (Chen et al., 2019;

Harris et al., 2019; Leinweber et al., 2017). Given that clozapine increased locomotion onset

responses in L5 IT neurons in V1, we wondered whether clozapine might more generally change the

long-range influence between L5 IT neurons. To investigate this possibility, we computed the

correlations of calcium activity between the six regions of interest across both hemispheres and all

recording conditions. We did this first in C57BL/6 mice that expressed GCaMP brain wide before

(**Figure 5A**) and after a single intraperitoneal injection of clozapine (**Figure 5B**). Overall, clozapine

resulted in a relatively modest decrease in correlations. On repeating this experiment in mice that

expressed GCaMP6 in Tlx3 positive L5 IT neurons, we found a larger decrease in correlations that

appeared to be stronger for regions that were further apart (**Figures 5C and 5D**). To quantify these

changes, we plotted the correlations as a function of linear distance between the regions as

measured in a top-down view of dorsal cortex (**Figure S6A**). Note, this is only an approximation of

the actual axonal path length between the regions. To visualize the distribution of correlations, we

interpolated the data and represented them as heatmaps (**Figure S6B**). The resulting representation

for the data from C57BL/6 mice that expressed GCaMP brain wide before and after clozapine

injection is shown in **Figures 6A** and **6B**, respectively. We then split the data into approximately

equal portions of short- and long-range correlations using a cutoff of 0.9 times the bregma-lambda

distance (approximately 3.8 mm). While both short- and long-range correlations were reduced, this

reduction was not significant in either case (**Figure 6C**). We also found no evidence of a difference

between short- and long-range correlation changes. In mice that expressed GCaMP6 in Tlx3 positive

L5 IT neurons, however, we found a significant reduction in both short- and long-range correlations

(**Figures 6D and 6E**), and both were more strongly reduced than in C57BL/6 mice that expressed

GCaMP brain wide (C57BL/6 vs Tlx3-Cre x Ai148, short-range: $p < 0.05$; long-range: $p < 10^{-5}$; rank-sum

test). Moreover, the reduction in activity correlations was stronger for long-range correlations than

it was for short-range correlations (**Figure 6F**). To control for possible clozapine induced changes in

hemodynamic occlusion, we repeated the experiment and analyses in Tlx3-Cre mice that expressed

eGFP in L5 IT neurons and found that alterations in hemodynamic occlusion could not account for

the decrease in correlations but resulted in a small increase of the same metric (**Figures S6C-S6E**). To

test whether the stress of injection, or simply passage of time, could explain the observed effects,

we conducted a separate set of experiments in which we injected saline in a cohort of Tlx3-Cre x

Ai148 mice and performed the same analysis. We found no evidence of a change in correlations with

saline injections (**Figures S6F-S6H**). To further dissect the spatial scale at which this decorrelation

occurred, we repeated the experiments in Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT

neurons, recording calcium activity using both a widefield and a two-photon microscope. We found

that while the activity correlations measured with the widefield imaging decreased (**Figures S7A-**

S7C), pairwise somatic activity correlations measured by two-photon imaging did not (**Figure S7D**).

This is consistent with the idea that decorrelation primarily occurs for long-range interactions.

Finally, to confirm that the clozapine induced reduction in correlation was stronger in deep cortical

layers, we repeated the experiment in a population of L2/3 and L4 excitatory neurons using Cux2-

CreERT2 x Ai148 mice. We found that clozapine also decreased correlations of cortical activity in

superficial excitatory neurons (**Figure S8**). However, this reduction was significantly weaker than the

reduction we observed in mice that expressed GCaMP6 in Tlx3 positive L5 IT neurons (Cux2-CreERT2

x Ai148 vs Tlx3-Cre x Ai148, short-range: $p < 0.005$, long-range: $p < 10^{-8}$, rank-sum test). Thus,

administration of the antipsychotic drug clozapine resulted in a decorrelation of activity across

dorsal cortex that was stronger in L5 IT neurons (Tlx3) than it was in either upper layer excitatory

neurons (Cux2) or in the brain wide average (C57BL/6).

To test whether the decorrelation of the activity of L5 IT neurons is specific to clozapine or might

more generally be a functional signature of antipsychotic drugs, we repeated the experiments with

two additional antipsychotic drugs, aripiprazole and haloperidol. We found that the principal effect

of decorrelation of the activity patterns of Tlx3 positive L5 IT neurons was preserved with both drugs

(**Figures 7A-7F**). These changes in correlation were absent upon injection of a psychostimulant,

amphetamine (**Figures 7G-7I**). Thus, it is possible that a decrease in long-range correlations between

cortical L5 IT neurons might be a common mechanism of action for antipsychotic drugs.

Changes in the correlation between cortical regions could either be driven by changes in correlated

external inputs or through changes in communication between cortical regions. If antipsychotic

drugs primarily reduce correlations by reducing the strength of communication between cortical

regions, independent of whether this communication is direct or indirect, we would expect to find

changes in the way responses spread between cortical regions. We used visuomotor prediction error

responses that originated in V1 (**Figure 3**) to quantify how the spread of these responses in Tlx3

positive L5 IT neurons was influenced by antipsychotic drugs. The computation of negative

prediction errors during movement is thought to be driven by an excitatory prediction of visual flow

mediated by long-range cortical input from L5 neurons in regions like A24b (Leinweber et al., 2017),

as well as an inhibitory bottom-up visual input. By contrast, the computation of a positive prediction

error during passive observation is thought to be driven by an excitatory bottom-up visual input and

an inhibitory top-down input. Thus, we would primarily expect to find a reduction in mismatch

responses in L5 IT neurons and a reduction in the spread of this signal to secondary visual regions.

This is indeed what we observed: Mismatch responses were partially reduced in V1 and almost

absent in V2am after injection of antipsychotic drugs (**Figure 8A**). While responses to onsets of

drifting gratings were also reduced in V2am, they were overall less affected by the antipsychotic

drugs than mismatch responses (**Figure 8B**). These results are consistent with the interpretation that

antipsychotic drugs reduce lateral communication in L5. Whether this communication is mediated by

direct long-range connections between L5 neurons, or indirectly via other cortical neurons or

subcortical loops, remains to be tested.

DISCUSSION

When interpreting our results, it should be kept in mind that the source of the widefield imaging

signal in these experiments is not entirely clear and is likely a combination of somatic, dendritic, and

axonal signals. Given that the signal attenuation half depth in cortex is on the order of 100 μm

(Aravanis et al., 2007), most signals likely come from the superficial 100 μm of tissue (Allen et al.,

2017). In the case of the Tlx3 recordings, this would mean that we were recording a mixture of

axonal and dendritic signals of the Tlx3 neurons that have their soma in L5 but have a dense net of

axons and dendrites in L1. In the case of the brain wide expression of GCaMP using an AAV-PHP.eB

in C57BL/6 mice, the signals we recorded likely also include contributions from thalamic axons. In

addition to this, there is a hemodynamic contribution to the signal (Allen et al., 2017; Ma et al.,

2016; Valley et al., 2020) that is the result of varying occlusion by blood vessels. We attempted to

correct for the hemodynamic response using a 405 nm wavelength isosbestic illumination of

GCaMP6 but could not recover the hemodynamic responses measured in eGFP experiments.

Accurate correction likely requires calibration on eGFP measurements (Valley et al., 2020). Given

that hemodynamic responses should not systematically differ between different mouse lines, and

that in the crystal skull preparation the contribution of hemodynamic occlusion to increases in

fluorescence was relatively small (**Figure S1**), these hemodynamic artefacts should not substantially

alter our conclusions. Lastly, it should be kept in mind that responses in these widefield recordings

inherently reflect population averages. Assuming there is some form of lateral inhibition among

neurons, population averages need not reflect responses of neurons that are tuned to the specific stimulus presented. For example, the suppression to grating stimuli we observed in V1 Tlx3 positive L5 IT neurons could be the result of a subset of Tlx3 neurons that are strongly responsive to the specific grating presented suppressing the rest of the population of Tlx3 neurons. The sum of both could then result in a net decrease in calcium activity. Thus, an understanding of the mechanism that results in opposing population responses of Tlx3 positive L5 IT neurons to positive and negative prediction errors will require measurements with single neuron resolution.

In our experiments we have focused on Tlx3 positive L5 IT neurons, but given that L5 IT neurons are densely connected to L5 pyramidal tract neurons (Anderson et al., 2010; Kiritani et al., 2012) and to L6 neurons (Xu et al., 2016; Yamawaki and Shepherd, 2015), it is not unlikely that these neuron types would follow similar patterns of activity and exhibit similar susceptibility to the influence of antipsychotic drugs. In the comparison of closed and open loop locomotion onsets, we do indeed find that Ntsr1 positive L6 excitatory neurons exhibit similar levels of dissimilarity as those observed in Tlx3 positive L5 IT neurons (**Figure 1L**). It is possible that responses in the Ntsr1 recordings, however, were noisier than those in the Tlx3 recordings (**Figure S4**), likely because of the lower dendritic and axonal volume of Ntsr1 positive neurons in L1. Thus, while we have focused on the Tlx3 positive L5 IT neurons mainly for reasons of experimental convenience, it is possible that both the differential activation in closed and open loop locomotion onsets as well as the susceptibility of long-range correlations to antipsychotic drugs is a general feature of cortical layers 5 and 6. Another possibility we currently cannot entirely rule out is that the decorrelation effect we observed in Tlx3 positive L5 IT neurons is specific to the Tlx3-Cre mouse line. However, this seems unlikely given that all mouse lines we used were maintained on the same C57BL/6 background.

It is still unclear what the relevant sites of action in the brain are of antipsychotic drugs. Antipsychotic drugs are often antagonists of the D2 receptors, which are most strongly expressed in striatum. Consequently, it is sometimes assumed that the relevant site of action of antipsychotics is the striatum. Consistent with this, an increase in striatal dopamine activates auditory cortex (Maia and Frank, 2017) and mediates hallucination-like perceptions in mice (Schmack et al., 2021). It is intriguing, however, to speculate that the effects we observed in L5 IT neurons, which provide a dominant input to striatum (Morita et al., 2019), may be an alternate site of action. While it is possible that the primary site of action of the antipsychotic drugs with respect to the L5 decorrelation phenotype we observed is striatum, this is unlikely. Changes to striato-thalamo-cortical loops could appear as changes in functional connectivity observed in cortex. However, feedback from striatum to cortex is exclusively mediated through thalamus, and neither L4, the

primary thalamo-recipient layer in cortex, nor our brain wide recordings exhibited any of the effects

we saw in L5 IT neurons. Thus, the effects we saw in L5 IT neurons are either not inherited from

striatum or are mediated by a separate set of thalamic neurons via direct thalamic input to deep

cortical layers (Constantinople and Bruno, 2013; Douglas and Martin, 1991). Thalamocortical

collaterals in deep layers, however, come from thalamic neurons that project primarily to superficial

layers (Oberlaender et al., 2012). Thus, we think it is unlikely that the observed effects in L5 IT

neurons were inherited from striatum. Independent of whether the treatment-relevant site of action

of antipsychotic drugs is indeed in L5 IT neurons, or whether the observed decorrelation of activity is

relevant to antipsychotic potency of these drugs, we speculate that the long-range decorrelation of

activity could be used as a basis for a functional screen for compounds with antipsychotic efficacy.

Given that in the brain wide recordings the decorrelation of activity was weaker compared to that

observed in Tlx3 positive L5 IT neurons, it might be possible to test for similar effects in human

subjects using layer specific fMRI recordings (Haarsma et al., 2020).

A rather surprising implication of our results is that the dominant mode of coupling in cortex might

be lateral, not vertical. Two findings would argue for this interpretation. First, while L5 was

differentially activated across cortical regions during closed and open loop locomotion onsets, L2/3

exhibited rather limited differences. Second, antipsychotics decorrelated L5 activity but had no

effect on L2/3. Both findings could be explained if the long-range lateral coupling within L5 is

stronger than the local vertical coupling between L2/3 and L5.

One of the diseases treated with antipsychotic drugs is schizophrenia. It has been speculated that

disrupted resting-state functional connectivity in brain networks involving cortex is central to the

pathogenesis of schizophrenia (Li et al., 2017; Northoff and Duncan, 2016). Treatment-naïve

schizophrenia patients predominantly exhibit increases in resting state functional connectivity

between cortical regions, while medicated schizophrenia patients exhibit more balanced fractions of

increased and decreased functional connections (Anticevic et al., 2015; Du et al., 2021; Li et al.,

2017). Auditory hallucinations, one of the hallmark symptoms of schizophrenia, are associated with

increased activation of diverse cortical regions (Catafau et al., 1994; Dierks et al., 1999; Shergill et al.,

2000). Thus, it is conceivable that antipsychotic drugs counteract this by reducing the strength of

long-range cortical connectivity mediated by L5 IT neurons, independent of whether the increased

activity in schizophrenic patients is indeed driven by aberrant increases in the strength of long-range

cortical connections or is a consequence of an aberrant increase in the strength of striato-thalamo-

cortical loops. However, a direct involvement of cortical circuits in the pathogenesis of schizophrenia

is consistent with the glutamate hypothesis of schizophrenia. This hypothesis is based on the finding

that NMDA receptor antagonists induce schizophrenia-like symptoms (Moghaddam and Javitt,

2012), and that disturbances of NMDA and AMPA receptor related gene expression are associated

with schizophrenia (Harrison and Weinberger, 2005; Singh et al., 2020). In V1, it has been shown that

NMDA receptors are necessary during visual development to establish normal prediction error

signaling (Widmer and Keller, 2021). Thus, it is conceivable that a disruption of glutamatergic

signaling in cortex that alters the way internal representations are activated by prediction errors is

central to the etiology of schizophrenia. If so, we should find impairments in all domains of cortical

function, including sensory processing. In the case of the sensory cortex, cortical dysfunctions should

be apparent earlier in development, concurrent with the critical period for plasticity in sensory areas

of cortex. While the core symptoms of schizophrenia typically appear in late adolescence, sensory

processing is indeed impaired in schizophrenic patients (Javitt, 2009) and is apparent, for example, in

the reduction of surround suppression in V1 (Yoon et al., 2009). Interestingly, some of these

disturbances are detectable already during childhood as visuo-perceptual and reading anomalies in

some subjects that later develop schizophrenia (Parellada et al., 2017). Positive symptoms of

schizophrenia can be explained in a predictive processing framework of cortical function as a

disturbance of a prediction error based update of internal representations (Fletcher and Frith, 2009;

Frith, 2005). There are two types of prediction error neurons, positive and negative, and, given that

cortex likely implements a non-hierarchical variant of predictive processing (Garner and Keller,

2022), two types of internal representation neurons. One set of internal representation neurons is

excited by positive prediction errors and inhibited by negative prediction errors, and the other one is

inhibited by positive prediction errors and excited by negative prediction errors (Keller and Mrsic-

Flogel, 2018). Tlx3 positive L5 IT neurons exhibited opposing responses to positive and negative

prediction errors (**Figure 3**), and, while several different interpretations are conceivable, these

responses would be consistent with those of internal representation neurons. Given that the effects

of antipsychotic drugs are predominantly observed in L5 IT neurons, and that anesthesia is thought

to decouple L5 neuron soma from their apical tuft and thereby prevent conscious perception (Suzuki

and Larkum, 2020), we speculate that Tlx3 positive L5 IT neurons may be one type of internal

representation neuron and one of the functionally relevant sites of antipsychotic action.

Figure 1. Activation patterns in deep cortical layers distinguished closed and open loop locomotion onsets more strongly than superficial layers.

(**A**) Schematic of GCaMP expression strategy. We either injected an AAV-PHP.eB virus retro-orbitally to express GCaMP brain wide (C57BL/6), in cortical excitatory neurons (Emx1-Cre) or in a subset of SST positive interneurons (see Methods and **Table S2**), or used the progeny of a cross of a cell type specific Cre driver line (Neuronal[Cre]: Cux2-CreERT2, Scnn1a-Cre, Tlx3-Cre, Ntsr1-Cre, PV-Cre, VIP-Cre or SST-Cre) with the Ai148 GCaMP6 reporter line. All mice were then implanted with a crystal skull cranial window prior to imaging experiments.

(**B**) Schematic of the experimental setup. We imaged GCaMP fluorescence under 470 nm LED illumination with an sCMOS camera through a macroscope. Mice were free to locomote on an air supported spherical treadmill while coupled (closed loop), uncoupled (open loop), or no (dark) visual flow in the form of movement along a virtual corridor was displayed on a toroidal screen placed in front of the mouse. Walls of the virtual corridor were patterned with vertical sinusoidal gratings. In a separate condition, we then presented drifting grating stimuli (grating session, see Methods).

(**C**) Average response in an example C57BL/6 mouse that expressed GCaMP6 brain wide during
closed loop locomotion onsets (top row, 83 onsets) and open loop locomotion onsets (bottom row,
153 onsets). Locomotion onsets in both conditions activated dorsal cortex similarly.

(**D**) As in **C**, but in an example Tlx3-Cre x Ai148 mouse that expressed GCaMP6 in L5 IT neurons
during closed loop locomotion onsets (top row, 88 onsets) and open loop locomotion onsets
(bottom row, 83 onsets). Note that activity decreased in posterior regions of dorsal cortex during
closed loop locomotion onsets.

(**E**) Example crystal skull craniotomy marking the 6 regions of interest in each hemisphere we
selected: primary visual cortex (V1, red), retrosplenial cortex (RSC, blue), antero-medial secondary
visual cortex (V2am, green), primary motor cortex (M1, yellow), anterior cingulate cortex (A24b,
purple), and secondary motor cortex (M2, cyan). The white cross marks bregma.

(**F**) Average response during closed loop locomotion onsets in C57BL/6 mice that expressed GCaMP
brain wide in the 6 regions of interest (activity was averaged across corresponding regions in both
hemispheres). Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical
bootstrap estimate for each time bin (see Methods and **Table S1**).

(**G**) As in **F**, but for open loop locomotion onsets.

(**H**) As in **F**, but for visual flow onsets in the open loop condition restricted to times when the mice
were not locomoting.

(**I**) Average response during closed loop locomotion onsets in Tlx3-Cre x Ai148 mice that expressed
GCaMP6 in L5 IT neurons, activity was averaged across corresponding regions in both hemispheres).
Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap
estimate for each time bin (see Methods and **Table S1**).

(**J**) As in **I**, but for open loop locomotion onsets.

(**K**) As in **J**, but for visual flow onsets during open loop sessions restricted to times when the mice
were not locomoting.

(**L**) Similarity of the average closed and open loop locomotion onset responses quantified as the
correlation coefficient between the two in a window -5 s to +3 s around locomotion onset (see
Methods). Error bars indicate SEM over the 12 (6 per hemisphere) cortical regions. Statistical
comparisons are against the Tlx3 data and were corrected for family-wise error rate (see Methods
and **Table S1**): adjusted significance thresholds, n.s.: not significant, *: $p < 0.05/9$, **: $p < 0.01/9$,
***: $p < 0.001/9$.

Figure 2. L5 IT neurons were differentially activated by locomotion depending on the type of visual feedback.

(**A**) Correlation between calcium activity and locomotion speed in the closed loop (left) and open loop (right) conditions, calculated for each pixel in the image for one example C57BL/6 mouse that expressed GCaMP brain wide. The black cross marks bregma. Activity in most of dorsal cortex correlates positively with locomotion speed in both closed and open loop conditions.

(**B**) Average correlation between calcium activity and locomotion speed in the 6 regions of interest in closed loop (red), open loop (blue) or dark (black) conditions in 6 C57BL/6 mice that expressed GCaMP brain wide (308, 316, and 68 5-minute recording sessions, respectively). Error bars indicate SEM over recording sessions. Bars above the plot indicate significant differences between conditions (compared conditions are indicated by colored line segments to the left, black: $p < 0.05$, gray: n.s., see **Table S1** for all information on statistical testing). On average, the correlation was highest in posterior dorsal cortex and was not different between conditions.

(**C**) As in **A**, but for an example Tlx3-Cre x Ai148 mouse that expressed GCaMP6 in L5 IT neurons. Correlations of calcium activity and locomotion speed were lower in the closed loop condition compared to the open loop condition, most prominently in posterior regions of dorsal cortex.

(**D**) As in **B**, but for 15 Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons (data from 420 closed loop, 394 open loop and 194 dark 5-minute recording sessions). Error bars indicate SEM over recording sessions. Correlation differed significantly between the closed and the open loop condition and was lowest in posterior dorsal cortex during the closed loop condition.

Figure 3. Visuomotor prediction error responses in dorsal cortex originated in V1 and activated L5 IT neurons differentially.

(**A**) Average responses to mismatches (top row, 230 onsets) and drifting gratings (bottom row, 86 onsets) in an example C57BL/6 mouse that expressed GCaMP brain wide.

(**B**) Left: Average response to mismatches in C57BL/6 mice that expressed GCaMP brain wide in 6 cortical regions (activity was averaged across corresponding regions in both hemispheres). Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**). Orange shading indicates duration of mismatch event. Right: Average response map to mismatch in 6 C57BL/6 mice (see Methods). Black cross marks bregma.

(**C**) As in **B**, but for drifting grating responses.

503   (**D**) As in **A**, but in an example Tlx3-Cre x Ai148 mouse that expressed GCaMP6 in L5 IT neurons (top

504   row: mismatch, 292 onsets; bottom row: drifting gratings, 171 onsets).

505   (**E**) As in **B**, but for Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons.

506   (**F**) As in **C**, but for Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons.

Figure 4. Clozapine increased locomotion related responses in L5 IT neurons in V1.

(**A**) Average response during closed loop locomotion onsets in C57BL/6 mice that expressed GCaMP brain wide (activity was averaged across corresponding regions in both hemispheres) after a single injection of the antipsychotic drug clozapine (see Methods). Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**). Inset: Comparison of responses for V1 in naive C57BL/6 mice (dashed red) and the same mice after clozapine injection (dark red, same data as in main panel).

(**B**) As in **A**, but for open loop locomotion onsets after a clozapine injection.

(**C**) As in **A**, but for open loop visual flow onsets restricted to times when the mice were not locomoting.

(**D**) Average response during closed loop locomotion onsets in Tlx3-Cre x Ai148 mice that expressed GCaMP6 in layer L5 IT neurons (activity was averaged across corresponding regions in both hemispheres) after a single injection of the antipsychotic drug clozapine (see Methods). Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**). Inset: Comparison of responses for V1 in naive Tlx3-Cre x Ai148 mice (dashed red) and the same mice after clozapine injection (dark red, same data as in main panel).

(**E**) As in **D**, but for open loop locomotion onsets.

(**F**) As in **D**, but for open loop visual flow onsets restricted to times when the mice were not locomoting.

Figure 5. Clozapine reduced activity correlations in dorsal cortex predominantly in L5 IT neurons.

(**A**) Average correlation of activity between the 12 regions of interest in 4 C57BL/6 mice that expressed GCaMP brain wide.

(**B**) As in **A**, but after a single injection of the antipsychotic drug clozapine in the same 4 C57BL/6 mice.

(**C**) Average correlation of activity between the 12 regions of interest in 5 Tlx3-Cre x Ai148 mice that expressed GCaMP6 in layer L5 IT neurons.

(**D**) As in **C**, but after a single injection of the antipsychotic drug clozapine in the same 5 Tlx3-Cre x Ai148 mice.

Figure 6. L5 IT neurons exhibited the strongest clozapine induced reduction of long-range correlations.

(**A**) Density map of correlation coefficients as a function of distance between the regions, normalized across mice by the bregma-lambda distance (see **Figure S6A and S6B**, and Methods), for 4 naive C57BL/6 mice that expressed GCaMP brain wide. Peak density is indicated at the bottom right of the plot and corresponds to the maximum value of the color scale (dark red). The white line is a contour line drawn at 50% of peak value.

(**B**) As in **A**, but for the same 4 C57BL/6 mice after a single injection of the antipsychotic drug clozapine. The white line is the contour line from panel **A** for comparison.

(**C**) Clozapine induced change in the activity correlation between regions, normalized to the correlation coefficient in the naive state, in 4 C57BL/6 mice. Data were split into short- and long-range activity correlations (see Methods). Boxes represent the median and the interquartile range. The open circle indicates the mean of the distribution. The whiskers mark 1.5 times the interquartile range. Dots are the individual data points (short-range: 122 pairs of regions, long-range: 142 pairs of regions). n.s.: not significant. See **Table S1** for all information on statistical testing.

(**D**) As in **A**, but for 5 Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons.

(**E**) As in **B**, but for 5 Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons. The white line is the contour line from panel **D** for comparison.

(**F**) As in **C**, but for 5 Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons (short-range: 182 pairs of regions, long-range: 148 pairs of regions). ***: $p < 0.001$.

Figure 7. Antipsychotic drugs aripiprazole and haloperidol mimicked the decorrelation of L5 IT activity observed with clozapine while psychostimulant amphetamine did not.

(**A**) Density map of correlation coefficients as a function of distance between the regions, normalized across mice by the bregma-lambda distance (see **Figure S6A and S6B**, and Methods), for 3 Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons. Peak density is indicated at the bottom right of the plot and corresponds to the maximum value of the color scale (dark red). The white line is a contour line drawn at 50% of peak value.

(**B**) As in **A**, for the same 3 Tlx3-Cre x Ai148 mice, but after a single injection of the antipsychotic drug aripiprazole. The white line is the contour line from panel **A** for comparison.

(**C**) Aripiprazole induced change in the activity correlation between regions, normalized to the correlation coefficient in the naive state, in 3 Tlx3-Cre x Ai148 mice. Data were split into short- and long-range activity correlations (see Methods). Boxes represent the median and the interquartile range. The open circle indicates the mean of the distribution. The whiskers mark 1.5 times the

interquartile range. Dots are the individual data points (short-range: 114 pairs of regions, long-range:

84 pairs of regions). *: $p < 0.05$, ***: $p < 0.001$. See **Table S1** for all information on statistical testing.

(**D**) As in **A**, but for 3 different Tlx3-Cre x Ai148 mice.

(**E**) As in **D**, for the same 3 Tlx3-Cre x Ai148 mice, but after a single injection of the antipsychotic drug

haloperidol. The white line is the contour line from panel **D** for comparison.

(**F**) As in **C**, but for the 3 mice that had received a single injection of the antipsychotic drug

haloperidol (short-range: 114 pairs of regions, long-range: 84 pairs of regions). *: $p < 0.05$; n.s.: not

significant.

(**G**) As in **A**, but for 3 different Tlx3-Cre x Ai148 mice.

(**H**) As in **G**, for the same 3 Tlx3-Cre x Ai148 mice, but after a single injection of the psychostimulant

amphetamine. The white line is the contour line from panel **G** for comparison.

(**I**) As in **D**, but for the 3 mice that had received a single injection of the psychostimulant

amphetamine (short-range: 114 pairs of regions, long-range: 84 pairs of regions). n.s.: not significant.

Figure 8. Antipsychotic drug treatment preferentially reduced responses to and propagation of negative prediction errors.

(**A**) Average responses to mismatches in Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons before (red: V1 naive, green: V2am naive, activity was averaged across corresponding regions in both hemispheres) and after (dark red: V1, dark green: V2am) injection of a single dose of an antipsychotic drug (data were averaged over all antipsychotics used: Clozapine: 5 mice, aripiprazole: 3 mice, and haloperidol: 3 mice). Orange shading indicates duration of the mismatch event. Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**).

(**B**) As in **A**, but for drifting grating onsets.

     SUPPLEMENTAL FIGURES

Figure S1. Fluorescence changes driven by hemodynamic occlusion. Related to Figure 1.

(**A**) Average responses during locomotion onsets in C57BL/6 mice that expressed eGFP brain wide using a crystal skull preparation or in similarly transfected mice using a clear skull preparation (see Methods). Top row shows the average activity of corresponding regions in dorsal cortex. Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**). The heatmaps in the bottom row show the responses for individual regions of interest. Heatmaps are scaled to the y-axis limits of the plot above (blue low, red high). Note, the fast onset transient apparent in the clear skull preparation was absent in the crystal skull preparation, while the slow decrease of activity was present in both. The increase in fluorescence at locomotion onset, which was primarily apparent in the clear skull preparation, is driven by hemodynamic occlusion.

(**B**) As in **A**, but for mismatches. Orange shading (top) or white dashed lines (bottom) indicate the duration of the mismatch stimulus.

(**C**) As in **A**, left, but for Tlx3-Cre that had been retro-orbitally injected with an AAV-PHP.eB-DIO-eGFP to express eGFP in L5 IT neurons.

616 (**D**) As in **B**, left, but for Tlx3-Cre that had been retro-orbitally injected with an AAV-PHP.eB-DIO-eGFP
617 to express eGFP in L5 IT neurons.

Figure S2. In V1, the sum of locomotion and visual flow onset could not explain the closed loop locomotion onset response of L5 IT neurons. Related to Figure 1.

(**A**) Average responses during closed (red) and open loop (blue) locomotion onsets, as well as open loop visual flow (turquoise) onsets, and the sum of open loop locomotion and open loop visual flow onsets (black) in C57BL/6 mice that expressed GCaMP brain wide (6 mice). Closed loop locomotion onset responses in V1 were larger than open loop locomotion onset responses, and part of this difference could be explained by the visual flow onset responses. Shading indicates SEM over mice.

(**B**) Average response during mismatch (red) and full-field drifting grating onsets (blue) in 6 C57BL/6 mice that expressed GCaMP brain wide. Same data as in **Figures 3B and 3C**. Shading indicates SEM over mice.

(**C**) As in **A**, but for Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons (15 mice). Closed loop locomotion onset responses in V1 could not be explained as the sum of open loop locomotion and visual flow onset responses.

(**D**) As in **B**, but for Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons (15 mice). Same data as in **Figures 3E and 3F**.

Figure S3. L5 IT soma recorded in V1 with two-photon imaging show similar patterns of activity as the widefield signal recorded at the surface of the dorsal cortex. Related to Figure 1.

(**A**) Average response of V1 recorded with the widefield macroscope during closed (solid) or open loop (dashed) locomotion onsets in Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons. Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**).

(**B**) Average response of L5 soma in V1, recorded with two-photon imaging in the same Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons as in **A**, during either closed loop (solid) or open loop (dashed) locomotion onsets. Shading indicates SEM over 8434 neurons.

(**C**) As in **A**, but for responses to mismatches (solid), open loop halts (dashed) or drifting grating onsets (dotted). Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**).

(**D**) As in **B**, but for responses to mismatches (solid), open loop halts (dashed) or drifting grating onsets (dotted). Shading indicates SEM over 8434 neurons.

Figure S4. L5 IT neurons had strikingly different responses during closed and open loop locomotion onsets compared to other cortical neuron types. Related to Figure 1.

(**A**) Average response during closed loop locomotion onsets (top) and open loop locomotion onsets (bottom) in Emx1-Cre mice that expressed GCaMP6 in excitatory cortical neurons. Mean (lines) and 90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**). Heatmaps are scaled to the y-axis limits of the plot above (blue low, red high).

(**B**) As in **A**, but for Cux2-CreERT2 x Ai148 mice that expressed GCaMP6 in upper layer excitatory
neurons.

(**C**) As in **A**, but for Scnn1a-Cre x Ai148 mice that expressed GCaMP6 in L4 excitatory.

(**D**) As in **A**, but for Tlx3-Cre x Ai148 mice that expressed GCaMP6 specifically in L5 IT neurons. Data
are the same as in **Figures 1I and 1K**.

(**E**) As in **A**, but for Ntsr1-Cre x Ai148 mice that expressed GCaMP6 in excitatory L6 neurons.

(**F**) As in **A**, but for C57BL/6 mice that expressed GCaMP brain wide. Data are the same as in **Figures**
1F and 1G.

(**G**) As in **A**, but for PV-Cre x Ai148 mice that expressed GCaMP6 in PV neurons.

(**H**) As in **A**, but for VIP-Cre x Ai148 mice that expressed GCaMP6 in VIP neurons.

(**I**) As in **A**, but for SST-Cre mice that expressed GCaMP6 in SST neurons.

(**J**) Similarity of the average closed and open loop locomotion onset responses in V1 quantified as the
correlation coefficient between the two in a window -5 s to +3 s around locomotion onset (left and
right V1 were averaged, see Methods). Error bars indicate SEM over mice that had an average
locomotion onset response of at least 1% ΔF/F in either the closed or the open loop condition (open
circles: individual data points; 6 C57BL/6 mice, 4 Emx1-Cre mice, 7 Scnn1a-Cre mice, 14 Tlx3-Cre
mice, 3 Ntsr1-Cre mice, 2 PV-Cre mice, 6 VIP-Cre mice, 6 Sst-Cre mice, 3 Cux2-CreERT2 mice).
Statistical comparisons are against the Tlx3 data and were corrected for family-wise error rate (see
Methods and **Table S1**): adjusted significance thresholds, n.s.: not significant, *: $p < 0.05/9$, **: $p <$
$0.01/9$, ***: $p < 0.001/9$. See **Table S1** for all information on statistical testing.

Figure S5. Additional information for antipsychotic drug data. Related to Figure 4.

(**A**) Average hemodynamic response during locomotion onsets in Tlx3-Cre mice that had been retro-
orbitally injected with an AAV-PHP.eB-DIO-eGFP to express eGFP in L5 IT neurons (responses were
averaged across corresponding regions in both hemispheres) after a single injection of the
antipsychotic drug clozapine (see Methods). Mean (lines) and 90% confidence interval (shading) are
calculated as a hierarchical bootstrap estimate for each time bin (see Methods and **Table S1**). Inset:
Comparison of responses to locomotion onsets in the same mice before clozapine injection (same
data as in **Figure S1C**, top panel).

(**B**) Average locomotion speed in 21 naive mice (black) and 22 mice that had been injected with
either of the antipsychotic drugs clozapine, aripiprazole, or haloperidol (gray), for the data
acquisition timepoint +1 h after injection. Shading indicates SEM over mice.

(**C**) Fraction of time spent locomoting above threshold (see Methods) for the 21 naive mice (black)
and 22 mice that had been injected with either of the antipsychotic drugs clozapine, aripiprazole or
haloperidol, for the data acquisition timepoint +1 h after injection (gray). Boxes represent the
median and the interquartile range. The open circles indicate the mean of the distributions. The
whiskers mark 1.5 times the interquartile range. Dots are the individual data points.

(**D**) Average correlation of activity and locomotion speed in dorsal cortex was increased after a single
injection of the antipsychotic drug clozapine in 4 C57BL/6 mice that expressed GCaMP brain wide
but decreased in 5 Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons, for all types of
visual feedback. Error bars indicate SEM over mice and corresponding dorsal cortex regions (see
Methods). n.s.: not significant, *: $p < 0.05$, ***: $p < 0.001$. See **Table S1** for all information on
statistical testing.

(**E**) Average response of V1 recorded with the widefield microscope during locomotion onsets in
closed loop (solid) or open loop (dashed) in Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT
neurons and that had received an injection of the antipsychotic drug clozapine. Mean (lines) and

90% confidence interval (shading) are calculated as a hierarchical bootstrap estimate for each time
bin (see Methods and **Table S1**).

(**F**) As in **E**, for the same 3 Tlx3 x Ai148 mice, but for responses recorded at L5 IT soma with two-
photon imaging. Shading indicates SEM over 5595 neurons.

Figure S6. Calculation of the distance-correlation heatmaps. Related to Figure 5.

(**A**) For each pair of dorsal cortex regions, we calculated the distance between the regions in a top-down view of dorsal cortex, normalized by the distance between bregma and lambda for each mouse (data are from an example C57BL/6 mouse that expressed GCaMP6 brain wide).

(**B**) For each pair, we then plotted the activity correlation against this distance as calculated in **A** (black dots). We then interpolated this distribution using a 40-by-40 2D grid to obtain the density plots (faded background), shown for all mice and pairs of regions in **Figures 6, 7, and S6-S8**.

(**C**) Density map of correlation coefficients as a function of distance between the regions, normalized across mice by the bregma-lambda distance (see panel **B** and Methods), for 6 Tlx3-Cre mice that had been retro-orbitally injected with an AAV-PHP.eB-DIO-eGFP to express eGFP in L5 IT neurons. Peak density is indicated at the bottom right of the plot and corresponds to the maximum value of the color scale (dark red). The white line is a contour line drawn at 50% of peak value.

(**D**) As in **C**, but for the same 6 Tlx3-Cre mice that expressed eGFP in L5 IT neurons after a single

injection of the antipsychotic drug clozapine. The white line is the contour from panel **C** for

comparison.

(**E**) Clozapine induced change in the fluorescence correlation between regions, normalized to the

correlation coefficient in the naive state, in 6 Tlx3-Cre mice that expressed eGFP in L5 IT neurons.

Data were split into short- and long-range activity correlations (see Methods). Boxes represent the

median and the interquartile range. The open circle indicates the mean of the distribution. The

whiskers mark 1.5 times the interquartile range. Dots are the individual data points (short-range:

204 pairs of regions, long-range: 192 pairs of regions). ***: p < 0.001, n.s.: not significant. See **Table**

S1 for all information on statistical testing.

(**F**) As in **C**, but for 3 Tlx3-Cre x Ai148 mice that expressed GCaMP6 in L5 IT neurons.

(**G**) As in **F**, but after a single injection of saline. The white line is the contour line from panel **F** for

comparison.

(**H**) Saline induced change in the activity correlation between regions, normalized to the correlation

coefficient in the naive state, in 3 Tlx3-Cre x Ai148 mice. Data were split into short- and long-range

activity correlations (see Methods). Boxes represent the median and the interquartile range. The

open circle indicates the mean of the distribution. The whiskers mark 1.5 times the interquartile

range. Dots are the individual data points (short-range: 93 pairs of regions, long-range: 105 pairs of

regions). n.s.: not significant. See **Table S1** for all information on statistical testing.

Figure S7. The clozapine induced decorrelation was not present in local somatic correlations.
Related to Figure 6.

(**A**) Density map of correlation coefficients as a function of distance between the regions, normalized
across mice by the bregma-lambda distance (see **Figures S6A and S6B**, and Methods), for 3 Tlx3-Cre
x Ai148 mice that expressed GCaMP6 in L5 IT neurons. Peak density is indicated at the bottom right
of the plot and corresponds to the maximum value of the color scale (dark red). The white line is a
contour line drawn at 50% of peak value.

(**B**) As in **A**, but for the same 3 Tlx3-Cre x Ai148 mice after a single injection of the antipsychotic drug
clozapine. The white line is the contour line from panel **A** for comparison.

(**C**) Clozapine induced change in the activity correlation between regions, normalized to the
correlation coefficient in the naive state, in 3 Tlx3-Cre x Ai148 mice. Data were split into short- and
long-range activity correlations (see Methods). Boxes represent the median and the interquartile
range. The open circle indicates the mean of the distribution. The whiskers mark 1.5 times the
interquartile range. Dots are the individual data points (short-range: 114 pairs of regions, long-range:
84 pairs of regions). ***, p < 0.001. See **Table S1** for all information on statistical testing.

(**D**) Distribution of pairwise neuronal activity correlations recorded with two-photon imaging at the
L5 soma, in the same 3 Tlx3-Cre x Ai148 mice as in panels **A-C**, before (naive, black, 1991 neurons
and 384 108 pairs of neurons) and after injection of the antipsychotic drug clozapine (dark gray,
7863 neurons and 885 596 pairs of neurons or pale gray, 1 field of view excluded, 7534 neurons and
831 640 pairs of neurons remaining). Dashed lines mark the mean of the respective distributions as
indicated by the arrows on top. Note that after clozapine injection, we observed aberrantly high
neuronal synchrony in 1 out of 25 recorded fields of view that substantially increased the average

763   pairwise correlation as indicated by the two curves that compare the clozapine data. In either
764   comparison, clozapine significantly increased local pairwise activity correlations.

Figure S8. The clozapine induced decorrelation of dorsal cortex activity was weaker in superficial cortical layers than in L5. Related to Figure 6.

(**A**) Average correlation of activity between the 12 regions of interest in 4 Cux2-CreERT2 x Ai148 mice that expressed GCaMP6 in upper layer excitatory neurons.

(**B**) As in **A**, but for the same 4 Cux2-CreERT2 x Ai148 mice after a single injection of the antipsychotic drug clozapine.

(**C**) Density map of correlation coefficients as a function of distance between the regions, normalized across mice by the bregma-lambda distance (see **Figures S6A and S6B**, and Methods), for 4 naive Cux2-CreERT2 x Ai148 mice that expressed GCaMP6 in upper layer excitatory neurons. Peak density is indicated at the bottom right of the plot and corresponds to the maximum value of the color scale (dark red). The white line is a contour line drawn at 50% of peak value.

(**D**) As in **C**, but for the same 4 Cux2-CreERT2 x Ai148 mice after a single injection of the antipsychotic drug clozapine. The white line is the contour line from panel **C** for comparison.

(**E**) Clozapine induced change in the activity correlation between regions, normalized to the correlation coefficient in the naive state, in 4 Cux2-CreERT2 x Ai148 mice. Data were split into short- and long-range activity correlations (see Methods). Boxes represent the median and the interquartile range. The open circle indicates the mean of the distribution. The whiskers mark 1.5 times the interquartile range. Dots are the individual data points (short-range: 137 pairs of regions, long-range: 127 pairs of regions). n.s.: not significant, *: $p < 0.05$, **: $p < 0.01$. See **Table S1** for all information on statistical testing.

     SUPPLEMENTAL TABLES

787   **Table S1. Statistical comparisons for bar and violin plots.**

788   Note, for **Figures 1L and S4J**, the significance thresholds were corrected for multiple comparisons

789   (see Methods).

Figure reference	Description (N$_1$ vs N$_2$)	Test	N$_1$ (mice)	N$_2$ (mice)	Unit	P-value
Figure 1F	Number of closed loop locomotion onsets in C57BL/6 mice	-	907 (6)	-	Onsets	-
Figure 1G	Number of open loop locomotion onsets in C57BL/6 mice	-	598 (6)	-	Onsets	-
Figure 1H	Number of open loop visual flow onsets in C57BL/6 mice	-	416 (6)	-	Onsets	-
Figure 1I	Number of closed loop locomotion onsets in Tlx3-Cre x Ai148 mice	-	1919 (15)	-	Onsets	-
Figure 1J	Number of open loop locomotion onsets in Tlx3-Cre x Ai148 mice	-	1125 (15)	-	Onsets	-
Figure 1K	Number of open loop visual flow onsets in Tlx3-Cre x Ai148 mice	-	1189 (15)	-	Onsets	-
Figure 1L	Tlx3 vs C57BL/6	Bootstrap	12 (15)	12 (6)	Regions	<10^{-4}
	Tlx3 vs Emx1	Bootstrap	12 (15)	12 (4)	Regions	0.00135
	Tlx3 vs Cux2	Bootstrap	12 (15)	12 (4)	Regions	<10^{-4}
	Tlx3 vs Scnn1a	Bootstrap	12 (15)	12 (7)	Regions	<10^{-4}
	Tlx3 vs Ntsr1	Bootstrap	12 (15)	12 (3)	Regions	0.4541
	Tlx3 vs PV	Bootstrap	12 (15)	12 (2)	Regions	<10^{-4}
	Tlx3 vs VIP	Bootstrap	12 (15)	12 (6)	Regions	0.0001
	Tlx3 vs SST	Bootstrap	12 (15)	12 (5)	Regions	0.0131
	ANOVA	ANOVA	108 (54)	N/A	Regions	< 10^{-5}
Figure 2B	V1					
	Closed loop vs open loop	Hierachical bootsstrap	308 (6)	316 (6)	Recording session	0.9856
	Closed loop vs dark	Hierachical bootsstrap	308 (6)	68 (6)	Recording session	0.9631
	Open loop vs dark	Hierachical bootsstrap	316 (6)	68 (6)	Recording session	0.8968
	RSC					
	Closed loop vs open loop	Hierachical bootsstrap	308 (6)	316 (6)	Recording session	0.3423
	Closed loop vs dark	Hierachical bootsstrap	308 (6)	68 (6)	Recording session	0.1492
	Open loop vs dark	Hierachical bootsstrap	316 (6)	68 (6)	Recording session	0.2362
	V2am					
	Closed loop vs open loop	Hierachical bootsstrap	308 (6)	316 (6)	Recording session	0.4576
	Closed loop vs dark	Hierachical bootsstrap	308 (6)	68 (6)	Recording session	0.4694
	Open loop vs dark	Hierachical bootsstrap	316 (6)	68 (6)	Recording session	0.4961
	M1					
	Closed loop vs open loop	Hierachical bootsstrap	308 (6)	316 (6)	Recording session	0.4417

		Closed loop vs dark	Hierachical bootsstrap	308 (6)	68 (6)	Recording session	0.06
		Open loop vs dark	Hierachical bootsstrap	316 (6)	68 (6)	Recording session	0.0788
	A24b						
		Closed loop vs open loop	Hierachical bootsstrap	308 (6)	316 (6)	Recording session	0.8310
		Closed loop vs dark	Hierachical bootsstrap	308 (6)	68 (6)	Recording session	0.2194
		Open loop vs dark	Hierachical bootsstrap	316 (6)	68 (6)	Recording session	0.0846
	M2						
		Closed loop vs open loop	Hierachical bootsstrap	308 (6)	316 (6)	Recording session	0.4593
		Closed loop vs dark	Hierachical bootsstrap	308 (6)	68 (6)	Recording session	0.2373
		Open loop vs dark	Hierachical bootsstrap	316 (6)	68 (6)	Recording session	0.2674
Figure 2D	V1						
		Closed loop vs open loop	Hierachical bootsstrap	420 (15)	394 (15)	Recording session	$<10^{-4}$
		Closed loop vs dark	Hierachical bootsstrap	420 (15)	194 (15)	Recording session	$<10^{-4}$
		Open loop vs dark	Hierachical bootsstrap	394 (15)	194 (15)	Recording session	0.2103
	RSC						
		Closed loop vs open loop	Hierachical bootsstrap	420 (15)	394 (15)	Recording session	$<10^{-4}$
		Closed loop vs dark	Hierachical bootsstrap	420 (15)	194 (15)	Recording session	0.0001
		Open loop vs dark	Hierachical bootsstrap	394 (15)	194 (15)	Recording session	0.0021
	V2am						
		Closed loop vs open loop	Hierachical bootsstrap	420 (15)	394 (15)	Recording session	$<10^{-4}$
		Closed loop vs dark	Hierachical bootsstrap	420 (15)	194 (15)	Recording session	$<10^{-4}$
		Open loop vs dark	Hierachical bootsstrap	394 (15)	194 (15)	Recording session	0.1337
	M1						
		Closed loop vs open loop	Hierachical bootsstrap	420 (15)	394 (15)	Recording session	$<10^{-4}$
		Closed loop vs dark	Hierachical bootsstrap	420 (15)	194 (15)	Recording session	0.3115
		Open loop vs dark	Hierachical bootsstrap	394 (15)	194 (15)	Recording session	0.0002
	A24b						
		Closed loop vs open loop	Hierachical bootsstrap	420 (15)	394 (15)	Recording session	$<10^{-4}$
		Closed loop vs dark	Hierachical bootsstrap	420 (15)	194 (15)	Recording session	0.3511
		Open loop vs dark	Hierachical bootsstrap	394 (15)	194 (15)	Recording session	0.0001
	M2						
		Closed loop vs open loop	Hierachical bootsstrap	420 (15)	394 (15)	Recording session	0.0005
		Closed loop vs dark	Hierachical bootsstrap	420 (15)	194 (15)	Recording session	0.2686

	Open loop vs dark	Hierachical bootsstrap	394 (15)	194 (15)	Recording session	0.0156
Figure 3B	Number of mismatch onsets in C57BL/6 mice	-	2512 (6)	-	Onsets	-
Figure 3C	Number of grating onsets in C57BL/6 mice	-	1858 (6)	-	Onsets	-
Figure 3E	Number of mismatch onsets in Tlx3-Cre x Ai148 mice	-	3297 (15)	-	Onsets	-
Figure 3F	Number of mismatch onsets in Tlx3-Cre x Ai148 mice	-	5318 (15)	-	Onsets	-
Figure 4A	Number of closed loop locomotion onset in clozapine treated C57BL/6 mice	-	273 (4)	-	Onsets	-
Figure 4A, inset	Number of closed loop locomotion onset in naïve C57BL/6 mice	-	720 (4)	-	Onsets	-
Figure 4B	Number of open loop locomotion onset in clozapine treated C57BL/6 mice	-	145 (4)	-	Onsets	-
Figure 4B, inset	Number of open loop locomotion onset in naive C57BL/6 mice	-	570 (4)	-	Onsets	-
Figure 4C	Number of open loop visual flow onsets in clozapine treated C57BL/6 mice that were not locomoting	-	99 (4)	-	Onsets	-
Figure 4C, inset	Number of open loop visual flow onsets in naïve C57BL/6 mice that were not locomoting	-	404 (4)	-	Onsets	-
Figure 4D	Number of closed loop locomotion onset in clozapine treated Tlx3-Cre x Ai148 mice	-	707 (5)	-	Onsets	-
Figure 4D, inset	Number of closed loop locomotion onset in naive Tlx3-Cre x Ai148 mice	-	1101 (5)	-	Onsets	-
Figure 4E	Number of open loop locomotion onset in clozapine treated Tlx3-Cre x Ai148 mice	-	350 (5)	-	Onsets	-
Figure 4E, inset	Number of open loop locomotion onset in naive Tlx3-Cre x Ai148 mice	-	348 (5)	-	Onsets	-
Figure 4F	Number of open loop visual flow onsets in clozapine treated Tlx3-Cre x Ai148 mice that were not locomoting	-	514 (5)	-	Onsets	-
Figure 4F, inset	Number of open loop visual flow onsets in naive Tlx3-Cre x Ai148 mice that were not locomoting	-	568 (5)	-	Onsets	-
Figure 6C	Short-range vs no change	T-test	122 (4)	-	Pairs of regions	0.3791
	Long-range vs no change	T-test	142 (4)	-	Pairs of regions	0.9035
	Short-range vs long-range	Ranksum	122 (4)	142 (4)	Pairs of regions	0.0511
Figure 6F	Short-range vs no change	T-test	182 (5)	-	Pairs of regions	$<10^{-5}$
	Long-range vs no change	T-test	148 (5)	-	Pairs of regions	$<10^{-10}$
	Short-range vs long-range	Ranksum	182 (5)	148 (5)	Pairs of regions	$<10^{-4}$
Figure 7C	Short-range vs no change	T-test	114 (3)	-	Pairs of regions	$<10^{-5}$
	Long-range vs no change	T-test	84 (3)	-	Pairs of regions	$<10^{-7}$
	Short-range vs long-range	Ranksum	114 (3)	84 (3)	Pairs of regions	0.0347
Figure 7F	Short-range vs no change	T-test	114 (3)	-	Pairs of regions	0.7058
	Long-range vs no change	T-test	84 (3)	-	Pairs of regions	0.0435

	Short-range vs long-range	Ranksum	114 (3)	84 (3)	Pairs of regions	0.0199
Figure 7I	Short-range vs no change	T-test	114 (3)	-	Pairs of regions	0.4865
	Long-range vs no change	T-test	84 (3)	-	Pairs of regions	0.2184
	Short-range vs long-range	Ranksum	114 (3)	84 (3)	Pairs of regions	0.1633
Figure 8A	Number of mismatch onsets in naïve Tlx3-Cre x Ai148 mice	-	2464 (11)	-	Onsets	-
	Number of mismatch onsets in Tlx3-Cre x Ai148 mice treated with antipsychotic drugs	-	2017 (11)	-	Onsets	-
Figure 8B	Number of grating onsets in naïve Tlx3-Cre x Ai148 mice	-	3942 (11)	-	Onsets	-
	Number of grating onsets in Tlx3-Cre x Ai148 mice treated with antipsychotic drugs	-	1645 (11)	-	Onsets	-
Figure S1A, left	Number of locomotion onsets in C57BL/6 mice that expressed GFP brain wide and were implanted with crystal skulls	-	96 (3)	-	Onsets	-
Figure S1A, right	Number of locomotion onsets in C57BL/6 mice that expressed GFP brain wide with clear skull cranial windows	-	615 (5)	-	Onsets	-
Figure S1B, left	Number of mismatch onsets in C57BL/6 mice that expressed GFP brain wide and were implanted with crystal skulls	-	229 (3)	-	Onsets	-
Figure S1B, right	Number of mismatch onsets in C57BL/6 mice that expressed GFP brain wide with clear skull cranial windows	-	228 (5)	-	Onsets	-
Figure S1C	Number of locomotion onsets in Tlx3-Cre mice that expressed GFP in L5 IT neurons	-	1880 (7)	-	Onsets	-
Figure S1D	Number of mismatch onsets in Tlx3-Cre mice that expressed GFP in L5 IT neurons	-	451 (7)	-	Onsets	-
Figure S3A	Number of closed loop locomotion onsets in Tlx3-Cre x Ai148 mice recorded with the widefield microscope	-	1012 (7)	-	Onsets	-
	Number of open loop locomotion onsets in Tlx3-Cre x Ai148 mice recorded with the widefield microscope	-	678 (7)	-	Onsets	-
Figure S3C	Number of mismatch onsets in Tlx3-Cre x Ai148 mice recorded with the widefield microscope	-	1038 (7)	-	Onsets	-
	Number of open loop halts in Tlx3-Cre x Ai148 mice recorded with the widefield microscope	-	410 (7)	-	Onsets	-
	Number of grating onsets in Tlx3-Cre x Ai148 mice recorded with the widefield microscope	-	2014 (7)	-	Onsets	-
Figure S4A	Number of closed loop locomotion onsets in Emx1-Cre mice	-	438 (4)	-	Onsets	-
	Number of open loop locomotion onsets in Emx1-Cre mice	-	286 (4)	-	Onsets	-
Figure S4B	Number of closed loop locomotion onsets in Cux2-CreERT2 x Ai148 mice	-	415 (4)	-	Onsets	-
	Number of open loop locomotion onsets in Cux2-CreERT2 x Ai148 mice	-	433 (4)	-	Onsets	-
Figure S4C	Number of closed loop locomotion onsets in Scnn1a x Ai148 mice	-	839 (7)	-	Onsets	-
	Number of open loop locomotion onsets in Scnn1a x Ai148 mice	-	313 (7)	-	Onsets	-

Figure S4D	Number of closed loop locomotion onsets in Tlx3-Cre x Ai148 mice	-	1919 (15)	-	Onsets	-
	Number of open loop locomotion onsets in Tlx3-Cre x Ai148 mice	-	1125 (15)	-	Onsets	-
Figure S4E	Number of closed loop locomotion onsets in Ntsr1-Cre x Ai148 mice	-	368 (3)	-	Onsets	-
	Number of open loop locomotion onsets in Ntsr1-Cre x Ai148 mice	-	112 (3)	-	Onsets	-
Figure S4F	Number of closed loop locomotion onsets in C57BL/6 mice	-	907 (6)	-	Onsets	-
	Number of open loop locomotion onsets in C57BL/6 mice	-	598 (6)	-	Onsets	-
Figure S4G	Number of closed loop locomotion onsets in PV-Cre x Ai148 mice	-	236 (2)	-	Onsets	-
	Number of open loop locomotion onsets in PV-Cre x Ai148 mice	-	110 (2)	-	Onsets	-
Figure S4H	Number of closed loop locomotion onsets in VIP-Cre x Ai148 mice	-	618 (6)	-	Onsets	-
	Number of open loop locomotion onsets in VIP-Cre x Ai148 mice	-	376 (6)	-	Onsets	-
Figure S4I	Number of closed loop locomotion onsets in Sst-Cre x Ai148 mice	-	747 (5)	-	Onsets	-
	Number of open loop locomotion onsets in Sst-Cre x Ai148 mice	-	558 (5)	-	Onsets	-
Figure S4J	Tlx3 vs C57BL/6	Bootstrap	14	6	V1 / Mice	$<10^{-4}$
	Tlx3 vs Emx1	Bootstrap	14	4	V1 / Mice	0.0042
	Tlx3 vs Cux2	Bootstrap	14	3	V1 / Mice	$<10^{-4}$
	Tlx3 vs Scnn1a	Bootstrap	14	7	V1 / Mice	0.0042
	Tlx3 vs Ntsr1	Bootstrap	14	3	V1 / Mice	0.4087
	Tlx3 vs PV	Bootstrap	14	2	V1 / Mice	$<10^{-4}$
	Tlx3 vs VIP	Bootstrap	14	6	V1 / Mice	0.0021
	Tlx3 vs SST	Bootstrap	14	6	V1 / Mice	0.0032
	ANOVA	ANOVA	54	N/A	V1 / Mice	0.0004
Figure S5A	Number of locomotion onsets in Tlx3-Cre mice	-	2331 (7)	-	Onsets	-
Figure S5C	Naive vs +1 h antipsy.	Ranksum	21	22	Mice	0.4735
Figure S5D	C57BL/6					
	Closed loop vs no change	T-test	24 (4)	-	Regions	0.0277
	Open loop vs no change	T-test	24 (4)	-	Regions	0.6384
	Dark vs no change	T-test	24 (4)	-	Regions	$<10^{-4}$
	Tlx3					
	Closed loop vs no change	T-test	30 (5)	-	Regions	0.0365
	Open loop vs no change	T-test	30 (5)	-	Regions	0.0004
	Dark vs no change	T-test	30 (5)	-	Regions	0.0104
Figure S5E	Number of closed loop locomotion onsets in Tlx3-Cre x Ai148 mice	-	376 (3)	-	Onsets	-
	Number of open loop locomotion onsets in Tlx3-Cre x Ai148 mice	-	245 (3)	-	Onsets	-
Figure S6E	Short-range vs no change	T-test	204 (7)	-	Pairs of regions	0.0006
	Long-range vs no change	T-test	192 (7)	-	Pairs of regions	$<10^{-4}$
	Short-range vs long-range	Ranksum	204 (7)	192 (7)	Pairs of regions	0.2540
Figure S6H	Short-range vs no change	T-test	93 (3)	-	Pairs of regions	0.7884
	Long-range vs no change	T-test	105 (3)	-	Pairs of regions	0.1176

	Short-range vs long-range	Rank-sum	93 (3)	105 (3)	Pairs of regions	0.2319
Figure S7C	Short-range vs no change	T-test	114 (3)	-	Pairs of regions	$<10^{-4}$
	Long-range vs no change	T-test	84 (3)	-	Pairs of regions	$<10^{-4}$
	Short-range vs long-range	Ranksum	114 (3)	84 (3)	Pairs of regions	$<10^{-4}$
Figure S7D	Naive vs clozapine	Ranksum	384108 (7)	885596	Pairs of neurons	$<10^{-4}$
	Naive vs clozapine 1 site excl.	Ranksum	384108 (7)	831640	Pairs of neurons	$<10^{-4}$
Figure S8E	Short-range vs no change	T-test	137 (4)	-	Pairs of regions	0.0262
	Long-range vs no change	T-test	127 (4)	-	Pairs of regions	0.0019
	Short-range vs long-range	Ranksum	137 (4)	127 (4)	Pairs of regions	0.9537

792 **Table S2. Indicator expression strategy in individual mice.**

Nbr. mice	Genotype	Virus	Figures
3	C57BL/6	AAV-PHP.eB-hSyn1-jGCaMP7f	**1C, 1F-1H, 1L, 2A and 2B, 3A-3C, 4A-4C, 5A and 5B, 6A-6C, S2, S4F, S4J, S5B-S5D, S6A and S6B**
3	C57BL/6	AAV-PHP.eB-EF1α-GCaMP6s	**1F-1H, 1L, 2B, 3B and 3C, 4A-4C, 5A and 5B, 6A-6C, S2, S4F, S4J, S5B-S5D**
8	C57BL/6	AAV-PHP.eB-EF1α-eGFP	**S1**
4	Emx1-Cre	AAV-PHP.eB-DIO-EF1α-GCaMP6s	**1L, S4A and S4J**
4	Cux2-CreERT2 x Ai148	-	**1L, S4B, S4J and S8**
7	Scnn1a-Cre x Ai148	-	**1L, S4C and S4J**
25	Tlx3-Cre x Ai148	-	**1D, 1I-1L, 2C and 2D, 3D-3F, 4D-4F, 5C and 5D, 6D-6F, 7, 8, S2, S3D, S4D, S4J, S5B-S5F, S6G-S6H, S7**
7	Tlx3-Cre	AAV-PHP.eB-EF1α-DIO-eGFP	**S1C and S1D, S5A, S6C-S6E**
3	Ntsr1-Cre x Ai148	-	**1L, S4E and S4J**
2	PV-Cre x Ai148	-	**1L, S4G and S4J**
6	VIP-Cre x Ai148	-	**1L, S4H and S4J**
4	SST-Cre	AAV-PHP.eB-DIO-EF1α-jGCaMP7f	**1L, S4I and S4J**
1	SST-Cre x Ai148	-	**1L, S4I and S4J**

  METHODS

  Key Resource Table

REAGENT or RESOURCE	Source	Identifier
Bacterial and virus strains		
AAV-PHP.eB-hSyn1-jGCaMP7f (10^{13} GC/ml)	FMI vector core	vector.fmi.ch
AAV-PHP.eB-EF1α-GCaMP6s (10^{13} GC/ml)	FMI vector core	vector.fmi.ch
AAV-PHP.eB-DIO-EF1α-GCaMP6s (10^{13} GC/ml)	FMI vector core	vector.fmi.ch
AAV-PHP.eB-DIO-EF1α-jGCaMP7f (10^{12} GC/ml)	FMI vector core	vector.fmi.ch
AAV-PHP.eB-EF1α-eGFP (10^{13} GC/ml)	FMI vector core	vector.fmi.ch
AAV-PHP.eB-EF1α-DIO-eGFP (10^{15} GC/ml)	FMI vector core	vector.fmi.ch
Chemicals, peptides, and recombinant proteins		
Fentanyl citrate	Actavis	CAS 990-73-8
Midazolam (Dormicum)	Roche	CAS 59467-96-8
Medetomidine (Domitor)	Orion Pharma	CAS 86347-14-0
Ropivacaine	Presenius Kabi	CAS 132112-35-7
Lidocaine	Bichsel	CAS 137-58-6
Buprenorphine	Reckitt Benckiser Healthcare	CAS 52485-79-7
Ophthalmic gel (Humigel)	Virbac	N/A
Flumazenil (Anexate)	Roche	CAS 78755-81-4
Atipamezole (Antisedan)	Orion Pharma	CAS 104054-27-5
N-Butyl-2-cyanoacrylate (Histoacryl)	Braun	CAS 6606-65-1
Dental cement (Paladur)	Heraeus Kulzer	CAS 9066-86-8
Metacam	Boehringer Ingelheim	CAS 71125-39-8
Clozapine (powder)	Novartis	CAS 5786-21-0
Aripiprazole (intramuscular injection solution 7.5 mg/ml)	Otsuka Pharmaceutical	CAS 129722-12-9
Haloperidol (intramuscular injection solution 5 mg/ml)	Janssen	CAS 52-86-8
Amphetamine (powder)	Hänseler	CAS 60-13-9
Tamoxifen food	Envigo	CAS 10540-29-1 (TD.55125)
Deposited data		
All data and code used to generate manuscript figures	This paper	data.fmi.ch
Experimental models: Organisms/strains		
Mus musculus: C57BL/6	Charles River	N/A
Mus musculus: Emx1^{tm1(cre)Krj}	Jackson Laboratories	RRID:IMSR_JAX:005628
Mus musculus: B6(Cg)-Cux2^{tm3.1(cre/ERT2)Mull}/Mmmh	MMRRC	RRID:MMRRC_032779-MU
Mus musculus: Scnn1a-cre3Aibs/J	Jackson Laboratories	RRID:IMSR_JAX:009613
Mus musculus: Tg(Tlx3-cre)PL56Gsat/Mmucd	MMRRC	RRID:MMRRC_041158-UCD
Mus musculus: Ntsr1^{GN220Gsat/Mmucd}	MMRRC	RRID:MMRRC_017266-UCD
Mus musculus: Pvalb^{tm1(cre)Arbr}	Jackson Laboratories	RRID:IMSR_JAX:008069
Mus musculus: Vip^{tm1(cre)Zjh}	Jackson Laboratories	RRID:IMSR_JAX:010908
Mus musculus: Sst^{tm2.1(cre)Zjh}	Jackson Laboratories	RRID:IMSR_JAX:018973
Mus musculus: Igs7^{tm148.1(tetO-GCaMP6f,CAG-tTA2)Hze}/J	Jackson Laboratories	RRID:IMSR_JAX:030328
Software and algorithms		
MATLAB (2021a)	The MathWorks	RRID:SCR_001622
LabVIEW	National Instruments	RRID:SCR_014325
Python	python.org	RRID:SCR_008394
Panda3D	panda3d.org	N/A

Mice. All animal procedures were approved by and carried out in accordance with guidelines of the Veterinary Department of the Canton Basel-Stadt, Switzerland. The mice used for cell-type specfic targeting and subsequent widefield imaging in this study were kept on a C57BL/6 background and were of the following genotype: 6 C57BL/6 (Charles River Laboratories), 4 Emx1-Cre to target expression to cortical excitatory neurons (Gorski et al., 2002), 4 Cux2-CreERT2 to target expression to L2/3 and L4 excitatory neurons (Franco et al., 2012), 7 Scnn1a-Cre to target expression to a subset of L4 excitatory neurons (Madisen et al., 2010), 25 Tlx3-Cre to target expression to a subset of L5 IT neurons (Gerfen et al., 2013), 3 Ntsr1-Cre to target expression to a subset of L6 excitatory neurons (Gong et al., 2007), and, to target inhibitory subpopulations, we used 2 PV-Cre (Hippenmeyer et al., 2005), 6 VIP-Cre (Taniguchi et al., 2011), and 5 SST-Cre (Taniguchi et al., 2011) mice. Ai148 ((Daigle et al., 2018), Jackson Laboratories, stock #030328) mice were used as breeders to drive GCaMP6f expression in Cre positive neurons. An additional 8 C57BL/6 and 7 Tlx3-Cre mice were used in experiments controlling for hemodynamic responses. To induce expression of Cre in the Cux2-CreERT2 line, mice were provided for at least 1 week with tamoxifen containing food (Envigo, 400 mg tamoxifen per kg of food) as their sole food source. Mice were group housed in a vivarium (light/dark cycle: 12/12 hours). Experimental mice were of either sex.

Surgery and virus injections. For all surgical procedures, mice were anesthetized using a mixture of fentanyl (0.05 mg/kg; Actavis), midazolam (5.0 mg/kg; Dormicum, Roche), and medetomidine (0.5 mg/kg; Domitor, Orion). In a subset of mice (6 C57BL/6, 4 Emx1-Cre, and 4 SST-Cre mice), we injected an AAV vector based on the PHP.eB capsid (Chan et al., 2017) retro-orbitally (6 µl each eye of at least 10^{13} GC/ml) to drive expression of GCaMP under either the EF1α or hSyn1 promoter pan-neuronally, or in a cell type specific manner using a double-floxed inverted open reading frame (DIO) construct. For hemodynamic response control experiments, we used an AAV-PHP.eB-EF1α-eGFP to pan-neuronally express eGFP (**Figures S1A and S1B)** or an AAV-PHP.eB-EF1α-DIO-eGFP to restrict eGFP expression to L5 IT neurons in Tlx3-Cre mice (**Figures S1C and S1D, S5A, S6C-S6E**). To improve optical access to the cortex, we implanted crystal skull cranial windows (Kim et al., 2016). Prior to removing the skull plate overlying dorsal cortex, we recorded the location of bregma relative to other landmarks on the skull. Superglue (Pattex) was used to glue the crystal skull in place in the craniotomy. In a subset of mice, for the comparison of hemodynamic responses (**Figure S1A and S1B**), we used a clear skull preparation (Guo et al., 2014). To do this, we attached the crystal skull coverslip directly onto the cleaned skull surface with a three-component polymer (C&B Metabond, Parkell). A custom-machined titanium head bar was attached to the skull using dental cement (Paladur, Heraeus). An epifluorescence overview image was taken to mark reference points on the

dorsal cortical surface. Anesthesia was antagonized by an intraperitoneal injection of a mixture of

flumazenil (0.5 mg/kg; Anexate, Roche) and atipamezole (2.5 mg/kg; Antisedan, Orion Pharma). For

peri- and post-operative analgesia we injected buprenorphine (0.1 mg/kg; Reckitt Benckiser

Healthcare (UK) Ltd.) and Metacam (5 mg/kg; Boehringer Ingelheim). Imaging commenced at the

earliest 1 week after head bar implantation or 3 weeks after retro-orbital AAV injection. Clozapine

(Novartis), aripiprazole (Otsuka Pharmaceutical), haloperidol (Janssen), amphetamine (Hänseler), or

saline (0.9% NaCl in H_2O) was intraperitoneally injected at 0.2 μg/g - 10 μg/g, 0.2 μg/g, 0.1 μg/g, 4

μg/g body weight, or undiluted, respectively.

Virtual reality setup and stimulus design. For all experiments we used a virtual reality setup as

previously described (Leinweber et al., 2014). Mice were head-fixed and free to run on a spherical,

air-supported Styrofoam ball. A virtual corridor was projected (using a Samsung SP-F10M projector)

onto a toroidal screen positioned in front of the mouse covering a field of view of approximately 240

degrees horizontally and 100 degrees vertically. Recordings were blocked into 5-minute sessions.

Experiments typically commenced with 4 sessions of a closed loop condition in which the rotation of

the spherical treadmill was coupled to the movement in a virtual corridor. To introduce mismatches,

we broke the coupling between locomotion and visual flow by briefly halting visual flow for 1 s at

random times. Closed loop sessions were followed by 4 sessions of an open loop condition, in which

rotation of the spherical treadmill and movement in the virtual corridor were decoupled. The visual

stimulus consisted of a replay of the visual flow recorded in the previous closed loop condition.

Following this we recorded 2 sessions of what we refer to as a dark condition in which the virtual

reality was switched off. Please note, this was not complete darkness, primarily because the

shielding of the blue excitation light was not light proof. Finally, we recorded activity in 2 sessions of

a grating condition in which we presented drifting grating stimuli covering the full toroidal screen

placed in front of the mouse (8 directions of 6 s ± 2 s (mean ± SD) duration each with an inter-grating

interval of 4.5 s ± 1.5 s (mean ± SD) gray screen, presented in randomized order). These sessions,

totaling approximately 1 hour of recording time per day, were acquired for 3 consecutive days. For

mice in which we went on to test the effect of drugs on dorsal cortex activity, the first post-drug

injection data were acquired +1 h after drug injection (+30 mins in the case of amphetamine because

of faster onset kinetics of the drug) and further data were collected at the +24 h and +48 h

timepoints. Data in the manuscript are shown as the combined data over either all data collected

before drug injection (labelled as naive) or after substance injection (labelled as clozapine,

aripiprazole, haloperidol, amphetamine, or saline, respectively).

Data acquisition. All widefield imaging experiments were performed on a custom-built macroscope

consisting of commercially available objectives mounted face-to-face (Nikon 85 mm/f1.8 sample

side, Nikon 50 mm/f1.4 sensor side). We used a 470 nm LED (Thorlabs) powered by a custom-built

LED driver for exciting GCaMP (or eGFP) fluorescence through an excitation filter (SP490, Thorlabs)

reflected off a dichroic mirror (LP490, Thorlabs) placed in the parfocal plane of the objectives. Green

fluorescence was collected through a 525/50 emission filter on an sCMOS camera (PCO edge 4.2).

Apertures on objectives were usually kept maximally open and the current at the LED driver was

used to adjust fluorescence intensity to a value that was kept below 25% of the maximum dynamic

range of the sensor. In cases where this was not possible (e.g., transfection of eGFP yielded

extremely bright fluorescence, often visible with the naked eye), objective apertures were gradually

reduced to avoid overexposure of the sensor. LED illumination was adjusted with a collimator

(Thorlabs SM2F32-A) to achieve homogenous illumination across the surface of the cranial window.

The resulting Gaussian profile of the illumination cone was further trimmed with black tape on the

sample side objective to avoid light shining directly into the mouse's eye. An Arduino board (Arduino

Mega 2560) was used to control LED onsets synced to the frame trigger signal of the camera. The

duty cycle of the 470 nm LED was 90%. Raw images were acquired at full speed (100 Hz; except for

eGFP controls, which were acquired at 50 Hz effective frame rate) and full dynamic range (16 bit) of

the sensor. The raw images were cropped on-sensor and the resulting data was streamed to disk

with custom written software in LabVIEW (National Instruments), resulting in an effective pixel size

of 60 μm^2 at a standardized imaging resolution of 1108 pixels x 1220 pixels (1.35 MP).

Two-photon calcium imaging was performed using a custom-built microscope based on a Thorlabs B-

scope (Leinweber et al., 2014). Illumination source was a tunable femtosecond laser (Coherent

Chameleon) tuned to 930 nm. Emission light was band-pass filtered using a 525/50 filter for GCaMP

and detected using a GaAsP photomultiplier (H7422, Hamamatsu). Photomultiplier signals were

amplified (DHPCA-100, Femto), digitized (NI5772, National Instruments) at 800 MHz, and band-pass

filtered around 80 MHz using a digital Fourier-transform filter implemented in custom-written

software on an FPGA (NI5772, National Instruments). The scanning system of the microscope was

based on a 12 kHz resonant scanner (Cambridge Technology). Images were acquired at a resolution

of 750 pixels x 400 pixels at a 60 Hz frame rate and a piezo-electric linear actuator (P-726, Physik

Instrumente) was used to move the objective (Nikon 16x, 0.8 NA) in steps of 15 μm between frames

to acquire images at 4 different depths. This resulted in an effective frame rate of 15 Hz. The field of

view was 350 μm x 350 μm.

Data processing. Off-line data processing and data analysis (section below) were done with custom-

written MATLAB (2021a) scripts. For widefield macroscope imaging, raw movie data was manually

registered across days by aligning subsequent mean projections of the data to the first recorded

image sequence. We placed regions of interest (ROIs) relative to readily identifiable anatomical

landmarks that had been previously noted during cranial window surgery (see above). This resulted

in the selection of six 20 pixels x 20 pixels ROIs per hemisphere. Of those ROIs, we calculated the

activity as the $\Delta F/F_0$, wherein F_0 was the median fluorescence of the recording (approximately

30 000 frames in a 5 min recording). $\Delta F/F_0$ was corrected for slow fluorescence drift caused by

thermal brightening of the LED using 8^{th} percentile filtering with a 62.5 s moving window similar to

what was described previously for two-photon imaging (Daniel A. Dombeck et al., 2007).

Raw two-photon imaging data were full-frame registered to correct for brain motion. Neurons were

manually selected based on mean and maximum fluorescence images. Raw fluorescence traces were

corrected for slow drift in fluorescence using an 8^{th} percentile filtering with a 15 s window (Daniel A

Dombeck et al., 2007). $\Delta F/F$ traces were calculated as mean fluorescence in a selected region of

every imaging frame and subsequently subtracted and normalized by the median fluorescence.

Data analysis. Locomotion and visual flow onsets were determined based on threshold crossing of

locomotion or visual flow speed. To select only well isolated onsets, we used a speed threshold of 30

cm/s and excluded all onsets with locomotion or visual flow in the 3 s window preceding the onset.

To increase the number of locomotion onsets that passed the speed threshold criterion for the eGFP

imaging data (**Figures S1, S5A**), we excluded only those onsets if there was locomotion in the 1 s

window preceding onset.

All stimulus response curves (**Figures 1F-1L, 3B and 3C, 3E and 3F, 4, 8, S1-S4, S5A and S5B, S5E and**

S5F) were baseline subtracted. The baseline subtraction window for unpredictable stimuli

(mismatches and grating onsets) was -200 ms to 0 ms before stimulus onset. For closed loop and

open loop locomotion onsets, and open loop visual flow onsets, to accommodate calcium indicator

offset dynamics and potential anticipatory activity, we placed the baseline subtraction windows for

these onsets at -2900 ms to -2700 ms for GCaMP imaging, or -900 ms to -700 ms for eGFP imaging.

Varying this window within the bounds of offset dynamics of the indicator and anticipatory activity

onset did not change the conclusions drawn in this manuscript. Analyses in which image sequences

were aligned to event onsets (**Figures 1C-1D and 3**) used the same parameters as above.

On rare occasions, we observed seizure-like activity after antipsychotic drug injection. These events

resulted in atypically high levels of correlation, and to avoid them from influencing our conclusions

we excluded these data from further analysis. We excluded sessions if they contained an event in

which average activity of all 12 ROIs remained elevated above 30% ΔF/F for more than 10 s. This

criterion led to the exclusion of 0.26% of data (10 out of 3801 of 5 min recording sessions).

To quantify the similarity of closed loop and open loop locomotion onset responses shown in **Figure**

1L we calculated the correlation coefficient between the averaged responses (onsets and mice, 1 s

moving average) in a window -5 s to +3 s, for each ROI. For **Figure S4J**, we first averaged over onsets

in either left or right V1, which resulted in an average locomotion onset response curve for left and

right V1, for each mouse and condition (closed and open loop). These curves were subsequently

used for calculating the correlation coefficient in the same window as above. The correlation

coefficients from left and right V1 are then averaged and reported as individual data points in the

figure panel. For this analysis, we excluded mice in which the responses were smaller than 1% ΔF/F

in either of the two conditions (3 of 54 mice), as the correlation analysis yields noise in the absence

of a clear response.

For the correlation analyses shown in **Figures 5-7, S6-S8**, we first calculated the correlation

coefficient of neuronal activity (or fluorescence change, in the case of Tlx3-Cre mice that had been

retro-orbitally injected with an AAV-PHP.eB-Ef1α-DIO-eGFP to express eGFP in L5 IT neurons) for all

possible pairs of the 12 regions of interest. We then determined the distance between each pair of

ROIs and normalized this value by the distance between bregma and lambda obtained from the

images during initial surgery for each mouse (**Figure S6A**). We plotted the correlation against the

distance for each pair of regions of interest and binned the data in a 40 x 40 grid and smoothed the

resulting image with a gaussian filter (**Figure S6B**), to obtain the heatmaps shown in **Figures 6A and**

6B, 6D and 6E, 7A and 7B, 7D and 7E, 7G and 7H, S6B-S6D, S6G and S6H, S7A and S7B, S8C and S8D.

Contour lines in the heatmaps were then drawn to 50% of the maximum value in the plot. The

boxplot quantifications (**Figures 6C, 6F, 7C, 7F, 7I, S6E, S6H, S7C, S8E**) were calculated as the change

in correlation coefficient of activity after and before treatment for all pairs of regions of interest,

normalized to the value before treatment and split into short- and long-range using a threshold of

0.9 of the bregma-lambda distance for each mouse. For pairwise neuronal correlations that had

been recorded with two-photon imaging (**Figure S7D**), we calculated the correlation coefficient

separately for all pairs of neurons within each field of view. We noticed extreme synchrony of

activity of L5 IT soma recorded in 1 field of view with two-photon imaging after clozapine injection

and show both distributions for all data as well as the distribution when this 1 field of view is

excluded.

For **Figure S2,** onset aligned data were first averaged for each mouse and then summed in the given

combinations. The SEM was calculated over mice.

To obtain the differences in the correlation between activity and locomotion after and before

clozapine injection (**Figure S5D**), we first averaged all correlation coefficients in the individual

sessions of a condition (closed loop, open loop, or dark) to generate one correlation coefficient per

mouse and ROI, for each condition. We then calculated and plotted the difference between these

correlation values after and before clozapine injection.

We used a hierarchical bootstrap (**Figures 2C and 2D**), or a bootstrap (where data were not nested,

Figure 1L and S4J), distribution to estimate p-values. For hierarchical bootstrap, we first resampled

the data (with replacement) at the level of mice, and then, from the selected mice, resampled for

recording sessions. In cases where data were not nested, bootstrap samples were drawn from the

distribution of data points. We then computed the mean of this bootstrap sample and repeated this

10 000 times to generate a bootstrap distribution of the mean estimate. The p-value was then

defined as the proportion of bootstrap samples higher (or lower, depending on the hypothesis) than

zero. For **Figures 1L and S4J**, to account for multiple comparisons, we used family-wise error

correction. The significance indicators in the figure panels therefore reflect an adjusted significance

threshold, α_{adj}, according to the formula $\alpha_{adj} = \alpha/m$, wherein α is 0.05, 0.01 or 0.001, and m denotes

the number of groups (genotypes, m = 9 in both figure panels).

For widefield imaging time course data (**Figures 1F-1L, 3B and 3C, 3E and 3F, 4, 8, S1, S3A and S3C,**

S4, S5A, S5E), we used a hierarchical bootstrap estimate of the mean (Saravanan et al., 2020) and the

90% confidence interval. This was done by generating 1000 bootstrap samples chosen hierarchically

(first over mice, then over onsets) for each time bin (10 ms) in the plot.

To compare the similarity of closed and open loop locomotion onset responses, we used a one-way

analysis of variance (ANOVA) with genotype (**Figure 1L**) or mice (**Figure S4J**) as factors (**Table S1**),

respectively, followed by a bootstrap-distribution based estimate of p-values as described above.

ACKNOWLEDGEMENTS

We thank Tingjia Lu for production of the AAV vectors, Rebecca Jordan for performing the clear skull

implants used in the hemodynamic response measurements, and members of the Keller lab for

discussion and support. This project has received funding from the Swiss National Science

Foundation, the Novartis Research Foundation, and the European Research Council (ERC) under the

European Union's Horizon 2020 research and innovation programme (grant agreement No 865617).

AUTHOR CONTRIBUTIONS

MH performed the experiments and analyzed the data. All authors wrote the manuscript.

CONFLICT OF INTEREST

Part of the results presented herein have been included in patent application EP22153051.1.

REFERENCES

Allen, W.E., Kauvar, I.V., Chen, M.Z., Richman, E.B., Yang, S.J., Chan, K., Gradinaru, V., Deverman,
B.E., Luo, L., Deisseroth, K., 2017. Global Representations of Goal-Directed Behavior in
Distinct Cell Types of Mouse Neocortex. Neuron 94, 891-907.e6.
https://doi.org/10.1016/j.neuron.2017.04.017
Anderson, C.T., Sheets, P.L., Kiritani, T., Shepherd, G.M.G., 2010. Sublayer-specific microcircuits of
corticospinal and corticostriatal neurons in motor cortex. Nat Neurosci 13, 739–744.
https://doi.org/10.1038/nn.2538
Anticevic, A., Hu, X., Xiao, Y., Hu, J., Li, F., Bi, F., Cole, M.W., Savic, A., Yang, G.J., Repovs, G., Murray,
J.D., Wang, X.-J., Huang, X., Lui, S., Krystal, J.H., Gong, Q., 2015. Early-course unmedicated
schizophrenia patients exhibit elevated prefrontal connectivity associated with longitudinal
change. J Neurosci 35, 267–286. https://doi.org/10.1523/JNEUROSCI.2310-14.2015
Aravanis, A.M., Wang, L.-P., Zhang, F., Meltzer, L.A., Mogri, M.Z., Schneider, M.B., Deisseroth, K.,
2007. An optical neural interface: in vivo control of rodent motor cortex with integrated
fiberoptic and optogenetic technology. J Neural Eng 4, S143-156.
https://doi.org/10.1088/1741-2560/4/3/S02
Attinger, A., Wang, B., Keller, G.B., 2017. Visuomotor Coupling Shapes the Functional Development
of Mouse Visual Cortex. Cell 169, 1291-1302.e14. https://doi.org/10.1016/j.cell.2017.05.023
Audette, N.J., Zhou, W., Schneider, D.M., 2021. Temporally precise movement-based predictions in
the mouse auditory cortex. https://doi.org/10.1101/2021.12.13.472457
Ayaz, A., Stäuble, A., Hamada, M., Wulf, M.A., Saleem, A.B., Helmchen, F., 2019. Layer-specific
integration of locomotion and sensory information in mouse barrel cortex. Nature
Communications 10, 1–14. https://doi.org/10.1038/s41467-019-10564-8
Bastos, A.M., Usrey, W.M., Adams, R.A., Mangun, G.R., Fries, P., Friston, K.J., 2012. Canonical
microcircuits for predictive coding. Neuron 76, 695–711.
https://doi.org/10.1016/j.neuron.2012.10.038
Catafau, A.M., Parellada, E., Lomeña, F.J., Bernardo, M., Pavía, J., Ros, D., Setoain, J., Gonzalez-
Monclús, E., 1994. Prefrontal and temporal blood flow in schizophrenia: resting and
activation technetium-99m-HMPAO SPECT patterns in young neuroleptic-naive patients with
acute disease. J Nucl Med 35, 935–941.
Chan, K.Y., Jang, M.J., Yoo, B.B., Greenbaum, A., Ravi, N., Wu, W.-L., Sánchez-Guardado, L., Lois, C.,
Mazmanian, S.K., Deverman, B.E., Gradinaru, V., 2017. Engineered AAVs for efficient
noninvasive gene delivery to the central and peripheral nervous systems. Nat Neurosci 20,
1172–1179. https://doi.org/10.1038/nn.4593
Chen, X., Sun, Y.-C., Zhan, H., Kebschull, J.M., Fischer, S., Matho, K., Huang, Z.J., Gillis, J., Zador, A.M.,
2019. High-Throughput Mapping of Long-Range Neuronal Projection Using In Situ
Sequencing. Cell 179, 772-786.e19. https://doi.org/10.1016/j.cell.2019.09.023
Constantinople, C.M., Bruno, R.M., 2013. Deep cortical layers are activated directly by thalamus.
Science (New York, N.Y.) 340, 1591–1594. https://doi.org/10.1126/science.1236425
Daigle, T.L., Madisen, L., Hage, T.A., Valley, M.T., Knoblich, U., Larsen, R.S., Takeno, M.M., Huang, L.,
Gu, H., Larsen, R., Mills, M., Bosma-Moody, A., Siverts, L.A., Walker, M., Graybuck, L.T., Yao,
Z., Fong, O., Nguyen, T.N., Garren, E., Lenz, G.H., Chavarha, M., Pendergraft, J., Harrington,
J., Hirokawa, K.E., Harris, J.A., Nicovich, P.R., McGraw, M.J., Ollerenshaw, D.R., Smith, K.A.,
Baker, C.A., Ting, J.T., Sunkin, S.M., Lecoq, J., Lin, M.Z., Boyden, E.S., Murphy, G.J., da Costa,
N.M., Waters, J., Li, L., Tasic, B., Zeng, H., 2018. A Suite of Transgenic Driver and Reporter
Mouse Lines with Enhanced Brain-Cell-Type Targeting and Functionality. Cell 174, 465-
480.e22. https://doi.org/10.1016/j.cell.2018.06.035

Dierks, T., Linden, D.E., Jandl, M., Formisano, E., Goebel, R., Lanfermann, H., Singer, W., 1999. Activation of Heschl's gyrus during auditory hallucinations. Neuron 22, 615–621. https://doi.org/10.1016/s0896-6273(00)80715-1

Dombeck, Daniel A., Khabbaz, A.N., Collman, F., Adelman, T.L., Tank, D.W., 2007. Imaging large-scale neural activity with cellular resolution in awake, mobile mice. Neuron 56, 43–57. https://doi.org/10.1016/j.neuron.2007.08.003

Dombeck, Daniel A, Khabbaz, A.N., Collman, F., Adelman, T.L., Tank, D.W., 2007. Imaging large-scale neural activity with cellular resolution in awake, mobile mice. Neuron 56, 43–57. https://doi.org/10.1016/j.neuron.2007.08.003

Douglas, R.J., Martin, K.A., 1991. A functional microcircuit for cat visual cortex. The Journal of Physiology 440, 735–769. https://doi.org/10.1113/jphysiol.1991.sp018733

Du, J., Palaniyappan, L., Liu, Z., Cheng, W., Gong, W., Zhu, M., Wang, J., Zhang, J., Feng, J., 2021. The genetic determinants of language network dysconnectivity in drug-naïve early stage schizophrenia. NPJ Schizophr 7, 18. https://doi.org/10.1038/s41537-021-00141-8

Eliades, S.J., Wang, X., 2008. Neural substrates of vocalization feedback monitoring in primate auditory cortex. Nature 453, 1102–6. https://doi.org/10.1038/nature06910

Fletcher, P.C., Frith, C.D., 2009. Perceiving is believing: a Bayesian approach to explaining the positive symptoms of schizophrenia. Nature reviews. Neuroscience 10, 48–58. https://doi.org/10.1038/nrn2536

Franco, S.J., Gil-Sanz, C., Martinez-Garay, I., Espinosa, A., Harkins-Perry, S.R., Ramos, C., Müller, U., 2012. Fate-Restricted Neural Progenitors in the Mammalian Cerebral Cortex. Science. https://doi.org/10.1126/science.1223616

Frith, C., 2005. The neural basis of hallucinations and delusions. Comptes rendus biologies 328, 169–75.

Garner, A.R., Keller, G.B., 2022. A cortical circuit for audio-visual predictions. Nat Neurosci 25, 98–105. https://doi.org/10.1038/s41593-021-00974-7

Gerfen, C.R., Paletzki, R., Heintz, N., 2013. GENSAT BAC Cre-Recombinase Driver Lines to Study the Functional Organization of Cerebral Cortical and Basal Ganglia Circuits. Neuron 80, 1368–1383. https://doi.org/10.1016/j.neuron.2013.10.016

Gong, S., Doughty, M., Harbaugh, C.R., Cummins, A., Hatten, M.E., Heintz, N., Gerfen, C.R., 2007. Targeting Cre Recombinase to Specific Neuron Populations with Bacterial Artificial Chromosome Constructs. J. Neurosci. 27, 9817–9823.

Gorski, J.A., Talley, T., Qiu, M., Puelles, L., Rubenstein, J.L.R., Jones, K.R., 2002. Cortical excitatory neurons and glia, but not GABAergic neurons, are produced in the Emx1-expressing lineage. J Neurosci 22, 6309–6314. https://doi.org/20026564

Guo, Z.V., Li, N., Huber, D., Ophir, E., Gutnisky, D., Ting, J.T., Feng, G., Svoboda, K., 2014. Flow of cortical activity underlying a tactile decision in mice. Neuron 81, 179–194. https://doi.org/10.1016/j.neuron.2013.10.020

Haarsma, J., Kok, P., Browning, M., 2020. The promise of layer-specific neuroimaging for testing predictive coding theories of psychosis. Schizophrenia Research. https://doi.org/10.1016/j.schres.2020.10.009

Harris, J.A., Mihalas, S., Hirokawa, K.E., Whitesell, J.D., Choi, H., Bernard, A., Bohn, P., Caldejon, S., Casal, L., Cho, A., Feiner, A., Feng, D., Gaudreault, N., Gerfen, C.R., Graddis, N., Groblewski, P.A., Henry, A.M., Ho, A., Howard, R., Knox, J.E., Kuan, L., Kuang, X., Lecoq, J., Lesnar, P., Li, Y., Luviano, J., McConoughey, S., Mortrud, M.T., Naeemi, M., Ng, L., Oh, S.W., Ouellette, B., Shen, E., Sorensen, S.A., Wakeman, W., Wang, Q., Wang, Y., Williford, A., Phillips, J.W., Jones, A.R., Koch, C., Zeng, H., 2019. Hierarchical organization of cortical and thalamic connectivity. Nature 575, 195–202. https://doi.org/10.1038/s41586-019-1716-z

Harrison, P.J., Weinberger, D.R., 2005. Schizophrenia genes, gene expression, and neuropathology:
on the matter of their convergence. Mol Psychiatry 10, 40–68.
https://doi.org/10.1038/sj.mp.4001558

Heindorf, M., Arber, S., Keller, G.B., 2018. Mouse Motor Cortex Coordinates the Behavioral Response
to Unpredicted Sensory Feedback. Neuron 99, 1040-1054.e5.
https://doi.org/10.1016/j.neuron.2018.07.046

Hippenmeyer, S., Vrieseling, E., Sigrist, M., Portmann, T., Laengle, C., Ladle, D.R., Arber, S., 2005. A
Developmental Switch in the Response of DRG Neurons to ETS Transcription Factor
Signaling. PLOS Biology 3, e159. https://doi.org/10.1371/journal.pbio.0030159

Javitt, D.C., 2009. Sensory Processing in Schizophrenia: Neither Simple nor Intact. Schizophr Bull 35,
1059–1064. https://doi.org/10.1093/schbul/sbp110

Jiang, L.P., Rao, R.P.N., 2021. Predictive Coding Theories of Cortical Function. arXiv:2112.10048 [q-
bio].

Jordan, M.I., Rumelhart, D.E., 1992. Forward Models: Supervised Learning with a Distal Teacher.
Cognitive Science 16, 307–354. https://doi.org/10.1207/s15516709cog1603_1

Jordan, R., Keller, G.B., 2020. Opposing Influence of Top-down and Bottom-up Input on Excitatory
Layer 2/3 Neurons in Mouse Primary Visual Cortex. Neuron 108, 1194-1206.e5.
https://doi.org/10.1016/j.neuron.2020.09.024

Keller, G.B., Bonhoeffer, T., Hübener, M., 2012. Sensorimotor mismatch signals in primary visual
cortex of the behaving mouse. Neuron 74, 809–15.
https://doi.org/10.1016/j.neuron.2012.03.040

Keller, G.B., Hahnloser, R.H.R., 2009. Neural processing of auditory feedback during vocal practice in
a songbird. Nature 457, 187–90. https://doi.org/10.1038/nature07467

Keller, G.B., Mrsic-Flogel, T.D., 2018. Predictive Processing: A Canonical Cortical Computation.
Neuron 100, 424–435. https://doi.org/10.1016/j.neuron.2018.10.003

Kim, T.H., Zhang, Y., Lecoq, J., Jung, J.C., Li, J., Zeng, H., Niell, C.M., Schnitzer, M.J., 2016. Long-Term
Optical Access to an Estimated One Million Neurons in the Live Mouse Cortex. Cell Rep 17,
3385–3394. https://doi.org/10.1016/j.celrep.2016.12.004

Kiritani, T., Wickersham, I.R., Seung, H.S., Shepherd, G.M.G., 2012. Hierarchical connectivity and
connection-specific dynamics in the corticospinal-corticostriatal microcircuit in mouse motor
cortex. J Neurosci 32, 4992–5001. https://doi.org/10.1523/JNEUROSCI.4759-11.2012

Le Bihan, D., Turner, R., Zeffiro, T.A., Cuénod, C.A., Jezzard, P., Bonnerot, V., 1993. Activation of
human primary visual cortex during visual recall: a magnetic resonance imaging study.
Proceedings of the National Academy of Sciences of the United States of America 90, 11802–
5.

Leinweber, M., Ward, D.R., Sobczak, J.M., Attinger, A., Keller, G.B., 2017. A Sensorimotor Circuit in
Mouse Cortex for Visual Flow Predictions. Neuron 95, 1420-1432.e5.
https://doi.org/10.1016/j.neuron.2017.08.036

Leinweber, M., Zmarz, P., Buchmann, P., Argast, P., Hübener, M., Bonhoeffer, T., Keller, G.B., 2014.
Two-photon calcium imaging in mice navigating a virtual reality environment. Journal of
visualized experiments : JoVE e50885.

Li, T., Wang, Q., Zhang, J., Rolls, E.T., Yang, W., Palaniyappan, L., Zhang, L., Cheng, W., Yao, Y., Liu,
Zhaowen, Gong, X., Luo, Q., Tang, Y., Crow, T.J., Broome, M.R., Xu, K., Li, C., Wang, J., Liu,
Zhening, Lu, G., Wang, F., Feng, J., 2017. Brain-Wide Analysis of Functional Connectivity in
First-Episode and Chronic Stages of Schizophrenia. Schizophr Bull 43, 436–448.
https://doi.org/10.1093/schbul/sbw099

Ma, Y., Shaik, M.A., Kim, S.H., Kozberg, M.G., Thibodeaux, D.N., Zhao, H.T., Yu, H., Hillman, E.M.C.,
2016. Wide-field optical mapping of neural activity and brain haemodynamics:
considerations and novel approaches. Philosophical Transactions of the Royal Society B:
Biological Sciences 371, 20150360. https://doi.org/10.1098/rstb.2015.0360

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D.,
Hawrylycz, M.J., Jones, A.R., Lein, E.S., Zeng, H., 2010. A robust and high-throughput Cre
reporting and characterization system for the whole mouse brain. Nature Neuroscience 13,
133–140. https://doi.org/10.1038/nn.2467
Maia, T.V., Frank, M.J., 2017. An Integrative Perspective on the Role of Dopamine in Schizophrenia.
Biol Psychiatry 81, 52–66. https://doi.org/10.1016/j.biopsych.2016.05.021
Moghaddam, B., Javitt, D., 2012. From Revolution to Evolution: The Glutamate Hypothesis of
Schizophrenia and its Implication for Treatment. Neuropsychopharmacol 37, 4–15.
https://doi.org/10.1038/npp.2011.181
Morita, K., Im, S., Kawaguchi, Y., 2019. Differential Striatal Axonal Arborizations of the
Intratelencephalic and Pyramidal-Tract Neurons: Analysis of the Data in the MouseLight
Database. Frontiers in Neural Circuits 13.
Musall, S., Kaufman, M.T., Juavinett, A.L., Gluf, S., Churchland, A.K., 2019. Single-trial neural
dynamics are dominated by richly varied movements. Nat Neurosci 22, 1677–1686.
https://doi.org/10.1038/s41593-019-0502-4
Northoff, G., Duncan, N.W., 2016. How do abnormalities in the brain's spontaneous activity translate
into symptoms in schizophrenia? From an overview of resting state activity findings to a
proposed spatiotemporal psychopathology. Prog Neurobiol 145–146, 26–45.
https://doi.org/10.1016/j.pneurobio.2016.08.003
Oberlaender, M., Ramirez, A., Bruno, R.M., 2012. Sensory Experience Restructures Thalamocortical
Axons during Adulthood. Neuron 74, 648–655.
https://doi.org/10.1016/j.neuron.2012.03.022
Pakan, J.M., Lowe, S.C., Dylda, E., Keemink, S.W., Currie, S.P., Coutts, C.A., Rochefort, N.L., 2016.
Behavioral-state modulation of inhibition is context-dependent and cell type specific in
mouse visual cortex. eLife 5. https://doi.org/10.7554/eLife.14985
Parellada, M., Gomez-Vallejo, S., Burdeus, M., Arango, C., 2017. Developmental Differences Between
Schizophrenia and Bipolar Disorder. Schizophr Bull 43, 1176–1189.
https://doi.org/10.1093/schbul/sbx126
Rao, R.P.N., Ballard, D.H., 1999. Predictive coding in the visual cortex: a functional interpretation of
some extra-classical receptive-field effects. Nature Neuroscience 2, 79–87.
https://doi.org/10.1038/4580
Saleem, A.B., Ayaz, A., Jeffery, K.J., Harris, K.D., Carandini, M., 2013. Integration of visual motion and
locomotion in mouse visual cortex. Nature neuroscience 16, 1864–9.
https://doi.org/10.1038/nn.3567
Saravanan, V., Berman, G.J., Sober, S.J., 2020. Application of the hierarchical bootstrap to multi-level
data in neuroscience. Neuron Behav Data Anal Theory 3,
https://nbdt.scholasticahq.com/article/13927-application-of-the-hierarchical-bootstrap-to-
multi-level-data-in-neuroscience.
Schmack, K., Bosc, M., Ott, T., Sturgill, J.F., Kepecs, A., 2021. Striatal dopamine mediates
hallucination-like perception in mice. Science 372, eabf4740.
https://doi.org/10.1126/science.abf4740
Schneider, D.M., Sundararajan, J., Mooney, R., 2018. A cortical filter that learns to suppress the
acoustic consequences of movement. Nature 561, 391–395.
https://doi.org/10.1038/s41586-018-0520-5
Shergill, S.S., Brammer, M.J., Williams, S.C., Murray, R.M., McGuire, P.K., 2000. Mapping auditory
hallucinations in schizophrenia using functional magnetic resonance imaging. Arch Gen
Psychiatry 57, 1033–1038. https://doi.org/10.1001/archpsyc.57.11.1033
Singh, T., Neale, B.M., Daly, M.J., Consortium, on behalf of the S.E.M.-A. (SCHEMA), 2020. Exome
sequencing identifies rare coding variants in 10 genes which confer substantial risk for
schizophrenia. https://doi.org/10.1101/2020.09.18.20192815

Stanley, J., Miall, R.C., 2007. Functional activation in parieto-premotor and visual areas dependent
1189    on congruency between hand movement and visual stimuli during motor-visual priming.
1190    NeuroImage 34, 290–9. https://doi.org/10.1016/j.neuroimage.2006.08.043
1191 Stringer, C., Pachitariu, M., Steinmetz, N., Reddy, C.B., Carandini, M., Harris, K.D., 2019. Spontaneous
1192    behaviors drive multidimensional, brainwide activity. Science 364, 255.
1193    https://doi.org/10.1126/science.aav7893
1194 Suzuki, M., Larkum, M.E., 2020. General Anesthesia Decouples Cortical Pyramidal Neurons. Cell 180,
1195    666-676.e13. https://doi.org/10.1016/j.cell.2020.01.024
1196 Taniguchi, H., He, M., Wu, P., Kim, S., Paik, R., Sugino, K., Kvitsani, D., Fu, Y., Lu, J., Lin, Y., Miyoshi, G.,
1197    Shima, Y., Fishell, G., Nelson, S.B., Huang, Z.J., 2011. A Resource of Cre Driver Lines for
1198    Genetic Targeting of GABAergic Neurons in Cerebral Cortex. Neuron 71, 995–1013.
1199    https://doi.org/10.1016/j.neuron.2011.07.026
1200 Valley, M.T., Moore, M.G., Zhuang, J., Mesa, N., Castelli, D., Sullivan, D., Reimers, M., Waters, J.,
1201    2020. Separation of hemodynamic signals from GCaMP fluorescence measured with wide-
1202    field imaging. J Neurophysiol 123, 356–366. https://doi.org/10.1152/jn.00304.2019
1203 Wekselblatt, J.B., Flister, E.D., Piscopo, D.M., Niell, C.M., 2016. Large-scale imaging of cortical
1204    dynamics during sensory perception and behavior. Journal of Neurophysiology 115, 2852–
1205    2866. https://doi.org/10.1152/jn.01056.2015
1206 Widmer, F.C., Keller, G.B., 2021. Developmental plasticity in visual cortex is necessary for normal
1207    visuomotor integration and visuomotor skill learning. bioRxiv 2021.06.20.449148.
1208    https://doi.org/10.1101/2021.06.20.449148
1209 Xu, X., Olivas, N.D., Ikrar, T., Peng, T., Holmes, T.C., Nie, Q., Shi, Y., 2016. Primary visual cortex shows
1210    laminar-specific and balanced circuit organization of excitatory and inhibitory synaptic
1211    connectivity. The Journal of Physiology 594, 1891–1910. https://doi.org/10.1113/JP271891
1212 Yamawaki, N., Shepherd, G.M.G., 2015. Synaptic circuit organization of motor corticothalamic
1213    neurons. J Neurosci 35, 2293–2307. https://doi.org/10.1523/JNEUROSCI.4023-14.2015
1214 Yoon, J.H., Rokem, A.S., Silver, M.A., Minzenberg, M.J., Ursu, S., Ragland, J.D., Carter, C.S., 2009.
1215    Diminished orientation-specific surround suppression of visual processing in schizophrenia.
1216    Schizophr Bull 35, 1078–1084. https://doi.org/10.1093/schbul/sbp064
1217 Zmarz, P., Keller, G.B., 2016. Mismatch Receptive Fields in Mouse Visual Cortex. Neuron 92, 766–
1218    772. https://doi.org/10.1016/j.neuron.2016.09.057
1219