1 Constraint-based modeling identifies metabolic vulnerabilities during the epithelial to
2 mesenchymal transition

3   Scott Campit, Venkateshwar G. Keshamouni*, Sriram Chandrasekaran*
4
5   *- Corresponding authors
6

[1]Program in Chemical Biology, University of Michigan, Ann Arbor, MI, 48109, USA.

[2]Center for Bioinformatics and Computational Medicine, Ann Arbor, MI, 48109, USA.

[3]Department of Biomedical Engineering, University of Michigan, Ann Arbor, MI, 48109, USA.

[4]Rogel Cancer Center, University of Michigan Medical School, Ann Arbor, MI, 48109, USA.

[5]Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, University
of Michigan, Ann Arbor, MI 48109

Abstract

Epithelial-to-mesenchymal transition (EMT) is a developmentally conserved cellular process critical for tumor metastasis. EMT enables malignant epithelial cells to acquire mesenchymal-like migratory and invasive phenotype. During EMT cancer cells undergo extensive metabolic reprogramming that correlates with the suppression of proliferation, and stimulation of the energy-intensive migratory behavior. However, the causal relationship between metabolic changes and coordinated physiological phenotypes that occur during EMT is still unclear. We used bulk time-course transcriptomics and proteomics, and single-cell transcriptomics from five independent EMT studies in A549 lung adenocarcinoma cells to simulate metabolic network activity using constraint-based modeling. Model predictions were validated using literature mining, experimental studies and CRISPR-Cas9 essentiality screens. We uncovered temporal metabolic dependencies in glycolysis and glutamine metabolism reactions, and experimentally validated isoform-specific dependency on Enolase3 for cell survival during EMT. Together, our approach uncovered temporally regulated cell-state-specific metabolic dependencies in cells undergoing EMT.

Introduction

EMT is a reversible developmental process stimulated by extracellular signals that facilitate the transition from an epithelial (E) cell to a motile and invasive mesenchymal-like (M) cells, enabling circulating tumor cells to initiate metastasis (Pastushenko and Blanpain 2019). More importantly, EMT is not a binary process but occurs through a spectrum of distinct intermediate states with potential functional consequences. The cytokine - transforming growth factor β (TGF-β), is a potent inducer of EMT. Consistently, TGF-β levels are highly upregulated and directly correlate with tumor progression, enhanced invasion, metastasis, and poor survival in patients with non-small cell lung cancer (NSCLC) (Padua and Massagué 2009). Cell culture models of TGF-β -induced EMT serves as robust *in vitro* model to investigate mechanisms of metastasis. In addition to metastasis, the process of EMT-MET is implicated in several clinically relevant aspects including, tumor heterogeneity, stemness, and drug resistance (Ramesh et al. 2020). Understanding underlying regulatory mechanisms is essential to develop therapeutic strategies that can prevent EMT or promote MET to inhibit metastasis.

Extensive molecular and structural changes that occur during EMT can potentially induce robust metabolic reprogramming to support changing cellular phenotypes. Cancers show enhanced glycolytic pathway activity even in the presence of oxygen, via the Warburg effect (Vander Heiden et al. 2009). It is now clear that the Warburg effect is not restricted to cancer cells but it is an adaptive physiological program that occurs in many normal cell types, when cells need rapid ATP production, biomass synthesis and balancing of reactive oxygen species (Pålsson-McDermott and O'Neill 2020)(Kim et al. 2016), Studies have demonstrated an enhanced glycolytic activity during EMT. However, similar to EMT, metabolic reprogramming may also involve a spectrum of pathway combinations at a given cellular steady state. While several metabolic rewiring strategies have been observed in diverse cancers, a comprehensive systems-level characterization of metabolic reprogramming during the EMT has not been

carried out. Methods that can infer metabolic dysregulation using omics data will be invaluable

for understanding causal relationship between EMT and metabolic reprogramming.

Constraint-Based Optimization   and Reconstruction Analysis (COBRA)   is   a widely

used approach for simulating genome-scale metabolic fluxes using omics data. COBRA

simulates metabolic fluxes by using the metabolic network architecture, nutrient availability, and

omics data as constraints in an optimization problem representing a cellular objective, such as

maximizing biomass production (Orth et al. 2010). COBRA models have inferred metabolic

rewiring strategies in several cancer subtypes (Oruganty et al. 2020; Yizhak et al. 2014; Nilsson

et al. 2020). For instance, incorporating metabolomics data to identify synthetically lethal

metabolic genes in pancreatic cancer (Nelson et al. 2020). However, to our knowledge, no one

has applied COBRA to study metabolic network heterogeneity during the EMT and

characterized the metabolic properties of intermediate states during EMT.

We used COBRA to simulate metabolic activity and vulnerabilities during EMT using

diverse omics sources, including time-course transcriptomics, proteomics, and single-cell

transcriptomics data. Notably, this study applies constraint-based modeling to single-cell cancer

transcriptomics data to capture the metabolic heterogeneity during EMT. From our analysis, we

were able to identify known metabolic dependencies during EMT, such as uptake of glucose

and glutamine. We also predicted new metabolic dependencies including the enolase and

GOT1 reactions and those related to alpha-ketoglutarate metabolism. Surprisingly, many of

these dependencies were time-specific, suggesting that there is a narrow temporal window

during which the cells can be targeted with drugs that inhibit these pathways. We also found

metabolic changes that showed consistent trends based on model predictions derived from both

the bulk and single-cell studies. Together, our analysis provides a framework to integrate

multiple omics datasets to examine tumor metabolic heterogeneity and infer new drug targets.

Methods

Differential Expression (Bulk Studies)

We analyzed two transcriptomics (Hecker et al. 2009; Keshamouni et al. 2009) and two

proteomics (Keshamouni et al. 2006; Lu et al. 2019) EMT time-course studies with A549 as the

cell model undergoing TGF-β induction. All studies compared later time point after TGF-β

induction over day 0 to obtain differentially expressed genes and proteins. When possible,

authors' methods and provided datasets were used to obtain a list of up- and downregulated

gene sets. If no preprocessed data was provided (as in the case of GSE17518), *limma-voom*

(Law et al. 2014) was performed to determine differentially expressed genes between

conditions. Additionally, a GAM-LOESS model was used to determine differentially expressed

genes in GSE147405 (Cook and Vanderhyden 2020), aggregating single-cells at the time-

course level. The regression coefficients from the GAM-LOESS model were used to determine

the sign of regulation (up/down). P-values from *limma-voom* and a GAM-LOESS model were

adjusted using the Benjamini-Hochberg method and the significance threshold used was P-

value < 0.05. The expression matrix containing statistically significant normalized scores for all

metabolic genes across all 5 experiments can be found in **Supplementary Table 1**.

Individual Cell Differential Expression

We computed differentially expressed genes for individual cells without TGF-β removal

in GSE147405 to simulate individual cell fluxes and reaction knockout growth rates. Data

preprocessing included data scaling, removing contaminant artifacts such as mitochondrial

genes, and removing cells with low total gene counts. This was performed on the raw data

object. Further, we used the data imputation algorithm MAGIC (Van Dijk, D. et al., 2018) to fill in

drop out values. The MAGIC-imputed data was transformed to a Z-score using a Z-score

method that subtracts out the median and centers the data based on the median absolute

deviation (MAD). The formula for the robust Z-score for a specific gene *i* in a given cell *j* is

shown in **Equation 1**:

$$Z_{i,j} = \frac{x_{i,j} - median(x_i)}{1.486 \times MAD} \ [1]$$

107 Where 1.486 is a scaling constant. For any given cell, a gene was determined to be

108 upregulated/downregulated if the robust Z-score was positive/negative and the P-value < 0.05.

110 Prioritizing metabolic gene targets across multiple studies

111 We evaluated the robustness metabolic gene dysregulation across five EMT studies

112 using the following prioritization score η. The method to compute η shown in **Equation 2**:

113
114
$$\eta_i = M \sum_{i=1}^{N} abs(E_i), \ [2]$$
115
Where M is the number of studies where the gene was determined to be significant and E_i is the

gene effect size (log_2 fold change or Z-score) for gene i. The prioritization scores were ranked in

descending order and used to prioritize reactions for further investigation.

COnstraint-Based Reconstruction and Analysis (COBRA)

Flux balance analysis (FBA; Orth et al. 2010) was used to simulate metabolic activity

using the human metabolic reconstruction RECON1 (Duarte et al. 2007). Cells were assumed to

maximize biomass production as the objective function. Differentially expressed metabolic

genes that intersected with RECON1 were used as biological constraints to maximize

(upregulation) or minimize (downregulation) metabolic flux using a modified form of the iMAT

algorithm (Zur et al. 2010; Shen et al 2019). Parsimonious enzyme usage (Lewis et al. 2010)

was an additional assumption to obtain a unique metabolic flux distribution and to minimize

fluxes that did not contribute to biomass formation. Metabolic fluxes and growth rates from

single gene and reaction knockout simulations were obtained using COBRA. To ensure a

feasible growth rate was calculated, we removed genes/reactions that were upregulated in the

knockout and set the percent knockout to be 99% to promote a feasible flux solution.

Differential Metabolic Activity and Knockout Sensitivity Analysis

To determine differentially active metabolic reactions, we used the priority score

described in equation 1 on the absolute value of the metabolic fluxes. Most reactions show zero

flux, and so reactions that showed metabolic activity were considered to be overactive metabolic

reactions.

To determine the impact metabolic genes have on the growth rate during different

stages of EMT, we computed a sensitivity score θ comparing EMT versus control growth rates

for each gene knockout. The equation to compute the bulk sensitivity score is shown in

Equation 3:

$$\theta_r = \text{mean}(\frac{\text{cancer}_{i,j} \ / \ \text{mean}(\text{cancer}_j)}{\text{control}_{i,j} \ / \ \text{mean}(\text{control}_j)}) - 1 \ [3]$$

The flux or growth rate was mean-normalized for the control (day 0) and the TGF-β

treatment (all other days). Then, the final score was taken as the ratio of the TGF-β treated

growth rate over the average control growth rate for a given reaction knockout. The average

ratio across all cells was taken as the score to identify differentially sensitive metabolic

reactions. The score was centered at 0. The intuition behind θ is as follows: if the score is 0, the

gene/reaction knockout has no difference between cancer and control. If the score is less than

0, the knockout impacts the cancer cell more than control, and is considered to be essential for

cellular growth. To rank and prioritize metabolic targets for experimental validation, we used the

same prioritization score as we did to rank differentially expressed genes.

Classifying cancer cell line states

Cancer cell lines from the Cancer Cell Line Encyclopedia (CCLE) were annotated by

their source from a primary tumor or metastatic tumor. To also classify whether a cell was in the

epithelial versus mesenchymal state, we calculated the Z-score and p-value for all genes in the

CCLE and mapped them to known EMT markers. Upregulated genes were classified as having

a positive Z-score and a significance threshold of p-value < 0.05, while downregulated genes

were classified as having a negative Z-score with the same significance threshold.

EMT markers (with up- and downregulated signatures) were taken from MSigDB

(Liberzon et al. 2015) across three studies from different tissues of origin induced using TGF-β.

We further filtered this list with NSCLC markers without up/downregulation annotations from

EMTome (Vasaikar et al. 2021). The final number of markers obtained for EMT was 14 genes,

which were used to classify cancer cell lines (CCLs). We chose to classify cell lines using upper

and lower quantiles of gene makers. Up- and downregulated genes from NSCLC CCLs were

cross referenced to the list of EMT signatures and labeled as E if the number of signatures was

less than 5 genes or M if the number of signatures was greater than 8.

CRISPR-Cas9 Analysis

We analyzed batch corrected CERES Scores (Pacini et al. 2021) for metabolic genes

that were predicted by COBRA to have increased metabolic activity or resulted in a reduction of

growth rate from knockout. CERES Scores were separated based on their association with

metastatic (Met) and primary (Prim) cell lines, which was determined based on the CCLE

metadata (Barretina et al. 2012). We also compared the CCLE annotations against our own

Epithelial (Epi) and Mesenchymal (Mes) annotations, methods described above.

To evaluate how well our model predictions related to CERES Scores, we calculated the

Pearson correlation coefficient between our predicted growth scores and ratios of Mes / Epi or

Met / Prim CERES Scores. Further, we compared the growth scores against different subsets of

the CERES Score data, including NSCLC only cell lines and all cell lines.

Identifying metabolic enzymes and EMT studies for systematic literature
validation

To be considered for our systematic literature validation, we pooled a list of metabolic

enzymes predicted from COBRA from bulk and single-cell reaction knockouts that had lethal

reactions in at least 2 studies (growth score < 0). The query was performed using PubMed and

Google with the following keywords using AND filtering: "EMT", "Metabolism", "A549",

"metastasis", "cancer", "cancer metabolism", and the individual gene of interest. The list of the

manually curated results can be found in **Supplementary Table 5**.

We expanded the scope of our literature search to encompass all cancer cell lines. The

query was manually curated to either support or refute the COBRA predictions. None of the

model predictions contradicted the literature. Reaction predictions and their confidence were

scored (1-3), where 1 has no evidence based on literature and 3 has strong A549 or lung

adenocarcinoma specific evidence. The rules to assign each score for each reaction prediction

are shown below:

● **1:** Prediction has no literature support.

● **2:** Prediction has literature evidence with general cancer lineages.

● **3:** Prediction has literature evidence either with specific experiments from A549 or

related lung adenocarcinoma tissue/cell lines.

Cell culture, siRNA transfection and EMT induction

A549 human lung adenocarcinoma cell line was obtained from the American Type

Culture Collection (Manassas, VA) and maintained in RPMI-1640 medium with glutamine

supplemented with 10% FBS, penicillin, and streptomycin at 37°C in 5% CO2. For inducing

EMT cells at 40-50% confluency in complete medium were serum starved for 24 hrs and treated

with TGF-β (5 ng/ml) for 72 hrs.

Isoform specific siRNA for enolases includes a pool of 4 SMART selection-

designed synthetic duplexes (Dharmocon's SMARTpool). A scrambled sequence from the

same company is used as a control. Cells at 40-50% confluency were transfected with siRNA

using Lipofectamine 2000 (Cat No: 18324-012, Invitrogen) and optiMEM medium (Cat No:

31985, Gibco) following the manufacturer's instructions. After 6 hours of transfection cells were

washed and allowed to recover from transfection in RPMI 1640 medium with 10% FBS before

inducing EMT as described above.

Apoptosis assays

Apoptosis was assessed by two independent methods; *1) AnnexinV/7-AAD staining (Kit*

from Biolegend Cat# 640922): At the end of the EMT experiment described above, all cells

(including floating cells) were collected, washed and resuspended in Annexin V binding buffer.

100 ul of cell suspension was stained with 5 ul of FITC-Annexin V, followed by 5 ul of 7-AAD

staining solution. After 30 min incubation at room temperature in dark, 400 ul of Annexin V

binding buffer is added and assessed for Annexin V and 7-AAD staining by flow cytometry. Both

Annexin V postive (early apoptotic) and Annexin V and 7-AAD double positive cells (late

apoptotic) are added together for assessing total apoptosis. *2) Assessing Caspase 3 activation:*

To assess casapase3 activation during EMT, an artificial caspase3 substrate coupled to a green

fluorescent DNA-binding dye (DEVD-Nucview) is added to the cell culture. When caspase3 is

activated, it cleaves the DNA-binding dye which enters the nuclei and labels an apoptotic cell

with green fluorescence allowing its imaging. Green fluorescent apoptotic cells were imaged

under a fluorescent microscope 48 hrs after TGF-β-induced EMT.

Results

COBRA reveals that cells undergoing EMT exhibit enhanced glycolysis during early and late stages

We performed a meta-analysis of differentially expressed genes and proteins across four bulk EMT datasets. Two were RNASeq-based datasets (GSE17708 and GSE17518), and two were proteomics-based datasets (Garcia and Keshamouni). To aggregate the results from multiple studies, we designed a prioritization score to rank the reactions based on effect size and whether or not the gene was significantly expressed in a given study (**Supplementary Table 1; Methods**).

We simulated the metabolic fluxes for each time-point using the transcriptomics and proteomics data to see how metabolic activity changes over time during EMT using Flux Balance Analysis (FBA; **see Methods**). FBA uses a linear optimization procedure with biological constraints, such as knowledge of the metabolic network structure (known as a stoichiometric (S) matrix) and expression levels as inputs to generate cell-state specific metabolic flux profiles (O'Brien et al. 2015). FBA assumes that the cell is maximizing an objective, usually its biomass production. While standard FBA outputs multiple flux profiles due to the rank deficiency of the S matrix (Orth et al. 2010), Parsimonious FBA or pFBA provides a unique flux distribution by assuming optimal enzyme efficiency by minimizing the overall metabolic flux throughout the metabolic network while maximizing biomass production (Lewis et al. 2010). pFBA identifies the smallest set of active reactions that best support biomass production.

Our predictions using pFBA reveal that there are more active reactions during the early and late phases of EMT. During the intermediate phases of EMT, metabolic activity goes down. As cells undergo dramatic structural rearrangements when transitioning to a mesenchymal cell, cells require energetic substrates such as ATP to facilitate these processes. Our metabolic model assumed that these cancer cells were optimizing for increased biomass production, and the reduction of fluxes for biomass production during the intermediate EMT stages suggests that

metabolic activity is being siphoned towards other processes such as motility. Our metabolic

flux profile data suggests that cancer cells upregulate metabolism initially to build up metabolic

substrate levels, and then divert all transcriptional resources towards other processes.

Samples within these time-points tend to have similar metabolic functions, as most

active reactions are found within central carbon metabolism (glycolysis/gluconeogenesis,

pentose phosphate pathway, folate metabolism) and nutrient exchange subsystems. These

metabolic pathways contribute to biomass formation. We visualized the top 50 reactions sorted

by prioritization scores (**Methods; S. Figure 1; Supplementary Table 2**). The prioritization

score takes into consideration the number of studies where a given metabolic gene(s) encoding

a reaction was determined to be significant and absolute value of the gene effect size ($\log_2$ fold

change or Z-score). Developing a prioritization score enabled us to filter through 3744 reactions

to provide a concise reaction list for downstream analyses.

267
Several glycolytic reactions were predicted to have increased metabolic activity and

priority scores (**Figure 1A**), which was expected given how cancer cells rewire glycolytic

activity, as evidenced by the Warburg effect (Vander Heiden et al. 2009). Several glycolytic

substrates play a role in both cellular survival and cancer proliferation. It has been well

established that TGF-β increases expression of several glycolytic enzymes (Jia et al. 2021). We

found that hexokinase, glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and enolase

were the top 3 glycolytic reactions that were highly active in both early and late EMT stages,

supporting previous studies that suggest glycolysis is directly impacted by TGF-β induction. The

timing of metabolic activity suggests that glycolysis is essential for initiating EMT and

establishing metastasis at later stages.

Genome-scale reaction knockout simulation identifies extensive
vulnerabilities in mesenchymal state

While our previous analysis focused on reaction fluxes, next we used FBA to simulate

the impact of reaction knockout on cellular growth in each time-points across five independent

A549 TGF-β induced EMT studies (**S. Figure 2**; **Supplementary Table 3**). Briefly, each

metabolic reaction encoded in the reconstruction was systematically shut off (upper and lower

bounds were set to 0) to simulate a "knockout", while the growth rate objective was optimized.

This method allows us to infer the impact systematic reaction knockouts have on cellular

growth. We analyzed the distribution of knockouts across bulk experiments and by time-course.

The later stage of EMT were predicted to have more vulnerabilities (932 reactions) than in the

early stage (874 reactions) and intermediate stage (660 reactions), suggesting mesenchymal

cells are more vulnerable to metabolic perturbation (**Figure 2; inset**).

We identified over 40 reactions that were specifically sensitive in specific EMT stages

and studies (**Figure 2**). These also highlight the technical and biological variance that is

observed in EMT studies across different omics modes. Notably, alpha-ketoglutarate (AKG)

transport between the cytosol was unique to the intermediate stage in two out of five

independent studies. From a mechanistic standpoint, AKG likely suppresses metastasis by

counteracting the effects of other oncometabolites such as 2-hydroxyglutarate, succinate, and

fumarate (Wei et al. 2020). While the impact of AKG and cellular differentiation / proliferation

has been observed through several nutrient perturbation studies in cancer and stem-cells, the

exact source and subcellular contribution of AKG and its impact on metastasis is difficult to

determine experimentally. Our computational model suggests that knockout of AKG transport

between the cytosol and mitochondria has a negative impact during EMT, providing clues about

cellular compartment dynamics and their impact on cancer metastasis. We hypothesize that

accumulation of AKG within the mitochondria counteracts oncometabolite effects through

additional regulatory mechanisms.

Two other reactions in central carbon metabolism had high priority scores, namely,

enolase (ENO) and lactate dehydrogenase (LDH_L), both predicted from the same EMT

proteomics data (Keshamouni et al. 2006). Upregulated enolase levels are associated with

promoting cell growth, migration, and invasion during EMT in various cancers (Song et al. 2014;

Zhao et al. 2015). Further, LDH is highly associated with cancer metastasis, and has been

shown to activate EMT in several cancers, including lung cancer, during metastasis (Hou et al.

2019; Zhang et al. 2018). While it is known that ENO and LDH upregulation and/or increased

activity are associated with poor patient prognosis, little is known about how changes in

metabolic activity over time leads to differential sensitivity in cancer. Our modeling approach

shows that enolase and lactate dehydrogenase are essential during later stages of EMT

compared to earlier stages, revealing information about time-dependent sensitivity of these well-

known targets for the first time.

In addition to nutrient exchange reactions, we found three metabolic reactions that

consistently decreased growth upon KO across all time points. Two metabolic enzymes were

involved in fatty acid metabolism: Fatty acid CoA ligase hexadecanoate and beta-ketoacyl

synthetase. Fatty acid synthase (FASN) is a potential therapeutic target for NSCLC, and beta-

ketoacyl synthetase is one component of FASN. Preclinical studies show that beta-ketoacyl

synthase inhibition induces apoptosis and stops proliferation in cancer cells *in vitro* and *in vivo*

(Menendez et al. 2004; Pizer et al. 1996).

Further, our model suggests two additional reactions that do not have literature backing

to be potential therapeutic targets, but are associated with metabolic pathways that are

frequently dysregulated across different cancers. Lipid metabolites concentrations have

prognostic value, and dysregulated fatty acid metabolism is associated with poor cancer patient

prognosis. While it is known that metabolic enzymes such as Fatty Acid CoA Ligases modify

ratios of these fatty acids, and that there is differential regulation and expression of these

metabolic enzymes in cancer, little is known about the balance of fatty acids and fatty acyl-CoAs

and its impact on cancer. Our model suggests that the fatty acid CoA ligase that specifically

modifies hexadecanoate contributes highly to EMT and suggests an interesting hypothesis that

needs to be validated experimentally, with potential for therapeutic intervention.

Additionally, we analyzed reactions that either had very strong effects on a single study

or were predicted to impact biomass in specific time points in at least two out of five studies (**S.**

Figure 2). This provided us information about reactions that show temporal-specificity or

robustness across datasets. Glucose and aspartate exchange reactions were predicted to be

sensitive across all time points and experiments, suggesting that cells in all stages of EMT are

sensitive to perturbations to these nutrients. It is well documented in the literature that high

glucose levels facilitate migration and invasion processes in EMT for several types of cancer

(Xu et al. 2019; Liu et al. 2016). Additionally, aspartate is crucial to cell proliferation and survival

in cancer (Birsoy et al. 2015; Alkan et al. 2018; Sullivan et al. 2015). Our model also captured

aspects of metabolic heterogeneity associated with glutamine metabolism in EMT. We found

that cells were dependent on glutamine exchange in early (1 hr) and late (48-72 hrs) time

points, while becoming insensitive to glutamine exchange during intermediate stages (8 - 24

hrs) (**S. Figure 2; top row**). Glutamine metabolism is essential for sustaining proliferation in

many tumor lineages including NSCLC, and the dysregulation of glutaminolysis is a hallmark of

cancer metabolism (Yang et al. 2017). Glutamine regulates the activation of STAT3, a critical

transcription factor associated with tumor growth and metastasis (Cacace et al. 2017; Yang et

al. 2014). Together, these results suggest that our COBRA models can accurately predict well-

known impact of nutrient perturbations in cancer and EMT.

Isoform-specific role of Enolase 3 in regulating cell survival during EMT

Reactions in glycolysis, especially enolase, was identified by both our flux and gene

knockout analysis to have high metabolic activity and sensitivity to knockout. During tumor

progression, cancer cells must increase glucose metabolism. Owing to the hypoxic tumor

microenvironments, cancer cells upregulate glycolytic enzymes, including Enolase (Eno), to

support anaerobic proliferation (Warburg effect). Enolase (Eno) is a key glycolytic enzyme that

catalyzes the dehydration of 2-phosphoglycerate to phosphoenolpyruvate. It occurs as 3

isoforms, Eno1 (ubiquitously expressed in all cells), Eno2 (neuronal specific) and Eno3 (muscle

specific) (Chang et al., 2006). Our transcriptomic analysis show that Eno3 the muscle specific

isoform which is catalytically more efficient, is 10 fold differentially expressed in cells undergoing

EMT (**S. Figure 3**) (Keshamouni et al., 2006). siRNA mediated inhibition of Eno3 selectively

induced apoptosis in cells undergoing EMT whereas, inhibition of the ubiquitously expressed

isoform, Eno1, did not, as assessed by Annexin-V/PI staining by flow cytometry (**Figure 3A**) and

Caspase8 activation assay (**Figure 3B**). These observations suggest that EMT induces

reprogramming of glycolysis to an Eno3 dependent pathway to meet the energy demands of

migratory and invasive cells. Inhibition of Eno3 will selectively kill cells undergoing EMT and

may prevent metastasis.

Single-cell knockout simulations reveal metabolic heterogeneity in a cell population undergoing EMT

To determine whether the variations observed in bulk dataset analysis are true reflection

of metabolic phenotypes at the single cell level, we next analyzed single cell transcriptomics

data of A549 cells induced with TGF-**β** (GSE147405; Cook & Vanderhyden, 2020). To capture

subtle metabolic differences as cells' transition from E to M states, we reconstructed separate

models for each cell based on its transcriptomic profile measured in the dataset. To ensure we

were observing the transition between E to M in this dataset, we visualized VIM and CDH1

expression levels in the UMAP embedding and found that the expression profiles are consistent

with what is observed in the literature (**Supplementary Figure 4**). We used the resulting models

for 644 individual cells across all time points and computed growth rates after genome-scale

reaction knockouts in each individual cell. For comparison with the bulk datasets, we

aggregated the cells, taking the average knockout scores across each time point (**Figure 4; S.**

Table 4). We found that many reactions predicted to impact growth in the single-cell analysis

were also sensitive in the bulk analysis (**Table 1, S. Table 5;** N = 95 intersected reactions).

To examine how reaction knockout sensitivity changes over EMT progression for the top 5

variable central carbon metabolism reactions, we plotted the growth scores for representative

reactions that had high variance onto the UMAP embedding (**Figure 5 A-F**). Reactions that

were predicted from our bulk knockout profiles including AKG-malate transport, Enolase,

Carnitine O-acetyltransferase, and ATP-Citrate Lyase show heterogeneous sensitivity across all

time points. Interestingly, each of these vulnerabilities affects a distinct subpopulation of cells.

Citrate Synthase was predicted to be sensitive across all time points, suggesting that this

reaction is critical in all stages of EMT. We also observed metabolic heterogeneity in glutamine

metabolism. We found that there was a positive correlation between the master regulator

STAT3, glutamine synthetase, and glutamine transporter levels in the single cell data (**S. Figure**

5A & B). This is consistent with studies that have observed that glutamine regulates the

activation of STAT3. Overall, our model identifies individual cells that are sensitive to specific

reaction knockouts, providing a granular metabolic dependency profile of a population of cells

undergoing EMT.

397
398　## Prioritization of metabolic targets during EMT using both bulk and single-
399　cell simulations

400
We performed extensive literature curation for genes that were found to show growth

reduction upon knockout in both the bulk and single-cell analysis (**Table 1, S. Table 5;** N = 95

intersected reactions). These reactions were prioritized based on the number of studies found

for each gene query and its relevance to cancer and EMT. Two high confidence predictions that

were found in both analyses included Pyruvate Carboxylase and Fructose-Bisphosphate

Aldolase, which were shown to contribute specifically to NSCLC progression and metastasis

(Sellers et al. 2015; Fu et al. 2020). Eight reaction predictions had some evidence of being

dysregulated in another cancer subtype, but not NSCLC. The remaining 10 reactions have not

been highlighted in the literature, and present opportunities for experimental validation.

To further assess our model predictions against experimental data, we compared our

bulk and single-cell knockout results against batch-corrected CRISPR-Cas9 essentiality

knockout screens integrated from the Broad and Sanger Institute (Pacini et al. 2021). Given the

limited availability of CRISPR-Cas9 screenings in EMT studies, we took NSCLC cancer cell

lines from the DepMap dataset, which annotated them to be derived either from a primary tumor

or a metastatic site. Further, we took EMT signatures from MSigDB and EMTome to classify

cancer cell lines from the cancer cell line encyclopedia (CCLE) into epithelial-like or

mesenchymal-like cell-lines (**Methods**). When comparing the classification of CCLE cancer

annotation with our EMT classification, we found that there was high agreement between cancer

cell lines obtained from a primary site and the epithelial state while there was low agreement

between the mesenchymal cell state and cell lines from a metastatic site **(Figure 6A)**.

The essentiality of metabolic enzymes identified from our model predictions were

interpreted using the CRISPR gene knockout (CERES) Scores, where a lower score is

associated with a higher likelihood that a given gene is essential for survival in a given cell line

(Meyers et al, 2017). A score of 0 was used as the threshold to indicate the median effect of

non-sensitive genes. We overlaid the CERES Scores for metabolic genes corresponding to

reactions predicted from our sensitivity analysis with the CCLE cancer cell line annotation and

our EMT annotation (**Figure 6B and C**). Overall, we found that the median values for both

classification methods agreed with each other for the most part. The alpha-ketoglutarate /

malate transporter SLC25A11, ENO1, ENO2, ENO3, IDH1, and LDH show lower median

CERES scores than the threshold of 0, supporting our model's findings. From this analysis, we

were able to identify isoform-specific sensitivity in NSCLC, analogous to our validation of Eno3

dependency (**Figure 3**). IDH1 gene depletion is associated with NSCLC essentiality, compared

to IDH2 and IDH3 depletion, suggesting that targeting IDH1 expression in NSCLC may be an

effective therapy to supplement existing therapeutics that target specific IDH1/2 mutations.

To evaluate how well our knockout growth score predictions performed against CRISPR-

Cas9 experimental data, we took the ratio of the CERES scores for the Primary site derived

NSCLC cell lines to the scores for the Metastatic site derived cell lines for each metabolic gene.

We found that the average growth scores and metabolic fluxes agreed significantly with the ratio

data with R = 0.31 and 0.20 respectively, P-value = 0.005 and 0.036 (**S.Table 6 & 7**). We

determined the correlation between our predictions against all cell lines as well, but found that

the correlations were not significant with the pan-cancer CERES Score data. These results

match our expectations, as our COBRA models were constrained using A549 transcriptomics

and proteomics data. We evaluated the quality of each dataset on COBRA predictions, and

found that the single-cell RNASeq data best matched the CERES Score ratios (KO R = 0.27,

KO P-value = 0.01; Flux R = 0.20, Flux P-value = 0.03; **S. Table 6 & 7**) while bulk

transcriptomics and proteomics data were weakly correlated (R < 0.1; P-value > 0.05; all bulk

experiments).

There are several confounding variables that could contribute to the reduced correlation

between our predictions and the experimental data. First, we assumed that cells derived from

primary tumor sites have similar metabolic attributes to epithelial-like cells, while metastatic cell

lines were similar to mesenchymal cells. We addressed this assumption by re-classifying cells

based on EMT gene markers. We obtained similar correlations and p-values from grouping cell

lines either using the CCLE annotation or our EMT classification method (**S. Table 8 & 9**).

While the transcriptomics and proteomics EMT datasets to build the metabolic models were

induced using TGF-β, the cancer cell lines used in the CRISPR screen were not induced with

an EMT inducer. Thus, the cell dynamics that occur during EMT are not captured in CRISPR

screening data. Finally, we used NSCLC CCLs to evaluate the statistical significance of our

results, while our simulations were performed on A549 exclusively. Due to these confounding

factors, in contrast to our siRNA knockdown experiment, we found that the CERES scores did

not distinguish between the three isoforms of enolase. Despite these assumptions and

considerations, we found that our simulations were correlated significantly with the CERES

Scores, suggesting that our model is able to extract relevant biological insights.

Discussion

Here we utilize constraint-based modeling informed by multiple omics data sources to

predict metabolic activity and knockout sensitivity during EMT. Our predictions are supported

from literature validation, siRNA knockout studies, and CRISPR-Cas9 essentiality panels. We

further provide a list of high confidence metabolic reaction dependencies during EMT for future

experimental validation. Our approach also provides insights into metabolic activity at the single-

cell level, which is not possible to infer with current experimental methodologies.

Our modeling identified metabolic enzymes that are novel as well as those with

experimental evidence in literature supporting their role in tumor progression. We identified

known metabolic reactions that contribute to cancer progression, such as glucose and

glutamine transport. We further identified metabolic reactions associated with fatty acid

metabolism that contribute to metastasis. We found that most glycolytic reactions were

overactive in the early and late stages of EMT. This time-dependent aspect of glycolytic activity

was intriguing and suggests a potential vulnerability during EMT. We further experimentally

validate the essential role of the enolase reaction in EMT. The enolase enzyme is implicated in

cancer progression for various tissue lineages, but so far has not been identified as a crucial

player in NSCLC metastasis. Our COBRA modeling approach identified reaction catalyzed by

Enolase as highly active during the early and late stages of EMT and predicted enolase

knockout to have a negative impact on cellular growth. Enolase has three isoforms with a

degree of cell type specific expression. Eno1 is ubiquitously expressed in all cells, Eno2 is

neuronal specific and Eno3 is a muscle specific isoform. In our transcriptomic data sets, we

observed expression of Eno1 and Eno3, but not Eno2. Even though COBRA analysis did not

distinguish between isoforms, we were able to experimentally demonstrate an isoform specific

function for Eno3 in cell survival during EMT. This is consistent with the kinetically more active

muscle specific Eno3 regulating energy-intensive migratory behavior of cancer cells. This

observation also fits with the broader trend between catalytic activity and over-expression

observed across various cancers (Oruganty et al. 2020).

Comparison of our model predictions against CRISPR knockout gene essentiality scores from

cancer cell lines revealed a significant correlation. Interestingly, single-cell knockout simulations

were more correlated with CRISPR-Cas9 gene knockout essentiality data than models derived

from bulk omics data. CRISPR-Cas9 essentiality screening is a promising high-throughput

approach to determine the contribution of individual genes on cell viability. The correlation

between our single-cell knockout simulations and CRISPR-Cas9 knockout essentiality data

suggests that our model captures vulnerabilities during EMT. In addition to Eno3, we found that

glutaminase (FTCD), 4-hydroxyphenylpyruvate oxidoreductase (HPD),

adenosylhomocysteinase (AHCY), and phosphatidylserine synthase (PTDSS1) to be novel

reactions that have no literature backing but have negative CERES Scores (i.e. impacts

viability) in NSCLC cancer cell lines. The reactions prioritized by our model are strong

candidates for drug development because they reduce cell growth in cells from later timepoints

(mesenchymal/metastatic-like) relative to earlier ones (epithelial/benign-like). In addition, our

model predicted ATP-Citrate lyase (ACLY) to be essential in mesenchymal-like cells. ACLY has

been implicated as a crucial metabolic enzyme that facilitates cancer progression and its

upregulation is associated with poor patient prognosis (Migita et al., 2008).

In summary, we present a computational model that captures metabolic activity and

gene essentiality during EMT. Our modeling approach can be applied to study metabolism at a

single-cell resolution and can capture the heterogeneity of other critical biological processes,

including tissue differentiation and development of disease states.

References

1. Pastushenko, I., & Blanpain, C. (2019). EMT Transition States during Tumor Progression and Metastasis. *Trends in Cell Biology*, *29*(3), 212–226.
2. Padua, D., & Massagué, J. (2009). Roles of TGFbeta in metastasis. *Cell Research*, *19*(1), 89–102.
3. Ramesh, V., Brabletz, T., & Ceppi, P. (2020). Targeting EMT in cancer with repurposed metabolic inhibitors. *Trends in Cancer*.
4. Vander Heiden, M. G., Cantley, L. C., & Thompson, C. B. (2009). Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science*, *324*(5930), 1029–1033.
5. Pålsson-McDermott, E. M., & O'Neill, L. A. J. (2020). Targeting immunometabolism as an anti-inflammatory strategy. *Cell Research*, *30*(4), 300–314.
6. Orth, J. D., Thiele, I., & Palsson, B. Ø. (2010). What is flux balance analysis? *Nature Biotechnology*, *28*(3), 245–248.
7. Oruganty, K., Campit, S. E., Mamde, S., Lyssiotis, C. A., & Chandrasekaran, S. (2020). Common biochemical properties of metabolic genes recurrently dysregulated in tumors. *Cancer & Metabolism*, *8*, 5.
8. Yizhak, K., Gaude, E., Le Dévédec, S., Waldman, Y. Y., Stein, G. Y., van de Water, B., Frezza, C., & Ruppin, E. (2014). Phenotype-based cell-specific metabolic modeling reveals metabolic liabilities of cancer. *Elife*, *3*, e03641.
9. Nilsson, A., Haanstra, J. R., Engqvist, M., Gerding, A., Bakker, B. M., Klingmüller, U., ... & Nielsen, J. (2020). Quantitative analysis of amino acid metabolism in liver cancer links glutamate excretion to nucleotide synthesis. *Proceedings of the National Academy of Sciences*, *117*(19), 10294-10304.
10. Nelson, B. S., Lin, L., Kremer, D. M., Sousa, C. M., Cotta-Ramusino, C., Myers, A., Ramos, J., Gao, T., Kovalenko, I., Wilder-Romans, K., Dresser, J., Davis, M., Lee, H.-J., Nwosu, Z. C., Campit, S., Mashadova, O., Nicolay, B. N., Tolstyka, Z. P., Halbrook, C. J., … Lyssiotis, C. A. (2020). Tissue of origin dictates GOT1 dependence and confers synthetic lethality to radiotherapy. *Cancer & Metabolism*, *8*, 1.
11. Hecker, L., Vittal, R., Jones, T., Jagirdar, R., Luckhardt, T. R., Horowitz, J. C., Pennathur, S., Martinez, F. J., & Thannickal, V. J. (2009). NADPH oxidase-4 mediates myofibroblast activation and fibrogenic responses to lung injury. *Nature Medicine*, *15*(9), 1077–1081.
12. Keshamouni, V. G., Jagtap, P., Michailidis, G., Strahler, J. R., Kuick, R., Reka, A. K., Papoulias, P., Krishnapuram, R., Srirangam, A., Standiford, T. J., Andrews, P. C., & Omenn, G. S. (2009). Temporal quantitative proteomics by iTRAQ 2D-LC-MS/MS and corresponding mRNA expression analysis identify post-transcriptional modulation of actin-cytoskeleton regulators during TGF-beta-Induced epithelial-mesenchymal transition. *Journal of Proteome Research*, *8*(1), 35–47.
13. Keshamouni, V. G., Michailidis, G., Grasso, C. S., Anthwal, S., Strahler, J. R., Walker, A., Arenberg, D. A., Reddy, R. C., Akulapalli, S., Thannickal, V. J., Standiford, T. J., Andrews, P. C., & Omenn, G. S. (2006). Differential protein expression profiling by iTRAQ-2DLC-MS/MS of lung cancer cells undergoing epithelial-mesenchymal transition reveals a migratory/invasive phenotype. *Journal of Proteome Research*, *5*(5), 1143–1154.
14. Lu, C., Sidoli, S., Kulej, K., Ross, K., Wu, C. H., & Garcia, B. A. (2019). Coordination between TGF-β cellular signaling and epigenetic regulation during epithelial to mesenchymal transition. *Epigenetics & Chromatin*, *12*(1), 11.
15. Law, C. W., Chen, Y., Shi, W., & Smyth, G. K. (2014). voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biology*, *15*(2), 1–17.
16. Cook, D. P., & Vanderhyden, B. C. (2020). Context specificity of the EMT transcriptional response. *Nature Communications*, *11*(1), 2142.
17. Van Dijk, D., Sharma, R., Nainys, J., Yim, K., Kathail, P., Carr, A. J., ... & Pe'er, D. (2018). Recovering gene interactions from single-cell data using data diffusion. *Cell*, *174*(3), 716-729.
18. Duarte, N. C., Becker, S. A., Jamshidi, N., Thiele, I., Mo, M. L., Vo, T. D., Srivas, R., & Palsson, B. Ø. (2007). Global reconstruction of the human metabolic network based on genomic and bibliomic data. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(6), 1777–1782.

19. Zur, H., Ruppin, E., & Shlomi, T. (2010). iMAT: an integrative metabolic analysis tool.
Bioinformatics , *26*(24), 3140–3142.

20. Shen, F., Boccuto, L., Pauly, R., Srikanth, S., & Chandrasekaran, S. (2019). Genome-scale
network model of metabolism and histone acetylation reveals metabolic dependencies of histone
deacetylase inhibitors. *Genome biology*, *20*(1), 1-15.

21. Lewis, N. E., Hixson, K. K., Conrad, T. M., Lerman, J. A., Charusanti, P., Polpitiya, A. D., Adkins,
J. N., Schramm, G., Purvine, S. O., Lopez-Ferrer, D., Weitz, K. K., Eils, R., König, R., Smith, R.
D., & Palsson, B. Ø. (2010). Omic data from evolved E. coli are consistent with computed optimal
growth from genome-scale models. *Molecular Systems Biology*, *6*, 390.

22. Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J. P., & Tamayo, P. (2015). The
molecular signatures database hallmark gene set collection. *Cell systems*, *1*(6), 417-425.

23. Vasaikar, S. V., Deshmukh, A. P., den Hollander, P., Addanki, S., Kuburich, N. A., Kudaravalli, S.,
... & Mani, S. A. (2021). EMTome: A resource for pan-cancer analysis of epithelial-mesenchymal
transition genes and signatures. *British journal of cancer*, *124*(1), 259-269.

24. Pacini, C., Dempster, J. M., Boyle, I., Gonçalves, E., Najgebauer, H., Karakoc, E., van der Meer,
D., Barthorpe, A., Lightfoot, H., Jaaks, P., McFarland, J. M., Garnett, M. J., Tsherniak, A., & Iorio,
F. (2021). Integrated cross-study datasets of genetic dependencies in cancer. *Nature
Communications*, *12*(1), 1661.

25. Barretina, J., Caponigro, G., Stransky, N., Venkatesan, K., Margolin, A. A., Kim, S., Wilson, C. J.,
Lehar, J., Kryukov, G. V., Murray, L., Morrissey, M. P., Sellers, W. R., Schlegel, R., & Garraway,
L. A. (2012). 22 The Cancer Cell Line Encyclopedia - Using Preclinical Models to Predict
Anticancer Drug Sensitivity. In European Journal of Cancer (Vol. 48, pp. S5–S6). doi:
10.1016/s0959-8049(12)70726-8

26. O'Brien, E. J., Monk, J. M., & Palsson, B. O. (2015). Using genome-scale models to predict
biological capabilities. *Cell*, *161*(5), 971-987.

27. Vander Heiden, M. G., Cantley, L. C., & Thompson, C. B. (2009). Understanding the Warburg
effect: the metabolic requirements of cell proliferation. *science*, *324*(5930), 1029-1033.

28. Jia, D., Park, J. H., Kaur, H., Jung, K. H., Yang, S., Tripathi, S., Galbraith, M., Deng, Y., Jolly, M.
K., Kaipparettu, B. A., Onuchic, J. N., & Levine, H. (2021). Towards decoding the coupled
decision-making of metabolism and epithelial-to-mesenchymal transition in cancer. *British Journal
of Cancer*. doi: 10.1038/s41416-021-01385-y

29. Wei, Q., Qian, Y., Yu, J., & Wong, C. C. (2020). Metabolic rewiring in the promotion of cancer
metastasis: mechanisms and therapeutic implications. *Oncogene*, *39*(39), 6139–6156.

30. Campit, S. E., Meliki, A., Youngson, N. A., & Chandrasekaran, S. (2020). Nutrient Sensing by
Histone Marks: Reading the Metabolic Histone Code Using Tracing, Omics, and
Modeling. *BioEssays*, *42*(9), 2000083.

31. Song, Y., Luo, Q., Long, H., Hu, Z., Que, T., Zhang, X. 'an, Li, Z., Wang, G., Yi, L., Liu, Z., Fang,
W., & Qi, S. (2014). Alpha-enolase as a potential cancer prognostic marker promotes cell growth,
migration, and invasion in glioma. *Molecular Cancer*, *13*, 65.

32. Zhao, M., Fang, W., Wang, Y., Guo, S., Shu, L., Wang, L., Chen, Y., Fu, Q., Liu, Y., Hua, S., Fan,
Y., Liu, Y., Deng, X., Luo, R., Mei, Z., Jiang, Q., & Liu, Z. (2015). Enolase-1 is a therapeutic target
in endometrial carcinoma. *Oncotarget*, *6*(17), 15610–15627.

33. Hou, X.-M., Yuan, S.-Q., Zhao, D., Liu, X.-J., & Wu, X.-A. (2019). LDH-A promotes malignant
behavior via activation of epithelial-to-mesenchymal transition in lung adenocarcinoma.
Bioscience Reports, *39*(1). doi: 10.1042/BSR20181476

34. Zhang, Y., Lin, S., Chen, Y., Yang, F., & Liu, S. (2018). LDH-Apromotes epithelial-mesenchymal
transition by upregulating ZEB2 in intestinal-type gastric cancer. *OncoTargets and Therapy*, *11*,
2363–2373.

35. Menendez, J. A., Vellon, L., Mehmi, I., Oza, B. P., Ropero, S., Colomer, R., & Lupu, R. (2004).
Inhibition of fatty acid synthase (FAS) suppresses HER2/neu (erbB-2) oncogene overexpression
in cancer cells. *Proceedings of the National Academy of Sciences of the United States of
America*, *101*(29), 10715–10720.

36. Pizer, E. S., Wood, F. D., Heine, H. S., Romantsev, F. E., Pasternack, G. R., & Kuhajda, F. P.
(1996). Inhibition of fatty acid synthesis delays disease progression in a xenograft model of
ovarian cancer. *Cancer Research*, *56*(6), 1189–1193.

37. Xu, X., Chen, B., Zhu, S., Zhang, J., He, X., Cao, G., & Chen, B. (2019). Hyperglycemia promotes Snail-induced epithelial–mesenchymal transition of gastric cancer via activating ENO1 expression. *Cancer Cell International*, *19*(1), 1–12.

38. Liu, M., Quek, L.-E., Sultani, G., & Turner, N. (2016). Epithelial-mesenchymal transition induction is associated with augmented glucose uptake and lactate production in pancreatic ductal adenocarcinoma. *Cancer & Metabolism*, *4*, 19.

39. Birsoy, K., Wang, T., Chen, W. W., Freinkman, E., Abu-Remaileh, M., & Sabatini, D. M. (2015). An Essential Role of the Mitochondrial Electron Transport Chain in Cell Proliferation Is to Enable Aspartate Synthesis. *Cell*, *162*(3), 540–551.

40. Alkan, H. F., Walter, K. E., Luengo, A., Madreiter-Sokolowski, C. T., Stryeck, S., Lau, A. N., Al-Zoughbi, W., Lewis, C. A., Thomas, C. J., Hoefler, G., Graier, W. F., Madl, T., Vander Heiden, M. G., & Bogner-Strauss, J. G. (2018). Cytosolic Aspartate Availability Determines Cell Survival When Glutamine Is Limiting. *Cell Metabolism*, *28*(5), 706–720.e6.

41. Sullivan, L. B., Gui, D. Y., Hosios, A. M., Bush, L. N., Freinkman, E., & Vander Heiden, M. G. (2015). Supporting Aspartate Biosynthesis Is an Essential Function of Respiration in Proliferating Cells. *Cell*, *162*(3), 552–563.

42. Yang, L., Venneti, S., & Nagrath, D. (2017). Glutaminolysis: a hallmark of cancer metabolism. *Annual review of biomedical engineering*, *19*, 163-194.

43. Cacace, A., Sboarina, M., Vazeille, T., & Sonveaux, P. (2017). Glutamine activates STAT3 to control cancer cell proliferation independently of glutamine metabolism. *Oncogene*, *36*(15), 2074-2084.

44. Yang, L., Moss, T., Mangala, L. S., Marini, J., Zhao, H., Wahlig, S., ... & Nagrath, D. (2014). Metabolic shifts toward glutamine regulate tumor growth, invasion and bioenergetics in ovarian cancer. *Molecular systems biology*, *10*(5), 728.

45. Sellers, K., Fox, M. P., Ii, M. B., Slone, S. P., Higashi, R. M., Miller, D. M., Wang, Y., Yan, J., Yuneva, M. O., Deshpande, R., Lane, A. N., & Fan, T. W.-M. (2015). Pyruvate carboxylase is critical for non–small-cell lung cancer proliferation. *The Journal of Clinical Investigation*, *125*(2), 687–698.

46. Fu, X.-G., Deng, J., Xu, W.-J., Chen, J.-Y., Sun, J., & Deng, H. (2020). Histidine decarboxylase-expressing PMN-MDSC-derived TGF-β1 promotes the epithelial-mesenchymal transition of metastatic lung adenocarcinoma. *International Journal of Clinical and Experimental Pathology*, *13*(6), 1361–1371.

47. Meyers, R. M., Bryan, J. G., McFarland, J. M., Weir, B. A., Sizemore, A. E., Xu, H., ... & Tsherniak, A. (2017). Computational correction of copy number effect improves specificity of CRISPR–Cas9 essentiality screens in cancer cells. *Nature genetics*, *49*(12), 1779-1784.

48. Migita, T., Narita, T., Nomura, K., Miyagi, E., Inazuka, F., Matsuura, M., ... & Ishikawa, Y. (2008). ATP citrate lyase: Activation and therapeutic implications in non–small cell lung cancer. *Cancer research*, *68*(20), 8547-8554.

Data and software availability statement

Bulk transcriptomics data was obtained from GSE17708 and GSE17518. Bulk proteomics data was obtained from Keshamouni et al., 2006 and Lu et al., 2019. Single-cell EMT transcriptomics data was obtained from GSE147405.

All COBRA data and meta-analyses performed can be found in the supplementary table.

All scripts used to analyze these datasets can be found in this GitHub repository.

Main Figures and Data Tables

Figure 1. A549 metabolism is predicted to be overactive during the early and late phases of EMT.

    A. Reactions are sorted based on the priority score, which is a function of the number of studies with significant metabolic genes that encode the reaction and the effect size (Z-score or log2 fold change). The metabolic fluxes were simulated using the RECON1 human metabolic reconstruction. Several metabolic reactions within the Glycolysis/Gluconeogenesis metabolic subsystem are overactive in the earlier stages (1hr) and late stages (48-72hrs) of EMT, based on the absolute value of the metabolic fluxes predicted by constraint-based modeling. The top 5 reactions in the Glycolysis/Gluconeogenesis subsystem have at least two studies supporting the flux predictions. Enolase is bolded as it was prioritized for experimental validation.

    B. Metabolic reactions within the Citric Acid Cycle are predicted to have more uniform activity across all time points relative to control (unconstrained flux distribution) and have lower priority scores compared to glycolysis.

    C. Metabolic reactions within the Oxidative Phosphorylation metabolic subsystem are also predicted to have more uniform activity relative to control across all time points.

689
690

Figure 2. A global view of cancer growth sensitivity to metabolic reaction knockout
during EMT. Normalized KO Growth Scores closer to a value of -1 confer a decrease in growth
relative upon simulated reaction knockout relative to the control (an unconstrained metabolic
reconstruction growth rate). Reactions are sorted based on priority scores.
The barplots (inset) show the distribution of lethal reactions (Growth Score < 0) for three
timepoints 1 hour, 24 hours, and 48-72 hours after TGF-β induction across all experiments in 3
time points. Reactions with red and bold text were predicted to be highly sensitive to knockout
and were prioritized for downstream analyses with CERES essentiality scores.
699

Figure 3. siRNA-mediated inhibition of Eno3, but not Eno1, triggers apoptosis in A549
cells undergoing EMT after 72 h TGF-b treatment.
A. Apoptosis is assessed by the percentage of AnnexinV & PI positive cells by Flow
cytometry.
B. Caspase activation is measured using a caspase8 specific substrate that fluoresces
after caspase 8-mediated cleavage.
708
709

Figure 4. Single-cell COBRA reaction knockout analysis is consistent with results from
bulk studies. Reaction knockout growth rates for the top 40 most variable reactions are shown
in the heatmap. Column1 shows averaged data across all cells in the single-cell simulations,
and data from cells grouped by time points are shown in subsequent columns (day 0, 8 hrs, day
1, day 3 and day 7). This list contains many reactions that were also found to be sensitive upon
knockout in bulk studies (Table 1).
720

Figure 5. Visualization of Single-cell COBRA reaction knockout data.
A. Shown is the UMAP visualization of the temporal trajectory of cells induced with TGF-B.
B – F. Single-cell knockout growth rates were overlaid onto a UMAP embedding for A549 TGF-B single cell data. Growth rates were scaled from 0 to 1, where 1 indicates no change in growth rate between the cancer cell and control, while 0 indicates cell death in the cancer cell relative to control. Selected single-cell growth rate profiles for reactions that were sensitive in bulk reaction knockout simulations are shown in B-E. Growth scores (g.s.) were discretized into sensitive (g.s. < 0; red), not sensitive (g.s. = 0; light gray), mildly enhancing (0 < g.s. < 0.3; dark gray) and enhancing (g.s. > 0.3; black). The citrate synthase reaction (panel F) was selected as a control as it is an essential metabolic reaction, and it correctly shows sensitivity across all time points.

A.

B.

C.

Figure 6. COBRA-prioritized NSCLC CERES Scores reveal metabolic vulnerabilities during EMT.

A. Frequency matrix for comparing the Cancer Cell Line Encyclopedia (CCLE) primary/metastatic annotations against our Epithelial/Mesenchymal annotations based on MSigDB and EMTome signatures. Cells from the CCLE were classified as epithelial or mesenchymal based on the number of genes were up/downregulated that matched the MSigDB/EMTome signatures (**Methods**).

B. Reactions predicted by COBRA to be sensitive were compared against CERES Scores. Cell lines were classified as primary or metastatic, and their distributions are shown on the violin plots. Overall, the average predicted growth scores and fluxes in hour 72 across all 5 experiments agreed with the Primary / Metastatic CERES Scores Ratios (R = 0.31 and 0.2; P-value = 0.005 and 0.035 respectively; **S. Table 6 and 7)**. These 10 metabolic genes were selected based on reactions of interest from our bulk COBRA knockout profiles and single-cell flux profiles (from **Figure 2**).

C. The same analysis was repeated with the epithelial/mesenchymal annotations for sensitive reactions. The predicted growth scores and fluxes from the single-cell simulations (GSE147405) agreed with the Epithelial / Mesenchymal CERES Scores Ratios (R = 0.28, p-value=0.01).

Main Tables and Legends

Table 1. Literature review of known and novel essential reactions predicted from both bulk and single-cell simulations**.**

For full list of all predicted essential reactions, see **S. Table 5.**

Reaction Name	Gene-Protein-Reaction Rule	Average NSCLC CERES Score	PMID
Strong evidence			
pyruvate carboxylase	(PC)	-0.5217	26070193 30005601 25607840 32530543
fructose-bisphosphate aldolase	(ALDOA) or (ALDOC) or (ALDOB)	-0.7012	28444969 31358528
Medium evidence			
ribulose 5-phosphate 3-epimerase	(RPE)	-1.3731	32365991
triose-phosphate isomerase	(TPI1) or (TPIP2)	-1.1346	27908734
formimidoyltransferase cyclodeaminase	(FTCD)	-0.0908	30784016
glyceraldehyde-3-phosphate dehydrogenase	(GAPDH) or (GAPDHS)	-0.7870	27878251
UTP-glucose-1-phosphate uridylyltransferase	(UGP2)	-0.2952	31243371
ATP-Citrate lyase	(ACLY)	-0.7807	23807225
acetone mitochondrial transport via proton symport	(SLC16A1)	-0.0556	31371390
aspartate transaminase	(GOT2)	-0.3341	23535601
Novel predictions			
glutaminase (mitochondrial)	(FTCD)	-0.0908	
4-Hydroxyphenylpyruvate:oxygen oxidoreductase	(HPD)	-0.0390	
adenosylhomocysteinase	(AHCY) or (AHCYL1)	-0.6535	
enolase	(ENO1) or (ENO3) or (ENO2)	-0.0015	
Phosphatidylserine synthase homo sapiens	(PTDSS1)	-0.2514	

757
758

Supplementary Figure 1. The top 50 reactions that are predicted to be overactive are ranked by priority score. (Related to figure 1)
The priority score is a function of the number of studies with significant metabolic genes that encode the reaction and the effect size (Z-score or log2 fold change). The metabolic fluxes were simulated using the RECON1 metabolic reconstruction. Several EMT associated metabolic reactions predicted by our model such as Glycolysis/Gluconeogenesis, Glutamine metabolism and Nucleic acid metabolism are commonly dysregulated in cancer.

771
772
773

Supplementary Figure 2. Top 50 reactions ranked by priority score predicted to be
sensitive to reaction KO. (related to figure 2) The heatmap shows the top 50 reactions
ranked by priority score predicted to reduce A549 growth rate upon reaction knockout in
RECON1 in all bulk experiments. In contrast, Figure 2 focuses on reactions sensitive in specific
EMT stages and studies, while this sensitivity profile shows the top 50 reactions by priority
score. Eight nutrient exchange reactions including glucose and aspartate exchange reactions
were predicted to have a negative impact on growth following knockout at different stages of
EMT. Glutamine exchange interestingly decreased growth upon KO during early and late EMT
timepoints but not in intermediate time-point (24hrs). Hexokinase and C100 fatty acid activation
were predicted to be essential across all time points. Pyruvate carboxylase was predicted to be
sensitive in early EMT (1 hour).

Supplementary Figure 3. mRNA expression fold change of ENO1 versus ENO3 over time
788

790 **Supplementary Figure 4. Single-cell RNASeq data of EMT biomarkers reveals cell states.**
791   A. Time-course trajectory of A549 cells induced with TGF-B.
792   B. VIM expression over time.
793   C. CDH1 expression over time.
794

A.

STAT3 and Glutamuine Synthetase A549 scRNASeq

R = 0.444 ; P-value = 0.002

y = 0.269 + 0.38 x

GLUL (Normalized Expression)

STAT3 (Normalized Expression)

B.

STAT3 and SLC7A5 A549 scRNASeq

R = 0.158 ; P-value = 0.017

y = 0.954 + 0.27 x

SLC7A5 (Normalized Expression)

STAT3 (Normalized Expression)

Supplementary Figure 5. Correlation analysis between STAT3 and glutamine metabolism-related genes Glutamine Synthetase and glutamine transporter SLC7A5.

A. Shown are cells that contain non-zero expression levels for both STAT3 and GLUL across cells with TGF-β induction. A significant positive correlation (R = 0.444; P-value = 0.002) was observed.

B. Shown are cells that contain non-zero expression levels for both STAT3 and SLC7A5 across cells with TGF-β induction. A significant but weaker positive correlation (R = 0.158; P-value = 0.017) was observed.