# Non-canonical scaffold-type ligase complex mediates protein UFMylation

Joshua J. Peter[1], Helge M. Magnussen[1], Paul Anthony DaRosa[2], David Millrine[1],

Stephen P Matthews[1], Frederic Lamoliatte[1], Ramasubramaniam Sundaramoorthy[3],

Ron R Kopito[2] and Yogesh Kulathu[1]*

**Affiliations –**

[1] Medical Research Council Protein Phosphorylation & Ubiquitylation Unit (MRC-

PPU), School of Life Sciences, University of Dundee, Dundee DD1 5EH, UK.

[2] Department of Biology, Stanford University, Stanford, CA 94305, USA

[3] Centre for Gene Regulation and Expression, School of Life Sciences, University of

Dundee, Dundee DD1 5EH, UK

**\*corresponding author –**

Dr Yogesh Kulathu, Medical Research Council Protein Phosphorylation &

Ubiquitylation Unit (MRC-PPU), School of Life Sciences, University of Dundee, Dow

Street, Dundee DD1 5EH, UK.

Tel: +44 1382 388163; E-mail: ykulathu@dundee.ac.uk

Running Title: Mechanism of UFMylation

**Abstract**

Protein UFMylation is emerging as a posttranslational modification essential for
endoplasmic reticulum and cellular homeostasis. Despite its biological importance,
we have a poor understanding of how UFM1 is conjugated onto substrates. Here, we
use a rebuilding approach to define the minimal requirements of protein UFMylation.
We find that the reported E3 ligase UFL1 is inactive on its own and identify UFBP1 to
bind UFL1 to form an active E3 ligase complex. While UFC1 is an intrinsically Cys-
reactive E2, we do not identify any catalytic cysteines on UFL1/UFBP1, suggesting a
scaffold-type E3 ligase mechanism. Interestingly, the E3 ligase complex consists of
winged-helix (WH) domain repeats that activates UFC1 for aminolysis. We identify
the adaptor protein CDK5RAP3 to bind to and regulate E3 ligase activity potentially
by preventing off-target UFMylation. In summary, our work identifies the minimal
requirements for UFMylation and reveals regulatory principles of this atypical E3
ligase complex.

UFM1 is a ubiquitin-like (Ubl) modifier that contains the characteristic β-grasp fold found in all UBLs and is highly conserved in multicellular eukaryotes [1,2]. Like other UBLs, UFM1 is covalently attached to lysine residues on substrates via an enzymatic cascade involving an E1 activating enzyme, UFM1-activating enzyme 5 (UBA5), E2 conjugating enzyme, UFC1, and an E3 ligase, UFL1 (UFM1 E3 Ligase 1) [3,4]. Recent studies have identified the ribosomal protein RPL26 to be the principal target of UFMylation which plays a critical role in endoplasmic reticulum (ER) homeostasis [5–7]. Besides ER associated roles, UFMylation has also been implicated in other cellular processes including protein translation, DNA damage response and nuclear receptor mediated transcription [8–13]. Further, loss of or mutation of components of the UFMylation machinery has been linked to many diseases such as cancer, type-2 diabetes, neurological disorders and cerebellar ataxia, where failure of ER homeostasis and protein quality control could be one of the major contributing factors [12,14–17]. Indeed, UFMylation is an essential posttranslational modification for animal development as loss of UFM1 or any of the UFMylation enzymes results in failure of erythropoiesis and embryonic lethality [18–21].

UFM1 is expressed in a precursor form consisting of 85 amino acids. It is then post-translationally processed by UFM1-specific proteases (UFSPs) which cleave off the last two residues at the C-terminus to generate mature UFM1 that contains an exposed C-terminal glycine, Gly[83] [22]. Mature UFM1 is activated by the E1, UBA5 via the formation of a high energy thioester bond between Cys[250] of UBA5 and Gly[83] of UFM1 [3,23]. Activated UFM1 is then transferred from UBA5 on to the catalytic Cys[116] of UFC1 [3,24]. UFC1 has a canonical UBC fold and in addition an N-terminal helical extension whose function is not known [25,26]. UFC1 lacks many of the features conserved in many E2s such as the catalytic HPN motif which suggests unique modes of action and regulation. Further, ubiquitin E2 enzymes such as UBE2D3 associate with ubiquitin via a backside interaction which mediates E2 dimerization and processivity of polyubiquitin formation [27,28]. For UFC1, it is unknown if similar mechanisms exist to regulate its activity as detailed biochemical characterization is lacking.

UFL1 is the only E3 ligase identified to date in the UFMylation pathway catalysing the final step of transfer of UFM1 from UFC1 to substrates [29]. Indeed, loss of UFL1 results in loss of UFMylation and mice lacking Ufl1 phenocopy loss of UFM1

exhibiting failure in haematopoiesis and embryonic lethality suggesting that it could

be the main E3 ligase [30]. While data implying UFL1 is the E3 ligase stem from

knockout studies and overexpression studies, direct biochemical evidence

demonstrating that UFL1 is an E3 ligase is insufficient [24]. Consequently, the

mechanism of how UFL1 functions as an E3 ligase is also not understood. In

general, E3 ligases are classified either as scaffold-type ligases that bring together a

UBL-charged E2 with substrate for direct transfer of the UBL from the E2 on to

substrate, or as Cys-dependent enzymes where the UBL is first transferred from the

E2 to the catalytic Cys of the E3 for subsequent transfer to the substrate [31].

Examples of scaffold-type E3s are RING family ligases and RANBP2 whereas

HECT, RBR and RCR enzymes typify the latter type [32–36]. Intriguingly, UFL1 does

not possess any conserved sequence or domain features found in known E3 ligases.

Hence it is not clear if UFL1 is a Cys-dependent E3 enzyme or uses a scaffolding

mechanism.

UFL1 is predominantly found anchored at the ER through its interaction with an

adaptor protein, DDRGK1/UFBP1 (UFM1 binding protein-1) which has also been

suggested to be one of the main substrates of UFL1 [29,37]. UFBP1 localizes to the ER

membrane via an N-terminal transmembrane segment and its function is unclear [5].

Intriguingly, loss of UFBP1 affects the stability and expression levels of UFL1 [20,29,37].

CDK5RAP3 is another poorly characterized protein commonly associated with UFL1.

Several anecdotal reports have led to speculations that UFL1, UFBP1 and

CDK5RAP3 together form an E3 ligase complex [24,38]. Only a handful of substrates of

UFL1 have been identified to date and many of them are ER localized [38,39]. In

addition to monoUFMylation, polyUFM1 chains can be assembled which are mainly

linked via K69 [12]. In the ubiquitin system, the type of polyUb linkage formed is

generally dictated by the last enzyme in the cascade that forms a thioester linkage

with ubiquitin before transfer to substrate. Scaffold-type E3s such as RING E3

ligases bind to charged E2 (E2~Ub) to mediate transfer of Ub onto susbtrate, and

hence the E2 enzymes are thought to determine the linkage type assembled. In

contrast, catalytic Cys containing E3 ligases form an E3~Ub thioester intermediate

and dictate the linkage type formed [31,34].

Given the lack of understanding of the molecular mechanism underpinning the UFM1

conjugation machinery and the mechanisms governing its regulation, here we adopt

a rebuilding approach with purified components to fully reconstitute UFMylation *in*
*vitro*. Using this reductionist approach, we reveal that UFL1 on its own is an unstable
protein that cannot support UFMylation. We demonstrate that UFL1 together with
UFBP1 forms a functional E3 ligase complex and provide in depth biochemical
characterization of the ligase complex. Structure prediction using Alphafold provides
an explanation for the underlying reason for complex formation between UFL1 and
UFBP1, and offers a rational framework for making truncations which help establish
the minimal regions on UFL1 and UFBP1 essential for ligase activity. Parallel
investigations of UFC1 reveal that it is an intrinsically Cys reactive enzyme that is
activated by the UFL1/UFBP1 ligase complex for aminolysis. Our analyses reveal
UFL1/UFBP1 to be a scaffold-type E3 ligase. Further, we identify CDK5RAP3 to bind
to the E3 ligase complex and inhibit UFMylation by preventing discharge of UFM1
from UFC1. Importantly, by reconstituting UFMylation of ribosomes *in vitro,* we find
that CDK5RAP3 has a regulatory function as it restricts UFMylation towards *bona*
*fide* substrates and lysine selectivity.  Lastly, we identify an N-terminal helical
extension on UFC1 to have an inhibitory role that is essential for CDK5RAP3 to
restrain UFMylation. Our mechanistic insights describe an atypical E3 ligase
complex and provide a foundation for investigating the catalytic mechanism,
regulation, and substrate specificity of this E3 ligase complex.

**Results**

**UFC1 has intrinsic cysteine reactivity**

The interaction of UFC1 with UBA5 and how UFM1 is transferred from UBA5 to the
catalytic Cys is reasonably well understood [23,24,40–42]. We therefore focussed on
subsequent events and first analysed the intrinsic reactivity and stability of
UFC1~UFM1 (~ denotes the thioester bond between UFM1 G83 and UFC1 C116) in
comparison to the well characterized ubiquitin E2s, UBE2D3 and UBE2L3 [43].
Intrinsic reactivity of E2s assayed using free amino acids can give insights into the
ability of E2s to transfer UBLs onto substrates and can provide clues about the type
of E3 ligase they work with and even the nature of the substrate [27,36]. For instance,
E2s such as UBE2D3 which function together with RING/scaffold like E3 ligases are
capable of aminolysis and are therefore reactive towards the free amino acid Lys
(**Fig 1A, B**) [43]. On the other hand, UBE2L3 which function only with Cys-dependent
E3 ligases is incapable of aminolysis and lack reactivity to free Lys on its own (**Fig
1B**) [43]. When discharge of UFM1 from UFC1~UFM1 onto free amino acids was
analysed, we observed preferential discharge onto Cys while no Lys discharge was
observed (**Fig 1C**). Under the same experimental conditions, UBE2D3 discharged
Ub in the presence of both Cys and Lys (**Fig 1C, Fig S1**). Further, using increasing
concentrations of free Lys or with prolonged incubation in a time course, we find that
UFC1 has negligible Lys reactivity (**Fig 1D-E**). These results suggest that UFC1 has
intrinsic Cys reactivity and may work with Cys-dependent E3 ligases.

**UFL1 together with UFBP1 forms an active E3 ligase complex**

To analyse how UFC1 works together with the E3 ligase UFL1 to transfer UFM1, we
aimed to reconstitute UFMylation using UFL1.  We expressed 6xHis-UFL1 in *E. coli*
and purified it by affinity chromatography as the first step.  Subsequent size
exclusion chromatography (SEC) analysis indicated UFL1 to elute in the void fraction
suggesting formation of soluble aggregates (**Fig S2A**). Our attempts to enhance
solubility by incorporating different solubility enhancing tags at the N-terminal or C-
terminal ends of UFL1 or with different buffers and additives were unsuccessful. We
therefore reasoned that UFL1 might require additional binding partners for its stability
and activity. Indeed, several studies have shown that expressing subunits of protein
complexes individually may result in aggregation [44,45]. To identify binding partners of

UFL1, we performed Yeast Two Hybrid (Y2H) screening using a cDNA library
derived from human placenta as prey (**Fig S2B**). Surprisingly, only three hits were
identified with UFM1 binding protein 1 (UFBP1, a.k.a DDRGK1) being the most
significant interactor. We therefore wondered if co-expressing UFBP1 with UFL1
might confer solubility and stability to UFL1. To achieve this, we used a co-
expression system to express full-length UFL1 together with UFBP1 lacking its
transmembrane sequence and purified the ligase complex with three enrichment
steps (**Fig S2C**). Indeed, UFL1 co-expressed with UFBP1 lacking its N-terminal
transmembrane segment (UFL1/UFBP1) is soluble, well behaved and no longer
formed soluble aggregates in SEC (**Fig 2A**). We confirmed the formation of a stable
UFL1/UFBP1 complex by analysing its precise molecular mass using a mass
photometer [46] and SEC-MALS (**Fig 2B, S2D**). This revealed UFL1/UFBP1 to have a
mass of 126 kDa which corresponds to that of a heterodimeric complex. Of note, the
mass photometry measurements are performed at low nanomolar concentrations
highlighting the stable nature of the complex.

Having obtained soluble UFL1/UFBP1 complexes, we next tested the catalytic
activity of UFL1 expressed alone and in complex with UFBP1 using *in vitro*
UFMylation assays containing UBA5, UFC1 and UFM1. Whereas UFL1 on its own
did not show any appreciable activity, the UFL1/UFBP1 complex is an active ligase
as evidenced by the formation of multiple UFMylated products and UFL1
autoUFMylation (**Fig 2C**). To check if UFL1/UFBP1 was capable of UFMylating
substrates *in vitro*, we used TRIP4/ASC1, a protein reported to be UFMylated in cells
[12]. Indeed, UFL1/UFBP1 can efficiently UFMylate TRIP4 *in vitro* (**Fig 2C**).
Intriguingly, exogenous addition of purified UFBP1 to UFL1 expressed on its own
does not change the elution profile of UFL1 to match that of co-expressed
UFL1/UFBP1 or in its E3 ligase activity (**Fig S2E-F**). Taken together, these results
show that UFL1 and UFBP1 form an obligate heterodimer required for E3 ligase
activity.

Since polyUFMylated products are formed in UFMylation reactions, we next
analysed the reaction products by mass spectrometry to determine the linkage types
assembled. Interestingly, LC/MS analyses revealed that polyUFM1 chains are
formed which are predominantly linked via K69 with some K3 and K7 linkages also

observed (**Fig 2D**). To validate these observations, we generated a series of K-R
mutants where every Lys in UFM1 was individually mutated to Arg. While mutation of
K3, K7, K19, K34 or K41 to Arg did not affect overall UFMylation, UFM1 K69R
resulted in near complete loss of polyUFMylated species and unanchored diUFM1,
closely resembling reactions with UFM1 K0 in which all Lys were mutated to Arg (**Fig**
**2E** lanes 7&8**, S2G**). Interestingly, western blotting for UFL1 revealed only
monoUFMylated UFL1 in the presence of UFM1 K69R suggesting that UFL1
autoUFMylation involves formation of polyUFM1 chains on UFL1 that are linked via
K69 (**Fig 2E**). We next generated $K_{only}$ UFM1 mutants that contain only a single Lys
while all others are mutated to Arg. UFMylation assays using these mutants confirm
that K69 is the preferred linkage type assembled (**Fig 2F**). Single Lys-retaining
mutants all abrogate polyUFMylation formation comparably to UFM1 K0 with the sole
exception of K69-only UFM1 (**Fig 2F**). In reactions with UFM1 K69R or K0, mainly
monoUFMylated products corresponding to modified UFL1, UFBP1 and UFC1 are
formed. In summary, our reconstitution experiments reveal that UFL1 together with
UFBP1 forms an active E3 ligase complex that can efficiently UFMylate substrates
and assemble polyUFM1 chains that are K69-linked.

**UFL1/UFBP1 is a scaffold type E3 ligase complex activating UFC1 for**
**aminolysis**

Our analyses find that UFC1 is an intrinsically Cys-reactive E2 suggesting that UFL1
may be a Cys-dependent enzyme (**Fig 1**). The N-terminal region of UFL1 contains
four Cys and sequence analysis show that all four Cys vary in their degree of
conservation with C32 being the most conserved (**Fig 3A**). Based on analysis of
predicted folding propensity (**Fig S3A**) and secondary structure prediction, we made
a C-terminal truncation of UFL1, 1-410, which when expressed on its own formed
soluble aggregates like full length UFL1 and required co-expression of UFBP1 to
obtain a heterodimeric complex (**Fig S3B**). Importantly, this UFL1$^{1-410}$/UFBP1
complex is an active ligase **(Fig S3C**). To identify the catalytic Cys in UFL1, we
mutated each Cys individually to Ala in the UFL1$^{1-410}$/UFBP1 complex. To our
surprise, *in vitro* UFMylation assays showed that none of the single mutants had any
impact on the autoUFMylation activity of UFL1 or UFMylation of substrates implying
that UFL1 does not contain a catalytic Cys (**Fig 3B**). Since UFBP1 does not have
any Cys residues, these results suggest that the UFL1/UFBP1 ligase complex may

instead employ a scaffolding mechanism to transfer UFM1 on to substrate. Further,
we do not observe any transthiolation products of UFL1/UFBP1 **(Fig S3D)**.

This raises the possibility that the UFL1/UFBP1 ligase complex could induce
aminolysis of UFC1~UFM1.   Hence, we compared discharge of UFM1 from
UFC1~UFM1 onto Lys in the absence and presence of the ligase complex. Whereas
UFC1 does not discharge UFM1 onto Lys on its own, UFM1 was readily discharged
onto Lys in the presence of UFL1/UFBP1 (**Fig 3D**). Based on these observations, we
suggest that UFL1/UFBP1 functions as a scaffold-type E3 ligase that binds to
charged UFC1 to promote aminolysis.

While inefficient, UFC1 on its own can assemble free UFM1 chains which is
significantly enhanced in the presence of the UFL1/UFBP1 ligase complex (**Fig S3E-**
**F**). Interestingly, UFC1 assembles mainly K69 linkages (**Fig 3D**) and this linkage
specificity is maintained in the presence of UFL1/UFBP1 (**Fig 2D**). Of note, this
reveals that the linkage specificity is determined by the E2 and is not altered by the
E3 ligase complex, a feature commonly observed in ubiquitin RING E3 ligases
[34,47,48]. These results further strengthen our conclusion that UFL1/UFBP1 is a
scaffold-type ligase complex.

**Tandem WH domains of UFL1/UFBP1 constitute minimal E3 ligase**

As UFL1 does not have any obvious sequence or domain features found in any of
the known E3 ligases, we attempted to define the minimal catalytic region on UFL1.
As our efforts to determine the structure of this complex were not successful, we
used the recently released Alphafold pipeline to predict the structure (**Fig 4A**) [49].
Structure prediction of the complex using Multimer had good PAE scores (**Fig S4A**)
and shows UFL1 and UFBP1 form a heterodimer (**Fig 4A-B**). Interestingly, UFL1 is
predicted to contain 5 winged-helix (WH) domain repeats and a C-terminal helical
domain (**Fig 3A** blue, S4B). The WH repeats are arranged such that a helix from
each WH domain forms a helical backbone. The fifth WH domain contains a stack of
α-helices at its C-terminus. These WH domains could be classified as PCI-like WH
domains, named so because it is commonly found in components of the proteasome
lid, the COP9 signalosome and the eukaryotic translation initiator, eIF3 [50,51]. In
addition, UFL1 is predicted to have an N-terminal helix (1-25) followed by a partial or
hemi-WH domain (pWH) (**Fig 4B,** blue). Following the transmembrane segment at its

N-terminus, UFBP1 has a long coiled-coil region (NTR) followed by a WH (WH')
domain and ends with a hemi-WH domain (pWH') (**Fig 4B, teal**). Interestingly, the
hemi-WH (pWH) from UFL1 complements a hemi WH (pWH') from the C-terminus of
UFBP1 to form a complete WH domain (**Fig 4B**), thus providing an explanation to
why UFBP1 is required to stabilize UFL1. Superposition of the WH domains of UFL1
and UFBP1 reveal them to have identical folds except for a β-strand which is lacking
in the composite WH domain (**Fig S4C-D**). The coiled-coil nature of UFBP1 and the
helical backbone of UFL1 may explain the unusual migration of UFL1/UFBP1 by
SEC. To validate the predicted structure of the UFL1/UFBP1 complex, we deleted
the N-terminal helix along with pWH of UFL1 and checked for complex formation
with UFBP1. No soluble protein was obtained upon deletion of this region of UFL1.
Likewise, removal of C-terminal pWH domain of UFBP1 also resulted in poor
expression, highlighting the importance of the formation of the composite WH (pWH-
pWH') domain for complex formation and protein stability.

Guided by the insights from the Alphafold predictions, we made additional
truncations to map the minimal catalytic domain of the ligase complex (**Fig 4C**). The
truncated complexes expressed well and were purified to homogeneity (**Fig S3F**).
We first analysed the ability of the different truncated complexes to promote
discharge of UFM1 from UFC1~UFM1 onto Lys. This revealed that all the
truncations tested could discharge UFM1 and the smallest region able to promote
discharge contains one WH domain each from UFL1 and UFBP1 and the composite
WH domain (**Fig 4D,S5A-B**).

We then monitored the ability of the different UFL1/UFBP1 complexes to bind to
UFC1 for which we performed analytical SEC of the different UFL1/UFBP1
constructs with UFC1 on its own or UFC1$^{\text{-O-UFM1}}$ where UFM1 is linked to C116S via
an oxyester bond (**Fig S5 C-D**). UFC1$^{\text{-O-UFM1}}$ is more stable compared to thioster-
linked UFC1~UFM1 and therefore used for these analyses. Post preparation, we
incubated UFC1$^{\text{-O-UFM1}}$ with UFSP2 or 0.2N NaOH and observed complete collapse
only in alkaline conditions confirming that the purified UFC1$^{\text{-O-UFM1}}$ is indeed linked
via an oxyester linkage **(Fig S4E)**. Analysis of complex formation revealed that
UFL1/UFBP1 does not bind to UFC1 or UFM1 on its own but forms a stable complex
with UFC1$^{\text{-O-UFM1}}$ that is of high enough affinity to elute as a complex on SEC (**Fig
4E**). Interestingly, even the smallest of the UFL1/UFBP1 variants was able to bind to

UFC1$^{-O-UFM1}$ (**Fig 4F**). Taken together, the ligase complex containing the N-terminal
segment of UFL1 and the C-terminal region of UFBP1, which we refer to as
UFL1/UFBP1$^{min}$ (denoted as complex I in **Fig 4C**) is sufficient for binding to charged
UFC1 and may represent the minimal ligase domain.

Next, we assayed ligase activity using UFMylation assays monitoring diUFM1
formation. Like full length UFL1, the different C-terminal truncations co-expressed
with UFBP1 did not exhibit any loss of activity. Thus, ligase activity requires the N-
terminal region of UFL1. Similarly, truncation of the N-terminus region (NTR) of
UFBP1 did not affect diUFM1 formation. This suggests that the minimal ligase
domain is sufficient to form diUFM1 chains. Lastly, we checked the impact of
truncations of different regions on UFL1/UFBP1 complex on their ability to modify the
substrates such as Histone H4 and MRE11 (**Fig 4G, S4G**). Intriguingly, while
deletion of regions outside of WH1, WH2 did not alter diUFM1 formation, these
deletions affected substrate modification. Further, deletion of the N-terminal region
(NTR) of UFBP1 significantly impacted substrate UFMylation suggesting roles for
these regions of the ligase complex in substrate recognition **(Fig 4G-H)**. Hence,
using systematic truncations based on sequence and Alphafold predictions, we here
make the surprising discovery that three tandem WH domains of UFL1/UFBP1
mediate E3 ligase activity.

**CDK5RAP3 restricts the E3 ligase activity of UFL1/UFBP1**

Our Y2H screen identified a second hit with low significance, CDK5 regulatory
subunit-associated protein 3 (CDK5RAP3, also known as LZAP), an evolutionarily
conserved 53 kDa protein that lacks any known functional domains or motifs.
Secondary structure analysis predicts CDK5RAP3 to contain a central unstructured
region flanked by α-helical stretches at the N and C terminus (**Fig S5**). To determine
if CDK5RAP3 can interact with UFL1 in the context of the UFL1/UFBP1 complex, we
incubated recombinant full length CDK5RAP3 with UFL1/UFBP1 and analysed
complex formation by SEC. Analysis of UV chromatograms and the fractions
obtained, confirmed that CDK5RAP3 interacts with UFL1/UFBP1 *in vitro* and forms a
stable complex (**Fig 5A**). We further analysed the complex by mass photometry
which confirmed the presence of a stable ternary complex with an experimental
molecular mass of 192 kDa (**Fig 5B**).

Since CDK5RAP3 forms an integral complex with UFL1/UFBP1 and CDK5RAP3 has
recently been suggested to function as a substrate adaptor [52], we wondered if it
could influence E3 ligase activity or substrate UFMylation  . We therefore performed
*in vitro* UFMylation assays and monitored the E3 ligase activity of UFL1/UFBP1 in
the presence of increasing concentrations of CDK5RAP3. Surprisingly, incubation of
CDK5RAP3 with UFL1/UFBP1 impaired E3 ligase activity and UFMylation in a
concentration    dependent    manner    **(Fig   5C)**.    Moreover,    preassembled
UFL1/UFBP1/CDK5RAP3 complexes purified by gel filtration also showed no ligase
activity (**Fig S6A**). The presence of CDK5RAP3 also inhibits UFL1 autoUFMylation
and MRE11 and histone H4 UFMylation (**Fig 5C,** bottom, **Fig S6E**). These
observations imply that CDK5RAP3 binds to inhibit the ligase complex or restricts its
activity towards specific substrates.

To gain insights into how CDK5RAP3 inhibits UFMylation, we tested if CDK5RAP3
inhibits complex formation between UFL1/UFBP1 and UFC1$^{-O-UFM1}$. When analysed
by pull down and analytical SEC, we find that CDK5RAP3 forms a complex together
with UFL1/UFBP1 and UFC1$^{-O-UFM1}$ (**Fig S6B-C**). Since CDK5RAP3 does not affect
UFC1~UFM1 binding, we next tested if CDK5RAP3 binding influences discharge of
UFM1 from UFC1~UFM1 by monitoring transfer of UFM1 onto free Lys. While UFM1
is readily discharged onto Lys in the presence of UFL1/UFBP1, this is completely
blocked in the presence of CDK5RAP3 (**Fig 5D-E**). These observations suggest that
CDK5RAP3 may bind to UFC1~UFM1 to inhibit the ligase complex.

While these *in vitro* experiments clearly demonstrate that CDK5RAP3 blocks UFC1
activation and UFMylation, it raises the question of the role of CDK5RAP3 in
substrate UFMylation, especially UFMylation of ribosomes, the best described
UFMylation substrate to date [5,6]. To address this question, we set up a cell-free
reconstitution of ribosome UFMylation. Strikingly, addition of UFL1/UFBP1 to purified
60S ribosomes leads to robust modification of RPL26 (**Fig 5F**), and in line with
previous observations, both mono- and di-UFMylated RPL26 are observed. When
increasing concentrations of CDK5RAP3 is titrated into this UFMylation reaction, the
di-UFMylation of RPL26 is completely abolished (**Fig 5F**). Importantly, even at the
highest concentration of CDK5RAP3, monoUFMylation of RPL26 is not affected.
Hence, we propose CDK5RAP3 to be a specificity determinant directing activity of

the ligase complex towards substrate Lys by limiting spurious and off-target
UFMylation.

**Regulation of UFC1 activity by its N-terminus**

In addition to its UBC fold, UFC1 has at its N-terminus a highly conserved $\alpha$-helix
($\alpha$0) whose function is unknown (**Fig 6A**). Further, UFC1 lacks the oxyanion hole
stabilizing Asn as part of the highly conserved HPN motif found in E2 enzymes,
which is instead replaced by a TAK motif at this position (**Fig 6B**). To determine if $\alpha$0
has a role in regulating UFMylation, we first compared the activity of UFC1$^{WT}$ and
UFC1 lacking $\alpha$0 (UFC1$^{\Delta N}$) in UFMylation assays containing UFC1 with or without
UFL1/UFBP1. Surprisingly, UFC1$^{\Delta N}$ showed stronger overall UFMylation compared
to UFC1$^{WT}$ in the presence of UFL1/UFBP1 (**Fig 6C**). These results suggest an
inhibitory role for the N-terminal helical extension of UFC1. Next, we compared
discharge of UFM1 from UFC1~UFM1 and UFC1$^{\Delta N}$~UFM1 onto free Lys. In the
absence of E3 ligase, UFC1$^{\Delta N}$ rapidly discharges UFM1 onto Lys (**Fig 6D & E**). In
the presence of UFL1/UFBP1, discharge of UFM1 onto Lys is enhanced when $\alpha$0 of
UFC1 is deleted (**Fig 6D & E**). Taken together with the increase in UFMylation in the
presence of UFL1/UFBP1, these results further cement an inhibitory role for the N-
terminal helix of UFC1.

Since we identified the N-terminal $\alpha$0 of UFC1 to restrain UFMylation, we wondered
if inhibition by CDK5RAP3 was mediated via this helix. Indeed, UFMylation assays
comparing activity of UFC1$^{WT}$ and UFC1$^{\Delta N}$ reveal that the inhibition of E3 ligase
activity by CDK5RAP3 requires the N-terminal helix of UFC1 (**Fig 6D & F, S8**).
Based on these results we propose that CDK5RAP3 may interact with both
UFL1/UFBP1 and $\alpha$0 of UFC1 to clamp the complex in an inhibited state. Together
these analyses reveal a previously unappreciated regulatory role for the N-terminal
helix of UFC1 in modulating UFMylation.

**Discussion**

In this study, we establish a robust *in vitro* reconstitution system using purified active components of the UFM1 enzymatic pathway to reveal the minimal requirements for UFMylation and mechanistic insights into the ligase machinery. Previous reports have provided conflicting views on the roles of UFBP1 and CDK5RAP3, with some suggesting that they are mainly substrates of UFMylation [13,29]. Further, it has been suggested that UFBP1 must be UFMylated first at K267 before it can associate with UFL1 to subsequently support UFL1 ligase activity [12]. Since both UFL1 and UFBP1 complexes are purified from bacteria, our work unequivocally demonstrates that UFL1 and UFBP1 can associate in the absence of any PTMs and the formation of this complex is essential for it to function as an E3 ligase.

It has long been debated as to whether UFL1 is a scaffold/adaptor-type E3 ligase or a Cys-based HECT-like enzyme. Scaffolding type E3 ligases such as RING E3 ligases bind to both E2 and substrate to bring them together for substrate transfer. Since UFL1 can bind to both UFC1 and its supposed substrate UFBP1 at the same time, it was suggested that UFL1 maybe a scaffold-type E3 ligase [3]. More recently, a UFMylation assay that relied on UFL1 present in mammalian cell extracts to which *in vitro* generated biotinylated E2~UFM1 thioesters was added, found that UFMylation occurred even when cell extracts were treated with cysteine-alkylating reagents, further suggesting that UFL1 is a scaffold-type E3 ligase [13]. Our mutational analysis showing that the UFL1/UFBP1 complex lacks a conserved catalytic Cys combined with the ability of the E3 ligase to promote UFC1~UFM1 aminolysis establish UFL1/UFBP1 is a scaffold-type E3 ligase. Further reinforcing this conclusion is the observation that K69-linkage specificity is imparted by the E2 in ligase free di-UFM1 formation assays and this K69-linkage specificity is maintained by the E3 ligase. Indeed, this is a feature observed in RING E3 ligases where linkage-specificity is determined by the E2 enzyme [31,34]. Since UFL1/UFBP1 promotes aminolysis, we speculate that UFL1/UFBP1 binding induces a closed UFC1~UFM1 conformation akin to RING and atypical SUMO E3 ligases [53–57].

UFC1 differs from prototypical E2s since it lacks the catalytic HPN motif, has an exposed catalytic Cys with a lower pKa, lacks the C-terminal α-helix observed in canonical E2s and has an additional α-helix at its N-terminus (α0) [25,40,58]. Further, it is unclear if the canonical "backside" interaction can occur in UFC1 [28,59]. These

unique features of UFC1 and the unconventional features of the ligase complex

makes it likely that the UFM1 machinery employs a unique mechanism to transfer

UFM1 from UFC1 onto substrate. Interestingly, the UBE2E enzymes also have an

intrinsically disordered N-terminal extension that have an inhibitory role restricting Ub

transfer thereby limiting polyUb chain formation [60]. Drawing analogy to RING E3

ligases whose binding induces a closed E2~Ub conformation, it remains to be seen

whether E3 binding induces a closed UFM1~UFM1 conformation and whether $\alpha 0$

impedes this process. Nevertheless, our data seem to suggest that $\alpha 0$ has a

regulatory role potentially preventing UFM1 discharge in the absence of the E3

ligase.

The recent advances in structure prediction enabled us to narrow the regions of

UFL1/UFBP1 essential for ligase activity to a tandem repeat of three WH domains -

a WH domain each from UFL1 and UFBP1 and the composite WH domain formed

by the two proteins [49]. As this region can bind to UFC1~UFM1 and activate the E2

for UFM1 discharge, we define this to be the minimal catalytic domain. Further

structural studies will be required to reveal why three WH domains are required to

form a functional ligase complex and the identification of linchpin residues in

UFL1/UFBP1 that play similar roles to RING E3 ligases to activate the E2 for

aminolysis [54,55]. Intriguingly, the anaphase promoting complex (APC) subunit APC2

contains a WHB domain that binds the "backside" of UCBH10/UBE2C, located at a

face opposite from the catalytic site and is an allosteric site in several E2s [61,62]. This

backside binding of the WHB domain is specific towards UBE2C and is important for

transfer of Ub onto specific substrates [62]. Our observations also raise the question of

the function of the other WH/PCI domains in UFL1/UFBP1 to ligase function. One

possibility is that the WH domains may mediate substrate recognition, binding to

UFM1 or serve to either regulate recruitment or processivity of chain formation.

Given that the WH domain of Cockayne syndrome group B (CSB) can bind to

ubiquitin, it is tempting to speculate that one of the WH domains of UFL1/UFBP1 can

bind to UFM1 [63]. WH domains commonly recognize nucleic acids and are typically

found in transcription factors or as protein interaction domains [64]. To our knowledge,

this is the first demonstration of catalytic activity mediated by WH domains [65],

leading us to propose moonlighting functions for other WH domain containing

proteins. It also raises the possibility that other WH domains could have E3 ligase

activity for UFMylation or other UBLs. Since UFL1 has been proposed to have

nuclear functions in telomere maintenance and DNA damage response [10,11,66], one

possibility is that the WH domains may mediate DNA binding. A further possibility is

the use of the WH domains of UFL1/UFBP1 to recognize rRNA or translating mRNA

during ribosome UFMylation.

Our *in vitro* studies show that formation of UFM1 chains and autoUFMylation is

inhibited by CDK5RAP3 (**Fig 2C**). Indeed, previous studies have observed an altered

pattern of UFMylation in the absence of CDK5RAP3 leading to the suggestion that it

may be a substrate adaptor [52]. Interestingly, CDK5RAP3 was suggested to function

as a sensor for ER stress to induce autophagic degradation of aberrant proteins

formed as a result of ribosomal stalling [67]. Several multidomain and multiprotein E3

ligases such as PARKIN and Cullin Ring Ligases (CRLs) are inhibited and their

activation is a carefully orchestrated multistep process [35,68]. Based on our

observations, we propose that ligase complexes containing UFL1, UFBP1 and

CDK5RAP3 represent an autoinhibited state. The surprising finding that ribosome

UFMylation is not abolished in the presence of CDK5RAP3 but rather restricted to

monoUFMylation leads us to suggest that CDK5RAP3 regulates ligase activity in the

following manner: (i) in the absence of substrate, binding of CDK5RAP3 inhibits E3

ligase activity; (ii) this autoinhibition is relieved when substrates such as the 60S

ribosome are encountered. This release from inhibition may involve conformational

changes induced upon recognition of structural features on the ribosome by UFL1,

UFBP1 or CDK5RAP3 to mediate substrate UFMylation. Such multi-layered

regulation possibly prevents spurious UFMylation and ensures ribosome UFMylation

only in the right context. While previous studies have suggested roles for UFMylation

for UFL1-UFBP1 interaction, and UFMylation and phosphorylation to enhance

UFMylation [10,12,19,29]. Our minimal reconstitution clearly demonstrates that the ligase

complex assembles in the absence of any PTM. While we cannot rule out a role for

phosphorylation in enhancing ligase activity, the rapid UFMylation of ribosomes by

the reconstituted ligase suggests that it may be unlikely.

Our rebuilding approach provides insights into the assembly, minimal requirements,

and mechanism of the UFL1/UFBP1/CDK5RAP3 ligase and reveals principles of

protein UFMylation. The *in vitro* reconstitution system we describe here using

purified components lays the foundation for future biochemical and structural studies

to understand the molecular mechanism of this unusual E3 ligase complex and

UFMylation. Further, the cell-free UFMylation system can be applied to understand

the logic of ribosome UFMylation and its relationship to ribosome quality control

pathways.

**Acknowledgements**

We thank members of the Kulathu lab for critical reading of the manuscript and helpful comments. We thank the Zeqiraj lab (Univ of Leeds) and Prof Hofmann (Univ of Köln) for helpful discussions and advice. We acknowledge MRC Reagents and Services for providing valuable reagents, especially Simone Weidlich and Rachel Toth for cloning the constructs used in this study. We thank Mr. Joby Varghese for help with Mass spectrometry data acquisition and preliminary analysis. We thank Christopher Lapointe (Puglisi lab, Stanford) for valuable reagents and discussion. We also thank Juraj Ahel (IMP Vienna) for his help with generating composite foldindex profile of UFL1. This research was supported by ERC Starting grant (RELYUBL, 677623), the Lister Institute of Preventive Medicine, BBSRC (BB/T008172/1) and MRC grant MC_UU_00018/3 to YK; Grant R01GM074874 from the National Institute of General Medical Sciences and a generous gift from the Becker Family Foundation to R.R.K; NIH Epilepsy Training Program T32 NS007280 to PAD.

**Author contributions**

YK and JJP conceived the study. JJP designed, performed and analysed all the experiments. HM designed, expressed and purified UFL1/UFBP1 constructs based on Alphafold. PADR performed *in vitro* ribosome UFMylation assays. DM and SPM contributed reagents and RS performed SEC MALS analysis. YK analysed data, supervised the study and secured funding. JJP and YK wrote the manuscript with input from all the authors.

**Declaration of interests**

The authors declare no competing interests.

**Supplemental information**

Supplemental information includes Figures S1-S8 and Table S1.

<sup>515</sup>

**Materials and Methods**

**Plasmids, Cloning and mutagenesis**

The details of cDNA constructs used in this study is given in Table S1. Cloning of most of the constructs was performed by DSTT (University of Dundee). Briefly, mutagenesis was carried out using Q5 site-directed mutagenesis (NEB) with KOD polymerase (Novagen) according to manufacturer's protocol. Following mutagenesis, cDNA constructs were amplified using *E. coli* DH5α and purified using Qiagen mini prep kit (Qiagen). All cDNA constructs were verified by DNA sequencing and services, University of Dundee using DYEnamic ET terminator chemistry (Amersham Biosciences) on Applied Biosystems automated DNA sequencers.

**Recombinant protein expression and purification**

Recombinant His$_6$-3C-UBA5, His$_6$-3C-UFM1$^{WT}$, His$_6$-3C-UFM1$^{Konly}$, His$_6$-3C-UFM1$^{KtoR}$, His$_6$-3C-Cys-UFM1, His$_6$-3C-UFC1$^{WT}$, His6-3C-UFC1$^{\Delta N}$ and His$_6$-3C-MRE11 were expressed in *E. coli* BL21 and purified using Ni$^{2+}$-NTA affinity chromatography as the first step. Briefly, *E. coli* BL21 cultures expressing His$_6$-tagged proteins were grown in 2xTY medium at 37ºC until OD$_{600}$ reached 0.6-0.8. Final concentration of 0.3 mM IPTG was added and the cultures were incubated at 18 °C for 16h. Cells were harvested, resuspended, and lysed in appropriate lysis buffer containing 25 mM Tris pH 8, 300 mM NaCl, 10% Glycerol, 2 mM DTT, 1 mM Benzamidine, 1 mM AEBSF, 1x protease inhibitor cocktail (Roche) by ultrasonication. Lysed cells were then clarified by centrifugation at 30000g for 30 mins at 4ºC. The clarified lysate was then incubated with pre-equilibrated Ni$^{2+}$-NTA Agarose beads (GE Life Sciences) for 2h in appropriate binding buffer containing 25 mM Tris pH 8, 300 mM NaCl, 10% Glycerol,10 mM Imidazole. The beads were then washed extensively using wash buffer containing 25 mM Tris pH 8, 300 mM NaCl, 10% Glycerol, 2 mM DTT and 20 mM Imidazole. Finally, the bound protein was eluted using elution buffer containing 25 mM Tris pH 8, 300 mM NaCl, 10% Glycerol, 2 mM DTT and 300 mM Imidazole. Wherever necessary, His6-tags were cleaved off by incubating tagged proteins with PreScission protease at 4ºC overnight. A final size exclusion chromatography step was performed using Superdex 75 HiLoad$^{TM}$ 16/60 Superdex$^{TM}$ 75pg and HiLoad$^{TM}$ 16/60 Superdex$^{TM}$ 200 pg columns (GE Healthcare Life Sciences) with the mobile phase buffer as 25 mM Tris pH 8.0, 150 mM NaCl, 10%

548 Glycerol, 2 mM DTT. The purified proteins were then concentrated using Amicon$^{TM}$ Ultra

549 15 concentrators (MERCK Millipore) and stored in -80°C.

**Purification of GST-tagged proteins**

GST-TEV-TRIP4 was expressed in *E. coli* BL21(DE3) strain as described above.

Cells were harvested and lysed in lysis buffer containing 25 mM Tris pH 7.5, 300 mM

NaCl, 10% Glycerol and 2 mM DTT using ultrasonication. Pre-equilibrated

Glutathione 4B-sepharose (DSTT, University of Dundee) beads were incubated with

clarified lysate for 2 h. The beads were then washed with high salt buffer containing

25 mM Tris pH 7.5, 500 mM NaCl, 10% glycerol and 2 mM DTT. Further the beads

were washed with low salt buffer containing 25 mM Tris pH 8, 150 mM NaCl, 10%

glycerol and 2 mM DTT. The protein was then cleaved off the tag by incubation with

Precision protease at 4°C overnight. A second ion exchange chromatography step

was carried out using Resource Q (6ml) (GE Healthcare Life Sciences) column with

low salt buffer as 25mM Tris pH pH7.5, 150 mM NaCl, 10% glycerol,2 mM DTT and

high salt buffer as 25mM Tris pH 7.5, 500mM NaCl, 10% glycerol and 2 mM DTT. A

final size exclusion chromatography step was performed using HiLoad$^{TM}$ 16/60

Superdex$^{TM}$ 75pg (GE Healthcare Life Sciences) and the protein was buffer

exchanged in buffer 25 mM Tris pH 7.5, 150 mM NaCl, 10% glycerol and 2 mM DTT.

The purified proteins were then concentrated and stored in -80°C.

**Co-expression and purification of UFL1-UFBP1 complex**

His$_6$-TEV-UFL1 and StrepII-C3-UFBP1 were cloned in a pET Duet1 construct and

expressed in *E. coli* BL21 codon plus (RIPL) (Agilent) cells. Bacterial cultures were

grown in 2xTY medium at 37°C until OD$_{600}$ reached 0.6. Final concentration of 0.3

mM IPTG was added to induce the expression and the cultures were incubated at

18°C for 16 h. Cells were harvested, resuspended in buffer containing 25 mM Tris

pH 8.0, 300 mM NaCl, 2 mM DTT, 1mM Benzamidine, 1mM AEBSF, protease

inhibitor cocktails (Roche). Cells were lysed by High pressure homogenization using

French press (Avestin Emulsiflux C3, Wolflabs). The lysate was then clarified by

Ultracentrifugation at 30000g for 30 min at 4°C. Affinity purification was carried out in

two steps. In the first step, the clarified lysate (25 mM Tris pH 8.0, 300 mM NaCl, 20

mM Imidazole, 2 mM DTT) was passed through HisTrap$^{TM}$ FF (GE Healthcare Life

Sciences) column. After binding, the column was washed with 10 column volumes

(CV) of binding buffer containing 25 mM Tris pH 8.0, 300 mM NaCl, 20 mM

Imidazole, 2 mM DTT.   After washing, bound proteins were eluted using buffer containing 25 mM Tris pH 8.0, 300 mM NaCl, 300 mM Imidazole, 2 mM DTT by applying a concentration gradient of Imidazole. The eluted proteins were then passed through StrepTrap[TM] (GE Healthcare Life Sciences) column pre-equilibrated with buffer containing 25 mM Tris pH 8.0, 300 mM NaCl, 2 mM DTT. After 2 CV of washing with binding buffer, the proteins were eluted using 25 mM Tris pH 8.0, 300 mM NaCl, 2 mM DTT and 2.5 mM Desthiobiotin.    Finally, purified proteins were passed through HiLoad[TM] 16/60 Superdex[TM] 200pg (GE Healthcare Life Sciences). The purified protein was then stored in -80°C in buffer containing 25 mM Tris pH 8.0, 300 mM NaCl, 5% Glycerol, 2 mM DTT until further use.

**Preparation and labelling of IRDye800CW-UFM1**

Cys-UFM1[1-83] which contains a Cys residue upstream of M1 was purified and exchanged into a fresh buffer containing 50 mM HEPES, pH 7.5, 150 mM NaCl and 0.5 mM TCEP using CentriPure P10 columns (EMP Biotech). Protein was then mixed with IRDye® 800CW Maleimide (LI-COR®) at 5:1 molar ratio and incubated for 2 h at RT. Unreacted dye was removed by three-step buffer exchange process using CentriPure P10 columns (EMP Biotech) (once) and PD-10 Sephadex G-25 column (twice) in a sequential manner according to manufacturer's instruction. Finally, to remove residual unreacted dye material, the labelled protein was dialyzed into buffer containing 25 mM Tris pH 8, 150 mM NaCl, 2 mM DTT using dialysis membrane (Thermo Scientific) (MWCO 3000) overnight at 4°C and stored at -80°C until further use.

**Biochemical assays**

**Lysine discharge assays**

To analyse the intrinsic reactivity of UFC1, a substrate-independent single turnover discharge assay was employed. Firstly, UFC1 was charged with UFM1[WT] or labelled UFM1 by incubating 0.5 µM UBA5, 10 µM UFC1 and 20 µM UFM1 in a buffer containing 50 mM HEPES pH 7.5, 50 mM NaCl, 0.5 mM DTT and 10 mM $MgCl_2$. The charging reaction was initiated by addition of 10 mM ATP and incubation of the reaction mix at 37°C for 20-30 min. To stop further charging of UFM1 onto UFC1, the reaction was quenched by addition of 50 mM EDTA and subsequent incubation at RT for 10 mins. To check for discharge, the quenched mixture was mixed with 50 mM lysine (pH 8.0) or other amino acids namely Serine, Threonine, Arginine,

Cysteine and incubated at 37°C. The reaction was stopped at each of the time points

by mixing the reaction mix with SDS-PAGE loading dye without any reducing agent

and run on a 4-12% SDS-PAGE under non-reducing conditions followed by

Coomassie staining or visualization using LI-COR®.

In reactions involving the UBCH5C and UBCH7, 0.5 µM UBE1 was incubated with

10 µM UBCH5C/UBCH7 and 20 µM Ub in reaction buffer containing 50 mM HEPES

pH 7.5, 50 mM NaCl,10 mM ATP and 10 mM $MgCl_2$ at 37°C for 20 mins. The

reaction was quenched, and discharge was analysed as described above.

In discharge assays involving UFL1/UFBP1 and CDK5RAP3, the quenched reaction

mix was added to a cocktail containing 1 µM UFL1/UFBP1 full length or truncated

constructs and/or 1 µM or 2 µM of CDK5RAP3. Discharge was initiated by addition

of  50 mM lysine (pH 8.0) in buffer containing 50 mM HEPES pH 7.5, 50 mM NaCl

and 0.5 mM DTT and incubated at 37°C for indicated time duration. The reaction

was stopped and analysed as described above.

*In vitro* **UFMylation assays**

To check for UFMylation *in vitro,* 0.25 µM UBA5, 5 µM UFC1, 1 µM UFL1 and 20-30

µM UFM1 were incubated in reaction buffer containing 50 mM HEPES 7.5, 10 mM

$MgCl_2$ and 5 mM ATP for 1h at 37ºC. The reaction was stopped by the addition of 3x

SDS loading buffer containing a reducing agent. The reaction products were then

separated on a 4-12% SDS-PAGE gel under reducing conditions and analysed by

immunoblotting using indicated antibodies. In UFMylation assays involving

substrates, around 1 µM-2 µM of recombinant substrates were incubated with 0.25

µM UBA5, 5 µM UFC1, 1µM UFL1 and 30 µM of UFM1 in buffer containing 50 mM

HEPES 7.5,10 mM $MgCl_2$ and 5 mM ATP for 1h at 37°C. Following incubation, the

reaction was stopped and analyzed using SDS-PAGE or Immunoblotting as

described above.

**Pull down assays**

To analyse the interaction of the E3 ligase with the E2, pulldown assays were

performed.  Approximately 10 µM of untagged UFC1 or UFC1[-O-UFM1] was incubated

with 5 µM of full length  UFL1/UFBP1 in binding buffer containing 25 mM Tris pH 8.0,

300 mM NaCl, 2 mM DTT for 1 h at 4°C. Following incubation, 30 µl of pre-

equilibrated Streptavidin Sepharose beads (IBA Life Sciences) (50% slurry) was
added and further incubated for about 1 h at 4°C. Following binding, centrifuge tubes
containing the reaction mix was spun down at 500g for 3 mins and the supernatant
was discarded to remove unbound proteins. To further remove weakly bound
proteins, the beads were washed thrice with 1 ml of ice-cold binding buffer. Finally,
the bound proteins were eluted by addition of binding buffer containing 2.5 mM
Desthiobiotin (pH 8.0) and incubation at 4°C for 30 mins. The eluates were then
analysed on 4-12% SDS PAGE under reducing conditions followed by coomassie
staining.

**Analytical gel filtration chromatography analysis**
For analysing the interaction between UFL1/UFBP1 and CDK5RAP3, around 30 µgs
of UFL1/UFBP1 and 15 µgs of CDK5RAP3 were mixed and incubated for 1 h on ice.
Then, around 50 µls of sample was loaded on Superdex$^{TM}$ 200 Increase 3.2/300
column (GE Healthcare Lifesciences) pre-equilibrated with buffer containing 25 mM
Tris pH 8.0, 300 mM NaCl, 2 mM DTT. The fractions corresponding to each peak
were collected and further analysed on a 4-12% SDS PAGE under reducing
conditions followed by coomassie staining.

To check if UFL1/ UFBP1 could be reconstituted *in trans*, around 15 µgs of purified
His$_6$-UFL1 and StrepII-UFBP1 were mixed and incubated on ice for 2h. Around 50
µls of sample was then loaded on Superdex$^{TM}$ 200 Increase 3.2/300 column (GE
Healthcare Lifesciences) column pre-equilibrated with buffer containing 25 mM Tris
pH 8.0, 300 mM NaCl, 2 mM DTT and the chromatograms were analysed.

**Preparation of UFC1$^{-O-UFM1}$**
UFC1$^{-O-UFM1}$ was prepared by incubating 40 µM UFC1$^{C116S}$ with 40 µM UFM1, 1 µM
UBA5 in buffer containing 50 mM Tris pH 8.8, 10 mM MgCl$_2$, 5 mM ATP overnight at
25°C. The reaction was incubated briefly with 20 mM DTT to remove any non-
specific di-sulfide linked adducts formed and passed through HiLoad$^{TM}$ 16/60
Superdex$^{TM}$ 75 pg with buffer containing 25 mM Tris 8.0, 150 mM NaCl, 5%
Glycerol, 2 mM DTT. The fractions corresponding to UFC1$^{-O-UFM1}$ were collected and
analysed on 4-12% SDS PAGE gel. Finally, the pure fractions containing UFC1$^{-O-}$
$^{UFM1}$ were pooled, concentrated and stored in -80°C until further use.

**Alphafold Predictions**

The structure of UFL1/UFBP1 was predicted using the ColabFold Google Colab notebook "AlphaFold2 using MMseqs2" [49] (https://github.com/sokrypton/ColabFold). The predicted model with the highest IDDT score is shown in the main figure and the PAE scores for different models ranking 1 to 5 are shown in **Fig S4A**.

**DALI Analysis**

DALI server was used to perform structural similarity analysis [69]. WH1 domain of UFL1(52-115) was used as the query model and searched against PDB25 database. The top ten models in the order of the Z-score is shown in **Fig S4B**.

**Multiple sequence alignments**

The sequences of target proteins and their homologs were manually downloaded from UNIPROT database in fasta format. Multiple sequence alignment was performed using Jalview version 2.11.1.7 programme (MAFFT algorithm module using L-INS-I) [70,71].

**Mass Photometry Data Acquisition and Analysis**

Protein samples were prepared in a solution containing 25 mM Tris pH 8.0, 300 mM NaCl, 2 mM DTT and stored on ice before loading onto the Refeyn One$^{MP}$ instrument (Refeyn). Typically, a set of protein standards (NativeMark™ Unstained Protein Standard, Invitrogen) are used for calibration and to generate a standard curve. Following calibration, approximately 10 μl of diluted protein sample in the concentration range of 5–25 nM was introduced into the flow-chamber and movies of 60s duration were recorded. Data acquisition was performed using Acquire MP (Refeyn Ltd, v1.1.3) and analysed using Discover MP software. Data shown here is representation of at least 3 independent acquisitions ($n \geq 3$).

**SEC MALS data acquisition and analysis**

Size exclusion chromatography and multi angle light scattering (SEC–MALS) experiments were performed on a Dionex Ultimate 3000 HPLC system with an inline Wyatt miniDAWN TREOS MALS detector and Optilab T-rEX refractive index detector. In addition, the elution profile of the protein was monitored with UV 280 attached to the Dionex system. For size exclusion chromatography, Superdex 200 10/300 gl column (GE Healthcare LifeSciences) was used. 50 μl of the purified UFL1/UFBP1 stored in buffer containing 25 mM Tris pH 8.0, 300 mM NaCl, 2 mM

711 DTT was loaded into the SEC column with Dionex auto loader at a concentration of 4

712 mg/ml and a flow rate of 0.3ml/min was maintained throughout the experiment. Molar

713 masses spanning elution peaks were calculated using ASTRA v6.0.0.108 (Wyatt).

**Ribosome UFMylation assays**

Purified 60S ribosomes were a generous gift from the Puglisi lab, and purified as

described previously [72,73]. *In vitro* UFMylation of ribosomes was performed at 30 ºC

with 1 µM UFC1, 0.5 µM UBA5, 0.5 µM UFM1, 0 nM or 75 nM UFL1/UFBP1, 50 nM

purified 60S ribosomes, and increasing concentrations of CDK5RAP3 (0, 38, 75, 150

or 375 nM) in a reaction buffer of 25 mM HEPES pH 7.5, and 100 mM NaCl. The

reaction was quenched in 4 x SDS-Load buffer. Western blots show 10 min reaction

time; immunoblots for RPL26 (Abcam,  59567) and UFL1 (Bethyl, A303-455M) were

run on the same gel, which was cut and probed for these proteins separately.

Immunoblots for CDK5RAP3 Bethyl, A300-871A) and UFM1 (Abcam, Ab109307)

were run on separate gels.

**LC-MS/MS sample preparation, data acquisition and analysis**

First *in vitro* UFMylation reaction was performed to generate UFMylated products

including free UFM1 chains. Then, the reaction products were run on a 4-12% SDS

PAGE gel to separate the products based on electrophoretic mobility. Next, the

bands corresponding to di-UFM1 chains were excised and in-gel digestion was

performed [74]. Digested peptides were analysed by LC-MS/MS on an Exploris 480

coupled to an Ultimate 3000 nanoLC system (Thermo Fisher Scientific) for figure 2D

and on an Exploris 240  (Thermo Fisher Scientific) coupled to an Evosep One

(Evosep) for figure 4D. For the analysis performed on the Ultimate 3000, samples

were loaded on a 100 µm x 2 cm trap column (Thermo Fisher Scientific #164564-

CMD) and analysed on a 75 µm x 50 cm analytical column (Thermo Fisher Scientific

#ES903) using a gradient from 3% to 35% Buffer B (80% LCMS grade acetonitrile,

0.08% formic acid in water) over 53 minutes. The columns were then washed with

95% Buffer B for 2 minutes prior equilibration in 97% Buffer A (0.1% formic acid in

LCMS grade water). For the analysis performed on the Evosep One, samples were

loaded onto the Evotips as per manufacturer recommendations and analysed using

the 30 SPD Method.Peptides were then analysed in on either the Exploris 240 or

480 using data dependant with a MS1 resolution of 60000, AGC target of 300% and

maximum injection time of 25 or 28 ms. Peptide were then fragmented using TOP 2

s method, MS2 resolution of 15000, NCE of 30 or 32%, AGC of 100% and maximum

injection time of 100 ms. Peptide identification was performed in Mascot using a

restricted and frequently updated database containing ~2000 protein sequences of

interest (MRC db). Carbamidomethylation (C) was set at fixed modification and

Oxidation (M), Dioxidation (M) and the addition of the dipeptide Glycine-Valine (K)

were set as variable modifications. Peptides were searchd using an MS1 tolerance

of 10 ppm and MS2 tolerance of 0.06 Da and a maximum of 2 missed cleavages

were allowed. Only hits indentified with a FDR (False Discovery Rate) <1% were

selected and further analysed in Scaffold viewer V5. For semi-quantitative analysis

of VG modification on individual sites, Total Ion chromatogram (TIC) values obtained

from the LC/MS and represented graphically.

**References:**

1.    Van Der Veen, A. G. & Ploegh, H. L. Ubiquitin-like proteins. *Annu. Rev. Biochem.* **81**,
323–357 (2012).
2.    Sasakawa, H. *et al.* Solution structure and dynamics of Ufm1, a ubiquitin-fold modifier
1. *Biochem. Biophys. Res. Commun.* **343**, (2006).
3.    Komatsu, M. *et al.* A novel protein-conjugating system for Ufm1, a ubiquitin-fold
modifier. *EMBO J.* **23**, 1977–1986 (2004).
4.    Cappadocia, L. & Lima, C. D. Ubiquitin-like Protein Conjugation: Structures,
Chemistry, and Mechanism. *Chemical Reviews* vol. 118 889–918 (2018).
5.    Walczak, C. P. *et al.* Ribosomal protein RPL26 is the principal target of UFMylation.
*Proc. Natl. Acad. Sci. U. S. A.* **116**, 1299–1308 (2019).
6.    Wang, L. *et al.* UFMylation of RPL26 links translocation-associated quality control to
endoplasmic reticulum protein homeostasis. *Cell Res.* **30**, 5–20 (2020).
7.    Liang, J. R. *et al.* A Genome-wide ER-phagy Screen Highlights Key Roles of
Mitochondrial Metabolism and ER-Resident UFMylation. *Cell* **180**, 1160-1177.e20
(2020).
8.    Wang, Z. *et al.* MRE11 UFMylation promotes ATM activation. *Nucleic Acids Res.* **47**,
(2019).
9.    Liu, J. *et al.* UFMylation maintains tumour suppressor p53 stability by antagonizing its
ubiquitination. *Nat. Cell Biol.* **22**, 1056–1063 (2020).
10.   Qin, B. *et al.* UFL1 promotes histone H4 ufmylation and ATM activation. *Nat.*
*Commun.* **10**, (2019).
11.   Lee, L. *et al.* UFMylation of MRE11 is essential for maintenance of telomere length
and hematopoietic stem cell survival. (2019) doi:10.1101/846477.
12.   Yoo, H. M. *et al.* Modification of asc1 by ufm1 is crucial for erα transactivation and
breast cancer development. *Mol. Cell* **56**, 261–274 (2014).
13.   Gak, I. A. *et al.* UFMylation regulates translational homeostasis and cell cycle
progression. *bioRxiv* (2020) doi:10.1101/2020.02.03.931196.
14.   Xie, Z., Fang, Z. & Pan, Z. Ufl1/RCAD, a Ufm1 E3 ligase, has an intricate connection
with ER stress. *International Journal of Biological Macromolecules* vol. 135 760–767
(2019).
15.   Liu, J. *et al.* A critical role of DDRGK1 in endoplasmic reticulum homoeostasis via
regulation of IRE1α stability. *Nat. Commun.* **8**, (2017).
16.   Duan, R. *et al.* UBA5 mutations cause a new form of autosomal recessive cerebellar
ataxia. *PLoS One* **11**, 1–12 (2016).
17.   Nahorski, M. S. *et al.* Biallelic UFM1 and UFC1 mutations expand the essential role of
ufmylation in brain development. *Brain* **141**, 1934–1945 (2018).
18.   Tatsumi, K. *et al.* The Ufm1-activating enzyme Uba5 is indispensable for erythroid
differentiation in mice. *Nat. Commun.* **2**, (2011).
19.   Lemaire, K. *et al.* Ubiquitin fold modifier 1 (UFM1) and its target UFBP1 protect
pancreatic beta cells from ER stress-induced apoptosis. *PLoS One* **6**, (2011).
20.   Cai, Y. *et al.* UFBP1, a Key Component of the Ufm1 Conjugation System, Is Essential
for Ufmylation-Mediated Regulation of Erythroid Development. *PLoS Genet.* **11**,
(2015).
21.   Cai, Y., Singh, N. & Li, H. Essential role of Ufm1 conjugation in the hematopoietic
system. *Experimental Hematology* vol. 44 (2016).
22.   Sung, H. K. *et al.* Two novel ubiquitin-fold modifier 1 (Ufm1)-specific proteases,
UfSP1 and UfSP2. *J. Biol. Chem.* **282**, 5256–5262 (2007).
23.   Oweis, W. *et al.* Trans-Binding Mechanism of Ubiquitin-like Protein Activation
Revealed by a UBA5-UFM1 Complex. *Cell Rep.* **16**, 3113–3120 (2016).

24.   Banerjee, S., Kumar, M. & Wiener, R. Decrypting UFMylation: How proteins are
modified with UFM1. *Biomolecules* **10**, 1–14 (2020).
25.   Mizushima, T. *et al.* Crystal structure of Ufc1, the Ufm1-conjugating enzyme.
*Biochem. Biophys. Res. Commun.* **362**, 1079–1084 (2007).
26.   Liu, G. *et al.* NMR and X-RAY structures of human E2-like ubiquitin-fold modifier
conjugating enzyme 1 (UFC1) reveal structural and functional conservation in the
metazoan UFM1-UBA5-UFC1 ubiquination pathway. *J. Struct. Funct. Genomics* **10**,
127–136 (2009).
27.   Stewart, M. D., Ritterhoff, T., Klevit, R. E. & Brzovic, P. S. E2 enzymes: More than just
middle men. *Cell Res.* **26**, 423–440 (2016).
28.   Brzovic, P. S., Lissounov, A., Christensen, D. E., Hoyt, D. W. & Klevit, R. E. A
UbcH5/ubiquitin noncovalent complex is required for processive BRCA1-directed
ubiquitination. *Mol. Cell* **21**, 873–880 (2006).
29.   Tatsumi, K. *et al.* A novel type of E3 ligase for the Ufm1 conjugation system. *J. Biol.*
*Chem.* **285**, 5417–5427 (2010).
30.   Cai, Y. *et al.* Indispensable role of the Ubiquitin-fold modifier 1-specific E3 ligase in
maintaining intestinal homeostasis and controlling gut inflammation. *Cell Discov.* **5**,
(2019).
31.   Deol, K. K., Lorenz, S. & Strieter, E. R. Enzymatic Logic of Ubiquitin Chain Assembly.
*Front. Physiol.* **10**, 1–14 (2019).
32.   Pichler, A., Fatouros, C., Lee, H. & Eisenhardt, N. SUMO conjugation - A mechanistic
view. *Biomolecular Concepts* vol. 8 13–36 (2017).
33.   Lorenz, S. Structural mechanisms of HECT-type ubiquitin ligases. *Biol. Chem.* **399**,
127–145 (2018).
34.   Deshaies, R. J. & Joazeiro, C. A. P. RING domain E3 ubiquitin ligases. *Annu. Rev.*
*Biochem.* **78**, 399–434 (2009).
35.   Walden, H. & Rittinger, K. RBR ligase–mediated ubiquitin transfer: a tale with many
twists and turns. *Nat. Struct. Mol. Biol.* **25**, 440–445 (2018).
36.   Pao, K. *et al.* Ligase With Esterification Activity. *Nature* (2018).
37.   Wu, J., Lei, G., Mei, M., Tang, Y. & Li, H. A novel C53/LZAP-interacting protein
regulates stability of C53/LZAP and DDRGK domain-containing protein 1 (DDRGK1)
and modulates NF-κB signaling. *J. Biol. Chem.* **285**, 15126–15136 (2010).
38.   Witting, K. F. & Mulder, M. P. C. Highly specialized ubiquitin-like modifications:
Shedding light into the ufm1 enigma. *Biomolecules* vol. 11 (2021).
39.   Gerakis, Y., Quintero, M., Li, H. & Hetz, C. The UFMylation System in Proteostasis
and Beyond. *Trends in Cell Biology* vol. 29 (2019).
40.   Kumar, M. *et al.* Structural basis for UFM1 transfer from UBA5 to UFC1. *Nat.*
*Commun.* **12**, (2021).
41.   Padala, P. *et al.* Novel insights into the interaction of UBA5 with UFM1 via a UFM1-
interacting sequence. *Sci. Rep.* **7**, 1–12 (2017).
42.   Soudah, N. *et al.* An N-Terminal Extension to UBA5 Adenylation Domain Boosts
UFM1 Activation: Isoform-Specific Differences in Ubiquitin-like Protein Activation. *J.*
*Mol. Biol.* **431**, 463–478 (2019).
43.   Wenzel, D. M., Lissounov, A., Brzovic, P. S. & Klevit, R. E. UBCH7 reactivity profile
reveals parkin and HHARI to be RING/HECT hybrids. *Nature* **474**, 105–108 (2011).
44.   Kihm, A. J. *et al.* An abundant erythroid protein that stabilizes free α-haemoglobin.
*Nature* **417**, 758–763 (2002).
45.   Warner, J. R. In the absence of ribosomal RNA synthesis, the ribosomal proteins of
HeLa Cells are synthesized normally and degraded rapidly. *J. Mol. Biol.* **115**, 315–333
(1977).
46.   Sonn-Segev, A. *et al.* Quantifying the heterogeneity of macromolecular machines by

mass photometry. *Nat. Commun.* **11**, 1–10 (2020).

47.  Deng, L. *et al. Activation of the IB Kinase Complex by TRAF6 Requires a Dimeric Ubiquitin-Conjugating Enzyme Complex and a Unique Polyubiquitin Chain through covalent modification by Ub. Cell* vol. 103 (2000).

48.  Branigan, E., Plechanovová, A., Jaffray, E. G., Naismith, J. H. & Hay, R. T. Structural basis for the RING-catalyzed synthesis of K63-linked ubiquitin chains. *Nat. Struct. Mol. Biol.* **22**, 597–602 (2015).

49.  Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

50.  Stewart, M. Structural biology of the PCI-protein fold. *Bioarchitecture* vol. 2 118–123 (2012).

51.  Scheel, H. & Hofmann, K. Prediction of a common structural scaffold for proteasome lid, COP9-signalosome and eIF3 complexes. *BMC Bioinformatics* **6**, (2005).

52.  Yang, R. *et al.* CDK5RAP3, a UFL1 substrate adaptor, is crucial for liver development. *Dev.* **146**, (2019).

53.  Pichler, A., Knipscheer, P., Saitoh, H., Sixma, T. K. & Melchior, F. The RanBP2 SUMO E3 ligase is neither HECT- nor RING-type. *Nat. Struct. Mol. Biol.* **11**, 984–991 (2004).

54.  Plechanovov, A., Jaffray, E. G., Tatham, M. H., Naismith, J. H. & Hay, R. T. Structure of a RING E3 ligase and ubiquitin-loaded E2 primed for catalysis. *Nature* **489**, 115–120 (2012).

55.  Pruneda, J. N. *et al.* Structure of an E3:E2~Ub Complex Reveals an Allosteric Mechanism Shared among RING/U-box Ligases. *Mol. Cell* **47**, 933–942 (2012).

56.  Reverter, D. & Lima, C. D. Insights into E3 ligase activity revealed by a SUMO–RanGAP1–Ubc9–Nup358 complex. *Nature* **435**, 687–692 (2005).

57.  Cappadocia, L., Pichler, A. & Lima, C. D. Structural basis for catalytic activation by the human ZNF451 SUMO E3 ligase. *Nat. Struct. Mol. Biol.* **22**, 968–975 (2015).

58.  Gundogdu, M. & Walden, H. Structural basis of generic versus specific E2–RING E3 interactions in protein ubiquitination. *Protein Sci.* **28**, 1758–1770 (2019).

59.  Middleton, A. J., Wright, J. D. & Day, C. L. Regulation of E2s: A Role for Additional Ubiquitin Binding Sites? *J. Mol. Biol.* **429**, 3430–3440 (2017).

60.  Schumacher, F.-R., Wilson, G. & Day, C. L. The N-Terminal Extension of UBE2E Ubiquitin-Conjugating Enzymes Limits Chain Assembly. *J. Mol. Biol.* **425**, 4099–4111 (2013).

61.  Brzovic, P. S. *et al.* Binding and recognition in the assembly of an active BRCA1/BARD1 ubiquitin-ligase complex. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 5646–5651 (2003).

62.  Brown, N. G. *et al.* RING E3 mechanism for ubiquitin ligation to a disordered substrate visualized for human anaphase-promoting complex. *Proc. Natl. Acad. Sci.* **112**, 5272–5279 (2015).

63.  Takahashi, T. S. *et al.* Structural basis of ubiquitin recognition by the winged-helix domain of Cockayne syndrome group B protein. *Nucleic Acids Res.* **47**, 3784–3794 (2019).

64.  Harami, G. M., Gyimesi, M. & Kovács, M. From keys to bulldozers: Expanding roles for winged helix domains in nucleic-acid-binding proteins. *Trends in Biochemical Sciences* vol. 38 364–371 (2013).

65.  ARAVIND, L., ANANTHARAMAN, V., BALAJI, S., BABU, M. & IYER, L. The many faces of the helix-turn-helix domain: Transcription regulation and beyond. *FEMS Microbiol. Rev.* **29**, 231–262 (2005).

66.  Qin, B. *et al.* STK38 promotes ATM activation by acting as a reader of histone H4 ufmylation. *Sci. Adv.* **6**, 1–10 (2020).

67. Stephani, M. *et al.* A cross-kingdom conserved er-phagy receptor maintains
endoplasmic reticulum homeostasis during stress. *Elife* **9**, (2020).

68. Baek, K., Scott, D. C. & Schulman, B. A. NEDD8 and ubiquitin ligation by cullin-RING
E3 ligases. *Curr. Opin. Struct. Biol.* **67**, 101–109 (2021).

69. Holm, L. & Rosenström, P. Dali server: Conservation mapping in 3D. *Nucleic Acids*
*Res.* **38**, (2010).

70. Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M. & Barton, G. J. Jalview
Version 2--a multiple sequence alignment editor and analysis workbench.
*Bioinformatics* **25**, 1189–1191 (2009).

71. Katoh, K. & Toh, H. Parallelization of the MAFFT multiple sequence alignment
program. *Bioinformatics* **26**, 1899–1900 (2010).

72. Johnson, A. G. *et al.* RACK1 on and off the ribosome. *RNA* **25**, 881–895 (2019).

73. Lapointe, C. P. *et al.* Dynamic competition between SARS-CoV-2 NSP1 and mRNA
on the human ribosome inhibits translation initiation. *Proc. Natl. Acad. Sci.* **118**,
e2017715118 (2021).

74. Shevchenko, A., Tomas, H., Havliš, J., Olsen, J. V. & Mann, M. In-gel digestion for
mass spectrometric characterization of proteins and proteomes. *Nat. Protoc.* **1**, 2856–
2860 (2007).

**Figures**

## Figure 1

**A**

**B**

**D**

**C**

**E**

**Figure 1: - Nucleophilic reactivity profiling of UFC1**

**A.** Schematic showing the workflow for substrate independent single turnover
lysine discharge assays.

**B.** Assay to monitor discharge of Ubiquitin from UBCH5C and UBCH7 in the
presence of free amino acids. 0 indicates time 0.

**C.** Assay to check for discharge of UFM1 from UFC1 in the presence of free
amino acids. UBCH5C is used as a positive control for lysine and cysteine
discharge.

**D.** Lysine discharge assays using UBCH5C and UFC1 in the presence of
increasing concentration of free Lysine.

**E.** Time-dependent analysis of discharge of UFM1 from UFC1 in the presence of
high concentration of free lysine (150 mM).

**Figure 2**

945
946

**Figure 2: -** *In vitro* **reconstitution of the active UFM1 E3 ligase.**

**A.** Comparison of size exclusion chromatography profiles of UFL1 expressed
alone (Red) and co-expressed UFL1/UFBP1 complex (Black) run under
identical buffer conditions on a HiLoad^TM Superdex 10/300 gl column.
Molecular weight standards are shown in grey. Fractions corresponding to

each peak were collected and analysed on 4-12% denaturing SDS-PAGE
which is shown below.

**B.** Mass photometry analysis of co-purified UFL1/UFBP1 complex. The
theoretical and experimental molecular weights are indicated above.

**C.** Immunoblot comparing UFL1 autoUFMylation *(Top)* and UFM1 chain
synthesis *(Middle)* in the presence of UFL1 expressed alone and in complex
with UFBP1. *(Bottom)* Assays to monitor UFMylation of substrates using
purified TRIP4 in the presence of UFL1 alone and UFL1/UFBP1 complex.
Reaction products were run on a 4-12% SDS-PAGE gel and analysed by
immunoblotting using indicated antibodies.

**D.** Immunoblot showing formation of free UFM1 chains in the presence and
absence of UFL1/UFBP1. *(Bottom)* Schematic representation of the linkage
composition of di-UFM1 chains obtained from LC-MS/MS analysis.

**E.** *In vitro* UFMylation assay in the presence of Lys to Arg (K-R) and lysine
less(K0) mutants of UFM1 to check for polyUFMylation *(top)* and UFL1
autoUFMylation *(bottom)* .

**F.** *In vitro* UFMylation assay to check for formation of di-UFM1 chains in the
presence of single Lys and Lys less mutants of UFM1.

972
973

**Figure 3: - UFL1/UFBP1 is a scaffold type E3 ligase**

**A.** Multiple sequence alignment of 1-410 a.a. region of UFL1 from various organisms to highlight the degree of conservation of cysteine residues in this region.

**B.** Immunoblot showing *in vitro* UFMylation assays in the presence of C to A mutants of UFL1 to check for UFBP1 UFMylation *(top)*, UFL1 autoUFMylation *(middle)* and substrate UFMylation *(bottom)*.

**C.** Coomassie stained SDS-PAGE gel showing discharge assays in the presence and absence of full length UFL1/UFBP1 complex. Charged UFC1 was incubated with lysine (50 mM ) in the presence and absence of full length UFL1/UFBP1 complex and analysed for discharge on a 4-12% SDS PAGE gel under non-reducing conditions.

**D.** Graphical representation showing the composition of linkage forms of di-UFM1 chains formed by minimal reconstitution of UBA5 and UFC1.

988
989
990

**Figure 4: - UFL1/UFBP1 complex functions as a scaffold type E3 ligase**

**A.** Cartoon representation of predicted full length human UFL1/UFBP1 complex using Alphafold. The individual winged helix domains of UFL1 and UFBP1 is shown in blue and cyan respectively. The dimeric interface of UFL1 and UFBP1 is highlighted in the dotted box.

**B.** Enlarged view of the dimeric interface of UFL1/UFBP1 complex shown to highlight the formation of full WH domain bridged by two partial winged helix domains of UFL1 (shown in blue) and UFBP1 (shown in teal).

**C.** Schematic representation of UFL1/UFBP1 constructs with different domain boundaries to identify the minimal catalytic region of UFL1/UFBP1 complex required for aminolysis.

**D.** Quantitative representation of Lysine discharge assays to identify the minimal boundaries of UFL1 and UFBP1 required for activation of UFC1 (n=3, mean±SDs.

**E.** Size exclusion chromatography profiles showing that full length UFL1/UFBP1 complex interacts specifically with charged E2. UFL1/UFBP1 complex was incubated with UFC1$^{\text{-O-UFM1}}$ at the molar ratio of 1:1 for 20 min at 4°C and loaded on a Superdex 200 Increase 3.2/300 column. The fractions corresponding to each peak were collected and analysed on a 4-12% SDS PAGE gel followed by Coomassie staining.

**F.** Minimal catalytic region is sufficient for interaction with charged UFC1. UFL1$^{(1-179)}$/UFBP1$^{(1-116)}$ complex was incubated with UFC1$^{\text{-O-UFM1}}$ at the molar ratio of 1:1 for 20 min at 4°C and analysed by analytical size exclusion chromatography as described in E).

**G.** *In vitro* UFMylation assay to monitor formation of free di-UFM1 chains in the presence of UFL1/UFBP1 complexes bearing different domain boundaries.

**H.** Substrate UFMylations assays to check for UFMylation of purified Histone H4 in the presence of different UFL1/UFBP1 truncations.

**I.** Role of UFBP1 in substrate UFMylation. *(Top)*Comparison of UFMylation activities of UFL1/UFBP1$^{29\text{-end}}$ and UFL1/UFBP1$^{207\text{-end}}$ using Histone H4. *(Bottom)*Comparison of UFMylation activities of UFL1$^{1-179}$/UFBP1$^{29\text{-end}}$ and UFL1$^{1-179}$/UFBP1$^{207\text{-end}}$ using Histone H4.

**Figure 5**

1029
1030

**Figure 5: - CDK5RAP3 inhibits E3 ligase activity of UFL1/UFBP1 *in vitro***

**A.** CDK5RAP3 forms a complex with UFL1/UFBP1 *in vitro*. 30 µg of
UFL1/UFBP1 was mixed with 15 µg of CDK5RAP3 and loaded on an
analytical gel filtration column. (Bottom) Fractions were collected and
analysed on 4-12% SDS PAGE and visualized by Coomassie staining.

**B.** Mass photometry analysis showing the experimental molecular weight of
UFL1/UFBP1/CDK5RAP3 complex.
**C.** *In vitro* UFMylation assay in the presence of increasing concentrations of
CDK5RAP3 to monitor the E3 ligase activity of UFL1/UFBP1 complex. The
reaction products were run on a 4-12% SDS-PAGE gel and immunoblotting
was performed using indicated antibodies.
**D.** *In vitro* UFMylation assay to monitor the UFMylation of purified TRIP4 in the
presence and absence of CDK5RAP3.
**E.** Lysine discharge assays to check for activation of UFC1 by UFL1/UFBP1 in
the presence and absence of CDK5RAP3. The reaction products were run on
a 4-12% SDS PAGE and visualized by Oriole staining.
**F.** Quantification of discharge of UFM1 from UFC1 in the presence and absence
of CDK5RAP3 as seen in E). n=3, mean ±SDs.
1049

**Figure 6**

1050
1051

**Figure 6:- Role of the N-terminal helix of UFC1 in UFMylation**

**A.** Multiple sequence alignment of UFC1 homologs from various organisms. A graphical representation of the secondary structure is extracted from the crystal structure of UFC1 (PDB ID:2Z6O)

**B.** Crystal structures of UBE2D3 (PDB ID:5EGG) and UFC1 (PDB ID:2Z6O) are shown in cartoon representation. *(Inset)* Catalytic cysteines and critical residues required for catalytic activity of UBE2D3 (green) and UFC1(dark red) are labelled and shown in ball and stick representation.

**C.** *In vitro* UFMylation assays in the presence of Wild type UFC1 (UFC1$^{WT}$) and UFC1 lacking the N-terminal helix (UFC1$^{\Delta N}$). Schematic representation of domain features of UFC1$^{WT}$ and UFC1$^{\Delta N}$ is shown above.

**D.** Assay to compare Lys discharge activities of UFC1$^{WT}$ and UFC1$^{\Delta N}$ on its own, in the presence of UFL1/UFBP1 or in the presence of preassembled UFL1/UFBP1/CDK5RAP3 complex. Top gel: LiCOR scan of fluorescently labelled UFM1; bottom gel – Coomassie stained (Representative of three independent experiments).

**E.** Quantitative analysis of the intrinsic Lys reactivity (Lysine 25 mM) of UFC1$^{WT}$ and in UFC1$^{\Delta N}$ in the absence of E3 ligase.

**F.** Quantitative analysis of Lys reactivity of UFC1$^{WT}$ and in UFC1$^{\Delta N}$ in the presence of UFL1/UFBP1.

**G.** Quantitative analysis of Lys reactivity of UFC1$^{WT}$ and in UFC1$^{\Delta N}$ in the the presence of preassembled UFL1/UFBP1/CDK5RAP3. E – G: n=3, mean ±SD.

1076 **Supplementary Material**

1077

1078

# Figure S1

1079
1080

1081 **Figure S1: Analysis of E2 discharge**

1082    **A.** & **B.** Time-dependent analysis of discharge of Ubiquitin from UBCH5C and

1083        UBCH7 in the presence of 150 mM.

1084

**Figure S2**

1085
1086
1087

**Figure S2: UFL1 is only active in the presence of co-expressed UFBP1**

**A.** Size exclusion chromatography profile of UFL1 run on Superdex 200 Increase 10/300gl column (shown in red). Overlay of chromatogram of molecular weight standards of different sizes (shown in grey) run under same buffer conditions.

**B.** Results obtained from ULTIMATE Y2H[TM] screening (Hybrigenics) for binary interactions with UFL1. Asterisk denotes protein hits obtained with low confidence.

**C.** Schematic describing the strategy for co-expression and purification of UFL1/UFBP1 complex.

**D.** SEC-MALS analysis of UFL1/UFBP1 complex. The theoretical and observed molecular weights are indicated.

**E.** Analytical gel filtration chromatography analysis of UFL1/UFBP1 complex and its subunits run on Superdex 200 3.2/300.

**F.** UFBP1 does not activate UFL1 when added exogenously. *In vitro* UFMylation assays to compare the E3 ligase activity of UFL1 and UFBP1 expressed alone and together as a complex. 0.25 µM UBA5, 5 µM UFC1 and 10 µM UFM1 was incubated with 1 µM UFL1 or 1 µM UFBP1 or 1 µM UFL1/UFBP1 complex for 1 h at 37°C in the presence of 50 mM HEPES 7.5, 0.5 mM DTT, 10 mM $MgCl_2$ and 10 mM ATP. The reaction was stopped by addition of 3x SDS loading dye and run on a 4-12% denaturing SDS PAGE gel under reducing conditions and immunoblotting was performed using indicated antibodies.

**G.** $MS^2$ spectra showing the peptide derived from *in vitro* UFMylation assay showing the VG-remnant on K69 of UFM1.

**Figure S3: Mode of UFMylation by UFL1/UFBP1 complex**

**A.** Composite Foldindex profile of UFL1 showing folding propensity of different regions of UFL1 to aid in construct design for soluble protein expression.

**B.** Size exclusion chromatography profile of MBP-UFL1$^{1-410}$ (dark blue) and His$_6$-UFL1$^{1-410}$/UFBP1$^{29-end}$(orange) run on Superdex 200 3.2/300 column (shown in red). Approximately 20 µgs of sample was run on a Superdex 200 3.2/300 analytical gel filtration column and UV traces measured at 280 nm were recorded and analysed. (Right) Coomassie stained gel showing the purity of the proteins.

**C.** *In vitro* UFMylation assay to check for E3 ligase activity of UFL1$^{1-410}$/UFBP1$^{29-end}$. Full length UFL1/UFBP1 complex is used as a positive control.

**D.** Time course assay to check for transthiolation activity of UFL1. Reaction
products were analysed on a 4-12% SDS PAGE gel under reducing and non-
reducing conditions.
**E.** *In vitro* UFMylation assays to check for formation of di-UFM1 chains in the by
minimal reconstitution in the presence UBA5 and UFC1 alone.
**F.** Immunoblot to check for the presence of UFM1 chains with and without
treatment of UFSP2.
1134

**Figure S4: - Identifying the minimal ligase domain of UFL1/UFBP1 complex**

    **A.** Predicted Aligned Error (PAE) scores of predicted UFL1/UFBP1 models using Alphafold.

    **B.** List of structurally similar proteins from the PDB25 database predicted using UFL1 WH1(52-115) domain as query structure using DALI server.

    **C.** Superposition of the winged helix (WH) domains of UFL1 and UFBP1.

**D.** Comparison of composite winged helix domain formed by two hemi WH
domains of UFL1 (pWH) and UFBP1 (pWH') with the WH1 domain of UFL1.
**E.** Overlaid structures of Winged helix 1(WH1) domain of UFL1(52-115) and top
three hits obtained from DALI search shown in cartoon representation.
**F.** Coomassie stained gel showing purity of different UFL1/UFBP1 truncations.
**G.** Schematic showing the tandem WH domains which constitute the minimal
ligase domain.
1150
1151

**Figure S5: - UFL1/UFBP1 E3 ligase complex binds to UFC1~UFM1 and induces aminolysis**

**A.** Lysine discharge assays to check for activation of UFC1 in the presence of different UFL1 truncations. Top gel: LiCOR scan of fluorescently labelled UFM1; bottom gel – Coomassie stained (Representative of three independent experiments).

**B.** Lysine discharge assays as in A) in the absence of NTR of UFBP1(representative of 3 independent experiments).

**C.** Coomassie stained gel showing analysis of *in vitro* UFMylation reaction
products to check for the formation of stable oxy-ester linked UFC1-UFM1
conjugate (UFC1$^{\text{-O-UFM1}}$).

**D.** Chromatogram obtained from SEC analysis using HiLoad$^{\text{TM}}$ Superdex 75
16/60pg column. (Bottom left) The fractions collected were run on an SDS-
PAGE gel to identify fractions that contained pure UFC1$^{\text{-O-UFM1}}$. (Bottom right)
Coomassie stained gel showing analysis of purified UFC1$^{\text{-O-UFM1}}$ product to
check for homogeneity.

**E.** Quality check to analyse if UFC1-UFM1 conjugate is linked through an oxy-
ester linkage by alkaline hydrolysis.

**F.** Substrate UFMylations assays to check for UFMylation of MRE11 in the
presence of different UFL1/UFBP1 truncations.

1174

1175

1176

# Figure S6

**Figure S6: - Sequence and secondary structure analysis of CDK5RAP3**

**A. (Left)** Multiple sequence alignment of CDK5RAP3 homologs showing predicted secondary structure features using JPred[TM] secondary structure program module from Jalview 2.11.1.7.**(Right)** Schematic showing predicted secondary structure features of CDK5RAP3.

## Figure S7

**Figure S7:- Role of CDK5RAP3 in UFMylation**

**A.** *In vitro* UFMylation assay to compare the E3 ligase activity of UFL1/UFBP1 mixed with CDK5RAP3 and preassembled ternary E3 ligase complex containing UFL1/UFBP1/CDK5RAP3.

**B.** UV traces of gel filtration chromatogram showing co-migration of (UFL1/UFBP1)/CDK5RAP3/UFC1$^{-O-UFM1}$ complex. Approximately, 200 µl of

sample containing (UFL1/UFBP1)/CDK5RAP3/UFC1$^{-O-UFM1}$ in the molar ratio

of 1:1.5:3 was mixed and incubated at 4°C for 1 h and loaded onto a

Superdex 200 10/300 g/l column. The fractions were collected and analysed

on a 4-12% SDS PAGE and visualized by Coomassie staining.

**C.** Pulldown assay to check for interaction of UFL1/UFBP1 with charged UFC1 in

the presence of absence of CDK5RAP3. Around 10 µM of Untagged UFC1

and 10 µM OF UFC1-O-UFM1 were mixed with 5 µM of UFL1/UFBP1

complex in the presence and absence of CDK5RAP3.

**D.** Lysine discharge assays to check for UFC1 discharge in the presence of

UFL1/UFBP1      mixed      with      CDK5RAP3      and      preassembled

UFL1/UFBP1/CDK5RP3 complex. The reaction products were run on a 4-

12% SDS PAGE analysis and visualized by LiCor.

**E.** *In vitro* UFMylation assay to monitor UFMylation of purified substrates namely

TRIP4, MRE11A and Histone H4.

1207

1208

1209

1210 **Figure S8:- Role of the N-terminus of UFC1 in regulating UFMylation**

1211     **A.** *In vitro* UFMylation assay to compare the activites of UFC1$^{WT}$ and UFC1$^{\Delta N}$ on

1212     its own, in the presence of UFL1/UFBP1 or in the presence of preassembled

1213     UFL1/UFBP1/CDK5RAP3 complex (Representative of three independent

1214     experiments).