1

Quantifying changes in individual-specific template-based representations of center-of-mass
dynamics during walking with ankle exoskeletons using Hybrid-SINDy

Michael C. Rosenberg[1*], Joshua L. Proctor[1,2], Katherine M. Steele[1]

[1]Department of Mechanical Engineering, University of Washington, Seattle, United States of America
[2]Department of Applied Mathematics, University of Washington, Seattle, United States of America

* Corresponding author
Email: mcrose3@emory.edu (MCR)

Abstract

Ankle exoskeletons alter whole-body walking mechanics, energetics, and stability by altering center-of-mass (CoM) motion. Controlling the dynamics governing CoM motion is, therefore, critical for maintaining efficient and stable gait. However, how CoM dynamics change with ankle exoskeletons is unknown, and how to optimally model individual-specific CoM dynamics, especially in individuals with neurological injuries, remains a challenge. Here, we evaluated individual-specific changes in CoM dynamics in unimpaired adults and one individual with post-stroke hemiparesis while walking in shoes-only and with zero-stiffness and high-stiffness passive ankle exoskeletons. To identify optimal sets of physically interpretable mechanisms describing CoM dynamics, termed *template signatures*, we leveraged hybrid sparse identification of nonlinear dynamics (Hybrid-SINDy), an equation-free data-driven method for inferring sparse hybrid dynamics from a library of candidate functional forms. In unimpaired adults, Hybrid-SINDy automatically identified spring-loaded inverted pendulum-like template signatures, which did not change with exoskeletons ($p>0.16$), except for small changes in leg resting length ($p<0.001$). Conversely, post-stroke paretic-leg rotary stiffness mechanisms increased by 37-50% with zero-stiffness exoskeletons. While unimpaired CoM dynamics appear robust to passive ankle exoskeletons, how neurological injuries alter exoskeleton impacts on CoM dynamics merits further investigation. Our findings support Hybrid-SINDy's potential to discover mechanisms describing individual-specific CoM dynamics with assistive devices.

Introduction

Ankle exoskeletons are used to improve walking function and gait mechanics[1-3]. Personalized passive and powered ankle exoskeletons have shown promise to improve walking function and muscle coordination in unimpaired adults and individuals with neurological injuries[2,4,5]. However, changes in gait in response to ankle exoskeletons are highly individualized, especially following neurological injury, making device personalization critical to improving mobility. For example, in stroke survivors and children with cerebral palsy, ankle exoskeletons elicit diverse—and sometimes detrimental—impacts on gait mechanics, walking speed, step length, and the energetic cost of walking[1-4]. Quantifying and characterizing these responses remain challenging and hinders clinicians' and designers' abilities to customize exoskeletons to support walking function.

Despite observed changes in gait mechanics and energetics with ankle exoskeletons, little is known about how ankle exoskeletons impact the neural and biomechanical processes governing center-of-mass (CoM) motion (*i.e.*, CoM dynamics) to achieve task-level goals, like walking stably and efficiently[2,5-10]. For example, CoM energetics are altered in unimpaired adults walking with ankle exoskeletons and following neurological injuries (*e.g.,* post-stroke, the paretic leg exhibits reduced power generation and changes in leg power with exoskeletons are heterogenous)[2,5,11]. However, whether these changes in leg power are driven by changes in CoM dynamics or simply altered kinematics is unclear. Characterizing the changes in CoM dynamics with ankle exoskeletons may help explain heterogeneous exoskeleton impacts on task-level goals during walking.

Reduced-order representations of CoM dynamics, often termed *template* models, provide a foundation to quantify complex exoskeleton responses using interpretable mechanical elements[12-17]. For example, Full & Kodistchek (1999) used template models of CoM dynamics, such as the spring-loaded inverted pendulum (SLIP), to quantify strategies to stabilize the CoM in response to perturbations[15]. Such reduced-order representations of gait encode neural and biomechanical dynamics using a minimalist set of physics-based mechanisms. Specifically, common template models of CoM dynamics use a variety of mechanisms, such as SLIP leg springs or the rigid legs of an inverted pendulum walker, to describe relationships between leg kinematics and CoM accelerations during gait[10,13,15,16,18-22]. Each mechanism within a model, therefore, encodes a hypothesis about how neural and biomechanical subsystems interact to achieve task-level walking goals.

However, selecting a set of template mechanisms to optimally describe an individual's CoM dynamics remains challenging. Inverted pendulum templates have been useful in describing CoM energetics, the transition from walking to running[16,18], strategies for energetically efficient CoM acceleration[8,22,23] and lateral stabilization[10,24]. Higher-dimensional template structures, such as the bipedal SLIP further improved predictions of sagittal-plane ground reaction forces (GRFs)[20,25]. Extensions of the bipedal SLIP templates have included leg damping mechanisms, rotary springs, or curved feet to further improve model accuracy[19,26]. The breadth of templates proposed for human walking suggests that the mechanisms that best describe gait are individual-specific[20,26]. Further, model coefficient estimates may be sensitive to the mechanisms selected. Consequently, selecting an optimal template structure (*i.e.*, the mechanisms included in the template) to describe an individual's gait is non-trivial but important in quantifying exoskeleton impacts on CoM dynamics using template models.

To emphasize the individual-specific nature of templates, we denote the combination of mechanisms best describing individual-specific CoM dynamics as a *template signature*. Deviations of signature mechanisms or model coefficients between individuals or trials may provide insight into how exoskeletons impact CoM dynamics. For example, template signatures in children with hemiparetic cerebral palsy differed from typically developing children and were asymmetric, with increased stiffness

83 in the paretic leg[27,28]. Atypical and asymmetric CoM dynamics suggested that, following neurological
84 injury, people may adopt individual- and leg-specific strategies to accelerate the CoM.

86 However, characterizing changes in template signatures with exoskeletons or following neurological
87 injury requires addressing a major methodological challenge: Manually identifying optimal template
88 signature structure is a slow, *ad hoc* process that relies on first-principles knowledge of the system[19].
89 Therefore, comprehensively comparing candidate template signatures for even a moderate number of
90 template mechanisms is challenging: the number of comparisons increases combinatorially with the
91 number of candidate mechanisms. New approaches are needed to select mechanisms rapidly and
92 systematically from a literature-based library of candidate mechanisms.

94 Recent advances in data-driven modeling and machine learning provide powerful tools to identify
95 template signatures from walking data[29-32]. One such algorithm, **Hybrid-SINDy** (*SINDy: Sparse*
96 *identification of nonlinear dynamics*[29]), identifies sparse nonlinear dynamics in hybrid systems from time-
97 series data, making it particularly appropriate for identifying template models of walking, which often
98 have distinct dynamics based on foot contact configuration[28,31]. Importantly, Hybrid-SINDy identifies and
99 compares a large number of candidate dynamical models (*e.g.,* template signatures) from an arbitrary
100 library of possible functional forms (*i.e.,* mechanisms). The algorithm uses information criteria to
101 determine levels of support for each candidate model and selects only those that are parsimonious and
102 highly representative of the system[33]. Therefore, this approach enables rapid, systematic identification of
103 individual-specific template signatures while leveraging first-principles knowledge of the physics of
104 human gait.

106 The purpose of this study was to identify changes in template-signature-based representations of CoM
107 dynamics in response to ankle exoskeletons. We used the Hybrid-SINDy algorithm to identify physically
108 interpretable, low-dimensional template signatures describing CoM dynamics during walking in
109 unimpaired adults and evaluated how template signature coefficients changed with hinged and stiff ankle
110 exoskeletons. We hypothesized that the addition of ankle exoskeleton frame and stiffness would alter
111 template signature coefficients. Additionally, to examine the potential of Hybrid-SINDy-based template
112 signatures to encode changes in CoM dynamics with ankle exoskeletons for individuals with neurological
113 injuries, we present a case study evaluating altered template signatures in one individual with post-stroke
114 hemiparesis.

Methods

Data collection

We collected 3D marker trajectories using a ten-camera motion capture system (Qualisys AB, Gothenburg, SE) and GRFs using an instrumented treadmill (Bertec Inc., Columbus, USA) in twelve unimpaired adults (6M/6F; age = 23.9 ± 1.8 years; height = 1.69 ± 0.10 m; mass = 66.5 ± 11.7 kg) and one stroke survivor with lower-limb hemiparesis (*sex not disclosed*; age = 24 years; height = 1.70 m; mass = 68.0 kg). Participants walked at their self-selected steady-state speed on a fixed-speed treadmill in shoes-only and with bilateral passive ankle exoskeletons under two conditions: zero resistance to ankle flexion (*i.e.*, zero stiffness; K_0) and high dorsiflexion resistance (*i.e.*, high stiffness; K_H = 5.1 Nm/deg; Figure 1). The order of walking conditions was randomized. A detailed description of the acclimatization protocol and data preprocessing can be found in[34]. Briefly, in a second session following a practice session, data were collected while participants walked at their self-selected speed—determined in the practice session—for six minutes per condition, including two minutes to acclimate to the treadmill before data were recorded. Only the third and fourth minutes of data were used in this study. To mitigate fatigue, the post-stroke participant walked under the same protocol, but for only three minutes per condition[35]. This study was approved by the University of Washington Institutional Review Board (#47744). The study was performed in accordance with the approved protocol and University of Washington Institutional Review Board guidelines and regulations. All participants provided written informed consent prior to participating in the study.

Figure 1: Depictions of walking conditions, phase variables, and example template state variables.
A) Two-dimensional depictions of template model applied to human walking without and with ankle exoskeletons (left). The phase portrait (right) defined a phase variable, ψ, used to cluster kinematically similar measurements for model fitting. Colors denote gait phases corresponding to first and second double-limb support, single-limb support, and swing or the right leg. B) Stride-averaged global CoM position, velocity, and acceleration for an exemplary unimpaired adult (left). CoM states (middle) and leg states (right) are shown for walking in shoes-only (solid lines), zero-stiffness exoskeletons (K_0; dashed lines), and stiff exoskeletons (K_H; dotted lines). Template states used to fit the template signatures (right) were defined by sagittal- and frontal-plane leg angles, and leg length.

Estimating template signatures with Hybrid-SINDy

We used the Hybrid-SINDy algorithm to identify template signatures during walking with and without
ankle exoskeletons, separately for each walking condition.

Kinematic variable extraction

For each exoskeleton condition, we used OpenSim's *Body Kinematics* algorithm to estimate the CoM and
foot positions[36,37]. In this section, we describe the SINDy and Hybrid-SINDy algorithms in the context of
identifying template signatures, while more detailed explanations can be found in[29] and[31]. The 3D CoM

accelerations, $\ddot{\boldsymbol{q}}(t) \in \mathbb{R}^{m \times n}$, were described by continuous-time nonlinear dynamics, $f\big(q(t), \dot{q}(t)\big)$,
where m denotes the number of samples and $n = 3$ denotes the output variables:

$$\frac{d^2}{dt^2}\boldsymbol{q}(t) = \ddot{\boldsymbol{q}}(t) = f\big(\boldsymbol{q}(t), \dot{\boldsymbol{q}}(t)\big),\qquad\qquad 1$$

where time is denoted by $t \in \mathbb{R}^{m \times 1}$, and $q(t)$ and $\dot{q}(t)$ represent CoM positions and velocities relative to
the feet, respectively, in $\mathbb{R}^{m \times n}$, in the anterior-posterior, vertical, and mediolateral directions. We assume
that only a small number of functional forms (*i.e.*, mechanisms) in $f(q(t), \dot{q}(t))$ describe most of the
system's behavior. We omit the time notation in the remaining sections.

Sparse Identification of Nonlinear Dynamics (SINDy)
The SINDy algorithm[29] recovers sparse nonlinear dynamics from a library of candidate functional forms,
which may consist of arbitrary nonlinear functions of system measurements. Adopting the notation
from[31], we can rewrite the dynamics in equation (1) as:

$$\ddot{\boldsymbol{q}} = \boldsymbol{\Theta}(\boldsymbol{q}, \dot{\boldsymbol{q}})\boldsymbol{\Xi},\qquad\qquad 2$$

where $\boldsymbol{\Xi} \in \mathbb{R}^{p \times n}$, is a linear map from nonlinear function library encoding mechanisms potentially
describing CoM dynamics, $\boldsymbol{\Theta}(\boldsymbol{q}, \dot{\boldsymbol{q}}) \in \mathbb{R}^{m \times p}$, to CoM accelerations, $\ddot{\boldsymbol{q}}$. The coefficients in $\boldsymbol{\Xi}$, therefore,
describe how each template signature mechanisms accelerated the CoM, with non-zero coefficients
denoting the active terms constituting the template signature structure. In the context of the template
model investigated here, non-zero coefficients denote the mechanisms included in the model. We
included $p = 14$ functional forms in the function library (7 per leg), described below. The SINDy
algorithm promotes sparsity in the model using sequential least-squares regression with hard thresholding,
with the threshold defined by the sparsity parameter, λ (equation 3)[31]. This thresholding approach
penalizes the zero-norm of $\boldsymbol{\Xi}$ and solves:

$$\min_{\boldsymbol{\Xi}} || \boldsymbol{\Theta}(\boldsymbol{q}, \dot{\boldsymbol{q}})\boldsymbol{\Xi} - \ddot{\boldsymbol{q}}||_2 + \lambda\|\boldsymbol{\Xi}\|_0,\qquad\qquad 3$$

Hybrid-SINDy
Hybrid-SINDy extends SINDy in two important ways. First, Hybrid-SINDy uses clustering to generalize
SINDy to hybrid systems. For human walking, clustering enables unique dynamics to be identified in
each *gait phase*, defined by foot contact configuration (*i.e.,* single- and double-limb support)[9,14,18]. We
replace the term *hybrid regime*, used in the original Hybrid-SINDy manuscript, with *gait phase* for clarity
in the context of human walking[31]. Second, Hybrid-SINDy uses information criteria to automatically
select the system dynamics that best describe the data. This approach enables competing hypotheses about
the mechanisms describing CoM acceleration to be rapidly and systematically compared, thereby
highlighting mechanisms that are critical to describing CoM dynamics across individuals and mechanisms
unique to a subset of individuals. Note that Hybrid-SINDy does not compare entirely distinct sets of
governing equations (e.g., SLIP-like template vs. a passive dynamic walker with knees[12]). Rather, the
algorithm selects which mechanisms should be included in a SLIP-like template model to best describe
CoM dynamics.

Applying Hybrid-SINDy to walking
We applied the Hybrid-SINDy algorithm to human gait using the following steps for each participant and
walking condition (Figure 2). Note that within each gait phase, we expanded upon the original Hybrid-
SINDy algorithm by using multi-model inference to define a single template signature when multiple
signatures were plausible (Step 5)[31].

1. *Clustering* (Figure 2b): We used a clustering approach to increase robustness to measurement noise
and identify frequently occurring template model structures[31,38]. For the first 3600 samples (30s; ~25-
30 strides) in the training set (10800 samples were available for clustering in each trial), we generated
clusters of each sample's 800 nearest neighbors and identified the centroid of each cluster (3600
clusters, total). These clusters were used to estimate template model coefficients (Step 2). During
clustering, nearest neighbors were selected based on their continuous kinematic phase: the phase
angle of the right/non-paretic leg angle and angular velocity relative to vertical (Figure 1; top
right)[34,39]. We also used this phase variable to normalize stride progression. The last 3600 samples of
each dataset were withheld from training and used during model evaluation and selection (Step 3).

Some clusters contained data in both single- and double-limb support phases. However, these clusters
tended to have relatively large error during model evaluation, such that they would not be selected as
plausible in Step 3, below[31]. Further, we selected our cluster size to ensure that clusters were small
enough to contain data from only one gait phase: the average cluster width (800 samples) spanned
only 7.4% of the training data, smaller than the duration of double-limb support (10-12%)[40].

2. *Model estimation* (Figure 2b & c): For each training cluster, we used SINDy to estimate the
coefficients of multiple template signatures by sweeping 40 sparsity threshold values, ranging
logarithmically from 1-100% of largest magnitude coefficient in the full-dimensional model in each
cluster. This approach typically produced 5-15 unique signatures per cluster.

3. *Model evaluation and selection*: Using the 3600 samples of held-out data, we evaluated the ability of
each template signature to reconstruct CoM accelerations in the anterior-posterior, vertical, and
mediolateral directions. We computed the average reconstruction error of the held-out data over the
gait cycle (equation 4).

$$\text{error}_\psi = |\mathbf{\Theta}(\boldsymbol{q}, \dot{\boldsymbol{q}})\mathbf{\Xi} - \ddot{\boldsymbol{q}}|_\psi, \qquad\qquad 4$$

where ψ represents the continuous phase of the gait cycle, from 0-100% of a stride.

We selected template signatures based on two criteria: First, we discarded signatures that were
identified in less than 1% of training clusters. Frequently occurring template signatures are more
likely to be robust to measurement noise or stride-to-stride variability, making them better
representations of an individual's gait dynamics[31].

Second, for each gait phase—single- and double-limb support—we selected the frequently occurring
template signatures that had the highest likelihood according to the Akaike Information Criterion
(AIC)[33,41]. The AIC is widely used to compare candidate representations of a system (*e.g.*, template
signatures) according to their number of free parameters and log-likelihood[33]. According to the AIC, a
candidate representation that has a lower AIC score than competing representations is considered the
most statistically plausible (*i.e.*, best) candidate representation of the system. Adopting the
formulation in[31], assuming that model errors are independently, identically, and normally distributed,
the AIC can be written in terms of the number of free parameters, k, number of samples, ρ, and the
sum of squared residuals:

$$AIC = 2k + \rho \ln\left(\frac{\sum_{j=1}^{\rho}\sum_{i=1}^{I}\left(\hat{f}(\mathrm{q}) - \dot{q}\right)_{i,j}^2}{\rho}\right), \qquad\qquad 5$$

237 where in the outer summation is over $\rho = 3600$ samples and the inner summation, is over the $I = 3$
238 output states.

239
240   The AIC favors parsimonious, highly representative models, which is ideal for identifying minimalist
241   representations of gait dynamics. Like Mangan and colleagues[31], we used the AIC corrected for finite
242   sample sizes (AICc):
243

$$AICc = AIC + \frac{2(k+1)(k+2)}{\rho - k - 2}.$$

244
245
246   The correction term approaches zero as the number of samples, ρ, increases. We then determined the
247   relative plausibility of competing template signatures using their relative AICc score, $\Delta AICc_j$[31,33]:
248

$$\Delta AICc_j = AICc_j - AICc_{min},$$

249
where $\Delta AICc_j$ represents the relative AICc score for the j^{th} model. $AICc_{min}$ represents the AICc of the
model with the lowest AICc among the models compared within a gait phase. The best model
according to the relative AICc has a score of $\Delta AICc_j = 0$ and all other models had higher scores.
Burnham and Anderson noted that models with $\Delta AICc_j \leq 2$ have substantial support, while
$\Delta AICc_j > 7$ have low support[33]. We adopted the threshold of[31], deeming template signatures with
$\Delta AICc_j \leq 3$ to be *plausible*.
256
257 4. *Multi-model inference*: Since human gait dynamics are not strictly hybrid, multiple template signature
258   structures may be plausible in each gait phase. To construct a single template signature for each gait
259   phase, we computed a weighted-average signature using Akaike weights, ω_j, where j is the j^{th}
260   plausible model in the gait phase[33]. Note that we performed multi-model inference separately for each
261   gait phase. Akaike weights are defined as:
262

$$\omega_j = \frac{\exp\left(-\frac{\Delta AICc_j}{2}\right)}{\sum_{r=1}^{R} \exp\left(-\frac{\Delta AICc_r}{2}\right)},$$

263
264   where $\exp\left(-\frac{\Delta AICc_j}{2}\right)$ defines the likelihood of the j^{th} template signature given the observations[33]. The
265   denominator denotes the summation of exponentially scaled relative AIC scores over all R candidate
266   models. This approach weighs each signature based on its likelihood relative to the other plausible
267   signatures.
268
5. *Uncertainty estimation and model accuracy* (Figure 2d): To evaluate the robustness of the template
signatures to noise and stride-to-stride variations in the data, we performed 200 bootstrapped
estimates of each template signature coefficient in each gait phase separately for each trial. Each
bootstrapping iteration randomly selected 3600 samples to estimate template signature coefficients,
with replacement. We quantified the robustness of each template signature coefficient to variability in
the data using the coefficient of variation (CV) of each participant and condition[27]. Template
signature coefficients for each participant, condition, and gait phase, were defined by the mean of the
bootstrapped estimates. Figure 2d shows the estimated coefficients of the bootstrapping procedure
applied to a synthetic SLIP model (*see Supplemental – S2*). Synthetic SLIP model parameters and
simulation results (gray bars and trajectories, respectively) were used as ground truth to validate the
algorithm's ability to identify template models of walking (*Supplemental – S2*).

Figure 2: Overview of the Hybrid-SINDy algorithm, as applied to a ground-truth synthetic SLIP walking model. Details of model validation using a synthetic SLIP can be found in Supplemental – S2. A) Example of simulated CoM kinematics from the synthetic SLIP and a diagram of the full-dimensional template model used to estimate template signatures. Each coefficient in the diagram corresponds to one element of the function library, $\boldsymbol{\Xi}$, which maps between nonlinear Template states, $\boldsymbol{\Theta}(\boldsymbol{q}, \dot{\boldsymbol{q}})$ to reconstruct CoM accelerations, $\ddot{\boldsymbol{q}}$. Template state variables are described in Table 1. B) Clustering of kinematically similar samples. The phase portrait (left) of simulated data from the synthetic SLIP (gray) is analogous to that shown in Figure 1 for human walking data. Colors denote clusters in different gait phases. Example clusters (colors) are shown for each gait phase. Multiple candidate template signatures are estimated for each cluster (right; examples of signatures in two clusters). C) In each gait phase, template signatures were compared using the Akaike Information Criterion (AIC). The plots show synthetic SLIP CoM accelerations (gray) and template signature reconstructions (colors) in single-limb (top) and second double-limb (bottom) support. Model errors were low only within the gait phase containing the model's cluster centroid (purple). D) Model coefficients (top) can be compared between walking conditions. Bar plots show ground truth synthetic SLIP parameters (gray) and the corresponding estimated template signature coefficients (colors). The range of percent errors in coefficients are shown above each plot; analogously, differences in human template signatures can be compared between walking conditions. The average ($\pm$1SD) ground truth CoM accelerations from the synthetic SLIP (bottom; gray) and model predictions in each gait phase (colors). Model accuracies are quantified using the coefficient of determination (r^2).

Template signatures mechanisms and dynamics
To model three-dimensional CoM dynamics during walking, we created a function library of candidate
mechanisms based on prior literature (Figure 2a; Table 1). We included *leg springs*[13,19-21,25] and
dampers[26], which produce force along the leg. Leg springs are common energetically conservative
mechanisms used to describe walking and running dynamics and enable a double-limb support phase. Leg
dampers are less common, but have been used to capture non-conservative gait dynamics[26]. We also
included *rotary springs*[13,19] and *dampers* in the sagittal and frontal planes, which enable forcing transverse
to the leg axis. The addition of rotary springs has been shown to improve reconstructions of anterior-
posterior GRFs in a bipedal SLIP[19]. We did not identify rotary damping elements in prior literature but
included them as candidate mechanisms describing non-conservative forcing transverse to the leg.
293
294 We included only passive mechanical elements in the mechanism library because these elements can be
295 used in a hybrid modeling framework to approximate active control of walking, such as for lateral
296 stabilization or to inject and dissipate energy[19,42-44]. Active mechanisms or state-based controllers could
297 produce more parsimonious models but would be more challenging to interpret.
298
299 The mechanisms selected by the Hybrid-SINDy algorithm define the template signature *structure*, which
300 describes characteristic strategies to accelerate the CoM. The identified template signature *coefficients*
301 describe each mechanism's contribution to CoM accelerations.
302
303 The dynamics of a three-dimensional bipedal SLIP augmented with damping and rotary mechanisms may
304 be written as
305

$$M(\ddot{\boldsymbol{q}} - \boldsymbol{g}) = \sum_{j=R,L} \left(-[k_L(L - L_0) + c\dot{L}]\frac{\boldsymbol{q}}{L} - [k_s\theta + c_s\dot{\theta}]\frac{\boldsymbol{q}}{L_s^2} - [k_f\phi + c_f\dot{\phi}]\frac{\boldsymbol{q}}{L_f^2} \right)_j, \qquad 9$$

306
where M is body mass, $\boldsymbol{g}$ is the gravity vector, ϕ describes the traverse-plane leg angle, and θ describes
the leg angle from vertical in the direction defined by ϕ[19,25]. The summation represents the total force
generated by the legs on the CoM. The left-most brackets contain mechanisms that impart forces radially
along the leg: k_L is the leg stiffness, L is the instantaneous leg length, L_0 is the leg resting length, c_L is the
leg damping, $\dot{L}$ is the instantaneous leg velocity. We henceforth denote L_0 as *leg length* for clarity. The
middle bracket contains mechanisms that impart forces transverse to the leg axis in the sagittal plane: k_s
and c_s are the sagittal-plane rotary stiffness and damping, respectively. L_s denotes the sagittal-plane leg
projection. Analogously in the right-most brackets, k_f and c_f represent the frontal-plane rotary stiffness
and damping, respectively, and L_f denotes the frontal-plane leg projection. The derivation of system
dynamics can be found in *Supplemental – S1*.
317
Normalized template mechanisms
To account for inter-individual differences in walking speed and body size during analysis, we
normalized the template signatures (Table 1). Leg stiffness was normalized as in[17,19,31]. Leg resting length
normalized to the measured leg length[19,26]. Rotary stiffness was normalized according to[19]. All damping
terms were converted to damping ratios[26]. The normalized leg, sagittal-plane, and frontal-plane stiffness
mechanisms are denoted by κ_L, κ_s, and κ_f, respectively. The normalized leg, sagittal-plane, and frontal-
plane damping mechanisms are denoted by ζ_L, ζ_s, and ζ, respectively. Normalized leg length is denoted
$\tilde{L}_0$. We can rewrite equation (9) as a linear combination of our normalized coefficients and nonlinear
transformations of our states:
327

$$M(\ddot{\boldsymbol{q}} - \boldsymbol{g}) = [\kappa_L \quad \kappa_L\tilde{L}_0 \quad \zeta_L \quad \kappa_s \quad \zeta_s \quad \kappa_f \quad \zeta_f]_j \boldsymbol{\Theta}(\boldsymbol{q}, \dot{\boldsymbol{q}}), \quad j = \{Right, Left\} \qquad 10$$

329 The CoM position and velocity relative to the feet were used to compute candidate template signature
330 states: leg lengths and lengthening velocities, sagittal-plane leg angles and angular velocities relative to
331 vertical, and frontal-plane leg angles and angular velocities relative to vertical. The complete function
332 library can be found in *Supplemental – S1*.
333

Table 1 – Template signature functional forms.

Term	Symbol	Normalized form
Leg stiffness	κ_L	$\dfrac{k_L L_{bio}}{Mg}$
Leg resting length	$\tilde{L}_0$	L_0/L_{bio}
Leg damping	ζ_L	$\dfrac{c_L}{2\sqrt{k_L M}}$
Sagittal-plane rotary stiffness	κ_s	$\dfrac{k_s}{MgL_{bio}}$
Sagittal-plane rotary damping	ζ_s	$\dfrac{c_s}{\sqrt{\kappa_s M L_{bio}^2}}$
Frontal-plane rotary stiffness	κ_f	$\dfrac{k_f}{MgL_{bio}}$
Frontal-plane rotary damping	ζ_f	$\dfrac{c_f}{\sqrt{\kappa_f M L_{bio}^2}}$

c_L=*leg damping;* c_s=*sagittal-plane rotary damping;*
c_f=*frontal-plane rotary damping;* g = *gravitational*
acceleration; k_L=*leg stiffness;* k_s=*sagittal-plane*
rotary stiffness; k_f=*sagittal-plane rotary stiffness;*
L_{bio} = *biological leg length;* M = *body mass*

334
Evaluating Hybrid-SINDy's ability to select template signatures
336 We evaluated the Hybrid-SINDy algorithm's ability to accurately identify walking dynamics in the
337 presence of noise and an incomplete mechanism library (*i.e.*, missing functional forms relative to the true
338 system dynamics) using forward simulations of a bipedal SLIP walking model (example shown in Figure
339 2)[25]. The synthetic SLIP model analysis and results are described in *Supplemental-S2*.
340
Identifying template signatures in human gait
342 To quantify how well template signatures captured COM dynamics, we computed coefficients of
343 determination (r^2) between the measured CoM accelerations and those predicted by each participant's
344 template signatures, averaged over the anterior-posterior, vertical, and mediolateral directions.
345
346 To evaluate the extent to which each mechanism described CoM accelerations in unimpaired adults, we
347 determined the proportion of participants for whom each template signature coefficient was selected and
348 the average number of non-zero mechanisms in each gait phase. Template signature terms that are
349 identified across individuals may represent mechanisms fundamental to CoM dynamics, while
350 infrequently identified mechanisms may describe individual-specific features of CoM dynamics.
351
To determine if unimpaired CoM dynamics during shoes-only walking generalized to walking with ankle
exoskeletons, we evaluated the ability of shoe-walking template signature structures to reconstruct CoM
accelerations in the K_0 and K_H conditions. We used least-squares regression to estimate template signature

coefficients for the K_0 and K_H conditions using the shoes-only template signature structure. We compared
the AICc scores between these signature structures and those of the signature structures specific to the K_0
and K_H trials. To determine if shoes-only template signatures were less plausible than signature structures
selected for the K_0 and K_H conditions, we used one-sample right-tailed t-tests ($\alpha = 0.05$) to test for
differences in the average relative AICc scores between the shoes-only signature structures and those of
the K_0 and K_H conditions were greater than three (e.g., $AICc_{Shoe} - AICc_{K0} > 3$ for the K_0 condition)[33].
361
To determine if the ankle exoskeleton frame or mass impacted CoM dynamics, we compared template
signature coefficients between the K_0 and Shoe conditions. Similarly, to evaluate the impacts of
exoskeleton stiffness on CoM dynamics, we compared template signature coefficients in the K_H and K_0
conditions. For both comparisons, we used paired independent-samples t-tests with Holm-Sidak step-
down corrections for multiple comparisons ($\alpha = 0.05$)[45]. Because shoes-only template signature structures
were plausible for most unimpaired participants, we re-estimated template signatures in the exoskeleton
conditions using the shoes-only template signature structure to estimate signature coefficients across
walking conditions.
370
371   To determine if CoM dynamics may be altered post-stroke, we computed the percent difference in the
372   non-paretic and paretic leg template signature coefficients during shoes-only walking in one individual
373   with post-stroke hemiparesis. We also evaluated changes in post-stroke CoM dynamics with ankle
374   exoskeletons by computing percent changes in template signature coefficients for the K_0 condition
375   compared to the shoes-only and K_H conditions.
376

Results

When walking in shoes-only, template signatures reveal common and more individual-specific
representations of CoM dynamics across unimpaired participants. Unimpaired template signatures were
not significantly different between legs (paired 2-sample t-test; p > 0.080). In all gait phases, SLIP
mechanisms—leg stiffness and leg length—were selected (*i.e.*, had non-zero coefficients) in 100% of legs
(κ_L and L_0 in Figure 3). Rotary stiffness and damping mechanisms were selected in less than 30% of legs
in single-limb support and swing. On average across participants and legs, 2.3 ± 0.8 and 2.9 ± 1.3 terms
had non-zero coefficient in the stance and swing legs, respectively. More mechanisms were selected
during double-limb support phases: Rotary stiffness terms were selected in 79-83% of legs in the leading
leg and 67-79% of legs in the trailing leg. Damping mechanisms were selected most frequently (33-67%)
in the double-leg support phases (ζ terms in Figure 3). On average, 5.0 ± 1.8 and 4.6 ± 1.6 terms had non-
zero coefficients in first and second double-limb support, respectively.

**Figure 3: The percentage of unimpaired legs whose template signatures contained each
mechanism in each gait phase.** For 24 legs, the percentage of legs for which each template signature
mechanism was selected by the Hybrid-SINDy algorithm. Colors denote each gait phase. Mechanisms
selected in a larger percentage of legs suggest common representations of CoM dynamics, while less

frequently selected mechanisms reflect more individual-specific template features describing CoM dynamics.

For each unimpaired participant, shoes-only template signature coefficients were reliable, having low
bootstrapped coefficients of variation (CVs), during single-limb support and swing: stiffness and leg
length CVs were less than 0.02 and 0.06, respectively (Figure 4; top). In both double-limb support phases,
coefficient estimates were less reliable: CVs ranged from 0.06-4.34 across coefficients. During swing,
coefficient estimates were generally reliable (CV = 0.03-0.10).

Across unimpaired participants, single-limb support leg length was the least variable coefficient (0.97 ±
0.03), while leg stiffness values were more variable between participants (16.7 ± 5.8; Figure 4; top).
Interindividual variability in double-limb support leg and rotary stiffness was larger than single-limb
support coefficients. For example, leg stiffness (1.7 ± 6.0 in first double-limb support) was lower and
more variable than in single-limb support. During swing, leg stiffness was relatively small (-1.5 ± 0.8)
compared to single-limb support. Template signatures explained 83 ± 7% (range: 67-94%) of the variance
in participants' CoM (Example reconstruction of experimental data in Figure 4; bottom).

Figure 4: Unimpaired template signatures of walking in shoes-only. Normalized template signatures (top) and reconstructed CoM accelerations (bottom) for shoes-only walking in each gait phase. A) Bars denote the average template signature (+1SD) in single- and double-limb support. For unimpaired adults, the left and right leg coefficient estimates were grouped in each gait phase (small circles; up to 24 samples per bar). The single-limb support and swing plots contain coefficient estimates from single-support or swing of each leg, respectively. The first and second double-limb support phases show the coefficients of the leading/trailing leg, respectively. Note that we omitted

mechanisms that were selected in less than 25% of participants for clarity. The dashed lines truncate large terms for clarity. B): Experimental (gray) and reconstructed (colors) CoM accelerations from the test dataset of an example unimpaired participant.

Passive ankle exoskeletons elicited only small changes in unimpaired CoM dynamics: in single-limb
support, the shoes-only template signature structures reconstructed CoM dynamics with similar accuracy
to template signature structures selected specifically for the K_0 condition (mean difference in $AICc = $
-1.5 ± 8.4; p = 0.99) and the K_H condition (mean difference in $AICc = 6.8 \pm 14.6$; p = 0.10) (Figure 5;
top). Negative AICc scores indicate that the shoes-only template signature structure was more plausible
than the exoskeleton-specific signature structure, and can occur when the shoes-only template signature
structure is not the most plausible model within individual clusters for the exoskeleton conditions. In
double-limb support, shoes-only signatures did not consistently reconstruct K_0 or K_H CoM dynamics, with
relative AICc scores ranging from $-258 \leq \Delta AICc < 282$. Therefore, to compare signature coefficients
between exoskeleton conditions, we constrained each participant's template signature structures to that
selected in the shoes-only condition.

Using this approach, we found that the only significant difference in unimpaired template signature
coefficients was in the leg length coefficient during single-limb support, which differed between walking
in shoes-only and the zero-stiffness exoskeletons (K_0; p = 7.9e-4; α_{Sidak} = 9.2e-4). However, this change
was small ($\Delta \tilde{L}_0 = -0.01 \pm 0.01$), with leg length being slightly longer in the shoes-only condition.
Neither ankle exoskeleton mass and frame (K_0) nor stiffness (K_H) altered other template signature
coefficients in any gait phase (p > 0.16; α_{Sidak} = 9.3e-4) (Figure 5; bottom).

Figure 5: Changes in template signatures with ankle exoskeletons. A) Differences in AICc scores between template signatures identified specifically for the K_0 and K_H conditions, and signatures constrained to the shoes-only signature structures. Positive $\Delta AICc$ values indicate that shoes-only signature structures were less plausible than those identified specifically for each condition. $\Delta AICc$ scores are shown for single-limb support and swing (left) and double-limb support (right). $\Delta AICc \le$

3 (green lines) denote that shoe template signature structures were plausible for the exoskeleton conditions. For double-limb support, large $\Delta AICc$ scores are truncated at $\Delta AICc = \pm 75$ for clarity. B) Template signatures of walking in shoes only (solid bars), zero-stiffness (K_0; slashed bars) exoskeletons, and high-stiffness exoskeletons (K_H; dotted bars) during single- and double-limb support. Bars represent the average (+1SD) template signature across participants. Leg stiffness (κ_L) is on a separate subplot for clarity.

One stroke survivor's shoes-only template signature was symmetric in single-limb support, swing, and
first double-limb support, but was asymmetric in second double-limb support (Figure 6). In single-limb
support and swing, leg stiffness and leg length mechanisms were selected and reliably estimated for both
legs, with sagittal-plane rotary stiffness also selected in swing (all CV < 0.02). In single-limb support the
paretic leg ($\kappa_L = 11.0$; white bars in Figure 6) was slightly (6%) stiffer than the non-paretic leg ($\kappa_L =$
10.4; colored bars in Figure 6). During first double-limb support, template signature coefficients differed
between legs, but only the paretic leg coefficients were reliably estimated (CV = 0.00-0.06). Conversely,
in second double-limb support, the sagittal- and frontal-plane rotary stiffness mechanisms were selected
for the paretic, but not the non-paretic leg. Unlike first double=limb support, non-paretic leg stiffness and
resting length coefficients were more reliable (CV = 0.00-0.26) than those in the paretic leg (CV = 0.34-
8.01).
436

Figure 6: Inter-leg differences in template signatures in one stroke survivor. Non-paretic (colored bars) and paretic (white bars) template signatures for one individual with post-stroke hemiparesis. Each plot represents a different gait phase.

The exoskeleton mass and frame (Shoe vs. K_0) primarily impacted the stroke survivor's non-paretic and
paretic leg stiffness in single-limb support, and paretic leg rotary stiffness in second double-limb support
(Figure 7). Note that these coefficients were reliably estimated by Hybrid-SINDy. In single-limb support,
the zero-stiffness (K_0) exoskeleton template signatures had 33% greater leg stiffness in the paretic leg and
19% lower leg stiffness in the non-paretic leg compared to shoes-only signatures (Figure 7; slashed bars
vs. no bars). In second double-limb support, sagittal and frontal plane rotary stiffness were 37 and 50%
greater, respectively, in the paretic limb in the K_0 condition compared to shoes-only. Exoskeleton
stiffness (K_0 vs. K_H) had smaller impacts on paretic leg template signatures: in single-limb support,
paretic leg stiffness was 22% less than in the K_0 condition (Figure 7; dotted vs, slashed bars). Conversely,
non-paretic leg stiffness was 85% lower in double-limb support. Paretic leg sagittal- and frontal-plane
rotary stiffness were both 33% lower in second double-limb support than in the K_0 condition.

Figure 7: Template signatures of one stroke survivor walking with and without ankle exoskeletons. Bars denote template signatures in the shoes-only (solid color bars), zero-stiffness (K_0; dashed bars), and high-stiffness (K_H; solid black bars) ankle exoskeleton conditions for the non-paretic (left) and paretic (right) legs. Bars represent each leg's average template signatures over 200 bootstrapped model fitting iterations. Colors correspond to each gait phase.

Discussion

We evaluated the impacts of passive ankle exoskeletons on individual-specific template-based representations of CoM dynamics—described by template signatures—using a recently developed data-driven modeling framework, Hybrid-SINDy. Despite balancing model accuracy with parsimony, template signatures captured CoM dynamics with similar accuracy to prior work using pre-defined 2D template structures[20,26]. The symmetric and SLIP-like unimpaired template signatures automatically selected by Hybrid-SINDy during walking in shoes-only were consistent with prior template models of CoM dynamics during walking, but suggest that dynamics described by rotary mechanisms represent more individual-specific structures describing CoM accelerations[19,25]. Contrary to our hypothesis, template-based representations of unimpaired CoM dynamics were robust to the mechanical constraints of passive ankle exoskeletons: frame and mass (K_0), and dorsiflexion stiffness (K_H). Conversely, in our post-stroke case study, asymmetric shoes-only signatures and changes in template signatures with exoskeletons support Hybrid-SINDy's potential to identify interpretable representations of pathological CoM dynamics, motivating future investigation into how neurological injuries impact template-based representations of CoM dynamics with ankle exoskeletons[28].

Unimpaired and post-stroke template signatures highlight potential inter-individual differences in CoM dynamics. The selection of leg stiffness and resting length as active mechanisms (*i.e.*, terms with non-zero coefficients) in 100% of legs and their selection as the only mechanisms in single-limb support for most participants is consistent with common template walking models and supports the perspective that elastic legs are foundational mechanisms for describing CoM accelerations during unimpaired walking[12,20,21,25,46]. Conversely, our finding that rotary mechanisms were selected to describe CoM dynamics in only 33-83% of legs is consistent with their less frequent application in template walking models[13,19]. One interpretation of individual differences in selected mechanisms is that leg stiffness and resting length describe coordination patterns necessary for stable or efficient walking, while rotary mechanisms describe coordination patterns that have more individual-specific impacts on gait[47]. Alternatively, differences in the selected mechanisms may be due to covariation among template signature state variables. We observed moderate covariation between some variables, particularly in double-limb support (*Supplemental – S2: Covariation of template signature state variables*). However, Hybrid-SINDy penalizes the selection of strongly covarying states, such that both variables would not likely be selected unless they independently increase model likelihood[31,41].

Contrary to our primary hypothesis, unimpaired CoM dynamics are robust to altered ankle constraints due to passive ankle exoskeletons, despite observed changes in kinematics and muscle activity[34]. Similarly, Collins and colleagues (2015) observed small changes in total CoM power with passive ankle exoskeletons compared to walking in shoes-only[5]. Our findings suggest that, if changes in CoM dynamics in single-limb support and swing are captured by the template-based mechanisms in our function library, these small changes in CoM power are driven by changes in CoM kinematics, rather than changes in the underlying CoM dynamics. Note that these findings may not generalize to powered ankle exoskeletons, which elicit larger changes in gait kinematics and kinetics than did our passive exoskeletons and may yield larger changes in template-based representations of CoM dynamics[2,6,34]. Note that we cannot compare coefficient estimates in those gait phases: template signatures in double-limb support were not reliably estimated. More data may be needed to robustly estimate model coefficients in shorter gait phases.

However, changes—or a lack thereof—in template signature coefficients may be biased by an incomplete function library. Because template models are incomplete representations of gait, they may be sensitive to both measurement noise and unmodeled dynamics. Our analysis of a synthetic SLIP (*Supplemental – S2: Effects of measurement noise on algorithm performance*) suggests that Hybrid-SINDy can accurately identify template signature coefficients at measurement noise levels comparable to marker-based motion

capture. However, because template signatures accounted for less than 94% of the variance in human
CoM accelerations, changes in true CoM dynamics with exoskeletons are not completely represented by
our mechanism library. For example, our mechanism library did not include torso dynamics, which are
known to contribute to angular momentum regulation during post-stroke gait and may be altered with
exoskeletons[48,49]. Omitting functional forms from the mechanism library induces at least 5-10%
differences in template signature coefficients estimates (*Supplemental – S2: Effects of missing physics on*
algorithm performance). While we encoded common functional forms from literature, larger function
libraries or novel functional forms may increase the robustness of inter- or intra-individual differences in
CoM dynamics that are more robust to variations in kinematics between trials[13,19-21,25,26,50]. However, even
if a complete set of mechanisms is included in the library, Hybrid-SINDy may not select small but
important mechanisms needed to reconstruct CoM dynamics across tasks[31,41]. Future studies should
consider the tradeoff between the improved interpretability of more-parsimonious models with decreased
model accuracy across the tasks of interest.
514
In our case study of one stroke survivor, asymmetric template signatures suggest that inter-leg differences
in the mechanisms describing CoM dynamics can be automatically identified from data. Consistent with
template-based studies in children with cerebral palsy, the post-stroke participant's paretic leg was
slightly (6%) stiffer than the nonparetic leg in single-limb support, and rotary stiffness mechanisms were
selected only in the non-paretic leg in double-limb support[27,28]. These mechanisms may reflect a more
rigid paretic leg or reliance on proximal muscles for propulsion[51,52]. However, as discussed above, the
small difference in stiffness may be driven by unmodeled torso dynamics[48]. While only a case study,
these results highlight the potential interpretability of subject-specific template signatures to understand
how the legs contribute differentially to CoM accelerations following neurological injury.
524
Changes in our post-stroke case study's template signatures with exoskeletons support Hybrid-SINDy's
ability to identify interpretable impacts of ankle exoskeletons on template-based representations of CoM
dynamics. For example, increases in paretic leg stiffness (K_0 vs. shoes-only) may stem from the
exoskeleton frame restricting inversion of the paretic ankle, which the participant noted during data
collection. Conversely, reduced non-paretic leg stiffness and paretic leg rotary stiffness with stiff
exoskeletons (K_H vs. K_0) may reflect a compensatory strategy to avoid reliance on the paretic leg, if it did
not adapt effectively to the stiff exoskeleton[2]. Note that these findings represent a proof-of-concept that
will not generalize across stroke survivors. Larger studies are needed to understand how individual-
specific neural or biomechanical constraints may alter exoskeleton impacts on CoM dynamics[2,5,7,28,34,53].
534
The Hybrid-SINDy algorithm[31,32] was essential to discovering individual-specific and leg-specific
changes in CoM dynamics with ankle exoskeletons in the present study. Conversely, manually testing all
possible template signatures for our fourteen-dimensional mechanism library would require a
combinatorially large number of models to be generated, fit, and compared. Conversely, Hybrid-SINDy
automatically selected mechanistic representations of CoM dynamics that are consistent with the literature
in a fraction of the time required to manually compare each candidate model[13,19-21,25,26]. While Hybrid-
SINDy has largely been applied to synthetic systems, this work supports its ability to capture dynamics
from human gait data[31,32,54].
543
While alternative modeling approaches, such as principle components analysis (PCA) or non-negative
matrix factorization (NMF), could identify low-dimensional representations of CoM dynamics, Hybrid-
SINDy provides immediately interpretable insight into the structure of these dynamics[55,56]. Hybrid-SINDy
facilitates the interpretation of CoM dynamics by selecting a sparse set of readily interpretable template
variables derived from expert knowledge of human gait[12,13,19-21,25,26]. PCA or NMF-based template
signatures would be harder to interpret, as they would not be sparse in the space of the template variables:
each mode of PCA or NMF would likely contain non-zero coefficients for all state variables[13,19-21,25,26].
More-similar approaches to ours used stepwise regression to model ankle quasi-stiffness using

mechanics-based functional forms[57] or multi-layer optimization to identify human-like template dynamics
for robot controllers[58]. Hybrid-SINDy is distinct from these approaches in its use of clustering and the
AIC to evaluate the relative plausibility of competing representations of CoM dynamics, a limitation of
stepwise regression[41,59,60]. For all participants, Hybrid-SINDy identified a single plausible shoes-only
template signature structure in each gait phase, providing confidence that the identified signatures have
strong statistical support compared to alternative signatures (*see Supplemental - S1: Comparison to*
alternative modeling frameworks).
559
Additional limitations should constrain the interpretation of template signatures. First, human gait
dynamics are continuously phase-varying rather than hybrid, such that template signatures may vary
continuously over the gait cycle. Our preliminary analyses found that predicted CoM dynamics became
less accurate near the transitions between single and double-limb support. Further, our prior work shows
that phase-varying models of gait have higher predictive accuracy, but quantifying changes in the
coefficients of continuously phase-varying models is challenging[34,50]. Because our goal was to quantify
changes in dynamics rather than predict CoM motion with maximal accuracy, defining gait phases based
on contact configuration was reasonable and consistent with existing hybrid template models of
walking[19,20,25,31]. Second, we assumed that CoM dynamics were time-invariant, though they may change
with adaptation to exoskeletons[61]. We included a two-minute adaptation period for each walking
condition, but additional adaptation may elicit larger changes in participants' template signatures.
Identifying the plausibility of template signatures across or within trials could improve our understanding
of how template-based representations of CoM dynamics change during adaptation[26]. Third, our test set—
90-120s of each trial recording—contained 25-30 strides of data per trial and was not a rigorous
evaluation of model generalizability. Our approach identifies template signatures specific to a task and
walking condition, and we do not expect these signatures to generalize across tasks or to overground
walking. However, because template signature coefficients are known to vary with speed[26] and kinematics
(*Supplemental – S2: Effects of missing physics on algorithm performance*), testing on withheld speeds or
walking conditions was not practical. Finally, our limited sample size may have masked exoskeleton
impacts on unimpaired gait.

Conclusions

We quantified changes in individual-specific template model-based representations of CoM dynamics in
response to passive ankle exoskeletons using an interpretable physics-informed data-driven modeling
framework, Hybrid-SINDy. The template mechanisms describing salient features of unimpaired CoM
dynamics were insensitive to ankle exoskeleton frame or stiffness. Interpretable ankle exoskeleton
impacts on template representations of CoM dynamics in a case study of one individual post-stroke
support the utility of template signatures in quantifying post-stroke CoM dynamics with exoskeletons in
people with neurological injuries. These findings also support the potential of data-driven frameworks
like Hybrid-SINDy to accelerate the investigation of individual-specific representations of CoM dynamics
during walking.

Data Availability

All experimental data and modeling code used in this study are freely available at
https://simtk.org/projects/ankleexopred.

Competing interests

The authors declare no competing interests.

Funding

This material is based upon work supported by the National Science Foundation (https://nsf.gov/) under grant no. CBET-1452646 to Katherine. M. Steele, the National Science Foundation Graduate Research Fellowship Program (https://www.nsfgrfp.org/) under grant no. DGE-1762114 to Michael. C. Rosenberg, and a University of Washington Gatzert Child Welfare Fellowship (https://grad.uw.edu/graduate-student-funding/funding-information-for-students/fellowships/list-of-fellowships/gatzert-child-welfare-fellowship/) to Michael. C. Rosenberg. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

M.C.R., K.M.S. and J.L.P. contributed equally to the conception and design of this work and to the interpretation of results. M.C.R. collected data, conducted all analyses, created figures, and wrote the initial manuscript draft. J.L.P. provided theoretical guidance on the implementation of the Hybrid-SINDy algorithm. K.M.S. provided experimental resources and guidance on the analysis of gait with exoskeletons. All authors contributed to substantive revisions of the manuscript.

References

1. Kerkum, Y. L. *et al.* The effects of varying ankle foot orthosis stiffness on gait in children with spastic cerebral palsy who walk with excessive knee flexion. *PloS one* **10**, e0142878 (2015).

2. McCain, E. M. *et al.* Mechanics and energetics of post-stroke walking aided by a powered ankle exoskeleton with speed-adaptive myoelectric control. *Journal of neuroengineering and rehabilitation* **16**, 1-12 (2019).

3. Ries, A. J., Novacheck, T. F. & Schwartz, M. H. The efficacy of ankle-foot orthoses on improving the gait of children with diplegic cerebral palsy: a multiple outcome analysis. *PM&R* **7**, 922-929 (2015).

4. Lerner, Z. F., Harvey, T. A. & Lawson, J. L. A Battery-Powered Ankle Exoskeleton Improves Gait Mechanics in a Feasibility Study of Individuals with Cerebral Palsy. *Annals of biomedical engineering* **47**, 1345-1356 (2019).

5. Collins, S. H., Wiggin, M. B. & Sawicki, G. S. Reducing the energy cost of human walking using an unpowered exoskeleton. *Nature* **522**, 212-215 (2015).

6. Jackson, R. W., Dembia, C. L., Delp, S. L. & Collins, S. H. Muscle–tendon mechanics explain unexpected effects of exoskeleton assistance on metabolic rate during walking. *Journal of Experimental Biology* **220**, 2082-2095 (2017).

7. Sawicki, G. S. & Khan, N. S. A Simple Model to Estimate Plantarflexor Muscle–Tendon Mechanics and Energetics During Walking With Elastic Ankle Exoskeletons. *IEEE Transactions on Biomedical Engineering* **63**, 914-923 (2016).

8. Donelan, J. M., Kram, R. & Kuo, A. D. Mechanical work for step-to-step transitions is a major determinant of the metabolic cost of human walking. *Journal of Experimental Biology* **205**, 3717-3727 (2002).

9. Kuo, A. D. The six determinants of gait and the inverted pendulum analogy: A dynamic walking perspective. *Human movement science* **26**, 617-656 (2007).

10. Kuo, A. D. Stabilization of Lateral Motion in Passive Dynamic Walking. *The International Journal of Robotics Research* **18**, 917-930, doi:10.1177/02783649922066655 (1999).

11. Farris, D. J., Hampton, A., Lewek, M. D. & Sawicki, G. S. Revisiting the mechanics and energetics of walking in individuals with chronic hemiparesis following stroke: from individual limbs to lower limb joints. *Journal of neuroengineering and rehabilitation* **12**, 1-12 (2015).

12. Alexander, R. M. Simple models of human movement. (1995).

13. Davoodi, A., Mohseni, O., Seyfarth, A. & Sharbafi, M. A. From template to anchors: transfer of virtual pendulum posture control balance template to adaptive neuromuscular gait model increases walking stability. *Royal Society open science* **6**, 181911 (2019).

14. McGeer, T. Passive dynamic walking. *I. J. Robotic Res.* **9**, 62-82 (1990).

15. Full, R. J. & Koditschek, D. E. Templates and anchors: neuromechanical hypotheses of legged locomotion on land. *Journal of experimental biology* **202**, 3325-3332 (1999).

16. Cavagna, G. A., Heglund, N. C. & Taylor, C. R. Mechanical work in terrestrial locomotion: two basic mechanisms for minimizing energy expenditure. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **233**, R243-R261 (1977).

17. Holmes, P., Full, R. J., Koditschek, D. & Guckenheimer, J. The dynamics of legged locomotion: Models, analyses, and challenges. *SIAM review* **48**, 207-304 (2006).

18. Srinivasan, M. & Ruina, A. Computer optimization of a minimal biped model discovers walking and running. *Nature* **439**, 72 (2006).

19. Antoniak, G. *et al.* Spring-loaded inverted pendulum goes through two contraction-extension cycles during the single-support phase of walking. *Biology open* **8** (2019).

655   20   Lipfert, S. W., Günther, M., Renjewski, D., Grimmer, S. & Seyfarth, A. A model-experiment
656   comparison of system dynamics for human walking and running. *Journal of theoretical biology*
657   **292**, 11-17 (2012).
658   21   Maus, H.-M., Lipfert, S., Gross, M., Rummel, J. & Seyfarth, A. Upright human gait did not provide
659   a major mechanical challenge for our ancestors. *Nature communications* **1**, 1-6 (2010).
660   22   Garcia, M., Chatterjee, A., Ruina, A. & Coleman, M. The simplest walking model: stability,
661   complexity, and scaling. (1998).
662   23   Buczek, F. L. *et al.* Performance of an inverted pendulum model directly applied to normal
663   human gait. *Clinical Biomechanics* **21**, 288-296 (2006).
664   24   Donelan, J. M., Shipman, D. W., Kram, R. & Kuo, A. D. Mechanical and metabolic requirements
665   for active lateral stabilization in human walking. *Journal of biomechanics* **37**, 827-835 (2004).
666   25   Geyer, H., Seyfarth, A. & Blickhan, R. Compliant leg behaviour explains basic dynamics of walking
667   and running. *Proceedings of the Royal Society B: Biological Sciences* **273**, 2861-2867 (2006).
668   26   Kim, S. & Park, S. Leg stiffness increases with speed to modulate gait frequency and propulsion
669   energy. *Journal of Biomechanics* **44**, 1253-1258,
670   doi:https://doi.org/10.1016/j.jbiomech.2011.02.072 (2011).
671   27   Fonseca, S. r. T., Holt, K. G., Fetters, L. & Saltzman, E. Dynamic resources used in ambulation by
672   children with spastic hemiplegic cerebral palsy: relationship to kinematics, energetics, and
673   asymmetries. *Physical Therapy* **84**, 344-354 (2004).
674   28   Holt, K. G., Fonseca, S. T. & LaFiandra, M. E. The dynamics of gait in children with spastic
675   hemiplegic cerebral palsy: theoretical and clinical implications. *Human Movement Science* **19**,
676   375-405 (2000).
677   29   Brunton, S. L., Proctor, J. L. & Kutz, J. N. Discovering governing equations from data by sparse
678   identification of nonlinear dynamical systems. *Proceedings of the National Academy of Sciences*
679   **113**, 3932-3937 (2016).
680   30   Brunton, S. L. & Kutz, J. N. *Data-driven science and engineering: Machine learning, dynamical*
681   *systems, and control*. (Cambridge University Press, 2019).
682   31   Mangan, N. M., Askham, T., Brunton, S. L., Kutz, J. N. & Proctor, J. L. Model selection for hybrid
683   dynamical systems via sparse regression. *Proceedings of the Royal Society A* **475**, 20180534
684   (2019).
685   32   Mangan, N. M., Kutz, J. N., Brunton, S. L. & Proctor, J. L. Model selection for dynamical systems
686   via sparse regression and information criteria. *Proceedings of the Royal Society A: Mathematical,*
687   *Physical and Engineering Sciences* **473**, 20170009 (2017).
688   33   Anderson, D. & Burnham, K. Model selection and multi-model inference. *Second. NY: Springer-*
689   *Verlag* **63** (2004).
690   34   Rosenberg, M. C., Banjanin, B. S., Burden, S. A. & Steele, K. M. Predicting walking response to
691   ankle exoskeletons using data-driven models. *Journal of the Royal Society Interface* **17**,
692   20200487 (2020).
693   35   Genthe, K. *et al.* Effects of real-time gait biofeedback on paretic propulsion and gait
694   biomechanics in individuals post-stroke. *Topics in stroke rehabilitation* **25**, 186-193 (2018).
695   36   Delp, S. L. *et al.* OpenSim: open-source software to create and analyze dynamic simulations of
696   movement. *IEEE transactions on biomedical engineering* **54**, 1940-1950 (2007).
697   37   Rajagopal, A. *et al.* Full-body musculoskeletal model for muscle-driven simulation of human gait.
698   *IEEE transactions on biomedical engineering* **63**, 2068-2079 (2016).
699   38   Abdullah, F., Wu, Z. & Christofides, P. D. Handling noisy data in sparse model identification using
700   subsampling and co-teaching. *Computers & Chemical Engineering* **157**, 107628 (2022).

39   Villarreal, D. J., Poonawala, H. A. & Gregg, R. D. A robust parameterization of human gait
patterns across phase-shifting perturbations. *IEEE Transactions on Neural Systems and*
Rehabilitation Engineering **25**, 265-278 (2016).
40   Whittle, M. W. Clinical gait analysis: A review. *Human movement science* **15**, 369-387 (1996).
41   Akaike, H. in *Selected Papers of Hirotugu Akaike*   215-222 (Springer, 1974).
42   Dingwell, J. B. & Cusumano, J. P. Identifying stride-to-stride control strategies in human
treadmill walking. *PloS one* **10**, e0124879 (2015).
43   Dingwell, J. B. & Cusumano, J. P. Humans use multi-objective control to regulate lateral foot
placement when walking. *PLoS computational biology* **15**, e1006850 (2019).
44   Maus, H.-M. *et al.* Constructing predictive models of human running. *Journal of The Royal*
Society Interface **12**, 20140899 (2015).
45   Glantz, S.   (New York: McGraw-Hill, 2012).
46   Mochon, S. & McMahon, T. A. Ballistic walking. *Journal of biomechanics* **13**, 49-57 (1980).
47   Lee, C. R. & Farley, C. T. Determinants of the center of mass trajectory in human walking and
running. *Journal of experimental biology* **201**, 2935-2944 (1998).
48   Nott, C., Neptune, R. & Kautz, S. Relationships between frontal-plane angular momentum and
clinical balance measures during post-stroke hemiparetic walking. *Gait & posture* **39**, 129-134
(2014).
49   Vistamehr, A., Kautz, S. A. & Neptune, R. R. The influence of solid ankle-foot-orthoses on
forward propulsion and dynamic balance in healthy adults during walking. *Clinical biomechanics*
29, 583-589 (2014).
50   Ebers, M. R., Rosenberg, M. C., Kutz, J. N. & Steele, K. M. A machine learning approach to
quantify individual gait responses to ankle exoskeletons. *Journal of Biomechanics*, 111695
(2023).
51   Clark, D. J., Ting, L. H., Zajac, F. E., Neptune, R. R. & Kautz, S. A. Merging of Healthy Motor
Modules Predicts Reduced Locomotor Performance and Muscle Coordination Complexity Post-
Stroke. *Journal of Neurophysiology* **103**, 844-857, doi:10.1152/jn.00825.2009 (2010).
52   Allen, J. L., Kautz, S. A. & Neptune, R. R. The influence of merged muscle excitation modules on
post-stroke hemiparetic walking performance. *Clinical Biomechanics* **28**, 697-704 (2013).
53   Steele, K. M., Jackson, R. W., Shuman, B. R. & Collins, S. H. Muscle recruitment and coordination
with an ankle exoskeleton. *Journal of biomechanics* **59**, 50-58 (2017).
54   Novelli, N., Lenci, S. & Belardinelli, P. Boosting the Model Discovery of Hybrid Dynamical Systems
in an Informed Sparse Regression Approach. *Journal of Computational and Nonlinear Dynamics*
17, doi:10.1115/1.4053324 (2022).
55   Lee, D. D. & Seung, H. S. Learning the parts of objects by non-negative matrix factorization.
Nature **401**, 788 (1999).
56   Ivanenko, Y. P., Cappellini, G., Dominici, N., Poppele, R. E. & Lacquaniti, F. Coordination of
locomotion with voluntary movements in humans. *Journal of Neuroscience* **25**, 7238-7253
(2005).
57   Shamaei, K., Sawicki, G. S. & Dollar, A. M. Estimation of quasi-stiffness and propulsive work of
the human ankle in the stance phase of walking. *PloS one* **8**, e59935 (2013).
58   Chen, Y.-M. & Posa, M. in *2020 IEEE International Conference on Robotics and Automation*
(ICRA). 8753-8760 (IEEE).
59   Whittingham, M. J., Stephens, P. A., Bradbury, R. B. & Freckleton, R. P. Why do we still use
stepwise modelling in ecology and behaviour? *Journal of animal ecology* **75**, 1182-1189 (2006).
60   Anderson, D. R. & Burnham, K. P. Avoiding pitfalls when using information-theoretic methods.
The Journal of wildlife management, 912-918 (2002).

61   Seethapathi, N., Clark, B. & Srinivasan, M. Exploration-based learning of a step to step controller
749   predicts locomotor adaptation. *bioRxiv*, 2021.2003.2018.435986,
750   doi:10.1101/2021.03.18.435986 (2021).

751