LAG3 regulates antibody responses in a murine model of kidney transplantation.

Michael Nicosia[1*], Ran Fan[1], Juyeun Lee[2], Victoria Gorbacheva[1], José I. Valenzuela[1], Yosuke Yamamoto[1], Ashley Beavers[1], Nina Dvorina[1], Eduardo Chuluyan[3,4], Motoo Araki[5], William M. Baldwin III[1], Robert L. Fairchild[1], Booki Min[1,6#] & Anna Valujskikh[1#]

1 - Department of Inflammation and Immunity, Lerner Research Institute, Cleveland Clinic, Ohio

2 - Department of Cardiovascular and Metabolic Sciences, Lerner Research Institute, Cleveland Clinic, Ohio

3 - Universidad de Buenos Aires, Consejo Nacional de Investigaciones Científicas y Técnicas, Centro de Estudios Farmacológicos y Botánicos (CEFYBO), Facultad de Medicina, Buenos Aires, Argentina

4 - Universidad de Buenos Aires, Facultad de Medicina, Departamento de Microbiología, Parasitología e Immunología, Buenos Aires, Argentina

5 - Department of Urology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

6 - Current - Department of Microbiology and Immunology, Feinberg School of Medicine, Northwestern University, Chicago, Illinois

* - Corresponding author

 # - Contributed equally

Correspondence: Michael Nicosia, Department of Inflammation and Immunity, Lerner Research Institute, Cleveland Clinic, 9500 Euclid Avenue, NB3-77, Cleveland, OH, 44195, USA. Email address: nicosim@ccf.org. Telephone:

26   Conflict of Interest Statement: The authors have declared that no conflict of interest exists.

Abstract

Lymphocyte activation gene-3 (LAG3) is a coinhibitory receptor expressed by a range of

immune cells. While immunomodulatory potential of LAG3 is being actively explored in cancer

and autoimmunity fields, there is no information on how this pathway affects alloreactive

immune responses following organ transplantation. The goal of this study was to investigate the

functions of recipient LAG3 in a mouse model of renal allograft rejection. We found that mice

deficient in LAG3 expression have elevated heterologous immunity against a panel of

alloantigens prior to transplantation. Recipient LAG3 deficiency results in rapid rejection of

MHC-mismatched renal allografts that are spontaneously accepted by WT recipients, with graft

histology characteristic of antibody mediated rejection (ABMR). Depletion of recipient B cells but

not CD8[+] T cells significantly extended kidney allograft survival in LAG3[-/-] recipients further

supporting ABMR as the main mechanism of graft loss. Treatment of WT recipients with an

antagonistic LAG3 antibody enhanced anti-donor immune responses and induced kidney

damage associated with chronic rejection. The experiments using conditional LAG3 knockout

recipients demonstrated that LAG3 expression on either T or B cells is sufficient to regulate anti-

donor humoral immunity. These results are the first to identify LAG3 as a regulator of both T

and B cell responses to kidney allografts and a potential therapeutic target for ABMR prevention

and treatment.

Introduction

Lymphocyte activation gene 3 (LAG3, CD223) is an immune checkpoint inhibitor of the

immunoglobulin superfamily, closely related to CD4[1]. LAG3 is expressed on a number of

immune cells including CD4[+] and CD8[+] T cells, T regulatory cells, B lymphocytes[1-6],

plasmacytoid dendritic cells[7] (pDCs) and NK cells[1]. Similar to other coinhibitory receptors such

as CTLA4 or PD-1, LAG3 is not expressed by naïve T cells, but its expression is induced on

CD4[+] and CD8[+] T cells following TCR engagement[1]. Murine LAG3 shares 69.9% sequence

homology with the human protein preserving structure and function, and largely mirrors the

expression profile of human LAG3[8,9]. Analogous to CD4, LAG3 binds to major histocompatibility

complex II molecules (MHC-II) on antigen presenting cells (APCs)[10,11], and additionally interacts

with other ligands such as Galectin-3[12], liver sinusoidal endothelial cell lectin (LSECtin)[13], α-

synuclein fibrils (α-syn)[14], and fibrinogen-like protein 1 (FGL1)[15].

In T cells, LAG3 is co-localized with T cell receptor/CD3 complex within immune synapse and

inhibits TCR signaling leading to impaired cell activation, proliferation, differentiation, and

effector functions in the context of autoimmunity, infection and tumor[16-22], although the

underlying mechanisms remain unclear. It was recently reported that, acidification via a

conserved tandem glutamic acid-proline repeat in the LAG3 cytoplasmic tail lowers the local pH

at the immune synapse, and dissociates Lck from the CD4 or CD8 co-receptor, resulting in a

loss of co-receptor-TCR signaling and limited T cell activation[23]. In addition, LAG3 plays an

essential role in T regulatory cells (Treg)[6,24,25], and mice with Treg-specific LAG3 deficiency are

resistant to autoimmune diabetes[26]. In contrast, there is a paucity of studies investigating the

role of LAG3 in B lymphocytes. Activation of murine splenocytes *in vitro* induces the

endogenous expression of LAG3 on CD19[+] B cells in T cell and soluble factor dependent

manner [3]. However, the functional consequences of LAG3 up-regulation in B cells during

humoral immune responses remain to be investigated. LAG3 expression has been shown to be

a defining feature of an IL-10 producing subset of plasma cells[27]. While it was demonstrated that

these cells arise from BCR dependent signals and that their IL-10 production is dependent upon

innate signals such as TLRs, the role of LAG3 expression on this B cell subset is still poorly

understood.

Even though the relative importance of LAG3 on conventional T cells, Tregs, B cells and other

cell types during immune responses still remains to be determined in different settings it is a

molecule of great interest for clinical interventions. A number of monoclonal antibodies targeting

LAG3 with either antagonistic, depleting, or agonistic activities have been developed and are

currently in clinical trials for cancer and autoimmunity patients[28-37]. However, LAG3 is not

commonly considered as a therapeutic target in organ transplantation, largely due to the paucity

of information on the role of this pathway in experimental animal models. Lucas *et al.*[38] reported

that LAG3 blockade prevents tolerance induction in alloreactive CD8[+] T cells in a mouse model

of allogeneic bone marrow transplantation, but that this effect did not depend on LAG3

expression by CD8[+] T lymphocytes themselves. In a different study, the depletion of LAG3[+] T

cells with monoclonal antibody extended rat cardiac allograft survival, yet prevented tolerance

induction via donor-specific cell transfusion[39]. These findings suggest LAG3 involvement in

different aspects of allograft rejection and tolerance. Whereas the functions of CTLA4 and PD1

coinhibitory molecules in transplantation have been extensively studied, it is unclear whether

and how LAG3 regulates alloimmune responses to solid organ transplants.

In the current study, we investigated the contribution of LAG3 to alloimmune responses using a

mouse model of life-supporting renal transplantation and found that non-transplanted mice

deficient in LAG3 expression have elevated cellular and humoral heterologous immunity against

a panel of alloantigens. Recipient LAG3 deficiency resulted in rapid rejection of fully MHC-

mismatched renal allografts that are spontaneously accepted by wild type recipients. While

LAG3[-/-] recipients developed increased cellular and humoral anti-donor immune responses, the

renal graft tissue injury was characteristic of antibody-mediated rejection and was significantly

diminished by the depletion of B cells but not of CD8[+] T lymphocytes. Neither T cell- or B-cell

specific LAG3 conditional knockout recipients rejected renal allografts indicating that LAG3

deficiency in both cell types is required for dysregulated alloantibody production in the context of

transplantation. These data present the first evidence for LAG3 involvement in regulation of

humoral alloimmune responses and identify LAG3 as a potential target for diminishing

pathogenic alloimmunity.

Materials and Methods

Animals

Male C3H/HeJ (C3H, H-2[k]), C57BL/6J (B6, H-2[b]), BALB/cJ (Balb/c, H-2[d]) DBA1/J (DBA H-2[q])

and SJL/J-*Pde6b[rd1]* (SJL, H-2[s]), B6.129P2(C)-Cd19 tm1(cre)Cgn/J (CD19Cre), B6.LAG3[-/-] aged

6-8 week, were purchased from The Jackson Laboratory (Bar Harbor,ME). LAG3[fl/fl] mice were

commissioned by Biocytogen (Beijing, China) and bred to CD4Cre or CD19Cre mice to

generate T cell or B cell conditional LAG3[-/-] mice respectively. All the mouse protocols were

approved by the Institutional Animal Care and Use Committee (IACUC) of the Cleveland Clinic.

The mice were maintained under specific pathogen-free conditions at the Cleveland Clinic

Lerner Research Institute Biological Resource Institute.

Kidney Transplantation and Graft Evaluation

Transplantation of murine kidneys was performed as previously described[40]. In brief, the kidney

and its vascular supply and ureter were harvested *en bloc*, the donor vasculature were

anastomosed to the recipient abdominal aorta and inferior vena cava. Excised kidneys were

perfused with University of Wisconsin (UW) solution (320mOsm; Preservation Solutions, Elkorn,

WI) and stored on ice for 0.5 hours before transplantation. Urinary reconstruction was

performed as previously published[41]. The remaining native kidney was removed at the time of

transplant, ensuring that survival of the mouse was dependent upon functionality of the

transplanted kidney. Graft survival was assessed by daily examination of overall animal health.

Recipients were euthanized when low and slow mobility in addition to a hunched posture

indicated rejection. At time of euthanasia serum creatinine levels were measured using the

VetScan i-STAT1 Analyzer (Abaxis, Union City, CA). Grafts were collected and fixed in

methanol before embedding in paraffin for staining with hematoxylin and eosin or Gomori's

trichrome C, and the following antibodies: rabbit monoclonal anti-mouse/human Mac-2 (clone

M3/M8; Biolegend #125401), rabbit monoclonal anti-mouse CD4 (clone EPR19514; Abcam

#ab183685), rabbit monoclonal anti-mouse CD8 (clone EPR21769; Abcam #ab217344),

polyclonal rabbit anti-mouse C4d[42], rabbit polyclonal anti-Human/mouse vWf (Dako #A0082), rat

monoclonal anti-mouse/rat FOXP3 (clone FJK-16s; eBioscience #14-5773) and rat monoclonal

anti-mouse B220/CD45R (clone RA3-682; BD Pharmingen #550286). Each stain was evaluated

based on two to three complete cross sections from 4-5 grafts per group. Blood urea nitrogen

(BUN) levels were measured using BUN Colorimetric Detection Kit (Thermo Fisher Scientific,

Waltham, MA). Urine levels of TIM-1/KIM-1/HAVCR and Lipocalin-2/NGAL were measured with

Quantikine ELISA kits (R@D System Inc., Minneapolis, MN) as previously published

(37084848).

Recipient Treatment

To deplete CD8$^+$ T cells recipients were treated with monoclonal anti-mouse CD8 antibodies,

clones TIB105 and YTS169 (Bio X Cell, 0.2mg of each i.p. on days -3, -2, -1 prior to

transplantation and every 5 days posttransplant for the duration of the experiment). For B cell

depletion, recipients were treated with anti-mouse CD19 clone 1D3 and anti-mouse B220 clone

RA3.3A1/6.1 (TIB-146) monoclonal antibodies Bio X Cell, 0.2mg of each i.p. on day 3 and every

5 days posttransplant for the duration of the experiment). For LAG3 blockade, anti-LAG-3 mAb

(clone C9B7w, Bio X Cell)[9,38] was given i.p. at a dose of 100 µg on days 3, 5, 7, 9, 11, 13 and

15 posttransplant.

Measurement of Serum Alloantibody

Alloreactive serum antibody titers were measures as previously published[43]. In brief, serum samples were collected via tail-vein bleeding from naïve 10 weeks old mice or from kidney allograft recipients at d. 14 posttransplant and stored at -20°C. Flat-bottomed 96-well plates (Thermo Fisher Scientific) were coated overnight at 4°C with biotinylated class 1 D^d (folded with RPGGRAFVTI peptide), class 1 D^k (folded with RRLGRTLLL peptide), class 2 I-A^d (folded with PVSKMRMATPLLMQA class 2-associated invariant chain peptide), or class 2 I-A^k (folded with PVSKMRMATPLLMQA class 2- associated invariant chain peptide) MHC monomers at 1μg/ml in PBS. Peptide/MHC monomers were provided by the National Institutes of Health Tetramer Core Facility at Emory University (Atlanta, GA). Plates were washed with PBS and blockade with 1% BSA-PBS solution for 1h at room temperature. A total of 100μl of serum diluted from 1:100-1:500 were added to the plates and incubated overnight at 4°C. Plates were washed with PBS and PBS-Tween (0.25%), and then incubated with goat anti-mouse IgG -horse radish peroxidase (HRP) conjugate (1:20000 dilution; Thermo Fisher Scientific) or IgG1, IgG2a, IgG2b, IgG3-horse radish peroxidase (HRP) conjugate (1:5000 dilution; Thermo Fisher Scientific) for 1 hour at room temperature. After incubation with conjugate-bound antibodies, plates were washed and developed with ABTS Peroxidase Substrate (KPL, Gaithersburgh, MD). Absorbance was measured at 415nm using the Bio-Rad iMark Microplate Reader (Bio-Rad Laboratories, Hercules, CA).

ELISPOT Assay

Interferon gamma (IFNγ) ELISPOT assays were performed as previously described using capture and detecting antibody from BD Pharmingen[44]. Spleens from recipients or naïve non-transplanted mice were stimulated with mitomycin C-treated donor C3H or BALB/c, SJL or DBA spleen cells for 24 hours. The resulting spots were analyzed using an ImmunoSpot Series 4 Analyzer (Cellular Technology, Cleveland, OH).

Flow Cytometry

Fluorochrome conjugated antibodies were purchased from BD Biosciences (San Jose, CA) or

eBioscience (San Diego, CA). Cells were isolated from the spleen and stained as previously

described[45], ≥100,000 events per sample were acquired on a BD Biosciences LSRII or a BD

Biosciences LSRFortessa X-20 followed by data analysis using FlowJo software (TreeStar, Inc.,

Ashland, OR).

Statistical Analyses

Kidney allograft survival was compared between groups by Kaplan-Meier analysis. All other

results were analyzed by using a parametric unpaired t-test. The difference between groups

was considered significant if the p value was <0.05. Unless noted otherwise, the data are

presented as mean□±□SD values. Total numbers of animals in each experimental group are

indicated in respective figure legends.

Results

Mice deficient in LAG3 have elevated heterologous alloimmune responses

prior to transplantation.

As LAG3 was demonstrated to play critical role in immune tolerance, we initially evaluated the

immune cells from naïve non-transplanted 10 week old B6.WT and B6.LAG3$^{-/-}$ mice. LAG3

deficiency resulted in modest increase in splenic cellularity (data not shown) and in the

increased numbers of total and effector memory (TEff$_M$) CD4$^+$ and CD8$^+$ T cells, as well as

Tregs compared to WT mice. In addition, mice deficient in LAG3 had elevated numbers of T

follicular helper (Tfh) and T follicular regulatory (Tfr) cells in the spleen (**Fig. 1A**, **Fig. S1**). The

numbers of major spleen B cell subsets – follicular (FoB), marginal zone (MzB), and transitional

(TrB) B cells – were not significantly different between WT and LAG3$^{-/-}$ mice. Notably, mice

deficient in LAG3 had significantly increased numbers of germinal center B cells (GcB) and plasma cells (PC) in the spleen suggesting ongoing B cell activation (**Fig. 1B**, **Fig. S2**). There was a trend towards the increase in numbers of regulatory B cells (Bregs) defined as CD19$^+$CD1d^{hi}CD5$^+$ cells. It should be noted that despite the observed shifts towards activated immune cell phenotypes, unmanipulated LAG3$^{-/-}$ mice did not exhibit signs of autoimmune disease up to 6 months of age.

To assess the impact of this phenotypic shift on alloimmunity, we probed both B6.WT and B6.LAG3$^{-/-}$ for T cell reactivity against a panel of allogeneic strains (**Fig. 2A**). IFNγ ELISPOT assay demonstrated that compared to WT mice, naïve LAG3$^{-/-}$ mice have increased frequencies of preexisting memory T cells reactive against BALB/c (H2-D^d), C3H (H2-D^k), and to a lesser extent, SJL (H-2D^s) and DBA (H2-D^q) alloantigens (**Fig. 2A**). We also tested for the presence of alloreactive antibodies in the sera of non-transplanted mice (**Fig. 2B**). Some of the tested naïve LAG3$^{-/-}$ (H2-D^b) contained elevated levels of IgG antibodies against class I and class II MHC molecules from BALB/c (H2-D^d, I-A^d) and C3H mice (H2-D^k, I-A^k). These results suggest that LAG3 is an important regulator of immune cell homeostasis and as such may play a crucial role during immune responses to transplanted organs.

Recipient LAG3 deficiency enhances anti-donor alloimmune responses and elicits kidney allograft rejection

To test the functional consequences of LAG3 deletion, we used a previously described mouse model of kidney transplantation in which B6 (H-2^b) recipients undergo bilateral nephrectomy, and are then transplanted with a single fully MHC-mismatched kidney allograft[43]. In this model the recipient survival is dependent upon allograft functionality, which can be measured by serum creatinine levels. Consistent with previously published studies that used different donor

strains[43,46], B6.WT recipients spontaneously accept C3H renal allografts for >60 days (**Fig.**

3A&B). In contrast, most B6.LAG3$^{-/-}$ recipients rapidly rejected kidney allografts (median

survival time, MST, of 15d) (**Fig. 3B**) and had significantly increased serum creatinine levels at

d. 14 (1.3 mg/dl versus 0.1 mg/dl in B6.WT controls and <0.4 mg/dl in non-transplanted mice,

Fig. 3C). Immunohistochemistry of allografts recovered from LAG3$^{-/-}$ recipients around the time

of rejection (d. 14) showed decreased numbers of infiltrating T cells compared to grafts from

B6.WT recipients (**Fig. S3**). Instead, the rejecting grafts showed characteristic signs of antibody

mediated rejection including diffuse C4d staining, atrophic peritubular capillaries, endothelial cell

swelling and edema (**Fig. 3D and Fig. S3**). Flow cytometry analysis of graft infiltrating cells on d.

10 posttransplant (**Fig. 3E-F**) revealed no significant changes in CD4^{+} or CD8^{+} T cell infiltration,

with the only significant difference being in the increased numbers of infiltrating FoxP3^{+} T

regulatory cells. LAG3$^{-/-}$ recipients did demonstrate a modest increase in the infiltration of NK

cells, the majority of which were of the NK1.1^{hi} phenotype (**Fig. 3E-F** and data not shown). This

finding is consistent with antibody-mediated tissue injury of renal allografts as recently

demonstrated by Yagisawa and colleagues[47].

The analysis of the peripheral T cell pool revealed no significant differences in the proportion or

numbers of various spleen T cell subsets (**Fig. 4A** and **Fig. S4A**). Nevertheless, the absence of

recipient LAG3 resulted in elevated frequencies of donor specific T cells, as measured by IFNγ

ELIPSOT assay on d. 14 posttransplant (**Fig. 4C**). Analogous to the T cell compartment, spleen

subsets of follicular, marginal zone, transitional, and regulatory B cells were similar in WT and

LAG3$^{-/-}$ allograft recipients (**Fig. 4B** and **Fig. S4B**), whereas germinal center B cells and plasma

cells were increased in some but not all recipients. To assess anti-donor humoral immune

responses, recipient serum was collected at 14 d posttransplant and analyzed for the levels of

IgG antibodies against donor class I (D^{k}) and class II (I-A^{k}) MHC molecules. Consistent with our

published studies, WT renal allograft recipients had developed minimal donor specific

alloantbodies (DSA) at this timepoint. In contrast, LAG3$^{-/-}$ recipients had significantly elevated

levels of IgG against both D^k and I-A^k (**Fig. 4D**). Taken together these data demonstrate that

LAG3 plays an essential role in regulating both cellular and humoral immune responses to

transplanted allografts and contributes to spontaneous renal allografts acceptance in wild type

recipients.

B cells, but not CD8$^+$T cells, are essential for kidney allograft rejection by

LAG3$^{-/-}$ recipients.

Given that LAG3 is a well-studied regulator of T cell responses we anticipated that LAG3

deficiency primarily affects priming of alloreactive T cells and their pathogenic functions within

the kidney graft. However, the modest increase in donor-reactive T cells and the graft histology

(**Figs. 3-4 and Fig. S3)** suggested that alloantibody are the major mediators of graft tissue

injury in LAG3$^{-/-}$ recipients. To formally test the contributions of cellular vs antibody mediated

rejection, LAG3$^{-/-}$ recipients were treated with CD8$^+$ T cell depleting antibody prior to

transplantation of C3H kidney allografts (**Fig. 5A** and **Fig. S6B**). CD8$^+$ T cell depletion

antibodies failed to prolong kidney allograft survival in LAG3$^{-/-}$ recipients, with MST of 16 days

(**Fig. 5B**). Anti-donor humoral immune responses were assessed at 14 d posttransplant and

analyzed for the levels of IgG antibodies against donor class I and class II MHC molecules. WT

allograft recipients depleted of CD8 T cells had a low-grade DSA response at this timepoint. In

contrast, LAG3$^{-/-}$ recipients had significantly elevated levels of IgG against both donor class I

and class II alloantigens (**Fig. 5C**). Although CD8 T cell depletion reduced the frequencies of

IFNγ producing T cells in both groups, LAG3 deficient recipients still had modestly elevated

frequencies of donor-specific T cells on d. 14 posttransplant compared to WT (**Fig. 5D**). This is

likely due to a faster reconstitution of LAG3$^{-/-}$ cells following depletion, which has been

previously reported in memory CD4$^+$ T cells[48]. Nevertheless, histological analysis at the time of

rejection showed that grafts from CD8$^+$ T cell depleted LAG3$^{-/-}$ recipients had significant damage

to the kidney tubules such as tubular dilation, casts, tubular atrophy, and edema. C4d staining

of these grafts showed dilated capillaries, endothelial cell swelling, and heavy damage to the

tubules and to the glomeruli (**Fig. 5E**). In summary, the grafts rejected by LAG3$^{-/-}$ recipients

following CD8$^+$ T cell depletion displayed the typical features of antibody mediated graft damage

similar to or even exceeding those observed in non-depleted LAG3-deficient recipients.

To test the roles of B cells and antibodies in the observed rejection, LAG3$^{-/-}$ recipients of C3H

kidney allografts were treated with anti-mouse CD19 and B220 B cell depleting antibodies

starting on d. 3 posttransplant and throughout the experiment (**Fig. 6A** and **Fig. S6A**).

Remarkably, B cell depletion restored graft survival in the majority of LAG3$^{-/-}$ recipients (**Fig. 6B**).

Depletion of recipient B cells lead to the abrogation of the DSA responses in both WT and

LAG3$^{-/-}$ recipients (**Fig. 6C**). While B cell depletion reduced T cell alloresponses in both groups,

LAG3$^{-/-}$ recipients still had increased frequencies of donor-reactive IFNγ producing T cells

compared to WT (**Figs. 6D and 4C**). Histological analysis at d. 30 posttransplant showed that

recipient B cell depletion reduced allograft damage with marked decrease in T cell infiltrates and

antibody binding, demonstrated by the absence of C4d staining (**Fig. 6E**). Together, these

results demonstrate that the rejection observed in LAG3$^{-/-}$ mice is driven by B cell production of

DSA and not by cytotoxic T cells, and that the generated DSA is a major effector mechanism of

graft injury.

LAG3 is expressed by regulatory B cells and plasma cells in kidney

allograft recipients.

As our data revealed that B cell responses are affected by the loss of LAG3 mediated immune

regulation, we next analyzed LAG3 expression in different B cell subsets following

transplantation. LAG3 expression was not detectable by flow cytometry in B cells from naïve

mice (data not shown) consistent with previous reports that LAG3 expression on B cells is only

inducible in a T cell dependent manner[3]. Given the observed rejection kinetics (**Fig. 3B & 5B**),

we measured LAG3 expression on splenic B cell subsets on d10 posttransplant, using LAG3$^{-/-}$

recipients as a control. Of the cell populations tested, only 16.28% ($\pm$5.227 SD)

CD19$^+$CD1d^{hi}CD5$^+$ Bregs and 8.08% ($\pm$1.979 SD) CD138$^+$ plasma cells expressed cell surface

LAG3 (**Fig. 7A & B** and **Fig. S5**), suggesting differential expression in specialized subsets.

LAG3 deficiency on both T and B lymphocytes is required for kidney

allograft rejection.

The absence of LAG3 on T cells often induces pathogenesis in other mouse models [15]. To

address whether increased T cell help or defective T regulatory cell function was a major driver

of the observed allograft injury, we used T cell conditional knockout mice and littermate controls

as renal allograft recipients (**Fig. 8A**). B6.CD4Cre$^{+/-}$LAG3$^{fl/fl}$ which lack LAG3 on CD4$^+$ and

CD8$^+$ T cells spontaneously accepted kidney allografts for > 60 days (**Fig. 8B**) indicating that

LAG3 deficiency on T cells alone was not sufficient to induce rejection. Analysis of the immune

response in these recipients showed elevated DSA responses (**Figs. 8C**), but no significant

changes in the frequencies of IFNγ producing alloreactive T cells (**Fig. 8D**). Histological analysis

of the allografts at four weeks posttransplant revealed no major impact on fibrosis or graft tissue

injury Interestingly, despite the presence of serum DSA, T cell LAG3 deficiency resulted in

reduced C4d deposition within interstitial capillaries compared to LAG3$^{-/-}$ recipients (**Figs. 8E**

and 3D).

We next investigated the role of LAG3 using a B cell conditional knockout recipient model (**Fig.**

9A). Unexpectedly, B6.CD19Cre[+/-]LAG3[fl/fl] recipients with specific LAG3 deficiency in B

lymphocytes also failed to reject kidney allografts by d. 30 posttransplant (**Fig. 9B**). LAG3

deficiency on B cells of renal transplant recipients resulted in a modest increase in DSA levels

by d14 posttransplant (**Fig. 9C**). In contrast, LAG3 deficiency on B cells had no impact on the

frequencies of donor reactive T cells at d14 posttransplant (**Fig. 9D**), indicating that B cell LAG3

expression does not affect T cell priming. Graft histology at four weeks posttransplant showed

no significant differences in T cell infiltration. C4d staining was less intense in the CD19Cre[+/-]

LAG3[fl/fl] recipients compared to the littermate controls but showed a rarefaction of peritubular

capillaries, and endothelial cell swelling consistent with antibody mediated graft injury (**Fig. 9E**

and **Fig. S7**)[49]. Taken together these data show that LAG3 expression on either helper T cells

or B cells is sufficient to induce spontaneous acceptance of murine kidney allografts.

LAG3 regulates de novo alloresponses.

Our data show that non-transplanted LAG3 deficient mice have increased alloreactive immune

responses (**Fig. 2A&B**). To establish whether LAG3 is regulating *de novo* immune responses,

we treated WT kidney allograft recipients with a course of LAG3 blocking antibody and

monitored mice for signs of rejection (**Fig. 10A**). Blockade of LAG3 resulted in rejection in 2 out

9 mice within 14 days posttransplant (**Fig. 10B**). Despite the long term graft survival in the

remaining recipients, we found that LAG3 blockade did induce graft damage, as measured by

kidney injury markers NGAL and KIM1 in the urine, and blood urea nitrogen (BUN) in the serum

(**Fig. 10C-E**). Blockade of LAG3 also resulted in increased DSA production (**Fig. 10F&G**) and

increased frequencies of donor reactive T cells (**Fig. 10H**), indicating that LAG3 regulates de

novo alloresponses following transplantation. The increase in markers of kidney injury and

alloresponses suggested that LAG3 blockade may induce chronic graft injury. Indeed, histological analysis at d. 42 posttransplant (**Fig. 10I & Fig. S8**) revealed tubular atrophy, endothelial cell swelling and increased alpha-smooth muscle actin (aSMA) staining due to graft fibrosis, also reflected in the Trichrome staining (**Fig. S8**). C4d staining showed C4d deposition in both control IgG treated and anti-LAG3 treated recipients, but recipients that received LAG3 blockade had dilated peritubular capillaries with marginated mononuclear cells (red arrows, **Fig. 10I**). In addition, recipients that received LAG3 blockade had increased CD4$^+$ and CD8$^+$ T cell graft infiltrates. These findings identify LAG3 as an important regulator of *de novo* immune responses to a kidney allograft.

Discussion

Coinhibitory molecule LAG3 is best studied in regulating effector and regulatory T cell functions in autoimmunity, infection and cancer[17-22]. However, little is known about the contribution of this pathway in regulating humoral immune responses. Our results definitively demonstrate that LAG3 regulates alloantibody generation in response to solid organ transplantation, and suggest that LAG3 expression on both T and B lymphocytes plays a role in this process.

The absence of LAG3 in T cells leads to enhanced effector T cell responses and increased formation of T cell memory[26]. Consistent with this, naïve, non-transplanted LAG3-deficient mice have elevated levels of CD44^{hi} memory T cells and enhanced response to a panel of alloantigens compared to wild type counterparts (**Fig. 1 & Fig. 2**). In addition, we observed increased numbers of germinal center B cells and CD138$^+$ plasma cells in naïve 10 week old LAG3$^{-/-}$ mice, which suggested ongoing B cell activation. Based on previous studies, we initially proposed that LAG3 deletion in the recipient will result in dysregulated alloimmunity

characterized by exaggerated activation of alloreactive T cells and leading to T cell-mediated

rejection.

The model of renal transplantation was chosen for these experiments as even fully MHC

mismatched mouse kidney allograft typically do not undergo acute rejection, and are

spontaneously accepted in many donor/recipients strain combinations[50]. While functional C3H

kidney allografts survive for > 60 days in B6.WT recipients, they are rapidly rejected by

B6.LAG3$^{-/-}$ mice (**Fig. 2**). Long term survival and function of mouse renal allografts have been

previously correlated with the numbers of graft-infiltrating regulatory T cells, as FoxP3$^+$

regulatory T cell depletion results in rapid graft rejection[51,52]. In contrast to germline LAG3$^{-/-}$

mice, T cell specific LAG3 deletion did not result in acute graft rejection (**Fig. 8**). LAG3 is highly

expressed by FoxP3$^+$ Tregs and LAG3 is thought to play a role in regulatory T cell's optimal

suppressor activity[6,53,54]. Previous studies demonstrated that FoxP3 regulatory T cell depletion

in mouse renal allograft recipients resulted in T cell mediated rejection without affecting DSA

generation[52]. Moreover, LAG3$^{-/-}$ mice have increased numbers of FoxP3$^+$ T cells before and

after transplantation (**Figs. 1-3**). However, it is unlikely that the augmented alloimmunity and

graft rejection in our model are entirely due to dysfunctional Tregs as specific deletion of LAG3

in recipient T cells (including Tregs) was not sufficient to induce rejection (**Fig. 8**). [52].

While LAG3 expression in B cells was reported in 2005[3], its contribution to their functions has

not been extensively studied. Our results demonstrate that the lack of recipient LAG3 leads to

increased titers of IgG DSA antibodies and allograft rejection with characteristic features of

antibody-mediated injury. Furthermore, depletion experiments confirmed that B cells are

required for acute rejection of kidney allografts (**Fig. 6**), whereas the rejection still occurred with

the same kinetics in CD8 T cells-depleted LAG3$^{-/-}$ recipients (**Fig. 5**). To date, there are only a

few studies addressing the role of LAG3 in humoral immune response. Butler and colleagues[55],

reported that the combination treatment with anti-PD-L1 and anti-LAG3 mAb during established

malaria enhances Tfh responses, plasma cell formation and protective antibody generation in a

mouse model of malaria. However, the individual contributions of PD-L1 and LAG3 blockade

were not evaluated in this study. Another recent study identified a novel subset of LAG3[+] IL-10

secreting plasma cells with regulatory properties has recently been reported[27]. However, the

relative importance of LAG3 on helper T cells vs B cells during the initiation of B cell activation,

germinal center formation and differentiation into antibody-secreting cells remains to be

determined and is the focus of ongoing studies in our laboratory.

Our study does not entirely rule out the contribution of LAG3 expressed by cells other than T

and B lymphocytes. NK cells express LAG3 and are important mediators of antibody-mediated

injury of renal allografts[47]. While the effects of LAG3 on various NK cell functions is highly

controversial (reviewed in[56]), it is possible that dysregulated NK cells mediate rejection in LAG3[-]

[/-] recipients, secondary to enhanced T cell activation and DSA generation. The absence of

LAG3 on recipient antigen presenting cells (APCs) may also contribute to elevated T cell

responses. Initial analyses of LAG3 expression in dendritic cell (DC) subsets showed high LAG3

expression by plasmacytoid but not lymphoid or myeloid conventional DCs[7]. However, a more

recent study demonstrated a role for LAG3 on bone marrow derived DCs in optimal T cell

priming[57]. Due to these controversies, the impact of LAG3 deficiency in recipient APCs needs to

be carefully dissected in future studies.

It is important to note that, treatment of WT B6 recipients with a LAG3 blocking antibody

induced a pathology consistent with chronic rejection (**Fig. 10**). This was evident in the

increased kidney injury markers in the urine and serum (**Fig. 10C-E**), increased serum DSA

levels (**Fig.10F&G**), and increased histological signs of injury (**Fig. 10I** and **Fig. S8**). There are

a paucity of small animal chronic injury/rejection models in kidney transplantation, with some of

the current models relying on repeated administration of anti-MHC-I antibodies[58], or repeat

transfers of anti-donor antibody containing sera[59,60]. Our findings thus identify a new clinically

relevant model for studying the mechanisms of chronic injury and graft rejection in renal

transplantation.

To our knowledge, our study is one of the first to address the role of LAG3 during alloimmune

responses. Lucas *et al.* reported that blocking LAG3 with mAb prevents tolerization of CD8 T

cells following allogeneic bone marrow transplantation in mice[38]. In an earlier study, the

investigators used depleting anti-LAG3 mAb as an induction therapy in a rat model of cardiac

transplantation. Interestingly, the treatment extended heart allograft survival, yet abrogated the

tolerogenic effects of donor specific cell transfusion, which the authors attributed to Treg

depletion[39].

Despite many gaps and controversies in LAG3 biology, it is a molecule of therapeutic interest,

particularly in the cancer field. Antagonistic antibodies such as Relatlimab are given to patients

in combination with anti-PD-1 therapy in phase II clinical trials[61,62]. Another promising reagent

undergoing phase III trials is a bispecific antibody against LAG3 and PD-1, MGD013[28]. Anti-

LAG3 depleting antibodies have been developed and validated in non-human primates to target

overactivated immune cells during undesired immune response[63]. An agonistic anti-LAG3

antibody (IMP761) is currently under development for T cell mediated autoimmunity, and this

approach could have potential for clinical transplantation[20].

In conclusion, our study demonstrates for the first time the importance of LAG3 in regulating

pathogenic humoral immune responses to a transplanted solid organ. The results provide

rationale for investigating LAG3 impact on B cell activation, survival and differentiation, and

suggest LAG3 as a potential target for future therapeutic interventions for the prevention and

treatment of antibody-mediated rejection.

456 Author Contributions

Design of research study: MN, JL, WMB, EC, MA, RLF, BM & AV

Conducting Experiments: MN, RF, JL, VG, JIV, AB, & ND

Acquiring data: MN, JL, VG, JIV, & AB

Analyzing data: MN, JL, VG, JIV, & WMB

Providing reagents/mice: BM

Writing the manuscript: MN, WB, BM, & AV

464 Acknowledgements

465   Models were created with Biorender.com. This work was supported by funding from the

466   American Society of Transplantation (gCDX-221C0MN - MN), and the National Institutes of

467   Health (R01AI165513 – WMB, R01AI-74740 – RLF, R01AI125247 – BM, R01AI60740 &

468   R01AI113142-AV).

References

1.   Triebel F, Jitsukawa S, Baixeras E, Roman-Roman S, Genevee C, Viegas-Pequignot E, Hercend T. LAG-3, a novel lymphocyte activation gene closely related to CD4. *J Exp Med*. 1990;171:1393-1405. doi: 10.1084/jem.171.5.1393

2.   Andrews LP, Marciscano AE, Drake CG, Vignali DA. LAG3 (CD223) as a cancer immunotherapy target. *Immunol Rev*. 2017;276:80-96. doi: 10.1111/imr.12519

3.   Kisielow M, Kisielow J, Capoferri-Sollami G, Karjalainen K. Expression of lymphocyte activation gene 3 (LAG-3) on B cells is induced by T cells. *Eur J Immunol*. 2005;35:2081-2088. doi: 10.1002/eji.200526090

4.   Durham NM, Nirschl CJ, Jackson CM, Elias J, Kochel CM, Anders RA, Drake CG. Lymphocyte Activation Gene 3 (LAG-3) modulates the ability of CD4 T-cells to be suppressed in vivo. *PLoS One*. 2014;9:e109080. doi: 10.1371/journal.pone.0109080

5.   Pena J, Jones NG, Bousheri S, Bangsberg DR, Cao H. Lymphocyte activation gene-3 expression defines a discrete subset of HIV-specific CD8+ T cells that is associated with lower viral load. *AIDS Res Hum Retroviruses*. 2014;30:535-541. doi: 10.1089/AID.2012.0195

6.   Huang CT, Workman CJ, Flies D, Pan X, Marson AL, Zhou G, Hipkiss EL, Ravi S, Kowalski J, Levitsky HI, et al. Role of LAG-3 in regulatory T cells. *Immunity*. 2004;21:503-513. doi: 10.1016/j.immuni.2004.08.010

7.   Workman CJ, Wang Y, El Kasmi KC, Pardoll DM, Murray PJ, Drake CG, Vignali DA. LAG-3 regulates plasmacytoid dendritic cell homeostasis. *J Immunol*. 2009;182:1885-1891. doi: 10.4049/jimmunol.0800185

8.   Mastrangeli R, Micangeli E, Donini S. Cloning of murine LAG-3 by magnetic bead bound homologous probes and PCR (gene-capture PCR). *Anal Biochem*. 1996;241:93-102. doi: 10.1006/abio.1996.0382

9.   Workman CJ, Rice DS, Dugger KJ, Kurschner C, Vignali DA. Phenotypic analysis of the murine CD4-related glycoprotein, CD223 (LAG-3). *Eur J Immunol*. 2002;32:2255-2263. doi: 10.1002/1521-4141(200208)32:8<2255::AID-IMMU2255>3.0.CO;2-A

10. Huard B, Mastrangeli R, Prigent P, Bruniquel D, Donini S, El-Tayar N, Maigret B, Dreano M, Triebel F. Characterization of the major histocompatibility complex class II binding site on LAG-3 protein. *Proc Natl Acad Sci U S A*. 1997;94:5744-5749. doi: 10.1073/pnas.94.11.5744

11. Baixeras E, Huard B, Miossec C, Jitsukawa S, Martin M, Hercend T, Auffray C, Triebel F, Piatier-Tonneau D. Characterization of the lymphocyte activation gene 3-encoded protein. A new ligand for human leukocyte antigen class II antigens. *J Exp Med*. 1992;176:327-337. doi: 10.1084/jem.176.2.327

12. Kouo T, Huang L, Pucsek AB, Cao M, Solt S, Armstrong T, Jaffee E. Galectin-3 Shapes Antitumor Immune Responses by Suppressing CD8+ T Cells via LAG-3 and Inhibiting Expansion of Plasmacytoid Dendritic Cells. *Cancer Immunol Res*. 2015;3:412-423. doi: 10.1158/2326-6066.CIR-14-0150

13. Xu F, Liu J, Liu D, Liu B, Wang M, Hu Z, Du X, Tang L, He F. LSECtin expressed on melanoma cells promotes tumor progression by inhibiting antitumor T-cell responses. *Cancer Res*. 2014;74:3418-3428. doi: 10.1158/0008-5472.CAN-13-2690

14.   Mao X, Ou MT, Karuppagounder SS, Kam TI, Yin X, Xiong Y, Ge P, Umanah GE,
Brahmachari S, Shin JH, et al. Pathological alpha-synuclein transmission initiated by
binding lymphocyte-activation gene 3. *Science*. 2016;353. doi: 10.1126/science.aah3374
15.   Wang J, Sanmamed MF, Datar I, Su TT, Ji L, Sun J, Chen L, Chen Y, Zhu G, Yin W, et
al. Fibrinogen-like Protein 1 Is a Major Immune Inhibitory Ligand of LAG-3. *Cell*.
2019;176:334-347 e312. doi: 10.1016/j.cell.2018.11.010
16.   Hannier S, Triebel F. The MHC class II ligand lymphocyte activation gene-3 is co-
distributed with CD8 and CD3-TCR molecules after their engagement by mAb or
peptide-MHC class I complexes. *Int Immunol*. 1999;11:1745-1752. doi:
10.1093/intimm/11.11.1745
17.   Maruhashi T, Sugiura D, Okazaki IM, Okazaki T. LAG-3: from molecular functions to
clinical applications. *J Immunother Cancer*. 2020;8. doi: 10.1136/jitc-2020-001014
18.   Jha V, Workman CJ, McGaha TL, Li L, Vas J, Vignali DA, Monestier M. Lymphocyte
Activation Gene-3 (LAG-3) negatively regulates environmentally-induced autoimmunity.
PLoS One. 2014;9:e104484. doi: 10.1371/journal.pone.0104484
19.   Kim D, Le HT, Nguyen QT, Kim S, Lee J, Min B. Cutting Edge: IL-27 Attenuates
Autoimmune Neuroinflammation via Regulatory T Cell/Lag3-Dependent but IL-10-
Independent Mechanisms In Vivo. *J Immunol*. 2019;202:1680-1685. doi:
10.4049/jimmunol.1800898
20.   Angin M, Brignone C, Triebel F. A LAG-3-Specific Agonist Antibody for the Treatment of
T Cell-Induced Autoimmune Diseases. *J Immunol*. 2020;204:810-818. doi:
10.4049/jimmunol.1900823
21.   Grosso JF, Kelleher CC, Harris TJ, Maris CH, Hipkiss EL, De Marzo A, Anders R, Netto
G, Getnet D, Bruno TC, et al. LAG-3 regulates CD8+ T cell accumulation and effector
function in murine self- and tumor-tolerance systems. *J Clin Invest*. 2007;117:3383-3392.
doi: 10.1172/JCI31184
22.   Goding SR, Wilson KA, Xie Y, Harris KM, Baxi A, Akpinarli A, Fulton A, Tamada K,
Strome SE, Antony PA. Restoring immune function of tumor-specific CD4+ T cells during
recurrence of melanoma. *J Immunol*. 2013;190:4899-4909. doi:
10.4049/jimmunol.1300271
23.   Guy C, Mitrea DM, Chou PC, Temirov J, Vignali KM, Liu X, Zhang H, Kriwacki R,
Bruchez MP, Watkins SC, et al. LAG3 associates with TCR-CD3 complexes and
suppresses signaling by driving co-receptor-Lck dissociation. *Nat Immunol*.
2022;23:757-767. doi: 10.1038/s41590-022-01176-4
24.   Gagliani N, Magnani CF, Huber S, Gianolini ME, Pala M, Licona-Limon P, Guo B,
Herbert DR, Bulfone A, Trentini F, et al. Coexpression of CD49b and LAG-3 identifies
human and mouse T regulatory type 1 cells. *Nat Med*. 2013;19:739-746. doi:
10.1038/nm.3179
25.   Liang B, Workman C, Lee J, Chew C, Dale BM, Colonna L, Flores M, Li N,
Schweighoffer E, Greenberg S, et al. Regulatory T cells inhibit dendritic cells by
lymphocyte activation gene-3 engagement of MHC class II. *J Immunol*. 2008;180:5916-
5926. doi: 10.4049/jimmunol.180.9.5916
26.   Zhang Q, Chikina M, Szymczak-Workman AL, Horne W, Kolls JK, Vignali KM, Normolle
D, Bettini M, Workman CJ, Vignali DAA. LAG3 limits regulatory T cell proliferation and
function in autoimmune diabetes. *Sci Immunol*. 2017;2. doi:
10.1126/sciimmunol.aah4569
27.   Lino AC, Dang VD, Lampropoulou V, Welle A, Joedicke J, Pohar J, Simon Q, Thalmensi
J, Baures A, Fluhler V, et al. LAG-3 Inhibitory Receptor Expression Identifies
Immunosuppressive Natural Regulatory Plasma Cells. *Immunity*. 2018;49:120-133 e129.
doi: 10.1016/j.immuni.2018.06.007

28. LaMotte-Mohs R, Shah K, Smith D, Gorlatov S, Ciccarone V, Tamura J, Li H, Rillema J,
Licea M. MGD013, a bispecific PD-1 x LAG-3 Dual-Affinity Re-Targeting (DART®)
protein with T-cell immunomodulatory activity for cancer treatment. *Cancer Res*.
2016;76:3217.
29. Savitsky D, Ward R, Riordan C, Mundt C, Jennings S, Connolly J, Findeis M, Sanicola M,
Underwood D, Nastri H, et al. Abstract 3819: INCAGN02385 is an antagonist antibody
targeting the co-inhibitory receptor LAG-3 for the treatment of human malignancies.
Cancer Research. 2018;78:3819-3819. doi: 10.1158/1538-7445.Am2018-3819
30. Zettl M, Wurm M, Schaaf O, Tirapu I, Mostböck S, Reschke M, Schmidbauer S-M, Frego
L, Lorenz IC, Thibodeau M, et al. Abstract 4547: Characterization of the LAG-3 targeting
antibody BI 754111 in monotherapy and in combination with the anti-PD-1 antibody BI
754091. *Cancer Research*. 2018;78:4547-4547. doi: 10.1158/1538-7445.Am2018-4547
31. Hong DS, Schoffski P, Calvo A, Sarantopoulos J, Olza MOD, Carvajal RD, Prawira A,
Kyi C, Esaki T, Akerley WL, et al. Phase I/II study of LAG525 ± spartalizumab (PDR001)
in patients (pts) with advanced malignancies. *Journal of Clinical Oncology*.
2018;36:3012-3012. doi: 10.1200/JCO.2018.36.15_suppl.3012
32. Kraman M, Faroudi M, Allen NL, Kmiecik K, Gliddon D, Seal C, Koers A, Wydro MM,
Batey S, Winnewisser J, et al. FS118, a Bispecific Antibody Targeting LAG-3 and PD-L1,
Enhances T-Cell Activation Resulting in Potent Antitumor Activity. *Clin Cancer Res*.
2020;26:3333-3344. doi: 10.1158/1078-0432.CCR-19-3548
33. Avice MN, Sarfati M, Triebel F, Delespesse G, Demeure CE. Lymphocyte activation
gene-3, a MHC class II ligand expressed on activated T cells, stimulates TNF-alpha and
IL-12 production by monocytes and dendritic cells. *J Immunol*. 1999;162:2748-2753.
34. Buisson S, Triebel F. MHC class II engagement by its ligand LAG-3 (CD223) leads to a
distinct pattern of chemokine and chemokine receptor expression by human dendritic
cells. *Vaccine*. 2003;21:862-868. doi: 10.1016/s0264-410x(02)00533-9
35. Brignone C, Gutierrez M, Mefti F, Brain E, Jarcau R, Cvitkovic F, Bousetta N, Medioni J,
Gligorov J, Grygar C, et al. First-line chemoimmunotherapy in metastatic breast
carcinoma: combination of paclitaxel and IMP321 (LAG-3Ig) enhances immune
responses and antitumor activity. *J Transl Med*. 2010;8:71. doi: 10.1186/1479-5876-8-71
36. Legat A, Maby-El Hajjami H, Baumgaertner P, Cagnon L, Abed Maillard S, Geldhof C,
Iancu EM, Lebon L, Guillaume P, Dojcinovic D, et al. Vaccination with LAG-3Ig (IMP321)
and Peptides Induces Specific CD4 and CD8 T-Cell Responses in Metastatic Melanoma
Patients--Report of a Phase I/IIa Clinical Trial. *Clin Cancer Res*. 2016;22:1330-1340. doi:
10.1158/1078-0432.CCR-15-1212
37. Romano E, Michielin O, Voelter V, Laurent J, Bichat H, Stravodimou A, Romero P,
Speiser DE, Triebel F, Leyvraz S, et al. MART-1 peptide vaccination plus IMP321 (LAG-
3Ig fusion protein) in patients receiving autologous PBMCs after lymphodepletion:
results of a Phase I trial. *J Transl Med*. 2014;12:97. doi: 10.1186/1479-5876-12-97
38. Lucas CL, Workman CJ, Beyaz S, LoCascio S, Zhao G, Vignali DA, Sykes M. LAG-3,
TGF-beta, and cell-intrinsic PD-1 inhibitory pathways contribute to CD8 but not CD4 T-
cell tolerance induced by allogeneic BMT with anti-CD40L. *Blood*. 2011;117:5532-5540.
doi: 10.1182/blood-2010-11-318675
39. Haudebourg T, Dugast AS, Coulon F, Usal C, Triebel F, Vanhove B. Depletion of LAG-3
positive cells in cardiac allograft reveals their role in rejection and tolerance.
Transplantation. 2007;84:1500-1506. doi: 10.1097/01.tp.0000282865.84743.9c
40. Zhang Z, Schlachta C, Duff J, Stiller C, Grant D, Zhong R. Improved techniques for
kidney transplantation in mice. *Microsurgery*. 1995;16:103-109.
41. Han WR, Murray-Segal LJ, Mottram PL. Modified technique for kidney transplantation in
mice. *Microsurgery*. 1999;19:272-274.

42. Murata K, Fox-Talbot K, Qian Z, Takahashi K, Stahl GL, Baldwin WM, 3rd, Wasowska
BA. Synergistic deposition of C4d by complement-activating and non-activating
antibodies in cardiac transplants. *Am J Transplant*. 2007;7:2605-2614.
43. Gorbacheva V, Fan R, Fairchild RL, Baldwin WM, 3rd, Valujskikh A. Memory CD4 T
Cells Induce Antibody-Mediated Rejection of Renal Allografts. *J Am Soc Nephrol*. 2016.
doi: 10.1681/ASN.2015080848
44. Ayasoufi K, Kohei N, Nicosia M, Fan R, Farr GW, McGuirk PR, Pelletier MF, Fairchild RL,
Valujskikh A. Aquaporin 4 blockade improves survival of murine heart allografts
subjected to prolonged cold ischemia. *Am J Transplant*. 2018;18:1238-1246. doi:
10.1111/ajt.14624
45. Nicosia M, Miyairi S, Beavers A, Farr GW, McGuirk PR, Pelletier MF, Valujskikh A.
Aquaporin 4 inhibition alters chemokine receptor expression and T cell trafficking. *Sci
Rep*. 2019;9:7417. doi: 10.1038/s41598-019-43884-2
46. Bickerstaff A, Nozaki T, Wang JJ, Pelletier R, Hadley G, Nadasdy G, Nadasdy T,
Fairchild RL. Acute humoral rejection of renal allografts in CCR5(-/-) recipients. *Am J
Transplant*. 2008;8:557-566.
47. Yagisawa T, Tanaka T, Miyairi S, Tanabe K, Dvorina N, Yokoyama WM, Valujskikh A,
Baldwin WM, 3rd, Fairchild RL. In the absence of natural killer cell activation donor-
specific antibody mediates chronic, but not acute, kidney allograft rejection. *Kidney Int*.
2019;95:350-362. doi: 10.1016/j.kint.2018.08.041
48. Previte DM, Martins CP, O'Connor EC, Marre ML, Coudriet GM, Beck NW, Menk AV,
Wright RH, Tse HM, Delgoffe GM, et al. Lymphocyte Activation Gene-3 Maintains
Mitochondrial and Metabolic Quiescence in Naive CD4(+) T Cells. *Cell Rep*.
2019;27:129-141 e124. doi: 10.1016/j.celrep.2019.03.004
49. Li X, Sun Q, Zhang M, Xie K, Chen J, Liu Z. Capillary dilation and rarefaction are
correlated with intracapillary inflammation in antibody-mediated rejection. *J Immunol Res*.
2014;2014:582902. doi: 10.1155/2014/582902
50. Tse GH, Hughes J, Marson LP. Systematic review of mouse kidney transplantation.
Transpl Int. 2013;26:1149-1160. doi: 10.1111/tri.12129
51. Alessandrini A, Turka LA. FOXP3-Positive Regulatory T Cells and Kidney Allograft
Tolerance. *Am J Kidney Dis*. 2017;69:667-674. doi: 10.1053/j.ajkd.2016.10.027
52. Miyajima M, Chase CM, Alessandrini A, Farkash EA, Della Pelle P, Benichou G,
Graham JA, Madsen JC, Russell PS, Colvin RB. Early acceptance of renal allografts in
mice is dependent on foxp3(+) cells. *Am J Pathol*. 2011;178:1635-1645. doi:
10.1016/j.ajpath.2010.12.024
53. Workman CJ, Vignali DA. Negative regulation of T cell homeostasis by lymphocyte
activation gene-3 (CD223). *J Immunol*. 2005;174:688-695. doi:
10.4049/jimmunol.174.2.688
54. Camisaschi C, Casati C, Rini F, Perego M, De Filippo A, Triebel F, Parmiani G, Belli F,
Rivoltini L, Castelli C. LAG-3 expression defines a subset of CD4(+)CD25(high)Foxp3(+)
regulatory T cells that are expanded at tumor sites. *J Immunol*. 2010;184:6545-6551. doi:
10.4049/jimmunol.0903879
55. Butler NS, Moebius J, Pewe LL, Traore B, Doumbo OK, Tygrett LT, Waldschmidt TJ,
Crompton PD, Harty JT. Therapeutic blockade of PD-L1 and LAG-3 rapidly clears
established blood-stage Plasmodium infection. *Nat Immunol*. 2011;13:188-195. doi:
10.1038/ni.2180
56. Khan M, Arooj S, Wang H. NK Cell-Based Immune Checkpoint Inhibition. *Front Immunol*.
2020;11:167. doi: 10.3389/fimmu.2020.00167
57. Garcia Cruz D, Giri RR, Gamiotea Turro D, Balsbaugh JL, Adler AJ, Rodriguez A.
Lymphocyte Activation Gene-3 Regulates Dendritic Cell Metabolic Programing and T
Cell Priming Function. *J Immunol*. 2021. doi: 10.4049/jimmunol.2001188

58. Fukami N, Ramachandran S, Saini D, Walter M, Chapman W, Patterson GA,
Mohanakumar T. Antibodies to MHC class I induce autoimmunity: role in the
pathogenesis of chronic rejection. *J Immunol*. 2009;182:309-318. doi:
10.4049/jimmunol.182.1.309
59. Akiyoshi T, Hirohashi T, Alessandrini A, Chase CM, Farkash EA, Neal Smith R, Madsen
JC, Russell PS, Colvin RB. Role of complement and NK cells in antibody mediated
rejection. *Hum Immunol*. 2012;73:1226-1232. doi: 10.1016/j.humimm.2012.07.330
60. Chong AS, Alegre ML, Miller ML, Fairchild RL. Lessons and limits of mouse models.
Cold Spring Harb Perspect Med. 2013;3:a015495. doi: 10.1101/cshperspect.a015495
61. Tawbi HA, Schadendorf D, Lipson EJ, Ascierto PA, Matamala L, Castillo Gutierrez E,
Rutkowski P, Gogas HJ, Lao CD, De Menezes JJ, et al. Relatlimab and Nivolumab
versus Nivolumab in Untreated Advanced Melanoma. *N Engl J Med*. 2022;386:24-34.
doi: 10.1056/NEJMoa2109970
62. Yu X, Huang X, Chen X, Liu J, Wu C, Pu Q, Wang Y, Kang X, Zhou L. Characterization
of a novel anti-human lymphocyte activation gene 3 (LAG-3) antibody for cancer
immunotherapy. *MAbs*. 2019;11:1139-1148. doi: 10.1080/19420862.2019.1629239
63. Poirier N, Haudebourg T, Brignone C, Dilek N, Hervouet J, Minault D, Coulon F, de Silly
RV, Triebel F, Blancho G, et al. Antibody-mediated depletion of lymphocyte-activation
gene-3 (LAG-3(+))-activated T lymphocytes prevents delayed-type hypersensitivity in
non-human primates. *Clin Exp Immunol*. 2011;164:265-274. doi: 10.1111/j.1365-
2249.2011.04329.x

Figure Legends

Figure 1. LAG3 deficient mice have expanded lymphocyte subsets. *Naïve none-transplanted B6.WT and B6.LAG3$^{-/-}$ mice were sacrificed at 10 weeks of age.* ***A.*** *Splenic T cell composition: CD4 (CD3^{+}CD4^{+}), CD8 (CD3^{+}CD8^{+}), Tregs (CD3^{+}, CD4^{+}, FoxP3^{+}), CD4 TEff$_{M}$ (CD3^{+}, CD4^{+}, CD62L^{lo}, CD44^{hi}), CD8 TEff$_{M}$ (CD3^{+}, CD8^{+}, CD62L^{lo}, CD44^{hi}), Tfh (TCRb^{+}, CD4^{+}, FoxP3^{-}, PD-1^{+}, CXCR5^{+}) and Tfr cells (TCRb^{+}, CD4^{+}, FoxP3^{+}, PD-1^{+}, CXCR5^{+}).* ***B.*** *The composition of splenic B cell subsets. Subsets were defined as follows; FoB -B220^{+}, IgM^{int}, CD21/35^{int}, MZB - B220^{+}, IgM^{hi}, CD21/35^{hi},TrB - B220^{+}, IgM^{hi}, CD21/35^{lo}, Bregs - CD19^{+}, CD1d^{hi}, CD5^{+}, GCB - B220^{+}, GL7^{+}, CD38^{lo}, and PC –B220^{-},CD138^{hi}. The data represents at least two pooled experiments where each symbol represents an individual mouse. Student's T tests were performed and p<0.05 were considered significant.*

Figure 2. LAG3 deficient mice have increased alloreactivity. A. *Left: Representative*

ELISPOT wells demonstrating the frequencies of alloreactive IFNγ-secreting splenic T cells in

naïve non-transplanted B6.WT and B6.LAG3$^{-/-}$ mice against BALB/c, C3H, SJL and DBA

stimulator cells. Right: Quantification of basal frequencies of alloreactivity in B6.WT and

B6.LAG3$^{-/-}$ mice. **B.** *ELISA quantification of serum IgG reactive to allogenic MHC I and II*

molecules. The data represents at least two pooled experiments where each symbol represents

an individual mouse. Student's T tests were performed and p<0.05 were considered significant.

Figure 3. LAG3-deficient recipients acutely reject kidney allografts. *Groups of B6.WT and*

B6.LAG3$^{-/-}$ were transplanted with complete major histocompatibility complex-mismatched C3H

kidney allografts (n=4-5/group). ***A.*** *Renal allograft survival.* ***B.*** *Serum creatinine levels at d. 14*

posttransplant. ***C.*** *Renal allografts analyzed at the time of rejection (B6.LAG3$^{-/-}$) or on d. 14*

posttransplant (B6.WT) by Trichrome C and immunoperoxidase staining for CD4, CD8 and

complement component C4d. The photographs were taken at 200x and are representative of 4-

5 animals in each group. ***D & E.*** *Flow cytometry analysis of graft infiltrating immune cells: CD3^{+}*

(CD45^{+} CD3^{+}), CD4^{+} (CD45^{+} CD3^{+} CD4^{+}), CD8^{+} (CD45^{+} CD3^{+} CD8^{+}), Tregs (CD45^{+}, CD3^{+}

CD4^{+} FoxP3^{+}), B220^{+} (CD45^{+} B220^{+}), and NK (CD45^{+} CD3^{-} DX1^{+}) at d10 following

transplantation of C3H kidney allografts to B6.WT or B6.LAG3$^{-/-}$ recipients. The data represents

one of two experiments where each symbol represents an individual mouse. Statistical analysis

of allograft survival was measured using Mantel-Cox log-rank test and for other analyses

student's T tests were performed and p<0.05 were considered significant.

*Figure 4. Recipient LAG3 deficiency enhances anti-donor immune responses. Analyses of donor-reactive immunity in B6.WT and B6.LAG3$^{-/-}$ allograft recipients were performed at d. 10 posttransplant. **A.** The composition of spleen T cell subsets was defined as follows: CD3 – CD3$^+$, CD4 – CD3$^+$CD4$^+$, CD8 - CD3$^+$CD8$^+$, Tregs - CD3$^+$CD4$^+$ FoxP3$^+$, CD4 Naïve - CD3$^+$CD4$^+$CD62L^{hi}CD44^{lo}, CD4 C$_M$ - CD3$^+$CD4$^+$CD62L^{hi}CD44^{hi}, CD4 Eff$_M$ - CD3$^+$CD4$^+$CD62L^{lo}CD44^{hi}, CD8 Naïve - CD3$^+$CD8$^+$CD62L^{hi}CD44^{lo}, CD8 C$_M$ - CD3$^+$CD8$^+$CD62L^{hi}CD44^{hi}, CD8 Eff$_M$ - CD3$^+$CD8$^+$CD62L^{lo}CD44^{hi}, TFh - TCRb$^+$CD4$^+$FoxP3$^-$PD-1$^+$CXCR5$^+$ and TFreg - TCRb$^+$CD4$^+$FoxP3$^+$PD-1$^+$CXCR5$^+$. **B.** The composition of splenic B cell was defined as follows: B220 – B220$^+$, FoB - B220$^+$IgM^{int}CD21/35^{int}, MZB - B220$^+$IgM^{hi}CD21/35^{hi}, TrB - B220$^+$IgM^{hi}CD21/35^{lo}, Bregs - CD19$^+$CD1d^{hi}CD5$^+$, GCB - B220$^+$GL7$^+$CD38^{lo}, PCB - B220$^-$CD138^{hi}. **C.** The frequencies of donor reactive IFNγ-secreting splenocytes on d. 14 posttransplant. **D.** Serum levels of IgG against donor MHC-I (H2-D^k) and MHC-II (I-A^k). The data are pooled from two-three experiments, and each symbol represents an individual mouse. Student's T tests were performed and p<0.05 were considered significant.*

 Figure 5. Kidney allograft rejection in LAG3 deficient recipients is not mediated by CD8 T

cells. *A. B6.WT or B6.LAG3$^{-/-}$ recipients depleted of CD8$^+$ T cells prior to transplantation of C3H*

renal allografts. **B.** *Survival of renal allografts (n=5-6/group).* **C.** *Serum levels of IgG against*

donor MHC-I (H2-D^k) and MHC-II (I-A^k) in B6.WT and B6.LAG3$^{-/-}$ kidney allograft recipients. **D.**

The frequencies of donor reactive IFNγ-secreting splenocytes on d. 14 posttransplant. **E.** *Renal*

allografts harvested at the time of rejection and analyzed by Trichrome C and

immunoperoxidase staining for CD4, CD8 and complement component C4d. Images were taken

at 200x and are representative of 4-5 animals in each group. The data are pooled from two-

three experiments, and each symbol represents an individual mouse. Statistical analysis of

allograft survival was measured using Mantel-Cox log-rank test and for other analyses student's

T tests were performed and p<0.05 were considered significant.

Figure 6. Kidney allograft rejection by LAG3 deficient recipients is mediated by B cells. A. *B cells were depleted in B6.WT or B6.LAG3$^{-/-}$ recipients after transplantation of C3H renal allografts.* **B.** *Survival of renal allografts (n=5-6/group).* **C.** *Serum levels of IgG against donor MHC-I (H2-D^k) and MHC-II (I-A^k).* **D.** *The frequencies of donor reactive IFNγ-secreting splenocytes on d. 14 posttransplant.* **E.** *Renal allografts harvested at the time of rejection and analyzed by Trichrome C and immunoperoxidase staining for CD4, CD8 and complement component C4d. Images were taken at 200x and are representative of 4-5 animals in each group. The data are pooled from two-three experiments, and each symbol represents an individual mouse. Statistical analysis of allograft survival was measured using Mantel-Cox log-rank test and for other analyses student's T tests were performed and p<0.05 were considered significant.*

Figure 7. LAG3 is induced on B cell subsets following transplantation. _Analyses of LAG3_

expression on B cell subsets in B6.WT and B6.LAG3$^{-/-}$ renal allograft recipients on d. 10

posttransplant. **_A._** _Representative contour plots of LAG3 expression in Bregs and plasma cell_

pool. **_B._** _Quantification of LAG3 expression in different B cell subsets following transplantation._

The dashed line represents the mean MFI of the WT-FMO control. The data are pooled from

two experiments, and each symbol represents an individual mouse. Student's T tests were

performed and p<0.05 were considered significant.

Figure 8. T cell LAG3 expression is dispensable for kidney allograft acceptance. **A.**

Experimental design*.* ***B.*** *Survival of renal allografts (n=5-7/group).* ***C.*** *Serum levels of IgG*

against donor MHC-I (H2-D^k) and MHC-II (I-A^k) in B6.CD4Cre$^{+/-}$LAG3$^{fl/fl}$ or B6.CD4Cre$^{+/-}$

littermate control kidney allograft recipients. ***D.*** *The frequencies of donor reactive IFNγ-secreting*

splenocytes on d. 14 posttransplant. ***E.*** *Renal allografts harvested at the time of rejection and*

analyzed by Trichrome C and immunoperoxidase staining for CD4, CD8 and complement

component C4d. Images were taken at 200x and are representative of 4-5 animals in each

group. The data are pooled from two-three experiments, and each symbol represents an

individual mouse. Statistical analysis of allograft survival was measured using Mantel-Cox log-

rank test and for other analyses student's T tests were performed and p<0.05 were considered

significant.

Figure 9. B cell LAG3 expression is dispensable for kidney allograft acceptance. **A.**

Experimental design*. **B.** Survival of renal allografts (n=7-8/group).* **C.** *Serum levels of IgG*

against donor MHC-I (H2-D^k) and MHC-II (I-A^k) in B6.*CD19Cre$^{+/-}$ littermate controls or*

B6.CD19Cre$^{+/-}$LAG3$^{fl/fl}$ kidney allograft recipients. **D.** *The frequencies of donor reactive IFNγ-*

secreting splenocytes on d. 14 posttransplant. **E.** *Renal allografts harvested at the time of*

rejection and analyzed by Trichrome C and immunoperoxidase staining for CD4, CD8 and

complement component C4d. Images were taken at 200x and are representative of 4-5 animals

in each group. The data are pooled from two-three experiments, and each symbol represents an

individual mouse. Statistical analysis of allograft survival was measured using Mantel-Cox log-

rank test and for other analyses student's T tests were performed and p<0.05 were considered

significant.

Figure 10. LAG3 blockade enhances de novo alloresponses following kidney transplantation leading to chronic antibody mediated graft injury. *A. B6.WT treated with anti-LAG3 mAb (Clone C9B7W) or control IgG posttransplantation of C3H renal allografts. **B.** Survival of renal allografts (n=6-9/group). **C-D.** NGAL and KIM1 levels in the urine collected from kidney allograft recipients. **E.** Serum Blood Urea Nitrogen (BUN) levels. **F-G.** Serum levels of IgG against donor MHC-I (H2-D^k) and MHC-II (I-A^k). **H.** The frequencies of donor reactive IFNγ-secreting splenocytes on d. 42 posttransplant. **I.** Renal allografts harvested at the time of rejection and analyzed by immunoperoxidase staining for complement component C4d. Images were taken at 400x and are representative of 4-5 animals in each group. The data are pooled from two-three experiments, and each symbol represents an individual mouse. Statistical analysis of allograft survival was measured using Mantel-Cox log-rank test, for the time-course analysis of kidney injury markers and for DSA one-way ANOVA with Tukey's multiple comparison was performed and for ELISPOT analysis student's T tests were performed and p<0.05 were considered significant.*

A

C3H - H-2^k

WT or LAG3$^{-/-}$
B6 H-2^b

DAY −3 −2 −1 0 5 10 15 etc.

αCD8 mAbs on d −3, −2, −1, and every 5 days posttransplant

B CD8 T cell depletion

Percent survival vs Days posttransplant

- WT + αCD8 mAbs — MST> d100
- LAG3$^{-/-}$ + αCD8 mAbs — MST= d16
- LAG3$^{-/-}$ untreated

p=0.0227

C

Day14: anti-H2-D^k DSA

OD 415 nm, WT vs LAG3$^{-/-}$, p = 0.0290

Day14: anti-I-A^k DSA

OD 415 nm, WT vs LAG3$^{-/-}$, p = 0.0070

D Day14: ELISPOT

IFNγ spots/10^6 splenocytes, WT vs LAG3$^{-/-}$, p = 0.1031

E

	WT	LAG3$^{-/-}$
Trichrome C		
CD4		
CD8		
C4d		

A

C3H - H-2^k

WT or LAG3$^{-/-}$
B6 H-2^b

DAY 0 3 8 11 16 etc.

αB220 + αCD19 mAbs every 5 days posttransplant from d3

B B cell depletion

p=0.729

MST> d100

MST>d100

- ● WT + αB220 + αCD19 mAbs
- ■ LAG3$^{-/-}$ + αB220 + αCD19 mAbs
- △ LAG3$^{-/-}$ untreated

Percent survival

Days posttransplant

C

Day14: anti-H2-D^k DSA

OD 415 nm

p = 0.3331

WT   LAG3$^{-/-}$

Day14: anti-I-A^k DSA

OD 415 nm

p = 0.2126

WT   LAG3$^{-/-}$

D

Day14: ELISPOT

IFNγ spots/10^6 splenocytes

p = 0.0117

WT   LAG3$^{-/-}$

E

WT   LAG3$^{-/-}$

Trichrome C

CD4

CD8

C4d

A C3H – H-2^k

CD4Cre$^{+/-}$ or
CD4Cre$^{+/-}$ LAG3$^{-fl/fl}$
B6 H-2^b

B CD4^{Cre}LAG3$^{fl/fl}$

Percent survival

MST> d100

Days posttransplant

- CD4Cre$^{+/-}$ (n=7)
- CD4Cre$^{+/-}$ LAG3$^{fl/fl}$ (n=5)
- LAG3$^{-/-}$ (n=5)

C Day14: anti-H2-D^k DSA

OD 415 nm

p = 0.0167

CD4Cre$^{+/-}$ CD4Cre$^{+/-}$
LAG3$^{fl/fl}$

Day14: anti-I-A^k DSA

OD 415 nm

p = 0.0007

CD4Cre$^{+/-}$ CD4Cre$^{+/-}$
LAG3$^{fl/fl}$

D Day14: ELISPOT

IFNγ spots/10^6
splenocytes

p = 0.2086

CD4Cre$^{+/-}$ CD4Cre$^{+/-}$
LAG3$^{fl/fl}$

E CD4Cre   CD4CreLAG3$^{fl/fl}$

Trichrome C

CD4

CD8

C4d

A

C3H - H-2^k

CD19Cre$^{+/-}$ or
CD19Cre$^{+/-}$ LAG3$^{-fl/fl}$
B6 H-2^b

B

CD19^{Cre}.LAG3$^{fl/fl}$

- CD19Cre$^{+/-}$ (n=7)
- CD19Cre$^{+/-}$ LAG3$^{fl/fl}$ (n=8)
- LAG3$^{-/-}$ (n=5)

Days posttransplant

C

Day14: anti-H2-D^k DSA

OD 415 nm

p = 0.1724

CD19Cre$^{+/-}$ CD19Cre$^{+/-}$
LAG3$^{fl/fl}$

Day14: anti-I-A^k DSA

OD 415 nm

p = 0.0902

CD19Cre$^{+/-}$ CD19Cre$^{+/-}$
LAG3$^{fl/fl}$

D

Day14: ELISPOT

IFNy spots/10^6
splenocytes

p = 0.9143

CD19Cre$^{+/-}$ CD19Cre$^{+/-}$
LAG3$^{fl/fl}$

E

CD19Cre   CD19CreLAG3$^{fl/fl}$

Trichrome C

CD4

CD8

C4d

A

C3H - H-2^k

WT B6 H-2^b

DAY 0 3 5 7 9 11 13 15

αLAG3 mAbs on every other day from d3 - 15 posttransplant

B

LAG3 Blockade

Percent survival

MST> d100

MST>d100

- WT control IgG (n=6)
- WT + α-LAG3 (n=9)
- LAG3$^{-/-}$ untreated (n=5)

Days posttransplant

C

Urine - NGAL

NGAL pg/ml

p = 0.45973   p = 0.45973   p = 0.39643   p = 0.45973

Days posttransplant

- Control IgG
- αLAG3

D

Urine - KIM1

KIM1 pg/ml

0.60028   0.13894   0.51524   0.51524

Days posttransplant

- Control IgG
- αLAG3

E

Serum - BUN

BUN mg/dL

p = 0.21846   p = 0.46323   p = 0.07325   p = 0.38822

Naive B6

Days posttransplant

- Control IgG
- αLAG3

F

anti-H2-D^k

OD 415 nm

p = 0.02030   p = 0.39236   p = 0.00232   p = 0.39236

Days posttransplant

- Control IgG
- αLAG3

G

anti-I-A^k

OD 415 nm

p = 0.46795   p = 0.15650   p = 0.00992   p = 0.05823

Days posttransplant

- Control IgG
- αLAG3

H

Day 42: ELISPOT

IFNγ spots/10^6 splenocytes

p < 0.0001

Control IgG   αLAG3

I

Control   αLAG3

C4d