Rare and population-specific functional variation across pig lines

Roger Ros-Freixedes[1,2§], Bruno D. Valente[3], Ching-Yi Chen[3], William O. Herring[3], Gregor Gorjanc[1], John M Hickey[1], Martin Johnsson[1,4]

[1] The Roslin Institute and Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Easter Bush, Midlothian, Scotland, UK.

[2] Departament de Ciència Animal, Universitat de Lleida - Agrotecnio-CERCA Center, Lleida, Spain.

[3] The Pig Improvement Company, Genus plc, Hendersonville, TN, USA.

[4] Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, Uppsala, Sweden.

§Corresponding author: RRF roger.ros@roslin.ed.ac.uk

Abstract

Background

17 It is expected that missense and loss-of-function (LOF) variants are responsible for

18 phenotypic differences among breeds, genetic lines and varieties of livestock and crop

19 species that have undergone diverse selection histories. However, there is still limited

20 knowledge about the existing missense and LOF variation in livestock commercial

21 populations, in particular regarding population-specific variation.

Methods

We performed whole-genome re-sequencing of 7,848 individuals from nine

commercial pig breeding lines (average coverage: 4.1x) and imputed genotypes for

440,610 pedigree-related individuals. The variants were categorized according to

predicted functional annotation (from loss-of-function to intergenic) and prevalence

level (number of lines in which the variant segregated; from private to widespread).

Variants in each category were examined in terms of distribution along the genome,

minor allele frequency, Wright's fixation index (F_{ST}), individual load, and association

to production traits.

Results

Of the 46 million called variants, 28% were private (i.e., called in only one line) and

21% were widespread (i.e., called in all nine lines). Genomic regions with low

recombination rate were enriched with private variants. Low-prevalence variants were

enriched for lower allele frequencies, lower F_{ST}, and putatively functional and

regulatory variants (including loss-of-function and deleterious missense). Only a

small subset of low-prevalence variants was found at intermediate allele frequencies

and had large estimated effects on production traits. Individuals on average carried

less private deleterious missense alleles than expected compared to other predicted

consequence types. A small subset of low-prevalence variants with intermediate allele

frequencies and higher F_{ST} were detected as significantly associated to the production

traits and explained small fractions of phenotypic variance (up to 3.2%). These

associations were tagged by other more widespread variants, including intergenic

variants.

Conclusions

43 Most low-prevalence variants are kept at very low allele frequency and only a small

44 subset contributed detectable fractions of phenotypic variance. Low-prevalence

45 variants are therefore unlikely to hinder across-breed analyses, in particular when

46 predicting genomic breeding values using reference populations of a different genetic

47 background.

Introduction

Genetic variation is the basis of selective breeding in livestock and crop

species. From a molecular point of view, genetic variants that result in either altered

protein structures or altered gene expressions are believed to be responsible for much

of the existing variation in production traits. Missense variants change one amino acid

of the encoded protein. Loss-of-function variants (LOF) are predicted to disrupt

protein-coding transcripts in a way that they will not be translated into proteins or that

they will be translated into non-functional proteins. Loss-of-function variants may

change one amino acid codon into a premature stop codon (nonsense variants),

change the reading frame during translation (frameshift indels) or change mRNA

splicing (splicing variants). As such, potentially functional variants in protein-coding

regions are assumed to be easier to detect (e.g., by association analyses) than variants

that moderate gene expression. Thus, missense and LOF variants are typically

prioritised as putative causal variants for the traits of interest (e.g., [1–4]).

Missense and LOF mutations can be pathogenic. For instance, missense and

nonsense variants account for 57% of the entries in the Human Gene Mutation

Database [5] (accessed on 30 April 2021), while small indels account for 22% and

splicing variants account for another 9%. Similarly, in livestock species many

missense and LOF variants have been described as causal of genetic diseases ([6,7];

Online Mendelian Inheritance in Animals [8], accessed on 30 April 2021), embryonic

lethality [9,10] or product defects [11,12]. Deleterious missense and LOF variants are

subject to purifying selection and are more likely to be rare, as they are related to

disease risk or reduced fertility.

However, some missense and LOF mutations can be beneficial too [13].

Moreover, some alleles that would be detrimental in the wild may be preferred in

artificial selection settings. The artificial selection performed in livestock and plant

breeding programs is expected to increase the frequency of alleles that favourably

affect the traits included in the selection objectives. Therefore, it is also expected that

missense and LOF variants are responsible for differences among breeds, genetic lines

and varieties of livestock and crop species that have undergone diverse selection

histories. Identification of such functional variants would have direct applications in

gene-assisted and genomic selection [14–16]. Furthermore, strategies based on

genome editing have been theorized to either promote favourable alleles [17] or

remove deleterious alleles [18] in selection candidates. Nevertheless, there is still

limited knowledge about the existing missense and LOF variation in livestock

commercial populations, in particular regarding population-specific variation, often

referred to as 'private'.

Next-generation sequencing has great potential for animal breeding. One of its

main benefits is the power to detect large amounts of variants, many of which will be

specific to the population under study. Sequencing a large number of individuals is

necessary to achieve high variant discovery rates, particularly for low-frequency

variants [19,20]. There are several sequencing studies that profile the genomic

variation in pigs [21–23], cattle [24,25], or chicken [26]. These studies involved the

sequencing of a low number of individuals (up to a few hundreds) at intermediate or

high sequencing coverage. Alternatively, exploring lower coverage allows affordable

sequencing of a much larger number of individuals, which would enable the

identification of a much larger number of variants.

The objective of our study was to characterize the genetic variants detected in

nine intensely selected pig lines with diverse genetic backgrounds. Particular

emphasis was given to quantifying rare and population-specific functional variants, as

well as the number of missense and LOF variants that an average individual carried.

We also assessed the contribution of population-specific functional variants to the

variance of production traits.

Materials and Methods

Populations and sequencing strategy

We performed whole-genome re-sequencing of 7,848 individuals from nine

commercial pig lines (Genus PIC, Hendersonville, TN) with a total coverage of

approximately 32,114x. Sequencing effort in each of the nine lines was proportional

to population size. Approximately 1.5% (0.9-2.1%) of the pigs in each line were

sequenced. Most pigs were sequenced at low coverage, with target coverage of 1 or

2x. A subset of pigs in each line was sequenced at higher coverage of 5, 15, or 30x.

Thus, the average individual coverage was 4.1x, but the median coverage was 1.5x.

The number of pigs sequenced at different coverage levels for each line is

summarized in Table 1.

The sequenced pigs and their coverage were selected following a three-part

sequencing strategy with the objective of representing the haplotype diversity in each

line. First (1), top sires and dams with the largest number of genotyped progeny were

sequenced at 2x and 1x, respectively. Sires were sequenced at greater coverage

because they individually contributed more progeny than dams. Then (2), the

individuals with the greatest genetic footprint on the population (i.e., those that carry

more of the most common haplotypes) and their immediate ancestors were sequenced

at a coverage between 1x and 30x (AlphaSeqOpt part 1; [27]). The target sequencing

coverage was assigned by an algorithm that maximises the expected phasing accuracy

of the common haplotypes from the cumulated family information. Finally (3), pigs

that carried haplotypes with low cumulated coverage (below 10x) were sequenced at

1x (AlphaSeqOpt part 2; [28]). Sets (2) and (3) were based on haplotypes inferred

from marker array genotypes (GGP-Porcine HD BeadChip; GeneSeek, Lincoln, NE),

which were phased and imputed using AlphaPhase [29] and AlphaImpute [30].

Most sequenced pigs, as well as pedigree-related pigs in those lines, were also

genotyped either at low density (15k markers) using the GGP-Porcine LD BeadChip

(GeneSeek) or at high density (80k markers) using the GGP-Porcine HD BeadChip

(GeneSeek).

Sequencing and data processing

Tissue samples were collected from ear punches or tail clippings. Genomic

DNA was extracted using Qiagen DNeasy 96 Blood & Tissue kits (Qiagen Ltd.,

Mississauga, ON, Canada). Paired-end library preparation was conducted using the

TruSeq DNA PCR-free protocol (Illumina, San Diego, CA). Libraries for

resequencing at low coverage (1 to 5x) were produced with an average insert size of

350 bp and sequenced on a HiSeq 4000 instrument (Illumina). Libraries for

resequencing at high coverage (15 or 30x) were produced with an average insert size

of 550 bp and sequenced on a HiSeq X instrument (Illumina). All libraries were

sequenced at Edinburgh Genomics (Edinburgh Genomics, University of Edinburgh,

Edinburgh, UK).

DNA sequence reads were pre-processed using Trimmomatic [31] to remove

adapter sequences. The reads were then aligned to the reference genome *Sscrofa11.1*

(GenBank accession: GCA_000003025.6) using the BWA-MEM algorithm [32].

Duplicates were marked with Picard (http://broadinstitute.github.io/picard). Single

nucleotide polymorphisms (SNPs) and short insertions and deletions (indels) were

identified with GATK HaplotypeCaller (GATK 3.8.0) [33,34] using default settings.

Variant discovery was performed separately for each individual and then a joint

variant set for each population was obtained by extracting the variant positions from

all the individuals in it. Between 20 and 30 million variants were discovered in each

population.

We extracted the read counts supporting each allele directly from the aligned

reads stored in the BAM files using a pile-up function. This approach was set to avoid

biases towards the reference allele introduced by GATK when applied on low-

coverage whole-genome sequence data [35]. That pipeline uses the tool pysam

(version 0.13.0; https://github.com/pysam-developers/pysam), which is a wrapper

around htslib and the samtools package [36]. We extracted the read counts for all

biallelic SNP positions, after filtering variants in potential repetitive regions with

VCFtools [37]. Such variants were here defined as variants that had mean depth

values 3 times greater than the average realized coverage.

Genotype imputation

Genotypes were jointly called, phased and imputed for a total of 537,257

pedigree-related individuals using the 'hybrid peeling' method implemented in

AlphaPeel [38–40], which used all available marker array and whole-genome

sequence data. Imputation was performed separately for each line using its complete

multi-generational pedigree, which encompassed from 15,495 to 122,753 individuals

each (Table 1). We have previously published reports on the accuracy of imputation

in the same populations using this method [39]. The estimated average individual-

wise dosage correlation was 0.94 (median: 0.97). Individuals with low predicted

imputation accuracy were removed before further analyses. An individual was

predicted to have low imputation accuracy if itself or all of its grandparents were not

genotyped with a marker array or if it had a low degree of connectedness to the rest of

the population. These criteria were based on the analysis of simulated and real data on

imputation accuracy [39]. A total of 440,610 individuals remained, from 5,247 to

104,661 individuals for each line (Table 1). The expected average individual-wise

dosage correlation of the remaining individuals was 0.97 (median: 0.98) according to

our previous estimates. We accounted for the whole minor allele frequency spectrum

for our analyses. However, variants with a minor allele frequency lower than 0.023

had an estimated variant-wise dosage correlations lower than 0.90 [39].

Variant predicted consequence types

182   The frequency of the alternative allele was calculated based on the imputed

genotypes. We defined the 'prevalence level' of each variant as the number of lines in

which the variant segregated. To distinguish between allele frequency and prevalence

level we used the terms 'rare' and 'common' to refer to variants in terms of allele

frequency and 'private' and 'widespread' in terms of prevalence level, where private

variants were those called only in one line and widespread variants those called in all

studied lines. We calculated Wright's fixation statistic (F_{ST}) [41] for each variant

among the lines where the variant segregated as $F_{ST} = (H_T–H_S)/H_T$, where H_T is the

expected heterozygosity across the combined lines assuming Hardy-Weinberg

equilibrium and H_S is the average heterozygosity within lines assuming Hardy-

Weinberg equilibrium.

Variants were annotated using Ensembl Variant Effect Predictor (Ensembl

VEP; version 97, July 2019) [42] using both Ensembl and RefSeq transcript

databases. For variants with multiple predicted consequence types (e.g., in case of

multiple transcripts), the most severe predicted consequence type for each variant was

retrieved. Stop-gain, start-loss, stop-loss, splice donor, and splice acceptor variants

were classified as LOF variants. While frameshift indels are typically included in the

LOF category, we considered them as a separate category in order to be able to

quantify their impact separately. The SIFT scores [43] for missense variants were

retrieved from the Ensembl transcript database. Such variants were classified either as

'deleterious' when their SIFT score were less than 0.05, or 'tolerated' otherwise.

Missense variants according to the RefSeq transcript database were left unclassified.

For further analyses we considered the following nine predicted consequence types:

LOF, frameshift indels, deleterious missense, tolerated missense, unclassified

missense, synonymous, variants in the 5' and 3' untranslated regions (UTR), intronic,

and intergenic variants. We considered the predicted consequence types of LOF,

frameshift indel, deleterious missense, and tolerated missense as putatively functional

variants; the predicted consequence types of synonymous and UTR as putatively

regulatory variants; and the predicted consequence types of intronic and intergenic as

putatively neutral variants.

Load of putatively functional alleles

We used the imputed genotypes to estimate the number of alleles of each

predicted consequence type and prevalence level that an average individual carried.

For the most common predicted consequence types, that number was estimated from

50,000 variants sampled randomly. For tolerated missense variants, we used the

50,000 variants with the highest SIFT scores. To account for the different number of

variants within each predicted consequence type and prevalence level category, we

calculated 'heterozygosity' as the percentage of variants of each category that an

individual carried in heterozygosis, and the 'homozygosity for the alternative allele'

as the percentage of variants of each category that an individual carried in

homozygosis for the alternative allele.

Association to production traits

To further explore the association of variants by prevalence level and

functional annotation to selected traits, we performed genome-wide association

studies (GWAS) for the three largest lines. For each line, we performed GWAS for

three complex traits with moderate to high heritability. Most pigs with records were

born during the 2008–2020 period. Breeding values were estimated by line with a

linear mixed model that included polygenic and non-genetic (as relevant for each

trait) effects. Deregressed breeding values (dEBV) were obtained following the

method by VanRaden and Wiggans [44]. Only individuals in which the trait was

directly measured were retained for further analyses. We fitted a univariate linear

mixed model that accounted for the genomic relationship matrix as:

$$\mathbf{y} = \mathbf{x}_i \beta_i + \mathbf{u} + \mathbf{e},$$

where $\mathbf{y}$ is the vector of preadjusted phenotypes, $\mathbf{x}_i$ is the vector of genotypes for the

ith SNP coded as 0 and 2 if homozygous for either allele or 1 if heterozygous, β_i is

the additive effect of the ith SNP on the trait, $\mathbf{u} \sim N(0, \sigma_u^2 \mathbf{K})$ is the vector of polygenic

effects with the covariance matrix equal to the product of the polygenic additive

variance σ_u^2 and the genomic relationship matrix $\mathbf{K}$, and $\mathbf{e}$ is a vector of uncorrelated

residuals. Due to computational limitations, the genomic relationship matrix **K** was

calculated using only imputed SNP genotypes for the high-density marker array and

its single-value decomposition was taken. We used the FastLMM software [45,46] to

fit the model.

We considered the associations with a p-value equal or smaller than 10^{-6} as

significant. We calculated an enrichment score for each predicted consequence type

and prevalence level category as:

$$\log\left(\frac{\text{nSignCategory}/\text{nNotSignCategory}}{\text{nSignTotal}/\text{nNotSignTotal}}\right),$$

where nSignCategory was the number of variants with significant association (with at

least one trait in one of the three lines) in a given predicted consequence type and

prevalence level category, nNotSignCategory was the number of variants with no

significant association in the same category, and nSignTotal and nNotSignTotal were

the total numbers of variants with and without significant association, respectively.

Linkage disequilibrium is pervasive between nearby significant variants due to

the extremely high variant density of whole-genome sequence data. To account for

this, we defined haplotype blocks so that only a single variant per haplotype block

was considered as the putative driver of the association detected in that region. We

defined the haplotype blocks for each line separately using the *--blocks* function in

Plink 1.9 [47,48]. Pairs of variants within 5 Mb of each other were considered. Two

variants were considered to be in strong linkage disequilibrium if the bottom of the

90% confidence interval of D' was greater than 0.7 and the top of the confidence

interval was at least 0.9. If the top of the confidence interval was smaller than 0.7, it

was considered as strong evidence for historical recombination between the two

variants. The remaining pairs of variants were considered uninformative. Regions

where at least 90% of the informative pairs showed strong linkage disequilibrium

were defined as a haplotype block. Within each haplotype block, we selected one

'candidate variant' as the variant with the most severe predicted consequence type. If

there was more than one variant with the same predicted consequence type, the one

with the lowest p-value was selected. This process was performed separately for each

trait and line. Establishing which of the variants in linkage disequilibrium is the most

likely to be causal remains one of the greatest challenges for GWAS. Nevertheless,

keeping the most severe variant in a haplotype block is a common assumption made

by researchers for prioritisation of variants as candidate.

We calculated the additive variance explained by each variant as $2pqa^2$, where

p and q were the allele frequencies and a was the estimated allele effect of the variant.

We expressed the additive variance explained as a proportion of the phenotypic

variance of each trait. We calculated the maximum percentage of phenotypic variance

explained by an individual variant in each predicted consequence type and prevalence

level category. Finally, we calculated the median F_{ST} of the candidate variants within

each predicted consequence type and prevalence level category. We compared the

median F_{ST} of the candidate variants to the median F_{ST} of the same category as the

logarithm of the ratio of the former to the latter.

Results

Prevalence of variants

A large number of variants were widespread across all nine lines. Private

variants represented a much smaller proportion of the variants called within each line.

However, when counted across lines, they cumulatively predominated over the

prevalent ones. Most variants were neither private nor widespread. A total of

46,344,624 biallelic variants were called and passed quality control criteria across all

lines. The distribution of these variants by line is shown in Table 2. Most variants

(38,642,777) were SNPs, of which 10,595,681 were called in a single line (27%;

366,486 to 2,743,965 within each line) and 8,377,578 (22%) were called in all nine

lines. The remaining 7,701,847 variants were indels, of which 2,436,674 were called

in a single line (32%; 121,525 to 506,149 in each line) and 1,560,353 (20%) were

called in all nine lines.

Initially, 24,394,763 additional variant positions were called across all nine

lines (adding up to 70,739,387 unique variant positions) but failed to meet quality

control criteria. Of these, 148,825 variants were discarded because they had mean

depth values 3 times greater than the average realized coverage, 1,927,221 were

multiallelic within line, and 1,673,219 were biallelic within line but multiallelic when

all lines were considered. The remaining variants were imputed for all pedigreed

individuals, but 20,645,588 of them were fixed for the reference allele in the imputed

individuals that passed our accuracy quality control. This affected mostly variants that

had been called in only one line and indicated that the alternative allele segregated at

very low frequency. Although this could also be caused by false positives in variant

calling, this hypothesis seems unlikely to be the main cause as for more than 99% of

these variants we read the alternative allele in at least two individuals. Additionally,

we previously confirmed with high-coverage data 96.9% of the variants that were

called using low-coverage data [35].

Distribution of variants and relationship with recombination rate

The number of variants by chromosome was strongly correlated with

chromosome length (r=0.98, P<0.05; Table S1). Due to the recombination landscape

of each pig chromosome [49], the average variant density by chromosome was

negatively correlated with chromosome length (r=–0.87, P<0.05; Table S1). The

distribution of variants within chromosomes was positively correlated to

recombination rate (r=0.65, P<0.05, between variant density and recombination rate

in 1-Mb non-overlapping windows [49]; Figure 1a). For example, within line A, there

was on average one variant every 81 bp, but in the 5% 1-Mb windows with the lowest

and highest recombination rates there was on average one variant every 152 and 54

bp, respectively (2.8-fold more variants in windows with high recombination rate).

Across all lines, there was one variant every 49 bp on average, but in the 5% 1-Mb

windows with the lowest and highest recombination rates there was on average one

variant every 79 and 34 bp, respectively (2.3-fold more variants in windows with high

recombination rate).

The distribution of private and widespread variants along the genome also

differed. The distribution of widespread variants was more correlated with

recombination rate than that of private variants (Figures 1b and 1c). As a

consequence, private variants represented a larger proportion of the variation in

regions with low recombination rate, which were depleted of widespread variants.

Across all lines, in the 5% 1-Mb windows with the highest recombination rates there

was on average one private variant every 167 bp and one widespread every 148 bp

(1.1-fold more private variants relative to widespread). In the 5% 1-Mb windows with

the lowest recombination rates there was on average one private variant every 260 bp

and one widespread every 531 bp (2.0-fold more private variants relative to

widespread). There were no genomic regions that were enriched for private variants

across lines (Figure S1).

Frequency and fixation index

The prevalence level and alternative allele frequency were related, in a way

that less prevalent variants had also lower allele frequency (Figure 2) and lower F_{ST}

(Figure 3). Private variants had an average alternative allele frequency of 0.03 (SD

0.09) as opposed to widespread variants, which had an average alternative allele

frequency of 0.50 (SD 0.25). As a consequence of the less prevalent variants generally

having low frequencies in the lines where they segregated, these variants showed a

small degree of differentiation between the lines in which they segregated (F_{ST}=0.04,

SD=0.07). In contrast, the widespread variants allowed for the largest degree of

differentiation between lines (F_{ST}=0.21, SD=0.11).

Prevalence and frequency of putatively functional variants

The predicted consequence types of the variants are shown in Table 3. Half

(50.1%) of the variants were called in intergenic regions and another 47.0% of the

variants were called in intronic regions. Only 2.0% of the variants were called in the

5' and 3' untranslated regions. The coding variants comprised 0.9% of the total

variants, of which more than half were missense (45.5%), frameshift indels (3.1%) or

LOF (3.7%). The variant density of putatively functional variants was only weakly

correlated to recombination rate (Figures 1d).

The low-prevalence variants (i.e., the variants that were identified in one or

few lines) were enriched with missense and LOF variants, as well as potentially

regulatory variants such as those located in 5' and 3' untranslated regions and other

intronic variants. On the other hand, the high-prevalence variants (i.e., the variants

that were identified in many or all lines) were enriched with frameshift indels,

synonymous (non-significant correlation) and intergenic variants. Frameshift indels

are typically included in the LOF category. However, our results show that the LOF

category is very heterogeneous and the frameshift indels presented opposite patterns

to other LOF variants. Therefore, we studied them as a separate category.

Whereas the LOF variants had lower allele frequency than the intergenic

variants in low-prevalence levels, they had similar allele frequencies in high-

prevalence levels (Table 4). Thus, there was a set of LOF variants that were prevalent

across lines and also had particularly high frequencies within lines. The missense

variants, especially those classified as deleterious, had lower allele frequencies than

the intergenic variants for all prevalence levels. The low-prevalence missense variants

were enriched with a larger fraction of deleterious variants and lower SIFT scores

(Figure 4). As opposed to LOF and missense variants, the frameshift indels had

intermediate allele frequencies that were much higher than those of the intergenic

variants, which indicated that in many cases the minor allele was the reference one.

Within prevalence level, the LOF and deleterious missense variants had lower F_{ST}

than the intergenic variants (Table 5), probably because they were kept at low allele

frequencies due to negative selection pressure. The frameshift indels also had lower

F_{ST} despite their intermediate allele frequencies.

Load of putatively functional alleles by prevalence level

Most of the missense deleterious and LOF variants that an individual carried

in homozygosis for the alternative allele were high-prevalence variants. Only a small

proportion of these variants were private. An average individual carried 1,048 (SD 57)

LOF variants in homozygosis for the alternative allele, of which 713 (SD 36) were

widespread across all nine lines and only 20 (SD 7) were private. An average

individual carried 1,379 (SD 165) deleterious missense variants in homozygosis for

the alternative allele, of which 1,012 (SD 79) were widespread and only 4 (SD 3)

were private. An average individual carried 1,080 (SD 89) LOF and 2,632 (SD 235)

deleterious missense variants in heterozygosis.

We found signals of negative selection against deleterious missense variants,

in particular the private ones. Individuals proportionally carried less deleterious

missense variants in homozygosis for the alternative allele than variants of other

predicted consequence types, regardless of prevalence level (Figure 5). Individuals

also carried a proportionally lower number of private tolerated missense, synonymous

and LOF variants in homozygosis for the alternative allele than expected, but not in

heterozygosis.

Association of low-prevalence variants to production traits

Significant variants were enriched with putatively functional and regulatory

variants of different prevalence levels, and depleted of intergenic variants. A total of

108,109 variants were significantly associated to at least one trait in one line. Figures

6a and 6b summarise the enrichment scores for all significant variants. The predicted

consequence types that reached the greatest enrichment scores were LOF, frameshift

indels and unclassified missense variants, with various prevalence levels. Variants

with intermediate prevalence levels were amongst the most enriched. These trends

were accentuated after selecting candidate variants from haplotype blocks. In each

line we defined from 1,554 to 2,118 haplotype blocks. A total of 6,692 candidate

variants remained after accounting for linkage disequilibrium within each haplotype

block for all lines and traits. Figures 6c and 6d summarise the enrichment scores for the candidate variants. The enrichment scores based on the candidate variants revealed a stronger depletion of intergenic variants, as well as intronic (with the exception of high-prevalence), and a much stronger enrichment for LOF, frameshift indels and missense variants. For putatively functional variants, there were no clear trends of their enrichment scores across prevalence levels.

In general, the lower allele frequency of low-prevalence variants hindered the detection of significant associations for these markers. Candidate variants with low prevalence that were detected as significantly associated to the production traits actually had intermediate allele frequencies that were greater than expected for their prevalence level. Another consequence of this was that low-prevalence variants in general explained low percentages of variance (Figure 7), although there were some instances of low-prevalence variants that explained up to 3.2% of phenotypic variance. Significant variants had higher F_{ST} than other variants of the same predicted consequence type and prevalence level (Figure 8). This enrichment was especially strong for low-prevalence variants, which in some instances reached F_{ST} estimates around 0.15.

Discussion

Our results contextualize the importance of population-specific and low-prevalence genetic variants. Next, we will discuss: (1) the distribution and functional annotation of low-prevalence variants, (2) the load of putatively functional alleles by prevalence level, and (3) the association of low-prevalence variants to production traits.

Distribution and functional annotation of low-prevalence variants

The main difficulty for the study of low-prevalence genetic variants is that the

prevalence of a variant across several lines is strongly related to its allele frequency,

in a way that the low-prevalence variants are also rare within the lines where they

occur. This is possibly because low-prevalence variants are relatively recent and are

constrained by negative selection.

On one hand, the distribution of private variants was only weakly correlated to

recombination rate and, therefore, regions with low recombination rate were enriched

for private variants. Although the interplay between recurring sweeps, background

selection and other phenomena at play is not fully understood yet, it is generally

accepted that selection on linked variants leads to loss of variation in regions with low

recombination rates [50]. Our observation that regions with low recombination rate

were enriched for private variants suggests that private variants may have been less

affected by selective sweeps than widespread variants. This would be consistent with

previous observations of the younger age of rare and low-prevalence variants [51],

and suggests that many private variants arose more recently in time than widespread

variants, likely after line differentiation, and accumulated in low-recombining regions

due to the reduced efficacy of purifying selection in those regions [52,53].

On the other hand, the low-prevalence variants were enriched for putatively

functional variants. Variants that affect performance traits or that cause a detrimental

condition are under the action of directional selection and are therefore driven towards

low allele frequencies or fixation [54,55]. The low F_{ST} estimates for the low-

prevalence variants indicated that selection pressure keeps these variants at low minor

allele frequency even when they occur in several lines, especially if they are

putatively functional [56]. This could be caused by natural selection or similar

selection objectives across livestock populations. These observations were also

consistent with previous reports showing that most putatively functional variants were

private to single cattle breeds [24], that putatively functional variants were less likely

to have high frequency of the alternative allele across multiple chicken lines [26], and

that population-specific variants in non-African humans were enriched with putatively

functional variants [57].

The relationship between variant prevalence across lines and allele frequency

highlighted the suitability of using a low-coverage sequencing approach to study this

fraction of genetic variation. Nonetheless, bioinformatics pipelines for calling,

genotyping and even imputing such variants should account for the increased

uncertainty associated to their low allele frequency. We decided on using a very

relaxed variant calling strategy with little filtering to account for as many rare variants

as possible, but a sizeable fraction of these rare variants were discarded after

imputation because they were fixed for the imputed individuals that passed quality

control. Low-coverage sequencing is also unsuitable for other types of genetic

variants, such as structural variations (CNVs, tandem duplications and inversions),

which could also be putatively functional and population-specific [58]. Of course, the

number of called variants and the proportion that were private or widespread depends

of the number of sequenced lines [23,26] as well as the sequencing effort in each line.

Our results also suggest that what is typically grouped as LOF is actually a

heterogeneous category. In particular, frameshift indels showed patterns that did not

conform to the other predicted consequence types.

Load of putatively functional alleles by prevalence level

We found that an average individual carried a larger number of LOF and

missense deleterious than previously reported in other livestock species or in humans.

However, there is not yet a clear consensus on the number of LOF and deleterious

missense alleles that are present in the genome of an average individual. In humans, it

has been estimated that an average individual carries 100-150 LOF alleles [54,59–61]

and around 800 weakly deleterious mutations [62], most of which are rare. In

domestic livestock populations, the number of LOF and deleterious alleles carried on

average by individuals has been reported to be greater than in wild populations [63],

including estimates of 100 to 300 deleterious variants in domestic pigs [64], over 400

deleterious variants in domestic chicken [64], and 1,200-1,500 deleterious variants in

domestic yak [65]. Similar magnitudes have been reported in dogs [66], rice [67], and

sunflower [53].

It has been debated why healthy individuals carry a larger number of LOF

variants in homozygosis than expected [68,69]. These could be driven by the fact that

not all predicted LOF variants are detrimental and their functional impact should be

validated before being considered as such. Many predicted LOF variants are in fact

neutral, advantageous, or even may arise simply because of sequencing and

annotation errors [68]. This claim is supported by the large proportion of LOF

observed in homozygosis for the alternative allele compared to the other consequence

types, which casts doubt on the real impact of those variants. On the contrary,

individuals carried a lower proportion of alleles predicted to be deleterious missense

in homozygous state, which supports that variants predicted as such may have a real

impact on genetic variation of production traits and, therefore, be subjected to

selection pressure.

These observations have implications for the identification of variants to be

used for genomic prediction or subjected to genomic edition strategies such as PAGE

[17] or RAGE [18]. Efforts to promote or remove alleles should target variants that

make a substantial contribution to traits of interest, namely functional variants.

However, it is hard to computationally predict and statistically estimate the effects of

such variants, especially if they have low allele frequency. The number of LOF

variants in homozygosis for the alternative allele suggests that predicted loss of

function is not a good indicator that a variant is strongly deleterious in the context of

animal breeding. Similarly, bioinformatics predictors of missense variant effects

appear to be not very accurate [70,71]. High-throughput fine-mapping and variant

screening would be needed to ascertain variant causality and disentangle causality

from linkage disequilibrium.

Association of low-prevalence variants to production traits

Genome-wide association studies on three polygenic traits of economical

importance in the three largest lines revealed that the significant markers were

enriched for putatively functional variants, such as LOF, frameshift indels and

missense variants, and depleted of intergenic variants. This pattern of enrichment was

similar to previous reports from human datasets [72]. However, only a few of the

population-specific and low-prevalence variants were significantly associated to the

traits, even after accounting for linkage disequilibrium. Most of the significant

variants showed intermediate or high prevalence levels. These observations are

consistent with previous meta-analyses in cattle that showed that significant variants

are often common variants [73]. This could be explained by either the fact that real

quantitative trait nucleotides have intermediate or high allele frequencies or the fact

that most studies are underpowered to map rare causal variants. The latter scenario

still seems more likely given that the significant private and low-prevalence variants

had intermediate allele frequencies. This also supports that these significant variants

have biological functions that contributed to trait phenotypic variance rather that they

reached intermediate allele frequencies by drift or by hitchhiking with linked variants

under selection [74]. However, these amounted to a small number of variants that

explained small fractions of variance. Other more widespread variants, including

intergenic variants, successfully acted as tag variants for them and captured much

larger fractions of trait variance. This makes them more suitable for applications in

animal breeding, as is already the case with marker arrays. A similar result was found

in cattle, where splice site and synonymous variants explained the largest proportions

of trait variance, while missense variants explained almost null variance [75].

It can be hypothesized that some of the low-prevalence variants with low allele

frequency have non-negligible marker effects for traits of interest. Despite the large

amount of individuals included in this study, the large volume of variants and the

pervasiveness of linkage disequilibrium among them still make it very challenging to

disentangle their contribution to trait variance. While genome-wide association

studies involving more than one breed typically find multiple breed-specific

associations (e.g., [76]), based on our results it seems unlikely that breed-specific

associations arise from the low-prevalence variants. They would instead stem from

differences in allele frequency, linkage disequilibrium structure or genetic background

that affect the power to detect the effect of prevalent variants across different

populations. Significant variants were enriched with higher F_{ST} estimates than non-

significant variants, which is also consistent with previous reports [73]. Although the

enrichment was greater for low-prevalence variants, it remains unclear to which

degree these variants could relate to selection history or explain differences among

lines for the studied traits.

Conclusion

Low-prevalence variants are enriched for putatively functional variants,

including LOF and deleterious missense variants. However, most low-prevalence

variants are kept at very low allele frequency and are therefore unlikely to hinder

across-breed analyses. This is particularly relevant when predicting genomic breeding

values using reference populations of a different genetic background. Only a small

subset of low-prevalence variants was found at intermediate allele frequencies and

had large estimated effects on production traits.

Ethics approval and consent to participate

563 The samples used in this study were derived from the routine breeding activities of

564 PIC.

Consent for publication

565 Not applicable.

Availability of data and material

566 The software packages AlphaSeqOpt, AlphaPhase, AlphaImpute and AlphaPeel are

567 available from the AlphaGenes website (http://www.alphagenes.roslin.ed.ac.uk). The

568 datasets generated and analysed in this study are derived from the PIC breeding

569 programme and not publicly available.

Competing interests

570 The authors declare that they have no competing interests. BDV, CYC, and WOH are

571 employees of Genus PIC.

Funding

572 The authors acknowledge the financial support from the BBSRC ISPG to The Roslin

573 Institute (BBS/E/D/30002275), from Genus plc, Innovate UK (grant 102271), and

574 from grant numbers BB/N004736/1, BB/N015339/1, BB/L020467/1, and

575 BB/M009254/1. MJ acknowledges financial support from the Swedish Research

576 Council for Sustainable Development Formas Dnr 2016-01386.

Authors' contributions

577 RRF, MJ and JMH designed the study; RRF, MJ and GG performed the analyses;

578 RRF and MJ wrote the first draft; BDV, CYC, WHO, GG and JMH contributed to the

579 interpretation of the results and provided comments on the manuscript. All authors

580 read and approved the final manuscript.

Acknowledgements

581 This work has made use of the resources provided by the Edinburgh Compute and

582 Data Facility (ECDF) (http://www.ecdf.ed.ac.uk/).

References

585 1. Grobet L, Royo Martin LJ, Poncelet D, Pirottin D, Brouwers B, Riquet J, et al. A
586 deletion in the bovine myostatin gene causes the double-muscled phenotype in cattle.
587 Nat Genet. 1997;17:71–4.

588 2. Grisart B, Farnir F, Karim L, Cambisano N, Kim J-J, Kvasz A, et al. Genetic and
589 functional confirmation of the causality of the DGAT1 K232A quantitative trait

nucleotide in affecting milk yield and composition. Proc Natl Acad Sci. 2004;101:2398–403.

3. Óvilo C, FernáNdez A, Noguera JL, BarragáN C, LetóN R, RodríGuez C, et al. Fine mapping of porcine chromosome 6 QTL and *LEPR* effects on body composition in multiple generations of an Iberian by Landrace intercross. Genet Res. 2005;85:57–67.

4. Zhao H, Qin Y, Xiao Z, Li Q, Yang N, Pan Z, et al. Loss of Function of an RNA Polymerase III Subunit Leads to Impaired Maize Kernel Development. Plant Physiol. 2020;184:359–73.

5. Stenson PD, Ball EV, Mort M, Phillips AD, Shiel JA, Thomas NST, et al. Human Gene Mutation Database (HGMD®): 2003 update. Hum Mutat. 2003;21:577–81.

6. Drögemüller C, Tetens J, Sigurdsson S, Gentile A, Testoni S, Lindblad-Toh K, et al. Identification of the Bovine Arachnomelia Mutation by Massively Parallel Sequencing Implicates Sulfite Oxidase (SUOX) in Bone Development. Georges M, editor. PLoS Genet. 2010;6:e1001079.

7. Waide EH, Dekkers JCM, Ross JW, Rowland RRR, Wyatt CR, Ewen CL, et al. Not All SCID Pigs Are Created Equally: Two Independent Mutations in the *Artemis* Gene Cause SCID in Pigs. J Immunol. 2015;195:3171–9.

8. Nicholas FW. Online Mendelian Inheritance in Animals (OMIA): a record of advances in animal genetics, freely available on the Internet for 25 years. Anim Genet. 2021;52:3–9.

9. Derks MFL, Gjuvsland AB, Bosse M, Lopes MS, van Son M, Harlizius B, et al. Loss of function mutations in essential genes cause embryonic lethality in pigs. PLOS Genet. 2019;15:e1008055.

10. Mesbah-Uddin M, Hoze C, Michot P, Barbat A, Lefebvre R, Boussaha M, et al. A missense mutation (p.Tyr452Cys) in the CAD gene compromises reproductive success in French Normande cattle. J Dairy Sci. 2019;102:6340–56.

11. Ma J, Yang J, Zhou L, Ren J, Liu X, Zhang H, et al. A Splice Mutation in the PHKG1 Gene Causes High Glycogen Content and Low Meat Quality in Pig Skeletal Muscle. PLoS Genet. 2014;10:e1004710.

12. Lunden A. A Nonsense Mutation in the FMO3 Gene Underlies Fishy Off-Flavor in Cow's Milk. Genome Res. 2002;12:1885–8.

13. Joseph SB, Hall DW. Spontaneous Mutations in Diploid Saccharomyces cerevisiae. Genetics. 2004;168:1817–25.

14. Pérez-Enciso M, Rincón JC, Legarra A. Sequence- vs. chip-assisted genomic selection: accurate biological information is advised. Genet Sel Evol. 2015;47:43.

15. MacLeod IM, Bowman PJ, Vander Jagt CJ, Haile-Mariam M, Kemper KE, Chamberlain AJ, et al. Exploiting biological priors and sequence variants enhances

QTL discovery and genomic prediction of complex traits. BMC Genomics.
2016;17:144.

16. Lopez BIM, An N, Srikanth K, Lee S, Oh J-D, Shin D-H, et al. Genomic
Prediction Based on SNP Functional Annotation Using Imputed Whole-Genome
Sequence Data in Korean Hanwoo Cattle. Front Genet. 2021;11:603822.

17. Jenko J, Gorjanc G, Cleveland MA, Varshney RK, Whitelaw CBA, Woolliams
JA, et al. Potential of promotion of alleles by genome editing to improve quantitative
traits in livestock breeding programs. Genet Sel Evol. 2015;47:55.

18. Johnsson M, Gaynor RC, Jenko J, Gorjanc G, de Koning D-J, Hickey JM.
Removal of alleles by genome editing (RAGE) against deleterious load. Genet Sel
Evol. 2019;51:14.

19. Le SQ, Durbin R. SNP detection and genotyping from low-coverage sequencing
data on multiple diploid samples. Genome Res. 2011;21:952–60.

20. Martin AR, Atkinson EG, Chapman SB, Stevenson A, Stroud RE, Abebe T, et al.
Low-coverage sequencing cost-effectively detects known and novel variation in
underrepresented populations. Am J Hum Genet. 2021;108:656–68.

21. Molnár J, Nagy T, Stéger V, Tóth G, Marincs F, Barta E. Genome sequencing and
analysis of Mangalica, a fatty local pig of Hungary. BMC Genomics. 2014;15:761.

22. Choi J-W, Chung W-H, Lee K-T, Cho E-S, Lee S-W, Choi B-H, et al. Whole-
genome resequencing analyses of five pig breeds, including Korean wild and native,
and three European origin breeds. DNA Res. 2015;22:259–67.

23. Cai Z, Sarup P, Ostersen T, Nielsen B, Fredholm M, Karlskov-Mortensen P, et al.
Genomic diversity revealed by whole-genome sequencing in three Danish commercial
pig breeds. J Anim Sci. 2020;98.

24. Daetwyler HD, Capitan A, Pausch H, Stothard P, van Binsbergen R, Brondum RF,
et al. Whole-genome sequencing of 234 bulls facilitates mapping of monogenic and
complex traits in cattle. Nat Genet. 2014;46:858–65.

25. Das A, Panitz F, Gregersen VR, Bendixen C, Holm L-E. Deep sequencing of
Danish Holstein dairy cattle for variant detection and insight into potential loss-of-
function variants in protein coding genes. BMC Genomics. 2015;16:1043.

26. Gheyas AA, Boschiero C, Eory L, Ralph H, Kuo R, Woolliams JA, et al.
Functional classification of 15 million SNPs detected from diverse chicken
populations. DNA Res. 2015;22:205–17.

27. Gonen S, Ros-Freixedes R, Battagin M, Gorjanc G, Hickey JM. A method for the
allocation of sequencing resources in genotyped livestock populations. Genet Sel
Evol. 2017;49:47.

28. Ros-Freixedes R, Gonen S, Gorjanc G, Hickey JM. A method for allocating low-
coverage sequencing resources by targeting haplotypes rather than individuals. Genet
Sel Evol. 2017;49:78.

29. Hickey JM, Kinghorn BP, Tier B, Wilson JF, Dunstan N, van der Werf JH. A
combined long-range phasing and long haplotype imputation method to impute phase
for SNP genotypes. Genet Sel Evol. 2011;43:12.

30. Hickey JM, Kinghorn BP, Tier B, van der Werf JH, Cleveland MA. A phasing
and imputation method for pedigreed populations that results in a single-stage
genomic evaluation. Genet Sel Evol. 2012;44:9.

31. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30:2114–20.

32. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv. 2013;1303.3997v1 [q – bio.GN].

33. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A
framework for variation discovery and genotyping using next-generation DNA
sequencing data. Nat Genet. 2011;43:491–8.

34. Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van der
Auwera GA, et al. Scaling accurate genetic variant discovery to tens of thousands of
samples. bioRxiv. 2018;10.1101/201178.

35. Ros-Freixedes R, Battagin M, Johnsson M, Gorjanc G, Mileham AJ, Rounsley
SD, et al. Impact of index hopping and bias towards the reference allele on accuracy
of genotype calls from low-coverage sequencing. Genet Sel Evol. 2018;50:64.

36. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

37. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The
variant call format and VCFtools. Bioinformatics. 2011;27:2156–8.

38. Whalen A, Ros-Freixedes R, Wilson DL, Gorjanc G, Hickey JM. Hybrid peeling
for fast and accurate calling, phasing, and imputation with sequence data of any
coverage in pedigrees. Genet Sel Evol. 2018;50:67.

39. Ros-Freixedes R, Whalen A, Chen C-Y, Gorjanc G, Herring WO, Mileham AJ, et
al. Accuracy of whole-genome sequence imputation using hybrid peeling in large
pedigreed livestock populations. Genet Sel Evol. 2020;52:17.

40. Ros-Freixedes R, Whalen A, Gorjanc G, Mileham AJ, Hickey JM. Evaluation of
sequencing strategies for whole-genome imputation with hybrid peeling. Genet Sel
Evol. 2020;52:18.

41. Wright S. The genetical structure of populations. Ann Eugen. 1949;15:323–54.

42. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The
Ensembl Variant Effect Predictor. Genome Biol. 2016;17:122.

43. Ng PC, Henikoff S. SIFT: predicting amino acid changes that affect protein
function. Nucleic Acids Res. 2003;31:3812–4.

44. VanRaden PM, Wiggans GR. Derivation, Calculation, and Use of National
Animal Model Information. J Dairy Sci. 1991;74:2737–46.

45. Lippert C, Listgarten J, Liu Y, Kadie CM, Davidson RI, Heckerman D. FaST
linear mixed models for genome-wide association studies. Nat Methods. 2011;8:833–
5.

46. Widmer C, Lippert C, Weissbrod O, Fusi N, Kadie C, Davidson R, et al. Further
Improvements to Linear Mixed Models for Genome-Wide Association Studies. Sci
Rep. 2015;4:6874.

47. Taliun D, Gamper J, Pattaro C. Efficient haplotype block recognition of very long
and dense genetic sequences. BMC Bioinformatics. 2014;15.

48. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-
generation PLINK: rising to the challenge of larger and richer datasets. GigaScience.
2015;4.

49. Johnsson M, Whalen A, Ros-Freixedes R, Gorjanc G, Chen C-Y, Herring WO, et
al. Genetic variation in recombination rate in the pig. Genet Sel Evol. 2021;53.

50. Cutter AD, Payseur BA. Genomic signatures of selection at linked sites: unifying
the disparity among species. Nat Rev Genet. 2013;14:262–74.

51. Mathieson I, McVean G. Demography and the Age of Rare Variants. Novembre J,
editor. PLoS Genet. 2014;10:e1004528.

52. Charlesworth D, Morgan MT, Charlesworth B. Mutation Accumulation in Finite
Populations. J Hered. 1993;84:321–5.

53. Renaut S, Rieseberg LH. The Accumulation of Deleterious Mutations as a
Consequence of Domestication and Improvement in Sunflowers and Other
Compositae Crops. Mol Biol Evol. 2015;32:2273–83.

54. Gudbjartsson DF, Helgason H, Gudjonsson SA, Zink F, Oddson A, Gylfason A, et
al. Large-scale whole-genome sequencing of the Icelandic population. Nat Genet.
2015;47:435–44.

55. Sulem P, Helgason H, Oddson A, Stefansson H, Gudjonsson SA, Zink F, et al.
Identification of a large set of rare complete human knockouts. Nat Genet.
2015;47:448–52.

56. Mezmouk S, Ross-Ibarra J. The Pattern and Distribution of Deleterious Mutations
in Maize. G3 GenesGenomesGenetics. 2014;4:163–71.

57. Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. On the accumulation of
deleterious mutations during range expansions. Mol Ecol. 2013;22:5972–82.

58. Liu GE, Hou Y, Zhu B, Cardone MF, Jiang L, Cellamare A, et al. Analysis of
copy number variations among diverse cattle breeds. Genome Res. 2010;20:693–703.

59. The 1000 Genomes Project Consortium. An integrated map of genetic variation
from 1,092 human genomes. Nature. 2012;491:56–65.

60. MacArthur DG, Balasubramanian S, Frankish A, Huang N, Morris J, Walter K, et
al. A Systematic Survey of Loss-of-Function Variants in Human Protein-Coding
Genes. Science. 2012;335:823–8.

61. Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, et al.
Analysis of protein-coding genetic variation in 60,706 humans. Nature.
2016;536:285–91.

62. Chun S, Fay JC. Identification of deleterious mutations within three human
genomes. Genome Res. 2009;19:1553–61.

63. Makino T, Rubin C-J, Carneiro M, Axelsson E, Andersson L, Webster MT.
Elevated Proportions of Deleterious Genetic Variation in Domestic Animals and
Plants. Genome Biol Evol. 2018;10:276–90.

64. Bosse M, Megens H, Derks MFL, Cara ÁMR, Groenen MAM. Deleterious alleles
in the context of domestication, inbreeding, and selection. Evol Appl. 2019;12:6–17.

65. Xie X, Yang Y, Ren Q, Ding X, Bao P, Yan B, et al. Accumulation of deleterious
mutations in the domestic yak genome. Anim Genet. 2018;49:384–92.

66. Cruz F, Vila C, Webster MT. The Legacy of Domestication: Accumulation of
Deleterious Mutations in the Dog Genome. Mol Biol Evol. 2008;25:2331–6.

67. Lu J, Tang T, Tang H, Huang J, Shi S, Wu C-I. The accumulation of deleterious
mutations in rice genomes: a hypothesis on the cost of domestication. Trends Genet.
2006;22:126–31.

68. MacArthur DG, Tyler-Smith C. Loss-of-function variants in the genomes of
healthy humans. Hum Mol Genet. 2010;19:R125–30.

69. Rausell A, Luo Y, Lopez M, Seeleuthner Y, Rapaport F, Favier A, et al. Common
homozygosity for predicted loss-of-function variants reveals both redundant and
advantageous effects of dispensable human genes. Proc Natl Acad Sci.
2020;117:13626–36.

70. Pagel KA, Pejaver V, Lin GN, Nam H-J, Mort M, Cooper DN, et al. When loss-
of-function is loss of function: assessing mutational signatures and impact of loss-of-
function genetic variants. Bioinformatics. 2017;33:i389–98.

71. Pejaver V, Urresti J, Lugo-Martinez J, Pagel KA, Lin GN, Nam H-J, et al.
Inferring the molecular and phenotypic impact of amino acid variants with MutPred2.
Nat Commun. 2020;11.

72. Schork AJ, Thompson WK, Pham P, Torkamani A, Roddey JC, Sullivan PF, et al.
All SNPs Are Not Created Equal: Genome-Wide Association Studies Reveal a
Consistent Pattern of Enrichment among Functionally Annotated SNPs. Gibson G,
editor. PLoS Genet. 2013;9:e1003449.

73. van den Berg I, Xiang R, Jenko J, Pausch H, Boussaha M, Schrooten C, et al.
Meta-analysis for milk fat and protein percentage using imputed sequence variant
genotypes in 94,321 cattle from eight cattle breeds. Genet Sel Evol. 2020;52:37.

74. Chun S, Fay JC. Evidence for Hitchhiking of Deleterious Mutations within the
Human Genome. Pritchard JK, editor. PLoS Genet. 2011;7:e1002240.

75. Koufariotis LT, Chen Y-PP, Stothard P, Hayes BJ. Variance explained by whole
genome sequence variants in coding and regulatory genome annotations for six dairy
traits. BMC Genomics. 2018;19.

76. Purfield DC, Evans RD, Berry DP. Breed- and trait-specific associations define
the genetic architecture of calving performance traits in cattle. J Anim Sci. 2020;98.

Figures

Figure 1. Variant density in line A (black and grey bars) and recombination rate (red line). The correlation (r) between variant density and recombination rate in 1-Mb non-overlapping windows is reported.

795
796 **Figure 2.** Frequency of the alternative allele by prevalence level. Red dots indicate
797 means. In blue, values greater than 1.5 times the interquartile range.
798

799
800 **Figure 3.** Wright's fixation statistic (F_{ST}) by prevalence level. Red dots indicate
801 means. In blue, values greater than 1.5 times the interquartile range.
802

803
804 **Figure 4.** Classification of the missense variants and median SIFT score by
805 prevalence level.
806

807
808 **Figure 5.** Percentage of variants in homozygosis for the alternative allele or in
809 heterozygosis in an average individual by predicted consequence type, including
810 frameshift indels, and prevalence level. LOF: loss-of-function; UTR: untranslated
811 regions.
812

813
Figure 6. Enrichment scores for the significant variants in the genome-wide
association study by variant prevalence level and predicted consequence type. Either
all significant variants (panels a and b) or only the candidate variants after accounting
for linkage disequilibrium (panels c and d) were used. Prevalence level was
considered across all 9 lines (panels a and c) or only across the 3 lines included in the
genome-wide association study (panels b and d).
820

821
822 **Figure 7.** Maximum percentage of phenotypic variance explained by the individual
823 candidate variants within each prevalence level and predicted consequence type. Only
824 the candidate variants after accounting for linkage disequilibrium were used.
825 Prevalence level was considered across all 9 lines (panel a) or only across the 3 lines
826 included in the genome-wide association study (panel b).
827

Figure 8. Enrichment scores for the F_{ST} median of the candidate variants within each prevalence level and predicted consequence type. Only the candidate variants after accounting for linkage disequilibrium were used. Prevalence level was considered across all 9 lines.

Tables

834   **Table 1.** Number of sequenced and analysed pigs.

Line	Individuals sequenced	Individuals sequenced by coverage				Individuals used in analyses		
		1x	2x	5x	15–30x	Pedigree	Imputed	GWAS
A	1,856	1,044	649	73	90	122,753	104,661	88,342
B	1,491	628	728	54	81	84,420	66,608	56,173
C	1,366	685	545	44	92	88,964	76,230	64,285
D	760	394	274	27	65	50,797	41,573	-
E	731	362	311	16	42	79,981	60,474	-
F	701	351	255	28	67	52,470	39,263	-
G	445	217	176	15	37	21,129	17,224	-
H	381	193	137	16	35	35,309	29,330	-
I	321	111	158	18	34	15,495	5,247	-

836   **Table 2.** Number of variants by line.

Line	Biallelic variant sites (M)	SNPs			Indels		
		All biallelic (M)	Private (M)	Widespread (M)	All biallelic (M)	Private (M)	Widespread (M)
A	28.83	24.38	1.56	8.38	4.44	0.39	1.56
B	28.57	24.32	2.74	8.38	4.24	0.51	1.56
C	28.88	24.60	2.51	8.38	4.28	0.44	1.56
D	21.44	17.94	1.23	8.38	3.50	0.32	1.56
E	19.06	15.71	0.51	8.38	3.35	0.22	1.56
F	20.21	16.86	0.42	8.38	3.35	0.16	1.56
G	23.38	19.64	0.50	8.38	3.74	0.16	1.56
H	22.32	18.78	0.37	8.38	3.55	0.12	1.56
I	24.59	20.82	0.76	8.38	3.77	0.13	1.56
Total	46.30	38.64	10.60	8.38	7.70	2.44	1.56

Table 3. Predicted consequence types of the variants by prevalence level. The most sever consequence of each variant was used. The main Sequence Ontology (SO) terms are shown in order of severity (more severe to less severe) as estimated by Ensembl Variant Effect Predictor. The correlation (r) between the percentage of variants of each consequence type and prevalence is reported. In bold, categories that will be analysed in the next sections.

| Consequence type | Percentage of variants (%) by prevalence level | | | | | | | | | | r |
	1	2	3	4	5	6	7	8	9	Total	
Loss-of-function[1]	**0.061**	**0.035**	**0.026**	**0.021**	**0.019**	**0.017**	**0.019**	**0.018**	**0.019**	**0.032**	**-.76**
Splice acceptor/donor	0.038	0.023	0.014	0.010	0.009	0.007	0.008	0.008	0.008	0.018	-.79
Stop-gain	0.015	0.009	0.008	0.008	0.007	0.007	0.007	0.006	0.006	0.009	-.79
Stop-loss	0.005	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.003	0.003	-.36
Start-loss	0.004	0.002	0.002	0.002	0.001	0.001	0.002	0.002	0.002	0.002	-.47
Frameshift indel	**0.014**	**0.017**	**0.019**	**0.021**	**0.020**	**0.021**	**0.024**	**0.032**	**0.055**	**0.027**	**+.81**
In-frame indel	0.005	0.008	0.009	0.008	0.008	0.008	0.007	0.007	0.005	0.007	-.23
Missense	0.557	0.38	0.356	0.341	0.345	0.338	0.319	0.306	0.325	0.394	-.74
Deleterious	**0.201**	**0.092**	**0.074**	**0.069**	**0.065**	**0.062**	**0.054**	**0.048**	**0.040**	**0.096**	**-.78**
Tolerated	**0.223**	**0.171**	**0.165**	**0.166**	**0.173**	**0.167**	**0.161**	**0.159**	**0.177**	**0.183**	**-.53**
Splice region	0.105	0.098	0.088	0.081	0.083	0.081	0.080	0.081	0.085	0.090	-.76
Synonymous	**0.240**	**0.313**	**0.334**	**0.348**	**0.355**	**0.353**	**0.337**	**0.331**	**0.354**	**0.316**	**+.66**
Untranslated regions	**2.068**	**2.034**	**2.039**	**1.973**	**1.933**	**1.944**	**1.892**	**1.880**	**1.860**	**1.965**	**-.98**
5' UTR	0.633	0.594	0.581	0.584	0.562	0.588	0.546	0.545	0.547	0.582	-.88
3' UTR	1.435	1.440	1.458	1.389	1.372	1.357	1.346	1.335	1.313	1.383	-.94
Non-coding transcript exon	0.107	0.116	0.109	0.117	0.132	0.120	0.108	0.112	0.121	0.115	+.26
Intronic	**47.743**	**47.570**	**47.633**	**47.162**	**46.513**	**46.709**	**46.700**	**46.355**	**46.132**	**46.981**	**-.95**
Upstream gene	3.290	3.281	3.290	3.255	3.292	3.266	3.127	3.148	3.133	3.227	-.84
Downstream gene	2.660	2.679	2.740	2.747	2.746	2.704	2.700	2.707	2.676	2.692	+.04
Intergenic	**43.147**	**43.467**	**43.355**	**43.926**	**44.553**	**44.438**	**44.687**	**45.021**	**45.235**	**44.154**	**+.97**

[1]If frameshift indels were included in this category: r = -.06 (P>.05)
 Significant correlation (P<.05)

845   **Table 4.** Frequency of the alternative allele by predicted consequence type and prevalence level. Values are medians.

Consequence type	Frequency of the alternative allele by prevalence level									
	1	2	3	4	5	6	7	8	9	Total
Loss-of-function	.0010	.017	.048	.061	.089	.114	.151	.225	.489	.020
Frameshift indel	.4847	.758	.757	.420	.304	.260	.339	.456	.693	.634
Deleterious missense	.0006	.018	.043	.061	.078	.092	.125	.170	.350	.010
Tolerated missense	.0011	.027	.047	.066	.083	.106	.143	.202	.443	.074
Synonymous	.0037	.032	.049	.066	.086	.107	.151	.205	.447	.110
Untranslated regions	.0018	.034	.059	.077	.099	.122	.167	.226	.476	.102
Intronic	.0015	.035	.059	.080	.102	.126	.171	.235	.485	.110
Intergenic	.0015	.033	.058	.080	.105	.129	.173	.237	.483	.116

846   **Table 5.** Wright's fixation statistic (F_{ST}) by predicted consequence type and prevalence
847   level. Values are medians.

Consequence type	F_{ST} by prevalence level								
	2	3	4	5	6	7	8	9	Total
Loss-of-function	.003	.022	.047	.066	.094	.114	.144	.171	.071
Frameshift indel	.010	.043	.065	.081	.088	.120	.147	.148	.114
Deleterious missense	.005	.029	.055	.073	.087	.110	.131	.160	.068
Tolerated missense	.009	.036	.061	.084	.107	.127	.158	.184	.108
Synonymous	.013	.040	.062	.090	.110	.130	.158	.194	.117
Untranslated regions	.009	.036	.059	.086	.108	.132	.159	.190	.110
Intronic	.009	.037	.063	.089	.111	.136	.164	.195	.118
Intergenic	.009	.036	.066	.091	.112	.139	.167	.193	.121

849 # Additional files

850 **Table S1.** Number of analysed variants by chromosome.

Chromosome	Length (Mb)	SNPs (M)	Indels (M)	Variant density (thousands/Mb)
1	274.3	3.77	0.76	16.5
2	151.9	2.60	0.52	20.5
3	132.8	2.35	0.44	21.0
4	130.9	2.21	0.43	20.2
5	104.5	1.95	0.39	22.4
6	170.8	2.80	0.55	19.6
7	121.8	2.20	0.43	21.6
8	139.0	2.37	0.50	20.6
9	139.5	2.47	0.48	21.1
10	69.4	1.60	0.31	27.5
11	79.2	1.57	0.31	23.7
12	61.6	1.35	0.25	26.0
13	208.3	2.97	0.64	17.3
14	141.8	2.38	0.48	20.2
15	140.4	2.20	0.46	18.9
16	79.9	1.50	0.30	22.5
17	63.5	1.32	0.25	24.7
18	56.0	1.04	0.19	22.0
Total	2,501.9	38.64	7.70	18.5

852
853 **Figure S1.** Variant density for the private variants in each line.
854