Rare and population-specific functional variation across pig lines

Roger Ros-Freixedes[1,2§], Bruno D. Valente[3], Ching-Yi Chen[3], William O. Herring[3], Gregor Gorjanc[1], John M Hickey[1], Martin Johnsson[1,4]

[1] The Roslin Institute and Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Easter Bush, Midlothian, Scotland, UK.

[2] Departament de Ciència Animal, Universitat de Lleida - Agrotecnio-CERCA Center, Lleida, Spain.

[3] The Pig Improvement Company, Genus plc, Hendersonville, TN, USA.

[4] Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, Uppsala, Sweden.

§Corresponding author: RRF roger.ros@roslin.ed.ac.uk

Abstract

Background

It is expected that functional, mainly missense and loss-of-function (LOF), and

regulatory variants are responsible for phenotypic differences among breeds, genetic

lines, and varieties of livestock and crop species that have undergone diverse selection

histories. However, there is still limited knowledge about the existing missense and

LOF variation in livestock commercial populations, in particular regarding

population-specific variation and how it can affect applications such as across-breed

genomic prediction.

Methods

We re-sequenced the whole genome of 7,848 individuals from nine commercial pig

breeding lines (average sequencing coverage: 4.1x) and imputed whole-genome

genotypes for 440,610 pedigree-related individuals. The called variants were

categorized according to predicted functional annotation (from LOF to intergenic) and

prevalence level (number of lines in which the variant segregated; from private to

widespread). Variants in each category were examined in terms of distribution along

the genome, minor allele frequency, Wright's fixation index (F_{ST}), individual load,

and association to production traits.

Results

Of the 46 million called variants, 28% were private (called in only one line) and 21%

were widespread (called in all nine lines). Genomic regions with low recombination

rate were enriched with private variants. Low-prevalence variants (called in one or a

few lines only) were enriched for lower allele frequencies, lower F_{ST}, and putatively

functional and regulatory roles (including loss-of-function and deleterious missense

variants). Only a small subset of low-prevalence variants was found at intermediate

allele frequencies and had large estimated effects on production traits. Individuals on

average carried less private deleterious missense alleles than expected compared to

other predicted consequence types. A small subset of low-prevalence variants with

intermediate allele frequencies and higher F_{ST} were detected as significantly

associated to the production traits and explained small fractions of phenotypic

variance (up to 3.2%). These associations were tagged by other more widespread

variants, including intergenic variants.

Conclusions

45   Most low-prevalence variants are kept at very low allele frequency and only a small

46   subset contributed detectable fractions of phenotypic variance. Not accounting for

47   low-prevalence variants is therefore unlikely to hinder across-breed analyses, in

48   particular for genomic prediction of breeding values using reference populations of a

49   different genetic background.

Introduction

Genetic variation is the basis of selective breeding in livestock and crop
species. From a molecular point of view, genetic variants that result in either altered
protein structures or altered gene expressions are believed to be responsible for much
of the existing genetic variation in complex traits [1–4]. Missense variants change one
amino acid of the encoded protein. Loss-of-function variants (LOF) are predicted to
disrupt protein-coding transcripts in a way that they will not be translated into
proteins or that they will be translated into non-functional proteins. Loss-of-function
variants may change one amino acid codon into a premature stop codon (nonsense
variants), change the reading frame during translation (frameshift indels) or change
mRNA splicing (splicing variants). As such, potentially functional variants in protein-
coding regions are assumed to be easier to detect (e.g., by association analyses) than
variants that moderate gene expression [5–7]. Thus, missense and LOF variants are
typically prioritised as putative causal variants for the traits of interest (e.g., [8–11]).

Missense and LOF mutations can be pathogenic. For instance, missense and
nonsense variants account for 57% of the entries in the Human Gene Mutation
Database [12] (accessed on 30 April 2021), while small indels account for 22% and
splicing variants account for another 9%. Similarly, in livestock species many
missense and LOF variants have been described as causal of genetic diseases and
post-natal defects ([13–16]; Online Mendelian Inheritance in Animals [17], accessed
on 30 April 2021), embryonic lethality [18,19] or product defects [20,21]. Deleterious
missense and LOF variants are subject to purifying selection and are more likely to be
rare, because they are related to unfavourable phenotypes such as disease risk or
reduced fertility.

However, some missense and LOF mutations can be beneficial too [22].

Moreover, some alleles that would be detrimental in the wild may be preferred in

artificial selection settings. The artificial selection performed in livestock and crop

breeding programs is expected to increase the frequency of alleles that favourably

affect the traits included in the selection objectives. Therefore, it is also expected that

missense and LOF variants are responsible for differences among breeds, genetic

lines, and varieties of livestock and crop species that have undergone diverse selection

histories. Identification of such functional variants would have direct applications in

gene-assisted and genomic selection [23–25]. Furthermore, strategies based on

genome editing have been theorized to either promote favourable alleles [26] or

remove deleterious alleles [27] in selection candidates. Nevertheless, there is still

limited knowledge about the existing missense and LOF variation in commercial

livestock populations, in particular regarding population-specific variation, often

referred to as 'private', and how it can affect applications such as across-breed

genomic prediction.

Next-generation sequencing has great potential for livestock breeding. One of

its main benefits is the power to detect large amounts of variants, many of which will

be specific to the population under study. Sequencing a large number of individuals is

necessary to achieve high variant discovery rates, particularly for low-frequency

variants [28,29]. There are several sequencing studies that profile the genomic

variation in pigs [30–32], cattle [33,34], or chicken [35]. These studies involved the

sequencing of a low number of individuals (up to a few hundreds) at intermediate or

high sequencing coverage. Alternatively, low sequencing coverage allows affordable

sequencing of a much larger number of individuals, which would enable the

identification of a much larger number of variants.

The objective of this study was to characterize the genetic variants detected in

nine intensely selected pig lines with diverse genetic backgrounds. Particular

emphasis was given to quantifying rare and population-specific functional variants, as

well as the number of missense and LOF variants that an average individual carried.

We also assessed the contribution of population-specific functional variants to the

variance of production traits.

Materials and Methods

Populations and sequencing strategy

We re-sequenced the whole genome of 7,848 individuals from nine

commercial pig lines (Genus PIC, Hendersonville, TN) with a total sequencing

coverage of approximately 32,114x. Breeds of origin of the nine lines included Large

White, Landrace, Pietrain, Hampshire, Duroc and synthetic lines. Sequencing effort in

each of the nine lines was proportional to population size. The number of pigs that

were available in the pedigree of each line and the number of sequenced pigs, by

coverage, is summarized in Table 1. Approximately 1.5% (0.9-2.1%) of the pigs in

each line were sequenced. Most pigs were sequenced at low coverage, with target

coverage of 1 or 2x. A subset of pigs in each line was sequenced at higher coverage of

5, 15, or 30x. Thus, the average individual coverage was 4.1x, but the median

coverage was 1.5x. The population structure across the nine lines was assessed with a

principal component analysis using the sequenced pigs and is shown in Figure S1.

The sequenced pigs and their coverage were selected following a three-part

sequencing strategy with the objective of representing the haplotype diversity in each

line. First (1), top sires and dams with the largest number of genotyped progeny were

sequenced at 2x and 1x, respectively. Sires were sequenced at greater coverage

because they individually contributed more progeny than dams. Then (2), the

individuals with the greatest genetic footprint on the population (i.e., those that carry

more of the most common haplotypes) and their immediate ancestors were sequenced

at a coverage between 1x and 30x (AlphaSeqOpt part 1; [36]). The target sequencing

coverage was assigned by an algorithm that maximises the expected phasing accuracy

of the common haplotypes from the accumulated family information. Finally (3), pigs

that carried haplotypes with low cumulated coverage (below 10x) were sequenced at

1x (AlphaSeqOpt part 2; [37]). Sets (2) and (3) were based on haplotypes inferred

from marker array genotypes (GGP-Porcine HD BeadChip; GeneSeek, Lincoln, NE),

which were phased using AlphaPhase [38] and imputed using AlphaImpute [39].

Most sequenced pigs, as well as pedigree relatives, were also genotyped with

marker arrays either at low density (15k markers) using the GGP-Porcine LD

BeadChip (GeneSeek) or at high density (80k markers) using the GGP-Porcine HD

BeadChip (GeneSeek).

Sequencing and data processing

Tissue samples were collected from ear punches or tail clippings. Genomic

DNA was extracted using Qiagen DNeasy 96 Blood & Tissue kits (Qiagen Ltd.,

Mississauga, ON, Canada). Paired-end library preparation was conducted using the

TruSeq DNA PCR-free protocol (Illumina, San Diego, CA). Libraries for

resequencing at low coverage (1 to 5x) were produced with an average insert size of

350 bp and sequenced on a HiSeq 4000 instrument (Illumina). Libraries for

resequencing at high coverage (15 or 30x) were produced with an average insert size

of 550 bp and sequenced on a HiSeq X instrument (Illumina). All libraries were

sequenced at Edinburgh Genomics (Edinburgh Genomics, University of Edinburgh,

Edinburgh, UK).

DNA sequence reads were pre-processed using Trimmomatic [40] to remove

adapter sequences. The reads were then aligned to the reference genome *Sscrofa11.1*

(GenBank accession: GCA_000003025.6) using the BWA-MEM algorithm [41].

Duplicates were marked with Picard (http://broadinstitute.github.io/picard). Single

nucleotide polymorphisms (SNPs) and short insertions and deletions (indels) were

identified with GATK HaplotypeCaller (GATK 3.8.0) [42,43] using default settings.

Variant discovery was performed separately for each individual and then a joint

variant set for each population was obtained by extracting the variant positions from

all the individuals in it. Between 20 and 30 million variants were discovered in each

population.

We extracted the read counts supporting each allele directly from the aligned

reads stored in the BAM files using a pile-up function. This approach was set to avoid

biases towards the reference allele introduced by GATK when applied on low-

coverage whole-genome sequence data [44]. That pipeline uses pysam (version

0.13.0; https://github.com/pysam-developers/pysam), which is a wrapper around

htslib and the samtools package [45]. We extracted the read counts for all biallelic

variant positions, after filtering variants in potential repetitive regions with VCFtools

[46]. Such variants were here defined as variants that had mean depth values 3 times

greater than the average realized coverage. A total of 46,344,624 biallelic variants

passed quality control criteria across all lines (see Supplementary Methods).

Genotype imputation

Genotypes were jointly called, phased and imputed for a total of 537,257

pedigree-related individuals using the 'hybrid peeling' method implemented in

AlphaPeel [47–49], which used all available marker array and whole-genome

sequence data. Imputation was performed separately for each line using its complete

multi-generational pedigree, which encompassed from 15,495 to 122,753 individuals

each (Table 1). We have previously published reports on the accuracy of imputation

in the same populations using this method [48]. The estimated average individual-

wise dosage correlation was 0.94 (median: 0.97). Individuals with low predicted

imputation accuracy were removed before further analyses. An individual was

predicted to have low imputation accuracy if itself or all of its grandparents were not

genotyped with a marker array or if it had a low degree of connectedness to the rest of

the population. These criteria were based on the analysis of simulated and real data on

imputation accuracy [48]. A total of 440,610 individuals remained, from 5,247 to

104,661 individuals for each line (Table 1). The expected average individual-wise

dosage correlation of the remaining individuals was 0.97 (median: 0.98) according to

our previous estimates. We accounted for the whole minor allele frequency spectrum

in our analyses. However, variants with a minor allele frequency lower than 0.023 had

an estimated variant-wise dosage correlations lower than 0.90 [48].

Variant predicted consequence types

The frequency of the alternative allele was calculated based on the imputed

genotypes. We defined the 'prevalence level' of each variant as the number of lines in

which the variant segregated. To distinguish between allele frequency and prevalence

level we used the terms 'rare' and 'common' to refer to variants in terms of allele

frequency and 'private' and 'widespread' in terms of prevalence level, where private

variants were those called only in one line and widespread variants those called in all

nine studied lines. We calculated Wright's fixation statistic (F_{ST}) [50] for each variant

among the lines where the variant segregated as $F_{ST} = (H_T–H_S)/H_T$, where H_T is the

expected heterozygosity across the combined lines assuming Hardy-Weinberg

equilibrium and H_S is the average heterozygosity within lines assuming Hardy-

Weinberg equilibrium.

Variants were annotated using Ensembl Variant Effect Predictor (Ensembl

VEP; version 97, July 2019) [51] using both Ensembl and RefSeq transcript

databases. For variants with multiple predicted consequence types (e.g., in the case of

multiple transcripts), the most severe predicted consequence type for each variant was

retrieved. Stop-gain, start-loss, stop-loss, splice donor, and splice acceptor variants

were classified as LOF variants. While frameshift indels are typically included in the

LOF category, we considered them as a separate category in order to quantify their

impact separately. The SIFT scores [52] for missense variants were retrieved from the

Ensembl transcript database. Missense variants for which SIFT scores were available

were then classified either as 'deleterious' when their SIFT score were less than 0.05,

or 'tolerated' otherwise. We considered the predicted consequence types of LOF,

frameshift and in-frame indels, and missense as putatively functional variants. To

account for the regulatory role of promoters, we classified variants within 500 bp

upstream of the annotated transcription start site together with the variants in the 5'

untranslated region (UTR). This was motivated because both regions are likely to

contain regulatory elements that affect transcription and because the same variant can

be simultaneously in the promoter or in the 5' UTR of different annotated transcripts

for the same gene. With this action, 6.6% of the variants that were initially classified

by Ensembl VEP as 'variants upstream of gene' were reclassified as 'variants in

promoter regions'. For further analyses, variants in promoters and in the 5' and 3'

UTR were jointly considered (Promoter+UTR). Because some variants such as stop-

gain (LOF) variants or frameshift indels are considered more likely to be benign when

located towards the end of the transcripts (e.g., [53]), we analysed the relative position

of these variants within transcripts (i.e., position accounting for transcript length).

Load of putatively functional alleles

We used the imputed genotypes to estimate the number of alleles of each

predicted consequence type and prevalence level that an individual carried on

average. For the most common predicted consequence types, that number was

estimated from 50,000 variants sampled randomly. For tolerated missense variants,

we used the 50,000 variants with the highest SIFT scores. To account for the different

number of variants within each predicted consequence type and prevalence level

category, we calculated 'heterozygosity' as the percentage of variants of each

category that an individual carried in heterozygosis, and the 'homozygosity for the

alternative allele' as the percentage of variants of each category that an individual

carried in homozygosis for the alternative allele.

Association to production traits

To further explore the association of variants by prevalence level and

functional annotation to selected traits, we performed genome-wide association

studies (GWAS) for the three largest lines. For each line, we performed GWAS for

average daily gain, backfat thickness, and loin depth using all the called variants that

passed filtering (Table 2). These three traits were chosen because they are complex

traits with moderate heritability estimates (range: 0.21 to 0.38). The number of pigs

with records that were included in the GWAS are provided in Table 1. Most pigs with

records were born during the 2008–2020 period. Breeding values were estimated by

line with a linear mixed model that included polygenic and non-genetic (including

contemporary group, litter, and weight as relevant for each trait) effects. Deregressed

breeding values were obtained following the method of VanRaden et al. [54]. Only

individuals in which the trait was directly measured were retained for the GWAS. We

fitted a univariate linear mixed model that accounted for the genomic relationship as:

$$\mathbf{y} = \mathbf{x}_i\beta_i + \mathbf{u} + \mathbf{e},$$

where $\mathbf{y}$ is the vector of deregressed breeding values, $\mathbf{x}_i$ is the vector of genotypes for

the ith variant coded as 0 and 2 if homozygous for either allele or 1 if heterozygous,

β_i is the allele substitution effect of the ith variant on the trait, $\mathbf{u}{\sim}N(0, \sigma_u^2\mathbf{K})$ is the

vector of polygenic effects with the covariance matrix equal to the product of the

polygenic additive genetic variance σ_u^2 and the genomic relationship matrix $\mathbf{K}$, and $\mathbf{e}$

is a vector of uncorrelated residuals. Due to computational limitations, the genomic

relationship matrix $\mathbf{K}$ was calculated using only imputed genotypes for the high-

density marker array and its single-value decomposition was taken. We used the

FastLMM software [55,56] to fit the model.

We considered the associations with a p-value equal or smaller than 10^{-6} as

significant. We calculated an enrichment score for each predicted consequence type

and prevalence level category as:

$$263 \qquad \log\left(\frac{\text{nSignCategory}/\text{nNotSignCategory}}{\text{nSignTotal}/\text{nNotSignTotal}}\right),$$

where nSignCategory was the number of variants with significant association (with at

least one trait in one of the three lines) in a predicted consequence type and

prevalence level category, nNotSignCategory was the number of variants with no

significant association in the same category, and nSignTotal and nNotSignTotal were

the total numbers of variants with and without significant association, respectively.

Linkage disequilibrium is pervasive between nearby significant variants due to

the extremely high variant density of whole-genome sequence data. To account for

this, we defined haplotype blocks so that only a single variant per haplotype block

was considered as the putative driver of the association detected in that region. We

defined the haplotype blocks for each line separately using the *--blocks* function in

Plink 1.9 [57,58]. To define haplotype blocks, pairs of variants within 5 Mb of each

other were considered to be in strong linkage disequilibrium if the bottom of the 90%

confidence interval of D' was greater than 0.7 and the top of the confidence interval

was at least 0.9. If the top of the confidence interval was smaller than 0.7, it was

considered as strong evidence for historical recombination between the two variants.

The remaining pairs of variants were considered uninformative. Regions where at

least 90% of the informative pairs showed strong linkage disequilibrium were defined

as a haplotype block. Within each haplotype block, we selected one 'candidate

variant' as the variant with the most severe predicted consequence type. If there was

more than one variant with the same predicted consequence type, the one with the

lowest p-value was selected. This process was performed separately for each trait and

line. Establishing which of the variants in linkage disequilibrium is the most likely to

be causal remains one of the greatest challenges in genomics. Nevertheless, keeping

the most severe variant in a haplotype block is a common assumption for

prioritisation of candidate variants.

We calculated the additive genetic variance explained by each variant as

$2pq\beta^2$, where p and q were the allele frequencies and β was the estimated allele

substitution effect of the variant. We expressed the additive genetic variance

explained by each variant as a percentage of the phenotypic variance of each trait.

Finally, we calculated the median F_{ST} of the candidate variants within each predicted

consequence type and prevalence level category. We compared the median F_{ST} of the

candidate variants to the median F_{ST} of the same category as the logarithm of the ratio

of the former to the latter.

Results

Prevalence of variants

A large percentage (21%) of the 46,344,624 biallelic variants that passed

quality control criteria were widespread across all nine lines. Private variants

represented a much smaller percentage (2 to 11%) of the variants called within each

line. However, when counted across lines, private variants cumulatively predominated

(28%) over the widespread ones. Most variants were neither private nor widespread.

The distribution of these variants by line is shown in Table 2. Most variants

(38,642,777) were SNPs, of which 10,595,681 were called in a single line (27%;

366,486 to 2,743,965 within each line) and 8,377,578 (22%) were called in all nine

lines. The remaining 7,701,847 variants were indels, of which 2,436,674 were called

in a single line (32%; 121,525 to 506,149 in each line) and 1,560,353 (20%) were

called in all nine lines.

Distribution of variants and relationship with recombination rate

The number of variants by chromosome was strongly correlated with
chromosome length (r=0.98, P<0.05; Table S1). The average variant density by
chromosome was negatively correlated with chromosome length (r=–0.87, P<0.05;
Table S1). The distribution of variants within chromosomes was positively correlated
to recombination rate (r=0.65, P<0.05, between variant density and recombination
rate in 1-Mb non-overlapping windows [59]; Figure 1a). For example, within line A,
there was on average one variant every 81 bp, but in the 5% 1-Mb windows with the
lowest and highest recombination rates there was on average one variant every 152
and 54 bp, respectively (2.8-fold more variants in windows with high recombination
rate). Across all lines, there was one variant every 49 bp on average, but in the 5% 1-
Mb windows with the lowest and highest recombination rates there was on average
one variant every 79 and 34 bp, respectively (2.3-fold more variants in windows with
high recombination rate).

The distribution of private and widespread variants along the genome also
differed. The distribution of widespread variants was more correlated with
recombination rate than that of private variants (Figures 1b and 1c). As a
consequence, private variants represented a larger proportion of the variation in
regions with low recombination rate, which were depleted of widespread variants.
Across all lines, in the 5% 1-Mb windows with the highest recombination rates there
was on average one private variant every 167 bp and one widespread variant every
148 bp (1.1-fold more private variants relative to widespread). In the 5% 1-Mb
windows with the lowest recombination rates there was on average one private variant
every 260 bp and one widespread variant every 531 bp (2.0-fold more private variants

relative to widespread). There were no genomic regions that were enriched for private

variants across lines (Figure S2).

Frequency and fixation index

The prevalence level and alternative allele frequency were related, in a way

that less prevalent variants had also lower allele frequency (Figure 2) and lower F_{ST}

(Figure 3). Private variants had an average alternative allele frequency of 0.03 (SD

0.09) as opposed to widespread variants, which had an average alternative allele

frequency of 0.50 (SD 0.25). As a consequence of the less prevalent variants generally

having low frequencies in the lines where they segregated, these variants showed a

small degree of differentiation between the lines in which they segregated (F_{ST}=0.04,

SD=0.07). In contrast, the widespread variants allowed for the largest degree of

differentiation between lines (F_{ST}=0.21, SD=0.11).

Prevalence and frequency of putatively functional variants

The predicted consequence types of the variants are shown in Table 3. Half

(49.9%) of the variants were called in intergenic regions and another 47.0% of the

variants were called in intronic regions. Only 2.2% of the variants were called in the

promoter or 5' and 3' UTR. The coding variants comprised 0.9% of the total variants,

of which more than half were missense (45.5%), frameshift indels (3.1%) or LOF

(3.7%). The density of putatively functional variants was only weakly correlated to

recombination rate (Figures 1d).

The low-prevalence variants (i.e., the variants that were identified in one or

few lines) were enriched with missense and LOF variants, as well as potentially

regulatory variants such as those located in the promoter and 5' and 3' UTR and other

intronic variants. On the other hand, the high-prevalence variants (i.e., the variants

that were identified in many or all lines) were enriched with frameshift indels,

synonymous (non-significant correlation), and intergenic variants. Frameshift indels

are typically included in the LOF category. However, our results show that the LOF

category is very heterogeneous and the frameshift indels presented opposite patterns

to other LOF variants. Therefore, we studied them as a separate category.

Whereas the LOF variants had lower allele frequencies than the intergenic

variants in low-prevalence levels, they had similar allele frequencies in high-

prevalence levels (Table 4). Thus, there was a set of LOF variants that were prevalent

across lines and also had particularly high frequencies within lines. The missense

variants, especially those classified as deleterious, had lower allele frequencies than

the intergenic variants for all prevalence levels. The low-prevalence missense variants

were enriched with a larger fraction of deleterious variants and lower SIFT scores

(Figure 4). Low-prevalence stop-gain (LOF) variants and frameshift indels, unlike

missense or synonymous variants, were more likely to occur towards the start of the

transcripts (Figure 5). As opposed to LOF and missense variants, the frameshift and

in-frame indels had intermediate allele frequencies that were much higher than those

of the intergenic variants (Table 4), which indicated that in many cases the minor

allele was the reference one. Within prevalence level, the LOF and deleterious

missense variants had lower F_{ST} than the intergenic variants (Table 5), probably

because they were kept at low allele frequencies due to negative selection pressure.

The frameshift and in-frame indels also had lower F_{ST} than the intergenic variants

despite their intermediate allele frequencies.

Load of putatively functional alleles by prevalence level

Most of the missense deleterious and LOF variants that an individual carried

in homozygosis for the alternative allele were high-prevalence variants. Only a small

proportion of these variants were private. An individual carried on average 1,048 (SD

57) LOF variants in homozygosis for the alternative allele, of which 713 (SD 36)

were widespread across all nine lines and only 20 (SD 7) were private. An average

individual carried 1,379 (SD 165) deleterious missense variants in homozygosis for

the alternative allele, of which 1,012 (SD 79) were widespread and only 4 (SD 3)

were private. An average individual carried 1,080 (SD 89) LOF and 2,632 (SD 235)

deleterious missense variants in heterozygosis.

We found signals of negative selection against deleterious missense variants,

in particular the private ones. Individuals proportionally carried less deleterious

missense variants in homozygosis for the alternative allele than variants of other

predicted consequence types, regardless of prevalence level (Figure 6). Individuals

also carried proportionally less private tolerated missense, synonymous and LOF

variants in homozygosis for the alternative allele than expected, but not in

heterozygosis.

Association of low-prevalence variants to production traits

Significant variants were enriched with putatively functional and regulatory

variants of different prevalence levels, and depleted of intergenic variants. A total of

108,109 variants were significantly associated to at least one trait in one line. Figures

7a and 7b summarise the enrichment scores for all significant variants. The predicted

consequence types that reached the greatest enrichment scores were LOF, frameshift

indels, and unclassified missense variants, with various prevalence levels. Variants

with intermediate prevalence levels were amongst the most enriched. These trends

were accentuated after selecting candidate variants from haplotype blocks. In each

line we defined from 1,554 to 2,118 haplotype blocks. A total of 6,692 candidate

variants remained after accounting for linkage disequilibrium within each haplotype

block for all lines and traits. Figures 7c and 7d summarise the enrichment scores for

the candidate variants. The enrichment scores based on the candidate variants

revealed a stronger depletion of intergenic variants, as well as intronic (with the

exception of high-prevalence), and a much stronger enrichment for LOF, frameshift

indels, and missense variants. For putatively functional variants, there were no clear

trends of their enrichment scores across prevalence levels. The trends of the

enrichment scores between predicted consequence types and prevalence levels were

similar in the three tested traits.

In general, the lower allele frequency of low-prevalence variants hindered the

detection of significant associations for these markers. Low-prevalence variants that

were detected as significantly associated to the production traits actually had

intermediate allele frequencies that were greater than expected for their prevalence

level. Low-prevalence variants in general explained low percentages of variance

(Figure 8), although there were some instances of low-prevalence variants that

explained up to 3.2% of phenotypic variance. Significant variants had higher F_{ST} than

other variants of the same predicted consequence type and prevalence level (Figure

9). This enrichment was especially strong for low-prevalence variants, which in some

instances reached F_{ST} around 0.15.

Discussion

428   Our results contextualize the importance of population-specific and low-

429   prevalence genetic variants. Next, we will discuss: (1) the distribution and functional

430   annotation of low-prevalence variants, (2) the load of putatively functional alleles by

431   prevalence level, and (3) the association of low-prevalence variants to production

432   traits.

Distribution and functional annotation of low-prevalence variants

The main difficulty for the study of low-prevalence genetic variants is that the

prevalence of a variant across several lines is strongly related to its allele frequency,

in a way that the low-prevalence variants are also rare within the lines where they

occur. This is possibly because low-prevalence variants are relatively recent or are

constrained by negative selection.

On one hand, the distribution of private variants was only weakly correlated to

recombination rate and, therefore, regions with low recombination rate were enriched

for private variants. Although the interplay between recurring sweeps, background

selection and other phenomena at play is not fully understood yet, it is generally

accepted that selection on linked variants leads to loss of variation in regions with low

recombination rates [60]. Our observation that regions with low recombination rate

were enriched for private variants suggests that private variants may have been less

affected by selective sweeps than widespread variants. This would be consistent with

previous observations of the younger age of rare and low-prevalence variants [61],

and suggests that many private variants arose more recently in time than widespread

variants, likely after line differentiation, and accumulated in low-recombining regions

due to the reduced efficacy of purifying selection in those regions [62,63].

On the other hand, the low-prevalence variants were enriched for putatively

functional variants and with signs of a greater severity (stop-gain and frameshift

indels that occur earlier in the transcript, and missense variants predicted to be

deleterious). Variants that affect performance traits or that cause a detrimental

condition are under the action of directional selection and are therefore driven towards

loss or fixation [64,65]. The low F_{ST} estimates for the low-prevalence variants

indicated that selection pressure keeps these variants at low minor allele frequency

even when they occur in several lines, especially if they are putatively functional [66].

This could be caused by natural selection or similar selection objectives across

livestock populations. These observations were also consistent with previous reports

showing that some putatively functional variant categories (such as stop-gain and

deleterious missense) were enriched for variants that were private to single cattle

breeds [33], that putatively functional variants were less likely to have high frequency

of the alternative allele across multiple chicken lines [35], and that population-specific

variants in non-African humans were enriched with putatively functional variants

[67].

The relationship between variant prevalence across lines and allele frequency

highlighted the suitability of using a low-coverage sequencing approach to study this

fraction of genetic variation. Nonetheless, bioinformatics pipelines for calling,

genotyping and, even imputing such variants should account for the increased

uncertainty associated to their low allele frequency. We decided on using a very

relaxed variant calling strategy with little filtering to account for as many rare variants

as possible, but a sizeable fraction of these rare variants were discarded after

imputation because they were fixed for the imputed individuals that passed quality

control. Low-coverage sequencing is also unsuitable for other types of genetic

variants, such as structural variations (CNVs, tandem duplications, and inversions),

which could also be putatively functional and population-specific [68]. Of course, the

number of called variants and the proportion that were private or widespread depends

of the number of sequenced lines [32,35] as well as the sequencing effort in each line.

Our results also suggest that what is typically grouped as LOF is actually a

heterogeneous category. In particular, frameshift indels showed patterns that did not

conform to the other predicted consequence types.

Load of putatively functional alleles by prevalence level

We found that an average individual carried a larger number of LOF and

missense deleterious variants than previously reported in other livestock species or in

humans. However, there is not yet a clear consensus on the number of LOF and

deleterious missense alleles that are present in the genome of an average individual. In

humans, it has been estimated that an average individual carries 100-150 LOF alleles

[64,69–71] and around 800 weakly deleterious mutations [72], most of which are rare.

In domestic livestock populations, the number of LOF and deleterious alleles carried

on average by individuals has been reported to be greater than in wild populations

[73], including estimates of 100 to 300 deleterious variants in domestic pigs [74], over

400 deleterious variants in domestic chicken [74], and 1,200-1,500 deleterious

variants in domestic yak [75]. Similar magnitudes have been reported in dogs [76],

rice [77], and sunflower [63].

It has been debated why healthy individuals carry a larger number of LOF

variants in homozygosis than expected [78,79]. These could be driven by the fact that

not all predicted LOF variants are detrimental and their functional impact should be

validated before being considered as such. Many predicted LOF variants are in fact

neutral, advantageous (either in the wild or in controlled production environments), or

even may arise simply because of sequencing and annotation errors [78]. This claim is

supported by the large proportion of LOF observed in homozygosis for the alternative

allele compared to the other consequence types, which casts doubt on the real impact

of those variants. On the contrary, individuals carried a lower proportion of alleles

predicted to be deleterious missense in homozygosis, which supports that variants

predicted as such may have a real impact on genetic variation of production traits and,

therefore, be subjected to selection pressure.

These observations have implications for the identification of variants to be

used for genomic prediction or genomic edition strategies such as PAGE [26] or

RAGE [27]. Efforts to promote or remove alleles should target variants that make a

substantial contribution to traits of interest, namely functional variants. However, it is

hard to computationally predict and statistically estimate the effects of such variants,

especially if they have low allele frequency. The number of LOF variants in

homozygosis for the alternative allele suggests that predicted loss of function is not a

good indicator that a variant is strongly deleterious in the context of livestock

breeding. Similarly, bioinformatics predictors of missense variant effects appear to be

not very accurate [80,81]. High-throughput fine-mapping and variant screening would

be needed to ascertain variant causality and disentangle causality from linkage

disequilibrium.

Association of low-prevalence variants to production traits

Genome-wide association studies on three polygenic traits of economical

importance in the three largest lines revealed that the significant markers were

enriched for putatively functional roles, such as LOF, frameshift indels and missense

variants, and depleted of intergenic variants. This pattern of enrichment was similar to

previous reports from human datasets [82]. However, only a few of the population-

specific and low-prevalence variants were significantly associated to the traits, even

after accounting for linkage disequilibrium. Most of the significant variants showed

intermediate or high prevalence levels. These observations are consistent with

previous meta-analyses in cattle that showed that significant variants are often

common variants [83]. This could be explained by either the fact that quantitative trait

nucleotides have intermediate or high allele frequencies or the fact that most studies

are underpowered to map rare causal variants. The latter scenario still seems more

likely given that the significant private and low-prevalence variants had intermediate

allele frequencies. This also supports that these significant variants have biological

functions that contributed to trait phenotypic variance rather that they reached

intermediate allele frequencies by drift or by hitchhiking with linked variants under

selection [84]. However, these amounted to a small number of variants that explained

small fractions of variance. Other more widespread variants, including intergenic

variants, successfully acted as tag variants for them and captured much larger

fractions of trait variance. This makes them more suitable for applications in animal

breeding, as is already the case with marker arrays. A similar result was found in

cattle, where splice site and synonymous variants explained the largest proportions of

trait variance, while missense variants explained almost null variance [85]. It is worth

pointing out that even a variant with a large allele substitution effect will explain a

small percentage of variance if the alternative allele is rare.

It can be hypothesized that some of the low-prevalence variants with low allele

frequency have non-negligible effects for traits of interest. Despite the large amount

of individuals included in this study, the large volume of variants and the

pervasiveness of linkage disequilibrium among them still make it very challenging to

disentangle their contribution to trait variance. While genome-wide association

studies involving more than one breed typically find multiple breed-specific

associations (e.g., [86]), based on our results it seems unlikely that breed-specific

associations arise from the low-prevalence variants. They would instead stem from

differences in allele frequency, linkage disequilibrium structure or genetic background

that affect the power to detect the effect of prevalent variants across different

populations. Significant variants were enriched with higher F_{ST} estimates than non-

significant variants, which is also consistent with previous reports [83]. Although the

enrichment was greater for low-prevalence variants, it remains unclear to which

degree these variants could relate to selection history or explain differences among

lines for the studied traits.

Conclusion

Low-prevalence variants are enriched for putatively functional variants,

including LOF and deleterious missense variants. However, most low-prevalence

variants are kept at very low allele frequency. Only a small subset of low-prevalence

variants was found at intermediate allele frequencies and had large estimated effects

on production traits. Population-specific variants that were significantly associated to

complex traits had greater degrees of differentiation than non-significant variants in

the same category. However, more widespread variants, including intergenic variants,

successfully captured larger fractions of trait variance. Therefore, overall, not

accounting for population-specific and other low-prevalence variants is unlikely to

hinder across-breed analyses, such as the prediction of genomic breeding values using

reference populations of a different genetic background.

Ethics approval and consent to participate

576 The samples used in this study were derived from the routine breeding activities of

577 PIC.

Consent for publication

578 Not applicable.

Availability of data and material

579 The software packages AlphaPhase, AlphaImpute, and AlphaPeel are available from

580 https://github.com/AlphaGenes. The software package AlphaSeqOpt is available from

581 the AlphaGenes website (http://www.alphagenes.roslin.ed.ac.uk). The datasets

582 generated and analysed in this study are derived from the PIC breeding programme

583 and not publicly available.

Competing interests

584 BDV, CYC, and WOH are employed by Genus PIC. The remaining authors declare

585 that the research was conducted in the absence of potential conflicts of interest.

Funding

The authors acknowledge the financial support from the BBSRC ISPG to The Roslin

Institute (BBS/E/D/30002275), from Genus plc, Innovate UK (grant 102271), and

from grant numbers BB/N004736/1, BB/N015339/1, BB/L020467/1, and

BB/M009254/1. MJ acknowledges financial support from the Swedish Research

Council for Sustainable Development Formas Dnr 2016-01386. For the purpose of

open access, the authors have applied a Creative Commons Attribution (CC BY)

licence to any author accepted manuscript version arising from this submission.

Authors' contributions

593　RRF, MJ, and JMH designed the study; RRF and MJ performed the analyses; RRF

594　and MJ wrote the first draft; BDV, CYC, WHO, GG, and JMH contributed to the

595　interpretation of the results and provided comments on the manuscript. All authors

596　read and approved the final manuscript.

Acknowledgements

597　This work has made use of the resources provided by the Edinburgh Compute and

598　Data Facility (ECDF) (http://www.ecdf.ed.ac.uk/).

References

1. Xiang R, Berg I van den, MacLeod IM, Hayes BJ, Prowse-Wilkins CP, Wang M, et
al. Quantifying the contribution of sequence variants with regulatory and evolutionary
significance to 34 bovine complex traits. Proc Natl Acad Sci. 2019;116:19398–408.

2. Zhang F, Wang Y, Mukiibi R, Chen L, Vinsky M, Plastow G, et al. Genetic
architecture of quantitative traits in beef cattle revealed by genome wide association
studies of imputed whole genome sequence variants: I: feed efficiency and component
traits. BMC Genomics. 2020;21.

3. Wang Y, Zhang F, Mukiibi R, Chen L, Vinsky M, Plastow G, et al. Genetic
architecture of quantitative traits in beef cattle revealed by genome wide association
studies of imputed whole genome sequence variants: II: carcass merit traits. BMC
Genomics. 2020;21.

4. Pan Z, Yao Y, Yin H, Cai Z, Wang Y, Bai L, et al. Pig genome functional
annotation enhances the biological interpretation of complex traits and human disease.
Nat Commun. 2021;12.

5. Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, et al.
Finding the missing heritability of complex diseases. Nature. 2009;461:747–53.

6. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al.
The mutational constraint spectrum quantified from variation in 141,456 humans.
Nature. 2020;581:434–43.

7. Van Hout CV, Tachmazidou I, Backman JD, Hoffman JD, Liu D, Pandey AK, et
al. Exome sequencing and characterization of 49,960 individuals in the UK Biobank.
Nature. 2020;586:749–56.

8. Grobet L, Royo Martin LJ, Poncelet D, Pirottin D, Brouwers B, Riquet J, et al. A
deletion in the bovine myostatin gene causes the double-muscled phenotype in cattle.
Nat Genet. 1997;17:71–4.

9. Grisart B, Farnir F, Karim L, Cambisano N, Kim J-J, Kvasz A, et al. Genetic and
functional confirmation of the causality of the DGAT1 K232A quantitative trait
nucleotide in affecting milk yield and composition. Proc Natl Acad Sci.
2004;101:2398–403.

10. Óvilo C, FernáNdez A, Noguera JL, BarragáN C, LetóN R, RodríGuez C, et al.
Fine mapping of porcine chromosome 6 QTL and *LEPR* effects on body composition
in multiple generations of an Iberian by Landrace intercross. Genet Res. 2005;85:57–
67.

11. Zhao H, Qin Y, Xiao Z, Li Q, Yang N, Pan Z, et al. Loss of Function of an RNA
Polymerase III Subunit Leads to Impaired Maize Kernel Development. Plant Physiol.
2020;184:359–73.

12. Stenson PD, Ball EV, Mort M, Phillips AD, Shiel JA, Thomas NST, et al. Human
Gene Mutation Database (HGMD®): 2003 update. Hum Mutat. 2003;21:577–81.

13. Drögemüller C, Tetens J, Sigurdsson S, Gentile A, Testoni S, Lindblad-Toh K, et
al. Identification of the Bovine Arachnomelia Mutation by Massively Parallel
Sequencing Implicates Sulfite Oxidase (SUOX) in Bone Development. Georges M,
editor. PLoS Genet. 2010;6:e1001079.

14. Waide EH, Dekkers JCM, Ross JW, Rowland RRR, Wyatt CR, Ewen CL, et al.
Not All SCID Pigs Are Created Equally: Two Independent Mutations in the *Artemis*
Gene Cause SCID in Pigs. J Immunol. 2015;195:3171–9.

15. Derks MFL, Harlizius B, Lopes MS, Greijdanus-van der Putten SWM, Dibbits B,
Laport K, et al. Detection of a Frameshift Deletion in the SPTBN4 Gene Leads to
Prevention of Severe Myopathy and Postnatal Mortality in Pigs. Front Genet.
2019;10.

16. Matika O, Robledo D, Pong-Wong R, Bishop SC, Riggio V, Finlayson H, et al.
Balancing selection at a premature stop mutation in the myostatin gene underlies a
recessive leg weakness syndrome in pigs. Andersson L, editor. PLOS Genet.
2019;15:e1007759.

17. Nicholas FW. Online Mendelian Inheritance in Animals (OMIA): a record of
advances in animal genetics, freely available on the Internet for 25 years. Anim
Genet. 2021;52:3–9.

18. Derks MFL, Gjuvsland AB, Bosse M, Lopes MS, van Son M, Harlizius B, et al.
Loss of function mutations in essential genes cause embryonic lethality in pigs. PLOS
Genet. 2019;15:e1008055.

19. Mesbah-Uddin M, Hoze C, Michot P, Barbat A, Lefebvre R, Boussaha M, et al. A
missense mutation (p.Tyr452Cys) in the CAD gene compromises reproductive
success in French Normande cattle. J Dairy Sci. 2019;102:6340–56.

20. Ma J, Yang J, Zhou L, Ren J, Liu X, Zhang H, et al. A Splice Mutation in the
PHKG1 Gene Causes High Glycogen Content and Low Meat Quality in Pig Skeletal
Muscle. PLoS Genet. 2014;10:e1004710.

21. Lunden A. A Nonsense Mutation in the FMO3 Gene Underlies Fishy Off-Flavor
in Cow's Milk. Genome Res. 2002;12:1885–8.

22. Joseph SB, Hall DW. Spontaneous Mutations in Diploid Saccharomyces
cerevisiae. Genetics. 2004;168:1817–25.

23. Pérez-Enciso M, Rincón JC, Legarra A. Sequence- vs. chip-assisted genomic
selection: accurate biological information is advised. Genet Sel Evol. 2015;47:43.

24. MacLeod IM, Bowman PJ, Vander Jagt CJ, Haile-Mariam M, Kemper KE,
Chamberlain AJ, et al. Exploiting biological priors and sequence variants enhances
QTL discovery and genomic prediction of complex traits. BMC Genomics.
2016;17:144.

25. Lopez BIM, An N, Srikanth K, Lee S, Oh J-D, Shin D-H, et al. Genomic
Prediction Based on SNP Functional Annotation Using Imputed Whole-Genome
Sequence Data in Korean Hanwoo Cattle. Front Genet. 2021;11:603822.

26. Jenko J, Gorjanc G, Cleveland MA, Varshney RK, Whitelaw CBA, Woolliams
JA, et al. Potential of promotion of alleles by genome editing to improve quantitative
traits in livestock breeding programs. Genet Sel Evol. 2015;47:55.

27. Johnsson M, Gaynor RC, Jenko J, Gorjanc G, de Koning D-J, Hickey JM.
Removal of alleles by genome editing (RAGE) against deleterious load. Genet Sel
Evol. 2019;51:14.

28. Le SQ, Durbin R. SNP detection and genotyping from low-coverage sequencing
data on multiple diploid samples. Genome Res. 2011;21:952–60.

29. Martin AR, Atkinson EG, Chapman SB, Stevenson A, Stroud RE, Abebe T, et al.
Low-coverage sequencing cost-effectively detects known and novel variation in
underrepresented populations. Am J Hum Genet. 2021;108:656–68.

30. Molnár J, Nagy T, Stéger V, Tóth G, Marincs F, Barta E. Genome sequencing and
analysis of Mangalica, a fatty local pig of Hungary. BMC Genomics. 2014;15:761.

31. Choi J-W, Chung W-H, Lee K-T, Cho E-S, Lee S-W, Choi B-H, et al. Whole-
genome resequencing analyses of five pig breeds, including Korean wild and native,
and three European origin breeds. DNA Res. 2015;22:259–67.

32. Cai Z, Sarup P, Ostersen T, Nielsen B, Fredholm M, Karlskov-Mortensen P, et al.
Genomic diversity revealed by whole-genome sequencing in three Danish commercial
pig breeds. J Anim Sci. 2020;98.

33. Daetwyler HD, Capitan A, Pausch H, Stothard P, van Binsbergen R, Brondum RF,
et al. Whole-genome sequencing of 234 bulls facilitates mapping of monogenic and
complex traits in cattle. Nat Genet. 2014;46:858–65.

34. Das A, Panitz F, Gregersen VR, Bendixen C, Holm L-E. Deep sequencing of
Danish Holstein dairy cattle for variant detection and insight into potential loss-of-
function variants in protein coding genes. BMC Genomics. 2015;16:1043.

35. Gheyas AA, Boschiero C, Eory L, Ralph H, Kuo R, Woolliams JA, et al.
Functional classification of 15 million SNPs detected from diverse chicken
populations. DNA Res. 2015;22:205–17.

36. Gonen S, Ros-Freixedes R, Battagin M, Gorjanc G, Hickey JM. A method for the
allocation of sequencing resources in genotyped livestock populations. Genet Sel
Evol. 2017;49:47.

37. Ros-Freixedes R, Gonen S, Gorjanc G, Hickey JM. A method for allocating low-
coverage sequencing resources by targeting haplotypes rather than individuals. Genet
Sel Evol. 2017;49:78.

38. Hickey JM, Kinghorn BP, Tier B, Wilson JF, Dunstan N, van der Werf JH. A
combined long-range phasing and long haplotype imputation method to impute phase
for SNP genotypes. Genet Sel Evol. 2011;43:12.

39. Hickey JM, Kinghorn BP, Tier B, van der Werf JH, Cleveland MA. A phasing
and imputation method for pedigreed populations that results in a single-stage
genomic evaluation. Genet Sel Evol. 2012;44:9.

40. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30:2114–20.

41. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv. 2013;1303.3997v1 [q – bio.GN].

42. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A
framework for variation discovery and genotyping using next-generation DNA
sequencing data. Nat Genet. 2011;43:491–8.

43. Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van der
Auwera GA, et al. Scaling accurate genetic variant discovery to tens of thousands of
samples. bioRxiv. 2018;10.1101/201178.

44. Ros-Freixedes R, Battagin M, Johnsson M, Gorjanc G, Mileham AJ, Rounsley
SD, et al. Impact of index hopping and bias towards the reference allele on accuracy
of genotype calls from low-coverage sequencing. Genet Sel Evol. 2018;50:64.

45. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

46. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The
variant call format and VCFtools. Bioinformatics. 2011;27:2156–8.

47. Whalen A, Ros-Freixedes R, Wilson DL, Gorjanc G, Hickey JM. Hybrid peeling
for fast and accurate calling, phasing, and imputation with sequence data of any
coverage in pedigrees. Genet Sel Evol. 2018;50:67.

48. Ros-Freixedes R, Whalen A, Chen C-Y, Gorjanc G, Herring WO, Mileham AJ, et
al. Accuracy of whole-genome sequence imputation using hybrid peeling in large
pedigreed livestock populations. Genet Sel Evol. 2020;52:17.

49. Ros-Freixedes R, Whalen A, Gorjanc G, Mileham AJ, Hickey JM. Evaluation of
sequencing strategies for whole-genome imputation with hybrid peeling. Genet Sel
Evol. 2020;52:18.

50. Wright S. The genetical structure of populations. Ann Eugen. 1949;15:323–54.

51. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The
Ensembl Variant Effect Predictor. Genome Biol. 2016;17:122.

52. Ng PC, Henikoff S. SIFT: predicting amino acid changes that affect protein
function. Nucleic Acids Res. 2003;31:3812–4.

53. Torella A, Zanobio M, Zeuli R, del Vecchio Blanco F, Savarese M, Giugliano T,
et al. The position of nonsense mutations can predict the phenotype severity: A survey
on the DMD gene. Singh RN, editor. PLOS ONE. 2020;15:e0237803.

54. VanRaden PM, Van Tassell CP, Wiggans GR, Sonstegard TS, Schnabel RD,
Taylor JF, et al. Invited review: reliability of genomic predictions for North American
Holstein bulls. J Dairy Sci. 2009;92:16–24.

55. Lippert C, Listgarten J, Liu Y, Kadie CM, Davidson RI, Heckerman D. FaST
linear mixed models for genome-wide association studies. Nat Methods. 2011;8:833–
5.

56. Widmer C, Lippert C, Weissbrod O, Fusi N, Kadie C, Davidson R, et al. Further
Improvements to Linear Mixed Models for Genome-Wide Association Studies. Sci
Rep. 2015;4:6874.

57. Taliun D, Gamper J, Pattaro C. Efficient haplotype block recognition of very long
and dense genetic sequences. BMC Bioinformatics. 2014;15.

58. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-
generation PLINK: rising to the challenge of larger and richer datasets. GigaScience.
2015;4.

59. Johnsson M, Whalen A, Ros-Freixedes R, Gorjanc G, Chen C-Y, Herring WO, et
al. Genetic variation in recombination rate in the pig. Genet Sel Evol. 2021;53.

60. Cutter AD, Payseur BA. Genomic signatures of selection at linked sites: unifying
the disparity among species. Nat Rev Genet. 2013;14:262–74.

61. Mathieson I, McVean G. Demography and the Age of Rare Variants. Novembre J,
editor. PLoS Genet. 2014;10:e1004528.

62. Charlesworth D, Morgan MT, Charlesworth B. Mutation Accumulation in Finite
Populations. J Hered. 1993;84:321–5.

63. Renaut S, Rieseberg LH. The Accumulation of Deleterious Mutations as a
Consequence of Domestication and Improvement in Sunflowers and Other
Compositae Crops. Mol Biol Evol. 2015;32:2273–83.

64. Gudbjartsson DF, Helgason H, Gudjonsson SA, Zink F, Oddson A, Gylfason A, et
al. Large-scale whole-genome sequencing of the Icelandic population. Nat Genet.
2015;47:435–44.

65. Sulem P, Helgason H, Oddson A, Stefansson H, Gudjonsson SA, Zink F, et al.
Identification of a large set of rare complete human knockouts. Nat Genet.
2015;47:448–52.

66. Mezmouk S, Ross-Ibarra J. The Pattern and Distribution of Deleterious Mutations
in Maize. G3 GenesGenomesGenetics. 2014;4:163–71.

67. Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. On the accumulation of
deleterious mutations during range expansions. Mol Ecol. 2013;22:5972–82.

68. Liu GE, Hou Y, Zhu B, Cardone MF, Jiang L, Cellamare A, et al. Analysis of
copy number variations among diverse cattle breeds. Genome Res. 2010;20:693–703.

69. The 1000 Genomes Project Consortium. An integrated map of genetic variation
from 1,092 human genomes. Nature. 2012;491:56–65.

70. MacArthur DG, Balasubramanian S, Frankish A, Huang N, Morris J, Walter K, et
al. A Systematic Survey of Loss-of-Function Variants in Human Protein-Coding
Genes. Science. 2012;335:823–8.

71. Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, et al.
Analysis of protein-coding genetic variation in 60,706 humans. Nature.
2016;536:285–91.

72. Chun S, Fay JC. Identification of deleterious mutations within three human
genomes. Genome Res. 2009;19:1553–61.

73. Makino T, Rubin C-J, Carneiro M, Axelsson E, Andersson L, Webster MT.
Elevated Proportions of Deleterious Genetic Variation in Domestic Animals and
Plants. Genome Biol Evol. 2018;10:276–90.

74. Bosse M, Megens H, Derks MFL, Cara ÁMR, Groenen MAM. Deleterious alleles
in the context of domestication, inbreeding, and selection. Evol Appl. 2019;12:6–17.

75. Xie X, Yang Y, Ren Q, Ding X, Bao P, Yan B, et al. Accumulation of deleterious
mutations in the domestic yak genome. Anim Genet. 2018;49:384–92.

76. Cruz F, Vila C, Webster MT. The Legacy of Domestication: Accumulation of
Deleterious Mutations in the Dog Genome. Mol Biol Evol. 2008;25:2331–6.

77. Lu J, Tang T, Tang H, Huang J, Shi S, Wu C-I. The accumulation of deleterious
mutations in rice genomes: a hypothesis on the cost of domestication. Trends Genet.
2006;22:126–31.

78. MacArthur DG, Tyler-Smith C. Loss-of-function variants in the genomes of
healthy humans. Hum Mol Genet. 2010;19:R125–30.

79. Rausell A, Luo Y, Lopez M, Seeleuthner Y, Rapaport F, Favier A, et al. Common
homozygosity for predicted loss-of-function variants reveals both redundant and
advantageous effects of dispensable human genes. Proc Natl Acad Sci.
2020;117:13626–36.

80. Pagel KA, Pejaver V, Lin GN, Nam H-J, Mort M, Cooper DN, et al. When loss-
of-function is loss of function: assessing mutational signatures and impact of loss-of-
function genetic variants. Bioinformatics. 2017;33:i389–98.

81. Pejaver V, Urresti J, Lugo-Martinez J, Pagel KA, Lin GN, Nam H-J, et al.
Inferring the molecular and phenotypic impact of amino acid variants with MutPred2.
Nat Commun. 2020;11.

82. Schork AJ, Thompson WK, Pham P, Torkamani A, Roddey JC, Sullivan PF, et al.
All SNPs Are Not Created Equal: Genome-Wide Association Studies Reveal a
Consistent Pattern of Enrichment among Functionally Annotated SNPs. Gibson G,
editor. PLoS Genet. 2013;9:e1003449.

83. van den Berg I, Xiang R, Jenko J, Pausch H, Boussaha M, Schrooten C, et al.
Meta-analysis for milk fat and protein percentage using imputed sequence variant
genotypes in 94,321 cattle from eight cattle breeds. Genet Sel Evol. 2020;52:37.

84. Chun S, Fay JC. Evidence for Hitchhiking of Deleterious Mutations within the
Human Genome. Pritchard JK, editor. PLoS Genet. 2011;7:e1002240.

85. Koufariotis LT, Chen Y-PP, Stothard P, Hayes BJ. Variance explained by whole
genome sequence variants in coding and regulatory genome annotations for six dairy
traits. BMC Genomics. 2018;19.

86. Purfield DC, Evans RD, Berry DP. Breed- and trait-specific associations define
the genetic architecture of calving performance traits in cattle. J Anim Sci. 2020;98.

Figures

840
841 **Figure 1.** Variant density in line A (black and grey bars) and recombination rate (red
842 line). The correlation (r) between variant density and recombination rate in 1-Mb non-
843 overlapping windows is reported.
844

845
846   **Figure 2.** Frequency of the alternative allele by prevalence level. Red dots indicate
847   means. In blue, values greater than 1.5 times the interquartile range.
848

849
850 **Figure 3.** Wright's fixation statistic (F_{ST}) by prevalence level. Red dots indicate
851 means. In blue, values greater than 1.5 times the interquartile range.
852

853
854 **Figure 4.** Classification of the missense variants and median SIFT score by
855 prevalence level.
856

a) Stop-gain (loss-of-function) variants

b) Frameshift indels

c) Missense variants

d) Synonymous variants

857
858 **Figure 5.** Relative position within transcript by prevalence level of stop-gain,
859 frameshift indels, missense, and synonymous variants.
860

861
862 **Figure 6.** Percentage of variants in homozygosis for the alternative allele or in
863 heterozygosis in an average individual by predicted consequence type and prevalence
864 level. LOF: loss-of-function; UTR: untranslated regions.
865

866
Figure 7. Enrichment scores for the significant variants in the genome-wide
association study by variant prevalence level and predicted consequence type. Either
all significant variants (panels a and b) or only the most sever significant variants
within haplotype block (panels c and d) were used. Prevalence level was considered
across all 9 lines (panels a and c) or only across the 3 lines included in the genome-
wide association study (panels b and d).
873

874
875 **Figure 8.** Maximum percentage of phenotypic variance explained by the individual
876 candidate variants within each prevalence level and predicted consequence type. Only
877 the candidate variants after accounting for linkage disequilibrium were used.
878 Prevalence level was considered across all 9 lines (panel a) or only across the 3 lines
879 included in the genome-wide association study (panel b).
880

881
882 **Figure 9.** Enrichment scores for the F_{ST} median of the candidate variants within each
883 prevalence level and predicted consequence type. Only the candidate variants after
884 accounting for linkage disequilibrium were used. Prevalence level was considered
885 across all 9 lines.
886

Tables

887   **Table 1.** Number of sequenced and analysed pigs.

Line	Individuals sequenced	Individuals sequenced by coverage				Individuals used in analyses		
		1x	2x	5x	15–30x	Pedigree	Imputed	GWAS
A	1,856	1,044	649	73	90	122,753	104,661	88,342
B	1,491	628	728	54	81	84,420	66,608	56,173
C	1,366	685	545	44	92	88,964	76,230	64,285
D	760	394	274	27	65	50,797	41,573	-
E	731	362	311	16	42	79,981	60,474	-
F	701	351	255	28	67	52,470	39,263	-
G	445	217	176	15	37	21,129	17,224	-
H	381	193	137	16	35	35,309	29,330	-
I	321	111	158	18	34	15,495	5,247	-

889 **Table 2.** Number of variants by line.

Line	Biallelic variant sites (M)	SNPs			Indels		
		All biallelic (M)	Private (M)	Widespread (M)	All biallelic (M)	Private (M)	Widespread (M)
A	28.83	24.38	1.56	8.38	4.44	0.39	1.56
B	28.57	24.32	2.74	8.38	4.24	0.51	1.56
C	28.88	24.60	2.51	8.38	4.28	0.44	1.56
D	21.44	17.94	1.23	8.38	3.50	0.32	1.56
E	19.06	15.71	0.51	8.38	3.35	0.22	1.56
F	20.21	16.86	0.42	8.38	3.35	0.16	1.56
G	23.38	19.64	0.50	8.38	3.74	0.16	1.56
H	22.32	18.78	0.37	8.38	3.55	0.12	1.56
I	24.59	20.82	0.76	8.38	3.77	0.13	1.56
Total	46.30	38.64	10.60	8.38	7.70	2.44	1.56

891 **Table 3.** Predicted consequence types of the variants by prevalence level. The most sever consequence of each variant was used. The main
892 Sequence Ontology (SO) terms are shown in order of severity (more severe to less severe) as estimated by Ensembl Variant Effect Predictor.
893 The correlation (r) between the percentage of variants of each consequence type and prevalence is reported. In bold, categories that will be
894 analysed in the next sections.

Consequence type	Percentage of variants (%) by prevalence level										r
	1	2	3	4	5	6	7	8	9	Total	
Loss-of-function[1]	**0.061**	**0.035**	**0.026**	**0.021**	**0.019**	**0.017**	**0.019**	**0.018**	**0.019**	**0.032**	**-.76***
Splice acceptor/donor	0.038	0.023	0.014	0.010	0.009	0.007	0.008	0.008	0.008	0.018	-.79*
Stop-gain	0.014	0.009	0.008	0.008	0.007	0.007	0.007	0.006	0.006	0.009	-.82*
Stop-loss	0.005	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.003	0.003	-.36
Start-loss	0.004	0.002	0.002	0.002	0.001	0.001	0.002	0.002	0.002	0.002	-.47
Frameshift indel	**0.014**	**0.017**	**0.019**	**0.021**	**0.020**	**0.021**	**0.024**	**0.032**	**0.055**	**0.027**	**+.81***
In-frame indel	**0.005**	**0.008**	**0.009**	**0.008**	**0.008**	**0.008**	**0.007**	**0.007**	**0.005**	**0.006**	**-.23**
Missense	0.556	0.378	0.355	0.340	0.344	0.336	0.319	0.306	0.325	0.393	-.73*
Deleterious	**0.201**	**0.092**	**0.074**	**0.069**	**0.064**	**0.062**	**0.054**	**0.048**	**0.040**	**0.096**	**-.78***
Tolerated	**0.223**	**0.170**	**0.165**	**0.165**	**0.173**	**0.167**	**0.161**	**0.159**	**0.177**	**0.183**	**-.52**
Splice region	0.105	0.098	0.088	0.081	0.083	0.081	0.080	0.081	0.085	0.090	-.76*
Synonymous	**0.240**	**0.313**	**0.334**	**0.348**	**0.355**	**0.353**	**0.337**	**0.331**	**0.353**	**0.316**	**+.65**
Untranslated regions	**2.300**	**2.252**	**2.257**	**2.191**	**2.146**	**2.156**	**2.093**	**2.089**	**2.061**	**2.180**	**-.98***
Promoter + 5' UTR	0.879	0.825	0.812	0.812	0.787	0.813	0.759	0.766	0.759	0.810	-.90*
3' UTR	1.421	1.427	1.445	1.378	1.359	1.343	1.334	1.322	1.302	1.370	-.94*
Non-coding transcript exon	0.104	0.113	0.107	0.113	0.128	0.118	0.105	0.109	0.117	0.111	+.25
Intronic	**47.744**	**47.571**	**47.634**	**47.162**	**46.513**	**46.709**	**46.701**	**46.355**	**46.132**	**46.981**	**-.95***
Upstream of gene	3.062	3.066	3.075	3.041	3.083	3.056	2.929	2.943	2.936	3.015	-.81*
Downstream of gene	2.660	2.679	2.740	2.747	2.746	2.705	2.700	2.707	2.676	2.692	+.04
Intergenic	**43.148**	**43.468**	**43.355**	**43.927**	**44.553**	**44.439**	**44.687**	**45.021**	**45.235**	**44.154**	**+.97***

895 [1]If frameshift indels were included in this category: r = -.06 (P>.05)
896 *Significant correlation (P<.05)
897

898 **Table 4.** Frequency of the alternative allele by predicted consequence type and prevalence level. Values are medians.

Consequence type	Frequency of the alternative allele by prevalence level									
	1	2	3	4	5	6	7	8	9	Total
Loss-of-function	.0010	.017	.048	.062	.089	.114	.151	.223	.489	.020
Frameshift indel	.4816	.758	.757	.420	.302	.260	.339	.456	.693	.634
In-frame indel	.8893	.903	.910	.898	.812	.785	.702	.595	.572	.735
Deleterious missense	.0006	.018	.043	.061	.078	.092	.125	.170	.350	.010
Tolerated missense	.0011	.027	.047	.066	.083	.106	.143	.202	.443	.074
Synonymous	.0037	.032	.049	.066	.086	.107	.151	.205	.447	.110
Promoter+UTR	.0019	.034	.059	.078	.099	.122	.166	.226	.475	.102
Intronic	.0015	.035	.059	.080	.102	.126	.171	.235	.485	.110
Intergenic	.0015	.033	.058	.080	.105	.129	.173	.237	.483	.116

899   **Table 5.** Wright's fixation statistic (F_{ST}) by predicted consequence type and prevalence
900   level. Values are medians.

Consequence type	F_{ST} by prevalence level								
	2	**3**	**4**	**5**	**6**	**7**	**8**	**9**	**Total**
Loss-of-function	.003	.022	.047	.066	.094	.114	.145	.171	.071
Frameshift indel	.010	.042	.065	.081	.088	.120	.146	.148	.114
In-frame indel	.011	.035	.051	.070	.087	.105	.115	.130	.077
Deleterious missense	.005	.029	.055	.073	.087	.110	.131	.160	.068
Tolerated missense	.009	.036	.061	.084	.107	.127	.158	.184	.108
Synonymous	.013	.040	.062	.090	.110	.130	.158	.194	.117
Promoter+UTR	.009	.036	.060	.086	.108	.131	.158	.190	.110
Intronic	.009	.037	.063	.089	.111	.136	.164	.195	.118
Intergenic	.009	.036	.066	.091	.112	.139	.167	.193	.121

Additional files

Supplementary Methods

A total of 70,739,387 variants were called across all nine lines. Of these, 24,394,763 variants failed to meet quality control criteria. Of these, 148,825 variants were discarded because they had mean depth values 3 times greater than the average realized coverage, 1,927,221 were multiallelic within line, and 1,673,219 were biallelic within line but multiallelic when all lines were considered. The remaining variants were imputed for all pedigreed individuals, but 20,645,588 of them were fixed for the reference allele in the imputed individuals that passed our accuracy quality control. This affected mostly variants that had been called in only one line and for which the alternative allele segregated at very low frequency. The hypothesis that such variants arise from false positives in variant calling seems unlikely to be the main cause as for more than 99% of these variants we read the alternative allele in at least two individuals. Additionally, we previously quantified that 96.9% of the variants called from low-coverage data were confirmed by sequencing the same individuals at high coverage [1]. A total of 46,344,624 biallelic variants passed quality control criteria across all lines.

1. Ros-Freixedes R, Battagin M, Johnsson M, Gorjanc G, Mileham AJ, Rounsley SD, & Hickey JM. 2018. Impact of index hopping and bias towards the reference allele on accuracy of genotype calls from low-coverage sequencing. Genet Sel Evol, 50: 64.

925
926 **Figure S1.** Population structure of the sequenced pigs according to the two first principal
927 components. The colour clusters correspond to lines A to I.
928

929
930 **Figure S2.** Variant density for the private variants in each line.
931

932 **Table S1.** Number of analysed variants by chromosome.

Chromosome	Length (Mb)	SNPs (M)	Indels (M)	Variant density (thousands/Mb)
1	274.3	3.77	0.76	16.5
2	151.9	2.60	0.52	20.5
3	132.8	2.35	0.44	21.0
4	130.9	2.21	0.43	20.2
5	104.5	1.95	0.39	22.4
6	170.8	2.80	0.55	19.6
7	121.8	2.20	0.43	21.6
8	139.0	2.37	0.50	20.6
9	139.5	2.47	0.48	21.1
10	69.4	1.60	0.31	27.5
11	79.2	1.57	0.31	23.7
12	61.6	1.35	0.25	26.0
13	208.3	2.97	0.64	17.3
14	141.8	2.38	0.48	20.2
15	140.4	2.20	0.46	18.9
16	79.9	1.50	0.30	22.5
17	63.5	1.32	0.25	24.7
18	56.0	1.04	0.19	22.0
Total	2,501.9	38.64	7.70	18.5

933