Decoding rRNA sequences for improved metagenomics of sylvatic mosquito species

Cassandra Koh[1], Lionel Frangeul[1], Hervé Blanc[1], Carine Ngoagouni[2], Sébastien Boyer[3], Philippe Dussart[4], Nina Grau[5], Romain Girod[5], Jean-Bernard Duchemin[6] and Maria-Carla Saleh[1*]

[1] Viruses and RNA Interference Unit, Institut Pasteur, Université Paris Cité, CNRS UMR3569, Paris F-75015, France

[2] Medical Entomology Laboratory, Institut Pasteur de Bangui, Bangui PO Box 923, Central African Republic

[3] Medical and Veterinary Entomology Unit, Institut Pasteur du Cambodge, Phnom Penh 12201, Cambodia

[4] Virology Unit, Institut Pasteur du Cambodge, Phnom Penh 12201, Cambodia

[5] Medical Entomology Unit, Institut Pasteur de Madagascar, Antananarivo 101, Madagascar

[6] Vectopôle Amazonien Emile Abonnenc, Institut Pasteur de la Guyane, Cayenne 97306, French Guiana

* To whom correspondence should be addressed.

Present address: Philippe Dussart, Institut Pasteur de Madagascar, Antananarivo 101, Madagascar; Nina Grau, Sciences Economiques et Sociales de la Santé et Traitement de l'Information Médicale, Faculté de Médecine, Marseille 13005, France

Corresponding author: Maria-Carla Saleh, carla.saleh@pasteur.fr, +33 1 45 68 85 47.

ABSTRACT

As mosquito-borne virus epidemics are often preceded by undetected spillover events, surveillance

and virus discovery studies in non-urban mosquitoes informs pre-emptive and responsive public

health measures. RNA-seq metagenomics is a popular methodology but it is constrained by

overabundant rRNA. The lack of reference sequences for most mosquito species is a major

impediment against physical and computational removal of rRNA reads.

We describe a strategy to assemble novel rRNA sequences from mosquito specimens, producing

an unprecedented dataset of 234 full-length 28S and 18S rRNA sequences of 33 medically important

species from countries with known histories of mosquito-borne virus circulation (Cambodia, the

Central African Republic, Madagascar, and French Guiana). We also evaluate the utility of rRNA

sequences as molecular barcodes relative to the mitochondrial *cytochrome* c *oxidase I* (COI) gene.

We show that rRNA sequences can be used for species identification when COI sequences are

ambiguous or unavailable, revealing evolutionary relationships concordant with contemporary

mosquito systematics.

This expansion of the rRNA reference library improves mosquito RNA-seq metagenomics by

permitting the optimization of species-specific rRNA depletion protocols for a broader species range

and streamlined species identification by rRNA barcoding. In addition, rRNA barcodes could serve as

an additional tool for mosquito taxonomy and phylogeny.

Keywords: surveillance, RNA-seq, ribosomal RNA, barcode, metagenomics, mosquito

INTRODUCTION

Mosquitoes top the list of vectors for arthropod-borne diseases, being implicated in the transmission of many human pathogens responsible for arboviral diseases, malaria, and lymphatic filariasis (WHO, 2017). Mosquito-borne viruses circulate in sylvatic or urban transmission cycles driven by different mosquito species with their own distinct host preferences. Although urban mosquito species are chiefly responsible for amplifying epidemics in dense human populations, non-urban (sylvatic and peri-urban) mosquitoes maintain the transmission of these viruses among forest-dwelling animal reservoir hosts and are implicated in spillover events when humans enter their ecological niches (Valentine et al., 2019). Given that mosquito-borne virus emergence is preceded by such spillover events, continuous surveillance and virus discovery in non-urban mosquitoes is integral to designing effective public health measures to pre-empt or respond to mosquito-borne viral epidemics.

Metagenomics on field specimens is the most powerful method in the toolkit for understanding mosquito-borne disease ecology through the One Health lens (Webster et al., 2016). With next-generation sequencing becoming more accessible, such studies have provided unprecedented insights into the interfaces among mosquitoes, their environment, and their animal and human hosts. However, working with lesser studied mosquito species poses several problems.

First, metagenomics studies based on RNA-seq are bedevilled by overabundant ribosomal RNAs (rRNAs). These non-coding RNA molecules comprise at least 80% of the total cellular RNA population (Gale & Crampton, 1989). Due to their length and their abundance, they are a sink for precious next generation sequencing reads, decreasing the sensitivity of pathogen detection unless depleted during library preparation. Yet the most common rRNA depletion protocols require prior knowledge of rRNA sequences of the species of interest as they involve hybridizing antisense oligos to the rRNA molecules prior to removal by ribonucleases (Fauver et al., 2019; Phelps et al., 2021) or by bead capture (Kukutla et al., 2013). Presently, reference sequences for rRNAs are limited to only a handful of species from three genera: *Aedes*, *Culex*, and *Anopheles* (Ruzzante et al., 2019). The lack of reliable rRNA depletion methods could deter mosquito metagenomics studies from expanding their sampling diversity, resulting in a gap in our knowledge of mosquito vector ecology. The inclusion of lesser studied yet medically relevant non-urban species is therefore imperative.

Second, species identification based on morphology is notoriously complicated between members of species subgroups. This is especially the case among *Culex* subgroups. Sister species are often

sympatric and show at least some competence for a number of viruses, such as Japanese encephalitis virus, St Louis encephalitic virus, and Usutu virus (Nchoutpouen et al., 2019). Although they share many morphological traits, each of these species have distinct ecologies and host preferences, thus the challenge of correctly identifying vector species can affect epidemiological risk estimation for these diseases (Farajollahi et al., 2011). DNA molecular markers are often employed to a limited degree of success to distinguish between sister species (Batovska et al., 2017; Zittra et al., 2016).

To address the lack of full-length rRNA sequences in public databases, we sought to determine the 28S and 18S rRNA sequences of a diverse set of Old and New World non-urban (sylvatic and peri-urban) mosquito species from four countries representing three continents: Cambodia, the Central African Republic, Madagascar, and French Guiana. These countries, due to their proximity to the equator, contain high mosquito biodiversity (Foley et al., 2007) and have had long histories of mosquito-borne virus circulation. Increased and continued surveillance of local mosquito species could lead to valuable insights on mosquito virus biogeography. Using a unique score-based read filtration strategy to remove interfering non-mosquito rRNA reads for accurate *de novo* assembly, we produced a dataset of 234 novel full-length 28S and 18S rRNA sequences from 33 mosquito species, 30 of which have never been recorded before.

We also explored the functionality of 28S and 18S rRNA sequences as molecular barcodes by comparing their performance to that of the mitochondrial *cytochrome* c *oxidase subunit I* (COI) gene for molecular taxonomic and phylogenetic investigations. The COI gene is the most widely used DNA marker for molecular species identification and forms the basis of the Barcode of Life Data System (BOLD) (Hebert et al., 2003; Ratnasingham & Hebert, 2007). Presently, full-length rRNA sequences are much less represented compared to other molecular markers. We hope that our sequence dataset, with its species diversity and eco-geographical breadth, and the assembly strategy we describe would further facilitate the use of rRNA as barcodes. In addition, this dataset enables the design of species-specific oligos for cost-effective rRNA depletion for a broader range of mosquito species and streamlined molecular species diagnostics during RNA-seq.

RESULTS

Poor rRNA depletion using a non-specific depletion method

During library preparations of mosquito samples for RNA-seq, routinely used methods for depleting

rRNA are commercial kits optimised for human or mice samples (Belda et al., 2019; Bishop-Lilly et al.,

2010; Chandler et al., 2015; N. Kumar et al., 2012; Weedall et al., 2015; Zakrzewski et al., 2018) or

through 80–100 base pair antisense probe hybridisation followed by ribonuclease digestion (Fauver et

al., 2019; Phelps et al., 2021). In cases where the complete reference rRNA sequence of the target

species is not known, oligos would be designed based on the rRNA sequence of the closest related

species (25, this study). These methods should deplete the conserved regions of rRNA sequences.

However, the variable regions remain at abundances high enough to compromise RNA-seq output. In

our hands, we have found that using probes designed for the *Ae. aegypti* rRNA sequence followed by

RNase H digestion according to the protocol published by Morlan *et al.* (2012) produced poor

depletion in *Ae. albopictus*, and in Culicine and Anopheline species (Figure 1A). Additionally, the lack

of reference rRNA sequences compromises the *in silico* clean-up of remaining rRNA reads from

sequencing data, as reads belonging to variable regions would not be removed. To solve this and to

enable RNA-seq metagenomics on a broader range of mosquito species, we performed RNA-seq to

generate reference rRNA sequences for 33 mosquito species representing 10 genera from

Cambodia, the Central African Republic, Madagascar, and French Guiana. Most of these species are

associated with vector activity for various pathogens in their respective ecologies (Table 1).

Figure 1. (**A**) Proportion of rRNA reads found in mosquito specimen pools of 5 individuals depleted by

probe hybridisation followed by RNase H digestion. Probes were antisense to *Ae. aegypti* rRNA

sequences. (**B**) Read vs. score ratio plot of a representative specimen "Ae simpsoni CS S27". Green

line indicates 0.8 cut-off where only reads above this threshold are used in rRNA assembly.

Table 1. List of mosquito species represented in this study and their vector status.

Mosquito taxonomy*	Origin**	Collection site (ecosystem type)	Vector for***	Reference
Aedes (Fredardsius) vittatus	CF	rural (village)	DENV, ZIKV, CHIKV, YFV	(Diallo et al., 2020)
Aedes (Ochlerotatus) scapularis	GF	rural (village)	YFV	(Vasconcelos et al., 2001)
Aedes (Ochlerotatus) serratus	GF	rural (village)	YFV, OROV	(Cardoso et al., 2010; Gaillet et al., 2021)
Aedes (Stegomyia) aegypti	CF	urban	DENV, ZIKV, CHIKV, YFV	(Kraemer et al., 2019)
Aedes (Stegomyia) albopictus	CF, KH	rural (village, nature reserve)	DENV, ZIKV, CHIKV, YFV, JEV	(Auerswald et al., 2021; Kraemer et al., 2019)
Aedes (Stegomyia) simpsoni	CF	rural (village)	YFV	(Mukwaya et al., 2000)
Anopheles (Anopheles) baezai	KH	rural (nature reserve)	unreported	–
Anopheles (Anopheles) coustani	MG, CF	rural (village)	RVFV, malaria	(Mwangangi et al., 2013; Nepomichene et al., 2018; Ratovonjato et al., 2011)
Anopheles (Cellia) funestus	MG, CF	rural (village)	ONNV, malaria	(Lutomiah et al., 2013)
Anopheles (Cellia) gambiae	MG, CF	rural (village)	ONNV, malaria	(Brault et al., 2004)
Anopheles (Cellia) squamosus	MG	rural (village)	RVFV, malaria	(Ratovonjato et al., 2011; Stevenson et al., 2016)
Coquillettidia (Rhynchotaenia) venezuelensis	GF	rural (village)	OROV	(Gaillet et al., 2021)
Culex (Culex) antennatus	MG	rural (village)	RVFV	(Nepomichene et al., 2018; Ratovonjato et al., 2011)
Culex (Culex) duttoni	CF	rural (village)	unreported	–
Culex (Culex) neavei	MG	rural (village)	USUV	(Nikolay et al., 2011)
Culex (Culex) orientalis	KH	rural (nature reserve)	JEV	(Kim et al., 2015)
Culex (Culex) perexiguus	MG	rural (village)	WNV	(Vázquez González et al., 2011)
Culex (Culex) pseudovishnui	KH	rural (nature reserve)	JEV	(Auerswald et al., 2021; Maquart et al., 2021)
Culex (Culex) quinquefasciatus	MG, CF, KH	rural (village, nature reserve)	ZIKV, JEV, WNV, DENV, SLEV, RVFV, *Wuchereria bancrofti*	(Bhattacharya et al., 2016; Maquart et al., 2021; Ndiaye et al., 2016)
Culex (Culex) tritaeniorhynchus	MG, KH	rural (village, nature reserve)	JEV, WNV, RVFV	(Auerswald et al., 2021; Maquart et al., 2021)
Culex (Melanoconion) spissipes	GF	rural (village)	VEEV	(Weaver et al., 2004)
Culex (Melanoconion) portesi	GF	rural (village)	VEEV, TONV	(Talaga et al., 2021; Weaver et al., 2004)
Culex (Melanoconion) pedroi	GF	rural (village)	EEEV, VEEV, MADV	(Talaga et al., 2021; M. J. Turell et al., 2008)
Culex (Oculeomyia) bitaeniorhynchus	MG, KH	rural (village, nature reserve)	JEV	(Auerswald et al., 2021; Maquart et al., 2021)
Culex (Oculeomyia) poicilipes	MG	rural (village)	RVFV	(Ndiaye et al., 2016)
Eretmapodites intermedius	CF	rural (village)	unreported	–
Limatus durhamii	GF	rural (village)	ZIKV	(Barrio-Nuevo et al., 2020)

Mansonia (Mansonia) titillans	GF	rural (village)	VEEV, SLEV	(Hoyos-López et al., 2015; Michael J. Turell, 1999)
Mansonia (Mansonioides) indiana	KH	rural (nature reserve)	JEV	(Arunachalam et al., 2004)
Mansonia (Mansonioides) uniformis	MG, CF, KH	rural (village, nature reserve)	WNV, RVFV, *Wuchereria bancrofti*	(Lutomiah et al., 2013; Maquart et al., 2021; Ughasi et al., 2012)
Mimomyia (Etorleptiomyia) mediolineata	MG	rural (village)	unreported	–
Psorophora (Janthinosoma) ferox	GF	rural (village)	ROCV	(Mitchell et al., 1986)
Uranotaenia (Uranotaenia) geometrica	GF	rural (village)	unreported	–

*   () indicates subgenus

**   Origin countries are listed as their ISO alpha-2 codes: Central African Republic, CF; Cambodia, KH;

Madagascar, MG; French Guiana, GF.

**   dengue virus, DENV; Zika virus, ZIKV; chikungunya virus, CHIKV; Yellow Fever virus, YFV; Oropouche virus,

OROV; Japanese encephalitis virus, JEV; Rift Valley Fever virus, RVFV; O'Nyong Nyong virus, ONNV; Usutu

virus, USUV; West Nile virus, WNV; Saint Louis encephalitis virus, SLEV; Venezuelan equine encephalitis

virus, VEEV; Tonate virus, TONV; Eastern equine encephalitis virus, EEEV; Madariaga virus, MADV; Rocio

virus, ROCV.

rRNA reads filtering and sequence assembly

Assembling Illumina reads to reconstruct rRNA sequences from total mosquito RNA is not a

straightforward task. Apart from host rRNA, total RNA samples also contain rRNA from other

organisms associated with the host (microbiota, external parasites, or ingested diet). As rRNA

sequences share high homology in conserved regions, Illumina reads (150 bp) from non-host rRNA

can interfere with the contig assembly of host 28S and 18S rRNA.

Our score-based filtration strategy, described in detail in Methods, allowed us to bioinformatically

remove interfering rRNA reads and achieve successful *de novo* assembly of 28S and 18S rRNA

sequences for all our specimens. Briefly, for each Illumina read, we computed a ratio of BLAST

scores against an Insecta library over a Non-Insecta library. Reads were segregated into four

categories: (i) reads mapping only to the Insecta library, (ii) reads mapping better to the Insecta

relative to Non-Insecta library, (iii) reads mapping better to the Non-Insecta relative to the Insecta

library, and finally (iv) reads mapping only to the Non-Insecta library (Figure 1B, Supplementary

Figure S1). By applying a conservative threshold of 0.8 to account for the non-exhaustiveness of the

SILVA database, we removed reads that likely do not originate from mosquito rRNA. Notably, 15 of

our specimens were engorged with vertebrate blood, a rich source of non-mosquito rRNA

(Supplementary Table S1). The successful assembly of complete 28S and 18S rRNA sequences

demonstrates that this strategy performs as expected even with high amounts of non-host rRNA

reads. This is particularly important in studies on field-captured mosquitoes as females are often

sampled already having imbibed a blood meal or captured using the human landing catch technique.

We encountered challenges for three specimens morphologically identified as *Ma. africana*

(Specimen ID 33-35). COI amplification by PCR did not produce any product, hence COI barcoding

could not be used to confirm species identity. In addition, SPAdes was only able to assemble partial

length contigs, despite the high number of reads with high scores against the Insecta library. Among

other *Mansonia* specimens, the partial length contigs shared the highest similarity with contigs

obtained from sample "Ma uniformis CF S51". We then performed a guided assembly using the 28S

and 18S sequences of this specimen as references, which successfully produced full-length contigs.

In two of these specimens (Specimen ID 34 and 35), our assembly initially produced two sets of 28S

and 18S rRNA sequences, one of which was similar to mosquito rRNA with low coverage and another

with ten-fold higher coverage and 95% nucleotide sequence similarity to a water mite of genus

Horreolanus known to parasitize mosquitoes. Our success in obtaining rRNA sequences for mosquito

and water mite shows that our strategy can be applied to metabarcoding studies where the input

material comprises multiple insect species, provided that appropriate reference sequences of the

target species or of a close relative are available.

Altogether, we were able to assemble 122 28S and 114 18S full-length rRNA sequences for 33

mosquito species representing 10 genera sampled from four countries across three continents. This

dataset contains, to our knowledge, the first records for 30 mosquito species and for seven genera:

Coquillettidia, *Mansonia*, *Limatus*, *Mimomyia*, *Uranotaenia*, *Psorophora,* and *Eretmapodites*.

Individual GenBank accession numbers for these sequences and specimen information are listed in

Supplementary Table S1.

Comparative phylogeny of novel rRNA sequences relative to existing records

To verify the assembly accuracy of our rRNA sequences, we constructed a comprehensive

phylogenetic tree from the 28S rRNA sequences generated from our study alongside those publicly

available from GenBank (Figure 2). We applied a search criterion for GenBank sequences with at

least 95% coverage of our sequence lengths (~4000 bp), aiming to represent as many species or

genera as possible. Although we rarely found records for the same species included in our study, the

resulting tree showed that our 28S sequences generally clustered according to their respective

species and subgenera, supported by moderate to good bootstrap support at terminal nodes. Species

taxa generally formed monophyletic clades, with the exception of *An. gambiae* and *Cx.*

quinquefasciatus. *An. gambiae* 28S rRNA sequences formed a clade with closely related sequences

from *An. arabiensis*, *An. merus*, and *An. coluzzii,* suggesting unusually high interspecies homology for

Anophelines or other members of subgenus *Cellia.* Meanwhile, *Cx. quinquefasciatus* 28S rRNA

sequences formed a taxon paraphyletic to sister species *Cx. pipiens*.

0.05

Figure 2. Phylogenetic tree based on 28S sequences generated from this study and from GenBank

(3900 bp) as inferred using maximum-likelihood method and constructed to scale in MEGA X (S.

Kumar et al., 2018). Values at each node indicate bootstrap support (%) from 500 replications. For

sequences from this study, each specimen label contains information on taxonomy, origin (in 2-letter

country codes), and specimen ID. Labels in bold indicate GenBank sequences with accession

numbers shown. Label colours indicate genera: *Culex* in coral, *Anopheles* in purple, *Aedes* in dark

blue, *Mansonia* in dark green, *Culiseta* in maroon, *Limatus* in light green, *Coquillettidia* in light blue,

Psorophora in yellow, *Mimomyia* in teal, *Uranotaenia* in pink and *Eretmapodites* in brown. Scale bar

at 0.05 is shown.

28S sequence-based phylogenetic reconstructions (Figure 2, with GenBank sequences;

Supplementary Figure S3, this study only) showed marked incongruence to that of 18S sequences

(Figure 3). Although all rRNA trees show the bifurcation of *Culicidae* into subfamilies *Anophelinae* and

Culicinae, the recovered intergeneric phylogenetic relationships vary between the 28S and 18S trees

and are weakly supported. The 18S tree also exhibited several taxonomic anomalies: (i) the lack of

definitive clustering by species within the *Culex* subgenus (ii) the lack of distinction between 18S

sequences of *Cx. pseudovishnui* and *Cx. tritaeniorhynchus*; (iii) the placement of Ma sp. 3 CF S35

within a *Culex* clade; and (iv) the lack of a monophyletic *Mimomyia* clade. The topology of the 18S

tree suggests higher sequence divergence between the two *Cx. quinquefasciatus* taxa and between

the two *Mimomyia* taxa than in the 28S trees. However, 28S and 18S rRNA sequences are encoded

by linked loci in rDNA clusters and should not be analysed separately.

Figure 3. Phylogenetic tree based on 18S sequences (1900 bp) as inferred using maximum-likelihood method and constructed to scale in MEGA X (S. Kumar et al., 2018). Values at each node indicate bootstrap support (%) from 500 replications. Each specimen label contains information on taxonomy, origin (in 2-letter country codes), and specimen ID. Label colours indicate genera: *Culex* in coral, *Anopheles* in purple, *Aedes* in dark blue, *Mansonia* in dark green, *Limatus* in light green, *Coquillettidia* in light blue, *Psorophora* in yellow, *Mimomyia* in teal, *Uranotaenia* in pink and *Eretmapodites* in brown. Scale bar at 0.05 is shown.

Indeed, when concatenated 28S+18S rRNA sequences were generated from the same specimens

(Figure 4), the phylogenetic tree resulting from these sequences more closely resembles the 28S tree

(Figure 2) with regard to the basal position of the *Mimomyia* clade within the *Culicinae* subfamily with

good bootstrap support in either tree (84% in 28S tree, 100% in concatenated 28S+18S tree). For

internal nodes, bootstrap support values were higher in the concatenated tree compared to the 28S

tree. Interestingly, the 28S+18S tree formed an *Aedini* tribe-clade encompassing taxa from genera

Psorophora, Aedes, and *Eretmapodites*, possibly driven by the inclusion of 18S sequences.

Concatenation also resolved the anomalies found in the 18S tree and added clarity to the close

relationship between *Culex* and *Mansonia* taxa. Of note, relative to the 28S tree (Figure 2) the *Culex*

and *Mansonia* genera are no longer monophyletic in the concatenated 28S+18S tree (Figure 4).

Genus *Culex* is paraphyletic with respect to subgenus *Mansonoides* of genus *Mansonia* (Figure 2).

Ma. titillans and Ma sp. 4, which we suspect to be *Ma. pseudotitillans,* always formed a distinct branch

in 28S or 18S phylogenies, thus possibly representing a clade of subgenus *Mansonia*.

The concatenated 28S+18S tree recapitulates what is classically known about the systematics of

our specimens, namely (i) the early divergence of subfamily *Anophelinae* from subfamily *Culicinae*, (ii)

the division of genus *Anopheles* into two subgenera, *Anopheles* and *Cellia,* (iii) the division of genus

Aedes into subgenera *Stegomyia* and *Ochlerotatus*, (iv) the divergence of monophyletic subgenus

Melanoconion within the *Culex* genus (Harbach, 2007; Harbach & Kitching, 2016).

Figure 4. Phylogenetic tree based on concatenated 28S and 18S sequences generated from this

study (3900+1900 bp) as inferred using maximum-likelihood method and constructed to scale in

MEGA X (S. Kumar et al., 2018). Values at each node indicate bootstrap support (%) from 500

replications. For sequences from this study, each specimen label contains information on taxonomy,

origin (as indicated in 2-letter country codes), and specimen ID. Label colours indicate genera: *Culex*

in coral, *Anopheles* in purple, *Aedes* in dark blue, *Mansonia* in dark green, *Culiseta* in maroon,

Limatus in light green, *Coquillettidia* in light blue, *Psorophora* in yellow, *Mimomyia* in teal, *Uranotaenia*

in pink and *Eretmapodites* in brown. Scale bar at 0.05 is shown.

rRNA as a molecular marker for taxonomy and phylogeny

We sequenced a 621 bp region of the COI gene to confirm morphological species identification of our

specimens and to compare the functionality of rRNA and COI sequences as molecular markers for

taxonomic and phylogenetic investigations. COI sequences were able to unequivocally determine the

species identity in most specimens except for the following cases. *An. coustani* COI sequences from

our study regardless of specimen origin shared remarkably high nucleotide similarity (>98%) with

several other *Anopheles* species such as *An. rhodesiensis*, *An. rufipes*, *An. ziemanni*, *An. tenebrosus*,

although *An. coustani* remained the most frequent and closest match. In the case of *Ae. simpsoni*,

three specimens had been morphologically identified as *Ae. opok* although their COI sequences

showed 97–100% similarity to that of *Ae. simpsoni.* As GenBank held no records of *Ae. opok* COI at

the time of this study, we instead aligned the putative *Ae. simpsoni* COI sequences against two sister

species of *Ae. opok*: *Ae. luteocephalus* and *Ae. africanus.* We found they shared only 90% and 89%

similarity, respectively. Given this significant divergence, we concluded these specimens to be *Ae.*

simpsoni. Ambiguous results were especially frequent among *Culex* specimens belonging to the *Cx.*

pipiens or *Cx. vishnui* subgroups, where the query sequence differed with either of the top two hits by

a single nucleotide. For example, between *Cx. quinquefasciatus* and *Cx. pipiens* of the *Cx. pipiens*

subgroup, and between *Cx. vishnui* and *Cx. tritaeniorhynchus* of the *Cx. vishnui* subgroup.

Among our three specimens of *Ma. titillans*, two appeared to belong to a single species that is

different but closely related to *Ma. titillans*. We surmised that these specimens could instead be *Ma.*

pseudotitillans based on morphological similarity but were not able to verify this by molecular means

as no COI reference sequence is available for this species. These specimens are hence putatively

labelled as "Ma sp.4".

Phylogenetic reconstruction based on the COI sequences showed clustering of all species taxa

into distinct clades, underlining the utility of the COI gene in molecular taxonomy (Figure 5)(Hebert et

al., 2003; Ratnasingham & Hebert, 2007). However, species delineation among members of *Culex*

subgroups were not as clear cut, although sister species were correctly placed as sister taxa. This is

comparable to the 28S+18S tree (Figure 4) and is indicative of lower intraspecies distances relative to

interspecies distances.

To evaluate the utility of 28S and 18S rRNA sequences for molecular taxonomy, we used the

28S+18S rRNA tree to discern the identity of six specimens for which COI barcoding could not be

performed. These specimens include three unknown *Mansonia* species (Specimen ID 33–35), a *Ma.*

uniformis (Specimen ID 51), an *An. gambiae* (Specimen ID 47), and a *Ur. geometrica* (Specimen ID

113). Their positions in the 28S+18S rRNA tree relative to adjacent taxa confirms the morphological

identification of all six specimens to the genus level and, for three of them, to the species level (Figure

4).

The phylogenetic relationships indicated by the COI tree compared to the 28S+18S tree present

only a few points of congruence. COI-based phylogenetic inference indeed showed clustering of

generic taxa into monophyletic clades albeit with very weak bootstrap support, except for genera

Culex and *Mansonia* (Figure 5). Contrary to the 28S+18S tree (Figure 4), *Culex* subgenus

Melanoconion was depicted as a polyphyletic taxon with *Cx. spissipes* being a part of the greater

Culicini clade with members from subgenera *Oculeomyia* and *Culex* while *Cx. pedroi* and *Cx. portesi*

formed a distantly related clade. Among the *Mansonia* specimens, the two unknown Ma sp.4

specimens were not positioned as the nearest neighbours of *Ma. titillans* and instead appeared to

have diverged earlier from most of the other taxa from the *Culicidae* family. Notably, the COI

sequences of genus *Anopheles* is not basal to the other members of *Culicidae* and is instead shown

to be sister to Culex COI sequences (8% bootstrap support). This is a direct contrast to what is

suggested by the rRNA phylogenies (Figures 2–4), which suggests *Culex* rRNA sequences to be

among the most recently diverged. Bootstrap support for the more internal nodes of the COI trees

were remarkably low compared to those of rRNA-based trees.

In all rRNA trees, it is clear that the interspecific and intersubgeneric evolutionary distances within

the genus *Anopheles* are high relative to any other genera, indicating a greater degree of divergence.

This is evidenced by the longer branch lengths connecting Anopheline species-clades to the node of

the most recent common ancestor for subgenera *Anopheles* and *Cellia* (Figures 2-4, Supplementary

Figure S3). This feature is not evident in the COI tree, where the Anopheline interspecies distances

are comparable to those within the *Culex*, *Aedes*, and *Mansonia* taxa.

Figure 5. Phylogenetic tree based on COI sequences (621–699 bp) as inferred using maximum-likelihood method and constructed to scale in MEGA X (S. Kumar et al., 2018). Values at each node indicate bootstrap support (%) from 500 replications. Each specimen label contains information on taxonomy, origin (as indicated in 2-letter country codes), and specimen ID. Label colours indicate genera: *Culex* in coral, *Anopheles* in purple, *Aedes* in dark blue, *Mansonia* in dark green, *Limatus* in

light green, *Coquillettidia* in light blue, *Psorophora* in yellow, *Mimomyia* in teal, *Uranotaenia* in pink

and *Eretmapodites* in brown. Scale bar at 0.05 is shown.

On *Culex* subgroups

Culex (subgenus *Culex*) specimens of this study comprise several closely related sister species

belonging to the *Cx. vishnui* and *Cx. univittatus* subgroups, which are notoriously difficult to

differentiate based on morphology. Accordingly, in the 28S+18S rRNA (Figure 4) and COI (Figure 5)

trees these species and their known sister species were clustered together within the *Culex*

(subgenus *Culex*) clade: *Cx. tritaeniorhynchus* with *Cx. pseudovishnui* (*Cx. vishnui* subgroup); *Cx.*

perexiguus with *Cx. neavei* (*Cx. univittatus* subgroup).

The use of COI barcoding to distinguish between members of the *Culex* subgroups was limited.

For example, for the two *Cx. quinquefasciatus* samples in our taxonomic assemblage (Specimen ID

74 and 75), BLAST analyses of their COI sequences revealed they are a single nucleotide away from

Cx. pipiens or *Cx. quinquefasciatus* COI sequences (Supplementary Table S2). In the 28S rRNA tree

with GenBank sequences (Figure 2), two GenBank sequences of *Cx. pipiens* sequences formed a

clade sister to another containing three *Cx. quinquefasciatus* GenBank sequences and the "Cx

quinquefasciatus MG S74" sequence with 78% bootstrap support. This is in accordance with other

studies examining mitochondrial sequences (Sun et al., 2019) and morphological attributes (Harbach

et al., 2017). This shows that the 28S rRNA sequence can distinguish the two species and confirms

that "Cx quinquefasciatus MG S74" is indeed a *Cx. quinquefasciatus* specimen. However, "Cx

quinquefasciatus MG S75" is shown to be basal from other sequences within this *Cx. pipiens*

subgroup-clade with 100% bootstrap support. Given that *Cx. quinquefasciatus* and *Cx. pipiens* are

known to interbreed, it is plausible that this individual is a hybrid of the two species (Farajollahi et al.,

2011).

DISCUSSION

RNA-seq metagenomics on field-captured non-urban mosquitoes is a valuable tool for pre-empting

mosquito-borne virus emergences through surveillance and virus discovery. However, the lack of

reference rRNA sequences hinders good oligo-based depletion and efficient clean-up of RNA-seq

data. Additionally, *de novo* assembly of rRNA sequences is complicated due to regions that are highly

conserved across all distantly related organisms that could be present in a single specimen, i.e.,

microbiota, parasites, or vertebrate blood meal. Hence, we sought to establish a method to

bioinformatically filter out non-host rRNA reads for the accurate assembly of novel 28S and 18S rRNA

reference sequences.

We found that phylogenetic reconstructions based on 28S sequences or concatenated 28S+18S

rRNA sequences were able to correctly cluster mosquito taxa according to species and corroborate

current mosquito classification. This demonstrates that our bioinformatics methodology reliably

generates bona fide 28S and 18S rRNA sequences, even in specimens parasitized by water mites or

engorged with vertebrate blood. Further, we were able to use 28S+18S rRNA taxonomy for molecular

species identification when COI sequences were unavailable or ambiguous, thus supporting the use

of rRNA sequences as a barcode. They have the advantage of circumventing the need to additionally

isolate and sequence DNA from specimens, as RNA-seq reads can be directly mapped against

reference sequences. Post-depletion, in our hands there are sufficient numbers of remaining reads

(5–10% of reads per sample) for assembly of complete rRNA contigs (unpublished data).

Phylogenetic inferences based on 28S or 18S rRNA sequences alone do not recover the same

interspecific relationships. Relative to 28S sequences, we observed more instances where multiple

specimens have near-identical 18S rRNA sequences. This can occur for specimens belonging to the

same species, but also for conspecifics sampled from different geographic locations, such as *An.*

coustani, *An. gambiae*, or *Ae. albopictus.* More rarely, specimens from the same species subgroup,

such as *Cx. pseudovishnui* and *Cx. tritaeniorhynchus*, also shared 18S rRNA sequences. This was

surprising given that the 18S rRNA sequences in our dataset is 1900 bp long. Concatenation of 28S

and 18S rRNA sequences resolved this issue, enabling species delineation even among sister

species of *Culex* subgroups, with which morphological identification meets its limits.

In Cambodia and other parts of Asia, the *Cx. vishnui* subgroup includes *Cx*. *tritaeniorhynchus*, *Cx*.

vishnui, and *Cx*. *pseudovishnui*, which are important vectors of JEV (Maquart & Boyer, 2022). The

former two were morphologically identified in our study but later revealed by COI barcoding to be a

sister species. Discerning sister species of the *Cx. pipiens* subgroup is further complicated by

interspecific breeding, with some populations showing genetic introgression to varying extents (Cornel

et al., 2003). The seven sister species of this subgroup are practically indistinguishable based on

morphology and require molecular methods to discern (Farajollahi et al., 2011; Zittra et al., 2016).

Indeed, the 621 bp COI sequence amplified in our study did not contain enough nucleotide

divergence to allow clear identification, given that the COI sequence of *Cx. quinquefasciatus*

specimens differed from that of *Cx. pipiens* by a single nucleotide. Batovska et al. (2017) found that

even the Internal Transcribed Spacer 2 (ITS2) rDNA region, another common molecular marker,

could not differentiate the two species. Other DNA molecular markers such as nuclear Ace-2 or CQ11

genes (Aspen & Savage, 2003; Zittra et al., 2016) or *Wolbachia pipientis* infection status (Cornel et

al., 2003) are typically employed in tandem. In our study, 28S rRNA sequence-based phylogeny

(Figure 2) validated the identity of specimen "Cx quinquefasciatus MG S74" and suggested that

specimen "Cx quinquefasciatus MG S75" might have been a *pipiens-quinquefasciatus* hybrid. These

examples demonstrate how 28S rRNA sequences, concatenated with 18S or alone, contain enough

resolution to differentiate between *Cx. pipiens* and *Cx. quinquefasciatus*. rRNA-based phylogeny thus

allows for more accurate species identification and ecological observations in the context of disease

transmission. Additionally, tracing the genetic flow across hybrid populations within the *Cx. pipiens*

subgroup can inform estimates of vectorial capacity for each species. As only one or two members

from the *Cx. pipiens* and *Cx. vishnui* subgroups were represented in our taxonomic assemblage, an

explicit investigation including all member species of these subgroups in greater sample numbers is

warranted to further test the degree of accuracy with which 28S and 18S rRNA sequences can

delineate sister species.

Our study included French Guianese *Culex* species *Cx. spissipes* (group Spissipes*)*, *Cx. pedroi*

(group Pedroi), and *Cx. portesi* (group Vomerifer). These species belong to the New World subgenus

Melanoconion, section Spissipes, with well-documented distribution in North and South Americas

(Sirivanakarn, 1982) and are vectors of encephalitic alphaviruses EEEV and VEEV among others

(Talaga et al., 2021; M. J. Turell et al., 2008; Weaver et al., 2004). Indeed, our rooted rRNA and COI

trees showed the divergence of the three *Melanoconion* species from the major *Culex* clade

comprising species broadly found across Africa and Asia (Auerswald et al., 2021; Farajollahi et al.,

2011; Nchoutpouen et al., 2019; Takhampunya et al., 2011). The topology of the concatenated

28S+18S tree places the *Cx. portesi* and *Cx. pedroi* species-clades as sister groups (92% bootstrap

support), with *Cx. spissipes* as a basal group within the *Melanoconion* clade (100% bootstrap

support). This corroborates the systematics elucidated by Navarro and Weaver (2004) using the ITS2

marker, and those by Sirivanakarn (1982) and Sallum and Forattini (1996) based on morphology.

Curiously, in the COI tree, *Cx. spissipes* sequences were clustered with unknown species Cx. sp1,

forming a clade sister to another containing other *Culex* (*Culex*) and *Culex* (*Oculeomyia*) species,

albeit with very low bootstrap support. Previous phylogenetic studies based on the COI gene have

consistently placed *Cx. spissipes* or the Spissipes group basal to other groups within the

Melanoconion subgenus (Torres-Gutierrez et al., 2016, 2018). However, these studies contain only

Culex (*Melanoconion*) species in their assemblage, apart from *Cx. quinquefasciatus* to act as an

outgroup. This clustering of *Cx. spissipes* with non-*Melanoconion* species in our COI phylogeny could

be an artefact of a much more diversified assemblage rather than a true phylogenetic link.

395   Taking advantage of our multi-country sampling, we examined whether rRNA or COI phylogeny

can be used to discriminate conspecifics originating from different geographies. Our assemblage

contains five of such species: *An. coustani*, *An. funestus*, *An. gambiae*, *Ae. albopictus*, and *Ma.

uniformis*. Among the rRNA trees, the concatenated 28S+18S tree and 28S tree were able to

discriminate between *Ma. uniformis* specimens from Madagascar, Cambodia, and the Central African

Republic, and between *An. coustani* specimens from Madagascar and the Central African Republic

(100% bootstrap support). In the COI tree, only *Ma. uniformis* was resolved into geographical clades

comprising specimens from Madagascar and specimens from Cambodia (72% bootstrap support). No

COI sequence was obtained from one *Ma. uniformis* from the Central African Republic. The 28S+18S

rRNA sequences ostensibly provided more population-level genetic information than COI sequences

alone with better support. The use of rRNA sequences in investigating the biodiversity of mosquitoes

should therefore be explored with a more comprehensive taxonomic assemblage.

407   The phylogenetic reconstructions based on rRNA or COI sequences in our study are hardly

congruent, but two principal differences stand out. First, the COI phylogeny does not recapitulate the

early divergence of *Anophelinae* from *Culicinae* (Figure 5). This is at odds with other studies

estimating mosquito divergence times based on mitochondrial genes (Logue et al., 2013; Lorenz et

al., 2021) or nuclear genes (Reidenbach et al., 2009). The second notable feature in the rRNA trees is

the remarkably large interspecies and intersubgeneric evolutionary distances within genus *Anopheles*

relative to *Culicinae* genera (Figures 2–4, Supplementary Figure S4) but this is not apparent in the

COI tree. The hyperdiversity among *Anopheles* taxa may be attributed to the earlier diversification of

the *Anophelinae* subfamily in the early Cretaceous period compared to that of the *Culicinae* subfamily,

a difference of at least 40 million years (Lorenz et al., 2021). The differences in rRNA and COI tree

topologies indicate a limitation in using COI alone to determine evolutionary relationships. Importantly,

drawing phylogenetic conclusions from short DNA barcodes such as COI has been cautioned against

due to its weak phylogenetic signal (Hajibabaei et al., 2006). The relatively short length of our COI

sequences (621–699 bp) combined with the 100-fold higher nuclear substitution rate of mitochondrial

genomes relative to nuclear genomes (Arctander, 1995) could result in homoplasy (Danforth et al.,

2005), making it difficult to clearly discern ancestral sequences and correctly assign branches into

lineages, as evidenced by the poor nodal bootstrap support at genus-level branches. Indeed, in the

study by Lorenz et al. (2021), a phylogenetic tree constructed using a concatenation of all 13 protein-

coding genes of the mitochondrial genome was able to resolve ancient divergence events. This

affirms that while COI sequences can be used to reveal recent speciation events, longer or multi-gene

molecular markers are necessary for studies into deeper evolutionary relationships (Danforth et al.,

2005).

In contrast to Anophelines where 28S rRNA phylogenies illustrated higher interspecies divergence

compared to COI phylogeny, two specimens of an unknown *Mansonia* species, "Ma sp.4 GF S103"

and "Ma sp.4 GF S104", provided an example where interspecies relatedness based on their COI

sequences is greater than that based on their rRNA sequences in relation to "Ma titillans GF S105".

While all rRNA trees (Figure 2–4) placed "Ma titillans GF S105" as a sister taxon with 100% bootstrap

support, the COI tree placed Ma sp.4 basal to all other species except *Ur. geometrica*. This may hint

at a historical selective sweep in the mitochondrial genome, whether arising from geographical

separation, mutations, or linkage disequilibrium with inherited symbionts (Hurst & Jiggins, 2005),

resulting in the disparate mitochondrial haplogroups found in French Guyanese Ma sp.4 and *Ma.*

titillans. In addition, both haplogroups are distant from those associated with members of subgenus

Mansonoides. To note, the COI sequences of "Ma sp.4 GF S103" and "Ma sp.4 GF S104" share

87.12 and 87.39% nucleotide similarity, respectively, to that of "Ma titillans GF S105". Interestingly,

the endosymbiont *Wolbachia pipientis* has been detected in *Ma. titillans* sampled from Brazil (De

Oliveira et al., 2015), which may contribute to the divergence of "Ma titillans GF S105" COI sequence

away from those of Ma sp.4. This highlights other caveats of using a mitochondrial DNA marker in

determining evolutionary relationships (Hurst & Jiggins, 2005), which nuclear markers such as 28S

and 18S rRNA sequences may be immune to.

Conclusions

RNA-seq metagenomics is a valuable tool for surveillance and virus discovery in non-urban

mosquitoes but it is impeded by the lack of full-length rRNA reference sequences. Here we presented

a rRNA sequence assembly strategy and 234 newly generated 28S and 18S mosquito rRNA

sequences. Our work has expanded the current rRNA reference library by providing, to our

knowledge, the first full-length rRNA records for 30 species in public databases and paves the way for

the assembly of many more. These novel rRNA sequences can improve mosquito RNA-seq

metagenomics by expanding reference sequence data for the optimization of species-specific oligo-

based rRNA depletion protocols, for streamlined species identification by rRNA barcoding and for

improved RNA-seq data clean-up. In addition, RNA barcodes could serve as an additional tool for

mosquito taxonomy and phylogeny although further studies are necessary to reveal how they

compare against other nuclear or mitochondrial DNA marker systems (Batovska et al., 2017; Beebe,

2018; Behura, 2006; Ratnasingham & Hebert, 2007; Reidenbach et al., 2009; Vezenegho et al.,

2022).

We showed that phylogenetic inferences from a tree based on 28S rRNA sequences alone or

concatenated 28S +18S rRNA sequences largely agree with contemporary mosquito classification

and can be used for species diagnostics given a reference sequence. In analysing the same set of

specimens by COI or rRNA sequences, we found deep discrepancies in phylogenetic inferences. We

conclude that while COI-based phylogeny can reveal recent speciation events, rRNA sequences may

be better suited for investigations of deeper evolutionary relationships as they are less prone to

selective sweeps and homoplasy.

MATERIALS AND METHODS

Sample collection

Mosquito specimens were sampled from 2019 to 2020 by medical entomology teams from the Institut

Pasteur de Bangui (Central African Republic, Africa; CF), Institut Pasteur de Madagascar

(Madagascar, Africa; MG), Institut Pasteur du Cambodge (Cambodia, Asia; KH), and Institut Pasteur

de la Guyane (French Guiana, South America; GF). Adult mosquitoes were sampled using several

techniques including CDC light traps, BG sentinels, and human-landing catches. Sampling sites are

non-urban locations including rural settlements in the Central African Republic, Madagascar, and

French Guiana and national parks in Cambodia. Mosquitoes were identified using morphological

identification keys on cold tables before preservation by flash freezing in liquid nitrogen and

transportation in dry ice to Institut Pasteur Paris for analysis. A list of the 112 mosquito specimens

included in our taxonomic assemblage and their related information are provided in Supplementary

Table S1.

RNA and DNA isolation

Nucleic acids were isolated from mosquito specimens using TRIzol reagent according to

manufacturer's protocol (Invitrogen, Thermo Fisher Scientific, Waltham, Massachusetts, USA). Single

mosquitoes were homogenised into 200 µL of TRIzol reagent and other of the reagents within the

protocol were volume-adjusted accordingly. Following phase separation, RNA were isolated from the

aqueous phase while DNA were isolated from the remaining interphase and phenol-chloroform phase.

From here, RNA is used to prepare cDNA libraries for next generation sequencing while DNA is used

in PCR amplification and Sanger sequencing of the mitochondrial *cytochrome* c *oxidase subunit I*

(COI) gene as further described below.

Probe depletion of rRNA

We tested a selective rRNA depletion protocol by Morlan *et al.* (2012) on several mosquito species

from the *Aedes*, *Culex*, and *Anopheles* genera. We designed 77 tiled 80 bp DNA probes antisense to

the *Ae. aegypti* 28S, 18S, and 5.8S rRNA sequences. A pool of probes at a concentration of 0.04 µM

were prepared. To bind probes to rRNA, 1 µL of probes and 2 µL of Hybridisation Buffer (100 mM

Tris-HCl and 200 mM NaCl) was added to rRNA samples to a final volume of 20 µL and subjected to

a slow-cool incubation starting at 95 ºC for 2 minutes, then cooling to 22 ºC at a rate of 0.1 ºC per

second, ending with an additional 5 minutes at 22 ºC. The resulting RNA:DNA hybrids were treated

with 2.5 µL Hybridase™ Thermostable RNase H (Epicentre, Illumina, Madison, Wisconsin, USA) and

incubated at 37 ºC for 30 minutes. To remove DNA probes, the mix was treated with 1 µL DNase I

(Invitrogen) and purified with Agencourt RNAClean XP Beads (Beckman Coulter, Brea, California,

USA). The resulting RNA is used for total RNA sequencing to check depletion efficiency.

Total RNA sequencing

To obtain rRNA sequences, RNA samples were quantified on a Qubit Fluorometer (Invitrogen) using

the Qubit RNA BR Assay kit (Invitrogen) for concentration adjustment. Non-depleted total RNA was

used for library preparation for next generation sequencing using the NEBNext Ultra II RNA Library

Preparation Kit for Illumina (New England Biolabs, Ipswich, Massachusetts, USA) and the NEBNext

Multiplex Oligos for Illumina (Dual Index Primers Set 1) (New England Biolabs). Sequencing was

performed on a NextSeq500 sequencing system (Illumina, San Diego, California, USA). Quality

control of fastq data and trimming of adapters were performed with FastQC and cutadapt,

respectively.

28S and 18S rRNA assembly

To obtain 28S and 18S rRNA contigs, we had to first clean our fastq library by separating the reads

representing mosquito rRNA from all other reads. To achieve this, we used the SILVA RNA sequence

database to create 2 libraries: one containing all rRNA sequences recorded under the "Insecta" node

of the taxonomic tree, the other containing the rRNA sequences of many others nodes distributed

throughout the taxonomic tree, hence named "Non-Insecta" (Quast et al., 2013). Each read was

aligned using the nucleotide Basic Local Alignment Search Tool (BLASTn,

https://blast.ncbi.nlm.nih.gov/) of the National Center for Biotechnology Information (NCBI) against

each of the two libraries and the scores of the best high-scoring segment pairs from the two BLASTns

are subsequently used to calculate a ratio of Insecta over Non-Insecta scores (Altschul et al., 1990).

Only reads with a ratio greater than 0.8 were used in the assembly. The two libraries being non-

exhaustive, we chose this threshold of 0.8 to eliminate only reads that were clearly of a non-insect

origin. Selected reads were assembled with the SPAdes assembler using the "-rna" option, allowing

more heterogeneous coverage of contigs and kmer lengths of 31 , 51 and 71 bases (Bankevich et al.,

2012). This method successfully assembled rRNA sequences for all specimens, including a parasitic

Horreolanus water mite (122 sequences for 28S and 114 sequences for 18S).

Initially, our filtration technique had two weaknesses. First, there is a relatively small number of

complete rRNA sequences in the Insecta library from SILVA. To compensate for this, we carried out

several filtration cycles, each time adding in the complete sequences produced in previous cycles to

the Insecta library. Second, when our mosquito specimens were parasitized by other insects, it was

not possible to bioinformatically filter out rRNA reads belonging to the parasite. For these rare cases,

we used the " --trusted-contigs" option of SPAdes, giving it access to the 28S and 18S sequences of

the mosquito closest in terms of taxonomic distance. By doing this, the assembler was able to

reconstruct the rRNA of the mosquito as well as the rRNA of the parasitizing insect. All assembled

rRNA sequences from this study have been deposited in GenBank with accession numbers

OM350214–OM350327 for 18S rRNA sequences and OM542339–OM542460 for 28S rRNA

sequences.

COI amplicon sequencing

The mitochondrial COI gene was amplified from DNA samples using the universal "Folmer" primer set

LCO1490 (5'- GGTCAACAAATCATAAAGATATTGG -3') and HCO2198 (5'-

TAAACTTCAGGGTGACCAAAAAATCA-3'), as per standard COI barcoding practices, producing a

658 bp product (Folmer et al., 1994). PCRs were performed using Phusion High-Fidelity DNA

Polymerase (Thermo Fisher Scientific). Every 50 µL reaction contained 10 µL of 5X High Fidelity

buffer, 1 µL of 10 mM dNTPs, 2.5 µL each of 10 mM forward (LCO1490) and reverse (HCO2198)

primer, 28.5 µL of water, 5 µL of DNA sample, and 0.5 µL of 2 U/µL Phusion DNA polymerase. A 3-

step cycling incubation protocol was used: 98 ºC for 30 seconds; 35 cycles of 98 ºC for 10 seconds,

60 ºC for 30 seconds, and 72 ºC for 15 seconds; 72 ºC for 5 minutes ending with a 4 ºC hold. PCR

products were size-verified using gel electrophoresis and then gel-purified using the QIAquick Gel

Extraction Kit (Qiagen, Hilden, Germany). Sanger sequencing of the COI amplicons were performed

by Eurofins Genomics, Ebersberg, Germany.

COI sequence analysis

Forward and reverse COI DNA sequences were end-trimmed to remove bases of poor quality (Q

score < 30). At the 5' ends, sequences were trimmed at the same positions such that all forward

sequences start with 5'- TTTTGG and all reverse sequences start with 5'- GGNTCT. Forward and

reverse sequences were aligned using BLAST to produce a 621 bp consensus sequence. In cases

where good quality sequences extends beyond 621 bp, forward and reverse sequences were

assembled using Pearl (**https://www.gear-genomics.com/pearl/**) and manually checked for errors

against trace files (Rausch et al., 2019, 2020). We successfully assembled a total of 106 COI

sequences. All assembled COI sequences from this study have been deposited in GenBank with

accession numbers OM630610–OM630715.

COI validation of morphology-based species identification

We analysed assembled COI sequences with BLASTn against the nucleotide collection (nr/nt)

database to confirm morphology-based species identification. BLAST analyses revealed 32 cases

where top hits indicated a different species identity, taking <95% nucleotide sequence similarity as the

threshold to delineate distinct species (Supplementary Table S2). In these cases, the COI sequence

of the specimen was then BLAST-aligned against a GenBank record representing the morphological

species to verify that the revised identity is a closer match by a significant margin, i.e., more than 2%

nucleotide sequence similarity. All species names reported hereafter reflect identities determined by

COI barcoding except for cases where COI-based identities were ambiguous, in which case

morphology-based identities were retained. In cases where matches were found within a single genus

but of multiple species, specimens were indicated as an unknown member of their genus (e.g., *Culex*

sp.). Information of the highest-scoring references for all specimens, including details of ambiguous

BLASTn results, are recorded in Supplementary Table S2.

Within our COI sequences, we found six unidentified *Culex* species (including two that matched to

GenBank entries identified only to the genus level), four unidentified *Mansonia* species, and one

unidentified *Mimomyia* species. For *An. baezai*, no existing GenBank records were found at the time

this analysis was performed.

Phylogenetic analysis

Multiple sequence alignment (MSA) were performed on assembled COI and rRNA sequences using

the MUSCLE software (Supplementary Files 1–4) (Edgar, 2004; Madeira et al., 2019). As shown in

Supplementary Figure S2, the 28S rRNA sequences contain many blocks of highly conserved

nucleotides, which makes the result of multiple alignment particularly obvious. We therefore did not

test other alignment programs. The multiple alignment of the COI amplicons is even more evident

since no gaps are necessary for this alignment.

Phylogenetic tree reconstructions were performed with the MEGA X software using the maximum-

likelihood method (S. Kumar et al., 2018). Default parameters were used with bootstrapping with 500

replications to quantify confidence level in branches. For rRNA trees, sequences belonging to an

unknown species of parasitic water mite (genus *Horreolanus*) found in our specimens served as an

outgroup taxon. In addition, we created and analysed a separate dataset combining our 28S rRNA

sequences and full-length 28S rRNA sequences from GenBank totalling 169 sequences from 58

species (12 subgenera). To serve as outgroups for the COI tree, we included sequences obtained

from GenBank of three water mite species, *Horreolanus orphanus* (KM101004), *Sperchon fuxiensis*

(MH916807), and *Arrenurus* sp. (MN362807).

DECLARATIONS

Availability of data and materials

Raw RNA-seq fastq sequence data are available from the corresponding author upon reasonable request. Multiple sequence alignment files are included as supplementary files. All sequences generated in this study have been deposited in GenBank under the accession numbers OM350214–OM350327 for 18S rRNA sequences, OM542339–OM542460 for 28S rRNA sequences, and OM630610–OM630715 for COI sequences (Supplementary Table S1).

Competing interests

The authors declare that they have no financial or non-financial competing interests.

Funding

This work was supported by the Defence Advanced Research Projects Agency PREEMPT program managed by Dr. Rohit Chitale and Dr. Kerri Dugan [Cooperative Agreement HR001118S0017] (the content of the information does not necessarily reflect the position or the policy of the U.S. government, and no official endorsement should be inferred).

Acknowledgements

We thank members of the Saleh lab for valuable discussions and to Dr Louis Lambrechts for critical reading of the manuscript. We especially thank all medical entomology staff of IP Bangui, IP Cambodge (Sony Yean, Kimly Heng, Kalyan Chhuoy, Sreynik Nhek, Moeun Chhum, Kimhuor Sour and Pierre-Olivier Maquart), IP Madagascar, and IP Guyane for assistance in field missions, laboratory work, and logistics, and Inès Partouche from IP Paris for laboratory assistance. We are also grateful to Dr Catherine Dauga for advice on phylogenetic analyses, and to Amandine Guidez for providing a French Guiana-specific COI reference library.

SUPPLEMENTARY FILES

Supplementary Figure S1. Study workflow from specimens to sequences.

Supplementary Table S1. Taxonomic and sampling information on mosquito specimens and

associated accession numbers of their 28S, 18S, and COI sequences (XLSX).

Supplementary Table S2. COI sequence BLAST analyses summary (XLSX).

Supplementary Figure S2. Sequence conservation among 169 28S rRNA sequences obtained from

this study and from GenBank combined. Multiple sequence alignment was performed on 28S rRNA

sequences, 3900 bp in length. Each bar represents a 25-bp sliding window of the 28S rRNA

sequence alignment where the values are the lowest percentage identity found.

0.05

Supplementary Figure S3. Phylogenetic tree based on 28S sequences generated from this study

(3900 bp) as inferred using maximum-likelihood method and constructed to scale in MEGA X (S.

Kumar et al., 2018) in (A) and (B) radial format. Values at each node indicate bootstrap support (%)

from 500 replication. Each specimen label contains information on taxonomy, origin (as indicated in 2-

letter country codes), and specimen ID. Label colours indicate genera: *Culex* in coral, *Anopheles* in

purple, *Aedes* in dark blue, *Mansonia* in dark green, *Limatus* in light green, *Coquillettidia* in light blue,

Psorophora in yellow, *Mimomyia* in teal, *Uranotaenia* in pink and *Eretmapodites* in brown. Scale bar

at 0.05 is shown.

Supplementary Figure S4. Phylogenetic tree based on 28S sequences generated from this study

and from GenBank (3900 bp) as inferred using maximum-likelihood method and constructed to scale

in MEGA X (S. Kumar et al., 2018) in radial format to illustrate the distance of Anopheline sequences

relative to other Culicidae members. Values at each node indicate bootstrap support (%) from 500

replications. For sequences from this study, each specimen label contains information on taxonomy,

origin (in 2-letter country codes), and specimen ID. Labels in bold indicate GenBank sequences with

accession numbers shown. Label colours indicate genera: *Culex* in coral, *Anopheles* in purple, *Aedes* in dark blue, *Mansonia* in dark green, *Culiseta* in maroon, *Limatus* in light green, *Coquillettidia* in light blue, *Psorophora* in yellow, *Mimomyia* in teal, *Uranotaenia* in pink and *Eretmapodites* in brown. Scale bar at 0.05 is shown.

Supplementary File 1: Multiple sequence alignment of 169 28S rRNA sequences from this study and from GenBank (FASTA).

Supplementary File 2: Multiple sequence alignment of 122 28S rRNA sequences, including two sequences from *Horreolanus sp.* (FASTA).

Supplementary File 3: Multiple sequence alignment of 114 18S rRNA sequences, including two sequences from *Horreolanus sp*. (FASTA).

Supplementary File 4: Multiple sequence alignment of 106 COI sequences (FASTA).

REFERENCES

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Arctander, P. (1995). Comparison of a mitochondrial gene and a corresponding nuclear pseudogene. *Proceedings of the Royal Society B: Biological Sciences*, *262*(1363), 13–19. https://doi.org/10.1098/rspb.1995.0170

Arunachalam, N., Philip Samuel, P., Hiriyan, J., Thenmozhi, V., & Gajanana, A. (2004). Japanese encephalitis in Kerala, South India: Can Mansonia (Diptera: Culicidae) play a supplemental role in transmission? *Journal of Medical Entomology*, *41*(3), 456–461. https://doi.org/10.1603/0022-2585-41.3.456

Aspen, S., & Savage, H. M. (2003). Polymerase chain reaction assay identifies North American members of the Culex pipiens complex based on nucleotide sequence differences in the acetylcholinesterase gene Ace.2. *Journal of the American Mosquito Control Association*, *19*(4), 323–328.

Auerswald, H., Maquart, P. O., Chevalier, V., & Boyer, S. (2021). Mosquito vector competence for japanese encephalitis virus. *Viruses*, *13*(6), 1154. https://doi.org/10.3390/v13061154

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M.,

Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G.,

Alekseyev, M. A., & Pevzner, P. A. (2012). SPAdes: A new genome assembly algorithm and its

applications to single-cell sequencing. *Journal of Computational Biology*, *19*(5), 455–477.

https://doi.org/10.1089/cmb.2012.0021

Barrio-Nuevo, K. M., Cunha, M. S., Luchs, A., Fernandes, A., Rocco, I. M., Mucci, L. F., DE Souza, R.

P., Medeiros-Sousa, A. R., Ceretti-Junior, W., & Marrelli, M. T. (2020). Detection of Zika and

dengue viruses in wildcaught mosquitoes collected during field surveillance in an environmental

protection area in São Paulo, Brazil. *PLoS ONE*, *15*(10), e0227239.

https://doi.org/10.1371/journal.pone.0227239

Batovska, J., Cogan, N. O. I., Lynch, S. E., & Blacket, M. J. (2017). Using next-generation sequencing

for DNA barcoding: Capturing allelic variation in ITS2. *G3: Genes, Genomes, Genetics*, *7*(1),

19–29. https://doi.org/10.1534/G3.116.036145/-/DC1

Beebe, N. W. (2018). DNA barcoding mosquitoes: Advice for potential prospectors. *Parasitology*,

145(5), 622–633. https://doi.org/10.1017/S0031182018000343

Behura, S. K. (2006). Molecular marker systems in insects: current trends and future avenues.

Molecular Ecology, *15*(11), 3087–3113. https://doi.org/10.1111/J.1365-294X.2006.03014.X

Belda, E., Nanfack-Minkeu, F., Eiglmeier, K., Carissimo, G., Holm, I., Diallo, M., Diallo, D., Vantaux,

A., Kim, S., Sharakhov, I. V., & Vernick, K. D. (2019). De novo profiling of RNA viruses in

Anopheles malaria vector mosquitoes from forest ecological zones in Senegal and Cambodia.

BMC Genomics, *20*(1), 664. https://doi.org/10.1186/s12864-019-6034-1

Bhattacharya, S., Basu, P., & Sajal Bhattacharya, C. (2016). The Southern House Mosquito, Culex

quinquefasciatus: profile of a smart vector. *Journal of Entomology and Zoology Studies JEZS*,

4(2), 73–81.

Bishop-Lilly, K. A., Turell, M. J., Willner, K. M., Butani, A., Nolan, N. M. E., Lentz, S. M., Akmal, A.,

Mateczun, A., Brahmbhatt, T. N., Sozhamannan, S., Whitehouse, C. A., & Read, T. D. (2010).

Arbovirus detection in insect vectors by Rapid, high- throughput pyrosequencing. *PLoS*

Neglected Tropical Diseases, *4*(11), e878. https://doi.org/10.1371/journal.pntd.0000878

Brault, A. C., Foy, B. D., Myles, K. M., Kelly, C. L. H., Higgs, S., Weaver, S. C., Olson, K. E., Miller, B.

R., & Powers, A. M. (2004). Infection patterns of o'nyong nyong virus in the malaria-transmitting

mosquito, Anopheles gambiae. *Insect Molecular Biology*, *13*(6), 625–635.

https://doi.org/10.1111/j.0962-1075.2004.00521.x

Cardoso, J. da C., de Almeida, M. A. B., dos Santos, E., da Fonseca, D. F., Sallum, M. A. M., Noll, C.

A., Monteiro, H. A. d. O., Cruz, A. C. R., Carvalho, V. L., Pinto, E. V., Castro, F. C., Neto, J. P.

N., Segura, M. N. O., & Vasconcelos, P. F. C. (2010). Yellow fever virus in Haemagogus

leucocelaenus and Aedes serratus mosquitoes, Southern Brazil, 2008. *Emerging Infectious*

Diseases, *16*(12), 1918–1924. https://doi.org/10.3201/eid1612.100608

Chandler, J. A., Liu, R. M., & Bennett, S. N. (2015). RNA Shotgun Metagenomic Sequencing of

Northern California (USA) Mosquitoes Uncovers Viruses, Bacteria, and Fungi. *Frontiers in*

Microbiology, *6*, 185. https://doi.org/10.3389/fmicb.2015.00185

Cornel, A. J., Mcabee, R. D., Rasgon, J., Stanich, M. A., Scott, T. W., & Coetzee, M. (2003).

Differences in Extent of Genetic Introgression between Sympatric Culex pipiens and Culex

quinquefasciatus (Diptera: Culicidae) in California and South Africa. *Journal of Medical*

Entomology, *40*(1), 36–51. https://doi.org/10.1603/0022-2585-40.1.36

Danforth, B. N., Lin, C. P., & Fang, J. (2005). How do insect nuclear ribosomal genes compare to

protein-coding genes in phylogenetic utility and nucleotide substitution patterns? *Systematic*

Entomology, *30*(4), 549–562. https://doi.org/10.1111/J.1365-3113.2005.00305.X

De Oliveira, C. D., Gonçalves, D. S., Baton, L. A., Shimabukuro, P. H. F., Carvalho, F. D., & Moreira,

L. A. (2015). Broader prevalence of Wolbachia in insects including potential human disease

vectors. *Bulletin of Entomological Research*, *105*(3), 305–315.

https://doi.org/10.1017/S0007485315000085

Diallo, D., Fall, G., Diagne, C. T., Gaye, A., Ba, Y., Dia, I., Faye, O., & Diallo, M. (2020). Concurrent

amplification of Zika, chikungunya, and yellow fever virus in a sylvatic focus of arboviruses in

Southeastern Senegal, 2015. *BMC Microbiology*, *20*, 181. https://doi.org/10.1186/s12866-020-

01866-9

Edgar, R. C. (2004). MUSCLE: A multiple sequence alignment method with reduced time and space

complexity. *BMC Bioinformatics*, *5*, 113. https://doi.org/10.1186/1471-2105-5-113

Farajollahi, A., Fonseca, D. M., Kramer, L. D., & Marm Kilpatrick, A. (2011). "Bird biting" mosquitoes

and human disease: A review of the role of Culex pipiens complex mosquitoes in epidemiology.

Infection, Genetics and Evolution, *11*(7), 1577–1585. doi: 10.1016/j.meegid.2011.08.013

Fauver, J. R., Akter, S., Morales, A. I. O., Black, W. C., Rodriguez, A. D., Stenglein, M. D., Ebel, G.

D., & Weger-Lucarelli, J. (2019). A reverse-transcription/RNase H based protocol for depletion of

mosquito ribosomal RNA facilitates viral intrahost evolution analysis, transcriptomics and

pathogen discovery. *Virology*, *528*, 181–197. https://doi.org/10.1016/j.virol.2018.12.020

Foley, D. H., Rueda, L. M., & Wilkerson, R. C. (2007). Insight into Global Mosquito Biogeography from

Country Species Records. *Journal of Medical Entomology*, *44*(4), 554–567.

https://doi.org/10.1093/JMEDENT/44.4.554

Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for amplification of

mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular*

Marine Biology and Biotechnology, *3*(5), 294–299.

Gaillet, M., Pichard, C., Restrepo, J., Lavergne, A., Perez, L., Enfissi, A., Abboud, P., Lambert, Y.,

Ma, L., Monot, M., Demar, M., Djossou, F., Servas, V., Nacher, M., Andrieu, A., Prudhomme, J.,

Michaud, C., Rousseau, C., Jeanne, I., … Rousset, D. (2021). Outbreak of oropouche virus in

french guiana. *Emerging Infectious Diseases*, *27*(10), 2711–2714.

https://doi.org/10.3201/eid2710.204760

Gale, K., & Crampton, J. (1989). The ribosomal genes of the mosquito, Aedes aegypti. *European*

Journal of Biochemistry, *185*(2), 311–317. https://doi.org/10.1111/j.1432-1033.1989.tb15117.x

Hajibabaei, M., Singer, G. A. C., & Hickey, D. A. (2006). Benchmarking DNA barcodes: An

assessment using available primate sequences. *Genome*, *49*(7), 851–854.

https://doi.org/10.1139/G06-025/SUPPL_FILE/G06-025B.PDF

Harbach, R. E. (2007). The Culicidae (Diptera): A review of taxonomy, classification and phylogeny.

Zootaxa, *1668*(1), 591–638. https://doi.org/10.11646/zootaxa.1668.1.28

Harbach, R. E., Culverwell, C. L., & Kitching, I. J. (2017). Phylogeny of the nominotypical subgenus of

Culex (Diptera: Culicidae): insights from analyses of anatomical data into interspecific

relationships and species groups in an unresolved tree. *Systematics and Biodiversity*, *15*(4),

296–306. https://doi.org/10.1080/14772000.2016.1252439

Harbach, R. E., & Kitching, I. J. (2016). The phylogeny of Anophelinae revisited: Inferences about the

origin and classification of Anopheles (Diptera: Culicidae). *Zoologica Scripta*, *45*(1), 34–47.

https://doi.org/10.1111/zsc.12137

Hebert, P. D. N., Cywinska, A., Ball, S. L., & DeWaard, J. R. (2003). Biological identifications through

DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences*, *270*(1512), 313–321.

https://doi.org/10.1098/rspb.2002.2218

Hoyos-López, R., Soto, S. U., Rúa-Uribe, G., & Gallego-Gómez, J. C. (2015). Molecular identification

of saint louis encephalitis virus genotype IV in Colombia. *Memorias Do Instituto Oswaldo Cruz*,

110(6), 719–725. https://doi.org/10.1590/0074-02760280040

Hurst, G. D. D., & Jiggins, F. M. (2005). Problems with mitochondrial DNA as a marker in population,

phylogeographic and phylogenetic studies: The effects of inherited symbionts. *Proceedings of*

the Royal Society B: Biological Sciences, *272*, 1525–1534.

https://doi.org/10.1098/rspb.2005.3056

Kim, H., Cha, G. W., Jeong, Y. E., Lee, W. G., Chang, K. S., Roh, J. Y., Yang, S. C., Park, M. Y.,

Park, C., & Shin, E. H. (2015). Detection of Japanese encephalitis virus genotype V in Culex

orientalis and Culex pipiens (Diptera: Culicidae) in Korea. *PLoS ONE*, *10*(2), e0116547.

https://doi.org/10.1371/journal.pone.0116547

Kraemer, M. U. G., Reiner, R. C., Brady, O. J., Messina, J. P., Gilbert, M., Pigott, D. M., Yi, D.,

Johnson, K., Earl, L., Marczak, L. B., Shirude, S., Davis Weaver, N., Bisanzio, D., Perkins, T. A.,

Lai, S., Lu, X., Jones, P., Coelho, G. E., Carvalho, R. G., … Golding, N. (2019). Past and future

spread of the arbovirus vectors Aedes aegypti and Aedes albopictus. *Nature Microbiology*, *4*(5),

854–863. https://doi.org/10.1038/s41564-019-0376-y

Kukutla, P., Steritz, M., & Xu, J. (2013). Depletion of ribosomal RNA for mosquito gut metagenomic

RNA-seq. *Journal of Visualized Experiments*, *74*, 50093. https://doi.org/10.3791/50093

Kumar, N., Creasy, T., Sun, Y., Flowers, M., Tallon, L. J., & Dunning Hotopp, J. C. (2012). Efficient

subtraction of insect rRNA prior to transcriptome analysis of Wolbachia-Drosophila lateral gene

transfer. *BMC Research Notes*, *5*, 230. https://doi.org/10.1186/1756-0500-5-230

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular evolutionary

genetics analysis across computing platforms. *Molecular Biology and Evolution*, *35*(6), 1547–

1549. https://doi.org/10.1093/molbev/msy096

Logue, K., Chan, E. R., Phipps, T., Small, S. T., Reimer, L., Henry-Halldin, C., Sattabongkot, J., Siba,

P. M., Zimmerman, P. A., & Serre, D. (2013). Mitochondrial genome sequences reveal deep

divergences among Anopheles punctulatus sibling species in Papua New Guinea. *Malaria*

Journal, *12*(1), 1–11. https://doi.org/10.1186/1475-2875-12-64/FIGURES/3

Lorenz, C., Alves, J. M. P., Foster, P. G., Suesdek, L., & Sallum, M. A. M. (2021). Phylogeny and

temporal diversification of mosquitoes (Diptera: Culicidae) with an emphasis on the Neotropical

fauna. *Systematic Entomology*, *46*(4), 798–811. https://doi.org/10.1111/SYEN.12489

Lutomiah, J., Bast, J., Clark, J., Richardson, J., Yalwala, S., Oullo, D., Mutisya, J., Mulwa, F., Musila,

L., Khamadi, S., Schnabel, D., Wurapa, E., & Sang, R. (2013). Abundance, diversity, and

distribution of mosquito vectors in selected ecological regions of Kenya: Public health

implications. *Journal of Vector Ecology*, *38*(1), 134–142. https://doi.org/10.1111/j.1948-

7134.2013.12019.x

Madeira, F., Park, Y. M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey, A. R. N.,

Potter, S. C., Finn, R. D., & Lopez, R. (2019). The EMBL-EBI search and sequence analysis

tools APIs in 2019. *Nucleic Acids Research*, *47*(W1), W636–W641.

https://doi.org/10.1093/nar/gkz268

Maquart, P. O., & Boyer, S. (2022). Culex vishnui. *Trends in Parasitology*, *Vector of the Month*, 491–

492. https://doi.org/10.1016/J.PT.2022.01.003

Maquart, P. O., Sokha, C., & Boyer, S. (2021). Mosquito diversity (Diptera: Culicidae) and medical

importance, in a bird sanctuary inside the flooded forest of Prek Toal, Cambodia. *Journal of

Asia-Pacific Entomology*, *24*(4), 1221–1227. https://doi.org/10.1016/j.aspen.2021.08.001

Mitchell, C. J., Forattini, O. P., & Miller, B. R. (1986). Vector competence experiments with Rocio virus

and three mosquito species from the epidemic zone in Brazil. *Revista de Saúde Pública*, *20*(3),

171–177. https://doi.org/10.1590/s0034-89101986000300001

Morlan, J. D., Qu, K., & Sinicropi, D. V. (2012). Selective depletion of rRNA enables whole

transcriptome profiling of archival fixed tissue. *PLoS ONE*, *7*(8), e42882.

https://doi.org/10.1371/journal.pone.0042882

Mukwaya, L. G., Kayondo, J. K., Crabtree, M. B., Savage, H. M., Biggerstaff, B. J., & Miller, B. R.

(2000). Genetic differentiation in the yellow fever virus vector, Aedes simpsoni complex, in

Africa: Sequence variation in the ribosomal DNA internal transcribed spacers of anthropophilic

and non-anthropophilic populations. *Insect Molecular Biology*, *9*(1), 85–91. doi: 10.1046/j.1365-

2583.2000.00161.x

Mwangangi, J. M., Muturi, E. J., Muriu, S. M., Nzovu, J., Midega, J. T., & Mbogo, C. (2013). The role

of Anopheles arabiensis and Anopheles coustani in indoor and outdoor malaria transmission in

Taveta District, Kenya. *Parasites and Vectors*, *6*, 114. https://doi.org/10.1186/1756-3305-6-114

Navarro, J. C., & Weaver, S. C. (2004). Molecular phylogeny of the Vomerifer and Pedroi Groups in

the spissipes section of the subgenus Culex (Melanoconion). *Journal of Medical Entomology*,

41(4), 575–581. https://doi.org/10.1603/0022-2585-41.4.575

Nchoutpouen, E., Talipouo, A., Djiappi-Tchamen, B., Djamouko-Djonkam, L., Kopya, E., Ngadjeu, C.

S., Doumbe-Belisse, P., Awono-Ambene, P., Kekeunou, S., Wondji, C. S., & Antonio-Nkondjio,

C. (2019). Culex species diversity, susceptibility to insecticides and role as potential vector of

Lymphatic filariasis in the city of Yaoundé, Cameroon. *PLoS Neglected Tropical Diseases*,

13(4), e0007229. https://doi.org/10.1371/journal.pntd.0007229

Ndiaye, E. H., Fall, G., Gaye, A., Bob, N. S., Talla, C., Diagne, C. T., Diallo, D., Ba, Y., Dia, I., Kohl,

A., Sall, A. A., & Diallo, M. (2016). Vector competence of Aedes vexans (Meigen), Culex

poicilipes (Theobald) and Cx. quinquefasciatus Say from Senegal for West and East African

lineages of Rift Valley fever virus. *Parasites and Vectors*, *9*, 94. https://doi.org/10.1186/s13071-

016-1383-y

Nepomichene, T. N. J. J., Raharimalala, F. N., Andriamandimby, S. F., Ravalohery, J. P., Failloux, A.

B., Heraud, J. M., & Boyer, S. (2018). Vector competence of Culex antennatus and Anopheles

coustani mosquitoes for Rift Valley fever virus in Madagascar. *Medical and Veterinary*

Entomology, *32*(2), 259–262. https://doi.org/10.1111/mve.12291

Nikolay, B., Diallo, M., Boye, C. S. B., & Sall, A. A. (2011). Usutu virus in Africa. *Vector-Borne and*

Zoonotic Diseases, *11*(11), 1417–1423. https://doi.org/10.1089/vbz.2011.0631

Phelps, W. A., Carlson, A. E., & Lee, M. T. (2021). Optimized design of antisense oligomers for

targeted rRNA depletion. *Nucleic Acids Research*, *49*(1), e5.

https://doi.org/10.1093/nar/gkaa1072

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O.

(2013). The SILVA ribosomal RNA gene database project: Improved data processing and web-

based tools. *Nucleic Acids Research*, *41*(Database issue), D590–D596.

https://doi.org/10.1093/nar/gks1219

Ratnasingham, S., & Hebert, P. D. N. (2007). BOLD: The Barcode of Life Data System: Barcoding.

Molecular Ecology Notes, *7*(3), 355–364. https://doi.org/10.1111/j.1471-8286.2007.01678.x

Ratovonjato, J., Olive, M. M., Tantely, L. M., Andrianaivolambo, L., Tata, E., Razainirina, J.,

Jeanmaire, E., Reynes, J. M., & Elissa, N. (2011). Detection, isolation, and genetic

characterization of Rift Valley fever virus from anopheles (Anopheles) coustani, anopheles

(Anopheles) squamosus, and culex (Culex) antennatus of the haute matsiatra region,

Madagascar. *Vector-Borne and Zoonotic Diseases*, *11*(6), 753–759.

https://doi.org/10.1089/vbz.2010.0031

Rausch, T., Fritz, M. H. Y., Untergasser, A., & Benes, V. (2020). Tracy: Basecalling, alignment,

assembly and deconvolution of sanger chromatogram trace files. *BMC Genomics*, *21*(1), 230.

https://doi.org/10.1186/s12864-020-6635-8

Rausch, T., Hsi-Yang Fritz, M., Korbel, J. O., & Benes, V. (2019). Alfred: Interactive multi-sample

BAM alignment statistics, feature counting and feature annotation for long- and short-read

sequencing. *Bioinformatics*, *35*(14), 2489–2491. https://doi.org/10.1093/bioinformatics/bty1007

Reidenbach, K. R., Cook, S., Bertone, M. A., Harbach, R. E., Wiegmann, B. M., & Besansky, N. J.

(2009). Phylogenetic analysis and temporal diversification of mosquitoes (Diptera: Culicidae)

based on nuclear genes and morphology. *BMC Evolutionary Biology*, *9*(1), 1–14.

https://doi.org/10.1186/1471-2148-9-298/FIGURES/4

Ruzzante, L., Reijnders, M. J. M. F., & Waterhouse, R. M. (2019). Of Genes and Genomes: Mosquito

Evolution and Diversity. *Trends in Parasitology*, *35*(1), 32–51.

https://doi.org/10.1016/J.PT.2018.10.003/ATTACHMENT/B9BE6BC5-D73A-4FEF-A654-

CC1374C59925/MMC1.MP4

Sallum, M. A. M., & Forattini, O. P. (1996). Revision of the Spissipes Section of Culex (Melanoconion)

(diptera: Culicidae). *Journal of the American Mosquito Control Association*, *12*(3), 517–600.

Sirivanakarn, S. (1982). A review of the Systematics and a Proposed Scheme of Internal

Classification of the New World Subgenus Melanoconion of Culex (Diptera, Culicidae). *Mosquito*

Systematics, *14*(4), 265–333.

Stevenson, J. C., Simubali, L., Mbambara, S., Musonda, M., Mweetwa, S., Mudenda, T., Pringle, J.

C., Jones, C. M., & Norris, D. E. (2016). Detection of plasmodium falciparum infection in

anopheles squamosus (diptera: Culicidae) in an area targeted for malaria elimination, Southern

Zambia. *Journal of Medical Entomology*, *53*(6), 1482–1487. https://doi.org/10.1093/jme/tjw091

Sun, L., Li, T. J., Fu, W. B., Yan, Z. T., Si, F. L., Zhang, Y. J., Mao, Q. M., Demari-Silva, B., & Chen,

B. (2019). The complete mt genomes of Lutzia halifaxia, Lt. fuscanus and Culex pallidothorax

(Diptera: Culicidae) and comparative analysis of 16 Culex and Lutzia mt genome sequences.

Parasites and Vectors, *12*, 368. https://doi.org/10.1186/s13071-019-3625-2

Takhampunya, R., Kim, H. C., Tippayachai, B., Kengluecha, A., Klein, T. A., Lee, W. J., Grieco, J., &

Evans, B. P. (2011). Emergence of Japanese encephalitis virus genotype v in the Republic of

Korea. *Virology Journal*, *8*, 449. https://doi.org/10.1186/1743-422X-8-449

Talaga, S., Duchemin, J. B., Girod, R., & Dusfour, I. (2021). The Culex Mosquitoes (Diptera:

Culicidae) of French Guiana: A Comprehensive Review With the Description of Three New

Species. *Journal of Medical Entomology*, *58*(1), 182–221. https://doi.org/10.1093/JME/TJAA205

Thongsripong, P., Chandler, J. A., Kittayapong, P., Wilcox, B. A., Kapan, D. D., & Bennett, S. N.

(2021). Metagenomic shotgun sequencing reveals host species as an important driver of virome

composition in mosquitoes. *Scientific Reports*, *11*(1), 8448. https://doi.org/10.1038/s41598-021-

87122-0

Torres-Gutierrez, C., Bergo, E. S., Emerson, K. J., de Oliveira, T. M. P., Greni, S., & Sallum, M. A. M.

(2016). Mitochondrial COI gene as a tool in the taxonomy of mosquitoes Culex subgenus

Melanoconion. *Acta Tropica*, *164*, 137–149. https://doi.org/10.1016/j.actatropica.2016.09.007

Torres-Gutierrez, C., De Oliveira, T. M. P., Emerson, K. J., Bergo, E. S., & Sallum, M. A. M. (2018).

Molecular phylogeny of Culex subgenus Melanoconion (Diptera: Culicidae) based on nuclear

and mitochondrial protein-coding genes. *Royal Society Open Science*, *5*, 171900.

https://doi.org/10.1098/rsos.171900

Turell, M. J., O'guinn, M. L., Dohm, D., Zyzak, M., Watts, D., Fernandez, R., Calampa, C., Klein, T. A.,

& Jones, J. W. (2008). Susceptibility of Peruvian Mosquitoes to Eastern Equine Encephalitis

Virus. *Journal of Medical Entomology*, *45*(4), 720–725.

https://doi.org/10.1093/JMEDENT/45.4.720

Turell, Michael J. (1999). Vector competence of three Venezuelan mosquitoes (Diptera: Culicidae) for

an epizootic IC strain of Venezuelan equine encephalitis virus. *Journal of Medical Entomology*,

36(4), 407–409. https://doi.org/10.1093/jmedent/36.4.407

Ughasi, J., Bekard, H. E., Coulibaly, M., Adabie-Gomez, D., Gyapong, J., Appawu, M., Wilson, M. D.,

& Boakye, D. A. (2012). Mansonia africana and Mansonia uniformis are Vectors in the

transmission of Wuchereria bancrofti lymphatic filariasis in Ghana. *Parasites and Vectors*, *5*(1),

1–5. https://doi.org/10.1186/1756-3305-5-89

Valentine, M. J., Murdock, C. C., & Kelly, P. J. (2019). Sylvatic cycles of arboviruses in non-human

primates. *Parasites and Vectors*, *12*(1), 1–18. https://doi.org/10.1186/S13071-019-3732-0/TABLES/4

Vasconcelos, P. F. C., Costa, Z. G., Travassos da Rosa, E. S., Luna, E., Rodrigues, S. G., Barros, V. L. R. S., Dias, J. P., Monteiro, H. A. O., Oliva, O. F. P., Vasconcelos, H. B., Oliveira, R. C., Sousa, M. R. S., Barbosa Da Silva, J., Cruz, A. C. R., Martins, E. C., & Travassos Da Rosa, J. F. S. (2001). Epidemic of jungle yellow fever in Brazil, 2000: Implications of climatic alterations in disease spread. *Journal of Medical Virology*, *65*(3), 598–604. https://doi.org/10.1002/jmv.2078.abs

Vázquez González, A., Ruiz, S., Herrero, L., Moreno, J., Molero, F., Magallanes, A., Sánchez-Seco, M. P., Figuerola, J., & Tenorio, A. (2011). West Nile and Usutu viruses in mosquitoes in Spain, 2008-2009. *American Journal of Tropical Medicine and Hygiene*, *85*(1), 178–181. https://doi.org/10.4269/ajtmh.2011.11-0042

Vezenegho, S. B., Issaly, J., Carinci, R., Gaborit, P., Girod, R., Dusfour, I., & Briolant, S. (2022). Discrimination of 15 Amazonian Anopheline Mosquito Species by Polymerase Chain Reaction—Restriction Fragment Length Polymorphism. *Journal of Medical Entomology*, *59*(3), 1060–1064. https://doi.org/10.1093/JME/TJAC008

Weaver, S. C., Ferro, C., Barrera, R., Boshell, J., & Navarro, J. C. (2004). Venezuelan Equine Encephalitis. *Annual Review of Entomology*, *49*, 141–174. https://doi.org/10.1146/annurev.ento.49.061802.123422

Webster, J. P., Gower, C. M., Knowles, S. C. L., Molyneux, D. H., & Fenton, A. (2016). One health - an ecological and evolutionary framework for tackling Neglected Zoonotic Diseases. *Evolutionary Applications*, *9*(2), 313–333. https://doi.org/10.1111/eva.12341

Weedall, G. D., Irving, H., Hughes, M. A., & Wondji, C. S. (2015). Molecular tools for studying the major malaria vector Anopheles funestus: Improving the utility of the genome using a comparative poly(A) and Ribo-Zero RNAseq analysis. *BMC Genomics*, *16*(1), 931. https://doi.org/10.1186/s12864-015-2114-z

WHO. (2017). Global vector control response 2017–2030. In *World Health Organization*.

Zakrzewski, M., Rašić, G., Darbro, J., Krause, L., Poo, Y. S., Filipović, I., Parry, R., Asgari, S., Devine, G., & Suhrbier, A. (2018). Mapping the virome in wild-caught Aedes aegypti from Cairns and Bangkok. *Scientific Reports*, *8*(1), 4690. https://doi.org/10.1038/s41598-018-22945-y

941 Zittra, C., Flechl, E., Kothmayer, M., Vitecek, S., Rossiter, H., Zechmeister, T., & Fuehrer, H. P.

942  (2016). Ecological characterization and molecular differentiation of Culex pipiens complex taxa

943  and Culex torrentium in eastern Austria. *Parasites and Vectors*, *9*, 197.

944  https://doi.org/10.1186/s13071-016-1495-4

B Ae simpsoni CF S27 (182,419 reads)

>=0.8 ,Insecta = 175671 (96.3%)
<0.8 ,Others =325 (0.18%)
No hits =6423 (3.52%)

- - - hits only in the Insecta library
– – – hits higher-scoring in Insecta library
······ hits higher-scoring in Non-Insecta library
······ hits only in Non-Insecta library

0.05