Title:

Why SARS-CoV-2 Omicron variant is milder? A single high-frequency mutation of structural envelope protein matters.

Author:

Bingqing Xia[1,2,*], Yi wang[1,3], Xiaoyan pan[4], Xi Cheng[1,2], Hongying Ji[1], Xiaoli Zuo[1],Jia Li[1,2], Zhaobing Gao[1,2,*]

[1] CAS Key Laboratory of Receptor Research, Stake Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 555 Zuchongzhi Road, Shanghai, 201203, China.

[2] University of Chinese Academy of Sciences, 19A Yuquan Road, Beijing, 100049, China.

[3] State Key Laboratory of Medical Neurobiology, Institutes of Brain Science, Fudan University, Shanghai 200032, China.

[4] State Key Laboratory of Virology, Wuhan Institute of Virology, Center for Biosafety Mega-Science, Chinese Academy of Sciences, Wuhan, 430071, China.

These authors contributed equally: Bingqing Xia, Yi wang

*Corresponding author. Email: zbgao@simm.ac.cn (Z.G.), xiabingqing@simm.ac.cn (B.X.).

Abstract:

SARS-CoV-2 Omicron variant is highly transmissible and extensive morbidity, which

has raised concerns for antiviral therapy. In addition, the molecular basis for the

attenuated pathogenicity and replication capacity of Omicron remains elusive. Here, we

report for the first time that a high-frequency mutation T9I on 2-E of SARS-CoV-2

variant Omicron forms a non-selective ion channel with abolished calcium permeability

and reduced acid sensitivity compared to the WT channel. In addition, T9I caused less

cell death and a weaker cytokine production. The channel property changes might be

responsible for the Omicron variant releases less efficiently and induces a

comparatively lower level of cell damage in the infected cells. Our study gives valuable

insights into key features of the Omicron variant, further supporting 2-E is a promising

drug target against SARS-CoV-2 and providing critical information for the COVID-19

treatment.

Main Text:

Since November 2021, a new SARS-CoV-2 variant (B.1.1.529) emerged in South Africa and was designated as the fifth variant of concern (VOC), named as Omicron [1,2]. Omicron variant is highly transmissible and extensive morbidity, which has raised concerns for antiviral therapy. Omicron has largest number (>30) of substitutions, deletions, or insertions with mutations frequency higher than 1% in the spike protein (S). Mutation studies in the spike RBD domain have provided a plausible explanation for altered transmissibility and antibody resistance [3-5]. Recent reports revealed Omicron variant induces a longer cycle for virus shed, attenuates replication capacity, and produces substantially attenuated lung pathology, indicating that the pathogenic ability of Omicron variant is indeed milder [6-8]. However, the molecular basis for the attenuated pathogenicity and replication capacity remains elusive. The envelope protein of SARS-CoV-2 (2-E) forms a homo-pentameric cation channel that is important for virus pathogenicity [9,10]. It has been demonstrated that 2-E channel alone is sufficient to induce cell death, provoke cytokine storm and even cause acute respiratory distress syndrome (ARDS)-like damages *in vivo*. Its inhibitor exhibits excellent antiviral activity *in vivo*[9]. Therefore, we focus on the 2-E mutations of Omicron variant.

Three high quality E protein sequences of Omicron variants (B.1.1.529, BA1, BA2) in the CNCB database as of January 18 th, 2022 (https://ngdc.cncb.ac.cn/ncov/) were comprehensively analyzed and a mutation T9I caught our attention. Compared with the original Wuhan reference strain, among more than one hundred identified 2-E

mutations in Omicron variant, T9I shows a much higher frequency than the rest

mutations. In addition, the mutation probability of T9I in all statistical samples is higher

than 99.5%. In particular, for the most common Omicron strain B.1.1.529, the mutation

probability is 100% (Fig.1a, b). According to the solved NMR structure, T9 locates at

the top of the transmembrane domain (TMD) of E protein [10]. The influence of T9I has

not been studied yet completely.

First, we asked whether 2-E^{T9I} (T9I) still retains its channel activity. We purified the

T9I protein and reconstituted the protein on Planar Lipid Bilayer (BLM)

(Supplementary Fig.S1). The observed T9I-induced typical single-channel currents

supports that T9I was able to form ion channels also as wild-type proteins. However,

the reversal potential of T9I channels was shifted to the left, from 57 mV to 3 mV, under

asymmetric KCl solutions. Similar reversal potential shifting was also detected in

asymmetric NaCl solutions (Fig.1c.d, Supplementary Fig.S2). A reversal potential close

to 0 mV suggests that T9I may have lost its selectivity to cations. To determine whether

T9I can permeate chloride，the channel activity of T9I was further examined in

asymmetric 50:500 mM choline chloride solutions. Choline chloride is a quaternary

ammonium salt containing choline cation and chloride anion, which is always to detect

anion permeability. As expected, outward currents were indeed observed when the

holding voltage potentials were higher than the theoretical reversal potential for

chloride (E_{rev}> 70 mV). 2-E^{WT} (WT) channels are permeable to Ca^{2+}. As shown in

Figure S2c, for WT channels, the frequent and continuous inward potassium currents

in the asymmetric 50:500 mM (trans: cis) K^+ solutions indicated that the 2-E channels

were incorporated into the membranes. The membrane potential was then changed to

+65 mV (K^+ reverse potential) to eliminate the K^+ currents. Intriguingly, outward step-

like signals appeared when 20 mM (final concentration) Ca^{2+} was added to the *trans*

side. When the Ca^{2+} concentration increased to 20 mM, the outward currents increased.

In contrast, under the identical conditions, T9I channels did not induce detectable

calcium currents (Supplementary Fig.S2c). These results revealed that T9I owns a

different ion selectivity and permeability from WT. WT channels are pH sensitive [9].

The pH influences on T9I channels were evaluated on a same channel using titration

strategy. After the channels incorporated into the membranes, HCl was titrated into

either *cis* or *trans* side to alter the pH to 4 from 6. Consistent to our previous results,

reduction of pH, either in *cis* or *trans* side, induced significant increase of channel

amplitude and open probability. In contrast, pH reduction in cis side failed to increase

the channel activity of T9I. Although we didn't have direct evidences to explain how

the threonine (T9) site is sensitive to pH changes, the structure of 2-E channels

supported the residues around T9 were sensitive to pH change including Glu7 and Glu8

side chain carboxyl could be deprotonated at neutral pH and protonated at acidic pH [10].

It is reported that SARS-CoV-2 traffics to the lysosomes for egress by lysosome

deacidification, instead of using the conventional biosynthetic secretory pathway [11].

The E channel of coronavirus is localized to intracellular organelles, including

lysosome and Golgi apparatus [11, 12]. The cellular localization and expression of WT and

T9I were assessed by immunocytochemistry and confocal microscopy. We found that

both WT and T9I are co-localized with the lysosomal marker LAMP (Fig.1g,

supplementary Fig.S3). Given the localization on lysosome, the influences of WT and

T9I on the luminal pH of lysosome were compared. We used flow cytometric analysis

of a ratio metric pHluorin molecule. This pH probe consists of green fluorescent protein

(GFP) pHluorin molecule fused with lysosome-resident protein CD63. To generate a

pH calibration curve, un-transfected cells were subjected to treatment with buffers

ranging from pH 5.5 to 7.5 in the presence of the ionophores monensin and nigericin

prior to flow analysis (Fig.1h, i). Expression of WT channels robustly neutralized the

lysosome pH, whereas T9I channels exhibited less influences in luminal pH, which was

consistent with our electrophysiological results (Fig.1j). Therefore, the capability of

deacidification of 2-E channels in lysosomes was significantly impeded by the mutation

T9I. Based on above, whether the weakened capability of deacidification hampers virus

releasing was examined. We transfected the Vero E6 cells with WT or T9I plasmids at

day 1, then the cells were infected with SARS-CoV-2 virus at day 2. As determined by

qRT-PCR, pre-expression of mutant T9I significantly attenuated viral loadings

compared with WT (Fig.1k). It was logic to deduce that some exogenous [T9I] subunits

would co-assemble with the wild type subunits from the virus and form

heteromultimers with impaired pH sensitivity, resulting in less deacidification of

lysosomes and less SARS-CoV-2 virus releasing.

Finally, the channel activity of the 2-E protein is a determinant of virulence. We used

Annexin V/PI staining and CCK8 assays to evaluate the cell lethality of T9I in Vero E6

cells. We found that heterogeneously expression of T9I caused similar cell death with

WT (Fig. 1 l, Fig.1g, supplementary Fig.S4a). In the mouse macrophage cell line

(RAW264.7), T9I induced a similar increase of inflammatory cytokines with WT also

(Fig. 1m, Supplementary Fig.S4b). However, in the case of comparable cell lethality

and cytokine releasing level, the expression level of T9I was far exceeding the

expression level of WT (Fig. 1 n). The correlation between the cell lethality and the

expression level was further compared. As shown in Figure 1o, the killing cells

capability of T9I was dropped around 150 times as well as the cytokines production.

Notably, the 2-E expression in Omicron variant was decreased approximately 3-fold,

which may further suppress the virulence of Omicron variants [6]. Consistently, it has

been demonstrated that the ability of Omicron to induce cell death was indeed abolished

within 48 hours [6]. How much the weakened virulence of T9I confers the milder clinical

symptoms of Omicron warrants further investigation.

Overall, as SARS-CoV-2 infections are progressing, there are chances that virus

accumulates new mutations. Among existed antiviral targets, different from spike

protein, RNA-dependent RNA polymerase (Rdrp) and 3CL pro, which mediate virus

entry, protein cleavage and viral replication, the envelope protein (2-E) majorly

participating in the virus releasing [13-15]. Here we report for the first time that a high-

frequency mutation T9I on 2-E of SARS-CoV-2 variant Omicron forms a non-selective

ion channel with abolished calcium permeability and reduced acid sensitivity compared

to the WT channel. In addition, T9I caused less cell death and a weaker cytokine

production (Fig.1p). The channel property changes might be responsible for the

Omicron variant releases less efficiently and induces a comparatively lower level of

cell damage in the infected cells. According to 2-E NMR structures, although T9I is not

located in the pore domain of the channel, it may alter the channel configuration via interacting with the amino acids at positions L12 and T11lining the pore (Supplementary Fig.S5). Our study gives valuable insights into key features of the Omicron variant, further supporting 2-E is a promising drug target against SARS-CoV-2 and providing critical information for the COVID-19 treatment.

Acknowledgement:

We are grateful to the National Science Fund of Distinguished Young Scholars (81825021), Fund of Youth Innovation Promotion Association (2019285), the National Natural Science Foundation of China (81773707, 31700732), the National Key Research and Development Program of China (2020YFC0842000), the Strategic Leading Science and Technology Projects of Chinese Academy of Sciences (XDA12050308), Fund of National Science and Technology Major Project (2018ZX09711002-002-006) and the Hubei Science and Technology Project (2020FCA003) for financial support.

Author contributions:

Z. G., and B. X. conceived designed the project. Z. G., J. L., and B. X designed the experiments; Y. W. and B. X. performed the electrophysiological recordings; B. X. carried out the cell-based assays; P. X. carried out the virus assays in vitro; Y. W purified the proteins; H.Y. analyzed the sequence; all authors analyzed and discussed the data. Z. G., B. X. wrote the manuscript. All authors read and approved the manuscript.

Reference:

Karim SSA, Karim QA. Omicron SARS-CoV-2 variant: a new chapter in the COVID-19 pandemic. *Lancet (London, England)* 2021; **398**:2126-2128.

Organization WH. Classification of Omicron (B.1.1.529): SARS-CoV-2 Variant of Concern. <*https://wwwwhoint/news/item/26-11-2021-classification-of-omicron-(b11529)-sars-cov-2-variant-of-concern*>.

Cao Y, Wang J, Jian F *et al.* Omicron escapes the majority of existing SARS-CoV-2 neutralizing antibodies. *Nature* 2021.

Cele S, Jackson L, Khoury DS *et al.* Omicron extensively but incompletely escapes Pfizer BNT162b2 neutralization. *Nature* 2021.

Bojkova D, Widera M, Ciesek S, Wass MN, Michaelis M, Cinatl J, Jr. Reduced interferon antagonism but similar drug sensitivity in Omicron variant compared to Delta variant of SARS-CoV-2 isolates. *Cell research* 2022:1-3.

Shuai H, Chan JF, Hu B *et al.* Attenuated replication and pathogenicity of SARS-CoV-2 B.1.1.529 Omicron. *Nature* 2022.

Halfmann PJ, Iida S, Iwatsuki-Horimoto K *et al.* SARS-CoV-2 Omicron virus causes attenuated disease in mice and hamsters. *Nature* 2022.

Torjesen I. Covid-19: Peak of viral shedding is later with omicron variant, Japanese data suggest. *BMJ (Clinical research ed)* 2022; **376**:o89.

Xia B, Shen X, He Y *et al.* SARS-CoV-2 envelope protein causes acute respiratory distress syndrome (ARDS)-like pathological damages and constitutes an antiviral target. *Cell research* 2021; **31**:847-860.

Mandala VS, McKay MJ, Shcherbakov AA, Dregni AJ, Kolocouris A, Hong M.

Structure and drug binding of the SARS-CoV-2 envelope protein transmembrane domain in lipid bilayers. *Nature structural & molecular biology* 2020; **27**:1202-1208.

11 Ghosh S, Dellibovi-Ragheb TA, Kerviel A *et al.* β-Coronaviruses Use Lysosomes for Egress Instead of the Biosynthetic Secretory Pathway. *Cell* 2020; **183**:1520-1535.e1514.

12 Westerbeck JW, Machamer CE. The Infectious Bronchitis Coronavirus Envelope Protein Alters Golgi pH To Protect the Spike Protein and Promote the Release of Infectious Virus. *Journal of virology* 2019; **93**.

13 Yin W, Mao C, Luan X *et al.* Structural basis for inhibition of the RNA-dependent RNA polymerase from SARS-CoV-2 by remdesivir. *Science (New York, NY)* 2020; **368**:1499-1504.

14 Jin Z, Du X, Xu Y *et al.* Structure of M(pro) from SARS-CoV-2 and discovery of its inhibitors. *Nature* 2020; **582**:289-293.

15 Liu L, Wang P, Nair MS *et al.* Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2 spike. *Nature* 2020; **584**:450-456.

Figure:

Fig.1 The envelope protein mutation T9I of SARS-CoV-2 Omicron variant shows

less pathogenicity and weakens virus production.

a, Structure of the 2-E TM domain [10]. Illustration of the amino-acid sequence of SARS-

CoV-2 E (8-38) and the indicated mutants. Threonine residues (T) are marked as blue

and the T >I mutants are marked as red. **b,** 2-E protein amino-acid sequence logo of

SARS-CoV-2 Omicron variant. The I9 residues are marked as red. 2-E protein

sequences were downloaded from China National Center for Bioinformation (CNCB)

using the "Lineage Browse" option. Logos were generated using the WebLogo 3.7.4

online tool. **c,** Single-channel current recording of T9I after reconstitution in lipid

bilayers with PC/PS = 3:2 lipids at indicated potentials and solutions (Left). **d,** Left, all-

point current histograms for the trace in **c** (0 mV) ("C" means Closed; "O" means Open).

Right, *I-V* curves of T9I in the solutions of **c** (n ≥ 3). **e,** Single-channel current recording

in 500 :500 mM KCl solutions via proton titration result in different pH (*cis* or *trans*).

Red arrow represents titration of HCl. **f,** The heat map shows the transition of the

currents from panel **e** under proton titration. **g,** Cell immunofluorescence of T9I (Red)

and lysosome marker LAMP (Green). T9I is co-localized with lysosome. **h-j,** Lyso-pH

indicator was used to assess the pH of the lysosome via flow cytometry. Flow cytometry

pattern of the cells in buffers of known pH and contained ionophores to equilibrate the

extracellular and lysosome luminal pH (**h**). Calibration curves were generated from data

like those illustrated in h (**i**). **j,** The pH value of Vero E6 cells lysosome transfected with

WT, T9I or Mock. **k.** Flow chart of the experiment (Left), the virus copies in supernatant

determined by qRT-PCR assay (Right). MOI of SARS-CoV-2 for infection was 0.01. **l,**

Flow cytometry pattern of Vero E6 cells transfected with WT or T9I plasmids. Right,

flow cytometry analysis of Propidium iodide (PI) and Annexin V stained Vero E6 cells.

m, Serum cytokine levels at 24 h after transfection with WT, T9I or Mock detected by

RT-qPCR. **n,** Protein expression levels of WT and T9I in Vero E6 cells after transfection

(Left). The histogram shows the quantization of the result (Right). **o,** The relative

activity of WT, T9I in pH regulation, cell virulence, channel conductance and viral

copies normalized to the expression levels. **p,** Life cycle pattern diagram of SARS-

CoV-2 Omicron variant. T9I on envelope protein (2-E) of SARS-CoV-2 variant

Omicron forms a non-selective ion channel with weakened pathogenicity and virus

production. Experiments were independently performed thrice and similar results were

obtained. One set of reprehensive data is shown here (n ⩾ 3). *$p < 0.05$; **$p < 0.01$;

***$p < 0.001$; unpaired Student's t test. All error bars are SEM.

Method:

Plasmids and mutagenesis

Wild type SARS-CoV-2-E sequences were synthesized by the Beijing Genomics

Institute (BGI, China). The mutants 2-E^{T9I} sequence was generated by site-directed

mutagenesis and confirmed by sequencing (BGI, China). Vector pET28a was used for

protein purification; vector pcDNA5 and pcDNA3.1 were used for cell survival assay

and cell imaging.

Cell culture and treatment

Vero E6 and Raw 264.7 cells were grown in 90% DMEM basal medium (Gibco, USA)

supplemented with 10% fetal bovine serum (Gibco, USA) at 37°C under 5% CO_2.

Planar lipid bilayers recording

According to our previous methods, the purified protein 2-E^{WT} or 2-E^{T9I} were

incorporated into lipid bilayers to test their functionality. The lipid bilayers were

prepared by PC: PS= 3: 2 (Avanti Polar Lipids, USA). Proteins were added in *cis* side

and the solution was 50:500 mM (*trans*: *cis*) KCl or others, all solutions were buffered

by 5mM HEPES, pH 6.35. Membrane currents were recorded by headstage connected

with amplifier BC-535 (Warner Instruments, USA), filtered at 1–2 kHz and digitized

by Axon Digidata 1440A (Molecular Devices, USA). The recording frequency was 10

kHz. The data were processed using pClamp 10.2 software (Molecular Devices, US).

The single-channel conductance was determined by fitting to Gaussian functions (bin

width = 0.25 pA). Opening time less than 0.5–1.5 ms was ignored to avoid noise

interference. The equilibrium potential was calculated using Goldman–Hodgkin–Katz

flux equation:

$$E_m = \frac{RT}{F} \ln \left(\frac{P_{Na^+}[Na^+]_{out} + P_{Cl^-}[Cl^-]_{in} + P_{K^+}[K^+]_{out}}{P_{Na^+}[Na^+]_{in} + P_{Cl^-}[Cl^-]_{out} + P_{K^+}[K^+]_{in}} \right)$$

Cell viability and cytotoxicity assay

Cell viability was measured by the CCK-8 kit (40203ES60, Yeason, China) according

to the manufacturer's instructions. The absorbance was measure at 450 nm with Thermo

Scientific Microplate Reader (Thermo Fisher Scientific Inc., USA).

Western blotting and antibodies

Proteins were resolved in 12% SDS-PAGE, transferred to PVDF membranes (GE,

USA). The PVDF membranes were incubated with primary antibodies against HA-Tag

(C29F4) (3724, CST, USA) or GAPDH (30201ES20, Yeasen, China), then incubated

with second antibodies against IgG (Yeasen, China).

Flow cytometry of cell death

After treatment, cells were detached, collected by centrifugation and resuspended in 1×

binding buffer containing 100 μg/mL propidium iodide (PI) and Annexin V-FITC (1:20,

V13242, Life, USA) and incubated at room temperature for 15 min in the dark.

Subsequently, samples were added 400 μL of 1× binding buffer and kept on ice. Cells

were analyzed by flow cytometry using BD FACSCALIBUR 4 (BD COMPANY, USA).

The percentages of differently labeled cells were calculated by FlowJo 7.6.

Immunofluorescence microscopy.

Vero E6 cells were seeded as 1×10^4 cells per well in coverslips on chamber slides

(Thermo Fisher Scientific Inc., USA) overnight. Cells were transfected with WT or T9I

with lysosome marker (LAMP1,). After transfection for 24 h, cells were washed with

phosphate-buffered saline (PBS) and fixed in 4% paraformaldehyde for 10 min at room

temperature. Then the cells were permeabilized in 0.5% Triton X-100   in PBS for 3

min. The coverslips were washed twice with PBS and incubated with primary antibody

in 1% bovine serum albumin (BSA) (36101ES25, Yeasen, China) for 2 h at room

temperature. Anti-HA (HT301-01, Trans, China) and rabbit anti-LAMP (55273-1-AP,

Proteintech, USA) were used to incubate the cells at a 1:1,000 dilution. Then, second

antibodies FITC-AffiniPure Goat Anti-Rabbit IgG (33107ES60, Yeasen, China) and

Cy3-AffiniPure Goat Anti-Mouse IgG (33208ES60, Yeasen, China)were used to

incubate the cells at a 1:1,000 dilution. Images were captured using the Leica TCS-SP8

STED system with 100× oil objective.

Virus loading and quantitative real-time PCR (qRT-PCR) analysis

SARS-CoV-2 (nCoV-2019BetaCoV/Wuhan/WIV04/2019) was preserved at Wuhan

institute of virology, Chinese Academy of Sciences. It was propagated and titrated with

Vero E6 cells, and its associated operations were performed in a biosafety level 3 (BSL-

3) facility. Cell supernatants RNA was isolated with MiniBEST Viral RNA/DNA

Extraction Kit (Takara, Japan) as described in the instruction, and cDNA was

transcribed with PrimeScript^{TM} RT reagent Kit with gDNA Eraser (Takara, Japan). In

detail, 50 μL supernatant were collected for RNA isolation with a MiniBEST Viral

RNA/DNA Extraction Kit Ver.5.0 (Takara, AK41820A), and the total RNA was eluted

with 30 μL RNase-free water. cDNA was transcribed from 3 μL total RNA in 20 μL

reaction system with PrimeScriptTM RT reagent Kit with gDNA Eraser (Takara,

AK71648A). Viral copies were quantified from 1 μL template viral cDNA by a standard

curve method on ABI 7500 (Takara TB Green Premix Ex Taq II, AK81975A) with a

pair of primers targeting S gene. The forward primer (5'-3') is:

CAATGGTTTAACAGGCACAGG; the reverse primer (5'-3') is:

CTCAAGTGTCTGTGGATCACG.

Determination of lysosome pH

Vero cells were transfected with Lyso-pHluorin (addgene,70113) and 2-E plasmid.

After transfection, cells were trypsinzed and washed with Live Cell Imaging Solution

(LCIS). Then, replaced LCIS with cellular pH calibration Buffers containing 10 μM

Valinomycin and Nigericin (Intracellular pH Calibration Buffer Kit, P35379, Themo,

USA) and incubate at 37°C for at least 5 minutes. At last, cells were analyzed via Flow

cytometry with excited at 405 nm and 488 nm, and the emission signals were collected

with detection filters at 500 to 550 nm and 515 to 545 nm, respectively. Flow cytometric

data were collected and quantified using CytoExpert 2.4 software.

Homology modeling

The solid-state NMR structure of SARS-CoV-2 envelope protein (PDB code: 7k3g)

[(2020) Nat Struct Mol Biol 27: 1202-1208] was used to as the template to build

homology models of the mutant envelope protein. This solid-state data yield to ten

lowest-energy structures. These ten structures were employed to construct ten

homology models of the mutant envelope protein using Modeller [A. Sali, T.L. Blundell,

Comparative protein modelling by satisfaction of spatial restraints, J Mol Biol 234

(1993) 779-815. 10.1006/jmbi.1993.1626.]. The models with the lowest root mean

square deviations from their template structures were selected.

Statistics

All measurements were derived from distinct samples. Statistics were performed in

GraphPad Prism. Statistical significance was determined by one-way ANOVA and two-

way ANOVA followed by pairwise Student's t-test with Tukey's or Sidak's correction.

Two-tailed unpaired Student's t test was performed if only two conditions were

compared. Data in the text is presented as Mean $\pm$ SEM. Adjusted P values are reported

in the figures.

 Supplementary figure：

338 **Supplementary information, Fig. S1 Purification of T9I proteins.** Purification of

339 full-length 2-E^{T9I} protein with Ni-NTA affinity chromatography. 15% SDS-PAGE gel

340 with coomassie blue staining.

342 **Supplementary information, Fig. S2 Electrophysiological property of T9I.**

a. Single-channel current recording of T9I after reconstitution in lipid bilayers with PC/PS = 3:2 lipids at indicated potentials and solutions. Protein (5–50 ng/mL) was added to the cis side. All-point current histograms for the left trace (0, 150mV) ("C" means Closed; "O" means Open). **b.** *I-V* curves of T9I in indicated solutions (n ≥ 3). **c.** Pure Ca^{2+} currents recording in 50 :500 mM KCl solutions (*trans*: *cis*) via calcium titration as indicated concentrations, red arrow represents titration of $CaCl_2$. **d.** Amplitude (*I*) and open probability (P_o) of channel current at pH 4 were normalized to channel current at pH 6 of both sides (n ≥ 3).

Supplementary information, Fig. S3 Wild type envelope protein (E) is co-localized with lysosome. Cell immunofluorescence of WT (Red) and lysosome marker (Green). Scale bar,10 μM.

Supplementary information, Fig. S4 Cell viability of Vero E6 cells after transfection with 2-E plasmids and expression level of after transfection with WT

andT9I.

a. Cell viability of Vero E6 cells after transfection with WT, T9I or Vector (Mock) at

24 h. **b.** Expression level of 2-E after transfection with WT, T9I or Mock in Raw

264.7 cells detected by RT-qPCR.

Supplementary information, Fig. S5 Structure of WT and T9I.

a, Top view of ten lowest-energy solid-state NMR structures of wildtype SARS-CoV-2

E protein (PDB code: 7k3g). Three N-terminal residues T9, T11, L12 and a pore-lining

residue N15 are shown as sticks (T9 in *yellow*, T11 in *green*, L12 in *orange*, N15 in

light blue). The protein is shown as ribbon. **b,** Top view of ten homology models of2-

E^{T9I}, which built based on the solid-state NMR structures of wildtype SARS-CoV-2

envelope protein (PDB code: 7k3g). Three N-terminal residues I9, T11, L12 and a pore-

lining residue N15 are shown as sticks (I9 in *yellow*, T11 in *green*, L12 in *orange*, N15

in *light blue*).