**Single-cell RNA profiling of *Plasmodium vivax* liver stages reveals parasite- and host-**
**specific transcriptomic signatures and drug targets**

Anthony A Ruberto [a], Steven P Maher [a], Amélie Vantaux [b], Chester J Joyner [a,c], Caitlin Bourke [d], Balu
Balan [d,e] , Aaron Jex [d,e], Ivo Mueller [d] , Benoit Witkowski [b], Dennis E Kyle [a]

[a] Center for Tropical and Emerging Global Diseases, University of Georgia, Athens, GA, United States

[b] Malaria Molecular Epidemiology Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia

[c] Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia, Athens,
GA, USA

[d] Population Health & Immunity Division, Walter and Eliza Hall Health Institute, Parkville, Victoria,
Australia

[e] Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Parkville, Victoria,
Australia

Running Head: Single-cell RNA profiling of *P. vivax* liver stages

Address correspondence to dennis.kyle@uga.edu

## ABSTRACT

The resilience of *Plasmodium vivax*, the most widely distributed malaria-causing parasite in humans worldwide, is attributed to its ability to produce dormant liver forms known as hypnozoites, which can activate weeks, months, or even years after an initial mosquito bite. The factors underlying hypnozoite formation and activation are poorly understood, as is the parasite's influence on the host hepatocyte. Here, we shed light on transcriptome-wide signatures of both the parasite and the infected host cell by sequencing over 1,000 *P. vivax* liver forms at single-cell resolution. We distinguish between replicating schizonts and hypnozoites at the transcriptional level, identifying key differences in transcripts encoding for RNA-binding proteins associated with cell fate. In infected hepatocytes, we show that genes associated with energy metabolism and antioxidant stress response were upregulated, and those involved in the host immune response downregulated, suggesting both schizonts and hypnozoites alter the host intracellular environment. The transcriptional markers in schizonts, hypnozoites, and infected hepatocytes revealed here pinpoint potential factors underlying dormancy and can inform drug targets against *P. vivax* liver-stage infection.

## IMPORTANCE

In humans, malaria-causing *Plasmodium* parasites invade and replicate in the liver before mounting an infection in the blood. A few *Plasmodium* species, including *P. vivax*, can cause relapsing infections due to forms that persist in the liver known as hypnozoites. While interventions targeting the hypnozoite can prevent clinical malaria, little is known about their biology at the molecular level. Transcriptome-wide assessment of individual hypnozoites has not been accessible due to the technical challenges associated with liver stage infections. To overcome technical limitations, we performed single-cell transcriptional profiling of liver stages generated using an *in vitro* infection platform. In elucidating the transcriptional signatures of *P. vivax* liver forms, as well as hepatocytes in response to infection, our data provide new insights into these host-parasite interactions. An improved understanding of these interactions will allow for novel drug or vaccine interventions targeting either parasite- or hepatocyte-specific metabolic pathways essential for successful liver stage development and disease progression.

## INTRODUCTION

Malaria is a significant global health burden, with an estimated 421 million cases and 627,000 deaths per year (World Health Organization, 2021). At least five *Plasmodium* species are known to infect humans, with *P. vivax* being responsible for the majority of cases outside Africa (World Health Organization, 2021). All *Plasmodium* parasites that infect humans have an obligatory developmental stage in the liver wherein the parasite undergoes asexual reproduction known as schizogony within a

hepatocyte before releasing thousands of merozoites into the blood. In *P. vivax*, a proportion of parasites

known as hypnozoites forgo immediate division in hepatocytes, and persist in the liver for weeks,

months, or years before activating to initiate schizogony resulting in a relapse blood infection (Adams

and Mueller, 2017; Krotoski et al., 1982). Hypnozoites are thought to cause up to 90% of cases in certain

geographical regions (Adekunle et al., 2015; Commons et al., 2020; Robinson et al., 2015), and

mathematical models predict that eliminating *P. vivax* malaria will be difficult without interventions that

target hypnozoite reservoirs (White et al., 2018).

The factors influencing the biogenesis, persistence, and activation of hypnozoites are poorly

understood. Studies have found that hypnozoites in temperate regions have long latencies and that

tropical ones have short latencies, suggesting a correlation between geographical region and relapse

patterns ((Battle et al., 2014; Huldén et al., 2008; Krotoski et al., 1986; White, 2011). These findings

suggest that the developmental trajectory is genetically encoded, and support the hypothesis that

developmental fate is somehow pre-programmed in the sporozoite (Lysenko et al., 1977; Ungureanu et

al., 1976); but other contributing factors from the host could play a role in determining whether the

sporozoite becomes a schizont or hypnozoite (Schäfer et al., 2021).

Our understanding of *P. vivax* liver stage biology at the cellular and molecular level is in its

nascency for two overarching reasons. For a long time, protocols to cryopreserve and continuously

maintain erythrocytic stage parasites *in vitro* were lacking, which necessitated the collection of

gametocyte-infected blood from human volunteers in *P. vivax* malaria-endemic regions to generate the

sporozoites required to initiate an infection. New laboratory models have recently been developed which

can partially alleviate the need to obtain gametocytes from human infected blood  (Mikolajczak et al.,

2015; Schäfer et al., 2020). In addition to the sporozoite limitation, upon infecting the liver the high

ratio of noninfected versus infected hepatocytes has historically made it difficult to isolate and study

these forms in sufficient numbers. Robust *in vitro* pre-erythrocytic platforms capable of long-term

culture of sufficient numbers of *P. vivax* liver stages have only recently been developed (Maher et al.,

2021; Roth et al., 2018). Much of our knowledge of hypnozoite biology at the molecular level remains

limited to studies using *P. cynomolgi*—a parasite capable of causing relapsing infections non-human

primates, and for which molecular tools are available (Chua et al., 2019; Cubi et al., 2017; Voorberg-

van der Wel et al., 2017, 2020). However, it is unknown whether mechanisms regulating hypnozoite

formation and dormancy are consistent across different relapsing malaria parasites.

Transcriptome-wide profiling of *P. vivax* liver stages has previously been performed using a

bulk RNA sequencing strategy (Gural et al., 2018). While informative, the data only provide insights

into average transcript levels derived from either mixed (schizont and hypnozoite) or hypnozoite-only

populations, masking any transcriptional variation that might exist between individual parasites. Second,

the parasite's potential impact on the hepatocyte is not considered, barring analysis of host-pathogen

interactions. Nevertheless, these shortcomings highlight a need to better understand differences between

individual parasites, as well as how the parasite alters the hepatocyte. Advancements in these areas could

help elucidate parasite- and host- factors important for the parasite's liver stage development.

Single-cell RNA-sequencing (scRNA-seq) methods offer new ways of studying cell-to-cell

variation (Aldridge and Teichmann, 2020), and have already allowed for novel insights into host-

pathogen interactions in various disease contexts (Penaranda and Hung, 2019). In *Plasmodium* spp.,

scRNA-seq analyses have so far revealed that individual parasite forms exist on a spectrum

of transcriptomic states (Bogale et al., 2021; Howick et al., 2019; Poran et al., 2017; Real et al., 2021;

Reid et al., 2018; Ruberto et al., 2021a), and applying them to study the biology of *P. vivax* is a

burgeoning area of research (Mancio-Silva et al., 2021; Ruberto et al., 2021b; Sà et al., 2020). Droplet-

based techniques, such as 10x Genomics (Zheng et al., 2017), which use microfluidics, allow for the

sequencing of tens to hundreds of thousands of cells, increasing the chances of observing rare cell

populations (e.g., hepatocytes infected with *P. vivax*). We therefore presumed that applying single-cell

sequencing to *P. vivax* liver forms derived using an *in vitro* platform would prove a fruitful avenue of

inquiry into the parasite's biology cellular and molecular levels.

Using scRNA-seq, we characterize transcriptomic signatures of parasite and host in an *in vitro*

*P. vivax* liver stage model. We find differences in gene expression between replicating schizonts and

hypnozoites, and reveal variation, previously unobservable by bulk sequencing approaches, between

individual hypnozoites. In infected hepatocytes, we find that genes associated with energy metabolism

and antioxidant stress response are upregulated, and those involved in the host immune response

downregulated. Various host genes associated with these processes are found to be exclusively

upregulated during hypnozoite infection. Our study elucidates the transcriptional signatures among

infected hepatocytes and presents potential mechanisms *P. vivax* can use to either exploit its host for

rapid proliferation or to quietly persist, remaining undetected by the host immune system.

**RESULTS**

**Strategy used to study *P. vivax* liver stage biology**

Primary human hepatocytes were cultured and infected with *P. vivax* sporozoites on a 384-well plate

platform (Maher et al., 2021; Roth et al., 2018). **Figure 1A** outlines the workflow for the experiment.

Following infection with sporozoites, cultures were assessed at two endpoints: one containing both

schizonts and hypnozoites at day 5 post-infection, and the other containing solely hypnozoites at day 9

post-infection. Schizonts were eliminated from day 9 post-infection cultures by adding the

phosphatidylinositol 4-kinase (PI4K) inhibitor MMV390048 (1 µM) to the culture media on days 5, 6

and 7 post-infection (Gural et al., 2018; Paquet et al., 2017). For each collection time point, we

simultaneously initiated two sets of hepatocyte cultures: one infected with sporozoites for quantification

of immunofluorescent-labeled parasites by high-content imaging (HCI), and one infected with

sporozoites for transcriptional profiling using a high-throughput, droplet-based scRNA-seq workflow

(Zheng et al., 2017). As controls, a third set of naïve cultures were left uninfected and processed for

scRNA-seq alongside cultures at days 5 and day 9 post-infection. At each collection point, a target of

approximately 10,000 individual hepatocytes were partitioned into droplets prior to the generation of

RNA libraries. Each droplet contained primers with a unique cellular barcode that was ligated to

transcripts during the reverse transcription step, allowing for the identification of each transcript's

cellular origin post-sequencing. After sequencing the single-cell libraries, we mapped the resulting reads

onto the human (Ensembl, v103) and *P. vivax* PVP01 (PlasmoDB, v51) transcriptomes. We combined

the transcript information with the cellular barcodes to assess the number of transcripts in each cell, and

to determine whether a hepatocyte contained parasite material.

As expected, we detected hepatocytes infected with both forms of the parasite at 5 days post-

infection, and only hypnozoites at 9 days post-infection following exposure to MMV390048 (**Fig. 1B**

**and Fig. S1A**). **Figure 1C** displays the distribution of individual parasite growth areas, number of

parasites detected, and parasitemia levels at each endpoint. Across replicates, the parasitemia ranged

from 5.50-6.93% at 5 days post-infection and 1.76-2.77% at 9 days post-infection. Together, the high

infection rates coupled with the high-throughput capacity of 10x Genomics' single-cell isolation and

sequencing workflow allowed for sufficient material to be generated for successful profiling of both

parasite and host transcriptomes.

***P. vivax* schizonts and hypnozoites are distinguishable at single-cell resolution**

We first analyzed the reads mapping to the *P. vivax* transcriptome to identify schizonts and hypnozoites,

as well as to assess the extent to which these forms differed at single-cell resolution. In total, we

characterized 1,438 parasite transcriptomes across two independent experimental replicates (**Fig. S1B**).

The number of parasite transcriptomes profiled corroborated the parasitemia levels quantified using

high-content imaging (**Table S1**). Low dimensional representation of the data revealed two populations

of parasites in samples profiled at 5 days post-infection (**Fig. 2A, left panel**). In contrast, and as

expected, samples profiled at 9 days post-infection contained one population (**Fig. 2A, right panel**).

Given that the population of cells profiled at day 9 post-infection, from which schizonts were chemically

removed, overlapped with one of the two cell populations profiled at 5 days post-infection, we deduced

that the non-overlapping population of cells encoded for schizonts. Consistent with our reasoning, an

unsupervised clustering algorithm (Traag et al., 2019) unbiasedly assigned the cells to one of the two

clusters (**Fig. 2B and Fig. S1C**). As expected, number of transcripts detected, unique molecular
identifier (UMI) counts (a readout for absolute transcript abundances (Kivioja et al., 2011), and the
transcript levels of the early liver stage development marker liver stage-specific protein, LISP2
(PVP01_0304700) (Gupta et al., 2019), all displayed greater quantities in the schizont cluster (C2)
relative to the hypnozoite cluster (C1) (**Fig. 1C**). Together, these findings show that the transcriptomes
of *P. vivax* schizonts and hypnozoites can be distinguished using our experimental workflow.

**Hypnozoites exhibit distinct transcriptional signatures compared to schizonts**

We next performed differential gene expression analysis to identify transcriptome-wide differences
distinguishing schizonts and hypnozoites. Using stringent inclusion cutoffs (adjusted p value < 0.01 &
absolute (average $\log_2$ fold change [FC]) > 0.5), we found significant differences in ~18% (836/4722)
of all transcripts detected (**Fig. S2A; Data Set S1A**). Relative to hypnozoites, we found an increase in
the expression levels of 368 genes in schizonts. As expected, we observed a significant increase of
LISP2 in schizonts (average $\log_2$ FC: 2.32) compared to hypnozoites. Shortly after invading the
hepatocyte, non-relapsing *Plasmodium* spp. liver forms undergo immediate replication. Studies in
rodent malaria models have shown that these forms are metabolically active and that various genes
involved in energy metabolism are upregulated (Caldelari et al., 2019; Tarun et al., 2008; Toro-Moreno
et al., 2020). Consistent with previous data, we found an increased transcription associated with
glycolysis (PVP01_0816000, PVP01_1229700, PVP01_1244000) and the TCA cycle
(PVP01_0106100, PVP01_1332400) in schizonts (**Fig. S2B; Data Set S1A**). More broadly, we found
increased transcription of various metabolic processes, as well as motility, DNA-, RNA-, and protein-
processing (**Fig. S2C; Data Set S1A**). Transcripts encoding for various proteasome subunits displayed
greater transcription in schizonts (**Fig. S3A; Data Set S1A**). To further show the importance of this
protein complex, we tested a known proteosome inhibitor, Carfilzomib (Kortuem and Stewart, 2013),
in our standard 12-day *P. vivax* small molecule screening assay (Maher et al., 2021), and found it
effectively killed schizonts ($IC_{50}$ 288nM) and hypnozoites ($IC_{50}$ 511nM) with some selectivity over host
hepatocytes ($IC_{50}$ 3.32 µM) (**Figs. S3B and S3D**).

The detection of transcripts associated with energy metabolism in hypnozoites, albeit at a lower
level relative to schizonts, suggested that these forms are also metabolically active (**Fig. S2B**). Relative
to replicating schizonts, we found an increase in gene expression of 468 transcripts in hypnozoites (**Data
Set S1A**). As represented in **Figure 3A**, genes with greater expression in hypnozoites encoded for a
diverse set of ontologies—including proteases, membrane proteins, transcription factors, and DNA-
/RNA-regulating proteins. Genes encoding for proteases vivapain-1, vivapain-2 (PVP01_1248900,
PVP01_0916200), and plasmepsin IV (PVP01_1340900) displayed the greatest differential expression
in these forms compared to schizonts (**Figs. 3A and 3B; Data Set S1A**). While limited research has

reported the functionality of these gene products in *P. vivax* (Byoung-Kuk et al., 2004), work in other *Plasmodium* spp. has revealed that they localize to the food vacuole and play an important role in hemoglobin catabolism during the blood stages of the parasite's life cycle (Dame et al., 2003; Drew et al., 2008).

We also identified genes with greater expression associated with RNA regulatory mechanisms. Notably, we found greater expression of CCR4 (PVP01_1429000), a transcript encoding for a protein shown to have mRNA deadenylase activity and play a role in mRNA turnover (Tucker et al., 2001) (**Data Set S1A**). In *Plasmodium* spp., it has been described as important in regulating the expression of genes associated with invasion and egress in blood stages (Balu et al., 2011). RNA-binding proteins (RBPs) also govern gene expression at the post-transcriptional level by transiently storing transcripts until later processing (Hentze et al., 2018). Pumilio domain (PUF) proteins are a major family of RBPs that regulate cellular fating through translational repression in eukaryotes. We found that the transcript encoding for PUF1 (PVP01_1015200) was detected almost exclusively in hypnozoites (**Figs. 3A and 3B**; **Data Set S1A**). Furthermore, we found greater expression of 7-Helix-1 protein (PVP01_1008400) (**Figs. 3A and 3B**; **Data Set S1A**)—a stress granule component that interacts with the RNA-binding protein PUF2 and serves to be crucial for protein synthesis in *P. falciparum* (Miao et al., 2010). Numerous studies have shown that *Plasmodium* spp. use various post-transcriptional regulatory mechanisms to coordinate stage transitions and development (Baumgarten et al., 2019; Bunnik et al., 2016; Foth et al., 2011; le Roch et al., 2004; Tarun et al., 2008; Vembar et al., 2016). Our data corroborate these findings and suggest that similar strategies may be used in *P. vivax* to regulate liver stage development. Moreover, these data suggest that RBP-mediated cellular fating and translational repression may be involved in regulating hypnozoite dormancy.

Given the potential role in RBP-mediated cellular fating and translational repression governing latency in hypnozoites, we performed further analyses focused on RBP-mediated regulation. To this end, we scanned the *P. vivax* genome searching for genes encoding for cellular fating RBPs based on ortholog groups (OGs) of known human cellular fating RBPs (348 RBPs; 227 OGs), identifying 89 OGs (99 RBPs) in *P. vivax*. We detected 48 genes encoding for cellular fating RBPs with greater expression in schizonts (**Fig. 3C, left panel**; **Data Set S1B**). Notably, we identified eukaryotic translational initiation factors (PVP01_1238100, PVP01_0531200, PVP01_1431600, PVP01_0807600, PVP01_1005600, PVP01_0519700, PVP01_1329200, PVP01_0825000, PVP01_1303500), and splicing factors (PVP01_0716000, PVP01_0521000, PVP01_0804100, PVP01_1111800, PVP01_0607600, PVP01_1012200) suggesting active translational events. Gene Ontology (GO) term analysis revealed that these transcripts were associated with translation initiation (GO:0005852; GO:0033290; GO:0016282) (adjusted p value < 0.05, **Fig. 3C**; **Data Set S1C**).

In hypnozoites, we detected 51 genes encoding for cellular fating RBPs with increased expression relative to schizonts (**Fig. 3C, left panel**; **Data Set S1B**). These transcripts encode for prominent P body markers, namely DOZI/DDX6 (PVP01_0819400), CCR4-NOT complex proteins (PVP01_1429000, PVP01_1014400, PVP01_1331700, PVP01_1453400, PVP01_0929400), mRNA-decapping enzyme subunit 1 (PVP01_0617200), mRNA-decapping enzyme subunit 2 (PVP01_1409900), UPF1 (PVP01_0805200), UPF2 (PVP01_0724300), UPF3 (PVP01_1226700), PUF2 (PVP01_0526500), PABP (PVP01_1442500), Fibrillarin (PVP01_1341600) and trailer hitch homolog (PVP01_1269800) (**Data Set S1B**). Transcripts encoding for cellular fating RBPs in hypnozoite also represented GO terms corresponding to active biological condensate formations, such as the CCR4-NOT complex (GO:0030014; GO:0030015) and the P body (GO:0000932) (adjusted p value < 0.05, **Fig. 3C, right panel; Data Set S1C**). Together, these findings suggest that P body-mediated cell fating could be important in regulating hypnozoite persistence.

**Transcriptomic heterogeneity exists amongst individual *P. vivax* hypnozoites**

Having described the transcriptomic signatures distinguishing schizonts and hypnozoites, we next sought to assess the extent to which gene expression varied amongst hypnozoites. We thus parsed the data into only hypnozoite transcriptomes. In total, 1,147 transcriptomes were assessed (**Fig. 4A, left panel**). Low dimensional representation of the data revealed slight variation between samples collected at days 5 and 9 post-infection (**Fig. 4A, right panel**). After regrouping the data using the same unsupervised clustering algorithm described previously, we identified three clusters (**Fig. 4B, right panel**). The proportion of hypnozoites in each cluster varied across the day of collection: specifically, the proportion of hypnozoites in cluster 3 (C3) was greater in hypnozoites collected at day 9 compared to day 5 post-infection (**Fig. 4B, left panel; Fig. S4A**). Notably, we also found greater gene expression of LISP2 in C3 relative to the other two clusters (**Fig. 4C**).

We next identified markers that define the hypnozoites in each cluster using the FindAllMarkers function in Seurat (Stuart et al., 2019). We defined a marker as a transcript displaying greater than 0.5 FC (average $\log_2$) compared to the other clusters. In total, we detected 421 markers (adjusted p value < 0.05) (**Fig. S4B**; **Data Set S2A**). Notable markers in C1 included transcripts associated with drug resistance (PVP01_1447300, PVP01_1259100, and PVP01_1010900), energy metabolism (PVP01_1435000 and PVP01_1122800), protein modification (PVP01_1248900 and PVP01_0916200), and RNA-binding proteins (PVP01_0939900 and PVP01_0819400); in C2 those associated with translation (PVP01_1255500 and PVP01_0822500); and in C3 those associated with translation (PVP01_0213700 and PVP01_0203400), fatty acid metabolism (PVP01_1143000 and PVP01_1022800), protein processing (PVP01_1308000 and PVP01_0927700) and ion transport (PVP01_1407500) (**Fig. 4D**; **Data Set S2A**).

We next explored the data in the context of cellular fating to elucidate potential patterns in RBP

signatures. Among the hypnozoite markers found within our sub-clustering analysis, we identified 68

cellular fating RBPs, of which 54 RBPs were differentially expressed in at least in one of the three

clusters (adjusted p value < 0.05) (**Fig. 4E, top; Data Set S2B**). Approximately 4% (2/54) of the

differentially expressed genes in C3 encoded for cellular fating. In this cluster, the increased gene

expression of eukaryotic translation initiation factor 4E (PVP01_0203400) further suggests active

translation events may be occurring. C2 contained ~16% (9/54) of the cellular fating RBPs differentially

expressed. In this cluster, we found RBPs which are important determinants of cell proliferation from a

state of senescence such as eukaryotic translation initiation factor 5A (PVP01_1303500),

polypyrimidine tract binding protein (PVP01_1142400), GTP-binding nuclear protein RAN/TC4

(PVP01_0918300), transformer-2 protein homolog beta (PVP01_0802200). C1 contained the majority

of the differentially expressed RBPs (~80%, 43/54). Within this cluster, we identified prominent P body

markers with cellular fating ability, such as DOZI/DDX6 (PVP01_0819400), CCR4-NOT complex

proteins (PVP01_1429000, PVP01_1014400, PVP01_1331700, PVP01_1453400), UPF1

(PVP01_0805200), UPF2 (PVP01_0724300), PABP (PVP01_1442500), Fibrillarin (PVP01_1341600)

and trailer hitch homolog (PVP01_1269800). GO analysis revealed enrichment of cellular

compartments associated to biological condensate formations, namely CCR4-NOT complex

(GO:0030014), P granule (GO:0043186), and germplasm (GO:0060293) (**Fig. 4E; Data Set S2C**).

These findings, combined with the shift of hypnozoite proportions and increased transcript

levels of LISP2 from C1 to C3, allowed us infer the phenotype of hypnozoites in these clusters. As these

observations likely represent populations of hypnozoites in persisting and activating states, we defined

clusters C1, C2, and C3 to represent persisting, transitioning, and activating hypnozoites, respectively.

Together, these findings further our understanding of *P. vivax* hypnozoites through characterization of

gene usage and putative regulatory mechanisms in activating and persister forms. Furthermore, they

highlight a potential role of P body-mediated cellular fating governing persister hypnozoites.

**Strategy used to assess hepatocytes infected with *P. vivax* parasites**

After deciphering the identities of *P. vivax* transcriptomes representing schizonts and hypnozoites, we

next sought to characterize the transcriptional signatures of hepatocytes infected with either of these

forms. To this end, we integrated the data across all the conditions (**Figs. S5A and 5B**) and examined

whether distinct sub-populations of infected hepatocytes existed. As highlighted by the low dimensional

visualization of the data, we did not find a prominent pattern of infection (**Fig. 5**). Moreover, we detected

parasites in hepatocytes with different metabolic states (MacParland et al., 2018), as represented by their

transcriptional signatures associated with different hepatocyte zonation programs (**Figs. S5C and S5D**).

**Identification of transcriptional signatures in *P. vivax*-infected hepatocytes**

To identify host transcriptional signatures specific to infection status, we performed differential gene expression analysis, comparing infected to non-infected hepatocytes. Among the hepatocytes infected with schizonts, 106 genes displayed greater expression and 16 displayed decreased expression (**Data Set S3A**); while in hepatocytes infected with hypnozoites, 365 genes displayed greater expression and 41 genes displayed decreased expression (**Data Set S3B**) (adjusted p value < 0.05). As depicted in the volcano plots in **Fig. 6A**, the magnitude of change in expression in the infected versus non-infected cells (average $\log_2$ fold change) was greater in hepatocytes containing schizonts compared to those containing hypnozoites.

To gain greater insights into the biological processes and pathways that characterize a schizont- or hypnozoite-infected hepatocyte, we performed enrichment analysis (Raudvere et al., 2019)—scoring 18,594 gene sets against the genes differentially expressed in each of our transcriptional comparisons. When considering both the schizont- and hypnozoite-infected hepatocyte populations, we detected a total of 254 enriched gene sets (adjusted p value < 0.05) (**Fig. 6B**; **Data Set S3C**). Approximately 18% (45/254) of these gene sets overlapped in hepatocytes infected with either schizonts or hypnozoites (**Fig. S5E; Data Set S3D**). These data indicate that while some overlap exists in the gene networks the parasite alters during infection, the majority of processes and pathways altered are specific to schizont versus hypnozoite infected hepatocytes.

Using these gene sets, we then constructed network enrichment maps (Merico et al., 2010) to provide a global overview of the prominent biological themes associated with each infection condition (**Figs. 6C and 6D**). We detected upregulation of genes associated with pathways related to energy metabolism (oxidative phosphorylation, mitochondrial function, and lipid catabolism) in schizont- and hypnozoite-infected hepatocytes. Analysis of the most differentially expressed genes associated with these pathways included *MT-ND2*, *CHCHD10*, and *AKR1B10* (**Data Sets S3A and S3B**). Corroborating our transcriptomic data, we found an increase in protein levels of AKR1B10 in infected versus non-infected hepatocytes (**Figs. S6A-D**). AKR1B10 has been implicated in cell survival through its regulation of lipid metabolism and elimination of carbonyls (Martin and Maser, 2009; Qu et al., 2021; Wang et al., 2009) and has primarily been described as a biomarker of cancer (Díez-Dacal et al., 2011; Fukumoto et al., 2005; Jin et al., 2006). More recently, it has been shown that its expression is positively correlated with non-alcoholic fatty liver disease (Govaere et al., 2020). Its role during *Plasmodium* spp. liver stage infection warrants further research. In schizonts, the upregulation of genes linked to fatty acid metabolism is consistent with observations in hepatocytes harboring developing forms from non-relapsing *Plasmodium* spp. (Albuquerque et al., 2009; Itoe et al., 2014). We now reveal that infection

with hypnozoites also elicits enrichment of transcripts associated with fatty acid metabolism (**Fig. 6D**),
suggesting that these forms may be scavenging host resources to fuel their persistence.

Previous work assessing *Plasmodium* liver stage biology has revealed that the host is able to
detect the parasite (Liehl and Mota, 2012) and that type I interferon plays an integral role in this response
(Liehl et al., 2013; Miller et al., 2014). The decrease in transcripts associated with interferon signaling
(REAC:R-HSA-913531, GO:0034340, GO:0034341, GO:0071357, GO:0060337) in hepatocytes
infected with either schizonts or hypnozoites suggest that *P. vivax* manipulates the hepatocytes to evade
detection (**Figs. 6C and 6D; Data Sets S3C and S3D**). In hepatocytes infected with schizonts, we found
a decrease in expression of NFKB-regulated genes (*SOD2, TMEM45A, CCL20,* and *TPM1*) (**Data Set
S3A**), consistent with findings that *Plasmodium* spp. release signals to block proinflammatory response
(Singh et al., 2007). These findings, in addition to the decrease in the expression of *HLA-B*—a molecule
involved in antigen presentation to T cells that can trigger their activation when 'self' peptides are not
presented—and the neutrophil chemoattractant *CXCL8* suggest that replicating parasites alter
hepatocyte signaling pathways to minimize recruitment of innate immune cells and avoid detection by
T cells.

Similar to hepatocytes infected with schizonts, we found genes associated with interferon
signaling downregulated in hepatocytes infected with hypnozoites, namely: *UBB, UBC, HSP90AB1,*
*IFITM3, IFITM2,* and *HLA-A* (**Data Set S3B**). Furthermore, transcripts associated with the
downregulation of class major histocompatibility complex (*CD81* and *B2M*) were also decreased in
these hepatocytes (**Data Set S3B**). Together, these findings suggest that *P. vivax* infection of
hepatocytes may suppress the secretion of various chemokines and proinflammatory cytokines to thwart
the recruitment of inflammatory cells to the site of infection while simultaneously reducing the chances
they will be detected if these cells are recruited by downregulating MHC- I.

Enrichment analysis linking the changes in gene expression in infected hepatocytes with
transcriptional factors revealed distinct sets of putative transcriptional regulators. In total, we detected
35 transcription factors with potential roles in governing the gene expression changes in infected
hepatocytes (adjusted p value < 0. 05, **Fig. 7; Data Set S3E**). In infected hepatocytes, we found
enrichment of transcripts governed by NRF2 (**Fig. 7; Data Set S3E)**. The regulation of gene expression
through NRF2 in infected hepatocytes may serve to minimize reactive oxygen species in the cell (Hayes
and Dinkova-Kostova, 2014). Thus, in addition to the decreased expression of genes associated with
inflammatory response (**Data Set S3C**), NRF2-mediated regulation of antioxidant responsive genes may
protect hepatocytes from *Plasmodium*-induced stress and thus serve as a complementary means of
minimizing the host inflammatory response.

**DISCUSSION**

The infection of the liver by *P. vivax* constitutes an obligatory step in its life-cycle. *In vivo* studies have shown that within hepatocytes, *P. vivax* can exist in one of two phenotypically distinct forms: schizonts or hypnozoites (Cogswell, 1992; Mikolajczak et al., 2015). They have also been observed *ex vivo* by infection of hepatocyte monolayers (Gural et al., 2018; Hollingdale et al., 1985; Roth et al., 2018; Sattabongkot et al., 2006). A key problem hindering the study of *P. vivax* liver stage biology is the difficulty involved in obtaining a sufficient number of infected hepatocytes to make meaningful comparisons between them. We show that by coupling our *in vitro* liver stage model (Maher et al., 2021; Roth et al., 2018) with a high throughput single-cell capture technology (Klein et al., 2015; Macosko et al., 2015; Zheng et al., 2017) transcriptome-wide characterization of a large number of infected hepatocytes is feasible. Our approach therefore provides much needed insight into the molecular signatures of *P. vivax* hypnozoites and the host cells they infect.

First, we found clear transcriptional differences between schizonts and hypnozoites. Our transcriptome-wide assessment supports the notion that hypnozoites are metabolically active (Adams and Mueller, 2017), demonstrated by the transcription associated with glycolysis and fatty acid metabolism. Second, we found that hypnozoites exist in various transcriptomic states. We speculate that these differences might represent a spectrum of phenotypes, from persisting to activating. The differential expression of genes encoding for cellular fating RBPs supports the notion that these forms may use post-transcriptional mechanisms to control gene usage (Mueller et al., 2019). The biological significance of the enrichment of transcripts associated with translational repression, found exclusively in a subpopulation of hypnozoites, suggests that this mechanism inhibits activation, and warrants further investigation.

Second, we found distinct transcriptomic signatures of infected versus non-infected hepatocytes, and showed that signaling pathways associated with energy metabolism, antioxidant stress, and immune response are altered in infected hepatocytes. We found no clear pattern of infection, as both schizonts and hypnozoites are detected in hepatocytes with various metabolic signatures. The extent to which individual hepatocytes respond to *P. vivax* schizonts and hypnozoites has previously been unknown. A bulk RNA sequencing approach has been used to identify pathways altered in hepatocytes in response to *Plasmodium* infection (LaMonte et al., 2019); however, it is limited insofar as it averages gene expression from millions of cells, obscuring any parasite-specific effects on the infected cell. Targeted approaches have revealed that *P. vivax* liver forms recruit host factors to facilitate growth (LaMonte et al., 2019; Posfai et al., 2020), but how certain infected hepatocyte populations contribute to disease transmission, and how hepatocyte subpopulations support development, are intriguing topics for future studies (Vijayan et al., 2021). Our data identify potential host factors essential for *P. vivax*'s

384     successful liver stage development. Whether certain hepatocyte phenotypes influence the development,

385     especially activation of hypnozoites into schizonts, requires further investigation.

386           Several observations suggest protein degradation may be important for the parasite's liver

stages. Targeting this process may be an effective means of killing these forms. The ability of

hepatocytes to uptake and break down hemoglobin (Bissell et al., 1972) supports that this protein

substrate could be an essential amino acid source for the parasite. Our data suggest that several

mechanisms may allow *P. vivax* to utilize the amino acids derived from this protein. In hypnozoites, the

increased expression of genes encoding for proteases, such as vivapains, supports the idea that these

parasites digest host cytosolic proteins**.** The role of these proteins as hemoglobinases have been defined

extensively in *P. falciparum* blood stages (Drew et al., 2008; Rosenthal, 2011; Salas et al., 1995). A

similar role has been described for *P. vivax* (Byoung-Kuk et al., 2004), but there is limited information

about their significance in the liver stages. Vivapains are highly homologous to cathepsin protease L

(TgCpl) in the related apicomplexan *Toxoplasma gondii.* In these parasites, TgCpl is localized to its

plant-like vacuolar compartment—a lysosome-like organelle containing a variety of proteases (Parussini

et al., 2010)—and one of its functions is to aid in the digestion of host-derived proteins (Dou et al.,

2014). Genetic or chemical disruption of TgCpl negatively impacts the viability of bradyzoites, a

persisting form of this parasite (di Cristina et al., 2017). Given these orthologous proteins' roles in *P.*

*falciparum* and *T. gondii*, we speculate that the expression of vivapains may support the persistence of

*P. vivax* hypnozoites.

403           Persistence of hypnozoites is contingent on the longevity of the hepatocyte in which it resides.

The increased expression of genes associated with antioxidant stress (Scarpulla, 2002) is consistent with

the notion that hypnozoite infection induces changes in the hepatocyte to promote cell survival, in which

NRF-2 could be playing a regulatory role. We detected genes displaying greater expression in infected

hepatocytes related to NYFB-regulation, a transcription factor important for cell proliferation,

mitochondrial integrity, and cellular longevity (Tharyan et al., 2020). The increased expression of

mitochondria-related genes in hepatocytes infected with schizonts may be required to meet the parasite's

energy demands during replication, but the enrichment of mitochondria-related transcripts in cells

infected with hypnozoites could also aid in ensuring longevity of the hepatocyte. In addition to these

positive regulators of cell survival, we found negative regulation of transcripts through the decreased

expression of genes associated with cell death (*UBB, UBC,* and *HSP90AB1*). Their expression is thought

to be regulated by CEBPD, a transcription factor that has a role in governing cell death (Tsai et al., 2017;

Wang et al., 2019).

416           Persistence of hypnozoites also requires evading host detection, but how they are able to do so

417     has been elusive. Our analyses indicate that the parasite alters the host hepatocyte to evade detection by

immune cells. First, there is a decrease in expression of genes encoding for chemoattractants (*CCL20, CXCL8, IL32*) in infected hepatocytes, and an increase in expression of PTGR1, a negative regulator of the chemotactic factor leukotriene B4 (Yokomizo et al., 1993). These changes would result in decreased secretion of chemokines that recruit innate immune cells to the liver, thereby reducing recruitment of adaptive immune cells like CD8 T-cells that kill infected hepatocytes (Chakravarty et al., 2007). Interestingly, there was also a decrease in expression of MHC-1 molecule (*HLA-A and -B*). The presentation of parasite peptides to be detected by surveying CD8 T-cells would therefore be reduced (Neefjes et al., 2011), creating an additional layer of protection should innate immune cells begin to be recruited, for example upon lysis of the hepatocyte by schizonts. We propose that *P. vivax* infected hepatocytes are re-programmed to prevent recruitment of innate immune cells on the one hand, and to reduce the likelihood of their detection by the adaptive immune system on the other, all to enable progress through schizogony and for hypnozoites to persist undetected. Understanding the mechanisms leading up to this should be studied in the future, as their reversal may lead to improved clearance of hypnozoites, thus reducing relapses. Indeed, a vaccine would need to decrease relapses to have any impact on *P. vivax* transmission and disease (White et al., 2017). However, the decrease of MHC-I expression in infected hepatocytes may mean that standard focuses to generate CD8 and CD4 polyfunctional T-cells in the liver may not be effective in eliminating *P. vivax* or other relapsing parasites. Recent attempts to test anti-relapse vaccine candidates with *P. cynomolgi* did not prevent relapses despite higher inducing IFNγ producing cells, a common metric used as a correlate of protection (Kim et al., 2020), indicating that alternative strategies may be needed.

Our transcriptome-wide characterization of schizonts, hypnozoites, and the infected hepatocytes highlights transcripts altered during infection, and provides a framework for examining the role of infection dynamics of *P. vivax* liver stage infections. By examining two early liver stage time points at single-cell resolution, our study lays out proof of principle and a technical advance demonstrating the importance of *in vitro* models to shed light on the molecular basis of *P. vivax* liver stage development and infection.

**CONCLUSIONS**

The overall aim of our research was to elucidate the molecular signatures of *P. vivax* liver infection, a better understanding of which will aid in facilitating the development of new therapeutics to ultimately reduce transmission. We anticipate that interventions could involve parasite- and/or host- directed small molecule therapies, two approaches our group is actively pursuing. The comprehensive mapping of the transcriptional landscape of both the host and *P. vivax* described here provides an important framework for further investigation.

## MATERIALS AND METHODS

### Ethics statement

All research procedures were reviewed and approved by the Cambodian National Ethics Committee for Health Research (approval numbers: #113NECHR, #104NECHR and #089NECHR).

### Blood samples, mosquitos, and infections

The complete protocol used for mosquito rearing an infection has been published (Maher et al., 2021). In brief, blood samples from symptomatic patients infected with *P. vivax* were collected at local health facilities in Mondulkiri province in Cambodia from March 2020 to October 2021. Following a *P. vivax* gametocyte-containing blood meal, *An. dirus* mosquitoes were maintained at ambient temperature on a 12:12 light: dark cycle and fed 10% sucrose + 0.5% PABA solution. *An. dirus* found positive for *P. vivax* oocysts at six-days post-feeding were transported to the Institut Pasteur Cambodia Insectary Facility in Phnom Penh, Cambodia where they were maintained under the same conditions described above.

### Sporozoite isolation and primary human hepatocyte infections

The complete protocol used for seeding primary human hepatocytes, harvesting sporozoites, and infecting cultures has been published (Maher et al., 2021). In brief, *P. vivax* sporozoites were isolated from the salivary glands of female *An. dirus* mosquitoes 18 days after an infectious blood-meal. Primary human hepatocytes (Lot BGW, BioIVT) were seeded 2 days before infection with 30,000 sporozoites (replicate 1), or 15,200 sporozoites (replicate 2) per well. Media was replaced with fresh CP media (BioIVT) containing antibiotics (penicillin, streptomycin, and gentamycin) the day after infection and every 2-3 days thereafter. Hepatocytes containing solely hypnozoites at 9 days post-infection were treated with 1 μM MMV390048 on days 5, 6, and 7 post-infection to kill replicating schizonts. At days 5 and 9 post-infection, cultures were processed for single-cell sorting (described in the following section) or immunofluorescence assay (IFA). For the latter, hepatocytes were fixed with 4% paraformaldehyde in 1X PBS. Fixed cultures were then stained overnight at 4ºC with recombinant mouse anti-*P. vivax* UIS4 antibody (Schäfer et al., 2018) diluted 1:25,000 in a stain buffer (0.03% Triton X-100) and 1% (w/v) BSA in 1X PBS. The following day, cultures were washed three times with 1X PBS and then stained overnight at 4ºC with 1:1,000 rabbit anti-mouse AlexaFluor ™488-conjugated antibody diluted 1:1,000 in stain buffer. Cultures were washed three times with 1X PBS and counterstained with 1 mg/mL Hoechst 33342 (Thermo Fisher) to detect parasite and host DNA. High content imaging was performed on a Lionheart imaging system (Biotek). Quantification of the number

482   of nuclei and parasite liver forms was performed using Gen5 high content analysis software (Biotek).

483   Confocal images were obtained using an ImageXpress Micro (Molecular Devices).

**Single-cell partitioning, library preparation, and sequencing**

Hepatocyte cultures infected with *P. vivax* were processed prior to- and post- PI4K treatment (5- and 9-

days post-infection, respectively). Two replicate experiments, defined as an independent plating of

hepatocytes infected with sporozoites from a unique field isolate, were performed several months apart.

Additionally, and as a negative control, one replicate of hepatocyte cultures never exposed to *P. vivax*

(naïve) were processed similarly to those that were infected. At each endpoint for each replicate,

approximately 640,000 PHH hepatocytes (16 wells of a 384-well plate each having approximately

10,000 cells) were treated with trypsin (Corning 25-053-Cl) to facilitate their detachment from the plate.

Once detached, trypsin was inactivated with 1:1 complete media and the cell suspension was collected

into a 5mL protein-low bind centrifuge tube (Eppendorf). To remove residual trypsin, hepatocytes were

washed via three sequential rounds of pelleting (50 x g, 3 min, 4ºC) and resuspending in 1X PBS

containing 0.1% BSA. Cells were then passed through a 35µm strainer (Falcon 352235) to minimize or

remove hepatocyte clusters. Prior to partitioning cells on the Chromium controller (10x Genomics),

hepatocytes were counted and the cell concentration was assessed using a hemocytometer.

Approximately 10,000 hepatocytes were loaded on a Chromium Chip B (10x Genomics) as per

manufacturer's recommendations. Chips containing hepatocyte suspensions, Gel Beads in Emulsion

(GEMs), and reverse transcription reagents were loaded into the Chromium controller (10x Genomics)

for single-cell partitioning and cell barcoding. Barcoded cDNA libraries were generated according to

the Chromium Single Cell 3' gene expression protocol (version 3). In total, 6 cDNA libraries were

generated (two replicates at 5 days post-infection, two replicates at 9 days post-infection, one uninfected

control for day 5 cultures, and one uninfected control for day 9 cultures). Libraries were loaded on

individual flow cell lanes and sequenced at a depth of 400 million reads using a HiSeq X Ten platform

(Illumina) at Macrogen (Seoul, Korea).

**Computational analysis of single-cell RNA sequencing data**

Quality control checks on the raw sequencing data were performed using FASTQC (Andrews, 2010). A

multi-species transcriptome index—containing the transcriptomes for *P. vivax* P01 (PlasmoDB.org,

version 51) and *H. sapiens* GRCh8 (Ensembl version 103)—was created using the 'ref' function in kb-

python (0.26.3) (Bray et al., 2016; Melsted et al., 2021). Next, reads pseudoaligned and counted using

the 'count' function in kb-python with default parameters selected. The resulting unfiltered cell-gene

count matrices were read into R (v 4.1.1) for further processing.

**Filtering and normalization of scRNA-seq count matrices**

R scripts containing all computational analyses will be available on a dedicated GitHub repository upon publication.

To distinguish between droplets containing cells and those only with ambient RNA, we used emptyDrops function (Lun et al., 2019) with the lower total UMI count threshold (at or below which all barcodes are assumed to correspond to empty droplets) set to 1,000. The droplets with cells (FDR < 0.001), more specifically, those containing barcoded transcripts derived from either human, parasite, or a combination of the two—were kept for downstream processing.

**Assessment of *P. vivax* liver stage transcriptomes**

For the analyses of *P. vivax*, we filtered out the transcripts derived from human, leaving only transcripts derived from the parasite quantifiable. For each sample, cells with less than 200 UMIs were removed. Next, we filtered out low detected genes, retaining transcripts with 10 or greater UMIs. Post cell and gene filtering, the data were normalized using Seurat (Stuart et al., 2019) 'LogNormalize' function with the default parameters selected.

*Data integration and clustering*

The data were integrated using Seurat's SCTransform function (Hafemeister and Satija, 2019) with the following parameters indicated: variable.features.n = NULL, variable.features.rv.th = 1.3. Clustering was performed using Seurat's SNN-Leiden algorithm (Traag et al., 2019). The full description of this analysis is found in the extended methods.

*Differential gene expression analysis*

To detect differentially expressed transcripts, the Seurat function FinalAllMarkers and FindMarkers were used. For differential gene expression testing using the FindMarkers function, we used the following parameters: logfc.threshold = 0.0, ident.1 = '1', ident.2 = '2', test.use = "wilcox", min.pct = 0, only.pos = F, assay = "RNA". For differential gene expression testing using the FindAllMarkers function, we used the following parameters: test.use = "wilcox", only.pos = T, assay = "RNA", log2FC = 0.0.

*RBP analyses*

RBPs with cell fating potential reported in humans (Decker and Parker; Gerstberger et al., 2014; Guallar and Wang, 2014; Voronina et al., 2011; Youn et al., 2019) were manually curated. The

orthologue's counterparts of these cell fating RBPs from the *P. vivax* PVP01 genome were mined using

OrthoMCL (Release-6.8)(Chen et al., 2006). Within the differentially expressed genes in *P. vivax*,

transcripts encoding for cell fating RBPs were filtered for further assessment.

*GO enrichment analyses*

GO enrichment analysis was performed on the PlasmoDB (v55) website with the following

input parameters selected: P value cutoff = 0.05; Limit to GO Slim terms = No; Evidence: Computed,

curated. Enrichment from GO cellular compartment (CC), biological processes (BP), and molecular

function (MF) were all assessed.

**Assessment of host hepatocyte transcriptomes**

For the analyses of the host, we removed all transcripts derived from *P. vivax*, leaving only the

transcripts derived from hepatocytes. We retained transcripts detected in at least 30 cells. Next, cells

with less than 1,500 UMIs were removed from further analyses. Post-cell and gene filtering, the data

from each replicate were normalized using Seurat's 'LogNormalize' function with the default

parameters.

*Data integration and clustering*

Filtered, normalized data matrices were integrated in a manner described in in the Seurat

(version 4) vignette, *Introduction to scRNA-seq integration*, available on the Sajita Lab's website

(https://satijalab.org/seurat/articles/integration_introduction.html). Briefly, integration features were

identified in each replicate using the 'SelectIntegrationFeatures' function. Next, integration anchors

were identified using the 'FindIntegrationAnchors'. Last, using these anchors, the four datasets were

integrated using the 'IntegrateData' function. Clustering was performed using Seurat's SNN-Leiden

algorithm (Traag et al., 2019).

*Gene Ontology and Reactome enrichment analyses*

Enriched biological processes and pathways were determined using gProfiler (Raudvere et al.,

2019) with the following options selected: ordered query, statistical domain scope—only annotated

genes, significance threshold—g:SCS threshold, user threshold—0.05, Gene Ontology—GO biological

processes, No electronic GO annotations, biological pathways—Reactome. Transcripts that displayed

significant changes in expression in infected hepatocytes versus non-infected hepatocytes were used as

input. EnrichmentMap (v 1.1.0) (Merico et al., 2010) for CytoScape (v 3.8.2) (Shannon et al., 2003) was

used to create the enrichment networks. Significantly enriched GO biological processes and Reactome

pathways (adjusted p value < 0.05) were used as input. Pathways were clustered and annotated using

the AutoAnnotate function (Kucera et al., 2016). Node size corresponds to the number of transcripts in

the corresponding set, and edge widths correspond to the number of transcripts shared between sets.

*Transcription factor enrichment analysis*

We downloaded curated sets of known transcription factors from the website

(http://amp.pharm.mssm.edu/Enrichr/). Transcription factor—target gene sets were obtained by ChIP-

X experiments from the ChEA (Lachmann et al., 2010) and ENCODE (Project Consortium, 2004)

projects. TF-target relationships using Chip-seq data from ENCODE and CHEA were used to identify

overrepresented pathways and TFs. Differentially expressed genes (adjusted p value < 0.05) in each

condition versus non-infected cells were used as inputs. Gene sets with an adjusted p value < 0.05 were

considered significantly enriched.

**Carfilzomib dose-response assessment**

A carfilzomib dose-response against *P. vivax* liver forms was obtained from three independent

experiments, in which a different *P. vivax* clinical isolate was allowed to infect BGW human

hepatocytes. The 12-day assay was used (Maher et al., 2021). All independent experiments resulted in

sufficient hypnozoite formation within each well for dose-response calculations. Liver stage parasite

growth metrics and compound dilutions were loaded into CDD Vault (Collaborative Drug Discovery)

and growth data normalized such that zero (negative) inhibition was set as the average of DMSO wells

and 100% (positive) inhibition was set to the effective doses of a nigericin control. Normalized values

were then combined in Graphpad Prism to fit a dose-response curve and calculated $IC_{50}$ values from all

replicates.

**Quantification of AKR1B10 protein expression**

To analyze AKR1B10 expression and localization, *P. vivax*-infected PHHs were fixed 12 days post-

infection. Infected wells were then permeabilized for 20 minutes with 0.2% TritonX-100, washed thrice

with 1X PBS, and blocked in 3% BSA for one hour. Primary antibodies rabbit anti-HsAKR1B10 (PA5-

30773, Thermo Fisher) and mouse anti-PvUIS4 were sequentially added, diluted 1:100 and 1:500,

respectively. Antibodies were incubated overnight at 4ºC and washed 3 times with 1X PBS before

addition of 1:400 Alexa- Fluor 568 donkey anti-rabbit for HsAKR1B10 or 1:400 AlexaFluor 488 goat

anti-mouse for PvUIS4. Secondary antibodies were incubated overnight at 4ºC. Cells were washed thrice

with 1X PBS before adding 0.5 mg/mL DAPI for 10 minutes. Cells were washed 3 final times with 1X

PBS before image collection with an ImageXpress inverted confocal microscope (Molecular Devices).
Image analysis was completed using MetaXpress analysis software (Molecular Devices).

**Contact for reagent and resource sharing**

606    Further information and requests for resources and reagents should be directed to and will be fulfilled
607    by the Lead Contact, Dennis Kyle (email: dennis.kyle@uga.edu).

**Data availability**

609    Raw sequencing files (FASTQ), processed data, scripts, and all host- and parasite- differentially
610    expressed genes will be made publicly available upon publication of the manuscript.

**Acknowledgements**

We thank the patients of Mondulkiri Province, Cambodia, for participating in this study. We thank the
Institut Pasteur du Cambodge's field site manager (Saorin Kim) for logistical assistance, insectary staff
(Makara Pring, Koeun Kaing, Nora Sambath) for the *An. dirus* mosquito colony maintenance, and
laboratory staff (Eakpor Piv, Chansophea Chhin, Sreyvouch Phen, Chansovandan Chhun, Sivcheng
Phal, Baura Tat) for assistance with the mosquito dissections and the *in vitro* assays.

High-content imaging data was produced in part by the Biomedical Microscopy Core at the
University of Georgia, supported by the Georgia Research Alliance. A.R.J., B.B. and I.M. acknowledge
support from the Victorian State Government Operational Infrastructure Support and Australian
Government National Health and Medical Research Council Independent Research Institute
Infrastructure Support Scheme. This work was also supported by the Agence Nationale de la Recherche
(ANR-17-CE13-0025 to A.A.R and I.M.); an Australian National Health and Medical Research Council
Leadership Fellowship (APP1194330 to A.R.J.); the Bill & Melinda Gates Foundation (OPP1023601
to D.E.K.); and Medicines for Malaria Venture (RD/2017/0042 to B.W. and A.V., RD/16/1082 and
RD/15/0022 to S.P.M. and D.E.K.).

A.A.R., S.P.M., A.V., I.M., B.W. and D.E.K conceived and designed the study; A.A.R, S.P.M.
and A.V. performed the experiments; A.A.R., C.B., B.B. and C.J. curated the data; A.A.R., S.P.M., B.B.,
A.R.J. and C.J. performed the formal analyses; S.P.M., A.V. and I.M., B.W. and D.E.K. coordinated the
research activities; S.P.M., A.V., I.M., B.W. and  D.E.K. provided resources; S.P.M. and D.E.K.
acquired funding for the work; A.A.R wrote the initial draft of the article; all authors reviewed and made
contributions to the submitted article.

**REFERENCES**

Adams, J.H., and Mueller, I. (2017). The Biology of Plasmodium vivax. Cold Spring Harbor Perspectives in Medicine *7*, a025585.

Adekunle, A.I., Pinkevych, M., McGready, R., Luxemburger, C., White, L.J., Nosten, F., Cromer, D., and Davenport, M.P. (2015). Modeling the Dynamics of Plasmodium vivax Infection and Hypnozoite Reactivation In Vivo. PLOS Neglected Tropical Diseases *9*, e0003595.

Albuquerque, S.S., Carret, C., Grosso, A.R., Tarun, A.S., Peng, X., Kappe, S.H.I., Prudêncio, M., and Mota, M.M. (2009). Host cell transcriptional profiling during malaria liver stage infection reveals a coordinated and sequential set of biological events. BMC Genomics *10*, 1–13.

Aldridge, S., and Teichmann, S.A. (2020). Single cell transcriptomics comes of age. Nature Communications 2020 11:1 *11*, 1–4.

Andrews, S. (2010). FastQC: A quality control tool for high throughput sequence data. Available at: Http://Www.Bioinformatics.Babraham.Ac.Uk/Projects/Fastqc.

Balu, B., Maher, S.P., Pance, A., Chauhan, C., Naumov, A. v., Andrews, R.M., Ellis, P.D., Khan, S.M., Lin, J. wen, Janse, C.J., et al. (2011). CCR4-Associated Factor 1 Coordinates the Expression of Plasmodium falciparum Egress and Invasion Proteins. Eukaryotic Cell *10*, 1257.

Battle, K.E., Karhunen, M.S., Bhatt, S., Gething, P.W., Howes, R.E., Golding, N., van Boeckel, T.P., Messina, J.P., Shanks, G.D., Smith, D.L., et al. (2014). Geographical variation in Plasmodium vivax relapse. Malaria Journal *13*, 1–16.

Baumgarten, S., Bryant, J.M., Sinha, A., Reyser, T., Preiser, P.R., Dedon, P.C., and Scherf, A. (2019). Transcriptome-wide dynamics of extensive m6A mRNA methylation during Plasmodium falciparum blood-stage development. Nature Microbiology *4*, 2246.

Bissell, D.M., Hammaker, L., and Schmid, R. (1972). Hemoglobin and Erythrocyte Catabolism in Rat Liver: The Separate Roles of Parenchymal and Sinusoidal Cells. Blood *40*, 812–822.

Bogale, H.N., Pascini, T. v., Kanatani, S., Sá, J.M., Wellems, T.E., Sinnis, P., Vega-Rodríguez, J., and Serre, D. (2021). Transcriptional heterogeneity and tightly regulated changes in gene expression during

658 *Plasmodium berghei* sporozoite development. Proceedings of the National Academy of Sciences *118*,
659 e2023438118.

660 Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq
661 quantification. Nature Biotechnology *34*, 525–527.

662 Bunnik, E.M., Batugedara, G., Saraf, A., Prudhomme, J., Florens, L., and le Roch, K.G. (2016). The
663 mRNA-bound proteome of the human malaria parasite Plasmodium falciparum. Genome Biology *17*.

664 Byoung-Kuk, N.A., Shenai, B.R., Sijwali, P.S., Choe, Y., Pandey, K.C., Singh, A., Craik, C.S., and
665 Rosenthal, P.J. (2004). Identification and biochemical characterization of vivapains, cysteine proteases
666 of the malaria parasite Plasmodium vivax. Biochemical Journal *378*, 529.

667 Caldelari, R., Dogga, S., Schmid, M.W., Franke-Fayard, B., Janse, C.J., Soldati-Favre, D., and Heussler,
668 V. (2019). Transcriptome analysis of Plasmodium berghei during exo-erythrocytic development.
669 Malaria Journal *18*, 330.

670 Chakravarty, S., Cockburn, I.A., Kuk, S., Overstreet, M.G., Sacci, J.B., and Zavala, F. (2007). CD8+ T
671 lymphocytes protective against malaria liver stages are primed in skin-draining lymph nodes. Nature
672 Medicine 2007 13:9 *13*, 1035–1041.

673 Chen, F., Mackey, A.J., Stoeckert, C.J., and Roos, D.S. (2006). OrthoMCL-DB: querying a
674 comprehensive multi-species collection of ortholog groups. Nucleic Acids Research *34*.

675 Chua, A.C.Y., Jie, J., Ong, Y., Malleret, B., Suwanarusk, R., Kosaisavee, V., Zeeman, A.-M., Cooper,
676 C.A., Tan, K.S.W., Zhang, R., et al. (2019). Robust continuous in vitro culture of the Plasmodium
677 cynomolgi erythrocytic stages. Nature Communications *10*, 1–13.

678 Cogswell, F.B. (1992). The hypnozoite and relapse in primate malaria. Clinical Microbiology Reviews
679 *5*, 26–35.

680 Commons, R.J., Simpson, J.A., Watson, J., White, N.J., and Price, R.N. (2020). Estimating the
681 Proportion of Plasmodium vivax Recurrences Caused by Relapse: A Systematic Review and Meta-
682 Analysis. The American Journal of Tropical Medicine and Hygiene *103*, 1094–1099.

di Cristina, M., Dou, Z., Lunghi, M., Kannan, G., Huynh, M.-H., McGovern, O.L., Schultz, T.L.,
Schultz, A.J., Miller, A.J., Hayes, B.M., et al. (2017). Toxoplasma depends on lysosomal consumption
of autophagosomes for persistent infection.

Cubi, R., Vembar, S.S., Biton, A., Franetich, J.-F., Bordessoulles, M., Sossau, D., Zanghi, G., Bosson-
Vanga, H., Benard, M., Moreno, A., et al. (2017). Laser capture microdissection enables transcriptomic
analysis of dividing and quiescent liver stages of Plasmodium relapsing species. Cellular Microbiology
*19*, e12735.

Dame, J.B., Yowell, C.A., Omara-Opyene, L., Carlton, J.M., Cooper, R.A., and Li, T. (2003).
Plasmepsin 4, the food vacuole aspartic proteinase found in all Plasmodium spp. infecting man.
Molecular and Biochemical Parasitology *130*, 1–12.

Decker, C.J., and Parker, R. P-Bodies and Stress Granules: Possible Roles in the Control of Translation
and mRNA Degradation.

Díez-Dacal, B., Gayarre, J., Gharbi, S., Timms, J.F., Coderch, C., Gago, F., and Pérez-Sala, D. (2011).
Identification of aldo-keto reductase AKR1B10 as a selective target for modification and inhibition by
prostaglandin A(1): implications for antitumoral activity. Cancer Research *71*, 4161–4171.

Dou, C., Mcgovern, Z.L., Cristina, D. di, and Carruthers, M.B. (2014). Toxoplasma gondii ingests and
digests host cytosolic proteins. MBio *5*, 1188–1202.

Drew, M.E., Banerjee, R., Uffman, E.W., Gilbertson, S., Rosenthal, P.J., and Goldberg, D.E. (2008).
Plasmodium Food Vacuole Plasmepsins Are Activated by Falcipains. The Journal of Biological
Chemistry *283*, 12870.

Foth, B.J., Zhang, N., Chaal, B.K., Sze, S.K., Preiser, P.R., and Bozdech, Z. (2011). Quantitative time-
course profiling of parasite and host cell proteins in the human malaria parasite Plasmodium falciparum.
Molecular and Cellular Proteomics *10*.

Fukumoto, S.I., Yamauchi, N., Moriguchi, H., Hippo, Y., Watanabe, A., Shibahara, J., Taniguchi, H.,
Ishikawa, S., Ito, H., Yamamoto, S., et al. (2005). Overexpression of the aldo-keto reductase family
protein AKR1B10 is highly correlated with smokers' non-small cell lung carcinomas. Clinical Cancer
Research : An Official Journal of the American Association for Cancer Research *11*, 1776–1785.

Gerstberger, S., Hafner, M., and Tuschl, T. (2014). A census of human RNA-binding proteins. Nature
Reviews Genetics 2014 15:12 *15*, 829–845.

Govaere, O., Cockell, S., Tiniakos, D., Queen, R., Younes, R., Vacca, M., Alexander, L., Ravaioli, F.,
Palmer, J., Petta, S., et al. (2020). Transcriptomic profiling across the nonalcoholic fatty liver disease
spectrum reveals gene signatures for steatohepatitis and fibrosis. Science Translational Medicine *12*.

Guallar, D., and Wang, J. (2014). RNA-binding proteins in pluripotency, differentiation, and
reprogramming. Frontiers in Biology *9*, 389–409.

Gupta, D.K., Dembele, L., Voorberg-Van Der Wel, A., Roma, G., Yip, A., Chuenchob, V.,
Kangwanrangsan, N., Ishino, T., Vaughan, A.M., Kappe, S.H., et al. (2019). The plasmodium liver-
specific protein 2 (LISP2) is an early marker of liver stage development. ELife *8*.

Gural, N., Mancio-Silva, L., Miller, A.B., Galstian, A., Butty, V.L., Levine, S.S., Patrapuvich, R., Desai,
S.P., Mikolajczak, S.A., Kappe, S.H.I., et al. (2018). In Vitro Culture, Drug Sensitivity, and
Transcriptome of Plasmodium Vivax Hypnozoites. Cell Host & Microbe *23*, 395-406.e4.

Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. Genome Biology *20*, 1–15.

Hayes, J.D., and Dinkova-Kostova, A.T. (2014). The Nrf2 regulatory network provides an interface
between redox and intermediary metabolism. Trends in Biochemical Sciences *39*, 199–218.

Hentze, M.W., Castello, A., Schwarzl, T., and Preiss, T. (2018). A brave new world of RNA-binding
proteins. Nature Reviews Molecular Cell Biology 2018 19:5 *19*, 327–341.

Hollingdale, M.R., Collins, W.E., Campbell, C., and Schwartz, A.L. (1985). In vitro culture of two
populations (dividing and nondividing) of exoerythrocytic parasites of Plasmodium vivax. The
American Journal of Tropical Medicine and Hygiene *34*, 216–222.

Howick, V.M., Russell, A.J.C., Andrews, T., Heaton, H., Reid, A.J., Natarajan, K., Butungi, H., Metcalf,
T., Verzier, L.H., Rayner, J.C., et al. (2019). The Malaria Cell Atlas: Single parasite transcriptomes
across the complete Plasmodium life cycle. Science (New York, N.Y.) *365*, eaaw2619.

Huldén, L., Huldén, L., and Heliövaara, K. (2008). Natural relapses in vivax malaria induced by
Anopheles mosquitoes. Malaria Journal *7*, 1–11.

Itoe, M.A., Sampaio, J.L., Cabal, G.G., Real, E., Zuzarte-Luis, V., March, S., Bhatia, S.N., Frischknecht,

F., Thiele, C., Shevchenko, A., et al. (2014). Host Cell Phosphatidylcholine Is a Key Mediator of Malaria

Parasite Survival during Liver Stage Infection. Cell Host & Microbe *16*, 778.

Jin, J., Krishack, P., and Cao, D. (2006). Role of aldo-keto reductases in development of prostate and

breast cancer. Frontiers in Bioscience : A Journal and Virtual Library *11*, 2767–2773.

Kim, Y.C., Dema, B., Rodriguez-Garcia, R., López-Camacho, C., Leoratti, F.M.S., Lall, A., Remarque,

E.J., Kocken, C.H.M., and Reyes-Sandoval, A. (2020). Evaluation of Chimpanzee Adenovirus and

MVA Expressing TRAP and CSP from Plasmodium cynomolgi to Prevent Malaria Relapse in

Nonhuman Primates. Vaccines *8*, 1–14.

Kivioja, T., Vähärautio, A., Karlsson, K., Bonke, M., Enge, M., Linnarsson, S., and Taipale, J. (2011).

Counting absolute numbers of molecules using unique molecular identifiers. Nature Methods *9*, 72–74.

Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., Veres, A., Li, V., Peshkin, L., Weitz, D.A.,

and Kirschner, M.W. (2015). Droplet barcoding for single-cell transcriptomics applied to embryonic

stem cells. Cell *161*, 1187–1201.

Kortuem, K.M., and Stewart, A.K. (2013). Carfilzomib. Blood *121*, 893–897.

Krotoski, W.A., Collins, W.E., Bray, R.S., Garnham, P.C., Cogswell, F.B., Gwadz, R.W., Killick-

Kendrick, R., Wolf, R., Sinden, R., Koontz, L.C., et al. (1982). Demonstration of hypnozoites in

sporozoite-transmitted Plasmodium vivax infection. The American Journal of Tropical Medicine and

Hygiene *31*, 1291–1293.

Krotoski, W.A., Garnham, P.C.C., Cogswell, F.B., Collins, W.E., Bray, R.S., Gwasz, R.W., Killick-

Kendrick, R., Wolf, R.H., Sinden, R., and Hollingdale, M. (1986). Observations on early and late post-

sporozoite tissue stages in primate malaria. IV. Pre-erythrocytic schizonts and/or hypnozoites of

Chesson and North Korean strains of Plasmodium vivax in the chimpanzee. American Journal of

Tropical Medicine and Hygiene *35*, 263–274.

Kucera, M., Isserlin, R., Arkhangorodsky, A., and Bader, G.D. (2016). AutoAnnotate: A Cytoscape app

for summarizing networks with semantic annotations. F1000Research *5*.

Lachmann, A., Xu, H., Krishnan, J., Berger, S.I., Mazloom, A.R., and Ma'ayan, A. (2010). ChEA:
transcription factor regulation inferred from integrating genome-wide ChIP-X experiments.
Bioinformatics (Oxford, England) *26*, 2438–2444.

LaMonte, G.M., Orjuela-Sanchez, P., Calla, J., Wang, L.T., Li, S., Swann, J., Cowell, A.N., Zou, B.Y.,
Abdel-Haleem Mohamed, A.M., Villa Galarce, Z.H., et al. (2019). Dual RNA-seq identifies human
mucosal immunity protein Mucin-13 as a hallmark of Plasmodium exoerythrocytic infection. Nature
Communications *10*, 488.

Liehl, P., and Mota, M.M. (2012). Innate recognition of malarial parasites by mammalian hosts.
International Journal for Parasitology *42*, 557–566.

Liehl, P., Zuzarte-Luís, V., Chan, J., Zillinger, T., Baptista, F., Carapau, D., Konert, M., Hanson, K.K.,
Carret, C., Lassnig, C., et al. (2013). Host-cell sensors for Plasmodium activate innate immunity against
liver-stage infection. Nature Medicine 2013 20:1 *20*, 47–53.

Lun, A.T.L., Riesenfeld, S., Andrews, T., Dao, T.P., Gomes, T., and Marioni, J.C. (2019). EmptyDrops:
distinguishing cells from empty droplets in droplet-based single-cell RNA sequencing data. Genome
Biology *20*, 63.

Lysenko Ja., A., Beljaev, A.E., and Rybalka, V.M. (1977). Population studies of Plasmodium vivax: 1.
The theory of polymorphism of sporozoites and epidemiological phenomena of tertian malaria. Bulletin
of the World Health Organization *55*, 541.

Macosko, E.Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I., Bialas, A.R.,
Kamitaki, N., Marterstteck, E.M., et al. (2015). Highly Parallel Genome-wide Expression Profiling of
Individual Cells Using Nanoliter Droplets. Cell *161*, 1202–1214.

MacParland, S.A., Liu, J.C., Ma, X.-Z., Innes, B.T., Bartczak, A.M., Gage, B.K., Manuel, J., Khuu, N.,
Echeverri, J., Linares, I., et al. (2018). Single cell RNA sequencing of human liver reveals distinct
intrahepatic macrophage populations. Nature Communications *9*, 4383.

Maher, S., Vantaux, A., Cooper, C., Chasen, N., Cheng, W., Joyner, C., Manetsch, R., Witkowski, B.,
and Kyle, D. (2021). A Phenotypic Screen for the Liver Stages of Plasmodium vivax. BIO-PROTOCOL
*11*.

Mancio-Silva, L., Gural, N., Wadsworth, M.H., Butty, V.L., Hughes, T.K., March, S., Nerurkar, N.,
Roobsoong, W., Fleming, H.E., Whittaker, C., et al. (2021). Gene signatures and host-parasite
interactions revealed by dual single-cell profiling of Plasmodium vivax liver infection. BioRxiv
2021.09.01.454555.

Martin, H.J., and Maser, E. (2009). Role of human aldo–keto-reductase AKR1B10 in the protection
against toxic aldehydes. Chemico-Biological Interactions *178*, 145–150.

Melsted, P., Booeshaghi, A.S., Liu, L., Gao, F., Lu, L., Min, K.H. (Joseph), da Veiga Beltrame, E.,
Hjörleifsson, K.E., Gehring, J., and Pachter, L. (2021). Modular, efficient and constant-memory single-
cell RNA-seq preprocessing. Nature Biotechnology 2021 39:7 *39*, 813–818.

Merico, D., Isserlin, R., Stueker, O., Emili, A., and Bader, G.D. (2010). Enrichment Map: A Network-
Based Method for Gene-Set Enrichment Visualization and Interpretation. PLOS ONE *5*, e13984.

Miao, J., Li, J., Fan, Q., Li, X., Li, X., and Cui, L. (2010). The Puf-family RNA-binding protein PfPuf2
regulates sexual development and sex differentiation in the malaria parasite Plasmodium falciparum.
Journal of Cell Science *123*, 1039.

Mikolajczak, S.A., Vaughan, A.M., Kangwanrangsan, N., Roobsoong, W., Fishbaugher, M.,
Yimamnuaychok, N., Rezakhani, N., Lakshmanan, V., Singh, N., Kaushansky, A., et al. (2015).
Plasmodium vivax liver stage development and hypnozoite persistence in human liver-chimeric mice.
Cell Host & Microbe *17*, 526–535.

Miller, J.L., Sack, B.K., Baldwin, M., Vaughan, A.M., and Kappe, S.H.I. (2014). Interferon-Mediated
Innate Immune Responses against Malaria Parasite Liver Stages. Cell Reports *7*, 436–447.

Mueller, I., Jex, A.R., Kappe, S.H.I., Mikolajczak, S.A., Sattabongkot, J., Patrapuvich, R., Lindner, S.,
Flannery, E.L., Koepfli, C., Ansell, B., et al. (2019). Transcriptome and histone epigenome of
Plasmodium vivax salivary-gland sporozoites point to tight regulatory control and mechanisms for liver-
stage differentiation in relapsing malaria. International Journal for Parasitology *49*, 501–513.

Neefjes, J., Jongsma, M.L.M., Paul, P., and Bakke, O. (2011). Towards a systems understanding of
MHC class I and MHC class II antigen presentation. Nature Reviews Immunology 2011 11:12 *11*, 823–
836.

Paquet, T., le Manach, C., Cabrera, D.G., Younis, Y., Henrich, P.P., Abraham, T.S., Lee, M.C.S., Basak,

R., Ghidelli-Disse, S., Lafuente-Monasterio, M.J., et al. (2017). Antimalarial efficacy of MMV390048,

an inhibitor of Plasmodium phosphatidylinositol 4-kinase. Science Translational Medicine *9*.

Parussini, F., Coppens, I., Shah, P.P., Diamond, S.L., and Carruthers, V.B. (2010). Cathepsin L

Occupies a Vacuolar Compartment and is a Protein Maturase within the Endo/Exocytic System of

Toxoplasma gondii. Molecular Microbiology *76*, 1340.

Penaranda, C., and Hung, D.T. (2019). Single-Cell RNA Sequencing to Understand Host–Pathogen

Interactions. ACS Infectious Diseases *5*, 336–344.

Poran, A., Nötzel, C., Aly, O., Mencia-Trinchant, N., Harris, C.T., Guzman, M.L., Hassane, D.C.,

Elemento, O., and Kafsack, B.F.C. (2017). Single-cell RNA sequencing reveals a signature of sexual

commitment in malaria parasites. Nature *551*, 95–99.

Posfai, D., Maher, S.P., Roesch, C., Vantaux, A., Sylvester, K., Péneau, J., Popovici, J., Kyle, D.E.,

Witkowski, B., and Derbyshire, E.R. (2020). Plasmodium vivax Liver and Blood Stages Recruit the

Druggable Host Membrane Channel Aquaporin-3. Cell Chemical Biology.

Project Consortium, E. (2004). The ENCODE (ENCyclopedia of DNA Elements) Project. Science *306*,

636–640.

Qu, J., Li, J., Zhang, Y., He, R., Liu, X., Gong, K., Duan, L., Luo, W., Hu, Z., Wang, G., et al. (2021).

AKR1B10 promotes breast cancer cell proliferation and migration via the PI3K/AKT/NF-κB signaling

pathway. Cell & Bioscience *11*, 163.

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J. (2019). g:Profiler:

a web server for functional enrichment analysis and conversions of gene lists (2019 update). Nucleic

Acids Research *47*, W191–W198.

Real, E., Howick, V.M., Dahalan, F.A., Witmer, K., Cudini, J., Andradi-Brown, C., Blight, J., Davidson,

M.S., Dogga, S.K., Reid, A.J., et al. (2021). A single-cell atlas of Plasmodium falciparum transmission

through the mosquito. Nature Communications 2021 12:1 *12*, 1–13.

Reid, A.J., Talman, A.M., Bennett, H.M., Gomes, A.R., Sanders, M.J., Illingworth, C.J.R., Billker, O.,

Berriman, M., and Lawniczak, M.K.N. (2018). Single-cell RNA-seq reveals hidden transcriptional

variation in malaria parasites. ELife *7*, 1–29.

Robinson, L.J., Wampfler, R., Betuela, I., Karl, S., White, M.T., Li Wai Suen, C.S.N., Hofmann, N.E.,
Kinboro, B., Waltmann, A., Brewster, J., et al. (2015). Strategies for understanding and reducing the
Plasmodium vivax and Plasmodium ovale hypnozoite reservoir in Papua New Guinean children: a
randomised placebo-controlled trial and mathematical model. PLoS Medicine *12*.

le Roch, K.G., Johnson, J.R., Florens, L., Zhou, Y., Santrosyan, A., Grainger, M., Yan, S.F., Williamson,
K.C., Holder, A.A., Carucci, D.J., et al. (2004). Global analysis of transcript and protein levels across
the Plasmodium falciparum life cycle. Genome Research *14*, 2308–2318.

Rosenthal, P.J. (2011). Falcipains and Other Cysteine Proteases of Malaria Parasites. Advances in
Experimental Medicine and Biology *712*, 30–48.

Roth, A., Maher, S.P., Conway, A.J., Ubalee, R., Chaumeau, V., Andolina, C., Kaba, S.A., Vantaux, A.,
Bakowski, M.A., Thomson-Luque, R., et al. (2018). A comprehensive model for assessment of liver
stage therapies targeting Plasmodium vivax and Plasmodium falciparum. Nature Communications *9*,
1837.

Ruberto, A.A., Bourke, C., Merienne, N., Obadia, T., Amino, R., and Mueller, I. (2021a). Single-cell
RNA sequencing reveals developmental heterogeneity among Plasmodium berghei sporozoites.
Scientific Reports *11*, 4127.

Ruberto, A.A., Bourke, C., Vantaux, A., Maher, S.P., Jex, A., Witkowski, B., Snounou, G., and Mueller,
I. (2021b). Single-cell RNA sequencing of Plasmodium vivax sporozoites reveals stage- and species-
specific transcriptomic signatures. BioRxiv 2021.11.24.469176.

Sà, J.M., Cannon, M. v., Caleon, R.L., Wellems, T.E., and Serre, D. (2020). Single-cell transcription
analysis of Plasmodium vivax blood-stage parasites identifies stage- and species-specific profiles of
expression. PLOS Biology *18*, e3000711.

Salas, F., Fichmann, J., Lee, G.K., Scott, M.D., and Rosenthal, P.J. (1995). Functional expression of
falcipain, a Plasmodium falciparum cysteine proteinase, supports its role as a malarial hemoglobinase.
Infection and Immunity *63*, 2120.

Sattabongkot, J., Yimamnuaychoke, N., Leelaudomlipi, S., Rasameesoraj, M., Jenwithisuk, R.,
Coleman, R.E., Udomsangpetch, R., Cui, L., and Brewer, T.G. (2006). Establishment of a human
hepatocyte line that supports in vitro development of the exo-erythrocytic stages of the malaria parasites
Plasmodium falciparum and P. vivax. Am. J. Trop. Med. Hyg *74*, 708–715.

Scarpulla, R.C. (2002). Transcriptional activators and coactivators in the nuclear control of
mitochondrial function in mammalian cells. Gene *286*, 81–89.

Schäfer, C., Dambrauskas, N., Steel, R.W., Carbonetti, S., Chuenchob, V., Flannery, E.L., Vigdorovich,
V., Oliver, B.G., Roobsoong, W., Maher, S.P., et al. (2018). A recombinant antibody against
Plasmodium vivax UIS4 for distinguishing replicating from dormant liver stages. Malaria Journal *17*,
370.

Schäfer, C., Roobsoong, W., Sattabongkot, J., Mikolajczak, S.A., and Kappe, S.H.I. (2020). A
Humanized Mouse Model for Plasmodium vivax to Test Interventions that Block Liver Stage to Blood
Stage Transition and Blood Stage Infection. IScience *23*, 101381.

Schäfer, C., Zanghi, G., Vaughan, A.M., and Kappe, S.H.I. (2021). Plasmodium vivax Latent Liver
Stage Infection and Relapse: Biological Insights and New Experimental Tools. Annual Review of
Microbiology *75*.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B.,
and Ideker, T. (2003). Cytoscape: A Software Environment for Integrated Models of Biomolecular
Interaction Networks. Genome Research *13*, 2498–2504.

Singh, A.P., Buscaglia, C.A., Wang, Q., Levay, A., Nussenzweig, D.R., Walker, J.R., Winzeler, E.A.,
Fujii, H., Fontoura, B.M.A., and Nussenzweig, V. (2007). Plasmodium Circumsporozoite Protein
Promotes the Development of the Liver Stages of the Parasite. Cell *131*, 492–504.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y., Stoeckius, M.,
Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*, 1888-
1902.e21.

Tarun, A.S., Peng, X., Dumpit, R.F., Ogata, Y., Silva-Rivera, H., Camargo, N., Daly, T.M., Bergman,
L.W., and Kappe, S.H.I. (2008). A combined transcriptome and proteome survey of malaria parasite
liver stages. Proceedings of the National Academy of Sciences *105*, 305–310.

Tharyan, R.G., Annibal, A., Schiffer, I., Laboy, R., Atanassov, I., Weber, A.L., Gerisch, B., and Antebi,
A. (2020). NFYB-1 regulates mitochondrial function and longevity via lysosomal prosaposin. Nature
Metabolism *2*, 387–396.

Toro-Moreno, M., Sylvester, K., Srivastava, T., Posfai, D., and Derbyshire, E.R. (2020). RNA-Seq
Analysis Illuminates the Early Stages of Plasmodium Liver Infection.

Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden: guaranteeing well-
connected communities. Scientific Reports *9*, 1–12.

Tsai, H.H., Lai, H.Y., Chen, Y.C., Li, C.F., Huang, H.S., Liu, H.S., Tsai, Y.S., and Wang, J.M. (2017).
Metformin promotes apoptosis in hepatocellular carcinoma through the CEBPD-induced autophagy
pathway. Oncotarget *8*, 13832.

Tucker, M., Valencia-Sanchez, M.A., Staples, R.R., Chen, J., Denis, C.L., and Parker, R. (2001). The
transcription factor associated Ccr4 and Caf1 proteins are components of the major cytoplasmic mRNA
deadenylase in Saccharomyces cerevisiae. Cell *104*, 377–386.

Ungureanu, E., Killick-Kendrick, R., Garnham, P.C.C., Branzei, P., Romanescu, C., and Shute, P.G.
(1976). Prepatent periods of a tropical strain of Plasmodium vivax after inoculations of tenfold dilutions
of sporozoites. Transactions of the Royal Society of Tropical Medicine and Hygiene *70*, 482–483.

Vembar, S.S., Droll, D., and Scherf, A. (2016). Translational regulation in blood stages of the malaria
parasite Plasmodium spp.: systems-wide studies pave the way. Wiley Interdisciplinary Reviews. RNA
*7*, 772.

Vijayan, K., Wei, L., Glennon, E.K.K., Mattocks, C., Bourgeois, N., Staker, B., and Kaushansky, A.
(2021). Host-targeted Interventions as an Exciting Opportunity to Combat Malaria. Chemical Reviews
*121*, 10452–10468.

Voorberg-van der Wel, A., Roma, G., Gupta, D.K., Schuierer, S., Nigsch, F., Carbone, W., Zeeman,
A.M., Lee, B.H., Hofman, S.O., Faber, B.W., et al. (2017). A comparative transcriptomic analysis of
replicating and dormant liver stages of the relapsing malaria parasite plasmodium cynomolgi. ELife *6*.

Voorberg-van der Wel, A.M., Zeeman, A.M., Nieuwenhuis, I.G., van der Werff, N.M., Klooster, E.J.,
Klop, O., Vermaat, L.C., Kumar Gupta, D., Dembele, L., Diagana, T.T., et al. (2020). A dual fluorescent
Plasmodium cynomolgi reporter line reveals in vitro malaria hypnozoite reactivation. Communications
Biology *3*.

Voronina, E., Seydoux, G., Sassone-Corsi, P., and Nagamori, I. (2011). RNA Granules in Germ Cells.
Cold Spring Harbor Perspectives in Biology *3*.

Wang, C., Yan, R., Luo, D., Watabe, K., Liao, D.-F., and Cao, D. (2009). Aldo-keto Reductase Family
1 Member B10 Promotes Cell Survival by Regulating Lipid Synthesis and Eliminating Carbonyls. The
Journal of Biological Chemistry *284*, 26742–26748.

Wang, S.M., Lin, H.Y., Chen, Y.L., Hsu, T.I., Chuang, J.Y., Kao, T.J., and Ko, C.Y. (2019).
CCAAT/enhancer-binding protein delta regulates the stemness of glioma stem-like cells through
activating PDGFA expression upon inflammatory stimulation. Journal of Neuroinflammation *16*, 1–12.

White, N.J. (2011). Determinants of relapse periodicity in Plasmodium vivax malaria. Malaria Journal
*10*, 297.

White, M., Amino, R., and Mueller, I. (2017). Theoretical Implications of a Pre-Erythrocytic
Plasmodium vivax Vaccine for Preventing Relapses. Trends in Parasitology *33*, 260.

White, M.T., Walker, P., Karl, S., Hetzel, M.W., Freeman, T., Waltmann, A., Laman, M., Robinson,
L.J., Ghani, A., and Mueller, I. (2018). Mathematical modelling of the impact of expanding levels of
malaria control interventions on Plasmodium vivax. Nature Communications 2018 9:1 *9*, 1–10.

World Health Organization (2021). World malaria report 2021.

Yokomizo, T., Izumi, T., Takahashi, T., Kasamal, T., Kobayashis, Y., Satos, F., Taketanill, Y., and
Shimizu, T. (1993). Enzymatic Inactivation the Porcine Kidney O f Leukotriene B4 by a Novel Enzyme
Found in PURIFICATION AND PROPERTIES OF LEUKOTRIENE Bq 12-
HYDROXYDEHYDROGENASE*. THE JOURNAL OF BIOLOGICAL CHEMISTRY *268*, 18128–
18135.

Youn, J.Y., Dyakov, B.J.A., Zhang, J., Knight, J.D.R., Vernon, R.M., Forman-Kay, J.D., and Gingras,
A.C. (2019). Properties of Stress Granule and P-Body Proteomes. Molecular Cell *76*, 286–294.

Zheng, G.X.Y., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R., Ziraldo, S.B., Wheeler,
T.D., McDermott, G.P., Zhu, J., et al. (2017). Massively parallel digital transcriptional profiling of single
cells. Nature Communications *8*, 14049.

## SUPPLEMENTARY MATERIAL

**Supplementary Table S1.** Comparison of metrics derived from scRNA-seq and high content imaging data analyses.

**Figure S1. Assessment of *P. vivax* liver stages using high-content imaging and scRNA-seq.** (A) Representative high content images of primary hepatocytes infected with *P. vivax* on day 5 (left) and day 9 (right) post-infection. Images were obtained from one well from a 384-well plate with a 4x objective. Inset: one field of view (orange box) from the same well captured with a 20x objective. Cells were stained with DAPI (blue) and *Pv*UIS4 (green). White arrow: liver form assigned as a hypnozoite, yellow arrow: liver form assigned as a schizont. White bar represents 1mm, grey bar represents 200μm. Images are representative of replicate 2. (B) Number of *P. vivax* liver stage transcriptomes assessed post-cell and gene filtering. (C) Number of *P. vivax* liver stage parasites assigned to cluster 1 and cluster 2 (hypnozoites and schizonts, respectively).

**Figure S2. *P. vivax* schizonts and hypnozoites have distinct transcriptomic signatures. (A)** Scatterplot showing average $\log_2$ fold change versus the difference in the proportion of cells the transcript is detected in clusters encoding for hypnozoites and schizonts. Negative values: greater detection in hypnozoites; positive values: greater detection in schizonts. (B) Violin plots displaying the expression of genes encoding for TCA- and glycolysis-related proteins. (C) Dot plots showing transcripts with decreased expression in hypnozoites relative to schizonts. The size of the dot corresponds to the percentage of cells expressing the gene, colored by average expression. Differentially expressed transcripts were identified using Seurat's FindMarkers function. Wilcoxon rank-sum test, adjusted p values < 0.01. Scale: normalized expression; pct. exp: percent of cells expressing the gene. ldh: L-lactate dehydrogenase; gapdh: glyceraldehyde 3-phosphate dehydrogenase; atp-synthC: ATP synthase subunit C; gdh1: NADP-specific glutamate dehydrogenase 1.

**Figure S3. Carfilzomib inhibits *P. vivax* schizonts and hypnozoites.** (A) Violin plots displaying the distribution of expression of select genes encoding for proteasome subunits in schizonts and hypnozoites. (B) Table of proteasome-associated transcripts differentially expressed between schizonts and hypnozoites. Positive average log2FC: higher expression in schizonts. (C) Structure of carfilzomib. (D) Dose-response curve of carfilzomib-induced inhibition of *P. vivax* schizont growth area (yellow, IC50 288nM), hypnozoite quantity per well (blue, IC50 511nM), and hepatocyte nuclei quantity (orange, IC50 3.32 μM). Data shown are pooled from three independent experiments, each containing duplicate wells at each concentration. Bars represent ± SEM.

**Figure S4. Assessment of hypnozoite transcriptomes.** (A) UMAP of hypnozoites colored by cluster and faceted by day post-infection. (B) Violin plot showing the average $\log_2$FC of differentially expressed genes in each cluster. Markers were identified using Seurat's AllFindMarkers function. Transcripts with an adjusted p value < 0.05 (Wilcoxon rank-sum) and average $\log_2$FC > 0.5 were deemed markers. Red dashed line: average $\log_2$FC = 0.5.

**Figure S5. Analysis of host-pathogen transcriptional signatures in *P. vivax*- infected and non-infected hepatocytes.** (A) Number of hepatocytes assessed. Colored by infection status and faceted by sample (left); proportion of infected, exposed, or naive hepatocytes assessed on day 5 or day 9 post-infection. (B) t-SNE plot of hepatocytes colored by day. (C) t-SNE plots of hepatocytes colored by percent expression of zonation markers in human hepatocytes. Markers obtained from Macparland et al. (2019). (D) t-SNE plots of hepatocytes colored by hypnozoite infection status. Grey, uninfected (naive and exposed) hepatocytes; red, hepatocytes infected with C1 hypnozoites; blue, hepatocytes infected with C2; teal, hepatocytes infected with C3 hypnozoites. (E) Unique and overlapping gene sets in hepatocytes infected with schizonts or hypnozoites. Positive enrichment derived from genes displaying greater expression versus non-infected hepatocytes; negative enrichment derived from transcripts displaying decreased expression versus noninfected hepatocytes.

**Figure S6. Human AKR1B10 is upregulated in hepatocytes infected with *P. vivax*.** (A) Representative confocal images of *P. vivax* parasites on day 12 post-infection of hepatocytes. Cells were stained with DAPI (blue), *Pv*UIS4 (green), and HsAKR1B10 (red). White bar represents 100μm. (B) Violin plots displaying the distribution of *AKR1B10* transcript levels in infected (schizonts and hypnozoites) and non-infected (exposed and naive) hepatocytes. Black dot: mean expression ($\log_2$). (C) Violin plot displaying the distribution of AKR1B10 signal in hepatocyte cultures infected with *P. vivax*. (D) Associated with panel C, violins plots displaying the distribution of AKR1B10 signal across biological replicates. IHC: immunohistochemistry; SZ: hepatocyte containing schizont; HZ: hepatocyte containing hypnozoite; EX: hepatocyte not infected but exposed to *P. vivax*. ****: $p <= 0.0001$; *: $p <= 0.05$; n.s: not significant, $p > 0.05$.

**Data Set S1.** Data associated with schizonts versus hypnozoites analyses. (A) Differentially expressed genes defining schizonts and hypnozoites. Positive Average log 2 FC: Upregulated in hypnozoites relative to schizonts. (B) Differentially expressed genes identified as encoding for cellular fating RNA-binding proteins. (C) GO Term enrichment of Hypnozoite or Schizont transcripts identified as cellular fating RBPs. Rows highlighted in light green are terms with Bonferroni adjusted p value < 0.05. Input parameters: P value cutoff = 0.05; Limit to GO Slim terms = No; Evidence: Computed, curated; PlasmoDB v55.

**Data Set S2.** Data associated with analyses of hypnozoite-to-hypnozoite heterogeneity. (A) Markers

defining each hypnozoite cluster. Marker defined as: Average log 2 FC > 0.5, Adjusted p value < 0.05.

Rows colored green are transcripts that meet marker criteria. (B) Differentially expressed genes

identified as encoding for cellular fating RNA-binding proteins in hypnozoite clusters. (C) GO Term

enrichment of hypnozoite transcripts identified as cellular fating RBPs. Rows highlighted in light green

are terms with Bonferroni adjusted p value < 0.05. P value cutoff = 0.05; Limit to GO Slim terms = No;

Evidence: Computed, curated; PlasmoDB v55.

**Data Set S3.** Data associated with host-specific transcriptional signatures in response to *P. vivax*

infection. (A) Differentially expressed genes in hepatocytes containing schizonts versus non-infected

cells. Positive average log2FC: greater expression in infected hepatocytes. (B) Differentially expressed

genes in hepatocytes containing hypnozoites versus non-infected cells. Positive average log2FC: greater

expression in infected hepatocytes. (C) Enrichment of GO biological processes and Reactome metabolic

pathways. (D) Unique and overlapping enriched gene sets in hepatocytes infected with schizonts or

hypnozoites. (E) Transcription factor enrichment analysis. Rows highlighted in green are sets that are

significantly enriched, adjusted p value < 0.05.

**Figure 1. Design and validation of detection of *P. vivax* liver stages using high-content imaging and scRNA-seq.** (A) Schematic illustrating the sample preparation and data generation pipeline used to assess *P. vivax* liver stage infection rates and gene expression. (B) Representative high content images of primary hepatocytes infected with *P. vivax* on day 5 (left) and day 9 (right) post-infection (p.i). Images were captured from one well from a 384-well plate with a 4x objective. Inset: one field of view (white dashed box) from the same well captured with a 20x objective. Cells were stained with DAPI (blue) and *Pv*UIS4 (green). White bar represents 1mm, inset bars represent 10μm. Images are representative of replicate 1. (C) Histogram displaying the distribution of parasite growth areas fixed at day 5 or day 9 post-infection (left) and net quantity of parasites and hepatocyte nuclei (right). Net infection rate is indicated for each endpoint. Growth and infection metrics are shown for both replicates.

**Figure 2. Low dimensional representation of *P. vivax* schizont and hypnozoite transcriptomes** (A) t-SNE plots of integrated *P. vivax* liver stage scRNA-seq data faceted by parasites isolated on day 5 (left) and day 9 (right) post-infection. (B) t-SNE plot of *P. vivax* liver stage data colored by parasites encoding for putative hypnozoites (C1) and schizonts (C2). (C) t-SNE plots of *P. vivax* liver stage scRNA-seq data colored by number of transcripts detected (left), UMI counts (middle), and the expression of the schizont marker, LISP2 (PVP01_0304700) (right). Scales: absolute count (left, middle), normalized expression (right).

**Figure 3. *P. vivax* hypnozoites have distinct transcriptomic signatures**. (A) Dot plots showing transcripts that distinguish all hypnozoites from schizonts. The size of the dot corresponds to the percentage of cells that transcript was detected per condition, colored by average expression. Differentially expressed transcripts were identified using Seurat's FindMarker function (Wilcoxon rank-sum test, adjusted p values < 0.01). (B) tSNE plots of liver stages colored by transcripts displaying greater levels in hypnozoites. Colored boxes in A and B indicate the family or canonical biological process that the transcript is associated with. Scale: normalized expression; pct. exp: percent of cells expressing transcript. (C) Number of transcripts in schizonts and hypnozoites identified as cellular fating RBPs (left) and GO term enrichment of this subset. Adj P: Bonferroni adjusted p value.

**Figure 4. Transcriptomic signatures of reactivating and persister hypnozoites.** (A) Number of hypnozoites assessed (left) and UMAP of cells identified as hypnozoites colored by day of collection. (B) Proportion of cells in clusters on days 5 and 9 post-infection (left) and UMAP of hypnozoites colored by cluster. (C) Violin plot showing the expression of LISP2 in each cluster. (D) Dot plot showing markers that distinguish hypnozoites in each cluster. The size of the dot corresponds the percentage of cells in the expressing the gene of interest, colored by scaled normalized expression of the transcript in each cluster. Marker transcripts were identified using Seurat's AllFindMarkers function. Wilcoxon rank-sum test, average log2FC > 1, adjusted p value < 0.01. (E) Number transcripts in each cluster encoding for cellular fating RBPs sorted by their average log2FC relative to the other clusters (top) and GO Term enrichment of transcripts in each cluster (bottom). Adj P: Bonferroni adjusted p value.

**Figure 5. Analysis of host-pathogen transcriptional signatures in *P. vivax*- infected and non-infected hepatocytes.** t-SNE plots of hepatocytes faceted by infection status. Grey, noninfected (naive and exposed) hepatocytes; light blue, hepatocytes infected with schizonts; teal, hepatocytes infected with hypnozoites.

**Figure 6. Analysis of host transcriptional signatures in *P. vivax* infected hepatocytes.** (A) Volcano plots showing changes in gene expression in infected hepatocytes versus non infected (Wilcoxon rank sum, adjusted p value < 0.05). Positive fold change (FC) values represent genes with greater expression, and negative FC values represent genes with decreased expression. Dashed horizontal red lines: adjusted p value = 0.05; dashed vertical red lines: $\log_2 FC = 0.5$. (B) Summary of differential gene expression data shown in the volcano plots and of the enrichment analyses. Hep: hepatocyte. (C) Enrichment map of cellular processes and pathways associated with hepatocyte infection with *P. vivax* schizonts. (D) Enrichment map of cellular processes and pathways associated with hepatocyte infection with *P. vivax* hypnozoites. For the maps depicted panels C and D, node size is proportional to the number of genes identified in each gene set (minimum 3 genes/gene set); each node represents a distinct biological process or pathway derived from gene with decreased expression (red) or increased expression (blue) versus non-infected cells; and edges (blue lines) represent the number of genes overlapping between two processes or pathways.

**Figure 7. Transcription factor enrichment in hepatocytes infected with schizonts or hypnozoites.** Dot plot of transcription factors enriched in Hepatocyte[+hypnozoite] (left) or Hepatocyte[+schizont] sets (right). Positive (+) enrichment derived from genes displaying greater expression versus non-infected hepatocytes; negative (-) enrichment derived from genes displaying decreased expression versus noninfected hepatocytes. ENCODE (Feingold et al., 2004) and ChEA (Lachmann et al., 2010) consensus transcription factors from ChIP-X (downloaded from Enrichr (Kuleshov et al., 2016)).