IZUMO1 is a sperm fusogen

Nicolas G. Brukman[1*#], Kohdai P. Nakajima[2*], Clari Valansi[1], Kateryna Flyak[1], Xiaohui Li[1], Tetsuya Higashiyama[2,3,4#], Benjamin Podbilewicz[1#]

[1]Department of Biology, Technion- Israel Institute of Technology, Haifa 32000, Israel
[2]Division of Biological Science, Graduate School of Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan
[3]Institute of Transformative Bio-Molecules (WPI-ITbM), Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8601, Japan
[4]Department of Biological Sciences, Graduate School of Science, the University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

*These authors contributed equally.
#Correspondence should be addressed to N.G.B (nbrukman@technion.ac.il), T.H. (higashi@bio.nagoya-u.ac.jp) or B.P. (podbilew@technion.ac.il).

Summary

Mammalian sperm-egg adhesion depends on the trans-interaction between the sperm-specific type I glycoprotein IZUMO1 and its oocyte-specific GPI-anchored receptor JUNO. However, the mechanisms and proteins (fusogens) which mediate the following step of gamete fusion remain unknown. Using live imaging and content mixing assays in a heterologous system and structure-guided mutagenesis, we unveil an unexpected function for IZUMO1 in cell-to-cell fusion. We show that IZUMO1 alone is sufficient to induce fusion, and that this ability is retained in a mutant unable to bind JUNO. On the other hand, a triple mutation in exposed aromatic residues prevents this fusogenic activity without impairing JUNO interaction. Our findings suggest a second, crucial function for IZUMO1 as a unilateral mouse gamete fusogen.

Highlights

- IZUMO1 expression in somatic cells in culture induces cell-to-cell fusion
- The fusogenic activity of IZUMO1 is unilateral
- Cell fusion is independent of the binding of IZUMO1 to JUNO
- IZUMO1-mediated cell merger depends on its transmembrane domain, and three solvent-exposed aromatic residues

Keywords

Sperm, fertilization, oocyte, cell-to-cell fusion, fusogen, IZUMO1, JUNO/IZUMOR, sexual reproduction, GCS1/HAP2.

Introduction

The final steps of mammalian egg-sperm fusion remain a mechanistic enigma. Cell-to-cell fusion requires the action of specialized proteins, named fusogens, to overcome the energetic barriers that arise when two plasma membranes come into close proximity (Chernomordik and Kozlov, 2003). By definition, fusogens are both necessary in their system of origin (*in situ*) and sufficient to induce membrane merging in otherwise non-fusing heterologous systems (Segev et al., 2018). While a mystery in mammals, gamete fusion in flowering plants and protists is mediated by the fusogen GENERATIVE CELL-SPECIFIC 1/HAPLESS 2 (GCS1/HAP2). The essentiality of GCS1/HAP2 in gamete fusion was first demonstrated in *Arabidopsis thaliana*, being sperm-expressed and necessary for sperm-egg and sperm-central cell fusion (von Besser et al., 2006; Johnson et al., 2004; Mori et al., 2006). GCS1/HAP2 is also essential to fuse gametes in the malaria parasite *Plasmodium*, in the slime mold *Dictyostelium* and in the algae *Chlamydomonas* (Hirai et al., 2008; Liu et al., 2008; Okamoto et al., 2016). We have subsequently demonstrated that the expression of *Arabidopsis thaliana* GCS1/HAP2 is sufficient to fuse mammalian cells in culture (Valansi et al., 2017), thereby characterizing this protein as a bona fide fusogen. Remarkably, structural analysis of GCS1/HAP2 from different species showed a shared tertiary and quaternary organization with class II viral glycoproteins (such as those from dengue, rubella and zika viruses) (Fédry et al., 2017; Pinello et al., 2017; Valansi et al., 2017) and with Fusion Family (FF) proteins from nematodes and other organisms (Avinoam et al., 2011; Mohler et al., 2002; Pérez-Vargas et al., 2014; Sapir et al., 2007). This protein superfamily, termed Fusexins (Valansi et al., 2017), are widely distributed in multiple phyla, but to date no members have been identified in vertebrates (Brukman et al., 2022; Vance and Lee, 2020).

Unlike gamete fusogens, other mammalian somatic fusogens are known and well-characterized. For example, fusion of myoblasts to form and maintain muscle fibers requires the coordinate action of Myomaker (TMEM8c) and Myomerger (Myomixer/Minion/Gm7325) (Bi et al., 2017; Millay et al., 2013; Quinn et al., 2017; Zhang et al., 2017). Their expression in fibroblasts drives cell-to-cell fusion: Myomerger can work unilaterally from either one of the merging membranes, while Myomaker is required on both fusing cells (bilateral mechanism) (Leikina et al., 2018). During placenta formation, extensive trophoblast fusion is mediated by syncytins. Remarkably, syncytins are evolutionarily related to retroviral Class I glycoproteins (Lavialle et al., 2013). *Syncytin-A* and *-B* mutations in mice results in fusion defects during the formation of the syncytiotrophoblast (Dupressoir et al., 2009, 2011); while human Syncytin-1 or -2 expression is sufficient to induce heterologous fusion of cells in culture (Blond et al., 2000; Esnault et al., 2008). Interestingly, the fusogenic activity of Syncytin-1 and -2 was highly increased when the respective receptor, ASCT2 or MFSD2, was present (Blond et al., 2000; Esnault et al., 2008). A tight binding step to a receptor in the target cell preceding membrane fusion is a common mechanism among viral fusogens, with binding and fusion often mediated by different domains or

subunits within the fusogen (Podbilewicz, 2014; White et al., 2008).

A number of events preceding gamete fusion take place during fertilization. Sperm
must undergo a physiological process called capacitation, which includes the
exocytosis of the acrosome, a specialized vesicle in the head (Yanagimachi, 1994).
This allows the sperm to penetrate a proteinic coating which covers the oocyte, called
vitelline envelope or, in mammals, *zona pellucida* (ZP) (Wassarman, 1999;
Yanagimachi, 1994). Only after penetration of the ZP, the plasma membranes of both
gametes are able to bind to each other and finally fuse to complete fertilization (Bianchi
and Wright, 2020). Some proteins expressed in the oocyte or in the sperm were shown
to be essential for the last steps of fertilization (Deneke and Pauli, 2021). The oocyte
tetraspanins CD9 and CD81 are required for sperm-egg fusion (Kaji et al., 2000; Le
Naour et al., 2000; Miyado et al., 2000; Rubinstein et al., 2006) by regulating
membrane architecture and compartmentalization (Inoue et al., 2020; Runge et al.,
2007). Mutation of any of the sperm-specific proteins TMEM95, SPACA6, FIMP, SOF1
and DCST1/2 leads to male infertility due to defects in gamete fusion (Barbaux et al.,
2020; Fujihara et al., 2020; Inoue et al., 2021; Lamas-Toranzo et al., 2020; Lorenzetti
et al., 2014; Noda et al., 2020, 2022). While all these genes are essential for late
stages in fertilization, and loss-of-function mutations of any of them prevent gamete
fusion, it is not clear what specific step in the process is being directly affected
(Brukman et al., 2019). The only known pair of trans-interacting proteins are IZUMO1,
in the sperm, and JUNO/IZUMO1R, from the oocyte. IZUMO1 was the first
characterized sperm protein whose deletion was demonstrated to block gamete fusion
in mouse (Inoue et al., 2005), while JUNO was subsequently identified as the IZUMO1
receptor in the oocyte (Bianchi et al., 2014). The IZUMO1-JUNO interaction is required
for efficient sperm-egg binding (Matsumura et al., 2021) in a species-specific manner
(Bianchi and Wright, 2015). Nevertheless, a previous study failed to detect fusion
between mixed Human Embryonic Kidney HEK293T cells expressing JUNO or
IZUMO1 (Bianchi et al., 2014). Here, however, we present evidence that the
expression of mouse IZUMO1 is sufficient to mediate cell-to-cell fusion independently
of its binding to JUNO, thus underpinning an additional role for IZUMO1 in membrane
merger during fertilization.

Results

IZUMO1 is sufficient to induce syncytia formation in somatic cells

There are two main attributes that define a membrane fusogen: First, the molecule
has to be necessary for membrane fusion so when it is absent or mutated fusion fails
to occur; and second, the protein must be sufficient to merge cells that normally do not
fuse. Generally non-fusing cells in culture, such as Baby Hamster Kidney (BHK) cells,
have been widely used to test the fusogenic ability of proteins and illustrate fusogen
sufficiency (Figure S1A, (Avinoam et al., 2011; Bitto et al., 2016; Jeetendra et al., 2002;

Valansi et al., 2017; Waning et al., 2004; White et al., 1981)). To look for gamete
fusogen(s) in mammals, we aimed to evaluate candidate gamete-specific
transmembrane proteins using this system, with mouse egg JUNO and sperm
IZUMO1 as negative controls to account for trans-interacting proteins involved in
binding but not fusion. To our surprise, when we expressed IZUMO1, but not JUNO,
we observed syncytia formation at levels similar to those in a positive control using
GCS1/HAP2, a gamete fusogen from *Arabidopsis thaliana* (Figure 1A). We found that
multinucleation increased three-fold when IZUMO1 was expressed compared to the
negative control containing a membrane-bound myristoylated EGFP (myrGFP)
(Figure 1B). For both IZUMO1 and GCS1/HAP2, we found an increase in cells with
two to five-nuclei (Table S1). The presence of both IZUMO1 and JUNO, on the surface
of BHK cells was confirmed by immunostaining of non-permeabilized samples (Figure
S2).

IZUMO1 is sufficient to fuse cells

Following the observation that IZUMO1 can induce multinucleation, we aimed to
test the ability of IZUMO1 and JUNO to mediate cell-to-cell fusion by using content
mixing experiments (Avinoam et al., 2011; Valansi et al., 2017). In this assay, two
populations of BHK cells expressing either cytosolic EGFP (GFPnes) or nuclear H2B-
RFP are co-incubated, and the appearance of multinucleated cells containing both
fluorescent markers is registered as an indication of fusion (Figures 2A and S3). Under
these conditions, expression of JUNO alone failed to induce content mixing, consistent
with the syncytia formation assay. In contrast, IZUMO1 induced a 5.5-fold increase in
the content mixing of BHK cells compared to the controls employing the fluorescent
empty vectors; while a 4-fold increase was observed in the positive GCS1/HAP2
control (Figures 2B, S3 and S4A). To determine whether IZUMO1 can merge different
heterologous cells, we performed content mixing assays using human HEK293T cells
and obtained similar results (Figure S5). Thus, the mouse sperm protein IZUMO1 is
sufficient to fuse hamster and human cells; whereas the egg receptor of IZUMO1
(JUNO) is unable to fuse heterologous cells (Figure S1B).

IZUMO1-mediated fusion uses a viral-like unilateral mechanism

Some fusogens, like the hemagglutinin of influenza virus or the syncytins, can
mediate fusion when they are expressed in only one of the fusing membranes while
others, like EFF-1 and AFF-1 from nematodes, are required in both fusing membranes
acting in a bilateral way (Blond et al., 2000; Esnault et al., 2008; Podbilewicz et al.,
2006; White et al., 1982). As IZUMO1 is present only in the sperm we aimed to test
whether cells expressing IZUMO1 are able to fuse to cells transfected with an empty
vector or expressing JUNO. We found that content mixing was observed when
IZUMO1 was present only in one of the two populations of cells that were mixed,
implying a unilateral fusion mechanism like in viruses (Figures 2 and S3). Expressing
JUNO *in trans* of IZUMO1 appeared to increase the number of fusion events, which
may be due to increased association between cells from the two populations as a

result of IZUMO1-JUNO interaction. In summary, our results suggest that mouse
gamete fusion can be mediated by the viral-like unilateral fusogenic activity of IZUMO1
on the sperm that follows the docking mediated by the IZUMO1-JUNO interaction
(Figure S1B).

Time-lapse imaging reveals the dynamics of IZUMO1-mediated fusion

To independently study the apparent fusogenic activity of IZUMO1, we performed
live imaging experiments to track fusion events of BHK cells in real time using an
inducible system (Figure S1A). For this purpose, we transfected cells with a plasmid
encoding for cytoplasmic RFP (RFPcyto) alone, for the negative control, or together
with a mifepristone-inducible system to express our candidate proteins. As a positive
control we used EFF-1, a somatic fusogen from *Caenorhabditis elegans* (Avinoam et
al., 2011). Vectors for E-cadherin, GEX2 (Gamete EXpressed 2), IZUMO1 and JUNO
were similarly prepared and the cells were visualized upon induction. We observed
that cell-to-cell fusion occurred following EFF-1 expression (Figure 3A; Movie S1).
Cell-to-cell fusion events were quantified 12 h after induction. We found that EFF-1
expression significantly increased fusion levels (Figure 3B), whereas the adhesion
proteins (E-cadherin, GEX2 and JUNO) did not display fusogenic activity; significantly,
IZUMO1 induced fusion albeit to somewhat lower levels than EFF-1 (Figure 3B;
Movies S2 and S3). Because IZUMO1 was expressed fused to Venus in the C
terminus we could follow its expression; we often observed Venus fluorescence at time
points when fusion started taking place (Figure 3A; Movies S2 and S3). Taken
together, our results utilizing content mixing and live imaging support a model in which
IZUMO1 acts independently of JUNO in a unilateral mechanism to fuse mammalian
cells (Figure S1B).

Somatic cells expressing IZUMO1 attach to mouse oocytes but fail to fuse

Following our observation that IZUMO1 can mediate fusion of heterologous somatic
cells in culture, we then aimed to study a semi-heterologous system which retains the
endogenous components of the oocyte, but not sperm, membranes. To this end, BHK
cells expressing IZUMO1 along with the nuclear marker H2B-RFP were incubated with
mouse oocytes expressing the transgenic protein CD9-GFP on their membrane and
the occurrence of fusion was evaluated (Figure S1A). We found that IZUMO1
expression was sufficient to mediate the binding of BHK cells to oocytes but did not
induce their fusion (Figure S4B), consistent with previous reports (Chalbi et al., 2014;
Inoue et al., 2013, 2015). As a positive control, we used a powerful and promiscuous
viral fusogen, the Vesicular Stomatitis Virus G protein (VSV-G) (Florkiewicz and Rose,
1984). We co-expressed IZUMO1 together with VSV-G to mediate BHK attachment to
the oocytes, and lowered the pH to trigger VSV-G activity (Florkiewicz and Rose,
1984). Only in this condition we observed chromosomes from the BHK in the
cytoplasm of the oocytes (Figure S4B). Thus, while viral VSV-G can efficiently fuse
BHK cells to oocytes, we could not demonstrate a similar role for IZUMO1 under the
conditions employed, which might suggest a role for additional cofactors that may be

required to trigger fusion or bypass the mechanisms that block fusion to the egg.

Dissection of the JUNO-binding and membrane fusion activities of IZUMO1

Some viral fusogens accomplish apposing membrane docking and elicit their
merger using two different structural domains (White et al., 2008). We hypothesized
that IZUMO1 will similarly have different domains mediating its dual activities. To study
the structural features involved in IZUMO1 fusogenic activity, we designed and
generated a series of mutants based on the crystal structure of IZUMO1 and its
interactions with the docking receptor JUNO (Figures 4A and 4B). In order to assess
the functionality of these mutants for the dual role of IZUMO1, we confirmed the
expression of each of the mutant proteins by western blot (Figure 4C) and their
localization by immunostaining (Figure 4D) and compared their performance in binding
and content-mixing experiments. IZUMO1 contains the so-called Izumo domain
composed of a four-helix bundle in the N-terminal part and a β-hairpin or hinge that
links the Izumo domain to an Ig-like domain (Figure 4A; (Nishimura et al., 2016)).
Single-pass transmembrane proteins with Ig-like domains have been involved in cell-
to-cell fusion, like the members of the Fusexin superfamily (Martens and McMahon,
2008; Vance and Lee, 2020). We found that the deletion of the transmembrane domain
produced an inactive form of the protein (IZUMO^{Ecto}; Figure 5A) that was not detected
on the surface of the cells, suggesting that anchoring to the plasma membrane is
required for fusion. When the Ig-like domain of IZUMO1 was absent, the protein
IZUMO1^{ΔIg} was expressed intracellularly (Figure 4C) but not detected on the cellular
surface (Figure 4D), suggesting that this disruption interferes with the correct
localization of the protein. While not detected on the cell surface using
immunostaining, IZUMO1^{ΔIg} was able to induce significant levels of content-mixing
compared to the empty vector, although lower than wild-type IZUMO1 (Figure 5A).
These results suggest that some IZUMO1^{ΔIg} was able to reach the cell surface at low
levels or transiently and was still able to induce fusion, detected as content mixing.
Thus, IZUMO^{Ecto} cannot mediate cell-cell fusion suggesting that the transmembrane
domain is necessary for fusion. In contrast, the Ig-like domain is required for correct
trafficking and localization to the plasma membrane but does not appear to be
essential for fusion, since IZUMO1^{ΔIg} is still able to mediate cell-cell fusion.

To directly study whether the fusogenic activity of IZUMO1 is correlated with the
ability to bind JUNO, we analyzed a mutant bearing a mutation in a conserved
tryptophan residue in the hinge region that is essential for IZUMO1-JUNO interaction
(W148A, Figure 4A, (Aydin et al., 2016; Ohto et al., 2016)). We found that
IZUMO1^{W148A} had no effect on the levels of content mixing compared to wild-type
IZUMO1 (Figure 5A) but disrupted the binding of BHK cells to oocytes (Figures 5B and
5C). Unlike IZUMO^{Ecto} or IZUMO1^{ΔIg}, the W148A mutation did not affect its localization
on the surface (Figure 4D). Thus, the W148A residue on the β-hairpin of IZUMO1 is
required for binding the oocyte via JUNO but does not play an active role in cell-to-cell
fusion.

The four-helix bundle of the Izumo domain contains three exposed aromatic
residues (Figure 4A; (Nishimura et al., 2016)) which may be important for the
interaction of IZUMO1 to the membrane of the oocyte or to other proteins given that
JUNO is displaced from the fusion site shortly after the apposition of the fusing
membranes (Inoue et al., 2015). To test whether these residues are required for
efficient fusion, we generated a triple mutant F28A, W88A and W113A (FWW, Figure
4B). IZUMO1^{FWW} was detected on the surface of BHK cells (Figure 4D) and was still
able to mediate BHK binding to the oocyte (Figures 5B and 5C), however, this mutant
induced significantly lower levels of content mixing than wild-type IZUMO1 (Figure 5A).
To summarize, our results support the existence of two functional domains in IZUMO1,
with binding to JUNO mediated through W148 and cell-to-cell fusion dependent on
three exposed aromatic residues on the four-helix bundle of the Izumo domain. These
results allow us to uncouple the dual role of IZUMO1 as a JUNO-interacting binding
protein and a unilateral fusogen, which utilize separate structural motifs.

Discussion

Aromatic residues on the four-helix bundle are required for fusion
independently of the JUNO-IZUMO1 interactions

Our results show that IZUMO1 is sufficient to induce heterologous cell-to-cell
fusion. While IZUMO1 shares structural similarities with proteins involved in cell
invasion of malaria parasites (Nishimura et al., 2016), there is no obvious homology
between IZUMO1 and any known fusogen. Additionally, IZUMO1 lacks any clear
hydrophobic stretch that could work as a fusion peptide or loop which serves to anchor
to the opposing membrane (Aydin et al., 2016), a characteristic feature which is shared
by many, but not all, cellular and viral fusogens (Brukman et al., 2019). However, the
four-helix bundle of the Izumo domain contains solvent-exposed aromatic residues
(Nishimura et al., 2016); which are not involved in JUNO-IZUMO1 interaction (Aydin
et al., 2016; Ohto et al., 2016) but might be relevant for interacting with the egg
membrane. This is supported by our findings that the simultaneous mutation of these
residues did not affect the transinteraction of IZUMO1-JUNO mediating binding of BHK
cells to oocytes, but did reduce the fusogenic activity of IZUMO1 (Figure 5). While F28
is well conserved in rodents, W88 and W113 are present in most IZUMO1 mammalian
orthologs (Aydin et al., 2016; Ohto et al., 2016). Interestingly, monoclonal antibodies
against the N-terminal region of IZUMO1 were capable of inhibiting gamete fusion *in*
vitro without altering sperm binding to the oocyte (Inoue et al., 2013), supporting not
only an additional role of IZUMO1 in fusion besides binding, but also a relevance of
the Izumo domain in these later stages of gamete fusion. Alternatively, the aromatic
residues on the surface of the Izumo domain may be required for the interaction with
an unknown secondary receptor (Inoue et al., 2015), for oligomerization (Ellerman et
al., 2009) or for interaction with the other domains within IZUMO1 protein after a
conformational change (Inoue et al., 2015).

Spatial and temporal dissection of egg-sperm binding and fusion

Previous work has established that after sperm binding, JUNO is excluded from the interface between egg and sperm, while IZUMO1 is conversely enriched in this region (Inoue et al., 2015). These observations are consistent with a dual role of IZUMO1 as a fusogen. The concentration of IZUMO1 at the fusion zone is accompanied by a conformational change that depends on disulfide isomerase activity, and dimerization (Inoue et al., 2015). Furthermore, IZUMO1 has a more ancestral origin, being found in many vertebrate phyla, while JUNO is found in mammals only (Grayson, 2015). This evolutionary distribution, as well as IZUMO1's unilateral fusogenic activity, suggest a function for IZUMO1 in fertilization that is independent of its well-established interaction with JUNO. This is further supported by our finding that the IZUMO1^{W148A} mutant which fails to bind JUNO is still able to induce cell-to-cell fusion, and that a triple mutant of the exposed aromatic residues in the four-helix bundle is able to mediate robust BHK binding to oocytes while cell–to-cell fusion was significantly reduced. Additionally, our results suggest a role for the IZUMO1 Ig-like domain for correct localization of the protein to the membrane (Figure 4D), however, it is not essential for its fusogenic activity (Figure 5A). This change in the localization could be explained by the lack of the single glycosylation site at the Ig-like domain that was reported to be relevant for protecting the protein from degradation but not for its activity in fertilization (Inoue et al., 2008).

Are other egg and sperm membrane proteins involved in membrane merger?

While our results point to IZUMO1 as a *bona fide* fusogen that can fuse hamster and human cells in culture, this does not exclude the possibility that other fusogenic proteins work cooperatively with IZUMO1 to ensure the success and specificity of fertilization *in vivo* (Chalbi et al., 2014; Deneke and Pauli, 2021; Ellerman et al., 2009; Gaikwad et al., 2019; Inoue et al., 2013, 2015). This kind of cooperation can also be seen in plant fertilization, where the mutation of DMP8/9 proteins drastically reduces the activity of GCS1/HAP2 in vivo (Cyprys et al., 2019; Zhang et al., 2019). Proteins related to IZUMO1, for instance, IZUMO2-4, SPACA6 or TMEM95 may work as fusogens or potentiate IZUMO1's fusogenic activity (Deneke and Pauli, 2021; Ellerman et al., 2009). Notably, IZUMO1 can homo- and hetero-oligomerize (Ellerman et al., 2009; Gaikwad et al., 2019). The absence of these proteins may explain the inability of somatic cells expressing IZUMO1 to fuse to oocytes (Figure S4B, also see (Chalbi et al., 2014; Inoue et al., 2013, 2015)). We cannot exclude other factors, such as the movement of the flagella (Ravaux et al., 2016) or the exposure of phosphatidylserine on the sperm (Rival et al., 2019), that may also be required for a successful fusion with the oocyte.

Additionally, the fact that other eukaryotic fusogens (*e.g.* EFF-1, AFF-1 and GCS1/HAP2) were shown to be less efficient *in vitro* than their viral counterparts (Avinoam et al., 2011; Valansi et al., 2017) might also contribute to explain why previous studies failed to detect fusion of somatic cells expressing IZUMO1 and JUNO

324 (Bianchi et al., 2014).

IZUMO1 activity is regulated during fertilization

Despite IZUMO1 being a unilateral fusogen, sperm cells do not fuse with other
sperm or with other cell types in the male or female reproductive tracts. Furthermore,
gamete fusion is species-specific (Yanagimachi, 1988). These non-physiological
fusion events do not lead to a successful formation of a zygote and therefore reduce
the individual fitness. The changes in the localization of IZUMO1 during sperm transit
may partially explain this. In fresh sperm, IZUMO1 is not exposed on the outer
membrane, being localized to the interior of the acrosome (Inoue et al., 2005), a giant
specialized vesicle in the head, and therefore it cannot mediate cell-to-cell fusion. Only
after capacitation and the acrosomal exocytosis in the female tract, IZUMO1 is
exposed to the cell surface and migrates to the fusogenic region of the sperm plasma
membrane (Inoue et al., 2005; Satouh et al., 2012). In addition, the requirement of a
tight binding step to JUNO prior to fusion determines that the sperm cell will fuse to
oocytes expressing a species-matching IZUMO1 receptor (Bianchi and Wright, 2015;
Bianchi et al., 2014). This is a common mechanism used by many viruses to regulate
their cellular tropism even when carrying powerful unilateral fusogens, such as the
requirement of CD4 and other co-receptors for HIV infection mediated by the Env
glycoprotein (Melikyan, 2011; White et al., 2008). Another analogy can be made with
intracellular fusogens such as the SNARE complex that requires the regulatory
activities of Munc18, Munc13 and synaptotagmin for efficient fusion (Stepien and Rizo,
2021). After sperm-to-oocyte fusion occurs, JUNO is shed from the plasma membrane
preventing further sperm to bind, and therefore, contributing to a block to polyspermy
(Bianchi et al., 2014).

Concluding remarks

Up to this report, GCS1/HAP2 proteins were the only known fusogens involved in
fertilization (reviewed in (Brukman et al., 2022)); these proteins are structurally and
evolutionarily related to Class II fusogens from enveloped viruses and FF fusogens
from nematodes (Fédry et al., 2017; Pinello et al., 2017; Valansi et al., 2017).
Together, these fusogens form a superfamily called Fusexins. The involvement of
GCS1/HAP2 in gamete fusion seems to be a characteristic of eukaryotes (Pinello et
al., 2017; Valansi et al., 2017); however, many sexually reproducing organisms
including fungi and vertebrates lack a fusexin homologue. Considering their
distributions in the phylogenetic tree (Vance and Lee, 2020), it is possible that
IZUMO1-type proteins are chordate's innovations that replaced GCS1/HAP2 during
evolution as a strategy for gamete fusion. Combined with previous information
(Deneke and Pauli, 2021; Ellerman et al., 2009; Inoue et al., 2013, 2015)(Stepien and
Rizo, 2021), we hypothesize that IZUMO1 from the sperm first transiently binds JUNO
on the oocyte for docking and subsequently undergoes a conformational change and
oligomerization that induce sperm-egg fusion (Figure S1B). This role in membrane
merging makes IZUMO1 even more suited to its name, coined after the Japanese

shrine dedicated to marriage (Inoue et al., 2005).

Acknowledgments

We thank Gavin Wright for the plasmids containing *mIzumo1* and *mJuno* sequences
and Masahito Ikawa for the CD9-GFP transgenic mouse line. We thank Pablo Aguilar,
Dan Cassel, Masahito Ikawa, Yael Iosilevskii and Luca Jovine for critically reading the
manuscript. This work was funded by the Israel Science Foundation (ISF grants
257/17, 2462/18, 2327/19 and 178/20 to B.P.), by Grant-in-Aid for Scientific Research
on Innovative Areas (16H06465, 16H06464, and 16K21727 to T.H.), by JST, ERATO
(JPMJER1004 to T.H.) and CREST (JPMJCR20E5 to T.H.). This project has received
funding from the European Union's Horizon 2020 research and innovation programme
under the Marie Skłodowska-Curie (grant agreement No 844807 to N.G.B).

Author contributions

378 B.P. and T.H. conceived the study. N.G.B., K.P.N., C.V., X.L. and K.F. designed and
379 conducted experiments, and processed and analyzed the data supervised by B.P. and
380 T.H.; N.G.B. and B.P. wrote the initial draft of the manuscript. All authors participated
381 in discussions of results and manuscript editing.

Declaration of interests

384 The authors declare no competing interests.

References

Avinoam, O., Fridman, K., Valansi, C., Abutbul, I., Zeev-Ben-Mordehai, T., Maurer,
U.E., Sapir, A., Danino, D., Grünewald, K., White, J.M., et al. (2011). Conserved
eukaryotic fusogens can fuse viral envelopes to cells. Science *332*, 589–592.

Aydin, H., Sultana, A., Li, S., Thavalingam, A., and Lee, J.E. (2016). Molecular
architecture of the human sperm IZUMO1 and egg JUNO fertilization complex.
Nature *534*, 562–565.

Barbaux, S., Ialy-Radio, C., Chalbi, M., Dybal, E., Homps-Legrand, M., Do Cruzeiro,
M., Vaiman, D., Wolf, J.-P., and Ziyyat, A. (2020). Sperm SPACA6 protein is
required for mammalian Sperm-Egg Adhesion/Fusion. Sci. Rep. *10*, 5335.

von Besser, K., Frank, A.C., Johnson, M.A., and Preuss, D. (2006). Arabidopsis
HAP2 (GCS1) is a sperm-specific gene required for pollen tube guidance and
fertilization. Development *133*, 4761–4769.

Bi, P., Ramirez-Martinez, A., Li, H., Cannavino, J., McAnally, J.R., Shelton, J.M., Sánchez-Ortiz, E., Bassel-Duby, R., and Olson, E.N. (2017). Control of muscle formation by the fusogenic micropeptide myomixer. Science *356*, 323–327.

Bianchi, E., and Wright, G.J. (2015). Cross-species fertilization: The hamster egg receptor, Juno, binds the human sperm ligand, Izumo1. Philos. Trans. R. Soc. Lond. B Biol. Sci. *370(1661)*, 20140101.

Bianchi, E., and Wright, G.J. (2020). Find and fuse: Unsolved mysteries in sperm-egg recognition. PLoS Biol. *18*, e3000953.

Bianchi, E., Doe, B., Goulding, D., and Wright, G.J. (2014). Juno is the egg Izumo receptor and is essential for mammalian fertilization. Nature *508*, 483–487.

Bitto, D., Halldorsson, S., Caputo, A., and Huiskonen, J.T. (2016). Low pH and Anionic Lipid-dependent Fusion of Uukuniemi Phlebovirus to Liposomes. J. Biol. Chem. *291*, 6412–6422.

Blond, J.L., Lavillette, D., Cheynet, V., Bouton, O., Oriol, G., Chapel-Fernandes, S., Mandrand, B., Mallet, F., and Cosset, F.L. (2000). An envelope glycoprotein of the human endogenous retrovirus HERV-W is expressed in the human placenta and fuses cells expressing the type D mammalian retrovirus receptor. J. Virol. *74*, 3321–3329.

Brukman, N.G., Uygur, B., Podbilewicz, B., and Chernomordik, L.V. (2019). How cells fuse. J. Cell Biol. *218*, 1436–1451.

Brukman, N.G., Li, X., and Podbilewicz, B. (2022). Fusexins, HAP2/GCS1 and Evolution of Gamete Fusion. Front Cell Dev Biol *9*, 824024.

Chalbi, M., Barraud-Lange, V., Ravaux, B., Howan, K., Rodriguez, N., Soule, P., Ndzoudi, A., Boucheix, C., Rubinstein, E., Wolf, J.P., et al. (2014). Binding of sperm protein Izumo1 and its egg receptor Juno drives Cd9 accumulation in the intercellular contact area prior to fusion during mammalian fertilization. Development *141*, 3732–3739.

Chernomordik, L.V., and Kozlov, M.M. (2003). Protein-lipid interplay in fusion and fission of biological membranes. Annu. Rev. Biochem. *72*, 175–207.

Cyprys, P., Lindemeier, M., and Sprunck, S. (2019). Gamete fusion is facilitated by two sperm cell-expressed DUF679 membrane proteins. Nat Plants *5*, 253–257.

Deneke, V.E., and Pauli, A. (2021). The Fertilization Enigma: How Sperm and Egg Fuse. Annu. Rev. Cell Dev. Biol. *37*, 391–414.

Dunsing, V., Luckner, M., Zühlke, B., Petazzi, R.A., Herrmann, A., and Chiantia, S. (2018). Optimal fluorescent protein tags for quantifying protein oligomerization in living cells. Sci. Rep. *8*, 10634.

Dupressoir, A., Vernochet, C., Bawa, O., Harper, F., Pierron, G., Opolon, P., and Heidmann, T. (2009). Syncytin-A knockout mice demonstrate the critical role in placentation of a fusogenic, endogenous retrovirus-derived, envelope gene. Proc.

Natl. Acad. Sci. U. S. A. *106*, 12127–12132.

Dupressoir, A., Vernochet, C., Harper, F., Guégan, J., Dessen, P., Pierron, G., and
Heidmann, T. (2011). A pair of co-opted retroviral envelope syncytin genes is
required for formation of the two-layered murine placental syncytiotrophoblast. Proc.
Natl. Acad. Sci. U. S. A. *108*, E1164–E1173.

Ellerman, D.A., Pei, J., Gupta, S., Snell, W.J., Myles, D., and Primakoff, P. (2009).
Izumo is part of a multiprotein family whose members form large complexes on
mammalian sperm. Mol. Reprod. Dev. *76*, 1188–1199.

Esnault, C., Priet, S., Ribet, D., Vernochet, C., Bruls, T., Lavialle, C., Weissenbach,
J., and Heidmann, T. (2008). A placenta-specific receptor for the fusogenic,
endogenous retrovirus-derived, human syncytin-2. Proc. Natl. Acad. Sci. U. S. A.
105, 17532–17537.

Fédry, J., Liu, Y., Péhau-Arnaudet, G., Pei, J., Li, W., Tortorici, M.A., Traincard, F.,
Meola, A., Bricogne, G., Grishin, N.V., et al. (2017). The Ancient Gamete Fusogen
HAP2 Is a Eukaryotic Class II Fusion Protein. Cell *168*, 904–915.e10.

Florkiewicz, R.Z., and Rose, J.K. (1984). A cell line expressing vesicular stomatitis
virus glycoprotein fuses at low pH. Science *225*, 721–723.

Fujihara, Y., Lu, Y., Noda, T., Oji, A., Larasati, T., Kojima-Kita, K., Yu, Z., Matzuk,
R.M., Matzuk, M.M., and Ikawa, M. (2020). Spermatozoa lacking Fertilization
Influencing Membrane Protein (FIMP) fail to fuse with oocytes in mice. Proc. Natl.
Acad. Sci. U. S. A. *117*, 9393–9400.

Gaikwad, A.S., Anderson, A.L., Merriner, D.J., O'Connor, A.E., Houston, B.J., Aitken,
R.J., O'Bryan, M.K., and Nixon, B. (2019). GLIPR1L1 is an IZUMO-binding protein
required for optimal fertilization in the mouse. BMC Biol. *17*, 86.

Grayson, P. (2015). Izumo1 and Juno: the evolutionary origins and coevolution of
essential sperm–egg binding partners. Royal Society Open Science *2*.

Hirai, M., Arai, M., Mori, T., Miyagishima, S. ya, Kawai, S., Kita, K., Kuroiwa, T.,
Terenius, O., and Matsuoka, H. (2008). Male Fertility of Malaria Parasites Is
Determined by GCS1, a Plant-Type Reproduction Factor. Curr. Biol. *18*, 607–613.

Inoue, N., Ikawa, M., Isotani, A., and Okabe, M. (2005). The immunoglobulin
superfamily protein Izumo is required for sperm to fuse with eggs. Nature *434*, 234–
238.

Inoue, N., Ikawa, M., and Okabe, M. (2008). Putative sperm fusion protein IZUMO
and the role of N-glycosylation. Biochem. Biophys. Res. Commun. *377*, 910–914.

Inoue, N., Hamada, D., Kamikubo, H., Hirata, K., Kataoka, M., Yamamoto, M.,
Ikawa, M., Okabe, M., and Hagihara, Y. (2013). Molecular dissection of IZUMO1, a
sperm protein essential for sperm-egg fusion. Development *140*, 3221–3229.

Inoue, N., Hagihara, Y., Wright, D., Suzuki, T., and Wada, I. (2015). Oocyte-triggered
dimerization of sperm IZUMO1 promotes sperm-egg fusion in mice. Nat. Commun.

6, 8858.

Inoue, N., Saito, T., and Wada, I. (2020). Unveiling a novel function of CD9 in
surface compartmentalization of oocytes. Development *147 (15): dev189985*.

Inoue, N., Hagihara, Y., and Wada, I. (2021). Evolutionarily conserved sperm factors,
DCST1 and DCST2, are required for gamete fusion. Elife 10:e66313.

Jeetendra, E., Robison, C.S., Albritton, L.M., and Whitt, M.A. (2002). The
membrane-proximal domain of vesicular stomatitis virus G protein functions as a
membrane fusion potentiator and can induce hemifusion. J. Virol. *76*, 12300–12311.

Johnson, M.A., Besser, K. von, Zhou, Q., Smith, E., Aux, G., Patton, D., Levin, J.Z.,
and Preuss, D. (2004). Arabidopsis hapless Mutations Define Essential
Gametophytic Functions. Genetics *168*, 971 (2):971-82.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583–589.

Kaji, K., Oda, S., Shikano, T., Ohnuki, T., Uematsu, Y., Sakagami, J., Tada, N.,
Miyazaki, S., and Kudo, A. (2000). The gamete fusion process is defective in eggs of
Cd9-deficient mice. Nat. Genet. *24*, 279–282.

Kito, S., Hayao, T., Noguchi-Kawasaki, Y., Ohta, Y., Hideki, U., and Tateno, S.
(2004). Improved in vitro fertilization and development by use of modified human
tubal fluid and applicability of pronucleate embryos for cryopreservation by rapid
freezing in inbred mice. Comp. Med. *54*, 564–570.

Lamas-Toranzo, I., Hamze, J.G., Bianchi, E., Fernández-Fuertes, B., Pérez-
Cerezales, S., Laguna-Barraza, R., Fernández-González, R., Lonergan, P.,
Gutiérrez-Adán, A., Wright, G.J., et al. (2020). TMEM95 is a sperm membrane
protein essential for mammalian fertilization. Elife *9:e53913*.

Lavialle, C., Cornelis, G., Dupressoir, A., Esnault, C., Heidmann, O., Vernochet, C.,
and Heidmann, T. (2013). Paleovirology of "syncytins", retroviral env genes exapted
for a role in placentation. Philos. Trans. R. Soc. Lond. B Biol. Sci. *368*, 20120507.

Leikina, E., Gamage, D.G., Prasad, V., Goykhberg, J., Crowe, M., Diao, J., Kozlov,
M.M., Chernomordik, L.V., and Millay, D.P. (2018). Myomaker and Myomerger Work
Independently to Control Distinct Steps of Membrane Remodeling during Myoblast
Fusion. Dev. Cell *46*, 767–780.e7.

Le Naour, F., Rubinstein, E., Jasmin, C., Prenant, M., and Boucheix, C. (2000).
Severely reduced female fertility in CD9-deficient mice. Science *287*, 319–321.

Liang, L.F., Chamow, S.M., and Dean, J. (1990). Oocyte-specific expression of
mouse Zp-2: developmental regulation of the zona pellucida genes. Mol. Cell. Biol.
10, 1507–1515.

Liu, Y., Tewari, R., Ning, J., Blagborough, A.M., Garbom, S., Pei, J., Grishin, N.V.,
Steele, R.E., Sinden, R.E., Snell, W.J., et al. (2008). The conserved plant sterility

gene HAP2 functions after attachment of fusogenic membranes in Chlamydomonas
and Plasmodium gametes. Genes Dev. *22*, 1051-1068.

Lorenzetti, D., Poirier, C., Zhao, M., Overbeek, P.A., Harrison, W., and Bishop, C.E.
(2014). A transgenic insertion on mouse chromosome 17 inactivates a novel
immunoglobulin superfamily gene potentially involved in sperm-egg fusion. Mamm.
Genome *25*, 141–148.

Martens, S., and McMahon, H.T. (2008). Mechanisms of membrane fusion: disparate
players and common principles. Nat. Rev. Mol. Cell Biol. *9*, 543–556.

Matsumura, T., Noda, T., Satouh, Y., Morohoshi, A., Yuri, S., Ogawa, M., Lu, Y.,
Isotani, A., and Ikawa, M. (2021). Sperm IZUMO1 is required for binding preceding
fusion with oolemma in mice and rats. Front Cell Dev Biol *9*, 3849.

Melikyan, G.B. (2011). Chapter 4 - Membrane Fusion Mediated by Human
Immunodeficiency Virus Envelope Glycoprotein. In Current Topics in Membranes,
L.V. Chernomordik, and M.M. Kozlov, eds. (Academic Press), pp. 81–106.

Millay, D.P., O'Rourke, J.R., Sutherland, L.B., Bezprozvannaya, S., Shelton, J.M.,
Bassel-Duby, R., and Olson, E.N. (2013). Myomaker is a membrane activator of
myoblast fusion and muscle formation. Nature *499*, 301–305.

Miyado, K., Yamada, G., Yamada, S., Hasuwa, H., Nakamura, Y., Ryu, F., Suzuki,
K., Kosai, K., Inoue, K., Ogura, A., et al. (2000). Requirement of CD9 on the egg
plasma membrane for fertilization. Science *287*, 321–324.

Miyado, K., Yoshida, K., Yamagata, K., Sakakibara, K., Okabe, M., Wang, X.,
Miyamoto, K., Akutsu, H., Kondo, T., Takahashi, Y., et al. (2008). The fusing ability
of sperm is bestowed by CD9-containing vesicles released from eggs in mice.
Proceedings of the National Academy of Sciences *105*, 12921–12926.

Mohler, W.A., Shemer, G., del Campo, J.J., Valansi, C., Opoku-Serebuoh, E.,
Scranton, V., Assaf, N., White, J.G., and Podbilewicz, B. (2002). The type I
membrane protein EFF-1 is essential for developmental cell fusion. Dev. Cell *2*, 355–
362.

Moi, D., Nishio, S., Li, X., Valansi, C., Langleib, M., Brukman, N.G., Flyak, K.,
Dessimoz, C., Sanctis, D. de, Tunyasuvunakool, K., et al. (2021). Archaeal origins of
gamete fusion. bioRxiv 2021.10.13.464100.

Mori, T., Kuroiwa, H., Higashiyama, T., and Kuroiwa, T. (2006). GENERATIVE CELL
SPECIFIC 1 is essential for angiosperm fertilization. Nat. Cell Biol. *8*, 64–71.

Nakajima, K.P., Valansi, C., Kurihara, D., Sasaki, N., Podbilewicz, B., and
Higashiyama, T. (2021). A novel live imaging-based assay for visualising species-
specific interactions in gametic adhesion molecules. Research Square
10.21203/rs.3.rs-1174354/v1.

Nicolson, G.L., Yanagimachi, R., and Yanagimachi, H. (1975). Ultrastructural
localization of lectin-binding sites on the zonae pellucidae and plasma membranes of
mammalian eggs. J. Cell Biol. *66*, 263–274.

Nishimura, K., Han, L., Bianchi, E., Wright, G.J., de Sanctis, D., and Jovine, L.
(2016). The structure of sperm Izumo1 reveals unexpected similarities with
Plasmodium invasion proteins. Curr. Biol. *26*, R661–R662.

Noda, T., Lu, Y., Fujihara, Y., Oura, S., Koyano, T., Kobayashi, S., Matzuk, M.M.,
and Ikawa, M. (2020). Sperm proteins SOF1, TMEM95, and SPACA6 are required
for sperm-oocyte fusion in mice. Proc. Natl. Acad. Sci. U. S. A. *117*, 11493–11502.

Noda, T., Blaha, A., Fujihara, Y., Gert, K.R., Emori, C., Deneke, V.E., Oura, S.,
Panser, K., Lu, Y., Berent, S., et al. (2022). Sperm membrane proteins DCST1 and
DCST2 are required for sperm-egg interaction in mice and fish. Commun Biol *5*, 332.

Ohto, U., Ishida, H., Krayukhina, E., Uchiyama, S., Inoue, N., and Shimizu, T.
(2016). Structure of IZUMO1–JUNO reveals sperm–oocyte recognition during
mammalian fertilization. Nature *534*, 566–569.

Okamoto, M., Yamada, L., Fujisaki, Y., Bloomfield, G., Yoshida, K., Kuwayama, H.,
Sawada, H., Mori, T., and Urushihara, H. (2016). Two HAP2-GCS1 homologs
responsible for gamete interactions in the cellular slime mold with multiple mating
types: Implication for common mechanisms of sexual reproduction shared by plants
and protozoa and for male-female differentiation. Dev. Biol. *415* (1): 6–13.

Pérez-Vargas, J., Krey, T., Valansi, C., Avinoam, O., Haouz, A., Jamin, M., Raveh-
Barak, H., Podbilewicz, B., and Rey, F.A. (2014). Structural Basis of Eukaryotic Cell-
Cell Fusion. Cell *157*, 407–419.

Pinello, J.F., Lai, A.L., Millet, J.K., Cassidy-Hanley, D., Freed, J.H., and Clark, T.G.
(2017). Structure-function studies link class II viral fusogens with the ancestral
gamete fusion protein HAP2. Curr. Biol. 27(5):651-660.

Podbilewicz, B. (2014). Virus and Cell Fusion Mechanisms. Annu. Rev. Cell Dev.
Biol. *30*, 111–139.

Podbilewicz, B., Leikina, E., Sapir, A., Valansi, C., Suissa, M., Shemer, G., and
Chernomordik, L.V. (2006). The C. elegans developmental fusogen EFF-1 mediates
homotypic fusion in heterologous cells and in vivo. Dev. Cell *11*, 471–481.

Quinn, M.E., Goh, Q., Kurosaka, M., Gamage, D.G., Petrany, M.J., Prasad, V., and
Millay, D.P. (2017). Myomerger induces fusion of non-fusogenic cells and is required
for skeletal muscle development. Nat. Commun. *8*, 15665.

Ravaux, B., Garroum, N., Perez, E., Willaime, H., and Gourier, C. (2016). A specific
flagellum beating mode for inducing fusion in mammalian fertilization and kinetics of
sperm internalization. Scientific Reports 2016 6:1 *6*, 1–13.

Rival, C.M., Xu, W., Shankman, L.S., Morioka, S., Arandjelovic, S., Lee, C.S.,
Wheeler, K.M., Smith, R.P., Haney, L.B., Isakson, B.E., et al. (2019).
Phosphatidylserine on viable sperm and phagocytic machinery in oocytes regulate
mammalian fertilization. Nat. Commun. *10*.

Rubinstein, E., Ziyyat, A., Prenant, M., Wrobel, E., Wolf, J.-P., Levy, S., Le Naour,
F., and Boucheix, C. (2006). Reduced fertility of female mice lacking CD81. Dev.

Biol. *290*, 351–358.

Runge, K.E., Evans, J.E., He, Z.-Y., Gupta, S., McDonald, K.L., Stahlberg, H.,
Primakoff, P., and Myles, D.G. (2007). Oocyte CD9 is enriched on the microvillar
membrane and required for normal microvillar shape and distribution. Dev. Biol. *304*,
317–325.

Sapir, A., Choi, J., Leikina, E., Avinoam, O., Valansi, C., Chernomordik, L.V.,
Newman, A.P., and Podbilewicz, B. (2007). AFF-1, a FOS-1-regulated fusogen,
mediates fusion of the anchor cell in C. elegans. Dev. Cell *12*, 683–698.

Satouh, Y., Inoue, N., Ikawa, M., and Okabe, M. (2012). Visualization of the moment
of mouse sperm-egg fusion and dynamic localization of IZUMO1. J. Cell Sci. *125*,
4985–4990.

Segev, N., Avinoam, O., and Podbilewicz, B. (2018). Fusogens. Curr. Biol. *28*,
R378–R380.

Stepien, K.P., and Rizo, J. (2021). Synaptotagmin-1–, Munc18-1–, and Munc13-1–
dependent liposome fusion with a few neuronal SNAREs. Proc. Natl. Acad. Sci. U. S.
A. *118(4) e2019314118*.

Truffi, M., Dubreuil, V., Liang, X., Vacaresse, N., Nigon, F., Han, S.P., Yap, A.S.,
Gomez, G.A., and Sap, J. (2014). RPTPα controls epithelial adherens junctions,
linking E-cadherin engagement to c-Src-mediated phosphorylation of cortactin. J.
Cell Sci. *127*, 2420–2432.

Valansi, C., Moi, D., Leikina, E., Matveev, E., Graña, M., Chernomordik, L.V.,
Romero, H., Aguilar, P.S., and Podbilewicz, B. (2017). Arabidopsis HAP2/GCS1 is a
gamete fusion protein homologous to somatic and viral fusogens. J. Cell Biol. *216*,
571–581.

Vance, T.D.R., and Lee, J.E. (2020). Virus and eukaryote fusogen superfamilies.
Curr. Biol. *30*, R750–R754.

Waning, D.L., Russell, C.J., Jardetzky, T.S., and Lamb, R.A. (2004). Activation of a
paramyxovirus fusion protein is modulated by inside-out signaling from the
cytoplasmic tail. Proc. Natl. Acad. Sci. U. S. A. *101*, 9217–9222.

Wassarman, P.M. (1999). Mammalian fertilization: Molecular aspects of gamete
adhesion, exocytosis, and fusion. Cell *96*, 175–183.

White, J., Matlin, K., and Helenius, A. (1981). Cell fusion by Semliki Forest,
influenza, and vesicular stomatitis viruses. J. Cell Biol. *89*, 674–679.

White, J., Helenius, A., and Gething, M.J. (1982). Haemagglutinin of influenza virus
expressed from a cloned gene promotes membrane fusion. Nature *300*, 658–659.

White, J.M., Delos, S.E., Brecher, M., and Schornberg, K. (2008). Structures and
mechanisms of viral membrane fusion proteins: multiple variations on a common
theme. Crit. Rev. Biochem. Mol. Biol. *43*, 189–219.

Williams, R.M., Senanayake, U., Artibani, M., Taylor, G., Wells, D., Ahmed, A.A., and
Sauka-Spengler, T. (2018). Genome and epigenome engineering CRISPR toolkit for
in vivo modulation of cis-regulatory interactions and gene expression in the chicken
embryo. Development *145(4):dev160333*.

Yanagimachi, R. (1988). Chapter 1 Sperm–Egg Fusion. In Current Topics in
Membranes and Transport, F. Bronner, ed. (Academic Press), pp. 3–43.

Yanagimachi, R. (1994). Mammalian fertilization. In The Physiology of Reproduction,
E. Knobil, and J.D. Neill, eds. (New York: Raven press), pp. 189–317.

Zhang, J., Pinello, J.F., and Snell, W.J. (2019). Plant sperm need a little help. Nat
Plants *5*, 247–248.

Zhang, Q., Vashisht, A.A., O'Rourke, J., Corbel, S.Y., Moran, R., Romero, A.,
Miraglia, L., Zhang, J., Durrant, E., Schmedt, C., et al. (2017). The microprotein
Minion controls cell fusion and muscle formation. Nat. Commun. *8*, 15664.

Figure 1. IZUMO1 induces multinucleation of BHK cells

(A) Cells were transfected with either pCI::myrGFP::H2B-RFP (myristoylated EGFP), pCI::GCS1/HAP-V5::H2B-RFP, pCI::IZUMO1-V5:H2B-RFP or pCI::JUNO::H2B-RFP vectors. JUNO was co-transfected with a plasmid for myrGFP. Immunofluorescence was performed with anti-V5 antibodies for IZUMO1 and GCS1/HAP2, while the myrGFP signal is shown for JUNO and vector control (gray). Arrows show cells with more than one nucleus (magenta). Scale bars, 20 μm.

(B) The percentage of multinucleation (see Supplementary Information) is presented as individual data and means ± SEM of four independent experiments. The number of cells counted can be found in Table S1. Comparisons were made with one-way ANOVA followed by Dunett's test against the empty vector. ns = non-significant, ** $p < 0.01$, *** $p < 0.001$.

Figure 2. IZUMO1 induces fusion of BHK cells

(A) Representative images of mixed cells transfected with pCI::GFPnes or pCI::H2B-RFP empty vectors or containing the coding sequence for the expression of GCS1/HAP2, IZUMO1 and JUNO as indicated. Arrows show content-mixed cells containing both GFPnes (green cytoplasm) and H2B-RFP (red nuclei). DAPI staining is shown in blue. Scale Bars, 20 μm. See also Figure S3.

(B) Quantification of content-mixing experiments. The percentage of mixing was defined as the ratio between the nuclei in mixed cells (NuM) and the total number of nuclei in mixed cells and fluorescent cells in contact that did not fuse (NuC), as follows: % of mixing = (NuM/(NuM+NuC)) x 100. Bar chart showing individual experiment values (each corresponding to 1000 nuclei) and means ± SEM of four independent experiments. Comparisons by one-way ANOVA followed by Dunett's test against the empty vectors. ns = non-significant, ** $p < 0.01$, **** $p < 0.0001$.

Figure 3. Quantification of fusion activity using live imaging

(A) Time-lapse images from a fusion assay. BHK cells were transfected with plasmids for expression of cytoplasmic RFP (RFPcyto, magenta) and EFF-1 (co-transfection) or IZUMO1-Venus (green). For IZUMO1-Venus, two independent fusion events are shown. Arrowheads and arrows indicate contacting and fused cells, respectively. Time (h:min) after the start of observation (see Movies S1-S3). Scale bars, 50 μm.

(B) Quantification of live imaging experiments in which BHK cells express RFPcyto, E-cadherin, EFF-1, GEX2, IZUMO1-Venus or JUNO. The percentage of fusion was defined as the ratio between the number of fusion events (Fe) and the number of transfected cells (Tc), as follows: % of fusion events = (Fe/Tc) x 100. Bar chart showing individual experiments and means ± SEM of at least three independent experiments. Total Tc analyzed = 1001 (RFP), 1179 (E-cadherin), 930 (EFF-1), 415 (GEX2), 817 (IZUMO1) and 1265 (JUNO). Comparisons by one-way ANOVA followed by Dunett's test against RFPcyto. ns = non-significant, * $p < 0.05$, ** $p < 0.01$.

Figure 4. Mutagenesis of IZUMO1 and expression of mutant proteins

(A) Structure of IZUMO1 (PDB: 5B5K, (Nishimura et al., 2016)) showing the four-helix bundle (in orange) containing four solvent-exposed aromatic residues (in red, F28, W88 and W113); the hinge (in light green) with the JUNO-interacting W148 (in blue); and the Ig-like domain (in teal).

(B) Schematic representation of wild-type IZUMO1 (WT) and the mutants maintaining the color coding for the different domains as in (A). The mutants with the deletion of the transmembrane domain and cytoplasmic tail (Ecto), the deletion of the Ig-like domain (ΔIg), the point mutation W148A and the triple mutant (F28A+W88A+W113A, FWW) are represented.

(C) Representative Western blot of total protein extract from BHK cells transfected with empty vector, a plasmid encoding for WT IZUMO1, or the mutants shown in (B). The different variants were detected with an anti-V5 antibody and actin was used as a loading control.

(D) Representative images of cells transfected with the pCI::H2B-RFP vectors encoding for WT IZUMO1 or the different mutants and subjected to an immunostaining (in white) after permeabilization with Triton X-100 using an anti-V5 antibody or without permeabilization using an antibody against the Izumo domain (Mab120). The signals for H2B-RFP and DAPI are shown in magenta and blue, respectively. The proportion of non-permeabilized cells showing surface expression was: IZUMO1^{WT} (41.9%, n=1013), IZUMO1^{ΔIg} (0%, n=1000), IZUMO1^{Ecto} (0%, n=1000), IZUMO1^{W148A} (60.3%, n=1005), IZUMO1^{FWW} (61.7%, n=995). Scale bars, 20 µm.

Figure 5. Functional characterization of IZUMO1 mutants

(A) Quantification of content-mixing in populations of cells expressing vectors, wild-type IZUMO1 (WT) and its mutants (Ecto, ΔIg, W148A and FWW, see Figure 4). Bar chart showing means ± SEM. n = 1000 nuclei per experiment. Comparisons by one-way ANOVA followed by Bonferroni's test against the vector (black) and against WT (red). ns = non-significant, * $p<0.05$, *** $p < 0.001$, **** $p< 0.0001$.

(B-C) Analysis of the binding ability of IZUMO1 (WT and mutants) to oocytes.

(B) Quantification of the binding of BHK cells to oocytes. Cells were transfected with pCI::GFPnes empty vector or encoding for IZUMO1, IZUMO1[W148A] or IZUMO1[FWW] and incubated with wild-type oocytes. The number of BHK cells bound per oocyte was determined; n = total number of oocytes analyzed in parenthesis. Bar chart showing means ± SEM. Comparisons by one-way ANOVA followed by Dunnett's test against the vector. ns = non-significant, * $p<0.05$.

(C) Representative images of oocytes from one experiment in (**B**) taken under a dissecting microscope (upper row) and a wide-field illumination microscope showing the merged DIC and GFP channels (lower row). Scale bar, 20 µm.

Graphical abstract

734 **STAR★ Methods**

736 **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-V5 tag monoclonal antibody	Thermo Fisher Scientific	Cat# R96025, RRID: AB_2556564)
Anti-FLAG M2 monoclonal antibody	Sigma	Cat# F3165, RRID: AB_259529
Anti-IZUMO1 Mab120 monoclonal antibody	Merck Millipore	Cat# MABT1357
Anti-Actin clone C4 monoclonal antibody	MP Biomedicals	Cat# ICN691001, RRID: AB_2335127
Anti-mouse IgG secondary antibody, Alexa Fluor 488	Thermo Fisher Scientific	Cat# A-21202, RRID: AB_141607
Anti-mouse IgG secondary antibody, Alexa Fluor 647	Thermo Fisher Scientific	Cat# A-21235, RRID: AB_2535804
Anti-Rat IgG secondary antibody, Alexa Fluor 488	Thermo Fisher Scientific	Cat# A-21470, RRID: AB_2535873
Anti-mouse IgG secondary antibody, HRP	Jackson ImmunoResearch Labs	Cat# 115-035-003, RRID: AB_10015289
Bacterial and virus strains		
DH5α competent cells	Thermo Fisher Scientific	Cat# 18265017
Chemicals, peptides, and recombinant proteins		
jetPRIME® transfection reagent	Polyplus Transfection	Cat# 101000046
Gibson Assembly® Cloning Kit	New England Biolabs	Cat# E5510
FdUdr, 5-Fluoro-2'-deoxyuridine	Sigma	Cat# F0503, CAS: 50-91-9
PMSG, pregnant mare serum gonadotropin	Prospect	Cat# HOR-272, CAS: 9002-70-4
hCG, human chorionic gonadotropin	Sigma	Cat# CG5, CAS: 9002-61-3
Mifepristone	Thermo Fisher Scientific	Cat# H11001, CAS: 84371-65-3

BSA, bovine serum albumin	Sigma	Cat# A7906, CAS: 9048-46-8
Hyaluronidase	Sigma	Cat# H3506, CAS: 37326-33-3
Experimental models: Cell lines		
BHK-21, clone 13	ATCC	Cat# CCL-10, RRID: CVCL_1915
BHK-21, clone 13	RCB	Cat# RCB1423, RRID: CVCL_1915
HEK293T	ATCC	Cat# CRL-3216, RRID:CVCL_0063
Experimental models: Organisms/strains		
B6D2.C57BL/6-Tg(Zp3-EGFP/Cd9)1Osb mouse line	Dr. Masahito Ikawa, (Miyado et al. 2008)	N/A
Oligonucleotides		
See Table S2 for complete list of oligonucleotides used in this study		
Recombinant DNA		
pCI::H2B-RFP	(Williams et al., 2018)	Addgene #92398
pCI::GFPnes	(Moi et al., 2021)	N/A
myr-EGFP	(Dunsing et al., 2018)	N/A
pCMV6-AC-IZUMO1-GFP	Origene	Origene #MG222708
Mouse ZP2 (JD#147)	(Liang et al., 1990)	Addgene #14645
pIZT::mZP2-V5-6HIS	This paper	N/A
pCAGGS::mZP2-V5-6HIS	This paper	N/A
pCAGGS::IZUMO1-V5-6HIS	This paper	N/A
pExpress1-JUNO	(Bianchi et al., 2014)	N/A
pExpress1-JUNO-flag	This paper	N/A
pCI::AtGCS1/HAP2-V5::H2B-RFP	(Moi et al., 2021)	N/A
pCI::AtGCS1/HAP2-V5::GFPnes	This paper	N/A
pCI::JUNO::H2B-RFP	This paper	N/A
pCI::JUNO::GFPnes	This paper	N/A

pCI::IZUMO1-V5-6HIS::H2B-RFP	This paper	N/A
pCI::IZUMO1-V5-6HIS::GFPnes	This paper	N/A
pGene/V5-His	Thermo Fisher Scientific	Cat# K106001
pSwitch	Thermo Fisher Scientific	Cat# K106001
pRFPnes	(Avinoam et al., 2011)	N/A
Mouse E-cadherin GFP	(Truffi et al., 2014)	Addgene #67937
pGENE::mCherry-JUNO	This paper	N/A
pGENE::IZUMO1-Venus	This paper	N/A
pGENE::CeEFF-1	(Valansi et al., 2017)	Addgene #132961
pGENE::AtGEX2-Venus	This paper	N/A
pCI::IZUMO1^{ecto}-V5::H2B-RFP	This paper	N/A
pCI::IZUMO1^{ecto}-V5::GFPnes	This paper	N/A
pCI::IZUMO1$^{\Delta Ig}$-V5::H2B-RFP	This paper	N/A
pCI::IZUMO1$^{\Delta Ig}$-V5::GFPnes	This paper	N/A
pCI::IZUMO1^{W148A}-V5::H2B-RFP	This paper	N/A
pCI::IZUMO1^{W148A}-V5::GFPnes	This paper	N/A
pCI::IZUMO1^{FWW}-V5::H2B-RFP	This paper	N/A
pCI::IZUMO1^{FWW}-V5::GFPnes	This paper	N/A
pMD2.G - VSV-G encoding plasmid	Didier Trono	Addgene #12259
Software and algorithms		
GraphPad Prism 9	GraphPad Prism	RRID:SCR_002798
FIJI (ImageJ 1.53c)	Image J	RRID:SCR_002285
Photoshop CS6	Adobe	RRID:SCR_014199
Illustrator CS6	Adobe	RRID:SCR_010279
ZEN microscopy software 7.0.4.0	ZEISS	RRID:SCR_013672
MetaMorph image analysis software 7.8.1.0	Molecular Devices	RRID:SCR_002368
Biorender	Biorender	RRID:SCR_018361
AlphaFold v2.1.0	(Jumper et al., 2021)	N/A

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Benjamin Podbilewicz (podbilew@technion.ac.il).

Data and code availability

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell lines and DNA transfection

In this study we used BHK (CCL-10; ATCC, Virginia, USA) for multinucleation, content mixing and BHK to oocyte interaction experiments and for evaluation of expression by Western blot and immunostaining; BHK (RCB1423; RIKEN Cell Bank, Tsukuba, Japan), for live imaging assays and HEK293T (CRL-3216; ATCC) cells for content mixing experiments. BHK and HEK293T cells were grown and maintained in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum (FBS). Cells were cultured at 37°C in 5% CO_2. Plasmids were transfected into cells using 2 µl jetPRIME (PolyPlus-transfection, Illkirch-Graffenstaden, France) per µg of DNA in 100 µl of reaction buffer for every ml of medium. For experiments with HEK293T cells, a coating with poly-L-lysine hydrobromide (Sigma, 20 µg/ml) was applied to the plates.

Mice

All animal studies were approved by the Committee on the Ethics of Animal Experiments of the Technion, Israel institute of Technology. B6D2.C57BL/6-Tg(Zp3-EGFP/Cd9)1Osb mice line (Miyado et al., 2008) was obtained from Dr. Masahito Ikawa (Osaka University, Japan) and animals were bred and housed in the Technion animal facility under specific pathogen-free conditions with *ad libitum* access to food and water. The primers used for genotyping are outlined in Table S2. Transgenic and wild type females between 2-6 months were used for the experiments.

METHOD DETAILS

DNA constructs

For the multinucleation and content mixing assays: Mouse *Izumo1* and *Juno* coding sequences were amplified from pCMV6-IZUMO1-GFP (MG222708, Origene, Rockville, MD, USA) and pExpress1-JUNO (Clone B2) plasmids, respectively, kindly

provided by Gavin Wright. The GFP tag of IZUMO1 construct was replaced with V5-HIS tags during cloning. *Izumo1-V5* and *Juno* sequences, and *Arabidopsis thaliana GCS1HAP2*-V5 were subcloned by restriction cloning into pCI::H2B-RFP and pCI::GFPnes vectors separately employing enzymes from Thermo Fisher Scientific (Invitrogen, Waltham, MA, USA). These bicistronic vectors translate for a nuclear RFP (H2B-RFP) or cytoplasmic GFP (GFPnes) after an IRES element (Internal Ribosome Entry Site). For mutagenesis of IZUMO1: i) IZUMO1^{ecto}: the ectodomain of IZUMO1 (M1-P312) was amplified; ii) IZUMO1$^{\Delta Ig}$: the upstream and downstream of Ig domain (G166-L253) were amplified independently from the full length IZUMO1 and fused together by overlap PCR; iii) IZUMO1^{W148A}: The W148 was mutated to Alanine by overlapping PCR; iv) IZUMO1^{FWW}: F28, W88 and W113 located in the Izumo domain were mutated to Alanines by overlapping primers and fused together. The folding of the mutants was corroborated by AlphaFold (Jumper et al., 2021). All mutants were ligated into pCI::H2B-RFP and pCI::GFPnes vectors for mixing assay. JUNO was tagged in the C-terminus by inserting an annealed oligo containing the FLAG sequence after the signal peptide into the BlpI restriction site. Oligonucleotides were obtained from Sigma-Aldrich or IDT and all constructs were verified by DNA sequencing (Macrogen). For the live imaging experiments: *Arabidopsis thaliana GEX2*, *Caenorhabditis elegans eff-1* and mouse *Izumo* or *Juno* sequences were amplified from cDNAs. To visualize the proteins, the fragments corresponding to *GEX2* and *Izumo1* were fused to fluorescent protein sequences by PCR and then cloned into pGENE B after double digestion with restriction enzymes using the Gibson assembly (NEB, #E5510, MA, USA). pGENE::mCherry-JUNO was made by ligating mCherry and pGENE::JUNO fragments (Nakajima et al., 2021). For inducible expression using mifepristone in BHK cells, we used the GeneSwitch System (Invitrogen). The complete lists of plasmids and primers used in this study are shown in Key Resource Table and Table S2, respectively.

Immunostaining and analysis of localization

BHK cells were grown on 24-well glass bottom tissue-culture plates or in tissue-culture plates with coverslips. 24 h after plating, cells were transfected and incubated for additional 24 h before proceeding to the immunostaining. For IZUMO1 (WT) and its mutants (IZUMO1^{ecto}, IZUMO1$^{\Delta Ig}$, IZUMO1^{W148A} and IZUMO1^{FWW}) encoded in the pCI::H2B-RFP vector cells were fixed with 4% PFA in PBS and, when indicated, permeabilized with 0.1% Triton X-100 in PBS. To detect the proteins immunofluorescence was performed with anti-V5 (1:500, R96025, Thermo Fisher Scientific) or anti-IZUMO1, clone Mab120 (1:500, MABT1357, Merck Millipore) followed by the secondary antibodies Alexa Fluor 488 goat anti-mouse (1:500, A21202, Thermo Fisher Scientific) and Alexa Fluor 488 chicken anti-rat (1:500, A21470, Thermo Fisher Scientific), respectively. For JUNO localization, pExpress1-JUNO-flag plasmid was transfected together with the empty pCI::H2B-RFP and the incubation with the first antibody anti-flag (1:1000, F3165, Sigma) was performed before fixation (for detecting surface protein) or after fixation and permeabilization (for

detecting total protein). Then, JUNO was probed using the secondary Alexa Fluor 488 goat anti-mouse. In all cases, the nuclei were stained with 1 µg/ml DAPI and micrographs were obtained using wide-field illumination using an ELYRA system S.1 microscope (Plan-Apochromat 20X NA 0.8; Zeiss).

Evaluation of multinucleation

BHK cells were grown on 24-well glass bottom tissue-culture plates. 24 h after plating, cells were transfected with pCI::myrGFP::H2B-RFP (myristoylated EGFP), pCI::GCS1/HAP-V5::H2B-RFP, pCI::IZUMO1-V5:H2B-RFP or pCI::JUNO::H2B-RFP vectors encoding for myristoylated EGFP (myrGFP, gray), GCS1/HAP2-V5, IZUMO1-V5 or JUNO. For JUNO a plasmid for myrGFP (gray) was co-transfected. 24 h post-transfection, 20 µM 5-fluoro-2'-deoxyuridine (FdUrd) was added to the plates to arrest the cell cycle and 24 h later, the cells were fixed with 4% PFA in PBS and processed for immunofluorescence using an anti-V5 antibody, as explained before. Micrographs were obtained using wide-field illumination using an ELYRA system S.1 microscope (Plan-Apochromat 20X NA 0.8; Zeiss). Multinucleation percentage (Figure S1A) was determined as the ratio between the number of nuclei in multinucleated cells (NuM) and the total number of nuclei in multinucleated cells and expressing cells that were in contact but did not fuse (NuC) as follows: (Num/(Nuc + Num)) × 100.

Content mixing experiments

BHK or HEK293T cells at 70% confluence in 35 mm plates were transfected with 1 µg pCI::H2B-RFP or pCI::GFPnes (empty vectors as negative controls); pCI::GCS1/HAP2-V5::H2B-RFP or pCI::GCS1/HAP2-V5::GFPnes (positive control); pCI::IZUMO1-V5::H2B-RFP or pCI::IZUMO1-V5::GFPnes; pCI::JUNO::H2B-RFP or pCI::JUNO::GFPnes. 4 h after transfection, the cells were washed 4 times with DMEM with 10% serum, 4 times with PBS and detached using Trypsin (Biological Industries). The transfected cells were collected, resuspended in DMEM with 10% serum, and counted. Equal amounts of H2B-RFP and GFPnes cells (1-1.25x10^5 each) were mixed and seeded on glass-bottom plates (12-well black, glass-bottom #1.5H; Cellvis) and incubated at 37°C and 5% CO_2. For IZUMO1, pCI::IZUMO-V5::GFPnes cells were also mixed with pCI::H2B-RFP or pCI::JUNO::H2B-RFP transfected cells. 18 h after mixing, 20 µM FdUrd was added to the BHK cells. The mixed cells were co-incubated for a total of 48 h after which they were fixed with 4% PFA in PBS and stained with 1 µg/ml DAPI. Micrographs were obtained using wide-field illumination using an ELYRA system S.1 microscope (Plan-Apochromat 20X NA 0.8; Zeiss). The percentage of mixing was defined as the ratio between the nuclei in mixed cells (NuM) and the total number of nuclei in mixed cells and fluorescent cells in contact that did not fuse (NuC), as follows: % of mixing = (NuM/(NuM+NuC)) x 100 (Figure S1A). 1000 nuclei (NuM+NuC) were counted in each independent repetition (experimental point). For immunostaining after the content mixing, cells were treated as explained above, using as the secondary antibody the Alexa Fluor 647 goat anti-mouse (A21235, Thermo Fisher Scientific).

Live imaging experiments

To evaluate fusion by live imaging, we transfected BHK cells with pGENE and
pSWITCH. 24 h after transfection, BHK cells were cultured at 5.0×10^4 cells/ml. 4 h
after transfection, the expression was induced by addition of 10^{-4} mM mifepristone.
3–4 h post-induction, images of the cells were acquired every 6 min for 12 h to record
cell-to-cell fusion, using a spinning disk confocal system (CellVoyager CV1000;
Yokogawa Electric, Tokyo, Japan) at a magnification of 10X (NA 0.40, 10×UPLSAPO;
Olympus, Tokyo, Japan) dry objective. The number of transfected cells and the
occurrence of fusion were evaluated. Image analyses were performed using CV1000
software (Yokogawa Electric) and the FIJI online tool was used to adjust the brightness
and contrast. The percentage of fusion was defined as the ratio between the number
of fusion events (Fe) and the number of transfected cells (Tc), as follows: % of fusion
= Fe/Tc (Figure S1A).

Western blots

BHK cells at 70% confluence in 35 mm plates were transfected with 1 µg pCl::GFPnes
(empty plasmid as negative controls); pCl::IZUMO1-V5::GFPnes; pCl::IZUMO1^{Ecto}-
V5::GFPnes, pCl::IZUMO1^{ΔIg}-V5::GFPnes, pCl::IZUMO1^{W148A}-V5::GFPnes or
pCl::IZUMO1^{FWW}-V5::GFPnes. 24 h post-transfection, cells were mixed with reducing
sample buffer (#S3401, Sigma) and incubated 5 min at 95°C. Samples were loaded
on a 10% SDS-PAGE gel and transferred to PVDF membrane. After blocking,
membranes were incubated with primary antibody anti–V5 mouse monoclonal
antibody (1:5,000) or anti-actin (1:2,000, ICN691001, MP Biomedicals) for 1 h at room
temperature and HRP-conjugated goat anti-mouse secondary antibody (1:10,000,
115-035-003, Jackson ImmunoResearch Labs) 1 h at room temperature. Membranes
were imaged by the ECL detection system using FUSION-PULSE.6 (VILBER).

Fusion of BHK cells to oocytes

BHK cells at 70% confluence in 35 mm plates were transfected with 1 µg
pCl::IZUMO1::H2B-RFP alone or together with pMD2.G (encoding for VSV-G) 24 h
before the collection of the oocytes. To induce ovulation, transgenic females were
treated with an i.p. injection of pregnant mare serum gonadotropin (PMSG; 5 IU;
#HOR-272, Prospec, Israel). followed by an i.p. injection of human chorionic
gonadotropin (hCG; 5 IU, #CG5, Sigma) 48 h later. Cumulus-oocyte complexes were
collected from the ampullae of induced females 12–15 h post-hCG administration in
mHTF medium (Kito et al., 2004). The oocytes were denuded from the cumulus and
the ZP by sequential treatment with 0.3 mg/ml hyaluronidase (H3506; Sigma) and acid
Tyrode solution (pH 2.5) (Nicolson et al., 1975). BHK cells were harvested with 0.05%
EDTA in PBS and washed once with mHTF. 15-20 ZP-free oocytes were incubated
with $5x10^4$ BHK cells for 15 min with occasional mixing and washed. In the case of
VSV-G-expressing cells, the oocytes were treated for 30 s with acid Tyrode solution

(pH 2.5) and washed again with mHTF. In all cases, the oocytes-cells complexes were
incubated for 5 h at 37°C and 5% CO_2. Oocytes were then fixed, stained with DAPI
and evaluated for the presence of H2B-RFP chromosomes within the cytoplasm
(Figure S1A), using a CSU-X1 spinning disk confocal (Yokogawa) on a Nikon Eclipse
Ti inverted microscope (Plan-Apochromat 60X NA 1.4, Nikon). Images were obtained
using an iXon3 EMCCD camera (ANDOR) through MetaMorph (Molecular Devices,
version 7.8.1.0).

Binding of BHK cells to oocytes

BHK cells at 70% confluence in 35 mm plates were transfected with 1 µg pCI::GFPnes,
pCI::IZUMO1::GFPnes, pCI::IZUMO1^{W148A}::GFPnes or pCI::IZUMO1^{FWW}::GFPnes 24
h before the collection of the oocytes. Oocytes were obtained from wild type females
as described above, incubated with the 5×10^4 BHK cells with occasional mixing for 15
min, washed and fixed. The average number of BHK in direct contact with the oocyte
was determined for each group. Micrographs were obtained using wide-field
illumination as described above.
917
918   **QUANTIFICATION AND STATISTICAL ANALYSIS**
919
920   **Statistics and data analysis**
921
922   Results are presented as means ± SEM. For each experiment we performed at least
923   three independent biological repetitions. To evaluate the significance of differences
924   between the averages we used one-way ANOVA as described in the legends
925   (GraphPad Prism 9). Figures were prepared with Photoshop CS6 and Illustrator CS6
926   (Adobe), BioRender.com and FIJI (ImageJ 1.53c).